Mechanical coupling coordinates microtubule growth

Bonnibelle K. Leeds, Katelyn F. Kostello, Yuna Y. Liu, Christian R. Nelson, Sue Biggins, and Charles L.
Asbury

Abstract

During mitosis, kinetochore-attached microtubules form bundles (k-fibers) in which many
filaments grow and shorten in near-perfect unison to align and segregate each chromosome. However,
individual microtubules grow at intrinsically variable rates, which must be tightly regulated for a k-fiber
to behave as a single unit. This exquisite coordination might be achieved biochemically, via selective
binding of polymerases and depolymerases, or mechanically, because k-fiber microtubules are coupled
through a shared load that influences their growth. Here, we use a novel dual laser trap assay to show
that microtubule pairs growing *in vitro* are coordinated by mechanical coupling. Kinetic analyses show
that microtubule growth is interrupted by stochastic, force-dependent pauses and indicate persistent
heterogeneity in growth speed during non-pauses. A simple model incorporating both force-dependent
pausing and persistent growth speed heterogeneity explains the measured coordination of microtubule
pairs without any free fit parameters. Our findings illustrate how microtubule growth may be
synchronized during mitosis and provide a basis for modeling k-fiber bundles with three or more
microtubules, as found in many eukaryotes.

Introduction

In the mitotic spindles of many eukaryotes, each chromosome is attached to a bundle of
microtubules (Kiewisz et al., 2022; McDonald et al., 1992; O'Toole et al., 2020; VandenBeldt et al., 2006)
that grow and shorten in unison, acting as a single unit to produce the forces necessary to move the
chromosome (Cimini et al., 2004). Microtubules are microns-long protein polymers composed of αβ-
tubulin dimers with their 'minus' ends oriented toward the spindle poles and their more dynamic 'plus'
ends oriented toward the chromosomes. Microtubule plus-ends, or tips, attach to chromosomes via
protein complexes called kinetochores to form 'k-fiber' bundles. Kinetochore-attached tips within a
single k-fiber remain very close to one another (within ~0.2 µm along the bundle axis) (O'Toole et al.,
2020) even as the bundle grows and shortens (Cimini et al., 2004) by the addition and loss of tubulin
subunits at each tip (Maddox et al., 2003). The number of microtubules per k-fiber varies across species;
human RPE1 cells have ten to fifteen microtubules per k-fiber, while marsupial PTK2 cells have more
than twenty, and cervid Indian Muntjac cells have up to sixty (O'Toole et al., 2020; Drpic et al., 2018).
Budding yeast lack k-fiber bundles, having just a single microtubule per kinetochore (Winey et al., 1995),
although microtubules attached to sister kinetochores on opposite sides of the yeast spindle must still
coordinate with each other. Uncovering how microtubule tips maintain a high degree of synchrony

within k-fibers of up to sixty microtubules is critical to understanding how chromosomes are accurately
segregated during normal mitosis and why even modest perturbations of kinetochore-attached
microtubule dynamics can lead to mis-segregation (Bakhoum & Compton, 2012; Bakhoum et al., 2009;
Ertych et al., 2014; Stumpff et al., 2014).

The tightly coordinated growth of microtubules within a k-fiber bundle is remarkable because
individually, microtubule tips exhibit highly variable growth speeds and switch stochastically between
distinct states of growth and shortening (Brouhard, 2015). Microtubule growth speed varies
considerably in purified tubulin systems (Figure 1a) (Cleary et al., 2022; Gildersleeve et al., 1992;
Mahserejian et al., 2022), in cell extracts (Geisterfer et al., 2020), and in live cells (Cassimeris et al., 1988;
Gaudenz et al., 2010; Yamashita et al., 2015). Part of this heterogeneity can be attributed to pausing,
when microtubules stochastically stop and start growing during periods of net growth (Amoah-Darko &
White, 2022; Mahserejian et al., 2022). How the intrinsic heterogeneity of microtubule growth is
coordinated to maintain sub-micrometer proximity between k-fiber tips remains a mystery.

One possible mechanism that might coordinate k-fiber microtubules is selective binding of
microtubule-regulatory proteins (Maiato, Rieder, et al., 2003; McAinsh & Meraldi, 2011; Tirnauer et al.,
2002). Many microtubule-associated proteins that are known to regulate growth *in vitro* are implicated
in modulating k-fiber dynamics and kinetochore movements in cells, including TOG-family proteins such
as XMAP215 (Brouhard et al., 2008) and CLASP (Al-Bassam et al., 2010; Maiato, Fairley, et al., 2003;
Sousa et al., 2007), mitotic kinesins such as MCAK and Kif18A (Sanhaji et al., 2011; Stumpff et al., 2012),
and other plus-end-trackers such as EB1 (reviewed in (Amin et al., 2019)). The localization of some of
these regulators is enriched near kinetochores specifically when a kinetochore is moving anti-poleward
and its k-fiber is growing longer (Amaro et al., 2010; Castrogiovanni et al., 2021; Tirnauer et al., 2002).
However, whether such selective enrichment is sufficient to explain the very tight coordination across
many individual microtubule tips within a k-fiber is unclear.

Another, simpler mechanism is suggested by the fact that microtubule dynamics are also strongly
affected by mechanical force. Tension applied to individual microtubules *in vitro* biases them toward
growth (Figure 1a) (Akiyoshi et al., 2010), while compression biases them toward shortening (Dogterom
& Yurke, 1997; Janson et al., 2003). Theoretically, these mechanical effects should promote
synchronization of any group of microtubules whose tips, like the tips of kinetochore-attached
microtubules within a k-fiber, are all coupled to the same load (Banigan et al., 2015; Schwietert &
Kierfeld, 2020; Skibbens et al., 1993). To illustrate how, consider a hypothetical k-fiber composed of just
two microtubules sharing a total load, F_{TOT}. The load is transmitted to the microtubule tips through the
kinetochore, which has elasticity (Cojoc et al., 2016; Renda et al., 2020) that we represent as spring-like
connections from the kinetochore to each tip (Figure 1b). Initially, the two microtubules might have
equal lengths and share the load equally – each one carrying half of F_{TOT} (Figure 1b, step 1). If one tip
happens to grow more slowly than the other, then its spring will be stretched more, and it will start
carrying more tension than the other tip (Figure 1b, step 2). Conversely, the faster-growing tip will start
carrying less tension. Because tension accelerates microtubule growth, the lagging tip will tend to

accelerate, while the leading tip will tend to slow down, such that, on average, they return to having
equal lengths and tensions (Figure 1b, step 3).

Mechanical coupling was proposed decades ago as an explanation for coordination between sister
k-fibers (Gardner et al., 2005; Khodjakov et al., 1996; Skibbens et al., 1995; Skibbens et al., 1993; Wan et
al., 2012), but how effectively it can coordinate microtubules within a k-fiber has not been determined.
Compressive forces can cause a bundle of microtubules growing *in vitro* to switch cooperatively into a
shortening state (Laan et al., 2008), but k-fiber bundles are usually under tension, not compression
(Khodjakov & Rieder, 1996; Waters et al., 1996). Theoretical studies have explored the effects of shared
tensile and compressive loads on cooperative switching and bi-stability in microtubule bundles (Banigan
et al., 2015; Das et al., 2014; Ghanti & Chowdhury, 2015; Schwietert & Kierfeld, 2020; Zelinski & Kierfeld,
2013). However, these studies have generally neglected any differences in growth speed between
microtubules and therefore cannot address how much variability exists, nor how such variability is
suppressed. Furthermore, to our knowledge, the possibility that part of this variability (and suppression
thereof) can be attributed to tension-dependent pausing during microtubule growth has not been
explored.

Here, we use a novel dual laser trap assay to test directly how well a constant shared load can
coordinate a pair of microtubules growing *in vitro*. We show that when microtubule pairs are
mechanically coupled to a single load through a material of sufficient stiffness, their growth is tightly
coordinated. By re-analyzing recordings of individual microtubules growing under constant force (from
(Akiyoshi et al., 2010)) and examining their stochastic pausing kinetics, we show that pausing is
suppressed by tension. We then develop a theoretical model, based on the measured behaviors of
individual growing microtubules, to describe both the natural variability and force dependence of
growth using no free fit parameters. Simulations based on this model fit the dual-trap data well.
Altogether, this work illustrates how mechanical coupling may coordinate microtubule growth and
provides a basis for modeling k-fibers of three or more microtubules under a shared load.

Results

Dual-trap assay to measure coordination between paired microtubules *in vitro*

To study how mechanical coupling affects microtubule growth, we developed a novel technique
which we call the dual-trap assay (Figure 2). The dual-trap assay is based on our previously developed
single-trap (force-clamp) assay (Akiyoshi et al., 2010; Miller et al., 2016; Sarangapani et al., 2013), in
which dynamic microtubules are grown from stabilized seeds bound to a biotinylated coverslip. Using
the single laser trap, we attach an individual bead decorated with isolated yeast kinetochores to the
growing plus-end of a single microtubule. A computer then continuously measures the bead position
and adjusts the trap to exert a precise, constant level of tension on the microtubule via the kinetochore-
decorated bead. Under this persistent feedback-controlled tension, kinetochore-beads typically track
with the microtubule tips even as the tips stochastically grow and shorten.

Our new dual-trap assay uses two separate laser trapping microscopes, located adjacent to one
another in the same room and connected to a single computer. On each of the two instruments, we
attach a kinetochore-decorated bead to a dynamic microtubule plus-end. The computer then
simultaneously monitors and controls the forces on both microtubules. Rather than keeping the force
constant on each microtubule, the computer adjusts the forces dynamically to simulate an elastic
coupling of both plus-ends to a single shared load (Figure 2a). Thus far, we have simulated only purely
elastic couplers, where both coupling springs are linear (Hookean) with stiffness, κ (Figure 2b). In the
future, more complex coupling materials with viscoelasticity (Cojoc et al., 2016) or strain-stiffening
(Schwietert et al., 2022; Volkov et al., 2018) could also be simulated.

To begin a dual-trap experiment, we first choose the spring stiffness, κ, and the total shared
load, F_{TOT}, which are kept constant (Figure 2a, step 1). After a kinetochore-decorated bead is attached
to a growing plus-end on each of the two instruments, feedback-control is initiated and the two plus-
ends are arbitrarily considered to be parallel, with tips side-by-side (i.e., both at $x_1 = x_2 = 0$) and sharing
the load equally ($F_1 = F_2 = \frac{1}{2} \cdot F_{TOT}$) (Figure 2a, step 2). Because microtubule growth is intrinsically
variable, the two microtubules subsequently grow at different speeds (Figure 2a, step 3). The computer
then dynamically monitors the bead positions (Figure 2a, step 4) and adjusts the forces, F_1 and F_2
(Figure 2a, step 5), according to the elastic coupling model (Figure 2b). For purely elastic couplers, the
force difference across the two microtubules equals the tip separation, $(x_2 - x_1)$, multiplied by the
coupling stiffness (κ). When one microtubule grows more quickly than the other, tension on the leading
(faster-growing) microtubule decreases and tension on the lagging (slower-growing) microtubule
increases to maintain a constant total force on the pair, F_{TOT}. Forces in our dual-trap feedback system
are updated at a rate of 50 Hz, which is more than sufficient for accurately simulating the elastic
coupling between two microtubules (Figure S3-1). For all the experiments described below, $F_{TOT} = 8$ pN.
Thus, each microtubule generally experiences between 1 and 7 pN of tension, close to the estimated
average of 4 to 6 pN experienced by kinetochore-microtubules in budding yeast (Chacon et al., 2014;
Parmar et al., 2023).

Mechanical coupling coordinates microtubule pairs *in vitro*

To measure the effects of mechanical coupling, we recorded the growth of many pairs of
microtubules coupled to a constant shared load ($F_{TOT} = 8$ pN) either through relatively soft springs, with
$\kappa = 1$ pN·µm^{-1}, or through springs with five-fold greater stiffness, $\kappa = 5$ pN·µm^{-1}. These two stiffnesses
span a range near the minimum elasticities estimated for mitotic chromosomes (Nicklas, 1983; Poirier et
al., 2002), for pericentromeres (Chacon et al., 2014), and for elements of the kinetochore (Schwietert et
al., 2022; Volkov et al., 2018). When coupled through soft springs ($\kappa = 1$ pN·µm^{-1}), microtubule pairs
tended to grow at different rates and their tips often became separated by more than a micrometer
(Figure 3a, left, and Figure S3-1, left). After 300 s, the average tip separation for soft-coupled pairs grew
to 0.8 ± 0.2 µm (mean ± SEM from $N = 10$ pairs) (Figure 3b, left). When coupled through stiff springs ($\kappa =$
5 pN·µm^{-1}), growth speeds were better matched, and the microtubule tips remained closer together

(Figure 3a, right, and Figure S3-1, right). The average tip separation for stiff-coupled pairs after 300 s was
0.25 ± 0.06 μm (mean ± SEM from N = 10 pairs) (Figure 3b, right), significantly lower than for soft-
coupled pairs (p = 0.03 based on a Wilcoxon-Mann-Whitney two sample rank test).

Because the plus-ends of microtubules within k-fiber bundles are rarely separated by more than
0.8 μm from each other *in vivo* (O'Toole et al., 2020), we used Kaplan Meier survival analysis to ask how
often the pairs of coupled microtubule tips in our experiments exceeded this threshold separation.
Consistent with their lower average tip separation, stiff-coupled microtubule pairs were much less likely
to exceed a separation of 0.8 μm than their soft-coupled counterparts (p = 0.01, log-rank test) (Figure
3c), which sometimes separated by 0.8 μm in as little as 60 s. Even after 400 s, more than 80% of the
stiff-coupled pairs had tip separations that remained below 0.8 μm, while fewer than half of the soft-
coupled pairs had tip separations below 0.8 μm. (Similar survival analyses using thresholds of 0.4, 0.6,
and 1.2 μm are provided in Supplemental Figure S6-2.)

Even at shorter time periods, the average behavior of soft- and stiff-coupled microtubule pairs
was distinct. During the first 100 s of growth, the average rate of tip separation for soft-coupled pairs
was 0.23 ± 0.04 μm·min^{-1} (mean ± SEM from N = 23 pairs), while the average rate of tip separation for
stiff-coupled pairs was only 0.14 ± 0.02 μm·min^{-1} (mean ± SEM from N = 26 pairs; p = 0.03 based on a
Wilcoxon-Mann-Whitney two sample rank test). Notably, the average rate of tip separation for stiff-
coupled microtubule pairs decreased as their average tip separation increased, suggesting they were
approaching a steady-state level of separation (Figure 3b, right). In contrast, the average rate of tip
separation for soft-coupled microtubule pairs remained approximately constant throughout the 400 s
measurement, suggesting that soft-coupled pairs would require more time to approach a steady state.
Altogether, these observations show that the growth of microtubule pairs can be tightly coordinated,
even over relatively short (< 100 s) timescales, by mechanically coupling their tips to a shared load
through couplers of sufficient stiffness.

Models of microtubule growth variability and force sensitivity can be derived from recordings of
individual microtubules

Coordination of microtubule growth can be thought of as the suppression of variability in
microtubule growth speed. Understanding how mechanical coupling suppresses this variability requires
a realistic model of how the growth of individual microtubules varies over time and between
microtubules, as well as how growth behaviors are influenced by force. Previous simulations of
microtubule bundles (Banigan et al., 2015; Schwietert & Kierfeld, 2020), based on force-clamp
measurements of individual microtubules (Akiyoshi et al., 2010), have suggested that mechanical
coupling might coordinate the stochastic switching of microtubules between shortening and growth,
such that sister k-fibers are nearly always kept in opposite states. However, these models assumed
identical growth speeds for all microtubules under the same force, and therefore cannot address how
mechanical coupling might coordinate differences in growth speed. We sought to create a model of

microtubule growth that explains both the variability in microtubule growth speeds and how that
variability is suppressed by sufficiently stiff mechanical coupling, as observed in the dual-trap assay.

To characterize the variability and force dependence of microtubule growth, we re-examined
the force-clamp recordings by Akiyoshi et al., 2010. This large dataset (N = 356 recordings) was recorded
with an identical setup to our dual-trap experiments, except that each microtubule was grown
individually with a single laser trap exerting a constant level of tension (between 0.5 and 18 pN). Using
this dataset, we created two models derived from the natural variability of individual kinetochore-
attached microtubules and their average behaviors as a function of force. Below, we begin by briefly
describing a simple non-pausing model, which was developed first and served as a foundation for our
second, more successful pausing model. The pausing model reasonably explains our dual-trap results by
incorporating both stochastic pausing and intrinsic differences in microtubule growth speed.

A simple non-pausing model predicts larger tip separations than measured *in vitro*

When averaged across time and across many individual microtubules, mean growth speed
increases exponentially with applied tension (Akiyoshi et al., 2010), such that

$$v_g = v_o * exp\left(\frac{F}{F_g}\right) \qquad [1]$$

where v_o represents the unloaded growth speed, F represents the applied tension, and F_g represents
the sensitivity with which the growth speed increases with tension. However, individual microtubules
show considerable variability around this mean behavior, with some microtubules remaining
persistently slower than others for many minutes (Figure 1a). In our first model, we adopted the simple
assumptions that this variability stems entirely from intrinsic differences in unloaded microtubule
growth speed, v_o, and that all microtubules have identical force-sensitivity (with F_g = 8.4 pN, based on
the exponential curve that best fit the mean growth speed-vs-force data (Akiyoshi et al., 2010)). In
principle, such static heterogeneity in growth speed might arise from persistent differences in
microtubule tip structure (Cleary & Hancock, 2021; Mahserejian et al., 2022; Rai et al., 2021) or from
variation in the numbers of protofilaments per microtubule (Chretien et al., 1992; Rai et al., 2021).

For each simulated microtubule, we randomly chose an individual growth speed measurement
from the force-clamp dataset (Akiyoshi et al., 2010), then extrapolated the microtubule's intrinsic
unloaded speed (v_o) from the chosen point (v_g, F) using equation [1] with F_g = 8.4 pN. The growth
speed of this simulated microtubule then followed the extrapolated curve as a function of force. To
simulate a pair of coupled microtubules, we made two such random draws from the dataset, defining
two distinct exponential (v_g-versus-F) relations for the two microtubules. (Bundles of three or more
microtubules could be simulated in the same manner by making additional draws.) We used a simple
Monte Carlo-based method (Gardner & Odde, 2010) to simulate traces of tip position versus time for
the pair. As in our dual-trap experiments, the simulated tip pairs were coupled with either soft or stiff

elastic springs to a constant shared load, F_{TOT} = 8 pN. Both the variability and force-dependence of
growth in the model were derived directly from the experimental force-clamp data, with no adjustable
('free') parameters.

Despite its direct derivation from force-clamp data, the non-pausing model could not explain the
behavior of microtubule pairs measured in the dual-trap assay (Figure 4). Simulated microtubule pairs
developed much larger tip separations than real microtubule pairs over time, implying that this model
overestimates real microtubule variability. The simulated microtubule pairs showed especially poor
agreement with real microtubule pairs under the stiff-coupled condition (Figure 4c, red curves),
indicating that the non-pausing model also does not capture the degree to which real microtubules are
coordinated by mechanical coupling.

Microtubule growth is interrupted by stochastic, force-dependent pauses

Our simple non-pausing model lacked a notable behavior exhibited by microtubules that could
influence their response to tension: Microtubule growth is frequently interrupted by stochastic pauses
of varying durations, both in cells (Gaudenz et al., 2010; Sousa et al., 2007) and in purified tubulin
systems (Mahserejian et al., 2022; Stumpff et al., 2012). Indeed, pauses recorded during our force-clamp
(Figures 1a and S4-1b) and dual-trap assays (Figures 3a and S3-1) were frequent and often lasted tens of
seconds. Presumably, these pauses occur because the configuration of the growing microtubule tip
fluctuates and sometimes becomes transiently refractory to the incorporation of new tubulin subunits
(Cleary & Hancock, 2021; Mahserejian et al., 2022). We hypothesized that entry and exit into the pause
state might be force-dependent processes, and that pausing might therefore be a key aspect of
microtubule force-sensitivity.

To test whether pause entry and exit rates vary with force, we reanalyzed the force-clamp
dataset (Akiyoshi et al., 2010), applying a 2 s sliding window to identify intervals during each recording
(N = 356) when the instantaneous growth speed fell to near-zero, indicative of a pause (Figure S5-1; see
Materials and Methods for details about our pause marking procedure). The intervals in-between these
pauses were designated as 'runs.' Distributions of run and pause durations were approximately
exponential (Figure S5-2), consistent with a relatively simple two-state kinetic model of microtubule
growth (Figure S5-1a), in which pause entry and exit are considered Poisson processes. For each force
level, we estimated the pause entry and exit rates, k_{EN} and k_{EX}, by counting the total number of
observed entrances into and exits out of pause and dividing by the total time spent in run and pause,
respectively (Figure S5-1a).

This analysis revealed that tension delays pause entry and accelerates pause exit (Figure 5a).
Both effects tend to decrease the time spent in the pause state for growing microtubule tips under high
tension. In addition, the average growth speed during runs was accelerated by tension (Figure 5b). All
three rates, pause entry, pause exit, and average growth speed during runs, were well-fit by single
exponential functions of force (Figure 5; see Table 1 for fit parameters). These observations show that

stochastic pausing during microtubule growth is force-dependent and imply that pausing kinetics will
strongly influence the response of growing microtubules to mechanical coupling.

A pausing model can explain the variability between paired microtubule and their coordination by
mechanical coupling

To examine how force-dependent pausing might affect the behavior of coupled microtubules,
we created a second model in which growing microtubules switch stochastically between states of
pause and run. Pause entry and exit were considered Poisson processes with kinetic rates dictated by
the exponential fits of k_{EX} and k_{EN} versus force estimated from the force-clamp dataset (Figure 5a,
dashed curves; Table 1). Intrinsic heterogeneity among microtubules was incorporated by randomly
choosing an individual run speed measurement for each simulated microtubule from the force-clamp
dataset (i.e., by randomly choosing one point in Figure 5b). We then extrapolated the microtubule's
intrinsic unloaded run speed (v_{RUN}^o) from the chosen point (v_{RUN}, F) by assuming an exponential force-
dependence with sensitivity F_{RUN} = 14.8 pN (based on the dashed curve in Figure 5b; Table 1). The
growth speed during runs for this simulated microtubule then followed the extrapolated curve as a
function of force. We used a simple Monte Carlo-based method (again) to simulate traces of
microtubule tip position versus time. All model parameters were derived directly from the experimental
force-clamp data, with no adjustable ('free') parameters. As in our dual-trap experiments, simulated
microtubule tip pairs were coupled with either soft or stiff elastic springs to a constant shared load, F_{TOT}
= 8 pN. Simulations of individual microtubules growing under constant force closely resembled force-
clamp recordings of real microtubules *in vitro*. (Compare Figure S6-1 to Figure 1a.)

The pausing model of microtubule growth provided significantly better fits to our dual-trap data
than the non-pausing model. Like real microtubules in the dual-trap experiments, the simulated pairs of
tips tended to drift apart under soft-coupled conditions, while under stiff-coupled conditions they
stayed closer together (Figure 6a). After 300 s, the average tip separation for simulated soft-coupled
pairs grew to 0.79 ± 0.01 μm (mean ± SEM from N = 10,000 simulations), while simulated stiff-coupled
pairs had a lower average tip separation of 0.43 ± 0.005 μm (mean ± SEM from N = 10,000 simulations)
(Figure 6b), similar to the corresponding separations measured in dual-trap experiments (0.8 ± 0.2 μm
and 0.25 ± 0.06 μm, respectively). Simulated pairs of microtubule tips under stiff-coupled conditions
were also less likely to exceed a threshold separation of 0.8 μm (Figure 6c, dashed curves), similar to the
survival analyses of microtubule pairs measured with the dual-trap assay (Figure 6c, solid curves).
(Survival analyses of simulated microtubule pairs with threshold separations of 0.4, 0.6, and 1.2 μm also
resembled the survival analyses of microtubule pairs measured with the dual-trap assay. See Figure S6-
2.) In addition, the pausing model scored better than the non-pausing model when we used probability
to compare simulated to *in vitro* measurements (Figure S6-4). (To quantify the variability due to
stochastic pausing alone, we also compared the pausing model to a version in which we assumed that
microtubules were intrinsically identical ('ergodic'). The low variability predicted by this model (Figure
S6-3), particularly under the soft-coupled condition, indicates that stochastic pausing alone is

insufficient to generate the large and persistent differences in growth speed exhibited by real
microtubules). Overall, the strong agreement between the pausing model and the dual-trap recordings
suggests that mechanical coupling coordinates microtubule growth in at least two distinct ways: first, by
counteracting persistent (intrinsic) differences in microtubule growth speed, and second, by selectively
suppressing stochastic pausing in the lagging tip relative to the leading tip.

302   **Discussion**
Our work demonstrates the effectiveness of mechanical coupling for coordinating the growth of
microtubules and helps explain how numerous filaments within a k-fiber bundle can be tightly regulated
to grow in near-perfect unison. Using our dual-trap assay, we found that pairs of microtubule tips
growing *in vitro* are kept in close proximity when sharing a single load through a coupling material of
sufficient stiffness. Tip pairs in our experiments were kept within a fraction of a micrometer by couplers
of only κ = 5 pN·μm^{-1}, a modest stiffness within the ranges reported for elements of the kinetochore
(Schwietert et al., 2022; Volkov et al., 2018), pericentromeric chromatin (Chacon et al., 2014), and
mitotic chromosomes (Nicklas, 1983; Poirier et al., 2002). Because the kinetochore and its underlying
centromeric chromatin both appear to deform together to maintain close contacts with k-fiber plus-
ends *in vivo* (O'Toole et al., 2020), the plus-ends presumably exert forces on each other through a
composite material composed of both kinetochore and chromatin. We propose that this mechanical
coupling is a major coordinating influence on plus-end growth dynamics within k-fiber bundles.

Our dual-trap approach provides information that prior single-trap experiments, and theoretical
work based solely on these prior measurements, cannot provide. Microtubules in real k-fiber bundles
presumably adjust their growth speeds dynamically in response to the forces they exert on each other.
The dual-trap assay uniquely allows this kind of mutual dynamic influence to be examined. Because each
microtubule in the pair grows simultaneously in the same preparation of buffer, any differences in
growth speed represent true biological variability, without the potentially confounding influences of
prep-to-prep or sample-to-sample variability. While previous experiments with individual microtubules
implied that a shared load should tend to coordinate growing microtubules, the dual-trap experiments
are a direct, model-independent way to examine the extent to which load sharing can suppress the
intrinsic variability of microtubules growing under identical conditions. With the dual-trap assay, we can
test models and assumptions about microtubule tension-dependence and variability derived from
recordings of individual microtubules, as we did with the pausing and non-pausing models presented
here.

Since most known eukaryotes have k-fibers of more than two microtubules, an important area
of future work will be to study how effectively mechanical coupling can coordinate k-fibers with more
microtubules. While extending our *in vitro* laser trap methods to more than two microtubules is
impractical, our model can be easily extended to explore mechanical coordination within k-fibers of
three or more microtubules. Simulation parameters, such as microtubule growth speeds, can also be
changed to explore how well mechanical coupling coordinates microtubules in different cell types or cell
cycle stages with varying microtubule dynamics (Cassimeris et al., 1988; Gaudenz et al., 2010; Yamashita

et al., 2015; Rusan et al., 2001). Similarly, simulations can be used to study how the addition of
microtubule polymerases or depolymerases with known effects on microtubule dynamics influences
coordination.

A variety of microtubule-modifying proteins are implicated in regulating k-fiber dynamics (reviewed
in (Ou & Scholey, 2022) and (Amin et al., 2019)). These biochemical factors must work in concert with
mechanical effects, and we suggest that the two classes of mechanisms probably operate over different
length- and timescales. Certain tip-tracking proteins capable of influencing microtubule dynamics are
enriched at growing k-fiber ends specifically near kinetochores that are moving anti-poleward (Amaro et
al., 2010; Castrogiovanni et al., 2021; Tirnauer et al., 2002). Various kinesins are known to regulate the
size of the metaphase spindle (Goshima et al., 2005; Neahring et al., 2021), to influence k-fiber dynamics
(Maiato, Fairley, et al., 2003; Rizk et al., 2009; Stumpff et al., 2012; Wordeman et al., 2007), and to alter
microtubule dynamics *in vitro* in a length-dependent manner (Varga et al., 2006). However, it is unclear
whether the specific enrichment of such microtubule-modulators on longer or faster-growing tips could
occur quickly enough and with enough selectivity to slow the growth of a single tip that has grown
ahead of its neighbors by only a fraction of a micrometer, as required for tight coordination within a k-
fiber. On the other hand, mechanical forces would be transmitted essentially instantaneously between
neighboring tips through local deformations of the material through which the tips are coupled.
Therefore, mechanical coupling could account for the rapid, sub-micron responses required to
coordinate microtubule tips within a k-fiber, while microtubule-modifiers and spindle-wide phenomena,
like polar ejection forces and poleward flux, are likely to help set overall k-fiber length (Risteski et al.,
2021).

We found that tension accelerates microtubule growth not only by directly increasing growth speed,
but also by decreasing time spent in pause. More specifically, tension reduced the rate at which growing
tips entered a pause state. This finding implies that pause entry is associated with movement of tension-
bearing elements at the tip toward the minus end, opposite the direction of applied force and therefore
inhibited by the force. Tension also increased the rate at which paused tips resumed growth, which
implies that pause exit is associated with movement toward the plus-end, in the same direction as the
applied force and therefore assisted by the force. These tension-dependent dynamics may reflect
mechanical deformations of the microtubule tip; perhaps pauses occur when newly added tubulin
dimers at a growing tip adopt curved structures that disfavor lateral bonding (McIntosh et al., 2018).
Tension could help straighten these curved structures, thereby accelerating lateral bond formation and
favoring growth (Cleary & Hancock, 2021).

We derived a model of microtubule growth with no free parameters, based on constant-force
recordings of individual microtubules (from (Akiyoshi et al., 2010)), and found that a combination of
stochastic, force-dependent pausing and intrinsic differences in microtubule growth speed explained
both the variability and the coordination measured in our dual laser trap assay. The necessity of
including microtubules with different intrinsic growth speeds for our model's success implies structural
heterogeneity in the growing tips that must persist for many minutes, even as thousands of tubulin
subunits are being added. In principle, these persistent differences might arise from different or

mismatched protofilament numbers (Chretien et al., 1992; Rai et al., 2021) or the accumulation of other
lattice defects (Wieczorek et al., 2015) that result in varying degrees of tip taper (Coombes et al., 2013)
(or tip "raggedness" (Brouhard & Rice, 2018)) which are more or less conducive to microtubule growth.

For technical reasons, we focused exclusively here on microtubule tips in a state of net growth.
However, coordination of k-fiber microtubules *in vivo* presumably also requires tight regulation of tip
shortening speeds and rates of switching between growth and shortening (i.e., 'catastrophe' and
'rescue' rates) (Banigan et al., 2015; Schwietert & Kierfeld, 2020). These will be more difficult to study
with our dual-trap assay because switch events are stochastic and episodes of shortening are observed
less frequently than episodes of growth under our assay conditions. Nevertheless, the dual-trap
approach can be used in the future (potentially with a cutting laser to sever the microtubule tips to
induce shortening (Driver et al., 2017; Murray et al., 2022)) to study how mechanical coupling affects
microtubule pairs in which one or both filaments are shortening, as well as switch rates between
shortening and growth. Our dual-trap approach can also be adapted to examine mechanical coupling
through materials with complex mechanical properties, such as viscoelasticity and strain-stiffening
(Cojoc et al., 2016; Schwietert et al., 2022). In principle, the effect of mechanical coupling on pairs of
processive motor proteins (Holzbaur & Goldman, 2010) or on pairs of dynamic actin filaments linked to a
shared load (Garner & Theriot, 2022) can also be studied with the dual-trap approach. Overall, our work
demonstrates that mechanical coupling is a feasible mechanism that could coordinate k-fiber
microtubules and maintain their synchronous growth in the mitotic spindle.

Funding and Acknowledgements

We are grateful for feedback and critical reading of the manuscript from members of the
Biggins, Asbury, and Trisha Davis laboratories.

This work was supported by a Packard Fellowship 2006-30521 (to C.L. Asbury), National
Institutes of Health grants R01GM079373, P01GM105537, and R35GM134842 (to C.L. Asbury) and
R01GM064386 (to S. Biggins), and by the Genomics and Scientific Imaging and the Proteomics and
Metabolomics Shared Resources of the Fred Hutchinson/University of Washington Cancer Consortium
(P30 CA015704). S. Biggins is an investigator of the Howard Hughes Medical Institute. Research reported
in this publication was also supported by the National Institute of General Medical Sciences of the
National Institutes of Health under Award Number T32GM008268 (to B.K. Leeds).

Author contributions: Conceptualization (C.L. Asbury, B.K. Leeds), data curation (B.K. Leeds, Y. Liu),
formal analysis (B.K. Leeds, Y. Liu), funding acquisition (C.L. Asbury, S. Biggins, B.K. Leeds) investigation
(B.K. Leeds, K.F. Kostello, C.L. Asbury), project administration (C.L. Asbury), resources (C.L. Asbury, C.R.
Nelson, S. Biggins), software (B.K. Leeds, C.L. Asbury, Y. Liu), supervision (C.L. Asbury, S. Biggins, B.K.
Leeds), validation (B.K. Leeds, K.F. Kostello, Y. Liu, C.L. Asbury), visualization (B.K. Leeds, C.L. Asbury, K.F.
Kostello, Y. Liu), writing–original draft (B.K. Leeds, C.L. Asbury), writing–review and editing (B.K. Leeds,
C.L Asbury, K.F. Kostello, S. Biggins).

Methods

Spring-coupler model

For both *in vitro* and *in silico* experiments, the force on each microtubule plus-end was dictated
by a model in which plus-ends were coupled to a shared "kinetochore" via simple Hookean springs of
identical stiffnesses (Figure 2b). Because k-fibers grow relatively slowly in cells (Khodjakov et al., 1996;
Skibbens et al., 1993; Stumpff et al., 2012), we neglect viscous drag such that the sum of the forces on
individual k-fiber tips equals F_{TOT}. Microtubule pairs were considered to grow along a shared 1D axis.
Microtubules were assumed to be incompressible because microtubules have high axial stiffness
(Howard, 2001; Kis et al., 2002; Pampaloni et al., 2006) and are cross-linked in bundles (O'Toole et al.,
2020), thus forming the structural support required to maintain mitotic spindle integrity. Microtubules
were considered stationary except at the plus-end, so that only growth or shortening at the plus-end
affected plus-end position.

Kinetochore purification

Native kinetochore particles were purified from asynchronously grown *S. cerevisiae* SBY8253
cells (Genotype: MATa DSN1-6His-3Flag:URA3) grown in YPD medium (2% glucose) by modifying
previous protocols (Akiyoshi et al., 2010; Miller et al., 2016). Protein lysates were prepared using a
Freezer Mill (SPEX SamplePrep) submerged in liquid nitrogen. Lysed cells were resuspended in buffer H
(25 mM Hepes, pH 8.0, 2 mM MgCl2, 0.1 mM EDTA, 0.5 mM EGTA, 0.1% NP-40, 15% glycerol, and 150
mM KCl) containing phosphatase inhibitors (0.1 mM Na-orthovanadate, 0.2 µM microcystin, 2 mM β-
glycerophosphate, 1 mM Na pyrophosphate, and 5 mM NaF) and protease inhibitors (20 µg/ml
leupeptin, 20 µg/ml pepstatin A, 20 µg/ml chymostatin, and 200 µM PMSF). Lysates were
ultracentrifuged at 98,500 g for 90 min at 4°C. Protein G Dynabeads (Invitrogen) were conjugated with
an α-FLAG antibody (Sigma-Aldrich), and immunoprecipitation of Dsn1-6His-3Flag was performed at 4°C
for 3 h. Beads were washed once with lysis buffer containing 2 mM DTT and protease inhibitors, three
times with lysis buffer with protease inhibitors, and once in lysis buffer without inhibitors, and
kinetochore particles were eluted by gentle agitation of beads in elution buffer (buffer H + 0.5 mg/ml
3FLAG Peptide [Sigma-Aldrich]) for 30 min at room temperature.

Bead functionalization and slide preparation for laser trap experiments

To functionalize laser trapping beads with native kinetochore particles, streptavidin-coated
polystyrene beads (0.44 µm in diameter, Spherotech) were incubated with biotinylated anti-5His
antibodies (Qiagen) and stored for up to 6 months with continuous rotation at 4°C in BRB80 (80 mM
Pipes, 1 mM MgCl$_2$, and 1 mM EGTA, pH 6.9) supplemented with 8 mg·mL^{-1} bovine serum albumin and 1
mM DTT. Before each experiment, functionalized beads were sonicated for two minutes and decorated
with kinetochore particles by incubating 6 pM anti-5His beads for 1-2 hours at 4°C with purified

kinetochore material, corresponding to concentrations of Dsn1-His-Flag between 1.5 and 4 nM, well
within the single-kinetochore-particle condition described in (Akiyoshi et al., 2010).

Each flow chamber (~5-10 µL in volume) was constructed by placing two parallel strips of
double-stick tape between a glass slide and a KOH- or plasma-cleaned coverslip. Immediately before
every experiment, two flow chambers were each functionalized as follows. First, flow chambers were
incubated with 10 µL of 10 mg·mL^{-1} biotinylated bovine serum albumin (Vector Laboratories) for 15
minutes at room temperature. Chambers were then washed with 100 µL BRB80. Next, 20 µL of 333
mg·mL^{-1} avidin DN (Vector Laboratories) in BRB80 was added to the chamber and incubated for 3
minutes before washing again with 100 µL BRB80. Chambers were then incubated for 5 minutes with
GMPCPP-stabilized biotinylated microtubule seeds warmed for 5-10 minutes in BRB80. Next, chambers
were washed with 100 µL of growth buffer (1 mM GTP, 1 mg·mL^{-1} 🄰-casein, and 0.08 mg·mL^{-1}
biotinylated bovine serum albumin in BRB80). Finally, chambers were incubated with kinetochore-
decorated beads diluted approximately 8-fold into growth buffer augmented with 1.5 mg·mL^{-1} purified
bovine brain tubulin and an oxygen scavenging system (1 mM DTT, 500 µg·mL^{-1} glucose oxidase, 60
µg·mL^{-1} catalase, and 25 mM glucose). The flow chamber exit and entrance were then sealed with nail
polish and used for experiments at 23°C for up to 2 hours.

Performing the dual-trap assay

Laser trap experiments were performed using two custom-built instruments based on
commercial inverted microscopes (Nikon; TE2000U) modified to each incorporate a trapping laser
(Spectra-Physics; J20I-BL-106C-NSI), a computer-controlled piezo specimen stage (Physik Instrumente; P-
517.3CL), and a position-sensitive photodetector (Pacific Silicon Sensor; DL100-7-PCBA2). Experiments
were performed at 22° C with a trap stiffnesses of 0.35 – 0.91 pN·nm^{-1}. See (Franck et al., 2010) for
details regarding instrument setup and calibration.

The dual-trap assay was modified from previous force-clamp protocols (Akiyoshi et al., 2010;
Franck et al., 2010; Miller et al., 2016) and maintained constant total force on two bead-laser trap
systems, rather than one (Figure 2a). Both lasers were controlled by a single computer, which adjusted
the force on each bead to maintain force equilibrium on the modeled coupler given the measured bead
positions (Figure 2b). The bead positions served as accurate proxies for the microtubule tip positions,
because kinetochore-decorated beads are known to track with microtubule plus-ends under tensions
between 0.5 and 18 pN (Akiyoshi et al., 2010; Asbury et al., 2006; Franck et al., 2010; Miller et al., 2016;
Powers et al., 2009; Tien et al., 2010). In addition, the yeast kinetochores used to link the beads to the
microtubule tips are very small, only ~130 nm in length (Gonen et al., 2012), compared to the relevant
micrometer-scale tip displacements. The small size of the yeast kinetochore, as well as the slow change
in force on each tip (Figure S3-1), ensures that the kinetochore's (visco)elastic properties did not
significantly affect experimental results, because any force-dependent changes in its length had a
negligible effect compared to tip growth.

To calibrate the laser trap's zero-force position, we recorded the position of a free bead using
each laser-trap at the start of each 2-hour experiment, and as necessary between recordings. Feedback
to exert precisely the amount of force dictated by the relative bead positions and the spring-coupler

model was implemented with custom LabView software (National Instruments). Bead-trap separation
was sampled at 40 kHz, while stage position to maintain the desired bead-trap separation (and thus
force) was updated at 50 Hz. Bead and stage position data were decimated to 200 Hz before storing to
disk, a temporal frequency which is more than sufficient to capture pauses and runs of many seconds.

LabView software to perform force-clamp experiments is available at
https://github.com/casbury69/laser-trap-control-and-data-acquisition and the LabView software to
perform dual-trap experiments is available at https://github.com/casbury69/dual-trap-control-and-data-
acquisition.

Analyzing dual-trap recordings

We analyzed recordings from the dual-trap assay using custom software made in Igor Pro 9
(Wavemetrics). First, by hand, we marked the beginning and end of recorded segments in which both
microtubules were growing and separated these episodes into individual recordings. We then calculated
the tip separation between coupled microtubules for each excised recording by taking the absolute
value of the difference in position at 1 s intervals from 0 to 400 s. For each recording, total force on the
microtubule pair was computed by adding the force on both microtubules together at 0.005 s intervals
(Fig. S3-1).

To calculate the speed of tip separation, we fit the first 100 s to a line with the Levenberg-
Marquardt method. Uncertainties in tip separation speed were estimated from residuals as described in
(Press et al.).

SEMs for microtubule pair tip separation measured using the dual-trap assay (Figure 3b) were
calculated at 1 s intervals after the recording began using the following equation:

$$SEM_{tip\,sep}(t) = \frac{\sigma_t}{\sqrt{N_t}}$$

where σ_t is the standard deviation of the tip separations at t seconds and N_t is the number of
microtubule pairs which were still growing and tip-attached to a bead at t seconds.

Download files at https://doi.org/10.5061/dryad.xksn02vnb to view position- and force-versus-
time curves for each of the dual-trap recordings analyzed in this work. See also 'Dual-trap anlaysis.txt' at
https://doi.org/10.5281/zenodo.8433109 for the software used to analyze dual-trap recordings.

Creating Kaplan-Meier survival curves

Kaplan-Meier survival curves for both simulated and *in vitro* microtubule pairs (Figure 3c, Figure
4c, Figure 6c, Figure S6-2, and Figure S6-3c) were calculated at 1 s intervals after the recording began
using the following equation:

$$S(t) = \prod_{j=0}^{t} \frac{n_j - d_j}{n_j}$$

where $S(t)$ is the Kaplan-Meier survival probability at t seconds, t is the number of Kaplan-Meier estimates made between time 0 and t seconds, n_j is the number of microtubule pairs still growing, tip-attached to a bead, and within the threshold tip separation at j seconds, and d_j is the number of microtubule pairs that exceeded the threshold tip separation (0.4, 0.6, 0.8, or 1.2 μm) between time $j - 1$ and j seconds.

SEMs for Kaplan-Meier survival plots estimated using the dual-trap assay (Figure 3c and Figure S6-2) were calculated at 1 s intervals after the recording began using the following equation:

$$SEM_{survival}(t) = S(t) * \sqrt{\sum_{j=1}^{t} \frac{d_j}{n_j(n_j - d_j)}}$$

where $S(t)$ is the Kaplan-Meier survival probability at t seconds, t is the number of Kaplan-Meier estimates made between time 0 and t seconds, d_j is the number of microtubule pairs that exceeded the threshold tip separation (0.4, 0.6, 0.8, or 1.2 μm) between time $j - 1$ and j seconds, and n_j is the number of microtubule pairs still growing, tip-attached to a bead, and within the threshold tip separation at j seconds.

Segmenting constant force traces into periods of run and pause

To estimate pause entrance and exit rates as a function of force, we segmented N = 356 recordings of growing microtubules under constant force (from (Akiyoshi et al., 2010)) into pauses and fast growth ('runs,' Figure S5-1a and b). First, we estimated the instantaneous growth speed over the entirety of each recording to be the slope of the best fit line for a sliding 2 s window (Figure S5-1c). Next, we defined a unique threshold between pause and run for each recording by assembling a histogram of growth speeds throughout the recording and fitting that histogram with a sum of between two and six Gaussian peaks. We centered one Gaussian at or near 0 μm·min^{-1} and defined it as the 'pause' peak. Fitting the growth speed histograms to Gaussian peaks was performed using the 'Multipeak fitting' package in Igor Pro 9. The intersection of the pause peak and the Gaussian with the next highest mean growth speed was identified for each individual recording and defined as the threshold to distinguish pauses and runs for that recording.

To reduce the impact of high-frequency noise, we excluded pauses and runs that lasted less than 1 s. If the instantaneous growth speed fell below that recording's predefined threshold for at least 1 s, the microtubule was considered to have paused. If the instantaneous growth speed fell below the threshold for less than 1 s, the short interval was combined with the run intervals before and after to make a single run. Likewise, if the instantaneous growth speed exceeded the threshold for at least 1 s, the microtubule was considered to be in run. If the instantaneous growth speed exceeded the threshold

for less than 1 s, the short interval was combined with the pause intervals before and after to make a
single pause. If the defined threshold was never crossed, the recording was considered a single run
interval.

The overall trends of tension suppressing pause entrance and promoting pause exit were
maintained irrespective of our choices of minimum event duration and smoothing window size. While
the 1 s minimum event duration decreased the overall frequency of pause entrances and exits, it did not
significantly affect the measured force sensitivities of pause entrance (F_{EN}) and exit (F_{EX}). Re-analyzing
the constant force recordings with only a 0.01 s minimum event duration, we measured force
sensitivities of F_{EN} = -5.1 pN and F_{EX} = 14.3 pN, compared to F_{EN} = -4.6 pN and F_{EX} = 14.1 pN with
the 1 s minimum event duration. The size of the smoothing window had a significant effect on measured
pause exit force sensitivity, but not pause entrance force sensitivity—re-analyzing the constant force
recordings with a 5 s smoothing window, we measured force sensitivities of F_{EN} = -4.4 pN and
F_{EX} = 24.8 pN. Ultimately, we chose the 2 s smoothing window and 1 s minimum duration threshold
because these choices resulted in recording segmentation that agreed best with how we would
differentiate pauses and runs by eye.

Download files at https://doi.org/10.5061/dryad.6djh9w16v to view position- and velocity-
versus time curves for each force-clamp recording analyzed in this work, as well as how pauses were
marked for each recording. This file also includes additional shortening events which were not studied
here. See also 'Force-clamp viewer.txt' at https://doi.org/10.5281/zenodo.8433062 for the software
used to distinguish pauses from runs.

Quantifying growth speed during runs for individual recordings

For each pause-based model that we considered, microtubule growth speed during runs was
assumed to be time-invariant under constant force. Thus, we made the same assumption when we
quantified the growth speed during runs of individual microtubules *in vitro*, and calculated a unique
growth speed during runs for each recording (dots in Figure 5b). The growth speed during each run
interval was calculated by averaging together the instantaneous growth speeds comprising that interval,
measured with the 2 s sliding window described above. We calculated the overall growth speed during
runs for a single microtubule by averaging together the growth speed of all contained run intervals,
weighted by the duration of that interval, such that:

$$v_{run} = \frac{\sum v_i * t_i}{T}$$

where v_{run} is the overall growth speed during runs for a particular recording, v_i is the growth speed for
the i^{th} run interval, t_i is the duration of the i^{th} run interval, and T is the total time spent in the run state.

Quantifying the force dependence of pause entrance rates, exit rates and growth speed

To calculate average pause exit rates, entrance rates, and growth speeds as a function of force
we binned each recording by the constant tensile force to which the microtubule was subjected (Figure
5). For consistency, we used the same seven force bins used by (Akiyoshi et al., 2010) (see Table 2 for
bin boundaries). For both the pause entrance and exit rate, we excluded the highest force bin from our
analysis because of low statistics. For each force bin, we calculated the mean pause entrance and exit
rate by counting the number of times microtubules in that force bin entered and exited pause and
dividing by the total time microtubules in that bin spent in run and pause, respectively (Figure S5-1a).
Error bars in the y-direction represent counting uncertainties, calculated with the equation:

$$SEM_{rate}(i) = \frac{k_i}{\sqrt{N_i}}$$

where k_i is the rate of pause entrance or exit for the i^{th} force bin, and N_i is the number of pause
entrances or exits in the recordings in the i^{th} bin (N = 387 to 3502 pause entrances and N = 378 to 3501
pause exits per bin, see Table 2 for the number of pause entrances and exits by bin). Error bars in the x-
direction represent standard deviations of forces for the recordings in each bin.

We found the mean growth speed during runs for each force bin by averaging together the
growth speed during runs for all the recordings in that bin. Error bars in the y-direction represent the
SEMs of growth speeds, calculated with the equation:

$$SEM_{run\ speed}(i) = \frac{\sigma_i}{\sqrt{N_i}}$$

600　where σ_i is the standard deviation of the growth speeds during runs, averaged over each recording, for
601　the i^{th} force bin, and N_i is the number of recordings in the i^{th} bin (N = 9 to 85 recordings per bin, see
602　Table 2 for the number of recordings by bin). Once again, error bars in the x-direction represent the
603　standard deviations of forces for the recordings in each bin. Fit lines for pause exit and entrance rates
604　and growth speed were calculated with Levenberg-Marquardt least-squares method (Press et al.).

606　Simulating pairs of growing microtubules

Simulated recordings using each model followed the same general procedure. For simulations in
which the growth or run speed was different between microtubules (both non-ergodic models), the
'unloaded' growth or run speed was drawn from a distribution of measured growth or run speeds as
described in the results section. For the non-ergodic pausing model, intrinsic unloaded run speeds
($v_{intrinsic}$) of more than 30 nm/s were considered outliers and excluded from the distribution. For
microtubules simulated with the ergodic pausing model, both microtubules had the same growth speed
during runs, based on the equation and parameters for v_{run} in Table 1. Each simulation began with both
microtubules growing (in run, for pause-based models) from the same position under the same tensile
force. As microtubules stochastically grew or paused, the tensile force on each microtubule evolved
according to the spring-coupler model, just like they did in the dual-trap assay. As the force changed,
depending on the model type, the growth or run speed changed according to exponential relationships
with force estimated from (Akiyoshi et al., 2010) (Table 1).

A modified Gillespie algorithm was used to implement stochastic microtubule events: pause entrance and exit, detachment from the kinetochore, switches into shortening (catastrophe), and other event interruptions. Each of these event rates were assumed independent Poisson processes and were derived directly from the data summarized in (Akiyoshi et al., 2010). Each microtubule was assumed to have identical rates for all stochastic events, given the same external force. Detachment and catastrophe rates were calculated using the exponential relationships as a function of force estimated by (Akiyoshi et al., 2010).The rate of interruption by other events was also estimated from these force-clamp recordings but was force-independent. For simulations that included pausing, rates of pause entrance and exit were calculated from the fit equations in Table 1 (fit lines in Figure 5).

Because detachment, catastrophe, and pause rates are force-dependent, they can only be approximated as Poisson (time-invariant) rates for short time intervals over which the force on the tip is relatively constant. Consequently, we imposed a maximum timestep of no greater than 1 s for each simulation. If the timestep chosen by our Gillespie algorithm exceeded the maximum timestep, no event occurred and the microtubule continued growing during that 1 s interval. All simulations were performed using custom software in Igor Pro 9 (Wavemetrics).

Download 'Dual-trap simulations.pxp' at https://github.com/casbury69/simulations-of-coupled-microtubules to run dual-trap simulations as described in the results and see 'Dual-trap simulator.txt' at https://github.com/casbury69/simulations-of-coupled-microtubules for the code used to run these simulations.

Comparing pausing and non-pausing models with simple probability

To evaluate whether the pausing or non-pausing model was more likely given our dual-trap measurements, we compared histograms of tip separations at 100 and 200 s after the start of the recording or simulation (Figure S6-4). We used 0.1 μm bins for both simulated and dual-trap data and normalized all histograms to have an area of 1.

We calculated the likelihood that simulations performed using the pausing and non-pausing models would produce each *in vitro* tip separation:

$$L_{measurement} = w_{bin} * P_{normalized}$$

where w_{bin} is the bin width of the tip separation histogram (0.1 μm) and $P_{normalized}$ is the normalized probability of the given model at the measured tip separation (y-values of the dashed lines in Figure S6-4).

To compare the overall likelihood that the pausing and non-pausing simulations would produce all the tip separations measured *in vitro*, we multiplied together the likelihoods of each measurement for a given model at a given coupler stiffness and at a given time point. We calculated the log likelihood by taking the base 10 logarithm of each probability (reported in the right-hand column of each table in Figure S6-4).

Statistical analysis

656   All statistical analysis was performed using custom software written in Igor Pro 9 (Wavemetrics).

657   We calculated p-values to determine whether survival curves of stiff-coupled and soft-coupled
658 *in vitro* microtubule pairs were statistically different for each threshold tip separation using a log-rank
659 test. For each survival curve, at 1 s intervals after the recording began, we calculated an expected value
660 using the following equation:

$$E(t) = \frac{n_t}{n_{tot}}(d_{tot})$$

where n_t is the number of microtubule pairs comprising one survival curve that are still growing, tip-
attached to a bead, and within the threshold tip separation at t seconds, n_{tot} is the total number of
microtubule pairs comprising both survival curves being compared that are still growing, tip-attached to
a bead, and within the threshold tip separation at t seconds, and d_{tot} is the total number of microtubule
pairs that exceeded the threshold tip separation between time $t - 1$ and t seconds for both survival
curves being compared. Expected values for each pair of survival curves were used to calculate a chi-
square value for each tip separation threshold using the following equation:

$$\chi^2 = \sum_{j=1}^{2} \frac{\left(\sum_{t=1}^{T} S_j(t) - \sum_{t=1}^{T} E_j(t)\right)^2}{\sum_{t=1}^{T} E_j(t)}$$

where T is the total number of timepoints (400, in all cases), $S_j(t)$ is the Kaplan-Meier survival
probability of curve j at time t, and $E_j(t)$ is the expected value of Kaplan-Meier survival curve j at time
t. A chi-square table was then used to estimate the p-value for each tip separation threshold.

To determine whether tip separation and tip separation speed between soft and stiff couplers
were statistically significant, we calculated p-values with a Wilcoxon-Mann-Whitney two sample rank
test using built-in Igor Pro software (Cheung & Klotz, 1997; Klotz, 2006; Streitberg & Rohmel, 1986; Zar,
1999).
675

Figures

Figure 1. Intrinsic variability in microtubule growth could be coordinated by mechanical coupling. (**a**) Microtubules (MTs) grow at intrinsically variable rates that increase with force, on average. Example recordings show plus-end position versus time for individual growing microtubules subject to a constant tensile force of either 2 or 6 pN (thin green or magenta traces, respectively). At each force, 10 to 14 example recordings are shown, together with the mean position versus time (thick traces, computed from N = 25 or N = 10 individual recordings at 2 or 6 pN, respectively). Individual traces are from (Akiyoshi et al., 2010). (**b**) Mechanism by which mechanical coupling could coordinate growing microtubules. In this example, two microtubules share a total force, F_{TOT}, through two spring-like connections to their plus-ends. If one microtubule stochastically lags behind, it will experience more tension than the leading microtubule due to differential stretching of their spring-connections. Because tension tends to accelerate plus-end growth (as illustrated in panel a), the higher tension will tend to accelerate the growth of the lagging microtubule, causing it to catch up.

Figure 2. Dual-trap assay to measure the coordination of two real microtubules *in vitro* coupled together by simulated spring-like connections to a shared tensile load. (a) Schematic of the dual-trap assay, which measures the effects of the spring-coupler model (detailed in b) on real microtubules (MTs) growing *in vitro*. Two dynamic microtubules are grown from seeds anchored to two separate coverslips on two separate stages, each with their own microscope and laser trap. Each laser trap applies tension to its respective microtubule plus-end via a bead decorated with isolated yeast kinetochores, and each tension is adjusted dynamically under feedback control by a shared computer. Initially (steps 1 and 2), microtubule plus-ends are considered to have the same position ($x_1 = x_2 = 0$) so that each bears half of the total user-input tension (F_{TOT}). Over time, as microtubules grow at different rates (step 3), the computer dynamically adjusts the tension on each microtubule according to the spring-coupler model (steps 4 and 5), so that the higher the tip separation ($|x_1 - x_2|$) and stiffness (κ), the larger the difference in tension between the two microtubules. (b) Spring-coupler model used to represent the physical connection between two microtubule plus-ends via a kinetochore. Both springs have identical stiffnesses (κ). The kinetochore is assumed to be in force equilibrium such that the sum of the spring forces is equal to the total external force, F_{TOT}, which is kept constant in the current study.

704
Figure 3. Microtubule pairs growing *in vitro* are coordinated by mechanical coupling. (**a**) Example dual-trap assay recordings
showing real microtubule (MT) plus-end positions over time for representative microtubule pairs coupled via soft (1 pN·µm⁻¹,
left) or stiff (5 pN·µm⁻¹, right) simulated spring-couplers. All dual-trap data was recorded with F_{TOT} = 8 pN. (**b**) Tip separation
($|x_1 - x_2|$) between microtubule plus-end pairs that were coupled with either soft (left) or stiff (right) simulated spring-couplers
over 400 s of growth. Light colors show tip separations for 10 individual microtubule pairs with each coupler stiffness, while
dark colors show mean tip separation for all recorded pairs (N = 50 and N = 43 recordings with soft and stiff couplers,
respectively). (**c**) Fraction of microtubule pairs whose plus-ends remained within 0.8 µm of each other over time. (In other
words, the fraction of pairs at any given time whose tip separation had not exceeded the dashed line in (b).) Shaded regions in
(b) and (c) show SEMs from N = 50 and N = 43 recordings with soft and stiff couplers, respectively.

714
Figure S3-1. Additional dual-trap assay recordings showing tip positions and forces on pairs of growing microtubules. (a)
Example dual-trap assay recordings showing real microtubule (MT) plus-end positions (top) and forces (bottom) plotted against
time for representative microtubule pairs coupled via soft or stiff simulated spring-couplers. Plus-end position data in the top
graphs is recopied from Figure 3a. Black traces in the bottom graphs show total force on the two microtubules (F_{TOT}) and
demonstrate that a constant total force of F_{TOT} = 8 pN was maintained by the feedback-control. (**b**) Two additional examples of
dual-trap assay recordings, formatted as in (a).

Figure 4. Simple non-pausing simulations fail to recapitulate the coordinated growth of mechanically coupled microtubule pairs. (a) Example simulations of microtubule (MT) plus-end positions over time using the non-pausing model. Simulated microtubule pairs were coupled via soft or stiff spring-couplers and shared a total load F_{TOT} = 8 pN to match dual-trap assay conditions. (b) Tip separations between microtubule plus-end pairs that were simulated using the non-pausing model and coupled with either soft (left) or stiff (right) spring-couplers. Light colors show tip separations for N = 40 individual simulated microtubule pairs, while dark colors show mean tip separation for N = 10,000 simulated pairs. (c) Comparison between the fraction of in silico (simulated, dashed curves) and in vitro (real, solid curves) microtubule pairs whose plus-ends remained within 0.8 μm of each other over time. Shaded regions show SEMs from N = 50 and N = 43 in vitro recordings with soft and stiff couplers, respectively. Soft- and stiff-coupled microtubule pairs were each simulated N = 10,000 times over 400 s of growth in both (b) and (c). Simulation parameters for (a), (b), and (c) can be found in Table 1.

Figure 5. Increasing tension on individual growing microtubules suppresses pausing and accelerates growth during runs. (a)
As tension on a growing microtubule tip increases, the rate of pause entrance decreases, while the rate of pause exit increases.
Mean pause entrance (■) and exit (■) rates (k_{EN} and k_{EX}) were estimated as described in Figure S5-1 and are plotted as
functions of force. Error bars show counting uncertainties from N = 25 to 85 individual microtubule recordings and dashed
curves show least squares exponential fits. (**b**) As tension on a microtubule tip increases, the growth during runs (i.e., in-
between pauses) is accelerated. Points show run speeds measured from recordings of individual microtubules subject to
constant tensile forces (from (Akiyoshi et al., 2010)). Symbols (■) represent mean run speeds (v_{RUN}) grouped according to the
applied tensile force. Error bars show SEMs from N = 9 to 85 individual microtubule recordings and dashed curve shows the
least squares exponential fit. See Figure S5-1 and Materials and Methods for details and Table 1 for fit parameters.

Figure S5-1. Individual microtubules frequently pause during periods of overall growth. (a) Growing microtubules switch between states of 'pause,' when plus-end growth speed is nearly zero, and 'run,' when the plus-end is clearly growing. Pause entrance and exit rates, k_{EN} and k_{EX}, were quantified by counting the numbers of entrances or exits and dividing by the total amount of time spent in the run or pause states, respectively. (b) Example recording showing plus-end position versus time for an individual microtubule subject to a constant tensile force of 4 pN and exhibiting frequent pauses (denoted by purple shaded regions). This example recording is one of $N = 356$ force-clamp traces (from (Akiyoshi et al., 2010)) used to estimate the pause entrance and exit rates and growth speeds during runs as functions of force (as shown in Figure 5). (c) Classification of runs and pauses. Instantaneous growth speed was computed as a function of time (and plotted at right) from the recording shown in (b), using a 2 s sliding window, and then assembled into a histogram (middle plot, red) and fit with a sum of multiple Gaussian curves (middle and left plots, blue). One Gaussian was always centered at 0 µm·min⁻¹ and defined as the 'pause' peak. The intersection of the pause peak and the Gaussian having the next highest mean growth speed was identified for each individual recording and used as a threshold (dashed horizontal line) to distinguish pauses and runs for that recording. Pauses were defined as periods longer than 1 s when the instantaneous growth speed fell below the threshold, while runs were defined as periods longer than 1 s when growth speed was above the threshold (see Materials and Methods for details).

Figure S5-2. Pause and run durations are exponentially distributed. (a) Normalized histograms showing pause durations
measured from recordings of individual microtubules growing under constant levels of tension (force-clamp data from (Akiyoshi
et al., 2010)), either between 0 and 2 pN (top), or between 6 and 8 pN (bottom), corresponding to the lowest and third-highest
force bins in Table 2, respectively. **(b)** Normalized histograms showing run durations measured from recordings of individual
microtubules growing under constant levels of tension (force-clamp data from (Akiyoshi et al., 2010)), as in (a). For both (a) and
(b), least-squares regressions are shown as dashed curves. The indicated time constants (τ) were derived from these fits.

Figure 6. Simulations recapitulate the coordinated growth of mechanically coupled microtubule pairs. (a) Example simulations of microtubule (MT) plus-end positions over time using the pausing model. Simulated microtubule pairs were coupled via soft or stiff spring-couplers and shared a total load F_{TOT} = 8 pN to match dual-trap assay conditions. (b) Tip separations between microtubule plus-end pairs that were simulated using the pausing model and coupled with either soft (left) or stiff (right) spring-couplers. Light colors show tip separations for N = 40 individual simulated microtubule pairs, while dark curves show mean tip separation for N = 10,000 simulated pairs. (c) Comparison between the fraction of *in silico* (simulated, dashed curves) and *in vitro* (real, solid curves) microtubule pairs whose plus-ends remained within 0.8 μm of each other over time. Shaded regions show SEMs from N = 50 and N = 43 *in vitro* recordings with soft and stiff couplers, respectively. Soft- and stiff-coupled microtubule pairs were each simulated N = 10,000 times over 400 s of growth in both (b) and (c). Simulation parameters for (a), (b), and (c) are listed in Table 1.

Figure S6-1. Simulations of microtubule growth under constant force using the pausing model resemble real microtubule recordings. Examples using the pausing model to simulate plus-end position versus time for individual growing microtubules subject to a constant tensile force of either 2 or 6 pN (light green or magenta traces, respectively). At each force, N = 16 example simulations are shown, together with the corresponding mean positions versus time (thick traces, computed from the N = 16 individual simulations).

Figure S6-2. Simulations of microtubule pairs using the pausing model maintain tip separations similar to real microtubule
pairs. Comparison between the fraction of *in silico* (simulated, dashed curves) and *in vitro* (real, solid curves) microtubule (MT)
pairs whose plus-ends remained within (**a**) 0.4 μm, (**b**) 0.6 μm, (**c**) 0.8 μm, and (**d**) 1.2 μm of each other over time. Microtubule
pairs *in silico* and *in vitro* shared a total load of F_{TOT} = 8 pN and were coupled via soft or stiff spring couplers. Shaded regions
show SEM from N = 50 and N = 43 *in vitro* recordings with soft and stiff couplers, respectively. Soft- and stiff-coupled
microtubule pairs were each simulated N = 10,000 times over 400 s of growth. Simulation parameters can be found in Table 1.
P-values indicate comparisons between soft- and stiff-coupled microtubule pairs *in vitro* using a log-rank test.

Figure S6-3. Simulations using an ergodic version of the pausing model, without intrinsic heterogeneity, fail to recapitulate
the coordinated growth of mechanically coupled microtubules pairs. (a) Example simulations of microtubule (MT) plus-end
positions over time using the ergodic pausing model. Simulated microtubule pairs were coupled via soft or stiff spring-couplers
and shared a total load F_{TOT} = 8 pN to match dual-trap assay conditions. (b) Tip separations between microtubule plus-end
pairs that were simulated using the ergodic pausing model and coupled with either soft (left) or stiff (right) spring-couplers.
Light colors show tip separations for N = 40 individual simulated microtubule pairs while dark colors show mean tip separation
for N = 10,000 simulated pairs. (c) Comparison between the fraction of *in silico* (simulated, dashed curves) and *in vitro* (real,
solid curves) microtubule pairs whose plus-ends remained within 0.8 μm of each other over time. Shaded regions show SEMs
from N = 50 and N = 43 *in vitro* recordings with soft and stiff couplers, respectively. Soft- and stiff-coupled microtubule pairs

were each simulated N = 10,000 times over 400 s of growth in both (b) and (c). Simulation parameters for (a), (b), and (c) can be
found in Table 1. As expected, microtubule pairs simulated using this ergodic model remained much closer together than real
microtubule pairs, rarely separating by more than 0.5 μm irrespective of coupler stiffness. After 300 s, the average tip
separation for soft-coupled pairs was only 0.202 ± 0.002 μm (mean ± SEM from N = 10,000 simulations), barely different from
the average tip separation for stiff-coupled pairs of 0.134 ± 0.002 μm (mean ± SEM from N = 10,000 simulations). Combined
with the success of the non-ergodic pausing model in fitting our dual-trap data, this supports the idea that microtubule growth
can be considered non-ergodic during our 400 s experiments.
810

	Mean [μm]	SD [μm]	SEM [μm]	N	Log Likelihood
In vitro	0.39	0.29	0.06	23	N/A
Pausing	0.32	0.27	0.0009	100,397	-15.6
Non-pausing	0.56	0.49	0.002	100,654	-16.3

	Mean [μm]	SD [μm]	SEM [μm]	N	Log Likelihood
In vitro	0.23	0.17	0.03	26	N/A
Pausing	0.25	0.21	0.0007	100,000	-10.2
Non-pausing	0.45	0.37	0.001	100,000	-13.4

	Mean [μm]	SD [μm]	SEM [μm]	N	Log Likelihood
In vitro	0.60	0.54	0.17	15	N/A
Pausing	0.56	0.48	0.0015	100,000	-18.0
Non-pausing	1.05	0.91	0.0028	104,792	-19.2

	Mean [μm]	SD [μm]	SEM [μm]	N	Log Likelihood
In vitro	0.31	0.23	0.06	16	N/A
Pausing	0.37	0.29	0.00092	100,000	-14.5
Non-pausing	0.71	0.55	0.0017	100,000	-17.3

Figure S6-4. Probability analysis shows pausing model fits dual-trap recordings better than the non-pausing model. (Top of
each panel) Histograms of tip separation between microtubule pairs measured in the dual-trap assay and simulated using the
pausing and non-pausing model. Soft-coupled pairs are shown on the left ((**a**) and (**c**)), while stiff-coupled pairs are shown on
the right ((**b**) and (**d**)). Histograms of tip separations measured 100 s after the start of the recording or simulation are shown on
the top ((**a**) and (**b**)), while tip separations measured 200 s after the start of the recording or simulation are shown on the
bottom ((**c**) and (**d**)). Tip separation histograms produced by the pausing (purple) and non-pausing model (green) are
represented with dashed lines, while dual-trap data is represented by bars. (**Bottom of each panel**) Tables comparing the
means, standard deviations (SD), standard errors of the mean (SEM) and numbers of recordings (N) for *in vitro* microtubule
pairs and microtubule pairs simulated using the pausing and non-pausing models. The log likelihood probability for each model
given the *in vitro* data is shown in the righthand column. The greater the log likelihood, the greater the probability that the
given model could reproduce the measured *in vitro* data. At both 100 s and 200s, for soft and stiff couplers, the pausing model
scores higher than the non-pausing model, indicating that the pausing model is a better fit to our *in vitro* data.

Parameter name	Parameter symbol	Relevant equation	Parameter estimate	Applicable models	Reference
Growth speed force sensitivity	F_g	$v_g = v_{g0}\exp(F/F_g)$	8.4 pN	non-pausing	Akiyoshi et al., 2010 Figure 4e
Unloaded pause entrance rate	k_{EN0}	$k_{EN}= k_{EN0}\exp(F/F_{EN})$	19.4 min^{-1}	pausing	Figure 5a, red
Pause entrance rate force sensitivity	F_{EN}	$k_{EN}= k_{EN0}\exp(F/F_{EN})$	-4.6 pN	pausing	Figure 5a, red
Unloaded pause exit rate	k_{EX0}	$k_{EX}= k_{EX0}\exp(F/F_{EX})$	9.0 min^{-1}	pausing	Figure 5a, blue
Pause exit rate force sensitivity	F_{EX}	$k_{EX}= k_{EX0}\exp(F/F_{EX})$	14.1 pN	pausing	Figure 5a, blue
Unloaded run speed	v_0	$v_{RUN}= v_0\exp(F/F_{RUN})$	0.52 μm·min^{-1}	neither	Figure 5b
Run speed force sensitivity	F_{RUN}	$v_{RUN}= v_0\exp(F/F_{RUN})$	14.8 pN	pausing	Figure 5b

826 **Table 1. Parameter table for coupled microtubule pairs simulated for non-pausing and pausing models.** Relevant parameters
827 measured from constant force recordings from (Akiyoshi et al., 2010). These parameters were used to simulate coupled
828 microtubule pairs using each of the models described in the main text. The equation in which each parameter is used, as well as
829 which models of microtubule growth use that equation, are indicated.

Bin number	N	Lower bound [pN]	Upper bound [pN]	Pause entrances	Time in pause [s]	Pause exits	Time in run [s]
1	41	0.69	1.94	1834	10957	1830	7364
2	85	1.94	2.97	3502	18047	3501	18952
3	77	2.97	3.999	2376	12842	2374	18159
4	60	3.999	5.98	1937	9661	1932	16993
5	60	5.98	7.99	1439	5930	1438	18465
6	25	7.99	10.3	387	1298	378	6559
7	9	10.3	18	23	93	23	712

832 **Table 2. Number of recordings, bounds, and pause quantification from force-clamp dataset, binned by force**. Table describing
833 constant force recordings in (Akiyoshi et al., 2010) and the pause quantification performed in the current work. N indicates the
834 number of recordings in a particular bin. For consistency, these force bins are identical to those used in (Akiyoshi et al., 2010).

References

838 Akiyoshi, B., Sarangapani, K. K., Powers, A. F., Nelson, C. R., Reichow, S. L., Arellano-Santoyo, H., Gonen,
839 T., Ranish, J. A., Asbury, C. L., & Biggins, S. (2010). Tension directly stabilizes reconstituted kinetochore-
840 microtubule attachments. *Nature*, *468*(7323), 576-579. https://doi.org/10.1038/nature09594
841
842 Al-Bassam, J., Kim, H., Brouhard, G., van Oijen, A., Harrison, S. C., & Chang, F. (2010). CLASP promotes
843 microtubule rescue by recruiting tubulin dimers to the microtubule. *Dev Cell*, *19*(2), 245-258.
844 https://doi.org/10.1016/j.devcel.2010.07.016
845
846 Amaro, A. C., Samora, C. P., Holtackers, R., Wang, E., Kingston, I. J., Alonso, M., Lampson, M., McAinsh,
847 A. D., & Meraldi, P. (2010). Molecular control of kinetochore-microtubule dynamics and chromosome
848 oscillations. *Nature Cell Bio*, *12*(4), 319-329. https://doi.org/10.1038/ncb2033
849
850 Amin, M. A., Agarwal, S., & Varma, D. (2019). Mapping the kinetochore MAP functions required for
851 stabilizing microtubule attachments to chromosomes during metaphase. *Cytoskeleton*, *76*(6), 398-412.
852 https://doi.org/10.1002%2Fcm.21559
853
854 Amoah-Darko, F. L., & White, D. (2022). Modelling microtubule dynamic instability: Microtubule growth,
855 shortening and pause. *J Theor Biol*, *553*, 111257-111257. https://doi.org/10.1016/j.jtbi.2022.111257
856
857 Asbury, C. L., Gestaut, D. R., Powers, A. F., Franck, A. D., & Davis, T. N. (2006). The Dam1 kinetochore
858 complex harnesses microtubule dynamics to produce force and movement. *Proc Natl Acad Sci U S A*,
859 *103*(26), 9873-9878. https://doi.org/10.1073/pnas.0602249103
860
861 Bakhoum, S. F., & Compton, D. A. (2012). Kinetochores and disease: keeping microtubule dynamics *in*
862 *check! Curr Opinion in Cell Biol*, *24*(1), 64-70. https://doi.org/10.1016/j.ceb.2011.11.012
863
864 Bakhoum, S. F., Genovese, G., & Compton, D. A. (2009). Deviant kinetochore microtubule dynamics
865 underlie chromosome instability. *Curr Biol*, *19*(22), 1937-1942.
866 https://doi.org/10.1016/j.cub.2009.09.055
867
868 Banigan, E., Chiou, K., Ballister, E. R., Mayo, A. M., Lampson, M. A., & Liu, A. J. (2015). Minimal model for
869 collective kinetochore-microtubule dynamics. *PNAS*, *112*(41), 12699-12704.
870 https://doi.org/10.1073/pnas.1513512112
871
872 Brouhard, G., & Rice, L. M. (2018). Microtubule Dynamics: an interplay of biochemistry and mechanics.
873 *Nat Rev Mol Cell Biol*, *19*(7), 451-463. https://doi.org/10.1038/s41580-018-0009-y
874
875 Brouhard, G. J. (2015). Dynamic instability 30 years later: complexities in microtubule growth and
876 catastrophe. *MBoC*, *26*(7), 1207-1210. https://doi.org/10.1091/mbc.E13-10-0594
877
878 Brouhard, G. J., Stear, J. H., Noetzel, T. L., Al-Bassam, J., Kinoshita, K., Harrison, S. C., Howard, J., &
879 Hyman, A. A. (2008). XMAP215 is a processive microtubule polymerase. *Cell*, *132*(1), 79-88.
880 https://doi.org/10.1016/j.cell.2007.11.043
881

882 Cassimeris, L., Pryer, N. K., & Salmon, E. D. (1988). Real-time observations of microtubule dynamic
883 instability in living cells. *J Cell Biol*, *107*(6), 2223-2231. https://doi.org/10.1083/jcb.107.6.2223
884
885 Castrogiovanni, C., Inchingolo, A. V., Harrison, J. U., Dudka, D., Sen, O., Burroughs, N. J., McAinsh, A. D.,
886 & Meraldi, P. (2021). Evidence for a HURP/EB free mixed nucleotide zone in kinetochore-microtubules.
887 *Nature communications*, *13*(1). https://doi.org/10.1038/s41467-022-32421-x
888
889 Chacon, J. M., Mukherjee, S., Schuster, B. M., Clarke, D. J., & Gardner, M. K. (2014). Pericentromere
890 tension is self-regulated by spindle structure in metaphase. *J Cell Biol*, *205*(3), 313-324.
891 https://doi.org/10.1083/jcb.201312024
892
893 Cheung, Y. K., & Klotz, J. H. (1997). The Mann Whitney Wilcoxon distribution using linked lists. *Statistica*
894 *Sinica*, *7*, 805-813. https://www3.stat.sinica.edu.tw/statistica/oldpdf/a7n316.pdf
895
896 Chretien, D., Metoz, F., Verde, F., Karsenti, E., & Wade, R. H. (1992). Lattice defects in microtubules:
897 protofilament numbers vary within individual microtubules. *J Cell Biol*, *117*(5), 1031-1040.
898 https://doi.org/10.1083/jcb.117.5.1031
899
900 Cimini, D., Cameron, L. A., & Salmon, E. D. (2004). Anaphase spindle mechanics prevent mis-segregation
901 of merotelically oriented chromosomes. *Curr Biol*, *14*(23), 2149-2155.
902 https://doi.org/10.1016/j.cub.2004.11.029
903
904 Cleary, J. M., & Hancock, W. O. (2021). Molecular mechanisms underlying microtubule growth dynamics.
905 *Curr Biol*, *31*(10), R560-R573. https://doi.org/10.1016/j.cub.2021.02.035
906
907 Cleary, J. M., Kim, T., Cook, A. S. I., McCormick, L. A., Hancock, W. O., & Rice, L. M. (2022).
908 Measurements and simulations of microtubule growth imply stron longitudinal interactions and reveal a
909 role for GDP on the elongating end. *Elife*, *11*. https://doi.org/10.7554/eLife.75931
910
911 Cojoc, G., Roscioli, E., Zhang, L., Garcia-Ulloa, A., Shah, J. V., Berns, M. W., Pavin, N., Cimini, D., Tolic, I.
912 M., & Gregan, J. (2016). Laser microsurgery reveals conserved viscoelastic behavior of the kinetochore. *J*
913 *Cell Biol*, *212*(7), 767-776. https://doi.org/10.1083/jcb.201506011
914
915 Coombes, C. E., Yamamoto, A., Kenzie, M. R., Odde, D. J., & Gardner, M. K. (2013). Evolving tip structures
916 can explain age-dependent microtubule catastrophe. *Curr Biol*, *23*(14), 1342-1348.
917 https://doi.org/10.1016/j.cub.2013.05.059
918
919 Das, D., Das, D., & Padinhateeri, R. (2014). Force-induced dynamical properties of multiple cytoskeletal
920 filaments are distinct from that of single filaments. *PLoS one*, *9*(12).
921 https://doi.org/10.1371/journal.pone.0114014
922
923 Dogterom, M., & Yurke, B. (1997). Measurement of the force-velocity relation for growing microtubules.
924 *Science*, *278*(5339), 856-860. http://www.ncbi.nlm.nih.gov/pubmed/9346483
925
926 Driver, J. W., Geyer, E. A., Bailey, M. E., Rice, L. M., & Asbury, C. L. (2017). Direct measurement of
927 conformational strain energy in protofilaments curling outward from disassembling microtubule tips.
928 *Elife*, *6*. https://doi.org/10.7554/eLife.28433
929

Drpic, D., Almeida, A. C., Aguiar, P., Renda, F., Damas, J., Lewin, H. A., Larkin, D., Khodjakov, A., &
Maiato, H. (2018). Chromosome segregation is biased by kinetochore size. Curr Biol, 28(9), 1344-1356.
https://doi.org/10.1016/j.cub.2018.03.023

Ertych, N., Stolz, A., Stenzinger, A., Weichert, W., KaulfuB, S., Burfeind, P., Aigner, A., Wordeman, L., &
Bastians, H. (2014). Increased microtubule assembly rates influence chromosomal instability in
colorectal cancer cells. *Nature Cell Bio*, *16*(8), 779-791. https://doi.org/10.1038/ncb2994

Franck, A. D., Powers, A. F., Gestaut, D. R., Davis, T. N., & Asbury, C. L. (2010). Direct physical study of
kinetochore-microtubule interactions by reconstitution and interrogation with an optical force clamp.
Methods, *51*(2), 242-250. https://doi.org/10.1016/j.ymeth.2010.01.020

Gardner, M. K., & Odde, D. J. (2010). Stochastic simulation and graphic visualization of mitotic processes.
Methods, *51*(2), 251-256. https://doi.org/10.1016/j.ymeth.2010.01.021

Gardner, M. K., Pearson, C. G., Sprague, B. L., Zarzar, T. R., Bloom, K., Salmon, E. D., & Odde, D. J. (2005).
Tension-dependent regulation of microtubule dynamics at kinetochores can explain metaphase
congression in yeast. *Mol Biol Cell*, *16*(8), 3764-3775. https://doi.org/10.1091/mbc.E05-04-0275

Garner, R. M., & Theriot, J. A. (2022). Leading edge maintenance in migrating cells is an emergent
property of branched actin network growth. *Elife*, *11*. https://doi.org/10.7554/eLife.74389

Gaudenz, D., Matov, A., Wittman, T., Kumar, P., Thoma, C., Applegate, K., & Krek, W. (2010). Analysis of
microtubule dynamic instability using a plus-end growth marker. *Nature methods*, *7*(9), 761-768.
https://doi.org/10.1038/NMETH.1493

Geisterfer, Z. M., Zhu, D. Y., Mitchison, T. J., Oakey, J., & Gatlin, J. C. (2020). Microtubule growth rates
are sensitive to global and local changes in microtubule plus-end density. *Curr Biol*, *30*(15), 3016-3023.
https://doi.org/10.1016/j.cub.2020.05.056

Ghanti, D., & Chowdhury, D. (2015). Collective cargo hauling by a bundle of parallel microtubules: bi-
directional motion caused by load-dependent polymerization and depolymerization. *J Stat Mech*,
2015(1), 1008-1013. https://doi.org/10.1088/1742-5468/2015/01/P01008

Gildersleeve, R. F., Cross, A. R., Cullen, K. E., Fagen, A. P., & Williams, R. C. J. (1992). Microtubules grow
and shorten at intrinsically variable rates. *J Biol Chem*, *267*(12), 7995-8006.
https://www.jbc.org/article/S0021-9258(18)42399-X/pdf

Gonen, S., Akiyoshi, B., Iadanza, M. G., Shi, D., Duggan, N., Biggins, S., & Gonen, T. (2012). The structure
of purified kinetochores reveals multiple microtubule-attachment sites. Nat Struct Mol Biol, 19(9), 925-
929. https://doi.org/10.1038/nsmb.2358

Goshima, G., Wollman, R., Stuurman, N., Scholey, J. M., & Vale, R. D. (2005). Length control of the
metaphase spindle. *Curr Biol*, *15*(22), 1979-1988. https://doi.org/10.1016/j.cub.2005.09.054
Holzbaur, E. L. F., & Goldman, Y. E. (2010). Coordination of molecular motors: from *in vitro* assays to
intracellular dynamics. *Curr Opinion in Cell Biol*, *22*(1), 4-13. https://doi.org/10.1016/j.ceb.2009.12.014

Howard, J. (2001). *Mechanics of motor proteins and the cytoskeleton*. Sinauer Associates, Publishers.

Janson, M. E., de Dood, M. E., & Dogterom, M. (2003). Dynamic instability of microtubules is regulated
by force. *J Cell Biol*, *161*(6), 1029-1034. https://doi.org/10.1083/jcb.200301147

Khodjakov, A., Cole, R. W., Bajer, A. S., & Rieder, C. L. (1996). The force for poleward chromosome
motion in Haemanthus cells acts along the length of the chromosome during metaphase but only at the
kinetochore during anaphase. *J Cell Biol*, *132*(6), 1093-1104.
http://www.ncbi.nlm.nih.gov/pubmed/8601587

Khodjakov, A., & Rieder, C. L. (1996). Kinetochores moving away from their associated pole do not exert
a significant pushing force on the chromosome. *J Cell Biol*, *135*(2), 315-327.
http://www.ncbi.nlm.nih.gov/pubmed/8896591

Kiewisz, R., Fabig, G., Conway, W., Baum, D., Needleman, D., & Muller-Reichert, T. (2022). Three-
dimensional structure of kinetochore-fibers in human mitotic spindles. *Elife*, *11*.
https://doi.org/10.7554/eLife.75459

Kis, A., Kasas, S., Babic, B., Kulik, A. J., Benoit, W., Briggs, G. A., Schonenberger, C., Catsicas, S., & Forro, L.
(2002). Nanomechanics of microtubules. *Phys Rev Lett*, *89*(24), 248101.
https://doi.org/10.1103/PhysRevLett.89.248101

Klotz, J. H. (2006). *A Computational Approach to Statistics*.
https://www.mimuw.edu.pl/~pokar/Statystykal/Literatura/KlotzBook.pdf

1001

1002　Laan, L., Husson, J., Munteanu, E. L., Kerssemakers, J. W. J., & Dogterom, M. (2008). Force-generation
1003　and dynamic instability of microtubule bundles. *105*(26), 8920-8925.
1004　https://doi.org/10.1073/pnas.0710311105

1005

1006　Maddox, P., Straight, A., Coughlin, P., Mitchison, T. J., & Salmon, E. D. (2003). Direct observation of
1007　microtubule dynamics at kinetochores in Xenopus extract spindles: implications for spindle mechanics. *J
1008　Cell Biol*, *162*(3), 377-382. https://doi.org/10.1083/jcb.200301088

1009

1010　Mahserejian, S. M., Scripture, J. P., Mauro, A. J., Lawrence, E. J., Jonasson, E. M., Murray, K. S., Li, J.,
1011　Gardner, M., Alber, M., Zanic, M., & Goodson, H. V. (2022). Quantification of microtubule stutters:
1012　dynamic instability behaviors that are strongly associated with catastrophe. *MBoC*, *33*(3).
1013　https://doi.org/10.1091/mbc.E20-06-0348

1014

1015　Maiato, H., Fairley, E. A. L., Rieder, C. L., Swedlow, J. R., Sunkel, C. E., & Earnshaw, W. C. (2003). Human
1016　CLASP1 is an outer kinetochore component that regulates spindle microtubule dynamics. *Cell*, *113*(7),
1017　891-904. https://doi.org/10.1016/S0092-8674(03)00465-3

1018

1019　Maiato, H., Rieder, C. L., Earnshaw, W. C., & Sunkel, C. E. (2003). How do kinetochores CLASP dynamic
1020　microtubules? *Cell Cycle*, *2*(6), 511-514. https://doi.org/10.4161/cc.2.6.576

1021

1022　McAinsh, A. D., & Meraldi, P. (2011). The CCAN complex: Linking centromere specification to control of
1023　kinetochore-microtubule dynamics. *Seminars in Cell & Dev Biol*, *22*(9), 946-952.
1024　https://doi.org/10.1016/j.semcdb.2011.09.016

1025

1026 McDonald, K. L., O'Toole, E. T., Mastronarde, D. N., & McIntosh, J. R. (1992). Kinetochore microtubules in
1027 PTK cells. *J Cell Biol*, *118*(2), 369-383. http://www.ncbi.nlm.nih.gov/pubmed/1629239
1028
1029 McIntosh, J. R., O'Toole, E., Morgan, G., Austin, J., Ulyanov, E., Ataullakhanov, F., & Gudimchuk, N.
1030 (2018). Microtubules grow by the addition of bent guanosine triphosphate tubulin to the tips of curved
1031 protofilaments [Article]. *Journal of Cell Biology*, *217*(8), 2691-2708.
1032 https://doi.org/10.1083/jcb.201802138
1033
1034 Miller, M. P., Asbury, C. L., & Biggins, S. (2016). A TOG Protein Confers Tension Sensitivity to
1035 Kinetochore-Microtubule Attachments. *Cell*, *165*(6), 1428-1439.
1036 https://doi.org/10.1016/j.cell.2016.04.030
1037
1038 Murray, L. E., Kim, H., Rice, L. M., & Asbury, C. L. (2022). Working strokes produced by curling
1039 protofilaments at disassembling microtubule tips can be biochemically tuned and vary with species.
1040 *Elife*, *11*. https://doi.org/10.7554/eLife.83225
1041
1042 Neahring, L., Cho, N. H., & Dumont, S. (2021). Opposing motors provide mechanical and functional
1043 robustness in the human spindle. *Dev Cell*, *56*(21), 3006-3018.
1044 https://doi.org/10.1016/j.devcel.2021.09.011
1045
1046 Nicklas, R. B. (1983). Measurements of the force produced by the mitotic spindle in anaphase. *J Cell Biol*,
1047 *97*(2), 542-548. http://www.ncbi.nlm.nih.gov/pubmed/6885908
1048 Nicklas, R. B. (1988). The forces that move chromosomes in mitosis. *Annu Rev Biophys Biophys Chem*, *17*,
1049 431-449. https://doi.org/10.1146/annurev.bb.17.060188.002243
1050
1051 O'Toole, E., Morphew, M., & McIntosh, J. R. (2020). Electron tomography reveals aspects of spindle
1052 structure important for mechanical stability at metaphase. *Molecular Biology of the Cell*, *31*(3), 184-195.
1053 https://doi.org/10.1091/mbc.E19-07-0405
1054
1055 Ou, G., & Scholey, J. M. (2022). Motor cooperation during mitosis and ciliogenesis. *Annu Rev Cell Dev
1056 Biol*, *38*(1), 49-74. https://doi.org/10.1146/annurev-cellbio-121420-100107
1057
1058 Pampaloni, F., Lattanzi, G., Jonas, A., Surrey, T., Frey, E., & Florin, E. L. (2006). Thermal fluctuations of
1059 grafted microtubules provide evidence of a length-dependent persistence length. *Proc Natl Acad Sci U S
1060 A*, *103*(27), 10248-10253. https://doi.org/10.1073/pnas.0603931103
1061
1062 Parmar, S., Gonzalez, S. J., Heckel, J. M., Mukherjee, S., McClellan, M., Clarke, D. J., Johansson, M., Tank,
1063 D., Geisness, A., Wood, D., & Gardner, M. (2023). Robust microtubule dynamics facilitate low tension
1064 kinetochore detachment in metaphase. J Cell Biol, 222(8). https://doi.org/10.1083/jcb.202202085
1065
1066 Poirier, M. G., Eroglu, S., & Marko, J. F. (2002). The bending rigidity of mitotic chromosomes. *MBoC*,
1067 *13*(6), 2170-2179. https://doi.org/10.1091/mbc.01-08-0401
1068
1069 Powers, A. F., Franck, A. D., Gestaut, D. R., Cooper, J., Gracyzk, B., Wei, R. R., Wordeman, L., Davis, T. N.,
1070 & Asbury, C. L. (2009). The Ndc80 kinetochore complex forms load-bearing attachments to dynamic
1071 microtubule tips via biased diffusion. *Cell*, *136*(5), 865-875. https://doi.org/10.1016/j.cell.2008.12.045
1072 Press, W. H., Flannery, B. P., Teukolsky, S. A., & Vetterling, W. T. *Numerical Recipes in C: The Art of
1073 Scientific Computing*.

1074

1075 Rai, A., Liu, T., Katrukha, E. A., Estevez-Gallego, J., Manka, S. W., Paterson, I., Diaz, J. F., Kapitein, L. C.,
1076 Moores, C. A., & Akhmanova, A. (2021). Lattice defects induced by microtubule-stabilizing agents exert a
1077 long-range effect on microtubule growth by promoting catastophes. *PNAS*, *118*(51).
1078 https://doi.org/10.1073/pnas.2112261118

1079

1080 Renda, F., Magidson, V., Tikhonenko, I., Fisher, R., Miles, C., Mogilner, A., & Khodjakov, A. (2020). Effects
1081 of malleable kinetochore morphology on measurements of intrakinetochore tension. *Open Biol*, *10*(7).
1082 https://doi.org/10.1098/rsob.200101

1083

1084 Risteski, P., Jagric, M., Pavin, N., & Tolic, I. M. (2021). Biomechanics of chromosome alignment at the
1085 spindle midplane. *Curr Biol*, *31*(10), R574-R585. https://doi.org/10.1016/j.cub.2021.03.082

1086

1087 Rizk, R. S., Bohannon, K. P., Wetzel, L. A., Powers, J., Shaw, S. L., & Walczak, C. E. (2009). MCAK and
1088 paclitaxel have differential effects on spindle microtubule organization and dynamics. *MBoC*, *20*(6),
1089 1639-1651. https://doi.org/10.1091/mbc.e08-09-0985

1090

1091 Rusan, N. M., Fagerstrom, C. J., Yvon, A. M., & Wadsworth, P. (2001). Cell cycle-dependent changes in
1092 microtubule dynamics in living cells expressing green fluorescent protein-alpha tubulin. Mol Biol Cell,
1093 12(4), 971-980. http://www.ncbi.nlm.nih.gov/pubmed/11294900

1094

1095 Sanhaji, M., Friel, C. T., Wordeman, L., Louwen, F., & Yuan, J. (2011). Mitotic centromere-associated
1096 kinesin (MCAK): a potential cancer drug target. *Oncotarget*, *2*(12), 935-947.
1097 https://doi.org/10.18632/oncotarget.416

1098

1099 Sarangapani, K. K., Akiyoshi, B., Duggan, N. M., Biggins, S., & Asbury, C. L. (2013). Phosphoregulation
1100 promotes release of kinetochores from dynamic microtubules via multiple mechanisms. *Proc Natl Acad
1101 Sci U S A*, *110*(18), 7282-7287. https://doi.org/10.1073/pnas.1220700110

1102

1103 Schwietert, F., & Kierfeld, J. (2020). Bistability and oscillations in cooperative microtubule and
1104 kinetochore dynamics in the mitotic spindle. *New journal of physics*, *22*(5).
1105 https://doi.org/10.1088/1367-2630/ab7ede

1106

1107 Schwietert, F., Volkov, V. A., Pim J., H. V., Dogterom, M., Musacchio, A., & Kierfeld, J. (2022). Strain
1108 stiffening of Ndc80 complexes attached to microtubule plus ends. *Biophys J*, *121*(21), 4048-4062.
1109 https://doi.org/10.1016/j.bpj.2022.09.039

1110

1111 Skibbens, R. V., Rieder, C. L., & Salmon, E. D. (1995). Kinetochore motility after severing between sister
1112 centromeres using laser microsurgery: evidence that kinetochore directional instability and position is
1113 regulated by tension. *J Cell Sci*, *108 (Pt 7)*, 2537-2548. http://www.ncbi.nlm.nih.gov/pubmed/7593295

1114

1115 Skibbens, R. V., Skeen, V. P., & Salmon, E. D. (1993). Directional instability of kinetochore motility during
1116 chromosome congression and segregation in mitotic newt lung cells: a push-pull mechanism. *J Cell Biol*,
1117 *122*(4), 859-875. http://www.ncbi.nlm.nih.gov/pubmed/8349735

1118

1119 Sousa, A., Reis, R., Sampaio, P., & Sunkel, C. E. (2007). The *Drosophila* CLASP homologue, Mast/Orbit
1120 regulates the dynamic behaviour of interphase microtubules by promoting the pause state. *Cell Motility*,
1121 *64*(8), 605-620. https://doi.org/10.1002/cm.20208

1122
1123 Streitberg, B., & Rohmel, J. (1986). Exact distributions for permutations and rank tests: An introduction
1124 to some recently published algorithms. *Statistical Software Newsletter*, *12*, 10-17.
1125
1126 Stumpff, J., Ghule, P. N., Shimamura, A., Stein, J. L., & Greenblatt, M. (2014). Spindle microtubule
1127 dysfunction and cancer predisposition. *J Cell Phys*, *229*(12), 1881-1883.
1128 https://onlinelibrary.wiley.com/doi/abs/10.1002/jcp.24691
1129
1130 Stumpff, J., Wagenbach, M., Franck, A., Asbury, C. L., & Wordeman, L. (2012). Kif18A and
1131 chromokinesins confine centromere movements via microtubule growth suppression and spatial control
1132 of kinetochore tension. *Dev Cell*, *22*(5), 1017-1029. https://doi.org/10.1016/j.devcel.2012.02.013
1133
1134 Tien, J. F., Umbreit, N. T., Gestaut, D. R., Franck, A. D., Cooper, J., Wordeman, L., Gonen, T., Asbury, C. L.,
1135 & Davis, T. N. (2010). Cooperation of the Dam1 and Ndc80 kinetochore complexes enhances
1136 microtubule coupling and is regulated by aurora B. *J Cell Biol*, *189*(4), 713-723.
1137 https://doi.org/10.1083/jcb.200910142
1138
1139 Tirnauer, J. S., Canman, J. C., Salmon, E. D., & Mitchison, T. J. (2002). EB1 targets to kinetochores with
1140 attached, polymerizing microtubules. *MBoC*, *13*(12), 4308-4316. https://doi.org/10.1091/mbc.E02-04-
1141 0236
1142
1143 VandenBeldt, K. J., Barnard, R. M., Hergert, P. J., Meng, X., Maiato, H., & McEwen, B. F. (2006).
1144 Kinetochores use a novel mechanism for coordinating the dynamics of individual microtubules. *Curr Biol*,
1145 *16*(12), 1217-1223. https://doi.org/10.1016/j.cub.2006.04.046
1146
1147 Varga, V., Helenius, J., Tanaka, K., Hyman, A. A., Tanaka, T. U., & Howard, J. (2006). Yeast kinesin-8
1148 depolymerizes microtubules in a length-dependent manner. *Nat Cell Biol*, *8*(9), 957-962.
1149 http.://doi.org/10.1038/ncb1462
1150
1151 Volkov, V. A., Huis In 't Veld, P. J., Dogterom, M., & Musacchio, A. (2018). Multivalency of NDC80 in the
1152 outer kinetochore is essential to track shortening microtubules and generate forces. *Elife*, *7*.
1153 https://doi.org/10.7554/eLife.36764
1154
1155 Wan, X., Cimini, D., Cameron, L. A., & Salmon, E. D. (2012). The coupling between sister kinetochore
1156 directional instability and oscillations in centromere stretch in metaphase PtK1 cells. *Mol Biol Cell*, *23*(6),
1157 1035-1046. https://doi.org/10.1091/mbc.E11-09-0767
1158
1159 Waters, J. C., Skibbens, R. V., & Salmon, E. D. (1996). Oscillating mitotic newt lung cell kinetochores are,
1160 on average, under tension and rarely push. *J Cell Sci*, *109 (Pt 12)*, 2823-2831.
1161 http://www.ncbi.nlm.nih.gov/pubmed/9013330
1162
1163 Wieczorek, M., Bechstedt, S., Chaaban, S., & Brouhard, G. (2015). Microtubule-associated proteins
1164 control the kinetics of microtubule nucleation. *Nat Cell Biol*, *17*(7), 907-916.
1165 https://doi.org/10.1038/ncb3188
1166
1167 Winey, M., Mamay, C. L., O'Toole, E. T., Mastronarde, D. N., Giddings, T. H., Jr., McDonald, K. L., &
1168 McIntosh, J. R. (1995). Three-dimensional ultrastructural analysis of the Saccharomyces cerevisiae
1169 mitotic spindle. J Cell Biol, 129(6), 1601-1615. http://www.ncbi.nlm.nih.gov/pubmed/7790357

1170

1171 Wordeman, L., Wagenbach, M., & von Dassow, G. (2007). MCAK facilitates chromosome movement by
1172 promoting kinetochore microtubule turnover. *J Cell Biol*, *179*(5), 869-879.
1173 https://doi.org/10.1083/jcb.200707120

1174

1175 Yamashita, N., Morita, M., Legant, W. R., Chen, B., Betzig, E., Yokota, H., & Mimori-Kiyousue, Y. (2015).
1176 Three-dimensional tracking of plus-tips by lattice light-sheet microscopy permits the quantification of
1177 microtubule growth trajectories within the mitotic apparatus. *J Biomedical Optics*, *20*(10), 101206-
1178 101206. https://doi.org/10.1117/1.jbo.20.10.101206

1179

1180 Zar, J. H. (1999). *Biostatistical analysis* (4th ed.). Prentice Hall.

1181

1182 Zelinski, B., & Kierfeld, J. (2013). Cooperative dynamics of microtubule ensembles: Polymerization forces
1183 and rescue-induced oscillations. *Phys Rev*, *87*(1). https://doi.org/10.1103/physreve.87.012703