Decoupled evolution of the *Sex Peptide* gene family and *Sex Peptide Receptor* in

Drosophilidae

Ben R. Hopkins[1*], Aidan Angus-Henry[1], Bernard Y. Kim[2], Jolie A. Carlisle[3], Ammon Thompson[1],

Artyom Kopp[1]

[1] Department of Evolution and Ecology, University of California – Davis, CA, USA

[2] Department of Biology, Stanford University, CA, USA

[3] Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY, USA

*Corresponding author: brhopkins@ucdavis.edu

Abstract

Across internally fertilising species, males transfer ejaculate proteins that trigger wide-ranging

changes in female behaviour and physiology. Much theory has been developed to explore the

drivers of ejaculate protein evolution. The accelerating availability of high-quality genomes now

allows us to test how these proteins are evolving at fine taxonomic scales. Here, we use genomes

from 264 species to chart the evolutionary history of Sex Peptide (SP), a potent regulator of female

post-mating responses in *Drosophila melanogaster*. We infer that *SP* first evolved in the

Drosophilinae subfamily and has followed markedly different evolutionary trajectories in different

lineages. Outside of the *Sophophora-Lordiphosa*, *SP* exists largely as a single-copy gene with

independent losses in several lineages. Within the *Sophophora-Lordiphosa,* the *SP* gene family

has repeatedly and independently expanded. Up to seven copies, collectively displaying

extensive sequence variation, are present in some species. Despite these changes, *SP*

expression remains restricted to the male reproductive tract. Alongside, we document

considerable interspecific variation in the presence and morphology of seminal microcarriers that,

despite the critical role SP plays in microcarrier assembly in *D. melanogaster*, appear to be

independent of changes in the presence/absence or sequence of SP. We end by providing

evidence that SP's evolution is decoupled from that of its receptor, SPR, in which we detect no

evidence of correlated diversifying selection. Collectively, our work describes the divergent

evolutionary trajectories that a novel gene has taken following its origin and finds a surprisingly

weak coevolutionary signal between a supposedly sexually antagonistic protein and its receptor.

Significance

In insects, seminal fluid proteins (SFPs) induce dramatic changes in female behaviour and

physiology. How this degree of male influence evolves remains a central question in sexual

selection research. Here, we map the origin and diversification of the posterchild insect SFP, the

Drosophila Sex Peptide (SP), across 264 Diptera species. We show that *SP* first evolved at the

base of the subfamily *Drosophilinae* and followed markedly different evolutionary trajectories in

different lineages, including accelerated change in sequence, copy number, and genomic position

in the lineage leading to *D. melanogaster.* By contrast, we find only limited, uncorrelated change

in the sequence of its receptor, SPR, arguing against a sexually antagonistic coevolutionary arms

race between these loci on macroevolutionary time scales.

Introduction

Female post-mating changes are a taxonomically widespread – if not general – phenomenon in

internal fertilisers. Often mediated by non-sperm components of the male ejaculate, such as

seminal fluid proteins, the female traits subject to post-mating plasticity are numerous and diverse.

For example, immune systems can be modified (*Drosophila melanogaster*, 1 ; humans, 2),

ovulation stimulated (camelids, 3), and dietary preferences shifted following copulation (crickets,

4 ; *D. melanogaster*, 5). Evolutionary biologists have a long-standing interest in post-mating

changes as they bear intimate connections to reproductive success (e.g., 6–8), can form barriers

to hybridisation (e.g., 9, 10), and, through the involvement of males in their induction, can act as

a point of evolutionary tension between the fitness interests of males and females (e.g., 11–14).

Indeed, post-mating changes have provided one of the centrepieces around which much of the

discussion of interlocus sexual conflict has revolved, including broader consideration of the

relative roles of conflict and sexual selection by female choice in shaping the evolution of

reproductive traits (15–19).

Available data suggest that different taxa can use non-homologous proteins to induce common

– or at least overlapping – phenotypic endpoints in mated females. To reduce female sexual

receptivity, for example, the moth *Helicoverpa zea* uses pheromonostatic peptide (PSP, 20), the

mosquito *Aedes aegypti* uses Head Protein 1 (HP-1, 21), and *D. melanogaster* uses Sex Peptide

(SP, 22)*.* Not only are these three proteins non-homologous to one another, but none have clear

homologs in either of the other species' genomes. This pattern suggests that regulators of female

post-mating change might experience a high degree of evolutionary turnover, with new regulators

evolving and old regulators being lost from populations through time. If so, the questions then are

what evolutionary forces drive this turnover, how quickly does this process occur, and how are

new regulators born? More fundamentally, it makes seminal proteins an exceptional model for

studying how newly evolved, lineage-specific genes acquire and diversify their functions.

More than 60 years on from its discovery in chromatographic extracts of *D. melanogaster*

accessory glands (23, 24), SP remains the best characterised insect seminal protein. Consisting

of two exons separated by a 65bp intron, *D. melanogaster SP* (*DmelSP*) encodes a 36aa mature

protein synthesised via a 55aa signal peptide-containing precursor (22, 25). DmelSP is produced

in accessory gland main cells and secreted into the lumen, where it is stored on abundant, lipid-

containing microcarriers – structures for which DmelSP acts as a key factor governing assembly

and, once inside females, disassembly (26–28). Following transfer to females, DmelSP binds to

sperm, a process mediated by a suite of additional seminal fluid proteins, and is transported into

the female sperm storage organs (29–33). The gradual release of DmelSP from the surface of

stored sperm continues to stimulate a wide range of post-mating changes, including shifts in

memory formation and sleep patterns, elevating appetite and changing dietary preferences,

reducing sexual receptivity, stimulating egg-laying, increasing aggression, and changing gut,

metabolic, and immune activity (reviewed in 15). At least some of these changes, namely reduced

sexual receptivity and stimulated egg-laying, are mediated by DmelSP binding to the Sex Peptide

Receptor (SPR) in a subset of neurons that innervate the female reproductive tract (34–36).

Different domains of mature SP appear to contribute selectively to different functions in *D.*

melanogaster: the tryptophan-rich N-terminus binds to sperm and stimulates juvenile hormone

synthesis (33, 37, 38), the hydroxyproline-rich mid-section elicits the innate immune response

(39), and, through interactions with SPR, the disulphide bridge-containing C-terminus stimulates

the core post-mating responses of increased oviposition and reduced sexual receptivity (40–42).

Consequently, different portions of the *SP* coding sequence are likely to be evolving in response

to different selective pressures.

SP is not the only member of its gene family present in *D. melanogaster.* This species also

encodes the paralogous *Dup99b* with which SP shares a high degree of similarity in the amino

acid sequence of the C-terminus (43). Both stimulate the core post-mating responses of increased

oviposition and reduced sexual receptivity, but SP appears to be the 'key player' showing a higher

binding affinity for the female reproductive tract and nervous system (41) and, in *in vitro* assays,

activating SPR at lower concentrations than does Dup99b (36). There are further differences

between the paralogs, too. While *SP* is expressed in accessory gland main cells, *Dup99b* is

expressed in the ejaculatory duct (44). And, unlike SP, the N-terminus of Dup99b does not

stimulate juvenile hormone synthesis (38). Thus, SP and Dup99b show partial redundancy but

different sensitivities within one region of the protein and distinct activities in other regions,

suggesting a degree of functional separation between the two paralogs.

SP occupies an important place in contemporary evolutionary biology, having emerged as one of

the preeminent systems for experimental work on the genetic basis and fitness effects of sexual

conflict (6, 45–47). However, comparative data on how SP sequence and function has evolved

and diversified through time is sparse by comparison. Losses of *SP* have been reported in three

Drosophila species (*D. grimshawi, D. albomicans,* and *D. mojavensis*) and the gene's origin has

been traced as far back as the most recent common ancestor of *D. virilis* and *D. melanogaster;*

SP is apparently absent from mosquitoes and insect orders beyond Diptera (25, 48, 49). This

contrasts with its receptor, *SPR*, which is deeply conserved among members of the Ecdysozoa

and Lophotrochozoa, where it interacts with a similarly well conserved class of alternative ligands,

the myoinhibitory peptides (MIPs)(48). MIP-SPR interactions are known to regulate diverse

behaviours across species, including regulating larval settlement behaviour in marine annelids

(50). In *Drosophila,* MIP-SPR interactions appear to be neither necessary nor sufficient for driving

post-mating changes in females (48, 51), but they do fulfil other functions, including regulating

sleep behaviour (52). Despite SP predating the group, several features of the SP-SPR system

appear to be collectively restricted to the *melanogaster* species group, namely robust expression

of *SPR* in the female reproductive tract, the ability of DmelSP to bind to female reproductive tract

tissue, and a reduction in sexual receptivity upon injection of conspecific SP (25). Thus, despite

the presence of SP orthologs beyond the group, many of the defining features of SP in *D.*

melanogaster appear to be recently derived.

Taking advantage of newly available genomes for over 250 drosophilid species, here we report

that *SP* is a drosophilid innovation that originated in the lineage leading to the *Drosophilinae*

subfamily. We show that *SP* has subsequently followed markedly different evolutionary

trajectories in different branches of the phylogeny, including lineage-accelerated evolution in

sequence, copy number, and translocation frequency in the *melanogaster* group. Despite these

changes, *SP* expression remains restricted to the male reproductive tract. We further fail to find

support for the hypothesis that change in SP presence/absence or sequence is a significant driver

of evolutionary change in microcarrier morphology. Finally, we fail to find a signal of coevolution

between *SP* and the receptor through which it induces many of its effects in females, *SPR,* arguing

against a sexually antagonistic coevolutionary arms race between these loci on

macroevolutionary time scales.

Results

***Sex Peptide* first evolved in the *Drosophilinae* subfamily.** To pinpoint the origin of *SP,* we

designed a pipeline to identify *SP* orthologs in whole-genome sequences. Our approach

combined reciprocal blast of the *D. melanogaster* SP C-terminus sequence with protein

sequence, gene structure, and synteny analysis. As in a previous study (48), we failed to detect

SP in a genome from the mosquito *Aedes aegypti,* and additionally failed to detect *SP* in genomes

from two calyptrate members of the Brachycera suborder to which *Drosophila* belongs: *Musca*

domestica (53) and *Glossina morsitans* (54). Within the Acalyptratae, we found that *SP* was

restricted to the *Drosophilidae* – specifically to the subfamily *Drosophilinae* Rondani (Figure 1).

Within the *Drosophilinae*, *SP* was present in genomes from several species that predate the

Drosophilini, the tribe that includes all members of the *Drosophila* genus (55). These non-

Drosophilini species included members of the *Colocasiomyini* tribe, including species within the

Scaptodrosophila, Lissocephala, Chymomyza, and *Colocasiomyia* genera, but several secondary

losses of SP were apparent. We failed to detect *SP* in genomes from 10 members of the other

Drosophilidae subfamily, the *Steganinae* Hendel, or members of several closely related

outgroups, namely the genera *Liriomyza, Cirrula, Ephydra,* and *Diastata.* We therefore infer that

SP first evolved in the *Drosophilinae* subfamily.

Figure 1. *SP* first evolved in the *Drosophilinae* subfamily. A phylogeny of species from across the *Drosophilidae* and closely related lineages. The number of *SP* genes detected is given in coloured circles at the tip of each branch. A brown bar denotes the branch on which we infer *SP* to have first evolved.

Sex Peptide has been repeatedly lost and rarely duplicated outside of the *Sophophora-*

***Lordiphosa* radiation.** Within our phylogenetic sample, the *Drosophilini* splits into two lineages.

The first contains the *Lordiphosa* genus and *Sophophora* subgenus (which we collectively refer

to as the '*Sophophora-Lordiphosa* radiation'), the latter of which includes the *melanogaster,*

obscura, willistoni, and *saltans* groups (see *SI Appendix Fig. S1* for an overview of the

Sophophora taxonomic terminologies used in this paper)*. The second lineage includes, among

others, the genera *Scaptomyza* and *Zaprionus* and the Hawaiian*, virilis, repleta, immigrans,*

cardini, and *quinaria* groups (the *Drosophila* genus is paraphyletic (55, 56)).

Outside of the *Sophophora-Lordiphosa* radiation*, we observed several features of *SP'*s evolution.

First, *SP* has been repeatedly and independently lost—four times in our phylogenetic sample.

Once in a monophyletic lineage of 29 species covering the *annumilana, bromeliae*, *nannoptera*,

mesophragmatica, and *repleta* groups (Figure 2; *SI Appendix, Fig. S2*). The other three separately

covered all 55 Hawaiian species in our dataset (Figure 2; *SI Appendix, Fig. S3*), a monophyletic

lineage within the *Scaptomyza* (Figure 2; *SI Appendix, Fig. S3*), and a species-specific loss in

Hirtodrosophila duncani (Figure 3)*. The second trend was that duplications of *SP* were rare.

Among the 53 non-*Sophophora-Lordiphosa* species in the *Drosophilini* in which we did detect *SP,*

all but one had just a single copy (Figure 2). The exception was *D. paramelanica* (*melanica* group)

in which we detected a tandem duplication, with the two copies sharing 100% identity in predicted

protein sequence. Outside of the *Drosophilini,* only *Scaptodrosophila latifasciaeformis* showed an

expansion in *SP* copy number, bearing 3 tandemly arranged copies that diverged from one

another in predicted protein sequence. The third trend was that *SP* rarely translocated to new

genomic locations. Where *SP* was detected in non-*Sophophora-Lordiphosa* species in the

Drosophilini, in all but two we found that it mapped to a syntenic neighbourhood on Muller element

D, which we call 'Muller D1', that contained orthologs of *FoxK, NaPi-III,* and *mRpL2.* The

exceptions were *Hirtodrosophila trivittata,* where *SP* mapped to a distinct neighbourhood on

Muller element D that contained orthologs of *bruno3, CG3349,* and *CG17173,* and *D. repletoides*,

in which *SP* mapped to a neighbourhood on Muller element B containing orthologs of *halo* and

haf. Outside of the *Drosophilini,* we also detected *SP* in the Muller D1 neighbourhood in *S. lebanonensis* and *S. latifasciaeformis,* suggesting that this may be the ancestral position of *SP.*

Figure 2. *Sex Peptide* family genes predate the *Drosophilini* and have been repeatedly lost outside of the *Sophophora-Lordiphosa* radiation. (Legend continues on next page).

Figure 2. ***Sex Peptide*** **family genes predate the *Drosophilini* and have been repeatedly lost outside of the *Sophophora-Lordiphosa* radiation.** This figure focuses on the non-*Sophophora-Lordiphosa* members of the *Drosophilini* (see Figure 1 and *SI Appendix, Fig. S1* for overviews of drosophilid taxonomy). A selection of *Sophophora* species, shaded in blue, are included for comparison. Also included are four non-*Drosophilini* dipterans: *Aedes aegypti* and three non-*Drosophilini* members of the *Drosophilinae* subfamily: *Chymomyza costata, Scaptodrosophila lebanonensis,* and *S. latifasciaeformis.* The number of *SP* genes detected in a representative of each species' genome is given at the tree tips. Losses are marked with a pink bar. For each species, the structures of two syntenic gene neighbourhoods are plotted. The first, Muller D1, is the canonical neighbourhood in which *SP* genes are detected outside of the *Sophophora-Lordiphosa.* The second, Muller D2, is the canonical position in the *Sophophora-Lordiphosa.* Positions of each gene are given relative to the first gene in the neighbourhood (*FoxK* or *capricious*). Absence of a flanking neighbourhood gene (*e.g., FoxK* in *D. wassermani*) doesn't necessarily mean the gene has been lost – it more likely means that a contig breakpoint fell within the neighbourhood. Note that *SP* in *Hirtodrosophila trivittata* and, independently, *D. repletoides,* has translocated out of the Muller D1 neighbourhood. See *SI Appendix, Figs S2 and 3* for expanded views of the losses in the *repleta* and *Hawaiian* groups, respectively.

Sex Peptide **has repeatedly duplicated in the *Sophophora-Lordiphosa* radiation.** Within the

Sophophora-Lordiphosa, SP has followed a markedly different evolutionary trajectory. For one,

and despite denser taxon sampling in this part of the phylogeny, we detected at least one copy in

all species sampled—no species was entirely without *SP.* Second, we detected a clear uptick in

the frequency of duplication (Figure 3). In the earlier branching lineages*,* we detected apparently

independent duplications within a sublineage of the *Lordiphosa,* in *D. subobscura* (obscura group;

see also 57), and in *D. lowei* (*obscura* group). Within the *melanogaster* group we found much

greater variability in *SP* copy number, consistent with repeated, lineage-specific expansions and

contractions of gene family size. This variation was greatest in the "Oriental" lineage, which

includes *D. melanogaster,* and *ananassae* subgroup. As many as 7 tandemly arranged copies

were present in some *ananassae* subgroup species (Figure 4).

Figure 3. *Sex Peptide* copy number is markedly more variable in the *Sophophora-Lordiphosa* radiation than in other branches of the phylogeny. This figure focuses on the *Sophophora-Lordiphosa* radiation to which *D. melanogaster* belongs. Four non-*Sophophora-Lordiphosa* drosophilids, shaded in purple, are included as outgroups. The structures of the Muller D1 and Muller D2 neighbourhoods are plotted as in Figure 2. Missing flanking genes are likely indicative of contig breakpoints falling within the neighbourhood. The exceptions are the *Lordiphosa* species, where substantially elevated intergenic distances meant that the whole neighbourhood would not

fit within the plot limits. Unresolved SP genes, shown in red, indicate genes that passed the reciprocal blast criteria and fell within one of the conserved *SP*-containing gene neighbourhoods but where a SP-like amino acid sequence couldn't be resolved (*e.g.,* due to a premature stop codon, as in the case of *D. rhopaloa*). Note that all members of the *obscura* group have an inversion that flips the relative positions of *FoxK* and *mRpL2* in the Muller D1 neighbourhood. In a number of cases, some or all copies of *SP* were found to have translocated outside of the Muller D1 and Muller D2 neighbourhoods (an *obscura* group lineage, the *melanogaster* subgroup, *D. kanapiae, D. takahashii*, and *D.eugracilis*; summarised in *SI Appendix Fig. S5*). In the shorter read *montium* subgroup assemblies, short contigs meant that in some species we couldn't identify the neighbourhood in which *SP* was located. This was the case for some *SP* genes in *D. cf. chauvacae, D. burlai, D. leontia, D. nikananu, D. pectinifera, D. punjabiensis,* and *D.watanabe*i. The *SP* genes in *D. auraria* and *D. trapezifons* could be mapped to the Muller D2 neighbourhood based on flanking sequence around the *SP* gene, but the *SP*-containing contigs were too small to include any of the neighbourhood genes.

To resolve the evolutionary relationships between *SP* paralogs within the *Sophophora-Lordiphosa*

radiation, we constructed a tree of the predicted SP protein sequences (*SI Appendix, Fig. S4*).

The tree supports numerous recent duplications affecting single species or species pairs,

including in *D. elegans/D.gunungcola, D. takahashii, D.ficusphila, D. punjabiensis/D. watanabei,*

D. kanapiae, and *D. subobscura*. The tree also suggests that there have been three separate

expansions of *SP* copy number within the *ananassae* subgroup, one in the *bipectinata* complex

and another in each of *D. varians* and *D. ercepeae* (*SI Appendix, Fig. S4*, coloured orange and

blue, respectively), likely from an ancestral starting point of the two copies seen in *D. ananassae.*

We note, however, that the sequence similarity that we observe between putative paralog pairs

may be driven instead by concerted evolution.

Frequent translocations of *Sex Peptide* genes within the *Sophophora* subgenus. At the

base of the *Sophophora-Lordiphosa* radiation, *SP* appears to have translocated from the Muller

D1 neighbourhood to a new neighbourhood ~2.1Mb away on the same Muller element (Figure 3).

This syntenic neighbourhood, which we refer to as 'Muller D2', contains orthologs of *capricious,*

CG14111, and *CG17687.* Despite translocation, the configuration of the ancestral Muller D1 gene

neighbourhood remains intact in the *Sophophora-Lordiphosa*. Thus, the mechanism of

translocation did not lead to the breakup of the Muller D1 neighbourhood via a larger scale

rearrangement. Several further translocations are then present (summarised in *SI Appendix, Fig.*

S5; *e.g.,* in the *obscura* group, *SI Appendix, Fig. S6*). Each of *D. suzukii, D. subpulchrella,* and *D.*

biarmipes, which form a monophyletic clade within the "Oriental" lineage, bear SP copies in the

canonical Muller D2 position with an additional copy just the other side of *CG14111* within the

same neighbourhood (Figure 3). The protein tree supports a *Dup99b* identity for these copies that

have skipped to the other side of *CG14111* (*SI Appendix, Fig. S4*, shown in pink)*.* In *D. takahashii,*

the sister species to this clade, two of the three SP genes we detected mapped to a

neighbourhood on *D. melanogaster* Muller element B that contained *NLaz, robo2,* and *CG14346*

orthologs. The protein tree also supports a *Dup99b* identity for these translocated copies (*SI*

Appendix, Fig. S4, shown in pink). However, given the low support for internal nodes, we were

not able to accurately determine the timing of the initial duplication that gave rise to separate *SP*

and *Dup99b* copies.

Figure 4. Repeated duplication of *Sex Peptide* genes in the *ananassae* subgroup. A phylogeny of the *ananassae* subgroup species used in this study, with *D. melanogaster* as an outgroup. The number of *SP* genes identified in each species is given at the tip of each branch. 'Unresolved' SP sequences, shown in red, are those which passed the reciprocal blast tests and fell within the syntenic Muller D2 gene neighbourhood but for which we could not resolve an SP-like protein sequence (*e.g.,* due to a premature stop codon). The structure of the neighbourhood is plotted on the right-hand side of the figure. Note that one of *D. melanogaster's SP* copies, *Dup99b,* falls outside of the Muller D2 neighbourhood.

In *D. melanogaster, SP* falls within the Muller D2 neighbourhood while its paralog *Dup99b* maps

to Muller element E in a neighbourhood that contains *dmrt99b, gycalpha99b,* and *CG34296.* This

arrangement appears to be ancestral to the *melanogaster* subgroup (*SI Appendix, Fig. S7A*,*B*).

The losses of an *SP* gene within a subset of species in this subgroup, namely *D. teissieri* (strain

CT02; present in 273.3*), D. orena,* and *D. erecta,* affect the Muller element E *Dup99b* copy, rather

than the Muller D2 *SP* copy. In *D. eugracilis,* the *melanogaster* subgroup's sister species, *SP* is

present in the Muller D2 neighbourhood, with a second copy in a different position on Muller

element E that contains orthologs of *SmD2, CG18048,* and *Hr83*. Despite the lack of synteny, the

protein tree supports a *Dup99b* identity for this translocated copy (*SI Appendix, Fig. S4*).

Male reproductive tract-biased expression is a conserved feature of *SP* genes. Our

identification of *SP* genes was based on gene sequence data and synteny, leaving open the

question of whether and where they are expressed. For 19 species, we were able to test for

expression using RNA-seq datasets available through NCBI. 38 of the 42 *SP* genes we detected

across the 19 species were expressed, although many were un- or incorrectly annotated (*e.g.,* as

long non-coding RNAs) in the reference genomes (see *SI Appendix*). All 4 of those that weren't

expressed lacked SP-like protein sequences due to point mutations affecting either the start

codon or introducing premature stop codons, suggestive of pseudogenisation (*e.g.,* in *D.*

rhopaloa; *SI Appendix, Fig. S8*). Of the 38 *SP* genes we found to be expressed, all showed

strongly male-biased expression, including in the early branching *D. busckii* (*SI Appendix, Fig.*

S9). The one exception was detection of appreciable *SP* expression in a single *D. simulans* female

sample, which was due to sample contamination or mislabelling (*SI Appendix, Fig. S10*)*.*

In all 10 species where we had tissue-specific expression data, including the distantly related *D.*

virilis, we observed clearly enriched expression of *SP* family genes in the male reproductive tract

(*SI Appendix, Fig. S11*). Where datasets were available for sub-portions of the male reproductive

tract, expression was generally substantially higher in samples labelled as accessory gland or

non-gonadal reproductive tissues than in samples labelled as testes. The extent of testes

expression was variable between samples and between species, perhaps reflecting varying

degrees of contamination between these closely associated tissues during dissection. Based on

these data, we conclude that male reproductive tract-biased expression is a conserved feature of

the *SP* gene family and is therefore likely the ancestral expression pattern within the *Drosophilini.*

Accelerated evolution of Sex Peptide proteins in the *Sophophora-Lordiphosa* radiation.

Aligning 233 SP sequences from 148 genomes for the species shown in Figure 2 and 3, we find

that the C-terminus, which is responsible for stimulating post-mating responses in *D.*

melanogaster (33, 42)*,* is highly conserved both inside and outside of the *Sophophora-Lordiphosa*

radiation (*SI Appendix, Fig. S12A-C*). Several residues in this region, including the disulphide

bond forming cysteine residues, are present in almost all SP sequences in our dataset: within the

consensus sequence KWCRLNLGPAWGGRGKC, W_2, C_3, G_8, P_9, G_{12}, G_{13}, and C_{17} are each

conserved in >97% of sequences (*SI Appendix, Fig. S12A*). In contrast, the mid-section, which

has been implicated in stimulating innate immune responses (39), and the N-terminus (following

cleavage of the signal peptide), which is responsible for binding to sperm and stimulating juvenile

hormone synthesis (33), showed quite limited sequence conservation, suggesting more rapid

evolutionary change.

The predicted length of SP proteins showed elevated variability in the *Sophophora-Lordiphosa*

radiation (*SI Appendix, Fig. S12D*). These differences held after *in silico* cleavage of predicted

signal peptides and were largely due to the introduction of additional amino acids upstream of the

post-mating-response-stimulating C-terminus. For 38 genes across 19 species, we were able to

use the RNA-seq data to validate our annotation of exon/intron boundaries. In all 38 cases our

predicted boundaries matched those derived from the RNA-seq data. The expression data alone

therefore supports a change in pre-cleavage SP sequence length between *e.g., D. virilis* (47aa)

and *D. rhopaloa* (60aa), as well as *D. bipectinata* expressing a set of 6 SP proteins of variable

length (46aa, 49aa, 54aa, 62aa, 68aa, 72aa). Moving on to SP protein sequence, a PCA

generated from substitution matrix scores showed a high degree of dispersion among

Sophophora-Lordiphosa orthologs relative to those from outside of the radiation (*SI Appendix,*

Fig. S12E-G). This included clear separation from the remaining sequences of SP – but not

Dup99b – orthologs from the "Oriental" lineage (except for the most basal species we sampled

from this lineage, *D. oshimai*). Their distinct clustering may be driven by their N-terminus and

midsection sequences, which showed limited conservation with those of other SP proteins. More

generally, the high degree of dispersion between sequences in the *Sophophora-Lordiphosa,* and

particularly the *ananassae* subgroup, points to a high degree of sequence diversity within this

lineage.

Microcarrier morphology is not clearly linked to the copy number of *Sex Peptide*. We next

wanted to explore functional consequences of the diversity in the phylogenetic distribution and

sequence of *SP* genes*.* Recently, SP was shown in *D. melanogaster* to be a key factor influencing

the assembly, disassembly, and morphology of microcarriers, lipid-based structures that appear

to store and traffic seminal fluid proteins (28). Because of this relationship between SP and

microcarrier structure, it was suggested that variation in SP sequence might be associated with

inter-specific variation in microcarrier morphology (28).

Using the neutral lipid-specific dye LipidTox, which has previously been used to stain

microcarriers (28), we sought to examine the relationship between SP and microcarrier structure

on two levels. The first was to ask whether variation in *SP* copy number is associated with a shift

in microcarrier morphology. For this, the *ananassae* subgroup provides an ideal system, given

that its constituent species encode between 2 and 7 *SP* copies. Within the *bipectinata* species

complex, all of the species that we looked at (*D. parabipectinata, D. bipectinata, D. m.*

malerkotliana, D. m. pallens, D. p. nigrens), which each encode between 5 and 7 *SP* copies,

showed small, globular microcarriers (Figure 5A-F), similar to those seen in the *obscura* group

(28). Curiously, however, and unlike those seen in the *obscura* group, these microcarriers

appeared to carry a central indentation reminiscent of the biconcave disk shape of human red

blood cells. This morphology was clearly distinct from that of *D. melanogaster* microcarriers, which

appear as a heterogeneous mix of fusiform, ellipsoid, and thread-like structures (Figure 5L). The

bipectinata complex morphology was also distinct from those of three other *ananassae* subgroup

species: *D. ananassae* (2 copies), which had thread-like and spiral or doughnut shaped

microcarriers (Figure 5G); *D. ercepeae* (5 copies)*,* which had thread-like microcarriers (Figure

5H); and *D. varians* (4 copies), which displayed a highly divergent organization of the lumen's

contents (Figure 5I,J). In *D. varians*, LipidTox appeared to be excluded from vacuolar structures

that were filled with small, weakly stained droplets. The vacuoles appeared larger in the proximal

region of the gland, suggesting that they may fuse as they move towards the ejaculatory duct

(Figure 5J). We observed a similar pattern in the three-copy-encoding *D. takahashii*, a non-

ananassae subgroup species, although here the LipidTox staining was negligible (Figure 5N).

This contrasted with *D. takahashii*'s close relative *D. biarmipes,* which despite also encoding three

copies showed a unique staining pattern of strongly stained, tiny microcarriers that appeared to

aggregate (Figure 5M). The microcarriers of *D. biarmipes* adopted a conformation reminiscent of

D. melanogaster's following transfer to the female reproductive tract as they begin to break down

into smaller puncta (28). The conformation observed in *D. biarmipes* appears to be a derived state

as the more distantly related *D. carrolli* (Figure 5O) bears microcarriers that more closely resemble

those of *D. melanogaster, D. sechellia,* and *D. simulans* (Figure 5L; 23)*.* Consequently, while the

melanogaster group shows remarkable diversity in both microcarrier morphology and *SP* copy

number, there appears to be no clear relationship between them.

Figure 5. *Sex Peptide* is neither necessary nor sufficient for microcarriers. (A) A phylogeny of all species included in this figure. Branches coloured pink indicate species shown in B-T that demonstrate canonical (*i.e., D. melanogaster*-like) staining with LipidTox, a neutral lipid-specific dye used to selectively stain microcarriers (28). (B-T) Accessory glands stained with LipidTox and the nuclear stain DAPI (blue). The circled number in the bottom right-hand corner of each panel

indicates the number of *Sex Peptide* copies we detect in each species. (G) The arrow is highlighting a spiral/doughnut shaped microcarrier, a shape which is rare in comparison to the more common thread-like conformation in this species. (J) Prox. refers to the proximal region of the gland, *i.e.* the region that connects to the ejaculatory duct. (P) Arrows point to the ambiguous, sparse, and weakly stained material we observed in *D. ficusphila* glands. (S) An arrow points to a cut in the glandular epithelium, which was made to enhance dye penetration. In each case, glands were co-stained in the same well with those from *D. melanogaster* to act as a positive control.

Detection of microcarrier-like, but LipidTox⁻, structures within and beyond *Sophophora*.

Previous staining of accessory glands from the single-copy-encoding *D. virilis* demonstrated that

a copy of *SP* is not sufficient for LipidTox-stained microcarriers (28)*. D. virilis* instead displayed a

more uniform 'flocculence' within the gland's lumen that showed little evidence of LipidTox

staining. We observed a similar flocculent arrangement in the single-copy-encoding *D. immigrans*

(Figure 5Q). We also observed an essentially microcarrier-free glandular lumen in the four-copy-

encoding *D. ficusphila* (Figure 5P). In this species we observed only a handful of weakly stained

structures per gland, the rarity and structural inconsistency of which renders their classification

as microcarriers doubtful. Alongside these cases, we detected instances of microcarrier-like,

ellipsoid structures that failed to take up LipidTox in several species from diverse parts of the

drosophilid tree, namely the *montium* subgroup species *D. kikkawai* (Figure 5K) and the non-

Sophophora species *Scaptomyza hsui* (Figure 5S)*. All four of these species – *D.immigrans, D.*

ficusphila, D.kikkawai, and *S. hsui* – each encode at least one *SP* copy, providing further support

for the claim that a copy of *SP* is not sufficient for LipidTox⁺ microcarriers.

Microcarriers predate the *Sophophora*, but copies of *Sex Peptide* are neither necessary

nor sufficient for their presence. Staining glands from a species that we identified as having

lost the *SP* gene*, S. flava,* we observed small, globular microcarriers reminiscent of those from

the *obscura* group, albeit weaker in their staining (Figure 5R). Thus, *SP* is not necessary for

microcarriers. Moreover, the previous complement of species that had been stained suggested

that LipidTox[+] microcarriers were confined to the *obscura* and *melanogaster* groups. We now

show that they are present outside the *Sophophora.*

To better understand the distribution of microcarriers within the *Scaptomyza,* we also looked at

S. flava's single-copy-encoding sister species, *S. pallida*. This species showed strong LipidTox

staining, but the pattern was unlike any other species we looked at (Figure 5T). Rather than the

lumen being filled with large numbers of small microcarriers with well-defined shapes, the *S.*

pallida lumen was filled with substantial clouds of stained secretion that in many cases spanned

the full diameter of the gland's internal space. This pattern was reminiscent of that observed in

repeatedly mated *SP* null – but not wild-type – *D. melanogaster* males (28)*.* Thus, in the presence

of an *SP* ortholog we observe in *S. pallida* an apparent phenocopying of an *SP* null conformation,

further evidence that microcarrier morphology may, at a broad taxonomic scale, be largely

decoupled from evolutionary change in *SP.*

No clear signal of episodic diversifying selection in SPR sequence among drosophilids. If

SP is coevolving with the G-protein coupled receptor (GPCR) through which it induces many of

its effects, SPR (36), then we might predict that bursts of evolutionary change in *SP* copy number

and protein sequence correlate with similar bursts of change in SPR. Resolving *SPR* sequences

from 193 genomes (the species shown in Figures 2 and 3), we failed to detect a single instance

of duplication, suggesting that the mode of evolutionary change is decoupled between *SP* and

SPR. For 5 species, including two that had lost *SP* (*S. flava* and *S. montana*), we failed to resolve

an SPR sequence (*SI Appendix, Fig. S13*). Thus, while there is no phylogenetically repeatable

association in the copy number of the two genes, the loss of one can be accompanied by the loss

of the other.

Figure 6. SPR residues showing evidence of episodic positive selection are enriched in the ligand-facing domains. (A) A consensus sequence based on MAFFT alignment of the resolvable amino acid sequences of *SPR* coding sequences. The top plot gives the proportion of sequences with the consensus amino acid in the same position, while the bottom plot gives the proportion of sequences in which each position is occupied in the alignment. Each residue is coloured based on the functional domain to which it belongs based on the UniProt annotations. The red line indicates the start of the conserved region we use in the molecular evolution analyses. Beneath the plot, we show the corresponding exon that encodes each consensus residue. EC = extracellular, IC = intracellular (B) The AlphaFold prediction of the structure of *D. melanogaster* SPR as downloaded from UniProt (AF-Q8SWR3-F1) and coloured by the domain each residue belongs to based on positions listed in the UniProt 'Features' table. (C) A boxplot showing the proportion of residues at each position that matched the consensus residue (*i.e.,* the degree of conservation at each position). Residues are plotted separately according to their domain class of origin. (D) PCA plot based on BLOSUM62 substitution scores from the MAFFT-aligned SPR protein sequences. The percentage values in the axis titles reflect the proportion of variance explained by a given PC. Points are coloured based on whether they correspond to *Sophophora-Lordiphosa,* non-*Sophophora-Lordiphosa,* or *ananassae* subgroup species. (E) As (D) but after removing the non-conserved region of the N-terminus (*i.e.,* the region preceding the red line in (A)). (F) The same prediction shown in (B) but with residues coloured by the proportion of conserved residues per position (see also Supplementary movie 1). High values indicate high conservation. (G) The ColabFold top-ranked prediction of the interactions between

SP (shown in green) and SPR (residues coloured by domain). (H) As in (G) but with predicted contact residues coloured purple and predicted hydrogen bonds between SP and SPR residues shown with red-dotted lines (See also Supplementary movies 2 and 3). (I) A bar chart showing the proportion of sequences sharing the consensus residue for each predicted contact residue. Bars are coloured by whether the residue is also predicted to form a hydrogen bond with SP. Asterisks denote predicted contact residues for which we detected evidence of episodic positive selection using *MEME.* Alongside the plot, coloured bars, using the same colour scale as in (A-C), denote the functional domain the residue falls within. The two adjacent blue bars denote separate, consecutive transmembrane domains. Alongside are the corresponding amino acid residues in each position for each of the consensus ('Cons.'), *D. melanogaster* ('Dmel.'), *D. bipectinata* ('Dbip.'), *D. pseudoobscura* ('Dpse.'), *D. virilis* ('Dvir.'), and *A. aegypti* ('Aaeg.') sequences. Residues that depart from the mode among these plotted sequences are coloured. (J) As (G) but colouring only the 10 residues in SPR for which we detected evidence of positive selection using *MEME.* Asterisks denote predicted contact residues that show evidence of positive selection.

At the sequence level, we found that evolutionary change in SPR is concentrated in the

extracellular N-terminus domain, perhaps consistent with evolution under relaxed selection (see

SI Appendix, Supp. Text), with the remainder of the protein sequence showing much stronger

conservation (Figure 6A-C; *SI Appendix, Fig. S13*). PCA suggested that while the degree of

diversity among SPR sequences was apparently higher in the *ananassae* subgroup and "Oriental"

lineages relative to the *montium* subgroup, the overall diversity did not appear markedly elevated

in these lineages compared to the full spectrum of SPR sequences (Figure 6D, E). After removing

the poorly conserved N-terminus region, we found no evidence for episodic diversifying selection

in *SPR* sequences using branch- (aBSREL, 58) or gene- (BUSTED, 59) level tests for selection.

This held whether we formally tested all branches, only those in the *Sophophora-Lordiphosa*

radiation, or only those outside of the *Sophophora-Lordiphosa* radiation (see *SI Appendix* for how

to view the full output of these analyses). Thus, we fail to find evidence of a burst in evolutionary

change in *SPR* that correlates with the lineage-accelerated evolutionary changes we record for

SP.

SPR sites with evidence of episodic positive selection are disproportionately located in

predicted extracellular facing domains. Several residues in the extracellular loops and

extracellular facing transmembrane domains – regions likely to be critical for ligand-binding –

showed reduced conservation (Figure 6A, F; Supplementary movie 1). To assess whether these

sites are under selection, we used *FUBAR* (60) to test for evidence of pervasive diversifying

selection at individual sites in the N-terminus trimmed SPR sequence. We detected evidence of

pervasive diversifying selection at 0/377 sites and purifying selection at 357/377. We followed this

analysis with a test for *episodic* positive selection at individual sites, implemented through *MEME*

(61). In this analysis, we detected evidence of episodic positive selection at 10/377 sites. Running

the same analysis separately for the *Sophophora-Lordiphosa* (91 species) and non-*Sophophora-*

Lordiphosa species (89 species), we identified 4/371 and 5/376 sites respectively as showing

evidence of episodic positive selection (the identity of these positively selected sites did not

overlap between the two analyses). Therefore, we found no evidence that the proportion of sites

experiencing episodic positive selection was elevated in the *Sophophora*, which includes the

lineages in which *SP* showed greatest evolutionary change.

Intriguingly, of the 10 positively selected sites identified in the phylogeny-wide analysis, 9 fell

within extracellular domains: 3 in the N-terminus region close to the start of the first

transmembrane domain, 6 across the three extracellular loops, and then one in the third

intracellular loop. Using ColabFold (62–65) to generate a model of SP-SPR interactions (*SI*

Appendix, Fig. S14A-C) and ChimeraX (66), we detected 33 residues in SPR that were predicted

to interface with SP, of which 12 were additionally predicted to form hydrogen bonds with SP

residues (Figure 6G-I; Supplementary movies 2, 3). Most of these residues were highly conserved

across the 193 sequences: 12/33 were 100% conserved and 20/33 were >99% conserved (Figure

6I). But this level of conservation was not atypical among the extracellular facing residues: those

that neighboured the predicted contact residues were similarly well conserved (*SI Appendix, Fig.*

S14E). Of the 13 less well conserved residues, none showed clear evidence of concerted change

among *ananassae* subgroup, Oriental lineage, or *Sophophora* species. However, there was a

significant enrichment of sites showing significant evidence of episodic positive selection among

the 33 predicted contact residues (4/33; χ^2=12.56, *df* =1, *p*=0.0004; marked by asterisks in Figure

6I,J). Overall, therefore, while we do detect evidence that sites in the putative SP-binding pocket

of SPR have undergone episodic positive selection, the number of changes does not appear to

be elevated in the *Sophophora-Lordiphosa* radiation where the major genomic and functional

changes (e.g., 25) in SP have occurred.

A validated mutational route that SPR could take to decouple responses to SP and MIPs

remains unexploited in drosophilids. If receipt of SP is associated with a net reduction in

female fitness, the potentially deleterious effects of disrupting MIP-SPR interactions may

constrain SPR's ability to evolve to defend against SP binding. However, substitution of certain

residues in SPR can have decoupled effects on the receptor's sensitivity to its different ligands.

Specifically, replacing the QRY motif at the boundary between the second intracellular loop and

third transmembrane domain with the DRY motif more widely found in class A GPCRs is

associated with a decrease in the responsiveness of SPR to SP, but not the ancestral MIP ligands,

in *in vitro* assays (51). Yet in no drosophilid did we detect change at this position. Therefore, this

potential avenue through which substitution of a single amino acid, albeit requiring change at two

nucleotide positions, could reduce sensitivity to SP without affecting pre-existing ligand

interactions remains unexploited.

Discussion

Over the past few decades, we've built up a detailed understanding of the function of SP in *D.*

melanogaster. We know that it is required for the normal assembly and disassembly of seminal

storage and trafficking structures ("microcarriers", 28); that it triggers an extensive range of

physiological and behavioural changes in females, at least some of which are mediated by its

interactions with SPR in female reproductive tract neurons (34–36); and that SP's effects in

females are extended via its binding to the surface of sperm, a process facilitated by a network of

other male-derived proteins (29, 30, 32, 33). And yet, previous work has suggested that despite

its integral roles in *D. melanogaster* reproduction, and despite the complex sperm-binding

machinery with which it interacts, *SP* is restricted to drosophilids and perhaps, therefore, a

drosophilid innovation (48). Consequently, *SP* represents a powerful system in which to chart the

origin and diversification of function in a novel gene across different lineages. The cross-species

analysis of *SP* and *SPR* genes that we report here makes several contributions to this.

The first relates to the phylogenetic distribution of *SP* genes. Previous work traced *SP* as far back

as the split between *D. melanogaster* and *D. virilis* and identified three apparently independent

loss events in the non-*Sophophora* species *D. grimshawi, D. mojavensis,* and *D. albomicans* (25,

48, 49). The data we present here pushes the origin of *SP* back to the base of the *Drosophilinae.*

We also showed that *SP* is present in a genome of *D. albomicans* and that the losses in *D.*

grimshawi and *D. mojavensis* are not species-specific, but instead cover much larger radiations,

including all 55 members of the Hawaiian radiation that we sampled and, independently, the

lineage leading to the *annulimana*, *bromeliae*, *nannoptera,* and *repleta* groups. Alongside, we

detected evidence of additional losses in *H. duncani* and a lineage of *Scaptomyza*. Given the

critical role of *SP* in many aspects of *D. melanogaster* reproduction these losses are intriguing

and may reflect a lower functional importance of *SP* outside of the *Sophophora*. Consistent with

this, SP injection experiments suggest that the ability of SP to reduce female sexual receptivity is

restricted to the *melanogaster* group (25). Where *SP* has been lost it may be that its functions

have been taken over by non-homologous proteins. Indeed, there is a clear precedent for non-

homologous reproductive proteins being used to achieve similar phenotypic endpoints in different

insect species (20, 21).

The second is our detection of substantial variation in copy number and sequence in the

Sophophora-Lordiphosa radiation, a feature that was particularly pronounced in the "Oriental"

lineage and *ananassae* subgroup. Outside of the *Sophophora-Lordiphosa, SP* is almost invariably

a single (or 0) copy gene. But inside, where we detect repeated, independent duplication of *SP*

across lineages, the story is quite different. Repeated duplication is at its most extreme in the

ananassae subgroup, where we see as many as 7 copies present in *D. pseudoananassae*

nigrens. Importantly, in this subgroup the intraspecific paralogs are not identical in sequence.

Instead, they generally showed considerable variation in length and amino acid composition.

What, then, are the functional consequences of this intraspecific diversity? If these paralogs are

all interacting with SPR and varying to different degrees in their C-terminal sequences, do they

vary in the efficiency with which they bind SPR? Or are they specialised for different receptors?

We know, for example, that *D. melanogaster* SP can activate another GPCR, Methusaleh (Mth),

in vitro. However, SP-Mth interactions don't seem to be required for the post-mating increase in

egg production or reduction in sexual receptivity – at least in *D. melanogaster* (67). There is also

evidence that SP can induce some of its effects in *D. melanogaster* independently of *SPR,*

pointing to the potential existence of additional, unidentified receptors (68). Thus, it's possible that

SP might be evolving in some lineages to make use of a wider set of receptors, a process that

copy number amplification might facilitate.

But what about the other regions of SP proteins, the regions beyond the post-mating response

stimulating C-terminus? After all, it's the N-terminus and midsection regions that we show to be

most variable between homologs. In *D. melanogaster*, we know that SP binds to sperm at its N-

terminus; the N-terminal WEWPWNR motif then remains bound to SP after the rest of the peptide

is cleaved (33). Our data suggest that clear variants of this motif are restricted to at least one SP

copy in the Muller D2 neighbourhood of each member of the "Oriental" lineage (except for *D.*

oshimai). This raises two possibilities: (1) the ability of SP to bind sperm is restricted to SP copies

carrying variants of this motif in this lineage; (2) the protein network that underlies SP-sperm

binding can facilitate attachment using a wide set of N-terminus sequences. Such flexibility might

stem from rapid evolution of the sequence or identity of SP network proteins, or of any sperm

surface proteins that SP might interact with. Indeed, there is evidence that several sex peptide

network proteins have experienced recurrent positive selection in the *melanogaster* group (49).

Flexible use of N-terminus sequences may also be due to some inherent, accommodating

property of the sperm binding apparatus. Consistent with this, no such WEWPWNR motif is

present in *D. melanogaster* Dup99b, but it nevertheless binds to sperm, albeit only to the sperm

head and only during the first few hours after mating, unlike SP (69). But whether Dup99b is

relying on the same network of proteins as SP to bind to sperm, or perhaps interacting with distinct

proteins on the sperm surface, remains untested. Where multiple, divergent SP copies are

present, as in the *ananassae* subgroup for example, we may be seeing specialisation in the N-

terminus region that relates to a given peptide's mechanism of interacting with the female: while

some may be adapted to bind sperm, others might be adapted to enter into the hemolymph, as

D. melanogaster SP has been shown to do (70).

The third relates to the genomic distribution of *SP* genes. We observed that *SP* genes have

frequently translocated to new genomic locations, a feature that's particularly pervasive in

Sophophora. A key question here is to what extent these translocations have shaped the evolution

of SP function. For example, did the 'Muller D1' to 'Muller D2' translocation at the base of the

Sophophora-Lordiphosa lineage open the door to the novel evolutionary trajectories taken within

this lineage? We could imagine that this translocation placed *SP* within a new *cis*-regulatory

environment that changed either the strength, timing, or tissue-specificity of its expression,

thereby exposing it to new selective forces. Our data suggest against this at a global level, at least

in relation to tissue-specificity, as we observed accessory gland biased expression of *SP* in *D.*

virilis, a species that pre-dates the translocation. However, it's possible that subsequent

translocations within some branches of the *Sophophora-Lordiphosa* radiation are associated with

shifts in expression pattern to new subregions of the male reproductive tract. After all, the

translocated Muller element E copy of *SP* in *D. melanogaster* (*Dup99b*) is expressed not in the

accessory glands, like *SP,* but in the ejaculatory duct (44).

The fourth is our failure to detect a clear association between evolutionary change in *SP* and

microcarriers. The presence of LipidTox$^+$ microcarriers in *S. flava,* which lacks a copy of *SP*,

suggests that *SP* isn't necessary for microcarriers. Moreover, the absence of canonical LipidTox$^+$

microcarriers in *SP*-encoding species, such as *D. immigrans*, suggests that a copy of *SP* isn't

sufficient. Thus, SP playing a role in structuring microcarriers might itself be a derived trait, as

might its association with microcarriers more broadly (*e.g.,* as a microcarrier cargo), thereby

casting doubt on the idea that this could be SP's ancestral function (15). This association might

be relatively recent, as even among many *SP*-bearing *Sophophora* species variation in *SP* copy

number and sequence doesn't seem to be obviously connected to variation in microcarrier

morphology. Our stainings also raise the issue of what exactly defines a 'microcarrier'. The

detection of LipidTox$^-$, ellipsoid, microcarrier-like structures and LipidTox$^-$ flocculence raises the

question of whether these represent fundamentally different structures to microcarriers or whether

taking up LipidTox (an indicator that they contain large quantities of triglycerides and other

nonpolar lipids (28)) is a feature of some, but not all, microcarriers – a feature that our data

suggest appears to have been gained and lost repeatedly, perhaps in line with higher-level dietary

or metabolic changes.

Our data point to lineage-accelerated evolution of *SP* within the *Sophophora-Lordiphosa*

radiation, marked by repeated, independent rounds of gene family expansion. What forces are

driving this trend? And why does it appear to be so much more pervasive in these lineages? A

tempting response to the latter question is that it coincides with some evolutionary shift in the

activity of SP. A previous study of 11 drosophilid species found that the ability of conspecific SP

to reduce female sexual receptivity is confined to the *melanogaster* group (25). Intriguingly, this

gain in responsiveness to SP also appears to coincide with both the gain of robust expression of

SPR within the female reproductive tract and the ability of SP (derived from *D. melanogaster*) to

bind to female reproductive tract tissue (25). Viewed from a sexual conflict perspective, therefore,

we might be detecting the effects of a sexually antagonistic coevolutionary arms race that was

initiated after the acquisition of new functions by SP. If responding to SP is deleterious to female

fitness (*e.g.,* 43), then females might evolve resistance, in turn selecting for structurally divergent

copies of *SP* through which males can overcome that resistance. Indeed, the numerous,

structurally divergent copies of *SP* expressed by members of the *ananassae* subgroup are

consistent with theory suggesting that males might gain from transferring diverse sexually

antagonistic seminal fluid products simultaneously as a 'combination' strategy to overcome the

evolution of resistance (71).

But if females are evolving resistance, then we fail to find strong evidence that it is occurring

through *SPR*. The only region of SPR where we see extensive evolutionary change is in the N-

terminus. But there are reasons to believe that any role this region plays in ligand binding is

relatively minor (see *SI Appendix, Supp. Text*). And while we detect evidence of episodic, but not

pervasive, positive selection at 10 sites in SPR beyond the N-terminus, these changes were

distributed throughout the phylogeny and without enrichment in the lineages experiencing

accelerated evolution of *SP.* Why, then, are we not seeing SPR rapidly evolving in concert with

SP? One explanation is that the evolution of SPR is constrained by the need to maintain

interactions with additional ligands, such as MIPs. But we know from *in vitro* assays that there is

at least one mutational route for SPR that has decoupled effects on MIP and SP binding,

negatively affecting the latter more strongly than the former (51). And yet across 193 drosophilid

SPR sequences we failed to find any instance where this route has been exploited (though it

would require a double mutation). Another possibility is that any resistance to SP-mediated sexual

antagonism might be mediated up- or down-stream of SP-SPR binding, such as via mechanisms

that degrade SP, block it from binding, or in how the neural circuitry that SPR feeds into responds

to SP. One such potential action point is sperm cleavage: if females could block the cleavage of

SP from the surface of the sperm – the mechanism for which remains uncharacterised – then they

could markedly reduce the timeframe over which SP's effects are active (~10-14 days in D.

melanogaster; 30).

But limited evolutionary change in *SPR* is also consistent with another possibility: the role of

sexual conflict in driving the evolution of SP might be relatively weak. While the functions of SP

are often framed in terms of male manipulation of female reproductive decision-making or

collateral damage in the pursuit of improved performance in sperm competition (e.g., 72–75), the

evidence base for this is not strong (reviewed in 15). There is theoretical support for antagonistic

effects of seminal proteins in general (*e.g.,* 83) and empirical support in *D. melanogaster* for the

antagonistic effects of seminal proteins (77) and SP specifically (45, 78), but there is also empirical

support for positive, neutral, and context-dependent (*i.e.,* in relation to female nutritional state)

effects of SP on female fitness (46, 47, 79), as well as considerable uncertainty over the extent

to which fitness measurements made in these laboratory studies reflect those experienced by

wild-living populations (15, 80). Indeed, the recently demonstrated gain of both robust *SPR*

expression in a subset of female reproductive tract neurons and the ability of SP to bind to

reproductive tract tissue in the *melanogaster* group (25) suggests that females did, at least at

some point, benefit from responding to SP, perhaps as part of a mechanism through which the

receipt of sperm could be aligned with the induction of reproductive processes. Whether that may

have subsequently initiated conflict – and what the associated genomic consequences might have

been – remains to be resolved.

The high diversity we detect in SP coupled with the low diversity documented in the receptor

mirrors patterns that have been recorded at the morphological level. While there are instances of

correlated diversification between male-specific traits and female defensive traits that are

suggestive of coevolutionary arms races over mating rate, such as in the clasping and anti-

clasping structure of *Gerris* water striders (81, 82), there are also a great many male-specific traits

across arthropods that show rapid interspecific diversification without apparent change in the

female structures they contact (83). In such cases, the diversification of male traits has

traditionally been thought to be driven by sexual selection under female choice (17). Others,

notably Eberhard and Cordero, have argued that seminal proteins may similarly be evolving under

female choice (16, 18). In terms of SP, this may mean that a female gains from discriminating

between different males on the basis of the dose or sequence of SP she receives, tailoring her

use of sperm or overall reproductive investment in response (15, 84). This, in turn, may select for

diversification in SP sequence or increases in expression (perhaps via copy number amplification)

to meet female preferences (85).

Ultimately, sexual conflict and sexual selection by female choice needn't act entirely

independently (15, 86, 87). But disentangling the historical importance of these forces in driving

the evolution of SP-female interactions continues to present a major challenge, just as it has for

many other traits since work in this area began (83, 84). Our belief is that by charting the origin

and diversification of SP-female interactions at a mechanistic level, we may be able to draw more

robust inferences regarding the evolutionary forces that are shaping this recently evolved system

in *Drosophilinae*.

Methods

Detailed sample information and accession numbers for the 271 genomes used in this study are

provided at https://osf.io/24unk. SP sequences were identified by combining reciprocal blast of

the *D. melanogaster* SP C-terminus sequence with protein sequence, gene structure, and synteny

analysis. Signal peptides were removed from the translated protein sequences using SignalP-6.0

run on 'fast' mode and specifying 'Eukarya' as the organism (88). For the microcarrier stainings,

accessory glands were stained in 100µl of 1:50 LipidTox Deep Red Neutral Lipid Stain (Invitrogen,

H34477) in PBS for 1 hour with 1µl 1:100 DAPI added for the last 15 mins. Tests for selection in

SPR coding sequence were conducted using Datamonkey (89, 90). All code, extracted coding

and protein sequences, expression data, and protein models are available at https://osf.io/tzu6v/.

A detailed methods section is available in the *SI Appendix*.

Acknowledgements

We thank the reviewers and editor for thoughtful and valuable feedback on earlier drafts of this

manuscript; Clive Wilson and Mark Wainwright for helpful feedback on the microcarrier part of this

work; Julianne Peláez and Noah Whiteman for providing us with *Scaptomyza* stocks; and Barbara

Parker, Kathy Darragh, Pavitra Muralidhar, and Carl Veller for valuable discussion. This work was

supported by a long-term fellowship from the Human Frontier Science Program Organization

(LT000123/2020-L to B.R.H.), a UC Davis College of Biological Sciences Dean's Circle Summer

Undergraduate Research Program Fellowship to A. A-H., and National Institutes of Health F32

GM135998 to B.Y.K., R37-HD038921 to Mariana Wolfner, and R35 GM122592 to A.K.

Figure legends:

Figure 1. *SP* first evolved in the *Drosophilinae* subfamily. A phylogeny of species from across

the *Drosophilidae* and closely related lineages. The number of *SP* genes detected is given in

coloured circles at the tip of each branch. A brown bar denotes the branch on which we infer *SP*

to have first evolved.

Figure 2. *Sex Peptide* family genes predate the *Drosophilini* and have been repeatedly lost outside of the *Sophophora-Lordiphosa* radiation. This figure focuses on the non-*Sophophora-Lordiphosa* members of the *Drosophilini* (see Figure 1 and *SI Appendix, Fig. S1* for overviews of drosophilid taxonomy). A selection of *Sophophora* species, shaded in blue, are included for comparison. Also included are four non-*Drosophilini* dipterans: *Aedes aegypti* and three non-*Drosophilini* members of the *Drosophilinae* subfamily: *Chymomyza costata, Scaptodrosophila lebanonensis,* and *S. latifasciaeformis.* The number of *SP* genes detected in a representative of each species' genome is given at the tree tips. Losses are marked with a pink bar. For each species, the structures of two syntenic gene neighbourhoods are plotted. The first, Muller D1, is the canonical neighbourhood in which *SP* genes are detected outside of the *Sophophora-Lordiphosa.* The second, Muller D2, is the canonical position in the *Sophophora-Lordiphosa.* Positions of each gene are given relative to the first gene in the neighbourhood (*FoxK* or *capricious*). Absence of a flanking neighbourhood gene (*e.g., FoxK* in *D. wassermani*) doesn't necessarily mean the gene has been lost – it more likely means that a contig breakpoint fell within the neighbourhood. Note that *SP* in *Hirtodrosophila trivittata* and, independently, *D. repletoides,* has translocated out of the Muller D1 neighbourhood. See *SI Appendix, Figs S2 and 3* for expanded views of the losses in the *repleta* and *Hawaiian* groups, respectively.

Figure 3. *Sex Peptide* copy number is markedly more variable in the *Sophophora-Lordiphosa* radiation than in other branches of the phylogeny. This figure focuses on the *Sophophora-Lordiphosa* radiation to which *D. melanogaster* belongs. Four non-*Sophophora-Lordiphosa* drosophilids, shaded in purple, are included as outgroups. The structures of the Muller D1 and Muller D2 neighbourhoods are plotted as in Figure 2. Missing flanking genes are likely indicative of contig breakpoints falling within the neighbourhood. The exceptions are the *Lordiphosa* species, where substantially elevated intergenic distances meant that the whole neighbourhood would not fit within the plot limits. Unresolved SP genes, shown in red, indicate

genes that passed the reciprocal blast criteria and fell within one of the conserved *SP*-containing

gene neighbourhoods but where a SP-like amino acid sequence couldn't be resolved (*e.g.,* due

to a premature stop codon, as in the case of *D. rhopaloa*). Note that all members of the *obscura*

group have an inversion that flips the relative positions of *FoxK* and *mRpL2* in the Muller D1

neighbourhood. In a number of cases, some or all copies of *SP* were found to have translocated

outside of the Muller D1 and Muller D2 neighbourhoods (an *obscura* group lineage, the

melanogaster subgroup, *D. kanapiae, D. takahashii*, and *D.eugracilis*; summarised in *SI Appendix*

Fig. S5). In the shorter read *montium* subgroup assemblies, short contigs meant that in some

species we couldn't identify the neighbourhood in which *SP* was located. This was the case for

some *SP* genes in *D. cf. chauvacae, D. burlai, D. leontia, D. nikananu, D. pectinifera, D.*

punjabiensis, and *D.watanabe*i. The *SP* genes in *D. auraria* and *D. trapezifons* could be mapped

to the Muller D2 neighbourhood based on flanking sequence around the *SP* gene, but the *SP*-

containing contigs were too small to include any of the neighbourhood genes.

Figure 4. Repeated duplication of *Sex Peptide* genes in the *ananassae* subgroup. A

phylogeny of the *ananassae* subgroup species used in this study, with *D. melanogaster* as an

outgroup. The number of *SP* genes identified in each species is given at the tip of each branch.

'Unresolved' SP sequences, shown in red, are those which passed the reciprocal blast tests and

fell within the syntenic Muller D2 gene neighbourhood but for which we could not resolve an SP-

like protein sequence (*e.g.,* due to a premature stop codon). The structure of the neighbourhood

is plotted on the right-hand side of the figure. Note that one of *D. melanogaster's SP* copies,

Dup99b, falls outside of the Muller D2 neighbourhood.

Figure 5. *Sex Peptide* is neither necessary nor sufficient for microcarriers. (A) A phylogeny

of all species included in this figure. Branches coloured pink indicate species shown in B-T that

demonstrate canonical (*i.e., D. melanogaster*-like) staining with LipidTox, a neutral lipid-specific

731 dye used to selectively stain microcarriers (28). (B-T) Accessory glands stained with LipidTox and

732 the nuclear stain DAPI (blue). The circled number in the bottom right-hand corner of each panel

733 indicates the number of *Sex Peptide* copies we detect in each species. (G) The arrow is

734 highlighting a spiral/doughnut shaped microcarrier, a shape which is rare in comparison to the

735 more common thread-like conformation in this species. (J) Prox. refers to the proximal region of

736 the gland, *i.e.* the region that connects to the ejaculatory duct. (P) Arrows point to the ambiguous,

737 sparse, and weakly stained material we observed in *D. ficusphila* glands. (S) An arrow points to

738 a cut in the glandular epithelium, which was made to enhance dye penetration. In each case,

739 glands were co-stained in the same well with those from *D. melanogaster* to act as a positive

740 control.

742 **Figure 6. SPR residues showing evidence of episodic positive selection are enriched in**

743 **the ligand-facing domains.** (A) A consensus sequence based on MAFFT alignment of the

744 resolvable amino acid sequences of *SPR* coding sequences. The top plot gives the proportion of

745 sequences with the consensus amino acid in the same position, while the bottom plot gives the

746 proportion of sequences in which each position is occupied in the alignment. Each residue is

747 coloured based on the functional domain to which it belongs based on the UniProt annotations.

748 The red line indicates the start of the conserved region we use in the molecular evolution

749 analyses. Beneath the plot, we show the corresponding exon that encodes each consensus

750 residue. EC = extracellular, IC = intracellular (B) The AlphaFold prediction of the structure of *D.*

751 *melanogaster* SPR as downloaded from UniProt (AF-Q8SWR3-F1) and coloured by the domain

752 each residue belongs to based on positions listed in the UniProt 'Features' table. (C) A boxplot

753 showing the proportion of residues at each position that matched the consensus residue (*i.e.,* the

754 degree of conservation at each position). Residues are plotted separately according to their

755 domain class of origin. (D) PCA plot based on BLOSUM62 substitution scores from the MAFFT-

756 aligned SPR protein sequences. The percentage values in the axis titles reflect the proportion of

variance explained by a given PC. Points are coloured based on whether they correspond to

Sophophora-Lordiphosa, non-*Sophophora-Lordiphosa,* or *ananassae* subgroup species. (E) As

(D) but after removing the non-conserved region of the N-terminus (*i.e.,* the region preceding the

red line in (A)). (F) The same prediction shown in (B) but with residues coloured by the proportion

of conserved residues per position (see also Supplementary movie 1). High values indicate high

conservation. (G) The ColabFold top-ranked prediction of the interactions between SP (shown in

green) and SPR (residues coloured by domain). (H) As in (G) but with predicted contact residues

coloured purple and predicted hydrogen bonds between SP and SPR residues shown with red-

dotted lines (See also Supplementary movies 2 and 3). (I) A bar chart showing the proportion of

sequences sharing the consensus residue for each predicted contact residue. Bars are coloured

by whether the residue is also predicted to form a hydrogen bond with SP. Asterisks denote

predicted contact residues for which we detected evidence of episodic positive selection using

MEME. Alongside the plot, coloured bars, using the same colour scale as in (A-C), denote the

functional domain the residue falls within. The two adjacent blue bars denote separate,

consecutive transmembrane domains. Alongside are the corresponding amino acid residues in

each position for each of the consensus ('Cons.'), *D. melanogaster* ('Dmel.'), *D. bipectinata*

('Dbip.')*, D. pseudoobscura* ('Dpse.')*, D. virilis* ('Dvir.'), and *A. aegypti* ('Aaeg.') sequences.

Residues that depart from the mode among these plotted sequences are coloured. (J) As (G) but

colouring only the 10 residues in SPR for which we detected evidence of positive selection using

MEME. Asterisks denote predicted contact residues that show evidence of positive selection.

References:

1.   R. A. Schwenke, B. P. Lazzaro, Juvenile Hormone Suppresses Resistance to Infection in

Mated Female Drosophila melanogaster. *Curr. Biol.* **27**, 596–601 (2017).

2.   B. Abu-Raya, C. Michalski, M. Sadarangani, P. M. Lavoie, Maternal Immunological

Adaptation During Normal Pregnancy. *Front. Immunol.* **11**, 2627 (2020).

3.   M. A. Berland, *et al.*, Seminal Plasma Induces Ovulation in Llamas in the Absence of a

Copulatory Stimulus: Role of Nerve Growth Factor as an Ovulation-Inducing Factor.

Endocrinology **157**, 3224–3232 (2016).

4.   Y. Tsukamoto, H. Kataoka, H. Nagasawa, S. Nagata, Mating changes the female dietary

preference in the two-spotted cricket, Gryllus bimaculatus. *Front. Physiol.* **5 MAR**, 1–6

(2014).

5.   S. J. Walker, V. M. Corrales-Carvajal, C. Ribeiro, Postmating Circuitry Modulates Salt

Taste Processing to Increase Reproductive Output in Drosophila. *Curr. Biol.* **25**, 2621–

2630 (2015).

6.   S. Wigby, *et al.*, Seminal fluid protein allocation and male reproductive success. *Curr.*

Biol. **19**, 751–757 (2009).

7.   C. Fricke, S. Wigby, R. Hobbs, T. Chapman, The benefits of male ejaculate sex peptide

transfer in Drosophila melanogaster. *J. Evol. Biol.* **22**, 275–286 (2009).

8.   B. R. Hopkins, *et al.*, Divergent allocation of sperm and the seminal proteome along a

competition gradient in Drosophila melanogaster. *Proc. Natl. Acad. Sci. U. S. A.* **116**,

17925–17933 (2019).

9.   J. T. Patterson, A New Type of Isolating Mechanism in Drosophila. *Proc. Natl. Acad. Sci.*

U. S. A. **32**, 202–208 (1946).

10.   Y. H. Ahmed-Braimah, M. F. Wolfner, A. G. Clark, Differences in Postmating

Transcriptional Responses between Conspecific and Heterospecific Matings in

Drosophila. *Mol. Biol. Evol.* **38**, 986–999 (2021).

11.   A. Civetta, A. G. Clark, Correlated effects of sperm competition and postmating female

mortality. *Proc. Natl. Acad. Sci.* **97**, 13162–13165 (2000).

12.   W. R. Rice, Sexually antagonistic male adaptation triggered by experimental arrest of

female evolution. *Nature* **381**, 232–234 (1996).

13. B. Holland, W. R. Rice, Perspective: Chase-Away Sexual Selection: Antagonistic

Seduction Versus Resistance. *Evolution (N. Y).* **52**, 1 (1998).

14. B. Hollis, *et al.*, Sexual conflict drives male manipulation of female postmating responses

in Drosophila melanogaster. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 8437–8444 (2019).

15. B. R. Hopkins, J. C. Perry, The evolution of sex peptide: sexual conflict, cooperation, and

coevolution. *Biol. Rev.* **97**, 1426–1448 (2022).

16. W. G. Eberhard, C. Cordero, Sexual selection by cryptic female choice on male seminal

products - a new bridge between sexual selection and reproductive physiology. *Trends*

Ecol. Evol. **10**, 493–496 (1995).

17. W. G. Eberhard, Postcopulatory sexual selection: Darwin's omission and its

consequences. *Proc. Natl. Acad. Sci.* **106**, 10025–10032 (2009).

18. C. Cordero, Ejaculate substances that affect female insect reproductive physiology and

behavior: Honest or arbitrary traits? *J. Theor. Biol.* **174**, 453–461 (1995).

19. T. Pizzari, R. R. Snook, Perspective: Sexual conflict and sexual selection: Chasing away

paradigm shifts. *Evolution (N. Y).* **57**, 1223–1236 (2003).

20. T. G. Kingan, W. M. Bodnar, A. K. Raina, J. Shabanowitz, D. F. Hunt, The loss of female

sex pheromone after mating in the corn earworm moth Helicoverpa zea: identification of a

male pheromonostatic peptide. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 5082 (1995).

21. L. B. Duvall, N. S. Basrur, H. Molina, C. J. McMeniman, L. B. Vosshall, A Peptide

Signaling System that Rapidly Enforces Paternity in the Aedes aegypti Mosquito. *Curr.*

Biol. **27**, 3734-3742.e5 (2017).

22. P. S. Chen, *et al.*, A male accessory gland peptide that regulates reproductive behavior of

female D. melanogaster. *Cell* **54**, 291–298 (1988).

23. P. S. Chen, C. Diem, A sex-specific ninhydrin-positive substance found in the paragonia

of adult males of Drosophila melanogaster. *J. Insect Physiol.* **7**, 289–298 (1961).

24.   A. S. Fox, Chromatographic Differences between Males and Females in Drosophila

melanogaster and Role of X and Y Chromosomes. *Physiol. Zool.* **29**, 288–298 (1956).

25.   M. Tsuda, J. B. Peyre, T. Asano, T. Aigaki, Visualizing molecular functions and cross-

species activity of sex-peptide in Drosophila. *Genetics* **200**, 1161–1169 (2015).

26.   A. C. Majane, J. M. Cridland, D. J. Begun, Single-nucleus transcriptomes reveal

evolutionary and functional properties of cell types in the Drosophila accessory gland.

Genetics **220** (2022).

27.   D. Styger, "Molekulare Analyse des Sexpeptidgens aus Drosophila melanogaster,"

University of Zurich, Zurich, Switzerland. (1992).

28.   S. M. Wainwright, *et al.*, Drosophila Sex Peptide controls the assembly of lipid

microcarriers in seminal fluid. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2019622118 (2021).

29.   K. Ravi Ram, M. F. Wolfner, A network of interactions among seminal proteins underlies

the long-term postmating response in Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* **106**,

15384–15389 (2009).

30.   G. D. Findlay, *et al.*, Evolutionary rate covariation identifies new members of a protein

network required for Drosophila melanogaster female post-mating responses. *PLoS*

Genet. **10**, e1004108 (2014).

31.   D. Gligorov, J. L. Sitnik, R. K. Maeda, M. F. Wolfner, F. Karch, A Novel Function for the

Hox Gene Abd-B in the Male Accessory Gland Regulates the Long-Term Female Post-

Mating Response in Drosophila. *PLoS Genet.* **9**, e1003395 (2013).

32.   A. Singh, *et al.*, Long-term interaction between Drosophila sperm and sex peptide is

mediated by other seminal proteins that bind only transiently to sperm. *Insect Biochem.*

Mol. Biol. **102**, 43–51 (2018).

33.   J. Peng, *et al.*, Gradual Release of Sperm Bound Sex-Peptide Controls Female

Postmating Behavior in Drosophila. *Curr. Biol.* **15**, 207–213 (2005).

34.   M. Häsemeyer, N. Yapici, U. Heberlein, B. J. Dickson, Sensory Neurons in the Drosophila

Genital Tract Regulate Female Reproductive Behavior. *Neuron* **61**, 511–518 (2009).

35. C. Rezával, *et al.*, Neural circuitry underlying Drosophila female postmating behavioral

responses. *Curr. Biol.* **22**, 1155–1165 (2012).

36. N. Yapici, Y.-J. Kim, C. Ribeiro, B. J. Dickson, A receptor that mediates the post-mating

switch in Drosophila reproductive behaviour. *Nature* **451**, 33–37 (2008).

37. P. Moshitzky, *et al.*, Sex-peptide activates juvenile hormone biosynthesis in

theDrosophila melanogaster corpus allatum. *Arch. Insect Biochem. Physiol.* **32**, 363–374

(1996).

38. Y. Fan, *et al.*, Common functional elements of Drosophila melanogaster seminal peptides

involved in reproduction of Drosophila melanogaster and Helicoverpa armigera females.

Insect Biochem. Mol. Biol. **30**, 805–812 (2000).

39. E. V. Domanitskaya, H. Liu, S. Chen, E. Kubli, The hydroxyproline motif of male sex

peptide elicits the innate immune response in Drosophila females. *FEBS J.* **274**, 5659–

5668 (2007).

40. K. Moehle, A. Freund, E. Kubli, J. A. Robinson, NMR studies of the solution conformation

of the sex peptide from Drosophila melanogaster. *FEBS Lett.* **585**, 1197–1202 (2011).

41. Z. Ding, I. Haussmann, M. Ottiger, E. Kubli, Sex-peptides bind to two molecularly different

targets inDrosophila melanogaster females. *J. Neurobiol.* **55**, 372–384 (2003).

42. T. Schmidt, Y. Choffat, S. Klauser, E. Kubli, The Drosophila melanogaster sex-peptide: A

molecular analysis of structure-function relationships. *J. Insect Physiol.* **39**, 361–368

(1993).

43. P. Saudan, *et al.*, Ductus ejaculatorius peptide 99B (DUP99B), a novel Drosophila

melanogaster sex-peptide pheromone. *Eur. J. Biochem.* **269**, 989–997 (2002).

44. A. Rexhepaj, H. Liu, J. Peng, Y. Choffat, E. Kubli, The sex-peptide DUP99B is expressed

in the male ejaculatory duct and in the cardia of both sexes. *Eur. J. Biochem.* **270**, 4306–

4314 (2003).

45.   S. Wigby, T. Chapman, Sex peptide causes mating costs in female Drosophila

888   melanogaster. *Curr. Biol.* **15**, 316–321 (2005).

889   46.   C. Fricke, A. Bretman, T. Chapman, Female nutritional status determines the magnitude

890   and sign of responses to a male ejaculate signal in Drosophila melanogaster. *J. Evol.*

891   *Biol.* **23**, 157–165 (2010).

892   47.   K. U. Wensing, C. Fricke, Divergence in sex peptide-mediated female post-mating

893   responses in Drosophila melanogaster. *Proc. R. Soc. B Biol. Sci.* **285**, 20181563 (2018).

894   48.   Y.-J. Kim, *et al.*, MIPs are ancestral ligands for the sex peptide receptor. *Proc. Natl. Acad.*

895   *Sci.* **107**, 6520–6525 (2010).

896   49.   M. K. McGeary, G. D. Findlay, Molecular evolution of the sex peptide network in

897   Drosophila. *J. Evol. Biol.* **33**, 629–641 (2020).

898   50.   M. Conzelmann, *et al.*, Conserved MIP receptor-ligand pair regulates Platynereis larval

899   settlement. *Proc. Natl. Acad. Sci.* **110**, 8224–8229 (2013).

900   51.   J. Poels, *et al.*, Myoinhibiting peptides are the ancestral ligands of the promiscuous

901   Drosophila sex peptide receptor. *Cell. Mol. Life Sci.* **67**, 3511–3522 (2010).

902   52.   Y. Oh, *et al.*, A Homeostatic Sleep-Stabilizing Pathway in Drosophila Composed of the

903   Sex Peptide Receptor and Its Ligand, the Myoinhibitory Peptide. *PLoS Biol.* **12**,

904   e1001974 (2014).

905   53.   J. G. Scott, *et al.*, Genome of the house fly, Musca domestica L., a global vector of

906   diseases with adaptations to a septic environment. *Genome Biol.* **15**, 466 (2014).

907   54.   B. Vicoso, D. Bachtrog, Numerous Transitions of Sex Chromosomes in Diptera. *PLoS*

908   *Biol.* **13**, e1002078 (2015).

909   55.   C. Finet, *et al.*, DrosoPhyla: Resources for Drosophilid Phylogeny and Systematics.

910   *Genome Biol. Evol.* **13** (2021).

911   56.   A. Yassin, Phylogenetic classification of the Drosophilidae Rondani (Diptera): The role of

912   morphology in the postgenomic era. *Syst. Entomol.* **38**, 349–364 (2013).

913   57.   S. Cirera, M. Aguadé, The sex-peptide gene (Acp70A) is duplicated in Drosophila

914   subobscura. *Gene* **210**, 247–254 (1998).

915   58.   M. D. Smith, *et al.*, Less Is More: An Adaptive Branch-Site Random Effects Model for

916   Efficient Detection of Episodic Diversifying Selection. *Mol. Biol. Evol.* **32**, 1342–1353

917   (2015).

918   59.   B. Murrell, *et al.*, Gene-Wide Identification of Episodic Selection. *Mol. Biol. Evol.* **32**,

919   1365–1371 (2015).

920   60.   B. Murrell, *et al.*, FUBAR: A Fast, Unconstrained Bayesian AppRoximation for Inferring

921   Selection. *Mol. Biol. Evol.* **30**, 1196–1205 (2013).

922   61.   B. Murrell, *et al.*, Detecting Individual Sites Subject to Episodic Diversifying Selection.

923   *PLOS Genet.* **8**, e1002764 (2012).

924   62.   M. Mirdita, *et al.*, ColabFold: making protein folding accessible to all. *Nat. Methods* **19**,

925   679–682 (2022).

926   63.   M. Mirdita, *et al.*, Uniclust databases of clustered and deeply annotated protein

927   sequences and alignments. *Nucleic Acids Res.* **45**, D170–D176 (2017).

928   64.   M. Mirdita, M. Steinegger, J. Söding, MMseqs2 desktop and local web server app for fast,

929   interactive sequence searches. *Bioinformatics* **35**, 2856–2858 (2019).

930   65.   A. L. Mitchell, *et al.*, MGnify: the microbiome analysis resource in 2020. *Nucleic Acids*

931   *Res.* (2019) https:/doi.org/10.1093/nar/gkz1035.

932   66.   E. F. Pettersen, *et al.*, UCSF ChimeraX: Structure visualization for researchers,

933   educators, and developers. *Protein Sci.* **30**, 70–82 (2021).

934   67.   W. W. Ja, G. B. Carvalho, M. Madrigal, R. W. Roberts, S. Benzer, The Drosophila G

935   protein-coupled receptor, Methuselah, exhibits a promiscuous response to peptides.

936   *Protein Sci.* **18**, 2203–2208 (2009).

937   68.   I. U. Haussmann, Y. Hemani, T. Wijesekera, B. Dauwalder, M. Soller, Multiple pathways

938   mediate the sexpeptide- regulated switch in female Drosophila reproductive behaviours.

939   *Proc. R. Soc. B Biol. Sci.* **280**, 20131938 (2013).

940   69.   J. Peng, P. Zipperlen, E. Kubli, Drosophila sex-peptide stimulates female innate immune

941   system after mating via the toll and Imd pathways. *Curr. Biol.* **15**, 1690–1694 (2005).

942   70.   N. Pilpel, I. Nezer, S. W. Applebaum, Y. Heifetz, Mating-increases trypsin in female

943   Drosophila hemolymph. *Insect Biochem. Mol. Biol.* **38**, 320–330 (2008).

944   71.   T. Chapman, Sexual Conflict: Mechanisms and Emerging Themes in Resistance Biology.

945   *Am. Nat.* **192**, 217–229 (2018).

946   72.   R. A. Slatyer, B. S. Mautz, P. R. Y. Backwell, M. D. Jennions, Estimating genetic benefits

947   of polyandry from experimental studies: A meta-analysis. *Biol. Rev.* **87**, 1–33 (2012).

948   73.   M. Pröschel, Z. Zhang, J. Parsch, Widespread Adaptive Evolution of Drosophila Genes

949   With Sex-Biased Expression. *Genetics* **174**, 893–900 (2006).

950   74.   L. F. Bussiégre, J. Hunt, M. D. Jennions, R. Brooks, Sexual Conflict and Cryptic Female

951   Choice in the Black Field Cricket, Teleogryllus Commodus. *Evolution (N. Y).* **60**, 792–800

952   (2006).

953   75.   T. A. R. Price, Z. Lewis, D. T. Smith, G. D. D. Hurst, N. Wedell, Sex ratio drive promotes

954   sexual conflict and sexual coevolution in the fly drosophila pseudoobscura. *Evolution (N.*

955   *Y).* **64**, 1504–1509 (2010).

956   76.   R. A. Johnstone, L. Keller, How males can gain by harming their mates: Sexual conflict,

957   seminal toxins, and the cost of mating. *Am. Nat.* **156**, 368–377 (2000).

958   77.   T. Chapman, L. F. Liddle, J. M. Kalb, M. F. Wolfner, L. Partridge, Cost of mating in

959   Drosophila melanogaster females is mediated by male accessory gland products. *Nature*

960   **373**, 241–244 (1995).

961   78.   J. Tower, *et al.*, Mifepristone/RU486 acts in Drosophila melanogaster females to

962   counteract the life span-shortening and pro-inflammatory effects of male Sex Peptide.

963   *Biogerontology* **18**, 413–427 (2017).

79.   B. Rogina, The effect of sex peptide and calorie intake on fecundity in female Drosophila

melanogaster. *Sci. World J.* **9**, 1178–1189 (2009).

80.   W. G. Eberhard, *et al.*, Sexual conflict and female choice [1] (multiple letters). *Trends*

Ecol. Evol. **18**, 438–439 (2003).

81.   G. Arnqvist, L. Rowe, Antagonistic coevolution between the sexes in a group of insects.

Nature **415**, 787–789 (2002).

82.   G. Arnqvist, L. Rowe, Correlated evolution of male and female morphologies in water

striders. *Evolution (N. Y).* **56**, 936–947 (2002).

83.   W. G. Eberhard, Rapid Divergent Evolution of Sexual Morphology: Comparative Tests of

Antagonistic Coevolution and Traditional Female Choice. *Evolution (N. Y).* **58**, 1947

(2004).

84.   W. G. Eberhard, Cryptic female choice and other types of post-copulatory sexual

selection. *Cryptic Female Choice Arthropods Patterns, Mech. Prospect.*, 1–26 (2015).

85.   L. K. Sirot, M. F. Wolfner, "Who's zooming who? seminal fluids and cryptic female choice

in diptera" in *Cryptic Female Choice in Arthropods: Patterns, Mechanisms and Prospects*,

(Springer International Publishing, 2015), pp. 351–384.

86.   C. Cordero, W. G. Eberhard, Interaction between sexually antagonistic selection and

mate choice in the evolution of female responses to male traits. *Evol. Ecol.* **19**, 111–122

(2005).

87.   H. Kokko, M. D. Jennions, The Relationship Between Sexual Selection and Sexual

Conflict. *Cold Spring Harb. Perspect. Biol.*, 1–13 (2014).

88.   F. Teufel, *et al.*, SignalP 6.0 predicts all five types of signal peptides using protein

language models. *Nat. Biotechnol. 2022 407* **40**, 1023–1025 (2022).

89.   W. Delport, A. F. Y. Poon, S. D. W. Frost, S. L. Kosakovsky Pond, Datamonkey 2010: a

suite of phylogenetic analysis tools for evolutionary biology. *Bioinformatics* **26**, 2455–

2457 (2010).

90.   S. Weaver, *et al.*, Datamonkey 2.0: A Modern Web Application for Characterizing

Selective and Other Evolutionary Processes. *Mol. Biol. Evol.* **35**, 773–777 (2018).