Time is of the essence: A general framework for uncovering temporal structures of communities

Hannah Yin & Volker H. W. Rudolf*

Program of Ecology & Evolutionary Biology, BioSciences, Rice University,

Houston, TX, USA

*corresponding author: volker.rudolf@rice.edu

Running title: The temporal structure of communities

Keywords: Biodiversity, time series, temporal ecology, phenological shifts, community metrics, networks

Article type: Perspective

Word count: Abstract: 185; Main text: 5836 Text box 1: 719, Textbox 2: 521

No. References: 116

Figs: 3, **Text Boxes**: 2, **Table**: 2

Statement of authorship: VHWR conceived the general idea. VHWR developed the framework and analogies together with HY. VHWR and HY wrote the manuscript, and HY collected and analyzed the data.

Data accessibility: All code and date are freely available at

https://figshare.com/s/38f6820937b3118ac08f (we will provide public DOI upon acceptance)

Abstract

Ecological communities are inherently dynamic: species constantly turn over within years,

months, weeks, or even days. These temporal shifts in community composition determine

essential aspects of species interactions and how energy, nutrients, information, diseases, and

perturbations "flow" through systems. Yet, our understanding of community structure has relied

heavily on static analyses not designed to capture critical features of this dynamic temporal

dimension of communities. Here, we propose a conceptual and methodological framework for

quantifying and analyzing this temporal dimension. Conceptually, we split the temporal structure

into two definitive features, sequence and duration, and review how they are linked to key

concepts in ecology. We then outline how we can capture these definitive features using

perspectives and tools from temporal graph theory. We demonstrate how we can easily integrate

ongoing research on phenology into this framework and highlight what new opportunities arise

from this approach to answer fundamental questions in community ecology. As climate change

reshuffles ecological communities worldwide, quantifying the temporal organization of

communities is imperative to resolve the fundamental processes that shape natural ecosystems

and predict how these systems may change in the future.

The temporal dimension of community structure

Communities are organized across two fundamental dimensions: time and space. Research on

spatial patterns has increased rapidly in the past decades and transformed how we think about

natural communities by emphasizing the importance of spatial organization (structure) of species

(Brown 1995; Kareiva 1994; Leibold & Chase 2017; Tilman & Kareiva 2018). In comparison,

we still know surprisingly little about how communities are organized in time, which has only

recently gained renewed attention as a line of research (Post 2019; Wolkovich *et al.* 2014; Yang

2020). The ongoing and projected impacts of climate change on ecological communities, as well

as calls for their short- and long-term mitigation and adaptation plans (IPCC 2023), create an

urgency to develop frameworks explicitly focusing on the temporal dimension of community

structure to fill this conceptual gap (CaraDonna *et al.* 2021; Rudolf 2019; Takimoto & Sato

2020; Wolkovich *et al.* 2014; Yang 2020; Yang & Rudolf 2010).

Ecologists have long recognized that the temporal patterns and shifts in community

composition are a hallmark of any natural community (Elton 1927). Their importance manifests

in several critical ecological concepts, like the temporal niche (Carothers & Jaksić 1984; Jensen

et al. 2019; Loreau 1989a; Schoener 1974), succession (Connell & Slatyer 1977; HA 1927;

Poorter *et al.* 2023), the window of opportunity (Yang *et al.* 2008; Yang & Cenzer 2020), and

the seasonal timing of life history events (phenology) (Cohen *et al.* 2018; Miller-Rushing *et al.*

2010; Parmesan & Yohe 2003; Post 2019; Roslin *et al.* 2021). Yet, despite the widespread

appreciation of the dynamic nature of community structure, community ecology has traditionally

taken a largely static approach. For instance, with few notable exceptions (e.g., Kantsa *et al.*

2018; Ushio *et al.* 2018), the structure of ecological networks is typically assumed to be fixed

and time-invariant such that only the abundance of species may change over time (CaraDonna *et*

al. 2021; Ings *et al.* 2009; Montoya *et al.* 2006). When analyzing empirical patterns, studies of

community structure predominantly use a single data "snapshot" in time or aggregate time points

to calculate key biodiversity metrics, thereby potentially overlooking transient states and critical

ecological processes present at the community level, such as seasonality (McMeans *et al.* 2015).

Consequently, our current understanding of the temporal structure of natural communities

remains limited concerning its variations across ecosystems, space, and time and the factors that

drive such variations. Yet, answering these and many related questions about the temporal

structure of communities is fundamental to our ability to understand and predict the dynamics of

natural communities and how they will respond to environmental change (CaraDonna *et al.*

2021; Rudolf 2019; Wolkovich *et al.* 2014; Yang 2020).

Given that ecologists have long appreciated the dynamic structure of communities, what has

prevented us from answering these questions already? In large part, progress has been hampered

by the lack of a unified conceptual and methodological framework for analyzing the temporal

patterns and the logistical challenges in obtaining the fine-scaled temporal data required, which

can require a lot of resources or new technologies. To move toward filling these gaps, we

propose a general framework to quantify and analyze temporal structure in natural communities.

First, we define the concept of temporal community structure and discuss its potential role in

both well-studied and understudied ecological concepts. We then briefly introduce temporal-

ordered networks and suggest how and why these networks can theoretically capture the

definitive features of temporal structure in natural communities by applying perspectives from

temporal graph theory. We end by applying this framework to publicly available phenological

data and highlight some promising research directions and outstanding questions that this

approach helps address.

What is the temporal structure of communities?

Just like "spatial structure" generally denotes how species are organized in space,

"temporal structure" refers to the organization of species in time. However, unlike space, time is

unidirectional: it always moves forward, never stops, and we can't go back. This directionality

means that events and processes from the past can affect events and processes that happen in the

future, but not the other way around. Temporal structure is thus fundamentally about the

sequence (order) and *duration* of and between events (Wolkovich *et al.* 2014). Take a piece of

music, for example. Knowing all the notes played in a melody does not tell you how it sounds.

What makes any melody unique is the sequence of notes (relative timing), how long each note is,

and the pauses between them. Similarly, knowing what species exist in a community (aggregated

over time) doesn't reveal much about how the community looks at any given point in time or the

"pace" or "rhythm" of key ecological processes and change. Instead, if we are to truly understand

the ecological and evolutionary processes that shape the dynamics of natural communities and

how communities will respond to environmental change, we need to study their temporal

structure: we need to analyze the duration and sequence of when certain groups (e.g., species) are

active and their intervening times (gaps).

What can we learn from the temporal structure of communities?

The presence of species along the temporal axis is not random but rather an emergent property

that reflects its ecological and evolutionary responses to constantly changing environmental

conditions and interactions. Therefore, the temporal positions of species can tell us a lot about

the ecology of species and the processes that structure natural communities. For example,

specific patterns of temporal spacing between species presences could provide insights into prior

and ongoing competition and coexistence mechanisms (Anten & Hirose 1999; Blackford *et al.*

2020; Jensen *et al.* 2019; Schamp & Jensen 2019; Tan *et al.* 2013), or predict the invasion

success of species (Alexander & Levine 2019; Godoy & Levine 2014; Wolkovich & Cleland

2011; Zettlemoyer *et al.* 2019). Anomalies in temporal spacing may reveal when environmental

conditions create windows of opportunities or constraints (Yang 2020; Yang & Cenzer 2020;

Yang & Rudolf 2010) and indicate facilitative interactions or co-dependence (Dante *et al.* 2013).

The distribution of species in time also structures species interactions. For instance, many

interactions are only possible when species stay synchronized enough to overlap in time; a bee

can only pollinate a flower and a predator can only consume a prey when both are present

simultaneously (see **Box 1**). Temporal co-occurrence thus indicates, at minimum, what

interactions are possible and is a crucial driver of the structure of ecological networks

(CaraDonna *et al.* 2021; Morente-López *et al.* 2018; Yang *et al.* 2013). Furthermore, the

sequence of species arrivals in a community and the time interval between arrivals typically

determines the per-capita strength and outcome of interactions (see **Box 1**, **Fig. 1**) (Rudolf 2019).

Finally, how species are arranged along the temporal axis also determines the causal chain of

possible events (like the potential for indirect interactions between species, **Box 1**, **Fig. 1**) and

limits the pathways through which perturbations, infectious diseases, energy, nutrients, and

information can spread through an ecosystem (Blonder *et al.* 2012). Thus, temporal structure

simultaneously acts as an integrative response variable for communities and constrains possible

ecological outcomes.

Analyzing and comparing temporal structure across taxa, systems, environments, space, and

timescales helps us identify and understand many biological processes that shape natural

communities (**Table 1**). This understanding allows for more testable predictions on how

communities respond to environmental change and unlocks more well-rounded and nuanced

perspectives on the roles both space and time play in communities (Wolkovich *et al.* 2014; Yang

2020). However, because the past can shape the future, a mechanistic and comprehensive

understanding of temporal patterns must account for the sequence of events to capture causality.

As we will explain in more detail below, studying the temporal structure of communities also

provides new quantitative approaches to detecting phenological shifts (e.g., due to climate

change) and putting them in a community context.

The need for a general framework to quantify temporal community structure

Because various studies and related ecological concepts use the sequence and duration of (or

between) events to understand community dynamics (**Table 1**), we believe in the merits of

establishing a general quantitative framework designed to capture the critical features of

temporal community structure. Current approaches to analyzing the temporal structure of

communities fall mainly into a few research areas: phenology, species turnover (including

succession), and interaction networks. Each approach has strengths and limitations in capturing

the complexity and essential features of the temporal structure of communities.

To date, phenology studies have predominantly focused on the onset (timing) or duration of

phenological events of single species (CaraDonna *et al.* 2014; Cohen *et al.* 2018; Forchhammer

et al. 1998), pairwise correlations of these events (Ovaskainen *et al.* 2013; Thackeray *et al.*

2016), and the temporal overlap of interacting pairs of species (Carter *et al.* 2018; Renner &

Zohner 2018; Visser & Both 2005; Visser & Gienapp 2019). While these and other studies have

been fundamental in documenting the temporal coordination and variability of interactions over

time, many aspects of the temporal community structure only emerge from the community

context. For instance, the dynamic structure of communities could show periods of high activity

separated by periods of inactivity. We can only identify such bursts of activity by analyzing the

distribution of "gaps" between phenologies of all species. Distributions of multiple species'

phenologies and gaps between them could characterize distinct temporal modules, e.g., subsets

of interacting species that always co-occur or a set of species in a module to consistently appear

in the same chronological order. Such modules may help predict the presence/absence and

strength of interactions, symbiotic relationships, temporal guilds, temporal niche partitioning, or

identify temporal windows of opportunity and species that exploit similar resources or abiotic

conditions (**Table 1**). Similarly, the ecological consequences of phenological shifts may only

become fully apparent in a community context (e.g., when shifts create temporal partitioning,

Fig. 1). In short, the temporal features characteristic of species' phenologies may suggest

important community dynamics, but identifying these temporal features requires a holistic

approach that moves beyond analyses of single-species or pairwise metrics that we traditionally

use in phenological studies.

On the other hand, studies on temporal succession focus on turnover in species composition

(e.g., Connell & Slatyer 1977; Dornelas *et al.* 2014; Penny *et al.* 2023; Poorter *et al.* 2023; Van

Allen *et al.* 2017) and interactions (CaraDonna *et al.* 2021; e.g. Díaz-Castelazo *et al.* 2010;

Kantsa *et al.* 2018; Suzuki *et al.* 2023; Ushio *et al.* 2018) and thus prioritize a multi-species

perspective to capture how communities change over time. These studies revealed that species

compositions and interactions are highly dynamic and can rapidly turn over across years,

seasons, weeks, and even days, while some structural patterns at the network level (e.g.,

nestedness or connectivity) may change comparatively little (Kaartinen & Roslin 2012;

Petanidou *et al.* 2008; Schwarz *et al.* 2020). These studies also highlight how we can infer

important ecological and evolutionary processes that drive community dynamics (e.g., the

concept of "succession" or "community drift") from the temporal organization of species.

However, they still mainly rely on static metrics (either based on snapshots or time-aggregated

data) that, at best, only contain indirect information on the sequence and duration of events. For

instance, metrics for temporal turnover in species composition indicate that species come and go

(e.g., Dornelas *et al.* 2014; Penny *et al.* 2023), but these metrics don't provide information on the

duration and sequence of events, e.g., how long species are present, which ones might return,

how the current composition relates to previous events and if community states differed in how

long they last. To use the music analogy again, temporal turnover, for example, helps quantify

the tempo of a song but not the melody itself. Consequently, turnover can only capture a subset

of patterns of the temporal structure itself. Of course, it would be unrealistic to assume that any

single metric can comprehensively describe the complex temporal organization of communities.

Accomplishing this requires a comprehensive toolset: a set of temporally explicit metrics, with

each metric specialized to capture a different aspect of the dynamic temporal organization of

communities.

Thus, despite significant recent advances that helped develop a more temporally explicit view

of community ecology, we still have much to learn from the temporal structure of communities.

A general framework accounting for the pros and cons of existing approaches would bring

together different research areas in community ecology for a more comprehensive understanding

of temporal community structure. Especially in this age of unprecedented environmental change,

doing so should help the field better understand communities and anticipate how they will

respond.

A general approach to studying the temporal structure of natural communities: time-

ordered co-occurrence networks

Here, we propose that time-ordered co-occurrence networks provide a powerful unifying

framework to study the temporal structure of natural communities. Spatial ecology has proven

over the past decades that we can learn a lot from studying the (co-) occurrence patterns of

species (Araújo *et al.* 2011; Borthagaray *et al.* 2014; Freilich *et al.* 2018; Galiana *et al.* 2023;

Gao *et al.* 2022). The proposed framework takes a similar comparative approach for analyzing

the temporal organization or species co-occurrences, except that it accounts for the directionality

of time and keeps track of the possible causal chains of events.

Time ordered networks

Time-ordered networks (also called time-varying, temporal, or dynamic networks) differ from

traditional static networks because they are inherently dynamic and provide a complete record of

the sequence and duration of events (Holme & Saramäki 2012; Masuda & Lambiotte 2020). In

these networks, each node (vertex) and link (edge) is only active during specific time points

(shown explicitly in **Fig. 2**). The unidirectionality of time (time always moves forward) means

that the relationships between nodes have to follow a "time-respecting walk" of adjacent nodes

and links. Importantly, these time-respecting walks have to follow the natural temporal sequence

of events and thus lay out the possible causal chains contributing to community dynamics. For

instance, if the path between nodes *A* to *B* ends before the link between *A* to *C* becomes active

(**Fig. 2** scenario (a)), then node *B* could potentially indirectly affect node *C* (via *A*), but nothing

from *C* can propagate to *B*, since signals cannot travel back in time. Thus, analyzing the structure

of temporally-ordered networks helps us to better visualize and quantify the organization of

events in time.

The study of time-ordered networks is ongoing, but it is a rapidly growing interdisciplinary

field (Holme & Saramäki 2019; Masuda & Lambiotte 2020). Time-ordered networks, in general,

have been used to describe temporal patterns for a range of systems, from public transportation

and neurons in the brain to social networks (Holme & Saramäki 2019), but these networks have

received surprisingly little attention in community ecology. Yet, as we will show below, they are

ideally suited to study the temporal organization of natural communities, and recent advances

and development of analytical tools not only help overcome the limitations of previous

approaches discussed above but also provide unique opportunities for new temporally explicit

analyses.

Building time-ordered co-occurrence networks from phenology data

We will focus here on phenology data to demonstrate how we can apply this approach to

quantify the temporal organization of species occurrences (but see "Future directions" for how

this can be applied to other types of data). We already have a rich body of phenology data for

various animal and plant communities in marine, freshwater, and terrestrial systems that are

typically publicly available (e.g., USA National Phenology Network, Pan European Phenology

project, Environmental Data Initiative). These data sets often have a fine temporal resolution and

span decades, and they are rapidly growing with help from researchers, community-driven

science initiatives, and government-funded observation networks (e.g., NEON, GBIF).

Collecting this data is being increasingly facilitated by recent technological advances. For

instance, deploying cameras and machine learning to identify flowers and even pollinators at

very fine temporal scales (Besson *et al.* 2022) can significantly help reduce the workforce

needed to collect this type of data. Similarly, DNA approaches can quantify phenologies and

even interactions at similar or even finer temporal scales than direct observation (e.g., meta-

barcoding of airborne pollen DNA, pollinator DNA in flowers, or eDNA analysis in freshwater

and marine systems) (Besson *et al.* 2022; Jensen *et al.* 2022; Jønsson *et al.* 2023). Importantly,

phenology data provide essential information about the temporal structure of communities by

documenting when species are present and active in a given community.

Building time-ordered networks from phenology data (phenology networks) is relatively

intuitive and generally straight forward in practice (see **Fig. 2** for a worked example). Following

traditional approaches, we can build a network where a species represents a node, and its

phenology determines the node's time stamps (interval). Depending on the system, a node may

thus have a single timestamp (e.g., present/active for a single day), a time interval (e.g., present

from day x-y), or multiple time stamps if there is a temporal gap (pause) between activity periods

(e.g., multiple distinct breeding events in a year). Links show relationships between nodes to

represent when two species are active simultaneously in a given community. The time stamps for

links would be similar to those of nodes (e.g., single or multiple time stamps), except that they

represent the duration of the temporal overlap. In time-ordered networks, these links are

sometimes called "contacts" to emphasize a time-limited information exchange (Holme 2015).

Constructed in this way, these phenology networks capture co-occurrences and, thus, the

complete temporal structure of communities.

Of course, if additional information on species interactions is available (or inferred from the

time series itself (e.g., Ushio *et al.* 2018)), it is straight forward to further restrict the topology of

these co-occurrence networks to focus only on the direct interactions. However, this would shift

the focus from analyzing temporal co-occurrence patterns of species to analyzing temporal

patterns of direct interactions. Furthermore, species or individuals could be grouped in other

ways, e.g., by traits, functional roles, or life stages. A particularly intriguing approach is to use

this time-ordered network approach to track transitions in community states, with nodes

representing community states and timestamped links signaling transitions between these states

(see **Box 2 and Fig. 3** for an example). This approach is particularly promising in more complex

systems and can help identify cycles, critical transitions in community states, and which species

drive them. In general, this time-ordered network approach is very flexible, and the type of

structure used to define nodes and links ultimately depends on the specific questions of interest

and available information.

Quantifying time-ordered networks to understand natural communities

Converting species' phenologies into time-ordered networks opens the door to calculating a

range of informative metrics that could provide new and more profound insights into species'

biology, their temporal relationships, and the organization of natural communities. Some of these

metrics for time-ordered networks are unique to temporal analyses, while others build on static

network metrics to account for the unidirectionality of time. Here, we highlight and discuss a

subset of previously developed time-ordered network metrics (**Table 2**) that we believe are

among the most conceptually straightforward and biologically meaningful for analyzing

phenology network data. Existing software can already calculate most of these metrics (e.g., tsna

package (Bender-deMoll & Morris 2021), timeordered (Blonder 2015), and networkDynamic

(Butts *et al.* 2023) in R, and Straph in Python). Several metrics like burstiness and memory have

applications beyond network analysis and do not necessarily rely on network-specific software.

Altogether, these metrics for phenological data capture important network properties such as

connectivity, transmissibility, and persistence and the sequence and duration of events, the key

features of temporal structure.

To better understand these metrics' biological meaning and utility, it helps to think about

what information they capture. Connectance metrics like the "*upstream set"* and "*downstream*

set" (**Table 2**) (also called backward reachable set and forward reachable set) show the

relationships of species to other community members that appeared before or after them,

respectively (see **Fig. 2** for worked-out example). For instance, we can compare the *downstream*

set across species to identify which species have the highest potential to influence other members

within a given community or how infections might spread.

Connectivity also depends on transmissibility—how easily and fast a signal or substance can

travel across a community. In static networks, we typically measure this with the "*shortest path*",

i.e., the smallest number of nodes we must traverse to connect one node to another. In time-

ordered networks, this is sometimes called "*flow*" (Blonder *et al.* 2012), but distances between

nodes can be measured in two ways in time-ordered networks: the number of nodes that need to

be traversed (*distance*) or the actual time required to travel from one node to another (*latency*).

Comparing these metrics across species gives us a better idea of species' relative temporal

position in a given community. They could also help identify the potential for indirect effects

between species (Box 2), differences in the sensitivity of communities to environmental

perturbations (e.g., how fast perturbations or infectious diseases could spread in communities

and how many species may be affected), or at what point in time communities are most

vulnerable to perturbations. These connectance metrics depend on the sequence and duration of

and between events. Thus, they can give important insights into the functional role of species and

their relationships in time (**Table 2**).

Other metrics, such as *burstiness, memory, node,* and *link* persistency (**Table 2**), focus on the

duration of or between events to quantify the permanence/transience of time-ordered network

structures, including temporal turnover of interactions. A high level of burstiness can help

identify windows of opportunity (e.g., favorable environmental conditions, resource pulses) or

temporal guilds. In contrast, a low level of *burstiness* could indicate temporal niche partitioning

(Table 1). Link persistency can indicate the interaction potential of species (Castro☐Arellano *et*

al. 2010; Resasco *et al.* 2021), while node persistency could help identify different life history

strategies among species (e.g., seasonal specialists with short node persistency vs. generalists

with long persistency).

We want to emphasize that time-ordered networks have not been widely applied to

phenological data, so there is much room to grow and develop new metrics and tools. The

metrics presented here are just a starting point, and we hope they inspire the development of

complementary approaches for analyzing temporal communities. For instance, how can time-

ordered networks incorporate count data from phenological datasets as weights to predict

ecological consequences of co-occurrences better? How can we implement sequence in metrics

of overlap or account for differences in proportional overlap? Developing these and other metrics

is a promising new research area that will grow as we learn about temporal structure.

Practical considerations

The size of any time-ordered network scales with the length of the period analyzed. Results of

several temporally explicit metrics can differ based on the length of time over which network

structures get quantified. This difference is particularly important for metrics that rely on time-

respecting path lengths since they typically correlate positively with the period length (e.g.,

upstream and downstream set, distance, betweenness centrality in **Table 2**). Therefore, as best

practice, any analyses of temporal community structure should have a clearly defined time

window *t*. Fortunately, when analyzing phenology data, the calendar year or seasonal constraints

(e.g., start and end of snow cover) often set the temporal window. However, these boundaries are

sometimes blurred for some systems, e.g., when individuals or life stages stay active all year

round. In this case, researchers must identify other biologically meaningful time windows (e.g.,

Does the system show a cyclical pattern reflecting the timing of El Nino Southern Oscillation

events?) or stick with widely used time windows like the calendar year. In both cases,

researchers must clearly communicate the time windows used in their analyses and standardize

network metrics by the length of time windows when comparing time-ordered networks with

variable time windows.

Visual representation of time-ordered networks can become cumbersome when exceeding a

certain complexity. However, for metrics or network visualizations like those in **Fig. 3**, we can

adapt existing core, open-source software packages developed for static networks (R packages:

ggraph, igraph, network; Python packages: NetworkX). Given the increasing popularity of time-

ordered networks and the development of new comprehensive guides (Masuda & Lambiotte

2020), we anticipate that more user-friendly options that work more directly with time-ordered

networks will emerge over the next decade.

Finally, we must consider potential data limitations and caveats to help decide when this

approach will be most helpful. Like all network approaches, it becomes more powerful when

more nodes (species) are included. Similarly, very short or sparse time series may not contain

enough information to reveal specific temporal patterns. Some metrics, like memory, require a

minimum number of time points (**Table 2**). While phenology data is becoming increasingly

available, the data recorded varies across studies, which determines the types of possible

analyses. For instance, phenology datasets may only have start dates once per species per year

(e.g., first flowering, first emergence). Without a biological notion of start and end times, a time-

ordered network reduces to a temporal path or chronological sequence of events. This limits the

amount of information we can extract from the temporal pattern. However, we can still extract

some helpful information on temporal structure that can help put phenological shifts in a

community context (see **Supplement, Fig. S1, S2)** for an example of an empirical application

using a data set on the leafout of woody plants).

Applications and future directions

The acceleration of climate change has created a need for a more temporally explicit

approach to ecology in the Anthropocene to understand and predict how these changes will affect

natural ecosystems (Wolkovich *et al.* 2014; Yang 2020; Yang & Rudolf 2010). To predict how

systems will change, we first need to understand the temporal structure of communities and the

mechanisms that drive this structure. To date, progress in answering these questions has been

hampered by a lack of a unified framework and limited available methods.

Our framework aims explicitly to overcome these limitations by offering comprehensive

quantitative approaches that could scale across communities. With multiple metrics in hand,

researchers can visually and quantitatively assess and compare the temporal structures of

communities across timescales, space, and ecosystems (see **Box 1** and **Fig. 3** for a worked

example) and help identify the underlying mechanisms. We already mentioned some applications

above but would like to highlight some additional exciting new research avenues and

applications. Of course, these areas serve as starting points and do not represent a complete list

of potential applications.

Functional roles and relationships among species

One novel aspect that emerges from this framework is that it can provide new ways to

quantify and compare species' "temporal niche" and functional roles by putting them in a

temporally explicit community context. This analysis can reveal previously hidden functional

differences between species within and across communities. For example, species that co-occur

with the same number of species but differ in how long they are present in the community, when

they are present, or how their interactions are distributed over time could show different

responses to species extinctions and other environmental perturbations. To better understand

species' roles at the community level, we can perform robustness and sensitivity analyses (e.g.,

randomly removing species or species at risk of extinction) (Dunne & Williams 2009; Rudolf &

Lafferty 2011) on these time-ordered networks to identify keystone species (e.g. those that result

in most significant changes in time-ordered network metrics), or those species and ecosystems at

greatest risk (i.e. species that experience the largest or more frequent changes in node-specific

network metrics). These findings could help guide management and conservation efforts in a

rapidly changing world.

General patterns and environmental drivers of temporal community structure

Given that temporal structure is non-random and reflects a community's ecological and

evolutionary processes, we can expect to learn a lot from studying temporal patterns. For

instance, perhaps weak burstiness and memory of fruiting plant communities (**Fig. 3e**) could

arise from plant species starting but not ending their fruiting phenophases in the same month.

Such a pattern suggests a potential shared underlying driver that might be linked to changes in

environmental conditions or species' life histories. However, weak burstiness could also arise

from a temporal niche partitioning, e.g., to avoid competition for frugivores.

When we applied our approach to compare two very different communities (anuran calling

versus a plant fruiting community), we found striking differences in the temporal network

structure and some surprising similarities (see Box 1 and Fig. 3 for a complete example). This

comparison highlights the power of having a set of metrics at hand to detect generalities and

essential differences in the temporal structure across communities and environments that do not

necessarily share the same abiotic drivers. At the same time, our worked examples also highlight

significant gaps in our current knowledge of temporal community structure: ecology currently

lacks a general framework to predict what these structures should look like or why and how they

should vary across systems. Are those differences representative of the respective ecosystems or

different environments? Do the similarities indicate shared drivers and biological constraints, or

do they arise by chance? How will the patterns for a given type of ecosystem vary across

different environments or with different species combinations? What will other types of

communities look like? Answering these and many other related questions will significantly

advance our understanding of the ecological and evolutionary processes shaping natural

communities.

A crucial step toward answering these questions can involve comparing network metrics

(**Table 2**) across systems, time, and space to identify potential drivers that shape the temporal

structure of communities. For instance, our approach allows us to compare network metrics

across latitudinal or altitudinal gradients to test how differences in seasonality influence the

temporal organization of communities. Growing and reproductive seasons are often shorter in

colder climates, but we don't know how consistently this abiotic constraint shapes the temporal

structure of communities. Do communities with shorter breeding/growing seasons show higher

temporal overlap but shorter co-occurrences or the opposite? Does a shorter season alter the

connectivity of time-ordered networks and their temporal relationships (e.g., *upstream set* vs.

downstream set, **Table 2**)? Are changes uniform across species, or do they depend on their

temporal position (e.g., early vs. late in the season)?

We can also see great potential in using our framework with temporally explicit null

modeling approaches to narrow down potential mechanisms driving temporal structure. For

instance, we can randomize time stamps of nodes within a given community to break up

sequence and temporal correlations or randomize the duration of phenophases to decouple the

sequence and duration of events. The high degree of flexibility offered by these null or random

network models allows for ensembles of increasingly randomized time-ordered networks useful

for disentangling and identifying nested temporal patterns (Holme 2015). Comparing metrics

from reference null models to empirical patterns within or across ecosystems helps identify non-

random patterns in different aspects of temporal structure and better understand the processes

that control temporal network structure. Say the latency between a set of species is much shorter

than expected by chance. Then, there are processes causing temporal correlations between these

species (Blonder *et al.* 2012). Ultimately, just like with spatial patterns (Gotelli *et al.* 2010;

Gotelli & Graves 1996), we believe that confronting different null models with empirical data of

time-ordered networks offers a particularly powerful approach for detecting general patterns in

temporal community structure and for testing hypothesis to identify the underlying mechanisms

of community assembly.

Quantifying phenological shifts in a community context

Time-ordered networks also provide new tools to study the ecological consequences of

phenological shifts. We want to reiterate that while past studies have primarily focused on single

species or pairs, some effects may only become apparent in a community context. Small changes

in the phenology of a species may have significant consequences at the community level

depending on the temporal position of the species relative to other community members. For

instance, the advancing phenology of species A in **Fig. 2** fundamentally alters the network

structure and thereby affects the metrics (upstream and downstream set) of all species. In

contrast, a similar shift in species D would have little consequence. Our approach represents a

step toward identifying these species (e.g., species' phenologies most important for maintaining

the temporal structure and functioning of the existing ecosystem) and the groups likely to be the

most affected by phenological shifts. The relative consequences of phenological shifts predicted

from these networks generate *apriori* hypotheses that researchers could test experimentally.

Paleo data

Finally, we can also use time-ordered networks to analyze other data types. For instance,

recent advances in genetic methods have improved Quaternary paleo-records (Fordham *et al.*

2020), which have emerged as an important way to explore the long-term dynamics of

communities and how they are linked to climate change (Blois *et al.* 2010; Jackson & Blois

2015). Our approach could easily be used with this type of data by simply using the presence of

species in the archaeological record instead of phenologies to build time-ordered networks that

study how communities have changed in the distant past. Since both types of data document co-

occurrences, time-ordered networks could help bridge gaps between community ecology of the

present and distant past, thereby improving our ability to understand the potential consequences

of climate change (Fordham *et al.* 2020).

Conclusions

Like spatial ecology, integrating temporal structure into community ecology has the potential

to fundamentally alter our understanding of the critical forces and processes that shape natural

communities. To date, we still lack a general understanding of how temporal structures of

communities vary, what general patterns might exist in nature, and what the underlying drivers

are. We know even less about their consequences. Nevertheless, answering these questions is

fundamental to understanding the maintenance of ecological communities (Ushio et al. 2018)

and predicting how they will respond to future perturbations and climate change. Our framework

outlines how to take the next important steps toward building a temporally explicit framework

for community ecology and suggests new and exciting opportunities to help overcome some

previous limitations. Building this framework is particularly important to understand and predict

how natural ecosystems will change as climate change reshuffles the temporal structure of

natural communities worldwide. The next challenges include collecting, accessing, and

analyzing this temporal data to shed new light on this understudied dimension of natural systems.

Acknowledgements

We would like to thank A.E. Dunham, J. HilleRisLambers, A. Silva, and A. Simha for

feedback on the ideas, methods, and manuscript. This work was supported by NSF DEB-

1655626 to V.H.W. Rudolf and by the NSF Graduate Research Fellowship Program under Grant

No. 1842494. Any opinions, findings, and conclusions or recommendations expressed in this

material are those of the author(s) and do not necessarily reflect the views of the National

Science Foundation.

Box 1: *Why is the sequence and duration of events so important?*

The critical aspects of temporal structure—sequence and duration of and between events—play essential roles in structuring species interactions and ecosystem processes. The sequence and duration of species' phenophases determine which species co-occur and, thus, what interactions are possible. A predator cannot consume prey, and a bee cannot pollinate a flower if neither is present at the right stage at the right time(s). Understanding what interactions are possible through temporal co-occurrences can thus help predict and capture species interactions, even though co-occurrence does not necessarily imply a direct interaction between species (Blanchet *et al.* 2020; Thurman *et al.* 2019). For instance, the loss of temporal overlap of phenologies can explain the loss of plant-pollinator interactions over a century (Burkle *et al.* 2013). Similarly, species with longer phenophases tend to co-occur with more species and thus have more species interactions (Resasco *et al.* 2021) (**Fig. 1a**). However, in this case, interactions also typically occurred in sequence with only one or a few interactions at any given point in time. Thus, species that might have seemed like generalists were often "sequential specialists" (**Fig. 1b**). These examples highlight that the sequence and duration of species' activities help identify the possible interactions and reveal the forces that structured life histories and community composition.

The sequence of species arrival times and duration between arrivals can also determine the outcome and strength (i.e., per-capita effect) of species interactions (Rudolf 2019; Zou & Rudolf 2023) (**Fig. 1c**). For instance, reducing the overlap of flowering times can either increase or decrease the fitness of plant species in a community via recruitment or competition for pollinators respectively (Faust & Iler 2022; Mitchell *et al.* 2009). Early arrivers often have a competitive advantage over species that arrive later, and this advantage typically depends on the

difference in relative arrival times (Alexander & Levine 2019; Blackford *et al.* 2020; Rasmussen

et al. 2014; Rudolf 2018). Similarly, predation rates decline with a delay in salamander and

dragonfly predator arrival relative to its tadpole prey (Rudolf 2022a). Such timing effects (also

called priority effects or historical contingencies) are well documented in animal, plant, and

microbial systems and determine the dynamics and composition of natural communities (Fukami

2015; Zou & Rudolf 2023).

Furthermore, the sequence of species' arrivals in a community and the time interval between

arrivals determines the chain of causal events. In the absence of temporal overlap, species

present earlier in a community can affect those that appear later but not vice versa (**Fig. 1d**). For

instance, experiments in pond communities found that *Bufo* tadpoles that breed early in the

season can negatively affect later arriving *Hyla* tadpoles, even though the two species never co-

occurred in time (Wilbur & Alford 1985). Such "legacy effects" are common in natural

communities and can arise through a range of different mechanisms (Fukami 2015), e.g., by

altering the environment (Kostenko *et al.* 2012), consuming a competitor of a later arriving

species, or by changing the community composition that later arriving species experience

(Rudolf & Van Allen 2017). However, this potential to affect other community members

naturally declines towards the end of a season; e.g., species that arrive at the end of a season tend

to have fewer interactions (i.e., are more specialized) (Seifert *et al.* 2021). The species that arrive

last can only affect a limited number of species with which they co-occur. The relative temporal

position of species is, therefore, inherently linked to its potential impact on the community and

how it can be affected by other community members.

At the community and ecosystem level, the duration and sequence of temporal co-occurrences

limit the possible pathways through which diseases, energy, water, organic and inorganic

elements, and perturbations cascade through the ecosystem. Perturbations (e.g., fire, drought,

heatwave, chemical spill, or disease outbreak) that happen in the middle of a season can only

affect species that are already present or arrive later in the season, while species that arrived and

left before the perturbation will not be affected. Thus, explicitly accounting for the sequence and

duration of events is imperative to capture the mechanisms that drive community dynamics and

composition, ecosystem processes, and how they will respond to environmental change.

Box 2: *Applying the temporal framework: temporal patterns of anuran calling community vs. plant fruiting community*

To highlight some of the utility of our framework, we show an example of how we can use it to compare the temporal structure of communities using publicly available data sets on two very different systems: one on anuran calling in Texas (Carter *et al.* 2018) and the other on fruiting phenology in Thoreau's Concord (Ellwood *et al.* 2022a) (**Fig. 3**). Due to the complexity of the communities and to highlight the flexibility of our approach, we focused here on changes in community states. The approach to building time-ordered networks is still the same (see **Fig. 2**), except that here, nodes represent a unique combination of co-occurring species (community state), and links (relationships between nodes) show the transition between the different community states. These links have time stamps for when transitions occur, while nodes have time intervals for the duration of community states.

Just comparing the network visually, we can immediately see striking differences in their temporal structure (**Fig. 3a**). The anuran-calling time-ordered network shows many recurring community states (combination of species) but with considerable variation in how long each community composition lasts and how often it reappears (green histograms in **Fig. 3b** and c, respectively). This pattern arises because most species have more than a single time interval during which they are active and can vary in the length of their phenophases (i.e., few species with long vs many with short phenophases. Together, this creates a complex and highly dynamic network, where species with long phenophases (which form the long tail in the histogram in **Fig. 3b**) are more central and act as "hubs" connecting different community states. In contrast, the fruiting network suggests a sequential turnover of community states because species typically only fruit during a single period. Consequently, there is a much more explicit temporal hierarchy

with a simpler causal structure (who comes first and could influence later arrivers). Each

community state typically occurs only once (note that because in and out transitions count each

separately, this pattern is indicated by the step peak at 2 in **Fig. 3b**). In both networks, most

states (nodes) do not last long before transitioning to another state, but a minority of states in the

plant fruiting network last much longer and form a right-skewed distribution. When plotting the

distributions of how long each community state lasts on average (node persistence) and recurring

community states in the same plot (**Fig. 3d**), we see the networks follow very different

relationships reflective of differences in time-ordered network structure (i.e., strongly positive

relationship in frog network vs. weak to no relationship in plant network). Yet, despite these

differences, both communities show similar temporal distribution of events at the community

level, i.e., a weak signal of temporal clustering of community shifts (burstiness in **Fig. 3e**) and no

clear autocorrelation in times between community shifts (memory in **Fig. 3e**). These examples

illustrate just some of the various analyses that can be performed with our approach to

comprehend better the common and unique temporal structures and functions of natural

communities.

Table 1: Examples of ecological concepts and how they are related to temporal patterns

Ecological concept	Temporal pattern used to infer concepts	Example references
Mechanisms of community assembly	Reduced temporal variation & non-random temporal patterns (over time, e.g., years or habitats) were used to infer the strength of stochastic vs. deterministic processes, historical contingencies, & priority effects.	(Neale & Rudolf 2023; Purschke *et al.* 2013; Van Allen *et al.* 2017)
Niche partitioning	Lower temporal overlap than expected from a null model was used to infer temporal niche partitioning and competition.	(Jensen *et al.* 2019; Koide *et al.* 2007; Loreau 1989b; Schamp & Jensen 2019)
	Temporal restructuring (change in phenologies) after species removal as an indicator for temporal niche separation to avoid competition	(Dumandan *et al.* 2023)
Coexistence mechanisms	The sequence of arrivals and time between them determines the potential for coexistence (via correlated changes in fitness and niche differences)	(Alexander & Levine 2019; Blackford *et al.* 2020; Godoy & Levine 2014; Rudolf 2019)
Functional relationships	Temporal covariances of occurrences used to infer species	(Deutschmann *et al.* 2023; Steele *et al.*

of species	interactions	2011; Ushio *et al.* 2018)
	Temporal clustering was used to infer facilitation/positive indirect interactions between species.	(Dante *et al.* 2013)
	Phenological tracking can indicate strong inter-dependency of species (e.g., predator-prey, plant-pollinator)	(Visser & Gienapp 2019)
	Competitive dominance depends on the relative time between and order of species arrivals.	(Faust & Iler 2022; Rudolf 2018; Rudolf & McCrory 2018)
	Phenology timing determines predator-prey and host-parasitoid interactions and community structure.	(Alford 1989; Pardikes *et al.* 2022; Rudolf 2022b)
Network structure and stability	Loss of temporal overlap results in loss of interactions	(Burkle *et al.* 2013)
	Phenological attributes determine network robustness and persistence of species.	(Guzman *et al.* 2021; Ramos–Jiliberto *et al.* 2018)
	Phenology drives modularity	(Morente-López *et al.* 2018)
Invasion success	Invasion success/growth rates dependent on the relative timing of arrival (germination or flowering)	(Godoy & Levine 2014; Wolkovich & Cleland 2011; Zettlemoyer *et al.*

		2019)
Specialist vs. generalist	Species with narrow time windows (short presence) are more likely to specialize on temporary resources and certain environmental conditions (e.g., seasonal specialist vs. generalist)	(Humphries *et al.* 2017)
Potential pathways for and speed of disease, information, and energy transmission	Temporal structure predicts the possible pathways and transmission speed through a community (e.g., faster with higher average overlap, less time between arrival, etc.)	(Blonder *et al.* 2012)
Effect of perturbations	The effects of perturbations on species and communities depend on the specific timing of perturbation and temporal organization of species.	(Craine *et al.* 2012; Crawley 2004; Li & Pennings 2017; Shepherd *et al.* 2012)
Windows of opportunity	Temporal clustering and gaps in species presences can indicate favorable (& unfavorable) environmental conditions.	(Martinez Falcon et al 2010)

580 **Table 2: Metrics to quantify different aspects of the temporal structure of natural communities.** For consistency, the meaning

581 and applications column assumes the structure is a time-ordered species-co-occurrence (e.g., phenology) network. Although

582 definitions vary among studies, these are all previously developed metrics and can be calculated with current software packages (see

583 main text for details). Here, we largely follow the notations from (Holme & Saramäki 2019).

Temporal structure	Metric	Definition of metric	Suggested biological meaning & potential applications
Sequence of events	Time-respecting traversal/path	Indicates sequences of links (edges) that can be traversed in a time-ordered network under the constraint that the following link to be traversed is activated at some point in time after the current one.	This concept is a foundation for many time-ordered network metrics and establishes a potential **causal relationship**. It assures that the connection between nodes (e.g., species) and time-ordered network metrics account for the temporal sequence/order of events and their durations. It reflects that time always moves forward.
	Downstream set	The collection of all nodes that can be reached from a node *i* via	Indicates *which* and *how many species could be "influenced" by a focal species*. This includes species (nodes) that do not have direct temporal

		time-respecting paths.	overlap with the focal species. The distribution of downstream sets across all species identifies species that play a central role in connecting the network or could exert a disproportionate influence on others. (see **Fig. 2** for a worked-out example)
	Upstream set	Number of nodes that can reach a given node i via temporal paths	Indicates *which* and *how many species could affect a focal species*. Like a stream where water flows downhill and not uphill, nodes in the downstream set cannot affect nodes in the upstream set, due to the unidirectionality of time. Note that since this definition relies on temporal paths, the upstream set strings together a series of co-occurrences with no temporal gaps between co-occurrences from back in time up to the specified present time. The resulting downside is that the upstream set will not count species creating gaps in time between co-occurrences, even though the species and co-occurrences happened before the focal species. (see **Fig. 2** for a worked-out example)

			Combined with the downstream set, the upstream set helps determine how well-connected a species is. This information could be used to infer the temporal position of species relative to other community members or how likely it is to be influenced by events that happen before it even appears.
	Distance	The smallest number of nodes between node x and y following time respective traversal	Indicates *how many indirect contacts are needed to reach another species*. The **average distance** tells us *how spread out a network is* (related to temporal overlap but goes beyond pairwise overlap). It provides new insights into *the functional roles of species (or other groups) in a community* by resolving relationships between species in a temporal context. At the community and ecosystem level, the pace of phenological shifts this metric identifies suggests *how perturbations, energy, or infections "flow" through a community and at what time scale*.
	Time-dependent –	Measures the fraction of shortest	Measures how well connected a node is to others. It can *identify a node's relative level of importance in terms of its ability to act as an*

	betweenness centrality	paths that go through a node	*intermediary to other nodes in a network.*
Duration of events	**Node persistency**	Period a node is present	Essentially, it indicates *how long species are present/active*. It is likely to be correlated with connectivity metrics. (see **Fig. 3** for a worked-out example)
	Edge persistency	Period a link is present	Measures *how long species co-occur,* which indicates their **interaction potential**. Applies only to networks where edges can represent durations (e.g., networks in **Fig. 2**)
Duration between events	**Latency**	The shortest time interval between node x and y following time respective traversal	Time-specific equivalent to distance. Essentially tells us *how fast a signal could travel between two species, including species without temporal overlap*. **Average latency** (across all combinations) tells us *how long it takes on average for a signal to travel in a network.*

	Time-dependent centrality - closeness	The inverse of average latency	Measures how quickly a node reaches all others Essentially metric of the centrality of a node, i.e. *how closely is a node connected to all other nodes in a network.*
	Burstiness	Temporal distribution of consecutive inter-event times over a given time	Measures *how evenly events (e.g. contact, or overlap) are spread out over* time. Requires at least three time points. A time-ordered network with a high degree of *burstiness* indicates temporal clustering of events, e.g. when phenologies are highly synchronized across species. Emphasizes events themselves. (see **Fig. 3** for a worked-out example)
	Memory	Correlation between consecutive inter-event times	Determines whether short (long) inter-event intervals are usually followed by short (long) inter-event intervals. A measure of "predictability" of gaps. Requires at least four time points. (see **Fig. 3** for a worked-out example)

Figure 1. Temporal structure, as defined by the sequence and duration of and between

events, influences many aspects of species interactions. Our four examples (a-d) may happen

simultaneously or in different orders and were arranged arbitrarily along a cyclical rather than

linear timeline to emphasize that many natural events reoccur periodically. Line segments

indicate the presence of a species for a temporal (e.g., annual) cycle. Colors represent different

species. **(a)** How long species are present (length of phenophases) typically correlates positively

with the number of interactions per species (e.g., the black species has more interaction than the

green or the grey species). **(b)** Consumers who appear as generalists in static networks (coral)

often feed on resources that occur during different periods (purple, green, black) and thus are not

interchangeable. **(c)** The sequence of species' arrival times and duration between arrivals can

determine the strength of species' interactions. **(d)** The order of phenologies determines what

direct and indirect interactions are possible in a community: early-arriving species (black) can

affect later-arriving species, even after the early arrivers leave (e.g., via resource depletion,

habitat modification, or other community members), but not vice versa.

Figure 2. Steps to build, analyze, and compare time-ordered networks using phenology

data. The start and end date of a phenophase quantifies when species are present and active in a

community (black bars in left column). We can then calculate the species' temporal overlap/co-

occurrence (red bars). From a network perspective, the length of the phenophase (black bar) and

overlap (red bar) indicate the persistency of nodes and links, respectively. We use this phenology

data to build time-ordered networks of communities (middle column), where nodes are species

with corresponding time stamps that indicate when a species is present. Time stamps can be a

single time point, a time interval with start and end date, or multiple points or intervals if species

re-enter a community. The temporal co-occurrence of species gives links that connect nodes

(species) with their corresponding time stamps (red bars). Once these time-ordered networks are

constructed, we can calculate a range of time-ordered network metrics that explicitly account for

the temporal sequence of events (see **Table 2**). For instance, we can find the "downstream set"

and "upstream set" for each species (bar graphs on the right) to determine how species influence

each other given the temporal sequence of species occurrences/phenologies (see **Table 2** for

complete definitions). Circular inserts show a graphical example of how those two metrics are

calculated for focal species B and E, with a solid circle indicating a focal species and an open

circle indicating a corresponding (by color) source or set of influence species. These metrics are

sensitive to the relative temporal position of a species and the sequence of temporal overlap.

They provide new insights into the basic temporal structure of communities and can also be used

to gain a deeper understanding of the functional role of species. For instance, species that appear

earlier often have a more extensive downstream set of nodes but a smaller upstream set than

species that appear late (e.g., B vs E in the top right panel). However, this can change when

temporal gaps (no species present) are present, which create "subsets" of networks (top vs

bottom bar graph). (**a**) vs. (**b**) shows how a phenological shift in a single species can lead to

dramatic changes in network structure and metrics, especially when this results in the loss of a

link that connects temporally separated groups of species (e.g., a shift of species A in top

scenario (**a**) vs. scenario (**b**)), highlighting the utility of this approach for comparing temporal

structure of communities across space and time and quantifying phenological shifts in a

community context.

Figure 3: Changes in community structure over time in two different communities. (a)

Transition networks, where nodes represent unique community states (species combinations,

with each list of two-letter codes for a different species combination or an underscore for all

species absent). The node's size indicates its average duration (scaled to the respective

community) and is scaled on a power function to improve visualization. Directed edges or links

(arrows) indicate transitions between states. Note that we could have shown the same networks

in a more condensed form by having a list of time stamps per arrow (like in **Fig. 2**). This would

reduce the number of arrows drawn repeatedly between the same two community states, but we

decided against this to visually emphasize the difference between the networks. The colors of

links indicate the relative order of transitions (light green = near the end of the calendar year,

dark brown = near the start of the calendar year). The anuran network (left) is based on daily

calling data throughout a ~1-year breeding season at one pond in Texas, USA, in the early 2000s

(Carter *et al.* 2018), while the plant fruiting network (right) comes from Massachusetts, USA

based on estimates during 1850-1860 (Ellwood *et al.* 2022a). Frequency plots show the

distribution of **(b)** the duration (node persistency) of community states and **(c)** the number of

transitions in and out of a given community state in the respective network (note that incoming

and outgoing links count as two separate transitions), and **(d)** indicates how both metrics are

correlated with each other within a network. **(e)** show two metrics that quantify different aspect

of temporal distribution of events at the community level for each network: burstiness and

memory (see **Table 2** for metric details).

Figure 1

Figure 2

Figure 3

References

Alexander, J.M. & Levine, J.M. (2019). Earlier phenology of a nonnative plant increases impacts on native competitors. *Proceedings of the National Academy of Sciences*, 201820569.

Alford, R.A. (1989). Variation in predator phenology affects predator performance and prey community composition. *Ecology*, 70, 206-219.

Anten, N.P.R. & Hirose, T. (1999). Interspecific differences in above-ground growth patterns result in spatial and temporal partitioning of light among species in a tall-grass meadow. *Journal of Ecology*, 87, 583-597.

Araújo, M.B., Rozenfeld, A., Rahbek, C. & Marquet, P.A. (2011). Using species co-occurrence networks to assess the impacts of climate change. *Ecography*, 34, 897-908.

Bender-deMoll, S. & Morris, M. (2021). tsna: Tools for temporal social network analysis. R package version 0.3.5.

Besson, M., Alison, J., Bjerge, K., Gorochowski, T.E., Høye, T.T., Jucker, T. *et al.* (2022). Towards the fully automated monitoring of ecological communities. *Ecology Letters*, 25, 2753-2775.

Blackford, C., Germain, R.M. & Gilbert, B. (2020). Species Differences in Phenology Shape Coexistence. *The American Naturalist*, 195, E168-E180.

Blanchet, F.G., Cazelles, K. & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions. *Ecology Letters*, 23, 1050-1063.

Blois, J.L., McGuire, J.L. & Hadly, E.A.J.N. (2010). Small mammal diversity loss in response to late-Pleistocene climatic change. 465, 771-774.

Blonder, B. (2015). timeordered: Time-ordered and time-aggregated network analyses. R package version 0.9. 8.

Blonder, B., Wey, T.W., Dornhaus, A., James, R. & Sih, A. (2012). Temporal dynamics and

network analysis. *Methods in Ecology and Evolution*, 3, 958-972.

Borthagaray, A.I., Arim, M. & Marquet, P.A. (2014). Inferring species roles in metacommunity

structure from species co-occurrence networks. *Proceedings of the Royal Society B:*

Biological Sciences, 281, 20141425.

Brown, J.H. (1995). *Macroecology*. University of Chicago Press.

Burkle, L.A., Marlin, J.C. & Knight, T.M. (2013). Plant-pollinator interactions over 120 years:

loss of species, co-occurrence, and function. *Science*, 339, 1611-1615.

Butts, C., Leslie-Cook, A., Krivitsky, P.N., Bender-deMoll, S., Almquist, Z., Hunter, D.R. *et al.*

(2023). DynamicExtensionsforNetworkObjects.

CaraDonna, P.J., Burkle, L.A., Schwarz, B., Resasco, J., Knight, T.M., Benadi, G. *et al.* (2021).

Seeing through the static: the temporal dimension of plant–animal mutualistic

interactions. *Ecology Letters*, 24, 149-161.

CaraDonna, P.J., Iler, A.M. & Inouye, D.W. (2014). Shifts in flowering phenology reshape a

subalpine plant community. *Proceedings of the National Academy of Sciences of the*

United States of America, 111, 4916-4921.

Carothers, J.H. & Jaksić, F.M. (1984). Time as a Niche Difference: The Role of Interference

Competition. *Oikos*, 42, 403-406.

Carter, S.K., Saenz, D. & Rudolf, V.H. (2018). Shifts in phenological distributions reshape

interaction potential in natural communities. *Ecology Letters*, 21, 1143-1151.

Castro☐Arellano, I., Lacher Jr, T.E., Willig, M.R. & Rangel, T.F. (2010). Assessment of

assemblage☐wide temporal niche segregation using null models. *Methods in ecology and*

evolution, 1, 311-318.

Cohen, J.M., Lajeunesse, M.J. & Rohr, J.R. (2018). A global synthesis of animal phenological

responses to climate change. *Nature Climate Change*, 8, 224-228.

Connell, J.H. & Slatyer, R.O. (1977). Mechanisms of succession in natural communities and

their role in community stability and organization. *The American Naturalist*, 111, 1119-

1144.

Craine, J.M., Nippert, J.B., Elmore, A.J., Skibbe, A.M., Hutchinson, S.L. & Brunsell, N.A.

(2012). Timing of climate variability and grassland productivity. *Proceedings of the*

National Academy of Sciences, 109, 3401-3405.

Crawley, M.J. (2004). Timing of disturbance and coexistence in a species-rich ruderal plant

community. *Ecology*, 85, 3277-3288.

Dante, S.K., Schamp, B.S. & Aarssen, L.W. (2013). Evidence of deterministic assembly

according to flowering time in an old-field plant community. *Functional Ecology*, 27,

555-564.

Deutschmann, I.M., Krabberød, A.K., Latorre, F., Delage, E., Marrasé, C., Balagué, V. *et al.*

(2023). Disentangling temporal associations in marine microbial networks. *Microbiome*,

11, 83.

Díaz-Castelazo, C., Guimaraes Jr, P.R., Jordano, P., Thompson, J.N., Marquis, R.J. & Rico-

Gray, V. (2010). Changes of a mutualistic network over time: reanalysis over a 10☐year

period. *Ecology*, 91, 793-801.

Dornelas, M., Gotelli, N.J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C. *et al.* (2014).

Assemblage time series reveal biodiversity change but not systematic loss. *Science*, 344,

296-299.

Dumandan, P.K.T., Yenni, G.M. & Ernest, S.K.M. (2023). Shifts in competitive structures can

drive variation in species' phenology. *Ecology*, 104, e4160.

Dunne, J.A. & Williams, R.J. (2009). Cascading extinctions and community collapse in model

food webs. *Philosophical Transactions of the Royal Society of London B Biological*

Sciences, 364, 1711-1723.

Ellwood, E.R., Gallinat, A.S., McDonough MacKenzie, C., Miller, T., Miller☐Rushing, A.J.,

Polgar, C. *et al.* (2022a). Plant and bird phenology and plant occurrence from 1851 to

2020 (non☐continuous) in Thoreau's Concord, Massachusetts. Wiley Online Library.

Ellwood, E.R., Gallinat, A.S., McDonough MacKenzie, C., Miller, T., Miller☐Rushing, A.J.,

Polgar, C. *et al.* (2022b). Plant and bird phenology and plant occurrence from 1851 to

2020 (non☐continuous) in Thoreau's Concord, Massachusetts. *Ecology*, 103, e3646.

Elton, C. (1927). Time and Animal Communities. In: *Animal Ecology*. Sidgwick and Jackson

London, pp. 83-100.

Faust, M.N. & Iler, A.M. (2022). Pollinator-mediated reproductive consequences of altered co-

flowering under climate change conditions depend on abiotic context. *Climate Change*

Ecology, 3, 100043.

Forchhammer, M.C., Post, E. & Stenseth, N.C. (1998). Breeding phenology and climate ...

Nature, 391, 29-30.

Fordham, D.A., Jackson, S.T., Brown, S.C., Huntley, B., Brook, B.W., Dahl-Jensen, D. *et al.*

(2020). Using paleo-archives to safeguard biodiversity under climate change. *Science*,

369, eabc5654.

Freilich, M.A., Wieters, E., Broitman, B.R., Marquet, P.A. & Navarrete, S.A. (2018). Species co-

occurrence networks: Can they reveal trophic and non-trophic interactions in ecological

communities? *Ecology*, 99, 690-699.

Fukami, T. (2015). Historical contingency in community assembly: Integrating niches, species

pools, and priority effects. *Annual Review of Ecology, Evolution, and Systematics*, 46, 1-

23.

Galiana, N., Arnoldi, J.-F., Mestre, F., Rozenfeld, A. & Araújo, M.B. (2023). Power laws in

species' biotic interaction networks can be inferred from co-occurrence data. *Nature*

Ecology & Evolution.

Gao, C., Xu, L., Montoya, L., Madera, M., Hollingsworth, J., Chen, L. *et al.* (2022). Co-

occurrence networks reveal more complexity than community composition in resistance

and resilience of microbial communities. *Nature Communications*, 13, 3867.

Godoy, O. & Levine, J.M. (2014). Phenology effects on invasion success: insights from coupling

field experiments to coexistence theory. *Ecology*, 95, 726-736.

Gotelli, N.J., Dorazio, R.M., Ellison, A.M. & Grossman, G.D. (2010). Detecting temporal trends

in species assemblages with bootstrapping procedures and hierarchical models.

Philosophical Transactions of the Royal Society of London B: Biological Sciences, 365,

3621-3631.

Gotelli, N.J. & Graves, G.R. (1996). *Null models in ecology*.

Guzman, L.M., Chamberlain, S.A. & Elle, E. (2021). Network robustness and structure depend

on the phenological characteristics of plants and pollinators. *Ecology and Evolution*, 11,

13321-13334.

HA, G. (1927). Further views on the succession-concept. *Ecology*, 8, 299.

Holme, P. (2015). Modern temporal network theory: a colloquium. *The European Physical*

Journal B, 88, 1-30.

Holme, P. & Saramäki, J. (2012). Temporal networks. *Physics Reports*, 519, 97-125.

Holme, P. & Saramäki, J. (2019). *Temporal network theory*. Springer Cham.

Humphries, M.M., Studd, E.K., Menzies, A.K. & Boutin, S. (2017). To Everything There Is a

Season: Summer-to-Winter Food Webs and the Functional Traits of Keystone Species.

Integrative and Comparative Biology, 57, 961-976.

Ings, T.C., Montoya, J.M., Bascompte, J., Bluthgen, N., Brown, L., Dormann, C.F. *et al.* (2009).

Ecological networks - beyond food webs. *Journal of Animal Ecology*, 78, 253-269.

IPCC (2023). Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and

III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change In:

IPCC (eds. Lee, H & Romero, J) Geneva, Switzerland.

Jackson, S.T. & Blois, J.L. (2015). Community ecology in a changing environment: Perspectives

from the Quaternary. *Proceedings of the National Academy of Sciences*, 112, 4915-4921.

Jensen, A.M., Schamp, B.S. & Belleau, A. (2019). Evidence of temporal niche separation via

low flowering time overlap in an old-field plant community. *Oecologia*, 189, 1071-1082.

Jensen, M.R., Sigsgaard, E.E., Ávila, M.d.P., Agersnap, S., Brenner-Larsen, W., Sengupta, M.E.

et al. (2022). Short-term temporal variation of coastal marine eDNA. *Environmental*

DNA, 4, 747-762.

Jønsson, K.A., Thomassen, E.E., Iova, B., Sam, K. & Thomsen, P.F. (2023). Using

environmental DNA to investigate avian interactions with flowering plants.

Environmental DNA, 5, 462-475.

Kaartinen, R. & Roslin, T. (2012). High temporal consistency in quantitative food web structure

in the face of extreme species turnover. *Oikos*, 121, 1771-1782.

Kantsa, A., Raguso, R.A., Dyer, A.G., Olesen, J.M., Tscheulin, T. & Petanidou, T. (2018).

Disentangling the role of floral sensory stimuli in pollination networks. *Nature*

Communications, 9, 1041.

Kareiva, P. (1994). Special Feature: Space: The Final Frontier for Ecological Theory. *Ecology*,

75, 1-1.

Koide, R.T., Shumway, D.L., Xu, B. & Sharda, J.N. (2007). On temporal partitioning of a

community of ectomycorrhizal fungi. *New Phytologist*, 174, 420-429.

Kostenko, O., van de Voorde, T.F., Mulder, P.P., van der Putten, W.H. & Martijn Bezemer, T.

(2012). Legacy effects of aboveground–belowground interactions. *Ecology Letters*, 15,

813-821.

Leibold, M.A. & Chase, J.M. (2017). *Metacommunity Ecology, Volume 59*. Princeton University

Press.

Li, S. & Pennings, S.C. (2017). Timing of disturbance affects biomass and flowering of a

saltmarsh plant and attack by stem-boring herbivores. *Ecosphere*, 8, e01675.

Loreau, M. (1989a). Coexistence of temporally segregated competitors in a cyclic environment.

Theoretical Population Biology, 36, 181-201.

Loreau, M. (1989b). On testing temporal niche differentiation in carabid beetles. *Oecologia*, 81,

89-96.

Masuda, N. & Lambiotte, R. (2020). *A Guide to Temporal Networks*.

McMeans, B.C., McCann, K.S., Humphries, M., Rooney, N. & Fisk, A.T. (2015). Food Web

Structure in Temporally-Forced Ecosystems. *Trends in Ecology & Evolution*, 30, 662-

672.

Miller-Rushing, A.J., Høye, T.T., Inouye, D.W. & Post, E. (2010). The effects of phenological

mismatches on demography. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, 365, 3177-3186.

Mitchell, R.J., Flanagan, R.J., Brown, B.J., Waser, N.M. & Karron, J.D. (2009). New frontiers in

competition for pollination. *Annals of Botany*, 103, 1403-1413.

Montoya, J.M., Pimm, S.L. & Sole, R.V. (2006). Ecological networks and their fragility. *Nature*,

442, 259-264.

Morente-López, J., Lara-Romero, C., Ornosa, C. & Iriondo, J.M. (2018). Phenology drives

species interactions and modularity in a plant - flower visitor network. *Scientific Reports*,

8, 9386.

Neale, Z. & Rudolf, V.H.W. (2023). Predation and competition drive trait diversity across space

and time. *Ecology*, 104, e4182.

Ovaskainen, O., Skorokhodova, S., Yakovleva, M., Sukhov, A., Kutenkov, A., Kutenkova, N. *et*

al. (2013). Community-level phenological response to climate change. *Proceedings of the*

National Academy of Sciences, 110, 13434-13439.

Pardikes, N.A., Revilla, T.A., Lue, C.-H., Thierry, M., Souto-Vilarós, D. & Hrcek, J. (2022).

Effects of phenological mismatch under warming are modified by community context.

Global Change Biology, 28, 4013-4026.

Parmesan, C. & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts

across natural systems. *Nature*, 421, 37-42.

Penny, A., Dornelas, M. & Magurran, A. (2023). Comparing temporal dynamics of

compositional reorganization in long-term studies of birds and fish. *Ecology Letters*, 26,

1071-1083.

Petanidou, T., Kallimanis, A.S., Tzanopoulos, J., Sgardelis, S.P. & Pantis, J.D. (2008). Long-

term observation of a pollination network: fluctuation in species and interactions, relative

invariance of network structure and implications for estimates of specialization. *Ecology*

Letters, 11, 564-575.

Poorter, L., Amissah, L., Bongers, F., Hordijk, I., Kok, J., Laurance, S.G.W. *et al.* (2023).

Successional theories. *Biological Reviews*, 98, 2049-2077.

Post, E. (2019). *Time in ecology: a theoretical framework*. Princeton University Press.

Purschke, O., Schmid, B.C., Sykes, M.T., Poschlod, P., Michalski, S.G., Durka, W. *et al.* (2013).

Contrasting changes in taxonomic, phylogenetic and functional diversity during a long-

term succession: insights into assembly processes. *Journal of Ecology*, 101, 857-866.

Ramos–Jiliberto, R., de Espanés, P.M., Franco–Cisterna, M., Petanidou, T. & Vázquez, D.P.

(2018). Phenology determines the robustness of plant–pollinator networks. *Scientific*

reports, 8, 1-10.

Rasmussen, N.L., Van Allen, B.G. & Rudolf, V.H.W. (2014). Linking phenological shifts to

species interactions through size-mediated priority effects. *Journal of Animal Ecology*,

83, 1206-1215.

Renner, S.S. & Zohner, C.M. (2018). Climate Change and Phenological Mismatch in Trophic

Interactions Among Plants, Insects, and Vertebrates. *Annual Review of Ecology,*

Evolution, and Systematics, 49, 165-182.

Resasco, J., Chacoff, N.P. & Vázquez, D.P. (2021). Plant–pollinator interactions between

generalists persist over time and space. Wiley Online Library.

Roslin, T., Antão, L., Hällfors, M., Meyke, E., Lo, C., Tikhonov, G. *et al.* (2021). Phenological

shifts of abiotic events, producers and consumers across a continent. *Nature Climate*

Change, 11, 241-248.

Rudolf, V.H.W. (2018). Nonlinear effects of phenological shifts link interannual variation to

species interactions. *Journal of Animal Ecology*, 87, 1395-1406.

Rudolf, V.H.W. (2019). The role of seasonal timing and phenological shifts for species

coexistence. *Ecology Letters*, 22, 1324-1338.

Rudolf, V.H.W. (2022a). Temperature and nutrient conditions modify the effects of phenological

shifts in predator–prey communities. *Ecology*, n/a, e3704.

Rudolf, V.H.W. (2022b). Temperature and nutrient conditions modify the effects of phenological

shifts in predator–prey communities. *Ecology*, 103, e3704.

Rudolf, V.H.W. & Lafferty, K.D. (2011). Stage structure alters how complexity affects stability

of ecological networks. *Ecology Letters*, 14, 75-79.

Rudolf, V.H.W. & McCrory, S. (2018). Resource limitation alters effects of phenological shifts

on inter-specific competition. *Oecologia*, 188, 515-523.

Rudolf, V.H.W. & Van Allen, B.G. (2017). Legacy effects of developmental stages determine

the functional role of predators. *Nature Ecology & Evolution*, 1, 0038.

Schamp, B.S. & Jensen, A.M. (2019). Evidence of limiting similarity revealed using a

conservative assessment of coexistence. *Ecosphere*, 10, e02840.

Schoener, T.W. (1974). Resource Partitioning in Ecological Communities. *Science*, 185, 27-39.

Schwarz, B., Vázquez, D.P., CaraDonna, P.J., Knight, T.M., Benadi, G., Dormann, C.F. *et al.*

(2020). Temporal scale-dependence of plant–pollinator networks. *Oikos*, 129, 1289-1302.

Seifert, C.L., Jorge, L.R., Volf, M., Wagner, D.L., Lamarre, G.P., Miller, S.E. *et al.* (2021).

Seasonality affects specialisation of a temperate forest herbivore community. *Oikos*, 130,

1450-1461.

Shepherd, B.J., Miller, D.L. & Thetford, M. (2012). Fire Season Effects on Flowering

Characteristics and Germination of Longleaf Pine (Pinus palustris) Savanna Grasses.

Restoration Ecology, 20, 268-276.

Steele, J.A., Countway, P.D., Xia, L., Vigil, P.D., Beman, J.M., Kim, D.Y. *et al.* (2011). Marine

bacterial, archaeal and protistan association networks reveal ecological linkages. *The*

ISME Journal, 5, 1414-1425.

Suzuki, S.S., Baba, Y.G. & Toju, H. (2023). Dynamics of species-rich predator–prey networks

and seasonal alternations of core species. *Nature Ecology & Evolution*, 7, 1432-1443.

Takimoto, G. & Sato, T. (2020). Phenology in a community context: Toward a better

understanding of the causes and consequences of phenology in seasonal environments.

Ecological Research, 35, 442-444.

Tan, J., Kelly, C.K. & Jiang, L. (2013). Temporal niche promotes biodiversity during adaptive

radiation. *Nature Communications*, 4, 2102.

Thackeray, S.J., Henrys, P.A., Hemming, D., Bell, J.R., Botham, M.S., Burthe, S. *et al.* (2016).

Phenological sensitivity to climate across taxa and trophic levels. *Nature*, 535, 241-245.

Thurman, L.L., Barner, A.K., Garcia, T.S. & Chestnut, T. (2019). Testing the link between

species interactions and species co-occurrence in a trophic network. *Ecography*, 42,

1658-1670.

Tilman, D. & Kareiva, P. (2018). *Spatial ecology: the role of space in population dynamics and*

interspecific interactions (MPB-30). Princeton University Press.

Ushio, M., Hsieh, C.-h., Masuda, R., Deyle, E.R., Ye, H., Chang, C.-W. *et al.* (2018).

Fluctuating interaction network and time-varying stability of a natural fish community.

Nature, 554, 360-363.

Van Allen, B.G., Rasmussen, N.L., Dibble, C.J., Clay, P.A. & Rudolf, V.H.W. (2017). Top

predators determine how biodiversity is partitioned across time and space. *Ecology*

Letters, 20, 1004-1013.

Visser, M.E. & Both, C. (2005). Shifts in phenology due to global climate change: the need for a

yardstick. *Proceedings of the Royal Society of London Series B-Biological Sciences*, 272,

2561-2569.

Visser, M.E. & Gienapp, P. (2019). Evolutionary and demographic consequences of

phenological mismatches. *Nature ecology & evolution*, 3, 879-885.

Wilbur, H.M. & Alford, R.A. (1985). Priority Effects in Experimental Pond Communities:

Responses of *Hyla* to *Bufo* and *Rana*. *Ecology*, 66, 1106-1114.

Wolkovich, E.M. & Cleland, E.E. (2011). The phenology of plant invasions: a community

ecology perspective. *Frontiers in Ecology and the Environment*, 9, 287-294.

Wolkovich, E.M., Cook, B.I., McLauchlan, K.K. & Davies, T.J. (2014). Temporal ecology in the

Anthropocene. *Ecology Letters*, 17, 1365-1379.

Yang, L.H. (2020). Toward a more temporally explicit framework for community ecology.

Ecological Research, 35, 445-462.

Yang, L.H., Bastow, J.L., Spence, K.O. & Wright, A.N. (2008). What can we learn from

resource pulses? *Ecology*, 89, 621-634.

Yang, L.H. & Cenzer, M.L. (2020). Seasonal windows of opportunity in milkweed–monarch
interactions. *Ecology*, 101, e02880.
Yang, L.H. & Rudolf, V.H.W. (2010). Phenology, ontogeny, and the effects of climate change
on the timing of species interactions. *Ecology Letters*, 13, 1-10.
Yang, S., Albert, R. & Carlo, T.A. (2013). Transience and constancy of interactions in a
plant☐frugivore network. *Ecosphere*, 4, 1-25.
Zettlemoyer, M.A., Schultheis, E.H. & Lau, J.A. (2019). Phenology in a warming world:
differences between native and non-native plant species. *Ecology Letters*, 22, 1253-1263.
Zou, H.-X. & Rudolf, V.H.W. (2023). Bridging theory and experiments of priority effects.
Trends in Ecology & Evolution.