Effects of blood meal source and seasonality on reproductive traits of

***Culex quinquefasciatus* (Diptera: Culicidae)**

Kevin Alen Rucci[1,3,*], Gabriel Barco[1,3], Andrea Onorato[1,3], Mauricio Beranek[1], Mariana Pueta[2,3] &

Adrián Díaz[1,3*]

[1] Laboratorio de Arbovirus, Instituto de Virología "Dr. J. M. Vanella" (InViV), Facultad de Ciencias

Médicas (FCM), Universidad Nacional de Córdoba (UNC)

[2] Instituto de Investigaciones en Biodiversidad y Medioambiente (INIBIOMA), CONICET -

Universidad Nacional de Comahue (UNCo)

[3] Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina

*Corresponding authors:

Kevin Alen Rucci

Laboratorio de Arbovirus – Instituto de Virología "Dr. J. M. Vanella" (InViV), Facultad de Ciencias

Médicas (FCM), Universidad Nacional de Córdoba (UNC).

Enfermera Gordillo Gómez s/n. Ciudad Universitaria, X5000HUA, Córdoba capital, Córdoba,

Argentina.

E-mail addresses: kevin.rucci@mi.unc.edu.ar, kevinrucci@outlook.com

Adrián Díaz

Laboratorio de Arbovirus – Instituto de Virología "Dr. J. M. Vanella" (InViV), Facultad de Ciencias

Médicas (FCM), Universidad Nacional de Córdoba (UNC).

Enfermera Gordillo Gómez s/n. Ciudad Universitaria, X5000HUA, Córdoba capital, Córdoba,

Argentina.

E-mail addresses: adrian.diaz@fcm.unc.edu.ar, adrian.diaz@conicet.gov.ar

Abstract

Host selection by mosquitoes is a keystone to understanding viral circulation and predicting future infection outbreaks. *Culex* mosquitoes frequently feed on birds during spring and early summer, shifting into mammals towards late summer and autumn. This host switch may be due to changes in mosquito fitness. The aim of this study was to assess if the interaction effect of blood meal source and seasonality may influence reproductive traits of *Culex quinquefasciatus* mosquitoes. For this purpose, *Cx. quinquefasciatus* mosquitoes were reared in simulated summer and autumn conditions and fed on two different hosts, chickens and mice, in a factorial design. Fecundity, fertility, and hatchability during two consecutive gonotrophic cycles were estimated. We found greater fecundity and fertility for mosquitoes fed upon birds than mammals. Fecundity and fertility did not vary between seasons for chicken-fed mosquitoes, whereas in autumn they decreased for mouse-fed mosquitoes. These traits decreased in the second gonotrophic cycle for mouse-fed mosquitoes, whereas they did not vary between cycles for chicken-fed mosquitoes. There was no statistically significant variation of hatchability among treatments. These results indicate a statistically significant interaction effect of blood meal source and seasonality on fecundity and fertility. However, the pattern was opposite in relation to our hypothesis, suggesting that further studies are needed to confirm and expand our knowledge about mosquito biology and its relationship with seasonal host use shifting.

Key words: mosquito vector, fecundity, fertility, hatchability

Introduction

The southern house mosquito, *Culex quinquefasciatus* Say (Diptera: Culicidae), is a worldwide

distributed vector of numerous pathogens, including *Wuchereria bancrofti* Cobbold (Kasili et al.,

2009), *Dirofilaria immitis* Leidy (Labarthe et al., 1998) and avian malaria (Farajollahi et al., 2011).

Besides, this mosquito is also one of the main species responsible for the transmission of the

arboviruses West Nile (WNV) and St. Louis encephalitis (SLEV) viruses (Farajollahi et al., 2011).

Because this, this species is of critical concern due to its impact on both public and veterinary

health.

Since arboviruses are transmitted by the bite of infected mosquitoes between vertebrate hosts,

activity of these pathogens is determined, at least in part, by the blood feeding habits of these

vectors. According to its host feeding pattern range, *Culex* mosquitoes can be regarded as

specialists or generalists. Specialist species include mosquitoes feeding primarily on mammals

(Muturi et al., 2008), birds (Jansen et al., 2009), reptiles and amphibians (Janssen et al., 2015).

Generalists are mosquitoes fed on the most abundant available hosts (Molaei et al., 2007).

Additionally, some species could experience a seasonal shift in its feeding habits, from avian to

mammal hosts (Edman & Taylor, 1968; Hancock & Camp, 2022), being of greater concern from

an epidemiological perspective given that they can act as bridge vectors in the transmission of

some arboviruses, such as SLEV and WNV (Kilpatrick et al., 2006).

West Nile virus (WNV) and St. Louis encephalitis virus (SLEV) are zoonotic pathogens maintained

in nature through an enzootic network involving transmission among viremic birds and

ornithophagic mosquitoes (Kilpatrick et al., 2006). However, during late summer and early

autumn, these viruses increase their activity in humans and other mammals, which has led to

repeated epidemics in the US (Kilpatrick et al., 2006) and South America (Spinsanti et al., 2008).

Likewise, several *Culex* species, including *Cx. quinquefasciatus*, have exhibited a seasonal shift in

their feeding behaviour, feeding primarily on birds in spring and early summer and increasing

the proportion of mammal blood meals in late summer and autumn (Thiemann et al., 2011).

This seasonal shift suggests that viral activity might spill over from birds to mammals as a result

of the mosquito host switch (Edman & Taylor, 1968; Kilpatrick et al., 2006).

Two main contrasting hypotheses have been proposed to explain why primarily ornithophagic

mosquitoes increase their feeding on mammals later in the season. The *migration* hypothesis,

proposed by Kilpatrick et al. (2006) for *Cx. pipiens*, suggests that this shift in feeding pattern is

related to the abundance of the preferred avian host, the American robin (*Turdus migratorius*

L.). This hypothesis is based on their findings that as robin densities decrease, the proportion of

mammalian blood meals in *Cx. pipiens* populations increase. However, this hypothesis has

several limitations: (i) multiple studies have shown that American robins are not always the

preferred host for all *Cx. pipiens* populations (Patrican et al., 2007; Apperson et al., 2002); (ii) in

some areas (e.g., Tennessee) with documented seasonal host shifts, American robins are

present year-round (Savage et al., 2007); and (iii) this explanation applies only to mosquitoes

whose primary host is the American robin, a species found only in the Northern Hemisphere, or

another migratory bird. Therefore, it is not applicable to southern regions, such as Argentina,

where SLEV seasonality is documented (Spinsanti et al., 2008) but American robins are absent.

Because this, it seems that the hypothesis proposed by Kilpatrick et al. (2006) might be habitat

dependent (Caranci, 2010). On the other hand, the *defensive behaviour* hypothesis proposed by

Burkett-Cadena et al. (2011), suggests that host breeding cycles drive the host shift of

mosquitoes. According to this hypothesis, in hosts with parental care, during reproductive

seasons there is a greater investment of energy in assuring the offspring survivorship,

consequently producing an increase in susceptibility of being bitten by mosquitoes as a result of

a decrease in defensive behaviours. This event leads to the detection of peaks of host use during

periods of reproductive investment, in summer for birds and autumn for mammals (Burkett-

Cadena et al., 2011). While this hypothesis was proposed as an alternative to address the

limitations of the migration hypothesis, it also has drawbacks. It is based on a limited range of

host species and assumes a distinct seasonal phenology, which is not present for all mosquito

hosts. Because this, this hypothesis seems to be host dependent.

Explanations for the seasonal shift in mosquito host feeding focused on vector biology are

largely lacking in the literature. Numerous biological factors, such as stress, metabolic rate, and

blood meal source, along with environmental variables like temperature and photoperiod,

influence mosquito physiology and can affect reproductive traits such as fecundity,

development rate, and survivorship. These factors can give rise to new nutritional requirements

that may ultimately lead to seasonal variations in host selection by mosquitoes (Ciota et al.,

2014; Costanzo et al., 2015; Gervasi et al., 2016; Yan et al., 2017, 2018). While several studies

have examined the effects of these biological and environmental variables on mosquito

reproduction, the complex nature of mosquito biology suggests that multiple interactions

among these variables may produce novel responses that are not apparent when considered

individually. The aim of this study was to assess whether there is an interaction between the

source of the host blood meal and seasonality (defined by temperature and photoperiod) on

three reproductive traits of *Culex quinquefasciatus* mosquitoes: fecundity, fertility, and

hatchability. We hypothesize that the interaction between these two variables influence the

reproductive outcomes, potentially leading to a seasonal shift in host selection, driven by a

reproductive advantage. Given the reported seasonal changes in host use by *Cx.*

quinquefasciatus, in autumn we expect a greater number of eggs (fecundity) and larvae (fertility)

in mosquitoes after feeding on a mammal host compared to an avian host, and the opposite

trend in summer.

Materials and methods

Establishment and maintenance of mosquitoes

Egg rafts of *Culex quinquefasciatus* were collected from a drainage ditch at Universidad Nacional

de Córdoba Campus, Córdoba city, in February 2021. Each raft was individually maintained in

plastic containers with one liter of distilled water. The hatched larvae were fed with 100 mg of

liver powder three times per week until pupation. Pupae were then transferred to plastic

emerging cages (21 cm x 12 cm) covered with a tulle-like fabric, containing distilled water but

no food. Adults emerging from each raft were identified morphologically (Darsie, 1985) and

molecularly (Smith & Fonseca, 2004) to ensure they corresponded to *Cx. quinquefasciatus*, since

Cx. pipiens and its hybrids coexist sympatrically in Córdoba city (Branda et al., 2021). All adults

were reared in a cardboard cage of 22.5 liters (28 cm x 36.5 cm) and provided *ad libitum* with a

10% sugar solution soaked in cotton pads placed on plastic cups. For long-term maintenance of

the colony, 24-hour-starved mosquitoes were offered a blood meal from a restrained chicken

twice a month. Four days after feeding, a plastic container with distilled water was placed inside

the cage to allow engorged females to lay egg rafts. Batches of egg were collected and

transferred to plastic containers (30 cm x 25 cm x 7 cm) in a proportion of three rafts per

container, filled with three liters of distilled water. The hatched larvae were also fed with liver

powder at the same proportion described above. Pupae were transferred to emerging cages,

and adult mosquitoes were placed in the final cardboard cage. The colony has been maintained

for several generations in the Insectary of the Instituto de Virología "Dr. J.M. Vanella" (InViV).

Room conditions were controlled at 28 ± 1 °C, with a 12L:12D photoperiod and 70% relative

humidity.

Experimental design

Blood source. For experimental trials, avian hosts (live chicks of the species *Gallus gallus*) and

mammalian hosts (live mice of the species *Mus musculus*, strain C57BL/6) were used to evaluate

the effect of blood meal source on fecundity, fertility, and hatchability. Chicks were generously

donated by the Bartolucci poultry farm (Córdoba, Argentina), while mice were commercially

obtained from the Instituto de Investigación Médica Mercedes y Martín Ferreyra (CONICET -

Universidad Nacional de Córdoba). In each trial, 24-hour-starved adult female mosquitoes were

provided with a blood meal from restrained chicks or mice. Vertebrate hosts were offered to

mosquitoes one hour before the lights were turned off and were kept for 3 hours during 2

consecutive gonotrophic cycles.

The experimental use of animals (mosquitoes and vertebrates) was approved by the ethical

committee at Facultad de Ciencias Médicas, Universidad Nacional de Córdoba (FCM-UNC) in

compliance with the legislation regarding the use of animals for experimental and other

scientific purposes (accession code: CE-2022-00518476-UNC-SCT#FCM).

Seasonality. To assess the effect of seasonality (photoperiod + temperature) on fecundity,

fertility, and hatchability, a "typical" summer and autumn day from Córdoba city was simulated

in an incubator where mosquitoes were housed throughout the entire experiment. Summer

conditions were as follows: $T°_{min} = 22°C$, $T°_{max} = 28°C$, photoperiod = 14L:10D. Autumn conditions

were characterized by: $T°_{min} = 16°C$, $T°_{max} = 22°C$, and photoperiod = 10L:14D. The humidity levels

were maintained at 60-70% for both conditions. Data for simulated conditions were obtained

from Climate Data website (https://es.climate-data.org/).

Feeding trial design. The interaction effect of blood source (chicken and mouse) and seasonality

(autumn and summer) was evaluated during two consecutive gonotrophic cycles (I and II). This

combination produced eight colonies that were established using egg rafts collected from the

maintenance colony: mouse:autumn-I, mouse:autumn-II, chicken:autumn-I, chicken:autumn-II,

mouse:summer-I, mouse:summer-II, chicken:summer-I, chicken:summer-II. This set of 8

colonies was repeated in triplicate (1-3), generating a total of 24 treatments.

The experimental colonies were maintained under controlled conditions to ensure that all adults

were of the same size (Kauffman et al., 2017). Adult mosquitoes were placed in cardboard cages

of 8.3 liters (21 cm x 24 cm) and were provided with *ad libitum* access to a 10% sugar solution,

which was soaked in cotton pads placed on plastic cups.

Feeding trials were conducted at two time points: 5 days post-emergence (first cycle) and 14

days post-emergence (second cycle). Following each blood meal, female mosquitoes were

anesthetized using CO_2 and classified as fully engorged (VI Sella's stage), partially fed (I-V Sella's

stage), or unfed (I Sella's stage), following the classification by Silva Santos et al. (2019). Fully

engorged females were counted and separated into a separate cage to complete their

gonotrophic cycle. Four days after feeding, a cup containing distilled water was placed inside the

cage to facilitate oviposition. After each oviposition cycle, egg rafts were collected, counted, and

transferred to a 12-well plate containing 4 mL of distilled water. They were then photographed

to subsequently determine the number of eggs per raft. Rafts were maintained in each well until

they hatched into L1 larvae, at which point they were preserved with 2 mL of 96% ethanol, and

the number of L1 larvae per raft was counted.

Statistical analysis. The reproductive outputs measured for each experimental colony were

fecundity, fertility, and hatchability. All analyses were performed using R Studio statistic

software, v.4.2.1 (R Core Team, 2022).

Fecundity was defined as the number of eggs per raft and *fertility* as the number of L1 larvae

hatched per raft. To evaluate the effect of blood meal source and seasonality on these two

variables, a Generalized Linear Mixed Model (GLMM; *lme4* package) with negative binomial

error distribution and logarithmic link function was adjusted (Bates et al., 2015). Fecundity and

fertility served as the response variables, while the fixed explanatory variables included blood

source, seasonality, and gonotrophic cycle, all in a three-way interaction. Treatments (n = 24)

were set as a random intercept.

Hatchability (= hatching rate) was defined as the ratio of the number of larvae to the number of

eggs per raft. A Generalized Linear Model (GLM; *MASS* package) was applied using a

quasipoisson error distribution and a logarithmic link function (Venables et al., 2002). The

response variable was the number of larvae, while the explanatory variables included blood

source, seasonality, gonotrophic cycle and replicate as interaction effects. Additionally, the

logarithm of the number of eggs was incorporated as an offset in the model.

Multiple comparisons among treatments were conducted using Tukey's honestly significant

difference test (HSD Tukey), incorporating the Kramer (1956) correction for unbalanced data,

also known as the Tukey-Kramer method. This approach allows setting the family-wise error rate

to 0.05. These comparisons were performed using the *emmeans* package (Lenth, 2024).

Model validation was assessed using a simulation-based approach to generate randomized

quantile residuals to test goodness of fit (Q-Q plot), homoscedasticity, and outliers and

influential points (Cook's distance and Leverage plots). These analyses were conducted with the

DHARMa package (Hartig, 2022).

Additionally, the statistical power of the models was assessed using the *mixedpower* package

(Kumle et al., 2021), specifically designed to evaluate the power of Generalized Linear Mixed

Models (GLMMs). The analysis involved simulating 1,000 iterations with a sample size ranging

from 30 to 150 observations per group. The power analysis aimed to achieve a power threshold

of 0.8, indicating an 80% chance of detecting true effects if they exist.

Results

A total of 1,162 egg rafts were obtained for analysis, distributed across the eight treatments as

follows: chicken:autumn-I: 260, chicken:autumn-II: 174, chicken:summer-I: 162,

chicken:summer-II: 88, mouse:autumn-I: 146, mouse:autumn-II: 96, mouse:summer-I: 167, and

mouse:summer-II: 69. The full count of egg rafts for each replicate (24 treatments), along with

the means for fecundity, fertility, and hatchability, are summarized in Table 1. This sample size

fulfilled the statistical power of 0.8 (Figure SM1).

In both models—fecundity and fertility—the interaction between blood source and seasonality

was statistically significant (fecundity: LRT X^2 = 5.69, $p < 0.05$; fertility: LRT X^2 = 4.37, $p < 0.05$)

(Tables 2,3; Figures 1,2A). Mosquitoes that fed on chicken blood had the highest fecundity

(summer: 136.4 eggs/raft, autumn: 141.5 eggs/raft) and fertility (summer: 123.9 larvae/raft,

autumn: 126 larvae/raft) in both seasons. In summer, fecundity and fertility were 7% and 11%

higher, respectively, in chicken-fed mosquitoes compared to mouse-fed mosquitoes. In autumn,

these differences increased to 46% for fecundity and 52% for fertility in favor of chicken-fed

mosquitoes. No significant seasonal variation was observed in fecundity and fertility for chicken-

fed mosquitoes, but both measures were statistically significant lower by 24% and 25%,

respectively, in autumn compared to summer for mouse-fed mosquitoes.

In the fertility model, there was also a significant interaction between blood source and

gonotrophic cycle (LRT X^2 = 3.98, p < 0.05), indicating that the impact of blood type on larvae

production differed between the first and second cycles (Table 3, Figure 2B). Chicken-fed

mosquitoes showed no difference in fertility between cycles, while those fed on mouse blood

had 25% lower fertility in the second cycle compared to the first. During the second cycle,

fertility was 50% higher in chicken-fed mosquitoes compared to mouse-fed mosquitoes.

Figure 3 presents the predicted mean values of fecundity and fertility for the 24 treatments,

with no differences observed across the three replicates. The variance of random intercepts

among treatments was approximately 0.006 for fecundity and 0.007 for fertility, accounting for

6% and 4% of the total variance in the respective models.

There was no statistical difference among treatments and replicates in terms of hatchability,

given that no statistical effects of blood source, seasonality or gonotrophic cycle was observed

(Table 4). The mean hatchability for all treatments was 0.89, with a range between 0.84 for the

mouse:autumn-II (replicate 1) colony and 0.96 for mouse:autumn-I (replicate 1) (Table 1, Figure

4).

Discussion

Our results show there is an interaction effect between blood meal source and seasonality on

reproductive outputs of *Cx. quinquefasciatus*. While our model hypothesized that fecundity and

fertility would be higher in bird-fed mosquitoes than in mammal-fed mosquitoes during

summer, and the reverse would occur in autumn, our findings revealed the opposite pattern.

Regardless the season, fecundity and fertility in *Culex quinquefasciatus* were higher in bird-fed

mosquitoes compared to those fed on mammals. This pattern aligns with those findings from

previous studies (Akoh et al., 1993; Richards et al., 2012; Telang & Skinner, 2019). This trend is

not unique to *Cx. quinquefasciatus* but has also been seen in other *Culex* species (Shroyer &

Silvery, 1972) and *Aedes* mosquitoes (Harrison et al., 2021). The commonly accepted

explanation for this pattern suggests that the nutritional quality of avian blood, with its

nucleated and larger erythrocytes, provides greater protein and energy, thereby enhancing

these reproductive outputs (Alto et al., 2014). However, our findings offer a mixed perspective

by highlighting the interaction between blood meal source and seasonality that further might

influence fecundity and fertility.

The significant interaction observed between blood meal source and seasonality revealed that

mammal-fed mosquitoes had lower fecundity and fertility in autumn compared to summer,

while bird-fed mosquitoes maintained similar reproductive outputs across both seasons. This

finding diverges from the expectation of consistently higher fecundity and fertility in bird-fed

mosquitoes and consistently lower fecundity and fertility in mammal-fed mosquitoes. This

suggests that environmental factors such as temperature and photoperiod, which have been

shown to affect mosquito fitness (Mordecai et al., 2019; Abouzied, 2017; Mogi, 1992), may

interact with blood meal source in complex ways. Although the effects of temperature and

photoperiod are generally coupled and follow a unimodal trend with an optimal peak at 28°C

(Mordecai et al., 2019), our results indicate that the interaction between seasonality and blood

source might play a critical role in determining reproductive success.

Additionally, our study found a significant interaction between blood meal source and

gonotrophic cycle on fertility. For mammal-fed mosquitoes, fertility was significantly lower in

the second gonotrophic cycle compared to the first, a pattern commonly reported in other

studies where the first and second cycles are generally the most productive (Awahmukalah &

Brooks, 1985; Christiansen-Jucht et al., 2015). This declining fertility pattern has been

documented in *Cx. quinquefasciatus* fed on both birds and mammals (Richards et al., 2012;

Telang & Skinner, 2019). However, our results showed that this decline in fertility was not

present in bird-fed mosquitoes, indicating a potential role of blood meal source in moderating

reproductive output across gonotrophic cycles.

We also investigated whether hatchability could account for the observed variations in fertility

among different treatments. Our results indicate no difference in hatchability among

treatments.

While our findings corroborate an interaction effect between blood meal source and

seasonality, they are not without caveats, as same as other studies. First, our research was

conducted under controlled conditions in a laboratory incubator, which implies that our results

may differ from those of field studies. Some of the main limitations of this approach include the

use of artificial, constant light and temperature rather than a fluctuating environment, and the

use of constant humidity, which could affect mosquito performance. However, this is the first

study of its kind, and future investigations should aim for more realistic setups. A second

limitation, commonly encountered in ecological and behavioral studies, is determining an

appropriate sample size for each treatment, as this can impact the ability to detect desired

effects or interactions (Taborsky, 2010). To address this issue, we conducted an a priori power

analysis to determine the minimum required sample size and repeated the assay multiple times

(three replicates). Furthermore, the presence of interaction effects among our explanatory

variables suggests that our chosen sample size may have been adequate. Lastly, another aspect

of this research that warrants further discussion is the underlying hypothesis that *Culex*

mosquitoes change their feeding habits from birds to mammals. Given that most species are

ornithophagic, why is this shift not directed from one bird host to another? In fact, both

patterns—bird-to-mammal (Burkett-Cadena et al., 2011) and bird-to-bird (Kent et al., 2009)

shifts—are mentioned in the literature. Future studies are aimed to clarify the complex feeding

behaviour of *Culex* mosquitoes regarding seasonal switching, which may depend on several

factors, such as habitat, as evidenced in U.S. *Culex* populations (Caranci, 2010).

In summary, our study investigates the critical feature of host shifting patterns in *Culex*

mosquitoes, which holds significant importance due to their role as bridge vectors between bird

and human hosts. Although our study detects a significant interaction between blood meal

source and seasonality, we observed the opposite expected trend, suggesting a complex

interplay of factors affecting mosquito reproduction and its relationship with host use. We could

consider three possible explanations for our results. Firstly, it is possible that additional factors,

such as genetic mechanisms or variations in midgut microbiota composition play a role in

influencing host use patterns. Secondly, host switching might be explained by factors related to

vector biology other than fitness. Lastly, it remains a possibility that Argentinian *Cx.*

quinquefasciatus populations do not exhibit the same host-switching behaviour observed in US

populations, and the observed interaction effects may be linked to different mosquito

behaviours. However, Spinsanti et al. (2008) found a seasonal variation in WNV human cases,

and Beranek (2019) also observed that in autumn, the proportion of blood meals was higher

than in summer for *Culex quinquefasciatus*. Therefore, more in-field studies are needed to

assess seasonal host variation in *Culex* mosquitoes from Argentina.

Further investigations are warranted to corroborate our findings and gain a deeper

understanding of the interaction between mosquito feeding pattern and seasonality. Last, these

investigations will provide valuable insights into mosquito-borne disease transmission and
forecasting its emergence.

Data availability

319 All data generated and analyzed during the current study are available from the corresponding
320 author upon reasonable request.

Acknowledgments

322 We would like to thank Dr. Agustín Quaglia for providing valuable help in statistical analysis. In
323 addition, we greatly appreciate very helpful comments on the manuscript by anonymous
324 reviewers. This study was supported by grant PICT 2018-1172 (FONCYT) and Consolidar Secyt-
325 UNC (2018-2023).

Competing interests

327 The authors declare no competing interests.

Author contributions

Kevin Alen Rucci, Methodology, Software, Validation, Formal analysis, Investigation, Data
Curation, Writing – Original Draft, Writing – Review & Editing, Visualization; Gabriel Barco,
Investigation, Data Curation, Writing – Review & Editing; Andrea Onorato, Investigation, Data
Curation, Writing – Review & Editing; Mauricio Beranek, Investigation, Data Curation, Writing –
Review & Editing; Mariana Pueta, Conceptualization, Methodology, Validation, Writing – Review
& Editing, Visualization, Supervision; Adrián Diaz, , Conceptualization, Methodology, Validation,
Writing – Review & Editing, Visualization, Supervision, Project administration, Funding
acquisition.

References

Abouzied, E.M. (2017). Life table analysis of *Culex pipiens* under simulated weather conditions

in Egypt. *Journal of the American Mosquito Control Association*, 33(1), 16–24.

https://doi.org/10.2987/16-6608.1

Akoh, J.I., Aigbodion, F.I., & Onyeka, J.O. (1993). Effect of source and size of bloodmeal on egg

production by *Culex quinquefasciatus* (Diptera, Culicidae). *Medical Entomology and Zoology*,

44(1): 41–43. https://doi.org/10.7601/mez.44.41

Alto, B.W., Connelly, C.R., O'Meara, G.F., Hickman, D., & Karr, N. (2014). Reproductive biology

and susceptibility of Florida *Culex coronator* to infection with West Nile virus. *Vector-Borne and*

Zoonotic Diseases, 14(8): 606–614. https://doi.org/10.1089/vbz.2013.1501

Apperson, C.S., Harrison, B.A., Unnasch T.R., Hassan, H.K., Irby, W.S., Savage, H.M., Aspen, S.E.,

Watson, D.W., Rueda, L.M., Engber, B.R., & Nasci, R.S. (2002). Host-feeding habits of *Culex* and

other mosquitoes (Diptera: Culicidae) in the Borough of Queens in New York City, with

characters and techniques for identification of *Culex* mosquitoes. *Journal of Medical*

Entomology, 39(5): 777–85. https://doi.org/10.1603/0022-2585-39.5.777

Awahmukalah, D.S., & Brooks, M.A. (1985). Viability of *Culex pipiens pipiens* eggs affected by

nutrition and aposymbiosis. *Journal of Invertebrate Pathology*, 45(2): 225–230.

https://doi.org/10.1016/0022-2011(85)90012-6

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software*, 67(1): 1–48. https://doi.org/10.18637/jss.v067.i01

Beranek, M. (2019). Mosquitos del género *Culex* (Diptera: Culicidae) como vectores del virus

Saint Louis encephalitis (*Flavivirus*) en ambientes urbanos de la ciudad de Córdoba. PhD Thesis.

Universidad Nacional de Córdoba, Córdoba, Argentina.

Branda, M.F., Laurito, M., Visintin, A.M., & Almirón, W.R. (2021). Gonoactivity of *Culex* (*Culex*)

(Diptera: Culicidae) Mosquitoes During Winter in Temperate Argentina. *Journal of Medical*

Entomology, 58(3): 1454–1458. https://doi.org/10.1093/jme/tjaa295

Burkett-Cadena, N.D., McClure, C.J., Ligon, R.A., Graham, S.P., Guyer, C., Hill, G.E., Ditchkoff, S.S.,

Eubanks, M.D., Hassan, H.K., & Unnasch, T.R. (2011). Host reproductive phenology drives

seasonal patterns of host use in mosquitoes. *PLOS one*, 6(3): e17681.

https://doi.org/10.1371/journal.pone.0017681

Caranci, A. (2010). Host switching in *Culex* mosquitoes: effects of photoperiod and aging on host

choice. PhD Thesis. University of Delaware, Delaware, US.

Ciota, A.T., Matacchiero, A.C., Kilpatrick, A.M., & Kramer, L.D. (2014). The effect of temperature

on life history traits of *Culex* mosquitoes. *Journal of Medical Entomology*, 51(1): 55–62.

https://doi.org/10.1603/ME13003

Costanzo, K.S., Schelble, S., Jerz, K., & Keenan, M. (2015). The effect of photoperiod on life

history and blood-feeding activity in *Aedes albopictus* and *Aedes aegypti* (Diptera: Culicidae).

Journal of Vector Ecology, 40(1): 164–171. https://doi.org/10.1111/jvec.12146

Christiansen-Jucht, C.D., Parham, P.E., Saddler, A., Koella, J.C., & Basáñez, M.G. (2015). Larval

and adult environmental temperatures influence the adult reproductive traits of *Anopheles*

gambiae ss. *Parasites & vectors*, 8: 1–12. https://doi.org/10.1186/s13071-015-1053-5

Darsie, R.F., Jr. (1985). Mosquitoes of Argentina. Part I. Keys for identification of adult females

and fourth stage larvae in English and Spanish (Diptera: Culicidae). *Mosquito Systematics*, 17(3):

153–253.

Edman, J.D., & Taylor, D.J. (1968). *Culex nigripalpus*: Seasonal Shift in the Bird-Mammal Feeding

Ratio in a Mosquito Vector of Human Encephalitis. *Science*, 161(3836): 67–68.

https://doi.org/10.1126/science.161.3836.67

Farajollahi, A., Fonseca, D.M., Kramer, L.D., & Kilpatrick, A.M. (2011). "Bird biting" mosquitoes

and human disease: a review of the role of *Culex pipiens* complex mosquitoes in epidemiology.

Infection, genetics and Evolution, 11(7): 1577–1585.

https://doi.org/10.1016/j.meegid.2011.08.013

Gervasi, S.S., Burkett-Cadena, N., Burgan S.C., Schrey, A.W., Hassan, H.K., Unnasch, T.R., &

Martin, L.B. (2016). Host stress hormones alter vector feeding preferences, success, and

productivity. *Proceedings of the Royal Society B*, 283: 20161278.

https://doi.org/10.1098/rspb.2016.1278

Hancock, C., & Camp, J.V. (2022). Habitat-Specific Host Selection Patterns of *Culex*

quinquefasciatus and *Culex nigripalpus* in Florida. *Journal of the American Mosquito Control*

Association, 38(2): 83–91. https://doi.org/10.2987/21-7054

Harrison, R.E., Brown, M.R., & Strand, M.R. (2021). Whole blood and blood components from

vertebrates differentially affect egg formation in three species of anautogenous mosquitoes.

Parasites & Vectors, 14: 1–19. https://doi.org/10.1186/s13071-021-04594-9

Hartig, F. (2022). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models. R package version 0.4.6.

Jansen, C.C., Webb, C.E., Graham, G.C., Craig, S.B., Zborowski, P., Ritchie, S.A., Russell, R.C., &

van den Hurk, A.F. (2009). Blood sources of mosquitoes collected from urban and peri-urban

environments in eastern Australia with species-specific molecular analysis of avian blood meals.

American Journal of Tropical Medicine and Hygiene, 81(5): 849–857.

https://doi.org/10.4269/ajtmh.2009.09-0008

Janssen, N., Fernandez-Salas, I., Díaz González, E.E., Gaytan-Burns, A., Medina-de la Garza, C. E.,

Sanchez-Casas, R.M., Börster, J., Cadar, D., Schmidt-Chanasit, J., & Jöst, H. (2015).

Mammalophilic feeding behaviour of *Culex quinquefasciatus* mosquitoes collected in the cities

of Chetumal and Cancun, Yucatán Peninsula, Mexico. *Tropical Medicine & International Health*,

20(11): 1488-1491. https://doi.org/10.1111/tmi.12587

Kasili, S., Oyieke, F., Wamae, C., & Mbogo, C. (2009). Seasonal changes of infectivity rates of

Bancroftian filariasis vectors in Coast Province, Kenya. *Journal of Vector Borne Diseases*, 46: 219–

224.

Kauffman, E., Payne, A., Franke, M.A., Schmid, M.A., Harris, E., & Kramer, L.D. (2017). Rearing of

Culex spp. and *Aedes* spp. mosquitoes. *Bio-protocol*, 7(17): e2542-e2542.

https://doi.org/10.21769/BioProtoc.2542

Kent, R.J. (2009). Molecular methods for arthropod bloodmeal identification and applications to

ecological and vector-borne disease studies. Molecular Ecology Resources, 9(1): 4–18.

https://doi.org/10.1111/j.1755-0998.2008.02469.x

Kilpatrick, A.M., Kramer, L.D., Jones, M.J., Marra, P.P. & Daszak, P. (2006). West Nile Virus

Epidemics in North America Are Driven by Shifts in Mosquito Feeding Behavior. *PLoS Biology*,

4(4): e82. https://doi.org/10.1371/journal.pbio.0040082

Kramer, C.Y. (1956). Extension of multiple range tests to group means with unequal numbers of

replications. *Biometrics*, 12: 307–310. https://doi.org/10.2307/3001469

Kumle, L., Võ, M.L., & Draschkow, D. (2021). Estimating power in (generalized) linear mixed

models: an open introduction and tutorial in R. *Behavior Research Methods*, 53(6): 2528–243.

https://doi.org/10.3758/s13428-021-01546-0

Labarthe, N., Serrão, M.L., Melo, Y., José-Oliveira, S., and Lourenço-de-Oliveira, R. (1998).

Potential vectors of *Dirofilaria immitis* in Itacoatiara, Oceanic Region of Niterói Municipality,

state of Rio de Janeiro, Brazil. *Memorias do Instituto Oswaldo Cruz*, 93: 425–432.

https://doi.org/10.1590/S0074-02761998000400001

Lenth R (2024). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.10.3.

Mogi, M. (1992). Temperature and photoperiod effects on larval and ovarian development of

New Zealand strains of *Culex quinquefasciatus* (Diptera: Culicidae). *Annals of the Entomological*

Society of America, 85(1), 58–66. https://doi.org/10.1093/aesa/85.1.58

Molaei, G., Andreadis, T.G., Armstrong, P.M., Bueno, R., Dennett, J.A., Real, S.V., Sargent, C.,

Bala, A., Randle, Y., Guzman, H., da Rosa, A.T., Wuithranyagool & Tesh, R. B. (2007). Host feeding

pattern of *Culex quinquefasciatus* (Diptera: Culicidae) and its role in transmission of West Nile

virus in Harris County, Texas. *American Journal of Tropical Medicine and Hygiene*, 77(1): 73–81.

https://doi.org/10.4269/ajtmh.2007.77.73

Mordecai, E.A., Caldwell, J.M., Grossman, M.K., Lippi, C.A., Johnson, L.R., Neira, M., Rohr, J.R.,

Ryan, S.J., Savage, V., Shocket, M.S., Sippy, R., Stewart Ibarra, A.M, Thomas, M.B., & Villena, O.

(2019). Thermal biology of mosquito-borne disease. *Ecology letters*, 22(10): 1690–1708.

https://doi.org/10.1111/ele.13335

Muturi, E.J., Muriu, S., Shililu, J., Mwangangi, J.M., Jacob, B.G., Mbogo, C., Githure, J. & Novak,

R.J. (2008). Blood-feeding patterns of *Culex quinquefasciatus* and other culicines and

implications for disease transmission in Mwea rice scheme, Kenya. *Parasitology Research*,

102(6): 1329–1335. https://doi.org/10.1007/s00436-008-0914-7

Patrican, L.A., Hackett, L.E., Briggs, J.E., McGowan, J.W., Unnasch, T.R., & Lee, J.H. (2007). Host-

feeding patterns of *Culex* mosquitoes in relation to trap habitat. *Emerging Infection Diseases*,

13(12): 1921–19233. https://doi.org/10.3201/eid1312.070275

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Richards, S.L., Anderson, S.L., & Yost, S.A. (2012). Effects of blood meal source on the

reproduction of *Culex pipiens quinquefasciatus* (Diptera: Culicidae). *Journal of Vector Ecology*,

37(1): 1–7. https://doi.org/10.1111/j.1948-7134.2012.00194.x

Savage, H.M., Aggarwal, D., Apperson, C.S., Katholi, C.R., Gordon, E., Hassan, H.K., Anderson, M.,

Charnetzky, D., McMillen, L., Unnasch, E.A., & Unnasch, T.R. (2007). Host choice and West Nile

virus infection rates in blood-fed mosquitoes, including members of the *Culex pipiens* complex,

from Memphis and Shelby County, Tennessee, 2002-2003. *Vector-Borne and Zoonotic Diseases*,

7(3): 365–386. https://doi.org/10.1089/vbz.2006.0602

Shroyer, D.A., & Silvery, R.E. (1972). A comparison of egg production of *Culex pipiens pipiens* L.

fed on avian and mammalian hosts. *Mosquito News*, 32(4): 636–637.

Silva Santos, C., Pie, M.R., da Rocha, T.C., & Navarro-Silva, M.A. (2019). Molecular identification

of blood meals in mosquitoes (Diptera, Culicidae) in urban and forested habitats in southern

Brazil. *PloS one*, 14(2): e0212517. https://doi.org/10.1371/journal.pone.0212517

Smith, J.L. & Fonseca, D.M. (2004). Rapid assays for identification of members of the *Culex*

(*Culex*) *pipiens* complex, their hybrids, and other sibling species (Diptera: Culicidae). *The*

American Journal of Tropical Medicine and Hygiene, 40(4): 339–345.

https://doi.org/10.4269/ajtmh.2004.70.339

Spinsanti, L.I., Díaz, L.A., Glastein, N., et al. (2008). Human outbreak of St. Louis encephalitis

detected in Argentina, 2005. *Journal of Clinical Virology*, 42: 27–33.

https://doi.org/10.1016/j.jcv.2007.11.022

Taborsky, M. (2010). Sample size in the study of behaviour. *Ethology*, 116(3): 185-202.

https://doi.org/10.1111/j.1439-0310.2010.01751.x

Telang, A., & Skinner, J. (2019). Effects of host blood meal source on reproductive output,

nutrient reserves and gut microbiome of West Nile virus vector *Culex quinquefasciatus*. *Journal*

of Insect Physiology, 114: 15–22. https://doi.org/10.1016/j.jinsphys.2019.02.001

Thiemann, T.C., Wheeler, S.S., Barker, C.M., Reisen, W.K. (2011). Mosquito Host Selection Varies

Seasonally with Host Availability and Mosquito Density. *PLOS Neglected Tropical Diseases*, 5(12):

e1452. https://doi.org/10.1371/journal.pntd.0001452

Venables, W.N., & Ripley, B.D. (2002). Exploratory multivariate analysis. In: Modern applied

statistics with S (pp. 301-330). Springer, New York.

https://doi.org/10.1007/978-0-387-21706-2_11

Yan, J., Gangoso, L., Martínez-de la Puente, J., Soriguer, R. & Figuerola, J. (2017). Avian

phenotypic traits related to feeding preferences in two *Culex* mosquitoes. *The Science of Nature*,

104(9): 1–10. https://doi.org/10.1007/s00114-017-1497-x

Yan, J., Broggi, J, Martínez-de la Puente, J., Gutiérrez-López, R., Gangoso, L., Soriguer, R. &

Figuerola, J. (2018). Does Bird metabolic rate influence mosquito feeding preference? *Parasites*

& Vectors, 11(1): 1–9. https://doi.org/10.1186/s13071-018-2708-9

Table 1. Reproductive outcomes of *Culex quinquefasciatus* indicating average fecundity fertility and hatchability for both gonotrophic, accompanied with its respective standard deviation.

Replicate	Blood source	Season	Gonotrophic cycle	No. engorged females	No. egg rafts	Fecundity (eggs/raft)	Fertility (larvae/raft)	Hatchability (larvae/eggs)
1	chicken	autumn	I	163	76	147.47 ± 31.84	133.24 ± 38.19	0.89 ± 0.14
1	chicken	summer	I	116	74	125.64 ± 39.57	109.39 ± 43.66	0.85 ± 0.19
1	mouse	autumn	I	58	23	124.48 ± 20.85	119.96 ± 21.47	0.96 ± 0.04
1	mouse	summer	I	208	120	150.67 ± 29.41	140.38 ± 33.15	0.93 ± 0.10
1	chicken	autumn	II	73	36	142.14 ± 36.59	129.09 ± 43.62	0.87 ± 0.18
1	chicken	summer	II	38	27	138.23 ± 30.27	117.27 ± 37.99	0.85 ± 0.18
1	mouse	autumn	II	23	15	89.13 ± 19.33	73.29 ± 28.09	0.84 ± 0.24
1	mouse	summer	II	92	45	106.68 ± 31.97	99.16 ± 31.18	0.93 ± 0.10
2	chicken	autumn	I	178	89	138.23 ± 37.45	122.7 ± 41.55	0.90 ± 0.15
2	chicken	summer	I	78	51	128.75 ± 38.9	126.08 ± 33.97	0.89 ± 0.13
2	mouse	autumn	I	110	69	103.04 ± 19.01	91.32 ± 22.96	0.89 ± 0.16
2	mouse	summer	I	36	27	113.26 ± 30.66	103.83 ± 34.81	0.89 ± 0.20
2	chicken	autumn	II	86	66	127.4 ± 37.07	116.35 ± 39.9	0.89 ± 0.16
2	chicken	summer	II	48	34	143.68 ± 28.06	136.75 ± 28.05	0.92 ± 0.11
2	mouse	autumn	II	63	52	88.87 ± 19.65	77.86 ± 22.88	0.88 ± 0.15
2	mouse	summer	II	18	14	78.29 ± 25.77	67 ± 25.79	0.87 ± 0.18
3	chicken	autumn	I	150	96	147.22 ± 33.4	129.33 ± 39.45	0.88 ± 0.16
3	chicken	summer	I	67	38	145.03 ± 44.39	136.03 ± 45.49	0.90 ± 0.14
3	mouse	autumn	I	92	54	96.04 ± 23.78	85.76 ± 27.45	0.89 ± 0.18
3	mouse	summer	I	27	20	104.3 ± 21.42	98.35 ± 23.86	0.93 ± 0.09
3	chicken	autumn	II	94	73	144.07 ± 37.49	125.23 ± 39.42	0.87 ± 0.17
3	chicken	summer	II	35	28	156.36 ± 26.71	141 ± 39.53	0.90 ± 0.18
3	mouse	autumn	II	42	29	79.66 ± 19.94	68.76 ± 23.73	0.87 ± 0.21
3	mouse	summer	II	17	11	100.36 ± 22.82	96.1 ± 24.06	0.94 ± 0.06

508 **Table 2.** Analysis of deviance table for the generalized linear mixed model examining the effects
509 of blood source, seasonality, gonotrophic cycle, and their interactions on the fecundity
510 (eggs/raft) of *Culex quinquefasciatus* mosquitoes across 3 replicates. A random intercept for
511 treatment was included to account for variability among the 24 treatments (variance =
512 0.006169). Abbreviations: LRT X^2 = likelihood-ratio test; df = degrees of freedom.

Factor	LRT X^2	df	p-value
blood source	19.22	1	1.16×10^{-5}
seasonality	1.50	1	0.22
gonotrophic cycle	0.47	1	0.49
blood source × seasonality	5.69	1	0.02
blood source × gonotrophic cycle	2.74	1	0.1
seasonality × gonotrophic cycle	2.07	1	0.15
blood source × seasonality × gonotrophic cycle	1.65	1	0.20

513 **Table 3.** Analysis of deviance table for the generalized linear mixed model examining the effects
514 of blood source, seasonality, gonotrophic cycle, and their interactions on the fertility
515 (larvae/raft) of *Culex quinquefasciatus* mosquitoes across 3 replicates. A random intercept for
516 treatment was included to account for variability among the 24 treatments (variance =
517 0.007226). Abbreviations: LRT X^2 = likelihood-ratio test; df = degrees of freedom.

Factor	LRT X^2	df	p-value
blood source	13.30	1	0.00026
seasonality	0.36	1	0.54
gonotrophic cycle	0.28	1	0.59
blood source × seasonality	4.37	1	0.03
blood source × gonotrophic cycle	3.98	1	0.04
seasonality × gonotrophic cycle	0.92	1	0.34
blood source × seasonality × gonotrophic cycle	0.50	1	0.47

518 **Table 4.** Analysis of deviance table for the generalized linear model examining the effects of
519 blood source, seasonality, gonotrophic cycle, replicate and their interactions on the hatchability
520 (larvae/eggs) of *Culex quinquefasciatus* mosquitoes. Abbreviations: LRT X^2 = likelihood-ratio test;
521 df = degrees of freedom.

Factor	LRT X^2	df	p-value
blood source	2.50	1	0.11
seasonality	3.06	1	0.08
gonotrophic cycle	0.005	1	0.94
replicate	1.31	1	0.51
blood source × seasonality	0.12	1	0.72
blood source × gonotrophic cycle	3.64	1	0.06
seasonality × gonotrophic cycle	0.07	1	0.79
blood source × replicate	2.42	1	0.30
seasonality × replicate	2.96	1	0.23
gonotrophic cycle × replicate	0.07	1	0.96
blood source × seasonality × gonotrophic cycle	2.73	1	0.10
blood source × seasonality × replicate	0.02	1	0.99
blood source × gonotrophic cycle × replicate	2.48	1	0.29
seasonality × gonotrophic cycle × replicate	0.43	1	0.81
blood source × seasonality × gonotrophic cycle × replicate	2.57	1	0.28

Figure captions

Figure 1. Interaction plot showing the effect of blood meal source (chicken or mouse) and seasonality (autumn or summer) on fecundity (eggs/raft). Points represent the predicted marginal mean values and the bars corresponding to their confidence intervals. Means sharing the same letters are not statistically different (p > 0.05).

Figure 2. Interaction plots showing the effect of blood meal source (chicken and mouse) and seasonality (autumn and summer) (**A**) or blood meal source and gonotrophic cycle (first or second) (**B**) on fertility (eggs/raft) of *Culex quinquefasciatus*. Points represent the predicted marginal mean values and the bars corresponding to their confidence intervals. Means sharing the same letters are not statistically different (p > 0.05).

Figure 3. Dotplots showing the variation of predicted fecundity (eggs/raft) and fertility (larvae/raft) means along the 24 treatments, combination of blood source (chicken or mouse), seasonality (autumn or summer), gonotrophic cycle (first or second) and replicate (1 to 3).

Figure 4. Boxplots showing the mean hatchability for the 24 treatments, combining replicate (1-3), blood meal source (chicken and mouse), seasonality (summer and autumn) and gonotrophic cycle (first and second), of *Culex quinquefasciatus*.

Supplementary Material.

Figure SM1. Variation of statistical power for the fecundity and fertility models across several sample sizes.