Effects of blood meal source and seasonality on reproductive traits of

***Culex quinquefasciatus* (Diptera: Culicidae)**

Kevin Alen Rucci[1,3,*], Mariana Pueta[2,3] & Adrián Díaz[1,3*]

[1] Laboratorio de Arbovirus, Instituto de Virología "Dr. J. M. Vanella" (InViV), Facultad de

Ciencias Médicas (FCM), Universidad Nacional de Córdoba (UNC)

[2] Instituto de Investigaciones en Biodiversidad y Medioambiente (INIBIOMA), CONICET -

Universidad Nacional de Comahue (UNCo)

[3] Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina

*Corresponding authors:

Kevin Alen Rucci

Laboratorio de Arbovirus – Instituto de Virología "Dr. J. M. Vanella" (InViV), Facultad de

Ciencias Médicas (FCM), Universidad Nacional de Córdoba (UNC).

Enfermera Gordillo Gómez s/n. Ciudad Universitaria, X5000HUA, Córdoba capital, Córdoba,

Argentina.

E-mail addresses: kevin.rucci@mi.unc.edu.ar, kevinrucci@outlook.com

Adrián Díaz

Laboratorio de Arbovirus – Instituto de Virología "Dr. J. M. Vanella" (InViV), Facultad de

Ciencias Médicas (FCM), Universidad Nacional de Córdoba (UNC).

Enfermera Gordillo Gómez s/n. Ciudad Universitaria, X5000HUA, Córdoba capital, Córdoba,

Argentina.

E-mail addresses: adrian.diaz@fcm.unc.edu.ar, adrian.diaz@conicet.gov.ar

Abstract

Host selection by mosquitoes is a keystone to understanding viral circulation and predicting future infection outbreaks. *Culex* mosquitoes frequently feed on birds during spring and early summer, shifting into mammals towards late summer and autumn. This host switch may be due to changes in mosquito fitness. The aim of this study was to assess if the interaction effect of blood meal source and seasonality may influence reproductive traits of *Culex quinquefasciatus* mosquitoes. For this purpose, *Cx. quinquefasciatus* mosquitoes were reared in simulated summer and autumn conditions and fed on two different hosts, chickens and mice, in a factorial design. Fecundity, fertility, and hatchability during two consecutive gonotrophic cycles were estimated. We found greater fecundity and fertility for mosquitoes fed upon birds than mammals. Fecundity and fertility increased in autumn for chicken-fed mosquitoes, whereas they decreased for mouse-fed mosquitoes. These traits decreased in the second gonotrophic cycle for mouse-fed mosquitoes, whereas they did not vary between cycles for chicken-fed mosquitoes. The effect of hatchability on fertility was rather limited. These results indicate a statistically significant interaction effect of blood meal source and seasonality on fecundity and fertility. However, the pattern was opposite in relation to our hypothesis, suggesting that further studies are needed to confirm and expand our knowledge about mosquito biology and its relationship with seasonal host use shifting.

Key words: mosquito vector, fecundity, fertility, hatchability

Introduction

The southern house mosquito, *Culex quinquefasciatus* Say (Diptera: Culicidae), is a worldwide distributed vector of several pathogens, such as *Wuchereria bancrofti* Cobbold (Kasili et al., 2009), *Dirofilaria immitis* Leidy (Labarthe et al., 1998) and avian malaria (Farajollahi et al., 2011). Besides, this mosquito is also one of the main species responsible for the transmission of the arboviruses West Nile virus (WNV) and Saint Louis encephalitis virus (SLEV), among others (Farajollahi et al., 2011). Because of this, this species is of critical concern due to its impact on both public and veterinary health.

Since arboviruses are transmitted by the bite of infected mosquitoes between vertebrate hosts, activity of these pathogens is determined, at least in part, by the blood feeding habits of these vectors. Host preference of *Culex* mosquitoes is variable and it can be classified in four categories: (1) mammophilic (Muturi et al., 2008), (2) ornithophilic (Jansen et al., 2009), (3) herpetophilic (Janssen et al., 2015) and (4) generalist (Molaei et al., 2007). Additionally, some species may experience a host switch in its feeding habits, from avian to mammal hosts (Hancock & Camp, 2022). These species are of greater concern from an epidemiological standpoint because they can act as bridge vectors in the transmission of some arboviruses, such as SLEV and WNV (Kilpatrick et al., 2005).

Culex quinquefasciatus is generally considered an ornithophilic species in several areas around the globe (Takken & Verhulst, 2013). Despite its ornithophilic nature, this and other *Culex* species sometimes feed primarily on mammal hosts, including humans, in certain situations. Various studies have pointed out that in autumn there is an increased activity of SLEV and WNV in human populations (Kilpatrick et al., 2006; Spinsanti et al., 2008), suggesting that viral activity spills over to mammals from birds as a result of the mosquito host switch (Edman & Taylor, 1968; Kilpatrick et al., 2006). A few hypotheses have attempted to explain why primarily ornithophilic mosquitoes also feed upon mammals in some contexts. One of the

most widely accepted hypothesis states that the shift on feeding behaviour of *Culex* mosquitoes is a consequence of the autumn migration of American robin (*Turdus migratorius*), the principal host in the northeastern US, generating an opportunistic change into mammal hosts (Kilpatrick et al., 2006). In South America in general and in Argentina in particular, studies regarding host switch are scarce. Nevertheless, a few limited studies have indicated a seasonal variation throughout the seasons (Stein et al., 2013; Beranek, 2018). Despite that, Argentina has acknowledged seasonal variation in the activity of SLEV in humans, which can be attributed to variations in mosquito host-feeding pattern (Spinsanti et al., 2008). In Southern latitudes, species such as *T. migratorius* are lacking, hence the "migration hypothesis" is very unlikely for many austral areas, being restricted only for some regions of the US.

A second hypothesis proposed by Burkett-Cadena et al. (2011) suggests that host breeding cycles drive the host shift of mosquitoes. According to this hypothesis, in hosts with parental care, during reproductive seasons there is a greater investment of energy in assuring the offspring survivorship, consequently producing an increase in susceptibility of being bitten by mosquitoes as a result of a decrease in defensive behaviours. This event leads to the detection of peaks of host use during periods of reproductive investment, in summer for birds and autumn for mammals (Burkett-Cadena et al., 2011).

While some other minor hypotheses, unsupported by field data, have been proposed, none of them have been specifically focused on vector biology. Numerous biological factors, including stress, metabolic rate, and the source of a blood meal, as well as environmental variables such as temperature and photoperiod, exert an influence on mosquito physiology that might affect various reproductive traits such as fecundity, development rate or survivorship. This set of factors gives rise to new nutritional requirements that, ultimately, may lead to seasonal variation in host choice by mosquitoes (Ciota et al., 2014; Costanzo et al., 2015; Gervasi et al., 2016; Yan et al. 2017, 2018). In the literature, several studies have evaluated the impact of

these biological and environmental variables on mosquito reproduction, as mentioned before.

However, due to the complex nature of mosquito biology, there are likely multiple interactions

among variables that give rise to new responses that may not be observed individually.

Therefore, the aim of the present study was to assess whether exists an interaction effect

between the source of the host blood meal and seasonality (in terms of temperature and

photoperiod) on three reproductive traits of *Culex quinquefasciatus* mosquitoes, fecundity,

fertility, and hatchability. Our hypothesis states that the interaction between these two

variables influence the reproductive outcomes, potentially leading to a seasonal shift in host

selection, driven by a reproductive advantage. Considering the reported host preference

changes in *Cx. quinquefasciatus*, in autumn we expect a greater number of eggs (fecundity)

and larvae (fertility) in mosquitoes after feeding on a mammal host compared to an avian host,

and the opposite relationship in summer.

Materials and methods

Establishment and maintenance of mosquitoes

Egg rafts of *Culex quinquefasciatus* were collected from a drainage ditch at Universidad

Nacional de Córdoba Campus, Córdoba city, in February 2021. Each raft was individually

maintained in plastic containers with one liter of distilled water. The hatched larvae were fed

with 100 mg of liver powder three times per week until pupation. Pupae were then transferred

to plastic emerging cages (21 cm x 12 cm) covered with a tulle-like fabric, containing distilled

water but no food. Adults emerging from each raft were identified morphologically (Darsie,

1985) and molecularly (Smith & Fonseca, 2004) to ensure they corresponded to *Cx.*

quinquefasciatus, since *Cx. pipiens* and its hybrids are also present in Córdoba (Branda et al.,

2021). All adults were reared in a cardboard cage of 22.5 liters (28 cm x 36.5 cm) and provided

ad libitum with a 10% sugar solution soaked in cotton pads placed on plastic cups. For long-

term maintenance of the colony, 24-hour-starved mosquitoes were offered a blood meal from

a restrained chicken twice a month. Four days after feeding, a plastic container with distilled

water was placed inside the cage to allow engorged females to lay egg rafts. Batches of egg

were collected and transferred to plastic containers (30 cm x 25 cm x 7 cm) in a proportion of 3

rafts per container, filled with three liters of distilled water. The hatched larvae were also fed

with liver powder at the same proportion described above. Pupae were transferred to

emerging cages, and adult mosquitoes were placed in the final cardboard cage. The colony has

been maintained for over twenty generations in the Insectary of the Instituto de Virología "Dr.

J.M. Vanella" (InViV). Room conditions were controlled at 28°C, with a 12L:12D photoperiod

and 70% relative humidity.

Experimental design

Blood source. For experimental trials, avian hosts (live chicks of the species *Gallus gallus*) and

mammalian hosts (live mice of the species *Mus musculus*, strain C57BL/6) were used to

evaluate the effect of blood meal source on fecundity, fertility, and hatchability. Chicks were

generously donated by the Bartolucci poultry farm (Córdoba, Argentina), while mice were

commercially obtained from the Instituto de Investigación Médica Mercedes y Martín Ferreyra

(CONICET - Universidad Nacional de Córdoba). In each trial, 24-hour-starved adult female

mosquitoes were provided with a blood meal from restrained chicks or mice. Vertebrate hosts

were offered to mosquitoes one hour before the lights were turned off and were kept for 3

hours during 2 consecutive gonotrophic cycles.

The experimental use of animals (*G. gallus* and *M. musculus*) was approved by the ethical

committee at Facultad de Ciencias Médicas, Universidad Nacional de Córdoba (FCM-UNC) in

compliance with the legislation regarding the use of animals for experimental and other

scientific purposes (accession code: CE-2022-00518476-UNC-SCT#FCM).

Seasonality. To assess the effect of seasonality (photoperiod + temperature) on fecundity,

fertility, and hatchability, a "typical" summer and autumn day from Córdoba city was

simulated in an incubator, where mosquitoes were housed throughout the entire experiment.

Summer conditions were as follows: $T°_{min}$ = 22°C, $T°_{max}$ = 28°C, photoperiod = 14L:10D. Autumn

conditions were characterized by: $T°_{min}$ = 16°C, $T°_{max}$ = 22°C, and photoperiod = 10L:14D. The

humidity levels were maintained at 60-70% for both conditions. Data for simulated conditions

were obtained from Climate Data website (https://es.climate-data.org/).

Feeding trial design. The interaction effect of blood source (bird or mammal) and seasonality

(autumn and summer) was evaluated during two consecutive gonotrophic cycles (I and II).

Eight experimental colonies were established using egg rafts collected from the maintenance

colony: mammal-autumn-I (MA-I), mammal-summer-I (MS-I), bird-autumn-I (BA-I), bird-

summer-I (BS-I), mammal-autumn-II (MA-II), mammal-summer-II (MS-II), bird-autumn-II (BA-

II), and bird-summer-II (BS-II).

The experimental colonies were maintained under controlled conditions to ensure that all

adults were of the same size (Kauffman et al., 2017). Adult mosquitoes were placed in

cardboard cages of 8.3 liters (21 cm x 24 cm) and were provided with *ad libitum* access to a

10% sugar solution, which was soaked in cotton pads placed on plastic cups.

Feeding trials were conducted at two time points: 5 days post-emergence (first cycle) and 14

days post-emergence (second cycle). Following each blood meal, female mosquitoes were

anesthetized using CO_2 and classified as fully engorged (VI Sella's stage), partially fed (I-V

Sella's stage), or unfed (I Sella's stage), following the classification by Silva Santos et al. (2019).

Fully engorged females were counted and separated into a separate cage to complete their

gonotrophic cycle. Four days after feeding, a cup containing distilled water was placed inside

the cage to facilitate oviposition. After each oviposition cycle, egg rafts were collected,

counted, and transferred to a 12-well plate containing 3 mL of distilled water. They were then

photographed to subsequently determine the number of eggs per raft. Rafts were maintained

in each well until they hatched into L1 larvae, at which point they were preserved with 1 mL of

96% ethanol, and the number of L1 larvae per raft was counted.

Statistical analysis. The reproductive outputs measured for each experimental colony were

fecundity, fertility, and hatchability. All analyses were performed using R Studio statistic

software, v.4.2.1 (R Core Team, 2022).

Fecundity was defined as the number of eggs per raft and *fertility* as the number of L1 larvae

hatched per raft. To evaluate the effect of blood meal source and seasonality on these two

variables, a Generalized Linear Model (GLM; package *MASS*) with negative binomial error

distribution and logarithmic link function was adjusted (Venables & Ripley, 2002). Fecundity

and fertility served as the response variables, while the explanatory variables included blood

source, seasonality, and gonotrophic cycle, all in a three-way interaction.

Hatchability (= hatching rate) was defined as the ratio of the number of larvae to the number

of eggs per raft. A GLM was applied using a quasipoisson error distribution and a logarithmic

link function. The response variable was the number of larvae, while the explanatory variables

included blood source, seasonality and gonotrophic cycle as interaction effects. Additionally,

the logarithm of the number of eggs was incorporated as an offset in the model.

Multiple comparisons among treatments were conducted using the Tukey's honestly

significant difference test (HSD Tukey), incorporating the Kramer (1956) correction for

unbalanced data, also known as the Tukey-Kramer method. This approach allows to set the

family-wise error rate to 0.05. These contrasts were performed using the *multcomp* package

(Hothorn et al., 2008).

Model validation was assessed using a simulation-based approach to generate randomized

quantile residuals to test goodness of fit (Q-Q plot), homoscedasticity, and outliers and

influential points (Cook's distance and Leverage plots). These analyses were conducted with

the DHARMa package (Hartig, 2022).

Model accuracy via simulations for the fecundity and fertility models

To assess the accuracy of the results, a simulation approach was conducted using a Monte

Carlo algorithm to simulate data and model parameters according to the methodology

proposed by Carsey & Harden (2013). For this purpose, parameters including coefficients,

treatment means and dispersion parameter (referred to as theta, θ), were extracted from the

fecundity and fertility models. Initially, a new dataset consisting of 1000 values (sample size of

125 per treatment) was generated. This was achieved using the *rnbinom* function from the

MASS package, specifying the treatment means (computed from model coefficients) and the

theta parameter. Subsequently, a negative binomial GLM was fitted to the simulated data, and

the expected means for each of the eight treatments were obtained from the model. This

simulation process was repeated 1000 times. The ratio between the number of iterations in

which the simulated means fell within the observed 95% confidence interval divided by the

total number of simulations was computed as an approximation of the robustness of results.

Results

Effect of blood source and seasonality on fecundity and fertility

Two interaction effects were found to be statistically significant in both the fecundity and

fertility models (Tables 1-4). The first interaction effect was found between blood source and

seasonality (fecundity: LRT X^2 = 25.0, p = 7.72 x 10^{-7}; fertility: LRT X^2 = 13.82, p = 0.002),

meaning that the effect of blood meal source on the number of eggs and larvae per raft varied

with season (Tables 2, 3). In general, mosquitoes that fed on avian blood sources exhibited

greater fecundity and fertility than those that fed on mammalian blood source, regardless of

the season. When treatments were examined by pairwise comparisons, fecundity and fertility

were 13% and 18% greater in autumn compared to summer for bird-fed mosquitoes, while

they were 17% and 19% lower for mammal-fed mosquitoes (Fig. 1 A, C, Table 1). The second

statistically significant interaction was between blood meal source and the gonotrophic cycle

(fecundity: LRT X^2 = 9.29, p = 0.0023; fertility: LRT X^2 = 14.4, p = 0.00073). This indicated that

the effect of blood type on the number of eggs and larvae per raft differed between the first

and second feeding trial (Tables 2, 3). For mosquitoes that fed on mammalian blood, fecundity

and fertility were 29% lower in the second cycle compared to the first. Although the difference

was not statistically significant, fecundity and fertility showed a slight increase in the second

cycle for bird-fed mosquitoes. During the second cycle, these outcomes were 41% and 40%

higher, respectively, for bird-fed mosquitoes compared to mammal-fed mosquitoes (Fig. 1 B,

D, Table 1).

Effect of blood source and seasonality on hatchability

The overall hatchability reached 86%, with a range spanning from 76% for MA-II to a peak of

96% for MA-I, as outlined in Table 1 and depicted in Figure 2. The model found a statistically

significant effect of blood meal source (LRT X^2 = 73.06, p = 0.0012) and an interaction effect

between seasonality and gonotrophic cycle (LRT X^2 = 41.02, p = 0.0015) on hatching rates

(Fig. 3, Table 4). Pairwise comparisons revealed a statistically significant 15% decrease in

hatchability for mosquitoes from autumn from the first to the second gonotrophic cycle.

Additionally, mosquitoes fed on mammals had greater hatchability than those fed on birds. In

summer, hatchability during the first cycle was also higher than during the second cycle. These

two trends were not found to be statistically significant.

Model accuracy via simulations for the fecundity and fertility models

When comparing the simulated means for each of the eight treatments (combinations of blood

meal source, seasonality, and gonotrophic cycle) with the means calculated from the observed

data a high level of accuracy exceeding 90% was observed. Most of the simulated means fitted

within the original 95% confidence interval, with an accuracy rate greater than 98%. The

greatest mismatching among simulated and original data were found in the treatment MS-I,

with an accuracy of 93% and 95% for the fecundity and fertility model, respectively. Accuracy

remained high despite variations in sample sizes and data dispersion (Figs. 4, 5).

Discussion

Our results revealed that there exists an interaction effect of blood meal source and

seasonality on reproductive outputs of *Cx. quinquefasciatus*. Although we expected a higher

fecundity and fertility in summer for bird-fed mosquitoes and higher outputs in autumn for

mammal-fed mosquitoes, we found the opposite pattern.

Most studies have reported a greater fecundity and fertility in *Cx. quinquefasciatus* fed on

different bird species (Akoh et al.,1993; Richards et al., 2012; Telang & Skinner,2019). This

trend is also present in other species of *Culex* (Shroyer & Silvery, 1972) and *Aedes* mosquitoes

(Harrison et al., 2021). The widely accepted hypothesis that has emerged to explain why avian

blood produces a greater fecundity and fertility even in mosquitoes with a known mammal-

preferred host choice focuses on the nutritional quality of host blood. Usually, it is addresses

that because avian blood possesses nucleated and larger erythrocytes this offers greater

quantities of proteins and energy than mammalian blood (Alto et al., 2014). However, this

assumption lacks solid evidence due to studies regarding nutritional quality of avian blood and

its impact in mosquito fitness are scarce. Actually, some studies have shown that mammalian

blood are nutritionally richer than avian blood in terms of total protein content (Pearce 1977,

Sant'Anna et al., 2010). Pre-diuresis, a physiological process where red blood cells and proteins

are concentrated in the midgut by excreting large amounts of water, has been postulated as a

mechanism for explaining the higher fecundity and fertility observed in bird-fed mosquitoes

(Erram et al., 2022). Furthermore, avian blood has higher concentrations of isoleucine than

mammalian blood, a crucial limiting amino acid necessary for egg production (Greenberg,

1951, Harrison et al., 2021).

The effect of temperature on fitness follows a unimodal relationship with a narrow optimal

range. For mosquitoes of genus *Culex* and *Aedes*, this optimal band ranges from 20°C to 30°C,

with an optimal at 28°C for species such as *Cx. quinquefasciatus* (Mordecai et al., 2019).

Several studies have corroborated this trend, with higher fecundity and fertility towards the

optimal of 28°C (Abouzied, 2017; Mogi, 1992), regardless of the blood source (bird or

mammal). Regarding photoperiod, the effects are not as evident as temperature since data are

somewhat contradictory. While Mogi (1992) found that mid and long day lengths induced

greater fecundity, Costanzo et al. (2015) did not find differences of day length on fecundity.

Because temperature and photoperiod are always coupled, in general their effects follow the

same unimodal trend. Although our experiment was conducted within this optimal range, the

highest fecundity and fertility in optimal conditions (summer) was present only in mammal-fed

mosquitoes, while higher fitness in bird-fed mosquitoes was detected in autumn. This

difference may indicate an interaction of seasonality with blood source.

While our findings may have certain limitations due to moderate sample sizes in some

treatments and their reliance solely on laboratory experiments, they verify an interaction

effect between blood meal source and seasonality, which had not been previously investigated

or documented in any other study. Nevertheless, the precise reasons behind the observed

pattern remain inconclusive. The highest levels of fecundity and fertility observed in bird-fed

mosquitoes during autumn, when conditions are more challenging, may confer a reproductive

advantage, leading to increased offspring production as an adaptive response to these adverse

conditions. In contrast, during the more favorable conditions of summer, this pattern may not

be as readily discernible. This proposed hypothesis needs further studies to be confirmed.

A second interaction was observed between blood meal source and gonotrophic cycle. We

found that fecundity and fertility during the first cycle were similar between mosquitoes fed

on chickens and those fed on mammals, and this was also similar with bird-fed mosquitoes

from the second gonotrophic cycle, regardless of the season. The only difference we observed

was between the fitness of mammal-fed mosquitoes from the second gonotrophic cycle and

the earlier mentioned treatments. This pattern is not different from those found in other

studies, as the first and second cycles are generally the most productive (Awahmukalah &

Brooks, 1985; Christiansen-Jucht et al., 2015). Although we observed a slight increase of

fecundity and fertility in bird-fed mosquitoes between cycles, it was statistically non-

significant. To our knowledge, only two studies have reported this increasing pattern: an older

one by Bennett (1970) and unpublished data from Obholz (personal communication), both

conducted with *Ae. aegypti*. On the contrary, a decrease in fitness between cycles has been

reported in other studies for *Cx. quinquefasciatus* (Richards et al., 2012; Telang & Skinner,

2019) fed upon birds or mammals. This is consistent with our findings regarding mouse-fed

mosquitoes; however, because the fecundity and fertility of mouse-fed mosquitoes during the

second cycle yielded the fewest egg rafts, this decreasing pattern might also be an effect due

to the insufficient sample size.

In addition to assessing fecundity, we also investigated whether hatchability could account for

the variations in fertility observed among different treatments. The only statistically significant

trend we identified was a decrease in the percentage of hatched eggs from the first to the

second gonotrophic cycle, but this was observed only in autumn mosquitoes. This decline in

hatching rate may be attributed to a constant exposure to cooling conditions, due to the

challenging environmental conditions during this season. This temperature-related effect

aligns with findings from other studies (van der Linde et al., 1990). However, although our data

suggest that changes in fertility may be linked to variations in hatching rate, it becomes

apparent that fecundity has a more pronounced impact on larval production due to the effects

of blood meal source and seasonality than hatching rate. Hence, the influence of hatchability

on the number of larvae per raft is rather limited.

In summary, our study investigates the critical issue of host shifting patterns in *Culex*

mosquitoes, which holds significant importance due to their role as bridge vectors between

bird and human hosts. Our model suggests that changes in photoperiod and environmental

temperature induce a different physiological state that requires the mosquito to feed on a

blood meal source that results in higher fitness (increased fecundity and fertility). We

predicted that mosquitoes fed on mammals in autumn will perform higher fecundity and

fertility than those fed on birds in the same season. Unexpectedly, we observed the opposite

pattern. Despite this, our study confirmed a significant interaction between blood meal source

and seasonality, suggesting a complex interplay of factors affecting mosquito reproduction and

its relationship with host use. Due to the observed opposite trend, we can consider three

plausible explaining scenarios. Firstly, it is possible that additional factors, such as genetic

mechanisms or variations in midgut microbiota composition, play a role in influencing host use

patterns. Secondly, host switching might be explained by factors related to vector biology

other than fitness. Lastly, it remains a possibility that Argentinian *Cx. quinquefasciatus*

populations do not exhibit the same host-switching behaviour observed in US populations, and

the observed interaction effects may be linked to different mosquito behaviours. Further

investigations are warranted to corroborate our findings and gain a deeper understanding of

the interaction between mosquito feeding pattern and seasonality. Ultimately, these

investigations will provide valuable insights into mosquito-borne disease transmission and

forecasting its emergence.

Acknowledgments

We would like to thank Dr. Agustín Quaglia for providing valuable help in statistical analysis. In

addition, we greatly appreciate very helpful comments on the manuscript by anonymous

reviewers. This study was supported by grant PICT 2018-1172 (FONCYT) and Consolidar Secyt-

UNC (2018-2023).

References

References

Akoh, J.I., Aigbodion, F.I., & Onyeka, J.O. (1993). Effect of source and size of bloodmeal on egg

production by *Culex quinquefasciatus* (Diptera, Culicidae). *Medical Entomology and Zoology*,

44(1): 41–43. https://doi.org/10.7601/mez.44.41

Alto, B.W., Connelly, C.R., O'Meara, G.F., Hickman, D., & Karr, N. (2014). Reproductive biology

and susceptibility of Florida *Culex coronator* to infection with West Nile virus. *Vector-Borne*

and Zoonotic Diseases, 14(8): 606–614. https://doi.org/10.1089/vbz.2013.1501

Bennett, G.F. (1970). The influence of the blood meal type on the fecundity of *Aedes*

(Stegomyia) aegypti L. (Diptera: Culicidae). *Canadian Journal of Zoology*, 48(3): 539–543.

https://doi.org/10.1139/z70-090

Beranek, M. (2018). Mosquitos del género *Culex* (Diptera: Culicidae) como vectores del virus

Saint Louis encephalitis (*Flavivirus*) en ambientes urbanos de la ciudad de Córdoba. PhD Thesis.

Universidad Nacional de Córdoba, Córdoba, Argentina.

Branda, M.F., Laurito, M., Visintin, A.M. & Almirón, W.R. (2021). Gonoactivity of *Culex* (*Culex*)

(Diptera: Culicidae) Mosquitoes During Winter in Temperate Argentina. *Journal of Medical*

Entomology, 58(3): 1454–1458. https://doi.org/10.1093/jme/tjaa295

Burkett-Cadena, N.D., McClure, C.J., Ligon, R.A., Graham, S.P., Guyer, C., Hill, G.E., Ditchkoff,

S.S., Eubanks, M.D., Hassan, H.K. & Unnasch, T.R. (2011). Host reproductive phenology drives

seasonal patterns of host use in mosquitoes. *PLOS one*, 6(3): e17681.

https://doi.org/10.1371/journal.pone.0017681

Carsey, T.M. & Harden, J.J. (2013). Monte Carlo simulation and resampling methods for social

science. Sage Publications, Los Angeles.

Ciota, A.T., Matacchiero, A.C., Kilpatrick, A.M., & Kramer, L.D. (2014). The effect of

temperature on life history traits of *Culex mosquitoes*. *Journal of Medical Entomology*, 51(1):

55–62.

https://doi.org/10.1603/me13003

Costanzo, K.S., Schelble, S., Jerz, K., & Keenan, M. (2015). The effect of photoperiod on life

history and blood-feeding activity in *Aedes albopictus* and *Aedes aegypti* (Diptera: Culicidae).

Journal of Vector Ecology, 40(1): 164–171. https://doi.org/10.1111/jvec.12146

Darsie, R.F., Jr. (1985). Mosquitoes of Argentina. Part I. Keys for identification of adult females

and fourth stage larvae in English and Spanish (Diptera: Culicidae). *Mosquito Systematics*,

17(3), 153–253.

Edman, J.D. & Taylor, D.J. (1968). Culex nigripalpus: Seasonal Shift in the Bird-Mammal Feeding

Ratio in a Mosquito Vector of Human Encephalitis. *Science*, 161(3836): 67–68.

https://doi.org/10.1126/science.161.3836.67

Erram, D., Vincent Black, T., & Burkett-Cadena, N. (2021). Host Bloodmeal Source Has No

Significant Effect on the Fecundity and Subsequent Larval Development Traits of the Progeny

in *Culicoides furens* Poey (Diptera: Ceratopogonidae). *Journal of Medical Entomology*, 58(6):

2439–2445. https://doi.org/10.1093/jme/tjab085

Farajollahi, A., Fonseca, D.M., Kramer, L.D. & Kilpatrick, A.M. (2011). "Bird biting" mosquitoes

and human disease: a review of the role of *Culex pipiens* complex mosquitoes in epidemiology.

Infection, genetics and Evolution, 11(7), 1577–1585.

https://doi.org/10.1016/j.meegid.2011.08.013

Gervasi, S.S., Burkett-Cadena, N., Burgan S.C., Schrey, A.W., Hassan, H.K., Unnasch, T.R. &

Martin, L.B. (2016). Host stress hormones alter vector feeding preferences, success, and

productivity. *Proceedings of the Royal Society B*, 283: 20161278.

https://doi.org/10.1098/rspb.2016.1278

Greenberg, J. (1951). Some nutritional requirements of adult mosquitoes (*Aedes aegypti*) for

oviposition: Two figures. *The Journal of Nutrition*, 43(1): 27–35.

https://doi.org/10.1093/jn/43.1.27

Hancock, C., & Camp, J.V. (2022). Habitat-Specific Host Selection Patterns of *Culex*

quinquefasciatus and *Culex nigripalpus* in Florida. *Journal of the American Mosquito Control*

Association, 38(2), 83–91. https://doi.org/10.2987/21-7054

Harrison, R.E., Brown, M.R., & Strand, M.R. (2021). Whole blood and blood components from

vertebrates differentially affect egg formation in three species of anautogenous mosquitoes.

Parasites & Vectors, 14: 1–19. https://doi.org/10.1186/s13071-021-04594-9

Hartig, F. (2022). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models. R package version 0.4.6.

Hothorn, T., Bretz F. & Westfall P. (2008). Simultaneous Inference in General Parametric

Models. *Biometrical Journal*, 50(3), 346–363.

Jansen, C.C., Webb, C.E., Graham, G.C., Craig, S.B., Zborowski, P., Ritchie, S.A., Russell, R.C. &

van den Hurk, A.F. (2009). Blood sources of mosquitoes collected from urban and peri-urban

environments in eastern Australia with species-specific molecular analysis of avian blood

meals. *American Journal of Tropical Medicine and Hygiene*, 81(5), 849–857.

https://doi.org/10.4269/ajtmh.2009.09-0008

Janssen, N., Fernandez-Salas, I., Díaz González, E.E., Gaytan-Burns, A., Medina-de la Garza, C.

E., Sanchez-Casas, R.M., Börster, J., Cadar, D., Schmidt-Chanasit, J. & Jöst, H. (2015).

Mammalophilic feeding behaviour of *Culex quinquefasciatus* mosquitoes collected in the cities

of Chetumal and Cancun, Yucatán Peninsula, Mexico. *Tropical Medicine & International Health*,

20(11), 1488–1491. https://doi.org/10.1111/tmi.12587

Kasili, S., Oyieke, F., Wamae, C. & Mbogo, C. (2009). Seasonal changes of infectivity rates of

Bancroftian filariasis vectors in Coast Province, Kenya. *Journal of Vector Borne Diseases*, 46:

219–224.

Kauffman, E., Payne, A., Franke, M.A., Schmid, M.A., Harris, E., & Kramer, L.D. (2017). Rearing

of *Culex* spp. and *Aedes* spp. mosquitoes. *Bio-protocol*, 7(17): e2542–e2542.

https://doi.org/10.21769/bioprotoc.2542

Kilpatrick, A.M., Kramer, L.D., Campbell, S.R., Alleyne, E.O., Dobson, A.P. & Daszak, P. (2005).

West Nile Virus Risk Assessment and the Bridge Vector Paradigm. *Emerging Infectious*

Diseases, 11(3): 425–429. https://doi.org/10.3201/eid1103.040364

Kilpatrick, A.M., Kramer, L.D., Jones, M.J., Marra, P.P. & Daszak, P. (2006). West Nile Virus

Epidemics in North America Are Driven by Shifts in Mosquito Feeding Behavior. *PLoS Biology*,

4(4): e82. https://doi.org/10.1371/journal.pbio.0040082

Kramer, C.Y. (1956). Extension of multiple range tests to group means with unequal numbers

of replications. *Biometrics*, 12: 307–310. https://doi.org/10.2307/3001469

Labarthe, N., Serrão, M.L., Melo, Y., José-Oliveira, S., and Lourenço-de-Oliveira, R. (1998).

Potential vectors of *Dirofilaria immitis* in Itacoatiara, Oceanic Region of Niterói Municipality,

state of Rio de Janeiro, Brazil. *Memorias do Instituto Oswaldo Cruz*, 93: 425–432.

https://doi.org/10.1590/s0074-02761998000400001

Mogi, M. (1992). Temperature and photoperiod effects on larval and ovarian development of

New Zealand strains of *Culex quinquefasciatus* (Diptera: Culicidae). *Annals of the*

Entomological Society of America, 85(1), 58–66. https://doi.org/10.1093/aesa/85.1.58

Molaei, G., Andreadis, T.G., Armstrong, P.M., Bueno, R., Dennett, J.A., Real, S.V., Sargent, C.,

Bala, A., Randle, Y., Guzman, H., da Rosa, A.T., Wuithranyagool & Tesh, R. B. (2007). Host

feeding pattern of *Culex quinquefasciatus* (Diptera: Culicidae) and its role in transmission of

439 West Nile virus in Harris County, Texas. *American Journal of Tropical Medicine and Hygiene*,

440 77(1), 73–81.

441 https://doi.org/10.4269/ajtmh.2007.77.73

442 Mordecai, E.A., Caldwell, J.M., Grossman, M.K., Lippi, C.A., Johnson, L.R., Neira, M., Rohr, J.R.,

443 Ryan, S.J., Savage, V., Shocket, M.S., Sippy, R., Stewart Ibarra, A.M, Thomas, M.B. & Villena, O.

444 (2019). Thermal biology of mosquito-borne disease. *Ecology letters*, 22(10), 1690–1708.

445 https://doi.org/10.1111/ele.13335

446 Muturi, E.J., Muriu, S., Shililu, J., Mwangangi, J.M., Jacob, B.G., Mbogo, C., Githure, J. & Novak,

447 R.J. (2008). Blood-feeding patterns of *Culex quinquefasciatus* and other culicines and

448 implications for disease transmission in Mwea rice scheme, Kenya. *Parasitology Research*,

449 102(6), 1329–1335. https://doi.org/10.1007/s00436-008-0914-7

450 R Core Team (2022). R: A language and environment for statistical computing. R Foundation

451 for Statistical Computing, Vienna, Austria.

452 Richards, S.L., Anderson, S.L., & Yost, S.A. (2012). Effects of blood meal source on the

453 reproduction of *Culex pipiens quinquefasciatus* (Diptera: Culicidae). *Journal of Vector Ecology*,

454 37(1): 1–7. https://doi.org/10.1111/j.1948-7134.2012.00194.x

455 Shroyer, D.A. & Silvery, R.E. (1972). A comparison of egg production of *Culex pipiens pipiens* L.

456 fed on avian and mammalian hosts. *Mosquito News*, 32(4):636–637.

457 Silva Santos, C., Pie, M.R., da Rocha, T.C., & Navarro-Silva, M.A. (2019). Molecular

458 identification of blood meals in mosquitoes (Diptera, Culicidae) in urban and forested habitats

459 in southern Brazil. *PloS one*, 14(2): e0212517. https://doi.org/10.1371/journal.pone.0212517

460 Smith, J.L. & Fonseca, D.M. (2004). Rapid assays for identification of members of the *Culex*

461 (*Culex*) *pipiens* complex, their hybrids, and other sibling species (Diptera: Culicidae). *The*

462 *American Journal of Tropical Medicine and Hygiene*, 40(4), 339–345.

463 https://doi.org/10.4269/ajtmh.2004.70.339

464 Spinsanti, L.I, Díaz, L.A, Glastein, N., et al. (2008). Human outbreak of St. Louis encephalitis

465 detected in Argentina, 2005. *Journal of Clinical Virology*, 42: 27–33.

466 https://doi.org/10.1016/j.jcv.2007.11.022

467 Stein, M., Zalazar, L., Willener, J.A., Almeida, F.L., & Almirón, W.R. (2013). Culicidae (Diptera)

468 selection of humans, chickens and rabbits in three different environments in the province of

469 Chaco, Argentina. *Memórias do Instituto Oswaldo Cruz*, 108, 563–571.

470 https://doi.org/10.1016/j.actatropica.2015.12.002

471 Takken, W., & Verhulst, N.O. (2013). Host preferences of blood-feeding mosquitoes. *Annual*

472 *Review of Entomology*, 58, 433–453. https://doi.org/10.1146/annurev-ento-120811-153618

473 Telang, A., & Skinner, J. (2019). Effects of host blood meal source on reproductive output,

474 nutrient reserves and gut microbiome of West Nile virus vector *Culex quinquefasciatus*. *Journal*

475 *of Insect Physiology*, 114, 15–22. https://doi.org/10.1016/j.jinsphys.2019.02.001

476 van der Linde, T.C. de K., Hewitt, P.H., Nel, A. & van der Westhuizen, M.C. (1990). Development

477 rates and percentage hatching of *Culex* (*Culex*) *theileri* Theobald (Diptera: culicidae) eggs at

478 various constant temperatures. *Journal of the Entomological Society of Southern Africa*, 53(1):

479 17–26. https://doi.org/10.1093/jmedent/27.6.939

480 Venables, W.N., & Ripley, B.D. (2002). Exploratory multivariate analysis. In: Modern applied

481 statistics with S (pp. 301–330). Springer, New York.

482 https://doi.org/10.1007/978-0-387-21706-2_11

483 Yan, J., Gangoso, L., Martínez-de la Puente, J., Soriguer, R. & Figuerola, J. (2017). Avian

484 phenotypic traits related to feeding preferences in two *Culex mosquitoes*. *The Science of*

485 *Nature*, 104(9), 1–10. https://doi.org/10.1007/s00114-017-1497-x

486   Yan, J., Broggi, J, Martínez-de la Puente, J., Gutiérrez-López, R., Gangoso, L., Soriguer, R. &

487   Figuerola, J. (2018) Does Bird metabolic rate influence mosquito feeding preference? *Parasites*

488   *& Vectors*, 11(1), 1–9. https://doi.org/10.1186/s13071-018-2708-9

Table 1. Results of reproductive outputs of *Culex quinquefasciatus* for the first and second gonotrophic cycle. Mean values are accompanied with its respective standard deviation.

Bloodmeal source	Season	Full engorged females	Numer of egg rafts	Fecundty (eggs/raft)	Fertility (larvae/raft)	Hatchability (larvae/eggs)
First gonotrophic cycle (first feeding)						
Mouse	Autumn	58	23	124.5 ± 20.9	120.7 ± 21.5	0.96 ± 0.046
Mouse	Summer	208	120	150.7 ± 29.4	141.8 ± 33.9	0.92 ± 0.10
Chicken	Autumn	163	76	147.5 ± 31.8	133.4 ± 38.4	0.89 ± 0.14
Chicken	Summer	116	74	125.6 ± 39.6	109.9 ± 44.0	0.85 ± 0.20
Second gonotrophic cycle (second feeding)						
Mouse	Autumn	23	15	89.1 ± 19.3	73.3 ± 28.0	0.82 ± 0.25
Mouse	Summer	92	45	106.7 ± 32.0	98.8 ± 31.4	0.91 ± 0.10
Chicken	Autumn	73	36	142.1 ± 36.6	129.2 ± 43.7	0.88 ± 0.18
Chicken	Summer	38	27	138.2 ± 30.3	117.3 ± 38.1	0.84 ± 0.18

Table 2. Analysis of deviance table of the negative binomial model for the effect of blood meal source, seasonality and gonotrophic cycle on fecundity. Dispersion parameter (θ) = 18.75.

Factor	LRT X^2	df	p-value
Seasonality	15.72	1	7.361×10^{-5}
Blood source	8.12	1	0.00438
Gonotrophic cycle	0.55	1	0.45904
Seasonality:Blood source	25.00	1	7.717×10^{-7}
Seasonality:Gonotrophic cycle	3.11	1	0.07764
Blood source:Gonotrophic cycle	9.29	1	0.00230
Seasonality:Blood source:Gonotrophic cycle	1.42	1	0.23340

Table 3. Analysis of deviance table of the negative binomial model for the effect of blood meal source, seasonality and gonotrophic cycle on fertility. Dispersion parameter (θ) = 9.52.

Factor	LRT X^2	df	p-value
Seasonality	11.87	1	0.00057
Blood source	1.64	1	0.19964
Gonotrophic cycle	0.20	1	0.65066
Seasonality:Blood source	13.82	1	0.00020
Seasonality:Gonotrophic cycle	0.87	1	0.34972
Blood source:Gonotrophic cycle	11.40	1	0.00073
Seasonality:Blood source:Gonotrophic cycle	0.04	1	0.83479

500 **Table 4.** Analysis of deviance table of the quasipoisson model for the effect of blood meal
501 source, seasonality and gonotrophic cycle on hatchability.

502
503
504

Factor	LRT X^2	df	p-value
Seasonality	5.52	1	0.37623
Blood source	73.06	1	0.00129
Gonotrophic cycle	29.37	1	0.04137
Seasonality:Blood source	7.56	1	0.30066
Seasonality:Gonotrophic cycle	41.02	1	0.01593
Blood source:Gonotrophic cycle	5.53	1	0.37590
Seasonality:Blood source:Gonotrophic cycle	1.20	1	0.67981

Figure 1. Interaction plots showing the marginal effects of blood meal source-seasonality and
blood meal source-gonotrophic cycle on (**A**, **B**) fecundity and (**C**, **D**) fertility. Predicted mean
values and its corresponding confidence intervals are shown. Means sharing same letters are
not statistically different.

Figure 2. Boxplots showing the effects of both blood meal source and seasonality on
hatchability for the first and second gonotrophic cycle.

Figure 3. Interaction plot showing the marginal effects of seasonality-gonotrophic cycle on
hatchability.

Figure 4. Simulated means of fecundity (eggs per raft) of each treatment along 1,000
iterations. Abbreviations. B: avian blood; M: mammalian blood; A: autumn; S: summer; I: first
gonotrophic cycle; II: second gonotrophic cycle; Ac: model accuracy.

Figure 5. Simulated means of fertility (eggs per raft) of each treatment along 1,000 iterations.
Abbreviations same as Figure 4.