Effects of blood meal source and seasonality on reproductive traits of

Culex quinquefasciatus **(Diptera: Culicidae)**

Kevin Alen Rucci[1,3,*], Gabriel Barco[1,3], Andrea Onorato[1,3], Mauricio Beranek[1], Mariana Pueta[2,3] &

Adrián Díaz[1,3*]

[1] Laboratorio de Arbovirus, Instituto de Virología "Dr. J. M. Vanella" (InViV), Facultad de Ciencias

Médicas (FCM), Universidad Nacional de Córdoba (UNC)

[2] Instituto de Investigaciones en Biodiversidad y Medioambiente (INIBIOMA), CONICET -

Universidad Nacional de Comahue (UNCo)

[3] Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina

*Corresponding authors:

Kevin Alen Rucci

Laboratorio de Arbovirus – Instituto de Virología "Dr. J. M. Vanella" (InViV), Facultad de Ciencias

Médicas (FCM), Universidad Nacional de Córdoba (UNC).

Enfermera Gordillo Gómez s/n. Ciudad Universitaria, X5000HUA, Córdoba capital, Córdoba,

Argentina.

E-mail addresses: kevin.rucci@mi.unc.edu.ar, kevinrucci@outlook.com

Adrián Díaz

Laboratorio de Arbovirus – Instituto de Virología "Dr. J. M. Vanella" (InViV), Facultad de Ciencias

Médicas (FCM), Universidad Nacional de Córdoba (UNC).

Enfermera Gordillo Gómez s/n. Ciudad Universitaria, X5000HUA, Córdoba capital, Córdoba,

Argentina.

E-mail addresses: adrian.diaz@fcm.unc.edu.ar, adrian.diaz@conicet.gov.ar

Abstract

Host selection by mosquitoes is a keystone to understanding viral circulation and predicting future infection outbreaks. *Culex* mosquitoes frequently feed on birds during spring and early summer, shifting into mammals towards late summer and autumn. This host switch may be due to changes in mosquito fitness. The aim of this study was to assess if the interaction effect of blood meal source and seasonality may influence reproductive traits of *Culex quinquefasciatus* mosquitoes. For this purpose, *Cx. quinquefasciatus* mosquitoes were reared in simulated summer and autumn conditions and fed on two different hosts, chickens and mice, in a factorial design. Fecundity, fertility, and hatchability during two consecutive gonotrophic cycles were estimated. We found greater fecundity and fertility for mosquitoes fed upon birds than mammals. Fecundity and fertility did not vary between seasons for chicken-fed mosquitoes, whereas in autumn they decreased for mouse-fed mosquitoes. These traits decreased in the second gonotrophic cycle for mouse-fed mosquitoes, whereas they did not vary between cycles for chicken-fed mosquitoes. There was no statistically significant effect of blood meal source, seasonality or their interaction on hatchability, hence this variable was similar among treatments. Overall, these results indicate a statistically significant interaction effect of blood meal source and seasonality on fecundity and fertility. However, the pattern was opposite in relation to our hypothesis, suggesting that further studies are needed to confirm and expand our knowledge about mosquito biology and its relationship with seasonal host use shifting.

Key words: mosquito vector, fecundity, fertility, hatchability

Introduction

The southern house mosquito, *Culex quinquefasciatus* Say (Diptera: Culicidae), is a worldwide distributed vector of numerous pathogens, including *Wuchereria bancrofti* Cobbold (Kasili et al., 2009), *Dirofilaria immitis* Leidy (Labarthe et al., 1998) and avian malaria (Farajollahi et al., 2011). Besides, this mosquito is also one of the main species responsible for the transmission of the arboviruses West Nile (WNV) and St. Louis encephalitis (SLEV) viruses (Farajollahi et al., 2011). Because of this, this species is of critical concern due to its impact on both public and veterinary health.

Since arboviruses are transmitted by the bite of infected mosquitoes between vertebrate hosts, activity of these pathogens is determined, at least in part, by the blood feeding habits of these vectors. According to its host feeding pattern range, *Culex* mosquitoes can be regarded as specialists or generalists. Specialist species include mosquitoes feeding primarily on mammals, birds, or reptiles and amphibians. Generalists are mosquitoes fed on the most abundant available hosts (Fikrig and Harrington, 2021). Additionally, some species could experience a seasonal shift in their feeding habits, from avian to mammal hosts (Edman and Taylor, 1968; Hancock and Camp, 2022), being of greater concern from an epidemiological perspective given that they can act as bridge vectors in the transmission of some arboviruses, such as SLEV and WNV (Kilpatrick et al., 2006).

West Nile virus (WNV) and St. Louis encephalitis virus (SLEV) are zoonotic pathogens maintained in nature through an enzootic network involving transmission among viremic birds and ornithophagic mosquitoes (Kilpatrick et al., 2006). However, during late summer and early autumn, these viruses increase their activity in humans and other mammals, which has led to repeated epidemics in the US (Kilpatrick et al., 2006) and South America (Spinsanti et al., 2008). Likewise, several *Culex* species, including *Cx. quinquefasciatus*, have exhibited a seasonal shift in their feeding behaviour, feeding primarily on birds in spring and early summer and increasing

the proportion of mammal blood meals in late summer and autumn (Thiemann et al., 2011).

This seasonal shift suggests that viral activity might spill over from birds to mammals as a result

of the mosquito host switch (Edman and Taylor, 1968; Kilpatrick et al., 2006).

Two main contrasting hypotheses have been proposed to explain why primarily ornithophagic

mosquitoes increase their feeding on mammals later in the season. The *migration* hypothesis,

proposed by Kilpatrick et al. (2006) for *Cx. pipiens*, suggests that this shift in feeding pattern is

related to the abundance of the preferred avian host, the American robin (*Turdus migratorius*

L.). This hypothesis is based on their findings that as robin densities decrease, the proportion of

mammalian blood meals in *Cx. pipiens* populations increase. However, this hypothesis has

several limitations: (i) multiple studies have shown that American robins are not always the

preferred host for all *Cx. pipiens* populations (Apperson et al., 2002; Patrican et al., 2007); (ii) in

some areas (e.g., Tennessee) with documented seasonal host shifts, American robins are

present year-round (Savage et al., 2007); and (iii) this explanation applies only to mosquitoes

whose primary host is the American robin, a species found only in the Northern Hemisphere, or

another migratory bird. Therefore, it might not be applicable to southern regions, such as

Argentina, where WNV seasonality is documented (Spinsanti et al., 2008) but American robin is

absent. Because of this, it seems that the hypothesis proposed by Kilpatrick et al. (2006) might

be habitat dependent (Caranci, 2010). On the other hand, the *defensive behaviour* hypothesis

proposed by Burkett-Cadena et al. (2011), suggests that host breeding cycles drive the host shift

of mosquitoes. According to this hypothesis, in hosts with parental care, during reproductive

seasons there is a greater investment of energy in assuring the offspring survivorship,

consequently producing an increase in susceptibility of being bitten by mosquitoes as a result of

a decrease in defensive behaviours. This event leads to the detection of peaks of host use during

periods of reproductive investment, in summer for birds and autumn for mammals (Burkett-

Cadena et al., 2011). While this hypothesis was proposed as an alternative to address the

limitations of the migration hypothesis, it also has drawbacks. It is based on a limited range of

host species, assumes a distinct seasonal phenology—which is not present for all mosquito

hosts—and remains untested, leaving its validity uncertain.

Explanations for the seasonal shift in mosquito host feeding focused on vector biology are

largely lacking in the literature. Numerous biological factors, such as stress, metabolic rate, and

blood meal source, along with environmental variables like temperature and photoperiod,

influence mosquito physiology and can affect reproductive traits such as fecundity,

development rate, and survivorship. These factors can give rise to new nutritional requirements

that may ultimately lead to seasonal variations in host selection by mosquitoes (Ciota et al.,

2014; Costanzo et al., 2015; Gervasi et al., 2016; Yan et al., 2018, 2017). While several studies

have examined the effects of these biological and environmental variables on mosquito

reproduction, the complex nature of mosquito biology suggests that multiple interactions

among these variables may produce novel responses that are not apparent when considered

individually. The aim of this study was to assess whether there is an interaction effect between

the source of the host blood meal and seasonality (defined by temperature and photoperiod)

on three reproductive traits of *Culex quinquefasciatus* mosquitoes: fecundity, fertility, and

hatchability. We hypothesize that the interaction between these two variables influences the

reproductive outcomes, potentially leading to a seasonal shift in host selection, driven by a

reproductive advantage. Given the reported seasonal changes in host use by *Cx.*

quinquefasciatus, in autumn we expect a greater number of eggs (fecundity) and larvae (fertility)

in mosquitoes after feeding on a mammal host compared to an avian host, and the opposite

trend in summer.

Materials and methods

Establishment and maintenance of mosquitoes

Egg rafts of *Culex quinquefasciatus* were collected from a drainage ditch at Universidad Nacional

de Córdoba Campus, Córdoba city, in February 2021. Each raft was individually maintained in

plastic containers with one liter of distilled water. The hatched larvae were fed with 100 mg of

liver powder three times per week until pupation. Pupae were then transferred to plastic

emerging cages (21 cm x 12 cm) covered with a tulle-like fabric, containing distilled water but

no food. Adults emerging from each raft were identified morphologically (Darsie, 1985) and

molecularly (Smith and Fonseca, 2004) to ensure they corresponded to *Cx. quinquefasciatus*,

since *Cx. pipiens* and its hybrids coexist sympatrically in Córdoba city (Branda et al., 2021). All

adults were reared in a cardboard cage of 22.5 liters (28 cm x 36.5 cm) and provided *ad libitum*

with a 10% sugar solution soaked in cotton pads placed on plastic cups. For long-term

maintenance of the colony, 24-hour-starved mosquitoes were offered a blood meal from a

restrained chicken twice a month. Four days after feeding, a plastic container with distilled water

was placed inside the cage to allow engorged females to lay egg rafts. Batches of egg were

collected and transferred to plastic containers (30 cm x 25 cm x 7 cm) in a proportion of three

rafts per container, filled with three liters of distilled water. The hatched larvae were also fed

with liver powder at the same proportion described above. Pupae were transferred to emerging

cages, and adult mosquitoes were placed in the final cardboard cage. The colony has been

maintained for over twenty generations in the Insectary of the Instituto de Virología "Dr. J.M.

Vanella" (InViV). Room conditions were controlled at 28 ± 1 °C, with a 12L:12D photoperiod and

70% relative humidity.

Experimental design

Blood source. For experimental trials, avian hosts (live chicks of the species *Gallus gallus*) and

mammalian hosts (live mice of the species *Mus musculus*, strain C57BL/6) were used to evaluate

the effect of blood meal source on fecundity, fertility, and hatchability. Chicks were generously

donated by the Bartolucci poultry farm (Córdoba, Argentina), while mice were commercially

obtained from the Instituto de Investigación Médica Mercedes y Martín Ferreyra (CONICET -

Universidad Nacional de Córdoba). In each trial, 24-hour-starved adult female mosquitoes were

provided with a blood meal from restrained chicks or mice. Vertebrate hosts were offered to

mosquitoes one hour before the lights were turned off and were kept for 3 hours during 2

consecutive gonotrophic cycles.

The experimental use of animals (mosquitoes and vertebrates) was approved by the ethical

committee at Facultad de Ciencias Médicas, Universidad Nacional de Córdoba (FCM-UNC) in

compliance with the legislation regarding the use of animals for experimental and other

scientific purposes (accession code: CE-2022-00518476-UNC-SCT#FCM).

Seasonality. To assess the effect of seasonality (photoperiod + temperature) on fecundity,

fertility, and hatchability, a "typical" summer and autumn day from Córdoba city was simulated

in an incubator, where mosquitoes were housed throughout the entire experiment. Summer

conditions were as follows: $T°_{min}$ = 22°C, $T°_{max}$ = 28°C, photoperiod = 14L:10D. Autumn conditions

were characterized by: $T°_{min}$ = 16°C, $T°_{max}$ = 22°C, and photoperiod = 10L:14D. The humidity levels

were maintained at 60-70% for both conditions. Data for simulated conditions were obtained

from Climate Data website (https://es.climate-data.org/).

Feeding trial design. The interaction effect of blood source (chicken or mouse) and seasonality

(autumn or summer) was evaluated during two consecutive gonotrophic cycles (I and II). This

combination produced eight colonies that were established using egg rafts collected from the

maintenance colony: mouse:autumn-I, mouse:autumn-II, chicken:autumn-I, chicken:autumn-II,

mouse:summer-I, mouse:summer-II, chicken:summer-I, chicken:summer-II. This set of eight

colonies was repeated in triplicate (1-3) at different time points, generating a total of 24

treatments.

The experimental colonies were maintained under controlled conditions to ensure that all adults

were of the same size (Kauffman et al., 2017). Adult mosquitoes were placed in cardboard cages

of 8.3 liters (21 cm x 24 cm) and were provided with *ad libitum* access to a 10% sugar solution,

which was soaked in cotton pads placed on plastic cups.

Feeding trials were conducted at two time points: 5 days post-emergence (first cycle) and 14

days post-emergence (second cycle). Following each blood meal, female mosquitoes were

anesthetized using CO_2 and classified as fully engorged (VI Sella's stage), partially fed (I-V Sella's

stage), or unfed (I Sella's stage), following the classification by Santos et al. (2019). Fully

engorged females were counted and separated into a separate cage to complete their

gonotrophic cycle. Four days after feeding, a cup containing distilled water was placed inside the

cage to facilitate oviposition. After each oviposition cycle, egg rafts were collected, counted, and

transferred to a 12-well plate containing 4 mL of distilled water. They were then photographed

to subsequently determine the number of eggs per raft. Rafts were maintained in each well until

they hatched into L1 larvae, at which point they were preserved with 2 mL of 96% ethanol, and

the number of L1 larvae per raft was counted.

Statistical analysis. The reproductive outputs measured for each experimental colony were

fecundity, fertility, and hatchability. All analyses were performed using R Studio statistic

software, v.4.2.1 (R Core Team, 2022).

Fecundity was defined as the number of eggs per raft and *fertility* as the number of L1 larvae

hatched per raft. To evaluate the effect of blood meal source and seasonality on these two

variables, a Generalized Linear Mixed Model (GLMM; *lme4* package) with negative binomial

error distribution and logarithmic link function was adjusted (Bates et al., 2015). Fecundity and

fertility served as response variables, while the fixed effect variables included blood source,

seasonality, and gonotrophic cycle, all in a three-way interaction. Treatment (n = 24) was set as

a random effect variable.

Hatchability (= hatching rate) was defined as the ratio of the number of larvae to the number of

eggs per raft. While the original intent was to apply a GLMM to all three variables, the validation

of GLMMs for hatchability across several distributions revealed poor fit. Consequently, a

Generalized Linear Model (GLM; MASS package) with a quasipoisson error distribution and

logarithmic link function was used instead (Venables and Ripley, 2002). In this model, the

response variable was the number of larvae, while the explanatory variables included blood

source, seasonality, gonotrophic cycle, and replicate as a four-way interaction. Additionally, the

logarithm of the number of eggs was incorporated as an offset in the model.

Multiple comparisons among treatments were conducted using the Tukey's honestly significant

difference test (HSD Tukey), incorporating the Kramer (1956) correction for unbalanced data,

also known as the Tukey-Kramer method. This approach allows to set the family-wise error rate

to 0.05. These comparisons were performed using the *emmeans* package (Lenth, 2017).

Model validation was assessed using a simulation-based approach to generate randomized

quantile residuals to test goodness of fit (Q-Q plot), homoscedasticity, and outliers and

influential points (Cook's distance and Leverage plots). These analyses were conducted with the

DHARMa package (Hartig, 2024).

Additionally, post-hoc statistical power was evaluated using the *mixedpower* package, a tool

specifically designed for simulating statistical power in Generalized Linear Mixed Models

(GLMMs). The analysis proceeded in three key steps: (1) simulation of new datasets based on

the parameters of an original fitted model (the fecundity and fertility models), (2) refitting the

model to each dataset and testing for statistical significance, and (3) calculating statistical power

as the proportion of simulated datasets where the effect of interest reached significance. Fixed

effects were simulated for sample sizes ranging from 30 to 150, with 1,000 datasets generated

at each sample size to ensure robust estimation. The variable treatment was designated as a

random effect, capturing variability associated with this factor. For further details on the

methodology, including considerations for simulation accuracy and model structure, refer to

Kumle et al. (2021).

Results

A total of 1,162 egg rafts were obtained for analysis, distributed across the eight treatments as

follows: chicken:autumn-I: 260, chicken:autumn-II: 174, chicken:summer-I: 162,

chicken:summer-II: 88, mouse:autumn-I: 146, mouse:autumn-II: 96, mouse:summer-I: 167, and

mouse:summer-II: 69. The full count of egg rafts for each replicate (24 treatments), along with

the means for fecundity, fertility, and hatchability, are summarized in Table 1. This sample size

was sufficient to detect the observed effect with a statistical power of 0.8, indicating an 80%

chance of detecting a true effect if it exists (Figure SM1).

In both models—fecundity and fertility—the interaction between blood source and seasonality

was statistically significant (fecundity: LRT X^2 = 5.69, p < 0.05; fertility: LRT X^2 = 4.37, p < 0.05)

(Tables 2,3; Figures 1,2A). Mosquitoes that fed on chicken blood had the highest fecundity

(summer: 136.4 eggs/raft, autumn: 141.5 eggs/raft) and fertility (summer: 123.9 larvae/raft,

autumn: 126 larvae/raft) in both seasons. In summer, fecundity and fertility were 7% and 11%

higher, respectively, in chicken-fed mosquitoes compared to mouse-fed mosquitoes. In autumn,

these differences increased to 46% for fecundity and 52% for fertility in favor of chicken-fed

mosquitoes. No significant seasonal variation was observed in fecundity and fertility for chicken-

fed mosquitoes, but both measures were statistically significant lower by 24% and 25%,

respectively, in autumn compared to summer for mouse-fed mosquitoes.

In the fertility model, there was also a significant interaction between blood source and

gonotrophic cycle (LRT X^2 = 3.98, p < 0.05), indicating that the impact of blood type on larvae

production differed between the first and second cycles (Table 3, Figure 2B). Chicken-fed

mosquitoes showed no difference in fertility between cycles, while those fed on mouse blood

had 25% lower fertility in the second cycle compared to the first. During the second cycle,

fertility was 50% higher in chicken-fed mosquitoes compared to mouse-fed mosquitoes.

Figure 3 presents the predicted mean values of fecundity and fertility for the 24 treatments,

with no differences observed across the three replicates. The variance of random intercepts

among treatments was approximately 0.006 for fecundity and 0.007 for fertility, accounting for

6% and 4% of the total variance in the respective models.

There was no statistical difference among treatments and replicates in terms of hatchability,

given that no statistical effects of blood source, seasonality or gonotrophic cycle were observed

(Table 4). The mean hatchability for all treatments was 0.89, with a range between 0.84 for the

mouse:autumn-II (replicate 1) colony and 0.96 for mouse:autumn-I (replicate 1) (Table 1, Figure

4).

Discussion

Understanding the factors influencing mosquito host selection is crucial for predicting arbovirus

transmission dynamics. Seasonal shift in host utilization have been widely attributed to changes

in host behaviour (Burkett-Cadena et al., 2011; Kilpatrick et al., 2006), yet the role of mosquito

biology in this process remains largely unexplored. Blood meal source and seasonality are known

to affect mosquito reproductive success, but their combined effect has received little attention.

Our study tested the hypothesis that the interaction between blood meal source and seasonality

could affect mosquito fitness, a key factor that could play a role in seasonal host shift. By

examining this interaction, we aimed to assess whether physiological processes like

reproduction could be considered a candidate for shaping seasonal feeding patterns in *Culex*

quinquefasciatus.

If seasonal host shift were driven by fitness advantages, we would expect mosquitoes to exhibit

higher reproductive output when feeding on the predominant host in each season—that is,

greater fecundity and fertility with bird blood in summer and with mammal blood in autumn.

However, our findings did not fully align with these expectations. While mosquitoes fed on birds

maintained high reproductive success in both summer and autumn, those fed on mammals

exhibited a significant decline in reproductive output during autumn. This suggests that, rather

than conferring a reproductive advantage in this season, mammalian blood may become a

suboptimal resource under autumn conditions, potentially due to metabolic constraints or

differences in nutrient assimilation.

The effect of blood meal on fitness observed in this study (greater in bird-fed mosquitoes) is

consistent with previous findings on *Cx. quinquefasciatus* (Richards et al., 2012; Telang and

Skinner, 2019) and other *Culex* (Demirci et al., 2014) and *Aedes* (Harrison et al., 2021) species.

Avian blood is known to be nutritionally richer than mammalian blood, particularly in proteins

and essential amino acids (e.g., isoleucine) required for egg development (Alto et al., 2014).

Beyond intrinsic differences in blood composition, environmental factors such as temperature

and photoperiod (seasonality) may further modulate how nutrients from a blood meal are

allocated. While most of the nutrients obtained from a blood meal are directed toward egg

production, a portion is also required for non-reproductive physiological functions (Rivera-Pérez

et al., 2017), and the extent to which these resources are utilized may depend on external

conditions such as temperature (Briegel and Timmermann, 2001).

Previous studies have shown that temperature regulates oogenesis and influences how

mosquitoes allocate proteins obtained from a blood meal. In *Aedes aegypti*, for example, the

proportion of blood-derived proteins allocated to egg production varies with temperature: at

low temperatures, approximately 50% of these proteins are used for oogenesis, whereas at

higher temperatures, this proportion drops to 25% (Briegel and Timmermann, 2001). If a similar

mechanism occurs in *Cx. quinquefasciatus*, it could partially explain why mammal-fed

mosquitoes exhibited a more substantial reproductive decline in autumn. Under lower

temperatures, a greater proportion of proteins may be required for egg development, which

could disproportionately impact individuals feeding on nutritionally poorer blood sources.

In addition to this blood meal-seasonal effect, we also found a significant interaction between

blood meal source and gonotrophic cycle. This result suggests that the impact of blood sources

is not uniform across cycles. If blood meal source were the only determinant of reproductive

success, we would expect its effects to be consistent across cycles, with mosquitoes fed on birds

maintaining higher reproductive output than those fed on mammals in a stable pattern.

However, our findings showed that while bird-fed mosquitoes sustained similar reproductive

output across cycles, those fed on mammals exhibited a significant decline in fertility in the

second cycle. Since each cycle involves a new blood meal, this suggests that factors beyond

blood nutritional quality contribute to the observed pattern, disproportionately affecting

mosquitoes feeding on mammals in later cycles.

A decline in fertility across gonotrophic cycles has also been documented in other *Cx.*

quinquefasciatus populations fed on both birds and mammals (Richards et al., 2012; Telang and

Skinner, 2019), suggesting that reproductive output varies not only with seasonality but also

with the number of successive cycles. One potential explanation for this decline is age-related

reproductive constraints. With each successive cycle, ovarian follicular degeneration increases,

leading to a reduction in the number of viable eggs per raft (Awahmukalah and Brooks, 1985).

Additionally, given that several days separate the first and second cycles, physiological aging

could contribute to this pattern (McCann et al., 2009). Interestingly, bird-fed mosquitoes

maintained stable fertility across cycles, a pattern that has also been observed in previous

studies (Bennett, 1970). While fertility often declines in later cycles, some studies have reported

no change or even an increase in egg production under certain conditions (Bennett, 1970). This

variability suggests that the effects of successive gonotrophic cycles on fertility may depend on

multiple factors, including blood meal quality. Since avian blood is known to be nutritionally

richer than mammalian blood, mosquitoes feeding on birds may have access to sufficient

resources to sustain egg production across cycles, even as they age. However, further research

is needed to determine whether specific nutrient components play a role in maintaining

reproductive output over time. Similarly, body size has been shown to influence reproductive

patterns in some mosquito species. In *Aedes aegypti*, for instance, larger females exhibit a rapid

decline in fecundity after the first cycle, whereas smaller females maintain consistent egg

production across early cycles before experiencing a significant drop (Briegel et al., 2002). While

we did not assess body size in this study, these findings underscore the complexity of

reproductive trade-offs in mosquitoes and the importance of considering multiple interacting

factors when examining reproductive dynamics across cycles.

Together, these results indicate that both blood meal source and seasonality influence mosquito

reproductive output. Moreover, hatchability did not differ among treatments, indicating that

differences in egg viability did not drive these effects. This reinforces the idea that the main

differences in reproductive output stem from factors affecting fecundity and fertility rather than

post-oviposition processes.

While our findings confirm an interaction effect between blood meal source and seasonality,

several caveats must be considered. First, our study was conducted under controlled laboratory

conditions, which may not fully capture the environmental variability experienced by

mosquitoes in the field. Factors such as fluctuating temperature, natural light cycles, and

variable humidity levels could influence mosquito reproductive performance in ways not

accounted for in our setup. However, as this is the first study of its kind, future research should

aim to incorporate more ecologically realistic conditions. A second limitation commonly

encountered in ecological and behavioural studies, is determining an appropriate sample size

for detecting biologically meaningful effects (Taborsky, 2010). To address this, we repeated the

assay across three replicates and conducted a post-hoc power analysis, which suggested that

our sample size was adequate given the presence of interaction effects among explanatory

variables. Finally, it remains possible that *Cx. quinquefasciatus* populations in Argentina do not

exhibit the same host-switching behaviour observed in US populations, potentially influencing

the relevance of our findings for different ecological contexts. However, previous studies have

reported seasonal shifts in host selection and arbovirus transmission patterns in Argentina

(Beranek, 2019; Spinsanti et al., 2008), suggesting that similar dynamics may be at play. More

field-based studies will be essential to assess seasonal host variation in *Cx. quinquefasciatus* and

its implications for arbovirus transmission.

In summary, our study investigates a critical aspect of mosquito reproductive success and its

potential link to seasonal host selection in *Culex* mosquitoes, a group of significant

epidemiological importance due to their role as bridge vectors between avian and human hosts.

While we detected a significant interaction between blood meal source and seasonality, our

findings do not support the idea that reproductive fitness alone explains seasonal host shifts.

Instead, they suggest that the relationship between mosquito reproduction and host selection

is more complex, likely influenced by multiple ecological and physiological factors beyond fitness

constraints. Given this complexity, alternative explanations for seasonal host shifts should be

considered. Genetic predisposition, midgut microbiota composition, physiological constraints,

or other biological mechanisms unrelated to fitness may also contribute to shaping host-use

patterns (Gervasi et al., 2016; Yan et al., 2018). Investigating these aspects will be crucial to fully

understand the drivers of host selection in mosquitoes. Future research should aim to unravel

the intricate relationship between feeding patterns, reproductive physiology, and

environmental conditions. Refining our understanding of these dynamics will improve vector

ecology models and enhance predictions of how seasonal variations in host selection may

influence pathogen transmission.

Data availability

365   All data generated and analyzed in this study, along with the accompanying code, are publicly

366   available on Figshare (https://doi.org/10.6084/m9.figshare.28555616).

Acknowledgments

368   We would like to thank Dr. Agustín Quaglia for providing valuable help in statistical analysis. In

369   addition, we greatly appreciate very helpful comments on the manuscript by anonymous

370 reviewers. This study was supported by grant PICT 2018-1172 (FONCYT) and Consolidar Secyt-

371 UNC (2018-2023).

Competing interests

373 The authors declare no competing interests.

Author contributions

Kevin Alen Rucci, Methodology, Software, Validation, Formal analysis, Investigation, Data

Curation, Writing – Original Draft, Writing – Review & Editing, Visualization; Gabriel Barco,

Investigation, Data Curation, Writing – Review & Editing; Andrea Onorato, Investigation, Data

Curation, Writing – Review & Editing; Mauricio Beranek, Investigation, Data Curation, Writing –

Review & Editing; Mariana Pueta, Conceptualization, Methodology, Validation, Writing – Review

& Editing, Visualization, Supervision; Adrián Diaz, , Conceptualization, Methodology, Validation,

Writing – Review & Editing, Visualization, Supervision, Project administration, Funding

acquisition.

References

Alto BW, Connelly CR, O'Meara GF, Hickman D, Karr N. 2014. Reproductive Biology and

Susceptibility of Florida *Culex coronator* to Infection with West Nile Virus. *Vector-Borne and*

Zoonotic Diseases **14**:606–614.

Apperson CS, Harrison BA, Unnasch TR, Hassan HK, Irby WS, Savage HM, Aspen SE, Watson DW,

Rueda LM, Engber BR, Nasci RS. 2002. Host-feeding habits of *Culex* and other mosquitoes

(Diptera: Culicidae) in the Borough of Queens in New York City, with characters and techniques

for identification of *Culex* mosquitoes. *Journal of Medical Entomology* **39**:777–785.

Awahmukalah DST, Brooks MA. 1985. Viability of *Culex pipiens pipiens* eggs affected by nutrition

and aposymbiosis. *Journal of Invertebrate Pathology* **45**:225–230.

Bennett GF. 1970. The influence of the blood meal type on the fecundity of *Aedes* (*Stegomyia*)

aegypti L. (Diptera: Culicidae). *Canadian Journal of Zoology* **48**:539–543.

Beranek MD. 2019. Mosquitos del género *Culex* (Diptera: Culicidae) como vectores del virus

Saint Louis encephalitis (*Flavivirus*) en ambientes urbanos de la ciudad de Córdoba. PhD Thesis.

Universidad Nacional de Córdoba, Córdoba, Argentina

Branda MF, Laurito M, Visintin AM, Almirón WR. 2021. Gonoactivity of *Culex* (*Culex*) (Diptera:

Culicidae) Mosquitoes During Winter in Temperate Argentina. *Journal of Medical Entomology*

58:1454–1458.

Briegel H, Hefti M, DiMarco E. 2002. Lipid metabolism during sequential gonotrophic cycles in

large and small female *Aedes aegypti*. *Journal of Insect Physiology* **48**:547–554.

Briegel H, Timmermann SE. 2001. *Aedes albopictus* (Diptera: Culicidae): physiological aspects of

development and reproduction. *Journal of Medical Entomology* **38**:566–571.

Burkett-Cadena ND, McClure CJW, Ligon RA, Graham SP, Guyer C, Hill GE, Ditchkoff SS, Eubanks

MD, Hassan HK, Unnasch TR. 2011. Host Reproductive Phenology Drives Seasonal Patterns of

Host Use in Mosquitoes. *PLoS One* **6**:e17681.

Caranci AT. 2010. Host switching in *Culex* mosquitoes: Effects of photoperiod and aging on host

choice. PhD Thesis. University of Delaware, Delaware, US.

Ciota AT, Matacchiero AC, Kilpatrick AM, Kramer LD. 2014. The effect of temperature on life

history traits of *Culex* mosquitoes. *Journal of Medical Entomology* **51**:55–62.

Costanzo KS, Schelble S, Jerz K, Keenan M. 2015. The effect of photoperiod on life history and

blood-feeding activity in Aedes albopictus and Aedes aegypti (Diptera: Culicidae). *Journal of*

Vector Ecology **40**:164–171.

Darsie RF, Jr. 1985. Mosquitoes of Argentina. Part I. Keys for identification of adult females and

fourth stage larvae in English and Spanish (Diptera: Culicidae). *Mosquito Systematics* **17**:153–

253.

Demirci B, Durmaz E, Alten B. 2014. Influence of Bloodmeal Source on Reproductive Output of

the Potential West Nile Vector, *Culex theileri* (Diptera: Culicidae). *Journal of Medical Entomology*

51:1312–1316.

Edman JD, Taylor DJ. 1968. *Culex nigripalpus*: Seasonal Shift in the Bird-Mammal Feeding Ratio

in a Mosquito Vector of Human Encephalitis. *Science* **161**:67–68.

Farajollahi A, Fonseca DM, Kramer LD, Marm Kilpatrick A. 2011. "Bird biting" mosquitoes and

human disease: A review of the role of *Culex pipiens* complex mosquitoes in epidemiology.

Infection, Genetics and Evolution **11**:1577–1585.

Fikrig K, Harrington LC. 2021. Understanding and interpreting mosquito blood feeding studies:

the case of *Aedes albopictus*. *Trends in Parasitology* **37**:959–975.

Gervasi SS, Burkett-Cadena N, Burgan SC, Schrey AW, Hassan HK, Unnasch TR, Martin LB. 2016.

Host stress hormones alter vector feeding preferences, success, and productivity. *Proceedings*

of the Royal Society B: Biological Sciences **283**:20161278.

Hancock C, Camp J V. 2022. Habitat-Specific Host Selection Patterns of *Culex quinquefasciatus*

and *Culex nigripalpus* in Florida. *Journal of the American Mosquito Control Association* **38**:89–

91.

Harrison RE, Brown MR, Strand MR. 2021. Whole blood and blood components from vertebrates

differentially affect egg formation in three species of anautogenous mosquitoes. *Parasites &*

Vectors **14**:1–19.

Kasili S, Oyieke F, Wamae A, Bisen C, Mbogo C. 2009. Seasonal changes of infectivity rates of

Bancroftian filariasis vectors in coast province, Kenya. *Journal of Vector Borne Diseases*

46:219224

Kauffman E, Payne A, Franke M, Schmid M, Harris E, Kramer L. 2017. Rearing of *Culex* spp. and

Aedes spp. Mosquitoes. *Bio-Protocol* **7**:e2542.

Kilpatrick AM, Kramer LD, Jones MJ, Marra PP, Daszak P. 2006. West Nile virus epidemics in

North America are driven by shifts in mosquito feeding behavior. *PLoS Biol* **4**:606–610.

Kumle L, Võ MLH, Draschkow D. 2021. Estimating power in (generalized) linear mixed models:

An open introduction and tutorial in R. *Behavior Research Methods* **53**:2528–2543.

Labarthe N, Serrão ML, Melo YF, De Oliveira SJ, Lourenço-de-Oliveira R. 1998. Mosquito

Frequency and Feeding Habits in an Enzootic Canine Dirofilariasis Area in Niterói, State of Rio de

Janeiro, Brazil. *Memórias do Instituto Oswaldo Cruz* **93**:145–154.

Lenth R V. 2017. emmeans: Estimated Marginal Means, aka Least-Squares Means. *CRAN:*

Contributed Packages.

McCann S, Day JF, Allan S, Lord CC. 2009. Age modifies the effect of body size on fecundity in

Culex quinquefasciatus Say (Diptera: Culicidae). *Journal of Vector Ecology* **34**:174-181.

Patrican LA, Hackett LE, Briggs JE, McGowan JW, Unnasch TR, Lee JH. 2007. Host-feeding

patterns of *Culex* mosquitoes in relation to trap habitat. *Emerging Infectious Diseases* **13**:1921–

1923.

R Core Team. 2022. R: A language and environment for statistical computing. Viena, Austria.

Richards SL, Anderson SL, Yost SA. 2012. Effects of blood meal source on the reproduction of

Culex pipiens quinquefasciatus (Diptera: Culicidae). *Journal of Vector Ecology* **37**:1-7

Rivera-Pérez C, Clifton ME, Noriega FG. 2017. How micronutrients influence the physiology of

mosquitoes. *Current Opinion in Insect Science* **23:**112–117.

Santos CS, Pie MR, da Rocha TC, Navarro-Silva MA. 2019. Molecular identification of blood meals

in mosquitoes (Diptera, Culicidae) in urban and forested habitats in southern Brazil. *PLoS One*

14:1–19.

Savage HM, Aggarwal D, Apperson CS, Katholi CR, Gordon E, Hassan HK, Anderson M, Charnetzky

D, McMillen L, Unnasch EA, Unnasch TR. 2007. Host choice and West Nile virus infection rates in

blood-fed mosquitoes, including members of the *Culex pipiens* complex, from Memphis and

Shelby County, Tennessee, 2002-2003. *Vector-Borne and Zoonotic Diseases* **7**:365–386.

Smith JL, Fonseca DM. 2004. Rapid assays for identification of members of the *Culex* (*Culex*)

pipiens complex, their hybrids, and other sibling species (Diptera: Culicidae). *American Journal*

of Tropical Medicine and Hygiene 70:339–345.

Spinsanti LI, Díaz LA, Glatstein N, Arselán S, Morales MA, Farías AA, Fabbri C, Aguilar JJ, Ré V,

Frías M, Almirón WR, Hunsperger E, Siirin M, Da Rosa AT, Tesh RB, Enría D, Contigiani M. 2008.

Human outbreak of St. Louis encephalitis detected in Argentina, 2005. *Journal of Clinical Virology*

42:27–33.

Taborsky M. 2010. Sample Size in the Study of Behaviour. *Ethology* 116:185–202.

Telang A, Skinner J. 2019. Effects of host blood meal source on reproductive output, nutrient

reserves and gut microbiome of West Nile virus vector *Culex quinquefasciatus*. *Journal of Insect*

Physiology **114**:15–22.

Thiemann TC, Wheeler SS, Barker CM, Reisen WK. 2011. Mosquito host selection varies

seasonally with host availability and mosquito density. *PLOS Neglected Tropical Diseases*

5:e1452.

Venables WN, Ripley BD. 2002. Modern Applied Statistics with S. Statistics and Computing.

Springer, New York.

Yan J, Broggi J, Martínez-De La Puente J, Gutiérrez-López R, Gangoso L, Soriguer R, Figuerola J.

2018. Does bird metabolic rate influence mosquito feeding preference? *Parasites & Vectors*

11:1–9.

Yan J, Gangoso L, Martínez-de la Puente J, Soriguer R, Figuerola J. 2017. Avian phenotypic traits

related to feeding preferences in two *Culex* mosquitoes. *The Science of Nature* **104**: 1–10.

Table 1. Reproductive outcomes of *Culex quinquefasciatus* indicating average fecundity, fertility and hatchability for both gonotrophic cycles, accompanied with its respective standard deviation.

Replicate	Blood source	Season	Gonotrophic cycle	No. engorged females	No. egg rafts	Fecundity (eggs/raft)	Fertility (larvae/raft)	Hatchability (larvae/eggs)
1	chicken	autumn	I	163	76	147.47 ± 31.84	133.24 ± 38.19	0.89 ± 0.14
1	chicken	summer	I	116	74	125.64 ± 39.57	109.39 ± 43.66	0.85 ± 0.19
1	mouse	autumn	I	58	23	124.48 ± 20.85	119.96 ± 21.47	0.96 ± 0.04
1	mouse	summer	I	208	120	150.67 ± 29.41	140.38 ± 33.15	0.93 ± 0.10
1	chicken	autumn	II	73	36	142.14 ± 36.59	129.09 ± 43.62	0.87 ± 0.18
1	chicken	summer	II	38	27	138.23 ± 30.27	117.27 ± 37.99	0.85 ± 0.18
1	mouse	autumn	II	23	15	89.13 ± 19.33	73.29 ± 28.09	0.84 ± 0.24
1	mouse	summer	II	92	45	106.68 ± 31.97	99.16 ± 31.18	0.93 ± 0.10
2	chicken	autumn	I	178	89	138.23 ± 37.45	122.7 ± 41.55	0.90 ± 0.15
2	chicken	summer	I	78	51	128.75 ± 38.9	126.08 ± 33.97	0.89 ± 0.13
2	mouse	autumn	I	110	69	103.04 ± 19.01	91.32 ± 22.96	0.89 ± 0.16
2	mouse	summer	I	36	27	113.26 ± 30.66	103.83 ± 34.81	0.89 ± 0.20
2	chicken	autumn	II	86	66	127.4 ± 37.07	116.35 ± 39.9	0.89 ± 0.16
2	chicken	summer	II	48	34	143.68 ± 28.06	136.75 ± 28.05	0.92 ± 0.11
2	mouse	autumn	II	63	52	88.87 ± 19.65	77.86 ± 22.88	0.88 ± 0.15
2	mouse	summer	II	18	14	78.29 ± 25.77	67 ± 25.79	0.87 ± 0.18
3	chicken	autumn	I	150	96	147.22 ± 33.4	129.33 ± 39.45	0.88 ± 0.16
3	chicken	summer	I	67	38	145.03 ± 44.39	136.03 ± 45.49	0.90 ± 0.14
3	mouse	autumn	I	92	54	96.04 ± 23.78	85.76 ± 27.45	0.89 ± 0.18
3	mouse	summer	I	27	20	104.3 ± 21.42	98.35 ± 23.86	0.93 ± 0.09
3	chicken	autumn	II	94	73	144.07 ± 37.49	125.23 ± 39.42	0.87 ± 0.17
3	chicken	summer	II	35	28	156.36 ± 26.71	141 ± 39.53	0.90 ± 0.18
3	mouse	autumn	II	42	29	79.66 ± 19.94	68.76 ± 23.73	0.87 ± 0.21
3	mouse	summer	II	17	11	100.36 ± 22.82	96.1 ± 24.06	0.94 ± 0.06

505 **Table 2.** Analysis of deviance table for the generalized linear mixed model examining the effects
506 of blood source, seasonality, gonotrophic cycle, and their interactions on the fecundity
507 (eggs/raft) of Culex *quinquefasciatus* mosquitoes across 3 replicates. A random intercept for
508 treatment was included to account for variability among the 24 treatments (variance =
509 0.006169). Abbreviations: LRT X^2 = likelihood-ratio test; df = degrees of freedom.

Factor	LRT X^2	df	p-value
blood source	19.22	1	1.16×10^{-5}
seasonality	1.50	1	0.22
gonotrophic cycle	0.47	1	0.49
blood source × seasonality	5.69	1	0.02
blood source × gonotrophic cycle	2.74	1	0.1
seasonality × gonotrophic cycle	2.07	1	0.15
blood source × seasonality × gonotrophic cycle	1.65	1	0.20

510 **Table 3.** Analysis of deviance table for the generalized linear mixed model examining the effects
511 of blood source, seasonality, gonotrophic cycle, and their interactions on the fertility
512 (larvae/raft) of Culex *quinquefasciatus* mosquitoes across 3 replicates. A random intercept for
513 treatment was included to account for variability among the 24 treatments (variance =
514 0.007226). Abbreviations: LRT X^2 = likelihood-ratio test; df = degrees of freedom.

Factor	LRT X^2	df	p-value
blood source	13.30	1	0.0026
seasonality	0.36	1	0.54
gonotrophic cycle	0.28	1	0.59
blood source × seasonality	4.37	1	0.03
blood source × gonotrophic cycle	3.98	1	0.04
seasonality × gonotrophic cycle	0.92	1	0.34
blood source × seasonality × gonotrophic cycle	0.50	1	0.47

515 **Table 4.** Analysis of deviance table for the generalized linear model examining the effects of
516 blood source, seasonality, gonotrophic cycle, replicate and their interactions on the hatchability
517 (larvae/eggs) of Culex *quinquefasciatus* mosquitoes. Abbreviations: LRT X^2 = likelihood-ratio
518 test; df = degrees of freedom.

Factor	LRT X^2	df	p-value
blood source	2.50	1	0.11
seasonality	3.06	1	0.08
gonotrophic cycle	0.005	1	0.94
replicate	1.31	1	0.51
blood source × seasonality	0.12	1	0.72
blood source × gonotrophic cycle	3.64	1	0.06
seasonality × gonotrophic cycle	0.07	1	0.79
blood source × replicate	2.42	1	0.30
seasonality × replicate	2.96	1	0.23
gonotrophic cycle × replicate	0.07	1	0.96
blood source × seasonality × gonotrophic cycle	2.73	1	0.10
blood source × seasonality × replicate	0.02	1	0.99
blood source × gonotrophic cycle × replicate	2.48	1	0.29
seasonality × gonotrophic cycle × replicate	0.43	1	0.81
blood source × seasonality × gonotrophic cycle × replicate	2.57	1	0.28

Figure captions

Figure 1. Interaction plot showing the effect of blood meal source (chicken or mouse) and seasonality (autumn or summer) on fecundity (eggs/raft). Points represent the predicted marginal mean values and the bars its corresponding confidence intervals. Means sharing same letters are not statistically different (p > 0.05).

Figure 2. Interaction plots showing the effect of blood meal source (chicken or mouse) and seasonality (autumn or summer) (**A**) or blood meal source and gonotrophic cycle (first or second) (**B**) on fertility (eggs/raft) of *Culex quinquefasciatus*. Points represent the predicted marginal mean values and the bars its corresponding confidence intervals. Means sharing same letters are not statistically different (p > 0.05).

Figure 3. Dotplots showing the variation of predicted fecundity (eggs/raft) and fertility (larvae/raft) means along the 24 treatments, with combination of blood source (chicken or mouse), seasonality (autumn or summer), gonotrophic cycle (first or second) and replicate (1 to 3).

Figure 4. Boxplots showing   predicted   mean hatchability for the 24 treatments, combining replicate (1-3), blood meal source (chicken or mouse), seasonality (summer or autumn) and gonotrophic cycle (first or second), of *Culex quinquefasciatus*.

Supplementary Material.

Figure SM1. Variation of statistical power for the fecundity and fertility models across several sample sizes.