Timed sequence task: a new paradigm to study motor learning and flexibility in mice

Anna Urushadze[1], Milan Janicek[2], Alice Abbondanza[1,3], Helena Janickova[1]

[1]Laboratory of Neurochemistry, Institute of Physiology of the Czech Academy of Sciences, Prague, 14200, Czech Republic

[2]Central Library, Charles University, Prague, 11636, Czech Republic

[3]Sorbonne Université, CNRS UMR 8246, INSERM U1130, Neuroscience Paris Seine-Institut de Biologie Paris Seine (NPS-IBPS), Paris, 75005, France.

Correspondence should be addressed to: Helena Janickova, Institute of Physiology of the Czech 18 Academy of Sciences, Videnska 1083, 14200, Prague 4, Czech Republic

helena.janickova@fgu.cas.cz

Conflict of interest

Authors report no conflict of interest.

Funding sources

This work was supported by Grant Agency of the Czech Republic Grant 19-07983Y.

Abstract

Motor learning and flexibility allow animals to perform routine actions efficiently while keeping them flexible. There is a number of paradigms used to test cognitive flexibility but not many of them focus specifically on learning of complex motor sequences and their flexibility. While many tests use operant or touchscreen boxes that offer high throughput and reproducibility, the motor actions themselves are mostly simple presses of a designated lever. To focus more on motor actions during the operant task and to probe the flexibility of these well-trained actions, we developed a new operant paradigm for mice, the "timed sequence task". The task requires mice to learn a sequence of lever presses that have to be emitted in precisely defined time limits. After training, the required pressing sequence and/or timing of individual presses is modified to test the ability of mice to alter their previously trained motor actions. We provide a code for the new protocol that can be used and adapted to common types of operant boxes. In addition, we provide a set of scripts that allow automatic extraction and analysis of numerous parameters recorded during each session. We demonstrate that the analysis of multiple performance parameters is necessary for detailed insight into animals' behavior during the task. We validate our paradigm in an experiment using the valproate model of autism as a model of cognitive inflexibility. We show that the valproate mice show superior performance at specific stages of the task, paradoxically due to their propensity to more stereotypic behavior.

Significance Statement

Cognitive flexibility impairment is a crucial component of many neurological disorders and it is frequently evaluated in animal models. As the commonly used tests usually do not focus on motor learning and the ability to adapt motor sequences, we designed a new paradigm to evaluate motor learning and its flexibility. The timed sequence task is automatized and easily accessible as it is based on widely available operant boxes. During the training, the task requires precise timing of each action to force stereotypic performance. Its relative complexity allows detailed analysis of multiple parameters and therefore detailed insight into animal's behavior. The task can be used to reveal and understand subtle differences in motor and operant learning and flexibility.

Graphical Abstract

Introduction

Acquisition of new motor actions and their successful automatization by repeating them is a crucial part of animals' behavior. The animals have to be able to not only learn and execute the motor action but also flexibly change it when required. Various tasks for testing of cognitive flexibility are performed in conventional operant or in touchscreen-based boxes (Mar et al., 2013). Alternatively, a variety of maze-based tasks including reversal Morris water maze and reversal T-maze task (Vorhees and Williams, 2006) can be used. In most of these paradigms, the original motor action is relatively simple (a lever press or a nose poke) and in the flexibility phase, it is simply transformed into its mirror image. In these paradigms, it may be difficult to capture subtle changes in animal's behavior as a simple motor action does not allow for a detailed insight (Keeler et al., 2014). In literature, several elegant paradigms can be found that are suitable for testing more complex motor learning and automatization (Keeler et al., 2014; Geddes et al., 2018; Dhawale et al., 2021; Janickova et al., 2021; Turner et al., 2022; Wolff et al., 2022). However, these tasks usually do not involve flexibility part (although they could be probably adapted to it) and, more importantly, they often require a specialized equipment and/or technical expertise which makes them less accessible. Notably, some of these more complex tasks have been so far performed only with rats and their adaptability to mice may not prove easy.

To overcome the aforementioned difficulties, we created a timed sequence task where mice perform a heterogeneous sequence of four presses in a highly stereotypic way. To ensure this, mice are taught to perform the individual presses within pre-defined time limits, to make the actions in individual trials highly uniform. Once the complex sequence is well trained, it can be repeatedly modified at different steps, to implement the flexibility component of the task. In addition, the task structure allows to collect and analyze numerous behavioral parameters. The present protocol can be realized with commonly available operant boxes equipped only with two non-retractable levers and a feeder. The protocol code is programmed in a proprietary but common programming language that is used with a common type of operant boxes. The code is open and can be easily modified to create new variations of the task. To shorten the tedious analysis of multiple task parameters, we also provide a set of scripts in Python for automatized processing of the data and for user-friendly manual checking of lengthy task logs if needed. As a first choice, we tested our new protocol in the valproate mouse model of autism (Nicolini and Fahnestock, 2018), a model

with a prominent social impairment, impairment of motor skills and a higher propensity to habitual and repetitive behavior (Nicolini and Fahnestock, 2018; Tartaglione et al., 2019). In this model, we show specific alterations in several parameters of the task, together providing evidence of increased habitual performance and slight motor impairment in these animals.

Materials and Methods

Animals, preparation and testing of the VPA model

Male mice of C57BL/6J genetic background were used in all experiments. Mice used for the production of VPA model and control animals were purchased from Charles River Laboratories. They were housed in groups of 2-5 per cage in a room with controlled temperature and 12-hour light cycle (lights ON 6am). All experimental procedures complied with the directive of the European Community Council on the use of laboratory animals (2010/63/EU) and were approved by the animal care committee of the Czech Academy of Sciences. The behavioral testing started at 2 months of age. For the food-motivated testing in the operant box, mice were mildly food restricted to 85-90 % of their free-food weight. Prenatal exposure to valproic acid (VPA) was used to induce autistic phenotype, according to the published protocol (Lucchina and Depino, 2014). Pregnant females were injected subcutaneously at the gestation day 12.5 with VPA sodium salt (600 mg/kg, Sigma) in a volume of 5 ml/kg or, for controls, with vehicle only (saline). In total, we produced 11 control males and 11 males exposed to VPA. To confirm the efficiency of the VPA protocol in inducing autistic phenotype (Ergaz et al., 2016), we tested social behavior by social preference test (Nadler et al., 2004), compulsive behavior by hole-board test (Martos et al., 2017) and dexterity by sticky tape test (Bouet et al., 2009).

Timed sequence task: overview of the procedure

The task can be performed in standard operant boxes equipped with fixed levers. In the timed sequence task, mice have to learn a sequence of lever presses that need to be precisely timed and performed without any error or interruption. The sequence is acquired in several stages as described below. After the reliable acquisition of the task, the sequence and/or the time intervals can be changed in multiple probe sessions to examine the ability to change the previously learned motor program. Each set of probe sessions is followed by baseline sessions to re-establish the

original performance. To perform the task, we used a standard operant box (MED-307A-B1, MedAssociates) placed in a sound-attenuating cubicle and operated by Med-PC V software (MedAssociates). The box was equipped with two fixed levers (left and right) and a feeder located between the levers, for the automatic delivery of reward pellets. Above each lever, a cue light was located indicating when the lever was active and should be pressed. The correct execution of each trial was rewarded with a chocolate cereal pellet (20 mg, Bio-Serv). The main house light in the box was always kept off throughout the whole training. Mice were trained for one session a day, 5 times per week and the completion of the whole training and testing required approximately 40 training/testing days (8 weeks) for an average control mouse. Based on our experience, the majority of animals were able to learn the task and complete the whole procedure without major difficulties. However, we defined criteria in individual stages that would lead to an exclusion of the animal from the study. The exclusion criteria are described in Table 1. The individual stages of the training and testing are listed in Figure 1 and Table 1 and in details described below. The respective codes for each stage are available at https://doi.org/10.5281/zenodo.7881104.

Figure 1: Schematic of the training stages during the timed sequence task.
The training consists of 3 main phases: 1/ basic operant training, 2/ acquisition of the timed LLRR sequence and 3/ alterations of the previously learned timed sequence. In the last stage, each alteration is followed by 1-3 sessions of the baseline timed sequence protocol, to ensure mice return to their baseline performance.

Table 1: Exclusion criteria used for the individual training stages of the task.

Training stage	Exclusion criteria
Habituation 1	No exclusion criteria
Habituation 2	Mouse does not reach performance criteria within 10 sessions
Habituation 3	
Operant training	
LR sequence	Mouse does not reach performance criteria within 10 sessions
LLR sequence	
LLRR sequence	
LLRR with baseline time limits	Mouse does not reach performance criteria within 10 sessions and performs less than 10 trials on the last (10th) session.
LLRR with stricter time limits	
LLRR alternating: 3x LLRR + 1x LLR	No exclusion criteria
LLRR baseline	Mouse performs less than 10 trials on the last (3rd) baseline session.
LLRR LEFT	No exclusion criteria
LLRR baseline	Mouse performs less than 10 trials on the last (3rd) baseline session.
LLRR RIGHT	No exclusion criteria
LLRR baseline	Mouse performs less than 10 trials on the last (3rd) baseline session.
LLRR MIDDLE	No exclusion criteria
LLRR baseline	Mouse performs less than 10 trials on the last (3rd) baseline session.
Extinction	No exclusion criteria

Food restriction

Two weeks before the training initiation, mice were weighed and the range of 85 - 90 % of their original weight was determined as the target weight. Mice were then weighed and fed daily in order to reach and maintain their target weight throughout the whole training and testing procedure.

Timed sequence task: description of individual stages

1st Phase, Operant training

Habituation 1

On the first day of training, mice were placed in the operant box with the feeder off for 10 minutes. No reward was placed in the feeder and all mice automatically advanced to the next training step on the next day.

Habituation 2

At the beginning of this session, four reward pellets were manually placed into the feeder and mice were left in the box for 10 minutes. At the end of the 10-minutes session the pellets were checked and only the mice that consumed all pellets advanced to the next training stage. Otherwise, Habituation 2 session was repeated.

Habituation 3

For the last habituation, mice stayed in the box for 30 minutes while the reward was automatically delivered every 2 minutes. Mice had to consume all 15 reward pellets in order to advance to the next stage.

Operant training

After habituation, mice had to associate lever presses and the delivery of the reinforcement. During the initial operant training, every (both left and right) lever press was rewarded and the levers became active immediately after the reward delivery. The session ended after mouse earned 40 rewards or after 30 minutes, whichever occurred first. After mice successfully earned 40 rewards within a session, they were moved to the next stage.

2nd Phase, LLRR Acquisition

LR sequence

To avoid any side bias that might cause problems in future training stages, mice were trained to perform a left-right sequence from the very beginning. In order to obtain a reward, mice had to complete the whole LR sequence with no errors (that is, sequences "LLR" or "RLR" were not rewarded). After every error, a punishment of 5s time out (TO) interval was introduced before the levers became active again. During the TO interval, potential lever presses were recorded but they had no consequences. During the session, cue light above the respective lever always indicated which lever should be pressed at the given moment. The session ended after 60 minutes or when the mouse earned 40 rewards, whichever occurred first. In order to proceed to the next training stage, mice had to earn 40 rewards within the session for three consecutive days.

LLR sequence

This stage was similar to the previous with the addition of another left press, forming the sequence LLR (left, left, right). The criterion for this training stage was again earning 40 rewards in three consecutive days.

LLRR sequence

The final sequence LLRR (left, left, right, right) consisted of four presses. Again, it had to be performed without any errors in order to be rewarded. Up to this point, intervals between the correct presses were not specified. The criterion was again 40 rewards in three consecutive days.

LLRR with baseline time limits

At this stage, time limits were introduced so that every press had to be emitted within a certain interval after the preceding press. We call the time limits introduced at this stage as "baseline time limits" and this particular timed sequence was used for the re-establishing of the baseline performance throughout the following testing. The example illustration of a correct and incorrect trial at this stage is shown in Figure 2. The time intervals we used were 0.5-5 s between the 1st and 2nd press (L-L), 0.5-20 s between the 2nd and 3rd press (L-R) and again 0.5-5 s between the 3rd and 4th press (R-R). To obtain the reward, the sequence had to be performed with no errors and interruption, within the pre-defined time limits. The cue lights indicated when the respective lever became active after the previous press. Subsequently, the cue light remained on even after the respective time limit for the lever press elapsed. Thus, the cue lights indicated only the beginning but not the end of each particular interval. As in the preceding training stages, the punishment 5 s TO was introduced after every either pressing or timing error. The criterion was defined as earning 40 rewards in three consecutive days.

Figure 2: Illustration of the correct and incorrect trial in the baseline timed LLRR sequence.
Only complete correct sequences performed within the pre-defined time intervals are rewarded. After each incorrect trial (resulting from an incorrect press or a timing error), 5s timeout is introduced.

LLRR with stricter time limits

At this stage, the mice had to perform the sequence within stricter time limits while all other parameters remained the same. The limits were set as 0.5-2 s (L-L), 0.5-10 s (L-R) and 0.5-2 s (R-R). The performance criterion remained the same, that is 40 rewards within a session for three consecutive days. At this stage, some mice started having difficulties to fulfil the criterion. If this was the case, we trained the animals for the maximum of ten days and after that, we moved them automatically to the next stage (also see Table 1 for exclusion criteria).

3rd Phase, LLRR alterations

LLRR alternating: 3x LLRR + 1x LLR

After training mice in the baseline timed sequence and pushing them to their performance limits with the stricter time intervals, we started alternating the pressing sequence and its timing in order to examine the cognitive flexibility of the animals. First, we tested the effect of an alternated sequence but rather than changing the whole sequence, we only changed every fourth sequence during the session to LLR. Therefore, in every fourth trial the last press had to be omitted and potential execution of the second right press was counted as an error. Hence, the mouse could choose either the earlier reward or, alternatively, the completion of the sequence before getting the pellet, in a stereotypic way. The time intervals between the presses were the same as in the baseline LLRR sequence. No performance criteria were set from this stage onwards and all mice were tested in the alternated sequence for three consecutive days. After that, they were run in 1-3 baseline session (LLRR sequence with baseline limits) until they earned 40 rewards at baseline or for the maximum of 3 days.

LLRR LEFT

After re-establishing the baseline performance, we started alternating the time limits for individual presses. First, we changed the interval between the 1st and 2nd press (L-L) to 1.5-5 s, forcing the mice to wait for additional 1 s before the second press. The other intervals between presses remained unchanged (0.5-20 s and 0.5-5 s, respectively). Mice were tested in this setting for three consecutive days and then let to re-baseline before the next stage as described before.

LLRR RIGHT

This time, we changed the last interval between the 3rd and 4th press (R-R) to 1.5-5 s. The other intervals remained the same. Mice were tested for 3 consecutive days before moving them to baseline session(s) and the next stage.

LLRR MIDDLE

Finally, we changed the interval between the two middle presses (L-R) to 3-20 s. After three days of testing, mice were moved to their last baseline session(s) and then to the final stage of the paradigm.

Extinction

On the last day of our paradigm, we submitted mice to the extinction session where no reinforcement was provided. The correct sequence was identical to the LLRR sequence with baseline time limits, including the cue lights and the 5 s TO after incorrect trials, however, no reward was delivered after completing a correct sequence.

Timed sequence task: data analysis

The boxes were operated by the MedPC-V software that was also used to record all events that were defined in the task code. The resulting task log shows all successive events in a row (one row per session) as decimal numbers: the number before decimal point indicates the time of the respective event that elapsed from the last recorded event. The number after the decimal point indicates the specific event using a respective code as defined in the task. The task events recorded in the log included individual lever presses, both correct and incorrect and presses during the TO interval, and reward delivery after the completion of correct sequences. Main summary parameters for each session (e.g. total number of presses) were recorded in a text file generated for each session. For more detailed analysis, we generated a comprehensive task log containing all sessions of the study and analyzed various performance parameters using the log and the analysis scripts created for that purpose. In addition to the individual analysis scripts, we also created a script allowing an easy and quick searching in the extensive task logs and visualization of defined time points during the session. The complete task log containing data from this study can be found at https://doi.org/10.5281/zenodo.7875357. Summary of all task codes and parameters we used is shown in Tables 2-4. The codes and the analysis scripts are available at https://doi.org/10.5281/zenodo.7881059.

Table 2: Task parameters used for the individual stages during the initial training. The depicted parameters follow the interface of the Med-PC V software (MedAssociates). They may slightly differ when adapted to a different type of software.

Setting Task Parameters (1st part)		
	Session time (min)	Maximum reinforcement
Habituation 1	10	
Habituation 2	10	
Habituation 3	30	15
Operant training	30	40

Table 3: Basic task parameters used for the individual stages during the advanced phase of the training. The depicted parameters follow the interface of the Med-PC V software (MedAssociates). They may slightly differ when adapted to a different type of software.

Setting Task Parameters (2nd part)										
	first correct lever (1=left, 2=right)	first correct lever (1=left, 2=right)	reinforcement device (1=pellet, 2=dipper, 3=drug)	reinforcement time (sec)	time out following reinforcement (sec)	session time (min)	first fixed ratio value	second fixed ratio value	maximum reinforcement	softcr data array (1=yes, 0=no)
LR sequence										
LLR sequence										
LLRR sequence										
LLRR with baseline time limits										
LLRR with stricter time limits	1,000	0,000	1,000	0,050	0,000	60,000	2,000	1,000	40,000	1,000
LLRR alternating: 3xLLRR + 1x LLR										
LLRR LEFT										
LLRR Right										
LLRR MIDDLE										
Extinction										

Table 4: Timing task parameters used in advanced stages of the training. The depicted parameters follow the interface of the Med-PC V software (MedAssociates). They may slightly differ when adapted to a different type of software.

Setting Tas Parameters (3rd part)								
	timeout low left	timeout high left	timeout low right	timeout high right	timeout low 1st right	timeout high 1st right	timeout low 2nd right	timeout high 2nd right
LR sequence	0.050 s	3600 s						
LLR sequence	0.050 s	3600 s	0.050 s	3600 s				
LLRR sequence					0.050 s	3600 s	0.050 s	3600 s
LLRR with baseline time limits	0.500 s	5.000 s			0.500 s	20 s	0.500 s	5.000 s
LLRR with stricter time limits	0.500 s	2.000 s				10 s		2.000 s
LLRR alternating: 3x LLRR + 1x LLR		5.000 s			0.500 s			5.000 s
LLRR LEFT	1.500 s					20 s		
LLRR RIGHT	0.500 s	5.000 s				20 s	1.500 s	5.000 s
LLRR MIDDLE	0.500 s				3.000 s		0.500 s	
Extinction					0.500 s		0.500 s	

Statistical analysis

In the experiment, 22 mice were used in total, 11 mice per group (control vs. VPA). Outliers were defined as values outside the range (average +/- 2 x standard deviation) and they were excluded from the analysis. The identified outliers are indicated in figure legends together with respective sample sizes. Statistical analysis was performed with GraphPad Prism version 8 (GraphPad Software). We used two t-test for comparison of two experimental groups and two-way or three-way ANOVA for evaluating the effects of two or three variables. Sidak's test was used for post-hoc comparison. Significance was set at $p < 0.05$ and data in graphs are presented as average +/- SEM. Results of statistical analyses are reported as 95 % confidence intervals for the difference between means (95 % CI) and respective p values.

Data and code accessibility

The code used to run the task is available at https://doi.org/10.5281/zenodo.7881104. The scripts for data analysis are available at https://doi.org/10.5281/zenodo.7881059. The raw data presented in the article are available at https://doi.org/10.5281/zenodo.7875357.

Results

Evaluation of the autism-like phenotype in the VPA mice

First, we used the social preference test, hole-board test and sticky tape test to confirm the mice prenatally exposed to VPA show the typical autism-like phenotype. Results indicating social and dexterity impairment are shown in Fig. 3.

Figure 3: VPA mice show social impairment and reduced dexterity.
(a) Interaction times during the social preference test in VPA and control mice. $n_{(VPA, ctrl)}$=11. Effect of treatment, 95 % CI [14.4; 49.0], p=0.0007; effect of stimulus, 95 % CI [-62.9; -28.3], p<0.0001; two-way ANOVA. (b) Probability of returning to the same hole during the hole board test in VPA and control mice.

n$_{(ctrl, VPA)}$=11. 95 % CI [-0.02; 0.12], p=0.157; two-tailed t test. (c) Time to remove tape in VPA and control mice during the tape removal task. n$_{(ctrl)}$=10 (1 outlier excluded), n$_{(VPA)}$=11. 95 % CI [22.9; 1119.7], p=0.0061; two-tailed t test. All data are shown as mean ± SEM. **, p < 0.01; ***, p < 0.001.

Both control and VPA mice can learn the timed sequence task and they improve during training

All mice finished the training and there was no significant difference between the control and VPA mice in the number of sessions needed to reach the training criteria (Fig. 4a). Both control and VPA mice also significantly improved their performance during training. We compared their performance in the 1st and the last session of the "LLRR baseline" timed paradigm, that is, in the 1st timed training session and in the last baseline session, just before the extinction. While mice were reaching the same number of rewards in both sessions, they were faster after training (Fig. 4b). Likewise, both control and VPA mice also significantly decreased the number of incorrect presses and presses emitted during timeout (TO presses) (Fig. 4c, d). Throughout the training, VPA mice performed less TO presses compared to controls (Fig. 4e, f). To understand the mice activity during the session, we quantified individual types of pressing sequences, correct and incorrect, performed in the 1st and last LLRR baseline session. Unlike the 1st session, in the last session mice performed significantly more correct sequences than any other type of sequence (Fig. 4g, h). During sessions, both control and VPA mice often alternated periods of activity (bouts) and inactivity (pauses). Most of the bouts and pauses were relatively short (less than 5 and 2 min, respectively) and their duration did not differ between control and VPA mice (Fig. 5a, b). We also analyzed intervals between the individual presses in the 1st and last session of the LLRR baseline paradigm, only in correctly executed sequences. Although the required intervals between the two left and the two right presses were identical, both control and VPA mice were significantly faster to emit the second right press, especially after training (Fig. 5c, d). In addition, initially VPA mice showed longer intervals between presses compared to controls but during training, the average inter-press intervals in VPA and controls became indistinguishable (Fig. 5c, d). Finally, we wanted to know if the mice's performance became more stereotypic during training. We checked variation coefficients of individual inter-press intervals during the 1st and the last session of the LLRR baseline paradigm and we found decrease in the variability in the middle interval (L-R) in VPA animals only (Fig. 5e-j).

Figure 4: VPA and control mice learn successfully all stages of the timed sequence task.
(a) Number of training sessions needed to reach performance criteria in individual training stages. (b-d) Cumulative graphs showing number of rewards earned (b), incorrect presses (c) and presses during timeout (effect of session, p<0.0001; time vs treatment interaction, p<0.0001) (d) emitted by VPA and control mice in the 1st (light symbols) and last (dark symbols) session of the baseline timed LLRR sequence. (e, f)

Number of left timeout presses emitted on the 1st (effect of treatment, 95 % CI [8.4; 30.6], p=0.0007) (e) and last (effect of treatment, 95 % CI [4.9; 30.9], p=0.0074) (f) sessions at different stages of training. (g, h) Different types of pressing sequences performed on the 1st (g) and last (h) session of the baseline timed LLRR sequence. Sequence abbreviations: LLRRP - correct sequence followed by a reward ("Pellet"); R - sequence started incorrectly with Right press; LR - second press incorrect Right; LLR̲R - fourth press too fast (below threshold); LLL - third press incorrect Left; LL̲ - second press too fast (below threshold); LL̅ - second press too slow (above threshold); LLRR̅ - fourth press too slow (above threshold); LLRL - fourth press incorrect Left. In all graphs, n(ctrl, VPA)=11. Data are shown as mean ± SEM and they were analyzed by two- (a, e-h) or three-way ANOVA (b-d). **, p < 0.01; ***, p < 0.001, ****, p < 0.0001.

Figure 5: VPA mice show longer inter-press intervals and decreased variability during specific stages of training.
(a-b) Duration of pressing bouts (a) and pauses (b) during the 1st and last session of the baseline timed LLRR sequence. (c-d) Duration of individual inter-press intervals during the 1st (effect of treatment, 95 % CI [-0.376; -0.011], p=0.0381; effect of interval, p=0.0037) (c) and last (effect of treatment, 95 % CI [-0.259; 0.187], p=0.750; effect of interval, p<0.0001) (d) session of the baseline timed LLRR sequence. (e-j) Comparison of the variability of inter-press intervals in VPA (f, h, j) (in h, 95 % CI [-0.452; -0.014], p=0.0391) and control (e, g, i) mice during the 1st and last session of the baseline timed LLRR sequence. In all graphs, n(ctrl, VPA)=11. Data are shown as mean ± SEM and they were analyzed by one- (a, b) or two-way (c, d) ANOVA or by paired t test (e-j). *, p < 0.05.

VPA mice show better adaptation to changed pressing times but not changed pressing sequence

After the mice showed reliable performance in the LLRR baseline paradigm, we manipulated either the sequence structure or the required inter-press intervals. First, we implemented LLRR *alternating* paradigm where in every 4th trial the LLRR sequence was shortened to LLR (Fig. 6a-

d). The occasional shortening of the pressing sequence affected control and VPA mice in a different way. During the 3 days of LLRR *alternating* paradigm, controls decreased the number of premature and increased the number of delayed second right presses while VPA mice did the opposite (Fig. 6c, d). After alternating the sequence, we changed the individual inter-press intervals, specifically by prolonging the lower time limit and forcing animals to wait longer before the next press. First, we increased the lower time limit between the two left presses to 1.5 s which had only modest effect on mice's performance and did not reveal any differences between control and VPA mice (data not shown). Next, we prolonged the lower time limit between the two right presses to 1.5 s. This affected mice's performance but unexpectedly, VPA mice adapted better to this change (Fig. 6e-q) and on the 3rd day, VPA mice emitted less incorrect presses (Fig. 6g, i, o). In particular, VPA mice were less prone to make premature second right press (Fig. 6i, q). Finally, we changed the interval between the left and right middle presses by increasing the lower limit to 3 s which proved to be the most challenging modification. Although VPA mice again adapted easier to this change, the difference between groups was not so prominent (Fig. 7). VPA mice did not make less premature first right presses compared to controls (Fig. 7i) but they were less likely to change the sequence structure by adding 3rd left press (Fig. 7h). Overall, our data suggest that VPA mice can easier adapt to changes in the inter-press intervals but they are less likely to change the number and order of presses in the originally learned sequence.

VPA mice are less affected by the omission of reinforcement in the extinction session

In the end of the experiment, we performed a single extinction session to test if VPA mice were more likely to maintain their performance in the absence of reward, which would be a sign of cognitive inflexibility (Mar et al., 2013). Control and VPA mice on average completed the same number of trials and a similar proportion of mice in both groups did not complete the maximum 40 trials (Fig. 8a, g). In line with that, the number of correct presses in both groups also did not differ (Fig. 8c, i). However, VPA mice emitted less incorrect presses (Fig. 8f, j). They were less prone to start a new sequence with the right press, after the reward failed to be delivered in the end of the sequence (Fig. 8m). Instead, VPA mice responded by prolongation of the last inter-press interval (Fig. 8l), from approximately 1.5 s (Fig. 5d) during the baseline performance to almost 2 s (Fig. 8l). In summary, similar to the previous sessions, VPA mice tended to maintain the original

order of presses in the sequence while having less difficulty to change the time intervals between them.

Figure 6: VPA mice show better adaptation to the change of timing but not when the pressing sequence itself is changed.

(a-d) Performance criteria during the LLRR alternating sessions. Number of left (effect of treatment, 95 % CI [4.69; 70.10], p=0.027) (a) and right (b) presses during timeout and number of fourth presses emitted too fast (c) and too slow (d). $n_{(ctrl)}$=10 (1 outlier excluded from a-d), $n_{(VPA)}$=8-10 (2 outliers excluded from a-b, 1 excluded from c and 3 excluded from d). (e-i) Performance criteria during the three LLRR *RIGHT*

sessions. Number of total (e), correct (f), incorrect (g) and timeout (h) presses in each session. (i) Number of the fourth presses emitted too fast in each session. $n_{(ctrl, VPA)}=11$. (j-q) Performance criteria during the last LLRR *RIGHT* session only. (j) Duration of individual inter-press intervals. (k, l) Number of left (95 % CI [-50.9; -15.9], p=0.0008) (k) and right (95 % CI [-120.1; -1.163], p=0.046) (l) timeout presses. (m-p) Cumulative graphs showing number of total (effect of treatment, 95 % CI [16.6; 234.1], p=0.026) (m), correct (effect of treatment, 95 % CI [2.12; 92.6], p=0.041) (n), incorrect (o) and timeout (effect of treatment, 95 % CI [6.09; 105.1], p=0.030) (p) presses. (q) Different types of pressing sequences emitted. Sequence abbreviations as in Fig. 4. $n_{(ctrl)}=10\text{-}11$ (1 outlier excluded from k), $n_{(VPA)}=10\text{-}11$ (1 outlier excluded from k). Effect of treatment, 95 % CI [1.52; 9.57], p=0.007. Data are shown as mean ± SEM and they were analyzed by two-way ANOVA (a-j, m-q) or t test (k, l). *, p < 0.05; **, p < 0.01; ***, p < 0.001.

Figure 7: VPA mice maintain the original sequence when timing of the presses is changed.
(a-d) Performance criteria during the three LLRR *MIDDLE* sessions. Number of total (a), correct (b), incorrect (c) and timeout (effect of treatment, 95 % CI [8.0; 112.9], p=0.026) (d) presses in each session. (e-n) Performance criteria during the 1st LLRR *MIDDLE* session only. (e) Duration of individual inter-

press intervals. (f, g) Number of left (95 % CI [-31.30; -0.79], p=0.040) (f) and right (95 % CI [-127.7; 15.0], p=0.115) (g) timeout presses. (h) Number of third presses incorrect Left. 95 % CI [-16.36; -3.46], p=0.004). (i) Number of third presses emitted too fast. 95 % CI [-20.69; 5.42], p=0.237. (j-m) Cumulative graphs showing number of total (j), correct (k), incorrect (l) and timeout (effect of treatment, 95 % CI [3.25; 129.9], p=0.040) (m) presses. (n) Different types of pressing sequences emitted during the 1st LLRR *MIDDLE* session. Sequence abbreviations as in Fig. 4. n_(ctrl)=10-11 (1 outlier excluded from f, g), n_(VPA)=11. Data are shown as mean ± SEM and they were analyzed by two-way ANOVA (a-e, j-n) or two-tailed t test (f-i). *, p < 0.05; **, p < 0.01.

Figure 8: VPA mice maintain the original sequence when the reward is omitted during extinction session.

(a-f) Performance parameters during extinction session. Number of completed trials (95 % CI [-10.75; 7.47], p=0.712) (a), total presses (95 % CI [-120.6; 2.10], p=0.058) (b), correct presses (95 % CI [-56.5; 32.3],

p=0.576) (c), left (95 % CI [-20.57; -5.63], p=0.002) (d) and right (95 % CI [-31.83; 0.195], p=0.053) (e) presses during timeout and incorrect presses (95 % CI [-32.6; -3.1], p=0.021) (f). (g-k) Cumulative graphs showing number of completed trials (g), total presses (h), correct presses (i), incorrect presses (effect of treatment, 95 % CI [1.25; 19.27], p=0.028) (j) and presses during timeout (effect of treatment, 95 % CI [4.64; 34.35], p=0.013) (k). (l) Duration of the last inter-press interval. 95 % CI [0.151; 0.667], p=0.0038. (m) Different types of pressing sequences emitted during the Extinction session. Effect of treatment, 95 % CI [0.368; 3.895], p=0.018. Sequence abbreviations as in Fig. 4. $n_{(ctrl)}$=10-11 (1 outlier excluded in d and l), $n_{(VPA)}$=10-11 (1 outlier excluded in d and l). Data are shown as mean ± SEM and they were analyzed by two-tailed t test (a-f, l) or two-way ANOVA (g-k, m). *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Discussion

Here we present a new behavioral paradigm that allows a detailed evaluation of motor learning and flexibility. The design of the new paradigm was motivated by the need to test a complex motor learning and behavior in an automated and reproducible manner, while using a commonly available equipment. We argue that in order to understand behavioral changes in various mouse models, it is necessary to evaluate multiple parameters in a given behavioral paradigm. However, in the operant tasks, often only a few primary parameters such as accuracy or total number of presses are commonly analyzed which may lead to disregard of more subtle differences in behavioral strategies or to overlook of potentially informative details in mice behavior. For instance, we showed that in the VPA mice, the total number of earned rewards was indistinguishable from controls at all stages of the training. Upon closer look however, the performance of the two groups was quite different, suggesting slower motor performance and decreased ability to adapt the originally learned sequence. Furthermore, possibly the most significant difference between the VPA and control mice was the increased number of TO presses, present at almost every stage of the training (Fig. 4e, f). Even though the difference is quite robust, it may easily escape attention as the behavior occurring between the trials is often ignored (Jeong et al., 2020). Of note, the inter-trial intervals themselves and their duration are known to have a major impact on learning in animals and human subjects, including those with autism (Koegel et al., 1980; Auchter et al., 2017; Jeong et al., 2020).

To obtain a detailed insight into mice learning and flexibility, we trained them in a relatively complex pressing sequence. In the timed sequence task, we take advantage of gradually increasing difficulty at different stages of the task to reveal potentially gross or more subtle behavioral impairment and we analyze multiple parameters to better understand its character. During the

training, mice have to employ basic operant learning, motor learning, time perception, habit formation, behavioral flexibility and motivation that can be analyzed at different stages of the task. Importantly, even with only two fixed levers, the task consists of reward-distal and reward-proximal elements and allows to analyze them separately (Keeler et al., 2014). Although the two left and the two right presses were seemingly symmetrical and the required pressing intervals between them were identical at baseline, mice were clearly able to distinguish their distinct reward proximity as the average inter-press interval between the two right presses was significantly shorter (Fig. 5d). In order to use the timed sequence task for the evaluation of flexibility, we need to make sure the mice performance is becoming automated and habitual during the training. To test this, we compared the variability of individual pressing intervals in the 1st and the last "LLRR baseline" session (Fig. 5e-j). Surprisingly, we did not find any decrease of variability in control animals, possibly because the pressing intervals were largely constricted by the pre-defined criteria and there was little space for further improvement. In addition to the low variability of individual motor actions, the automated character of behavior is indicated by an increase in speed and accuracy (Turner et al., 2022). In our experiment, we observed a significant increase in accuracy (Fig. 4c) and efficiency, as mice performed markedly less TO presses in later stages of the training (Fig. 4d).

There are currently several alternatives that allow a detailed testing of motor or operant learning and flexibility. Among those, the touchscreen-based systems offer a great number of pre-designed paradigms that can be modified and allows creating additional, custom made tasks (Horner et al., 2013; Mar et al., 2013; Heath et al., 2016; Janickova et al., 2021). However, a significant cost of such systems may be prohibitive for some users and it also makes it less attractive for occasional users that are not specializing in this kind of behavioral testing. Standard operant boxes, on the other hand, are easily accessible to most laboratories. We show here that even the most basic operant box equipped with two fixed levers and one feeder can be used to perform a relatively complex task and collect detailed behavioral data. We provide a computer code for the task itself and also a set of scripts for data analysis, allowing easy and fast processing of large data sets.

In summary, we provide a new easily applicable paradigm for detailed testing of motor learning and flexibility. We provide tools for data analysis, arguing that multiple parameters should be used to evaluate the animals' performance in a complex task. Finally, we provide an example showing

that our paradigm can reveal subtle differences in mice performance and help to identify general behavioral mechanisms underlying those differences.

References

Auchter A, Cormack LK, Niv Y, Gonzalez-Lima F, Monfils MH (2017) Reconsolidation-Extinction Interactions in Fear Memory Attenuation: The Role of Inter-Trial Interval Variability. Front Behav Neurosci 11:2.

Bouet V, Boulouard M, Toutain J, Divoux D, Bernaudin M, Schumann-Bard P, Freret T (2009) The adhesive removal test: a sensitive method to assess sensorimotor deficits in mice. Nat Protoc 4:1560–1564.

Dhawale AK, Wolff SBE, Ko R, Ölveczky BP (2021) The basal ganglia control the detailed kinematics of learned motor skills. Nat Neurosci 24:1256–1269.

Ergaz Z, Weinstein-Fudim L, Ornoy A (2016) Genetic and non-genetic animal models for autism spectrum disorders (ASD). Reprod Toxicol 64:116–140.

Geddes CE, Li H, Jin X (2018) Optogenetic Editing Reveals the Hierarchical Organization of Learned Action Sequences. Cell 174:32-43.e15.

Heath CJ, Phillips BU, Bussey TJ, Saksida LM (2016) Measuring Motivation and Reward-Related Decision Making in the Rodent Operant Touchscreen System. Current Protocols in Neuroscience, pp 8.34.1-8.34.20.

Horner AE, Heath CJ, Hvoslef-Eide M, Kent BA, Kim CH, Nilsson SRO, Alsi? J, Oomen CA, Holmes A, Saksida LM, Bussey TJ (2013) The touchscreen operant platform for testing learning and memory in rats and mice. Nat Protoc 8:1961–1984.

Janickova H, Kljakic O, Robbins TW, Saksida LM, Bussey TJ, Prado VF, Prado MAM (2021) Evaluating Sequential Response Learning in the Rodent Operant Touchscreen System. Curr Protoc 1:e268.

Jeong M-J, Lee C, Sung K, Jung JH, Pyo JH, Kim J-H (2020) Fear response-based prediction for stress susceptibility to PTSD-like phenotypes. Mol Brain 13:134.

Keeler JF, Pretsell DO, Robbins TW (2014) Functional implications of dopamine D1 vs. D2 receptors: A 'prepare and select' model of the striatal direct vs. indirect pathways. Neuroscience 282:156–175.

Koegel RL, Dunlap G, Dyer K (1980) Intertrial interval duration and learning in autistic children. J Appl Behav Anal 13:91–99.

Lucchina L, Depino AM (2014) Altered peripheral and central inflammatory responses in a mouse model of autism. Autism Res 7:273–289.

Mar AC, Horner AE, Nilsson SRO, Alsiö J, Kent BA, Kim CH, Holmes A, Saksida LM, Bussey TJ (2013) The touchscreen operant platform for assessing executive function in rats and mice. Nat Protoc 8:1985–2005.

Martos Y V., Braz BY, Beccaria JP, Murer MG, Belforte JE (2017) Compulsive Social Behavior Emerges after Selective Ablation of Striatal Cholinergic Interneurons. J Neurosci 37:2849–2858.

Nadler JJ, Moy SS, Dold G, Simmons N, Perez A, Young NB, Barbaro RP, Piven J, Magnuson TR, Crawley JN, Crawley JN (2004) Automated apparatus for quantitation of social approach behaviors in mice. Genes, Brain Behav 3:303–314.

Nicolini C, Fahnestock M (2018) The valproic acid-induced rodent model of autism. Exp Neurol 299:217–227.

Tartaglione AM, Schiavi S, Calamandrei G, Trezza V (2019) Prenatal valproate in rodents as a tool to understand the neural underpinnings of social dysfunctions in autism spectrum disorder. Neuropharmacology 159:107477.

Turner KM, Svegborn A, Langguth M, McKenzie C, Robbins TW (2022) Opposing Roles of the Dorsolateral and Dorsomedial Striatum in the Acquisition of Skilled Action Sequencing in Rats. J Neurosci 42:2039–2051.

Vorhees C V, Williams MT (2006) Morris water maze: procedures for assessing spatial and related forms of learning and memory. Nat Protoc 1:848–858.

Wolff SBE, Ko R, Ölveczky BP (2022) Distinct roles for motor cortical and thalamic inputs to striatum during motor skill learning and execution. Sci Adv 8:eabk0231.