Distinct functional constraints driving conservation of the cofilin N-terminal regulatory tail

Joel Sexton[1], Tony Potchernikov[2], Jeffrey P. Bibeau[2], Gabriela Casanova Sepúlveda[2], Wenxiang Cao[2], Hua Jane Lou[1], Titus J. Boggon[1,2], Enrique M. De La Cruz[2], and Benjamin E. Turk[1,*]

[1]Department of Pharmacology, Yale School of Medicine, New Haven, CT 06520

[2]Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT 06520

*Corresponding author. Email: ben.turk@yale.edu

Summary

Cofilin family proteins have essential roles in remodeling the cytoskeleton through filamentous actin depolymerization and severing. The short unstructured N-terminal region of cofilin is critical for actin binding and harbors the major site of inhibitory phosphorylation. Atypically for a disordered sequence, the N-terminal region is highly conserved, but the aspects of cofilin functionality driving this conservation are not clear. Here, we screened a library of 16,000 human cofilin N-terminal sequence variants for their capacity to support growth in *S. cerevisiae* in the presence or absence of the upstream regulator LIM kinase. Results from the screen and subsequent biochemical analysis of individual variants revealed distinct sequence requirements for actin binding and regulation by LIM kinase. LIM kinase recognition only partly explained sequence constraints on phosphoregulation, which were instead driven to a large extent by the capacity for phosphorylation to inactivate cofilin. The individual sequence requirements for cofilin function and regulation were remarkably loose when examined separately, but collectively restricted the N-terminus to sequences found in natural cofilins. Our results illustrate how a regulatory phosphorylation site can balance potentially competing sequence requirements for function and regulation.

Introduction

Phosphorylation is perhaps the most common reversible post-translational protein modification and has widespread roles in regulating cell behavior. Sites of phosphorylation evolve rapidly, in part because kinase recognition motifs can be introduced by a small number of mutations in flanking residues. However, regulatory sites of phosphorylation are necessarily under additional sequence constraints beyond those required for kinase recognition. For example, intracellular environments harbor many protein kinases – up to ~500 for human cells. Accordingly, just as the sequence surrounding the site must be tuned to its cognate kinase, there is also selective pressure to evade phosphorylation by the "wrong" kinases. Furthermore, regulatory phosphorylation often occurs in functionally important regions, so that variability in the local sequence may be limited to maintain basic function. Finally, the specific sequence context of a phosphorylation site may dictate how phosphorylation modifies protein behavior, for example by generating a binding site for a phosphopeptide interaction module. These factors are potentially in competition with one another, but there is currently limited insight into how they are balanced in any given phosphorylation site.

One example of a phosphoregulatory event occurring at a highly conserved site involves proteins in the cofilin/ADF (actin depolymerizing factor) family (hereafter referred to collectively as cofilin). Cofilin is a small actin-binding protein important for dynamic regulation of the actin cytoskeleton in eukaryotes[1-4]. Cofilin binds cooperatively to actin filaments, promoting severing at junctions between bare and decorated segments, as well as depolymerization, to produce new filament ends and a pool of actin monomers for reassembly[5-10]. Cofilin-dependent cytoskeletal remodeling is essential for multiple actin-

dependent processes, including cytokinesis, cell polarity, migration, neurite extension, and mechanosensing[3].

Due to its major impact on the actin cytoskeleton, cofilin activity is tightly regulated by several mechanisms. Principal among these is phosphorylation at Ser3 within its disordered N-terminal region, catalyzed in most animals and some protists by members of the LIM kinase family (LIMK1/2 and TESK1/2). The N-terminal tail is an essential part of the actin-binding interface, and Ser3 phosphorylation inactivates cofilin by dramatically decreasing its affinity for actin[11]. The phosphomimetic cofilin[S3D] mutant binds to filamentous actin weakly but with higher cooperativity than the WT protein, and severs filaments poorly even when bound[12].

The N-terminus of animal cofilins contains a conserved acetyl-Ala2-Ser3-Gly4-Val5 sequence (by convention, we use residue numbering based on the unprocessed precursor)[3]. Orthologs from other kingdoms vary in the sequence upstream of Ser3 and may have other β-branched or aliphatic residues in place of Val5, but the phosphoacceptor residue and Gly4 are invariant. Factors driving conservation of this sequence are currently unclear. Aside from full N-terminal deletion mutants that are non-functional, only the phosphomimetic S3D and phosphorylation resistant S3A variants have been examined with respect to their effects on actin in vitro or in cells[11-15]. Recognition by LIMK occurs through a docking interaction between the cofilin globular domain and a site adjacent to the kinase catalytic cleft on the kinase C-terminal lobe[16]. Consequently, the N-terminal region projects into the LIMK active site in a non-canonical orientation that lacks complementary binding interactions typically observed in other protein kinase-substrate complexes. Limited mutational analyses have suggested loose

sequence requirements surrounding the phosphorylation site, save for an unusually strict exclusion of Thr as a phosphoacceptor residue[17]. Attempts to identify a LIMK phosphorylation site sequence motif using peptide libraries have been unsuccessful, presumably because the docking interaction is essential for efficient phosphorylation[16,18].

A platform for evaluation of large numbers of N-terminal sequence variants would facilitate discovery of those features important for the various aspects of cofilin functionality. Here we exploit the widespread requirement for cofilin to establish a budding yeast system for high-throughput evaluation of sequence variants. *S. cerevisiae* lacking their sole cofilin ortholog, Cof1, are inviable[19,20], but ectopic overexpression of mammalian cofilin-1 can rescue growth[16]. In this context, we have reported that inducible expression of human LIMK1 suppresses growth in a manner dependent on Ser3 phosphorylation. With this system, we can evaluate in parallel the impact of a mutation on either core cofilin function or its regulation by LIMK based on its ability to support growth in the absence or presence of LIMK induction, respectively. Here, we have applied the system to screen a comprehensive collection of N-terminal sequence variants, and subsequently conducted biochemical analysis of a panel of mutants. We found unexpectedly loose sequence requirements for each of the various aspects of cofilin function and regulation, with each dominated by its own unique position within the N-terminal sequence. Overall conservation of the N-terminal region thus appears to be constrained by the collective requirements imposed by the full complement of cofilin functionality. These observations may have general implications for how regulatory phosphorylation sites emerge and are maintained through evolution.

Results

Dual-functionality cofilin library yeast competitive growth screen

To define cofilin N-terminal sequence requirements for actin severing and phosphoregulation, we leveraged the capacity of mammalian cofilin to substitute for yeast Cof1 in supporting growth.[21] We used a yeast strain in which the endogenous *COF1* gene is placed under control of a tetracycline-repressible promoter (TeTO$_7$-*COF1*).[22] This strain grew poorly in the presence of doxycycline (DOX) due to loss of endogenous Cof1 expression, but ectopic expression of either yeast Cof1 or human cofilin-1 rescued growth (**Fig. 1A**). Our previous work with a temperature sensitive *cof1* mutant strain demonstrated that ectopic expression of the LIMK1 catalytic domain (LIMK1^{CAT}) completely blocked growth in the context of human cofilin-1[16]. Likewise, we found that inducible expression of LIMK1^{CAT} inhibited growth of TeTO$_7$-*COF1* yeast expressing human cofilin-1 in a manner dependent on its phosphorylation at Ser3 (**Fig. S1**). Expression of full length (FL) LIMK1, which has lower cofilin kinase activity than LIMK1^{CAT}, caused only partial growth suppression (**Fig. 1B**). We chose to use FL LIMK1 for our screens as lower activity should allow us to better distinguish cofilin mutants that impact LIMK regulation to varying degrees. We next constructed a saturation mutagenesis library of the human cofilin-1 N-terminal region in which Ala2, Gly4, and Val5 were combinatorially randomized as all twenty potential canonical amino acids (**Fig. 1C**). Ser3 was maintained as either a serine or threonine residue to preserve a potential phosphorylation site, producing a library with 16,000 (20^3 x 2) unique sequences. The library was introduced into a constitutive yeast expression plasmid with an N-terminal His$_6$ tag included to prevent potential sequence-dependent post-translational modifications

(e.g. initiator methionine removal and acetylation) and to help normalize levels of expression. The His$_6$ tag did not prevent the ability of cofilin to rescue yeast growth, sever actin, or be phosphorylated by LIMK1 (**Fig. 1A, Fig. S2**). TeTO$_7$-*COF1* yeast harboring the LIMK1 plasmid were transformed with the cofilin library and expanded in liquid media with a portion of the culture reserved as a starting point sample. The remaining cells were grown in raffinose to derepress the LIMK1 expression construct and treated with DOX to repress Cof1 expression. The culture was then split and propagated in either glucose or galactose to repress or induce LIMK1 expression, respectively (**Fig. 1D**). At various times, the cultures were sampled for plasmid DNA extraction. The variable library sequence within the recovered plasmids was PCR-amplified and subjected to next-generation sequencing. This analysis indicated how the relative abundance of each component of the library changed over time (**Fig. 1E**, **Dataset S1**). In the context of competitive growth, DOX + glucose conditions should enrich for viable cells expressing functional library variants. Growth in DOX + galactose should deplete the culture of functional cofilin variants that could be phosphorylated and inhibited by LIMK1. We first analyzed our data from the DOX + glucose condition to identify core determinants of cofilin function.

Identification of functional constraints on the cofilin N-terminal sequence in supporting growth

Yeast grown in glucose were analyzed for enrichment of cofilin sequences that could presumably bind and sever actin filaments. We compared sequence representation between the beginning of culture in glucose and ~6 population doublings later (**Fig. 1E and 2A**), beyond which changes were less pronounced. We ranked each sequence by

its average log$_2$ fold change in normalized read count across three independently performed replicate screens (**Fig. 2A**). When ranked this way, 4,099 of the 16,000 library sequences were enriched in the population, suggesting that the majority of N-terminal substitutions were deleterious to cofilin function. As anticipated, native cofilin sequences performed well, including those from yeast (Arg-Ser-Gly-Val, rank 27), human (Ala-Ser-Gly-Val, rank 87) and *Arabidopsis* (Ala-Ser-Gly-Met, rank 262), though the most enriched sequence was non-native (Phe-Thr-Gly-Leu) (**Fig. 2A, Dataset S1**). Examination of the 400 most enriched sequences overall identified a Leu-X-Gly-Met consensus motif. (**Fig. S3A**). Within the N-terminal sequence, the fifth position was the most stringently selective, with aliphatic or β-branched amino acids (73% of enriched sequences) being apparently required for function (**Fig. 2B,C,D**) and Met being most commonly enriched (17% of enriched sequences). By comparison, other positions within the sequence were less selective. Despite being highly conserved in plants and animals, the second residue (Ala2 in human) was effectively indiscriminate, with modest enrichment for non-native hydrophobic over charged residues (**Fig. 2D**). Likewise, enriched sequences had roughly equal representation of Ser and Thr at the phosphoacceptor (third) position. Finally, selection at the fourth residue was intermediate. As found in all wild-type (WT) cofilin sequences, Gly was overrepresented at this position in enriched sequences (8% overall). Despite being absent from all native cofilin sequences, hydrophobic residues were generally selected at this position, with small residues and charged residues being depleted.

Because our combinatorial library includes all possible combinations of residues at the random positions, it has the potential to reveal cases where there is cooperativity

between specific pairs of residues. In such cases, two residues would be co-enriched more frequently than expected if they behaved independently. We found several instances where residues appeared to co-vary in this manner. For example, Leu and Gly residues are represented similarly at position 4 within the 400 most enriched sequences. However, Gly4 sequences showed enhanced selection of Ile or Val residues at position 5, while Leu4 sequences were more likely to be co-enriched with Met5 (**Fig. S3A-C**). Likewise, sequences with Val5 more prominently included Gly4 compared to those sequences containing Met5 (**Fig. S3D,E**). The Gly4-Val5 pair, which is found invariably in animals and fungi, was uniquely over-represented within these most highly enriched sequences (**Fig. S3F**) and may provide a fitness advantage over sequences combining independently favorable residues. We also noted covariation of Thr5 with Thr3 at the phosphoacceptor position, where there was a large difference in mean $\log_2$ fold change for Thr3/Thr5 variants compared with Ser3/Thr5 variants (1.7 vs. -0.28) (**Fig. S4**).

To confirm results from the screen, we selected a number of variants with varying degrees of enrichment for further analysis (**Table 1**). We initially assayed these mutants for their ability to support yeast growth on solid media, a format distinct from the competitive growth scheme used for our screen. Human cofilin-1 single or double point mutants were expressed in TeTO$_7$-*COF1* yeast and grown with or without DOX to control endogeneous Cof1 expression. In keeping with their enrichment in the screen, cofilin^{V5M} supported growth similarly to WT human cofilin (**Fig. 3A**), while yeast expressing cofilin^{V5E} failed to grow. In contrast, Ala2 mutations had little effect on yeast growth, with both cofilin^{A2L} and cofilin^{A2D} supporting growth in the presence of DOX. Growth of yeast expressing the "consensus" double mutant cofilin^{A2L,V5M}, was indistinguishable from that

of yeast expressing cofilin[WT] or cofilin[V5M]. Substitutions to Gly4 displayed a range of growth phenotypes that generally correlated with behavior in the screen (**Fig. 3B**). Yeast expressing cofilin[G4L] grew similarly to yeast with cofilin[WT]. Cofilin[G4A], which was slightly depleted in the screen, and cofilin[G4F], which was in the bottom half of enriched sequences, each supported growth at a rate ~5-fold slower than cofilin[WT]. The more substantially depleted variants, G4P, G4E and G4K, provided at best a marginal amount of growth. Overall, we found that among variants enriched in the context of competitive growth in liquid culture, apparent differences in fitness did not translate into overt differences in growth rate on solid media. Furthermore, variants that can support growth on solid media but at lower rates than cofilin[WT] were likely depleted in the screen due to being outcompeted by higher fitness variants. We note that the extent of growth rescue did not generally correlate with the level of cofilin expression (**Fig. S5**). However, one of the most damaging mutations, cofilin[V5E], expressed to very low levels suggesting that it and perhaps other non-functional Val5 substitutions may destabilize the protein.

We next examined how the ability of a cofilin variant to support growth correlated with its biochemical behavior *in vitro*. We expressed and purified cofilin mutants using a system that produces untagged protein with the native N-terminal sequence (initiating with Ala2 or the desired substitution). We initially evaluated the panel of mutants for binding to actin filaments measured in a fluorescence quenching assay, and further analyzed a subset of these mutants for their capacity to sever actin filaments in a TIRF microscopy-based assay. Phosphorylated cofilin[WT] and phospho-mimetic cofilin[S3D] were used as weak binding controls.[12,15,23] We used the cofilin concentration producing half maximal occupancy ($K_{0.5}$) as a proxy for relative overall binding affinity[24]. Cofilin[WT] bound

to actin filaments with a $K_{0.5}$ approximately 0.6 µM (**Fig. 3C-E, Table 1**) whereas cofilin[S3D] required ~8-fold higher concentrations to bind actin filaments. As anticipated, phosphorylated cofilin[WT] displayed almost no detectable binding at the concentrations tested. Almost all mutants that had supported growth on solid media, including the G4F, A2D, V5M and the double mutant A2L/V5M, displayed actin binding comparable to cofilin[WT] (**Fig. 3**), and where tested these mutants likewise severed actin filaments similarly to WT (**Fig. S6-7**). Cofilin[G4A], which supported growth at a lower rate than cofilin[WT], showed somewhat weaker actin binding ($K_{0.5}$ = 3.2 µM), as did the G4K variant that failed to support growth ($K_{0.5}$ = 3.5 µM). Of all variants tested, only cofilin[G4E] showed essentially no binding to actin filaments (**Fig. 3E**), and accordingly it had no detectable severing activity (**Fig. 6**). Consistent with its low level of expression in yeast, cofilin[V5E] formed soluble aggregates when expressed in bacteria, and we were unable to isolate monomeric protein. Overall, we found a general correlation between yeast growth rate and actin binding, with a threshold $K_{0.5}$ of ~2 µM being required to support maximal growth.

Identification of cofilin variants inhibited by LIMK1

To identify cofilin variants susceptible or resistant to LIMK1, we examined the relative abundance of functional variants in our screen following induction of the kinase by growth in galactose. We noted that depletion of the cofilin[WT] sequence from the culture lagged 1 – 2 population doublings behind induction of LIMK1. We therefore followed variant representation from this time point onward over the course of 10 population doublings in the presence of either glucose or galactose (**Fig. 1E**), and ranked sequences by the log$_2$-

fold change in abundance. We found that that approximately 1/3rd of functional cofilin sequences (those enriched in glucose) became depleted in the presence of galactose, suggesting that they were phosphorylated and inhibited by LIMK1, with the remaining 2/3rds persisting in the presence of galactose and thus presumably resistant to LIMK1 expression (**Dataset S1 and Fig. S8**). Functional sequences depleted in galactose overwhelmingly had Ser3 as the phosphoacceptor residue and were dominated by small residues at the Gly4 position, while conversely the majority of LIMK1-resistant sequences contained Thr3 and large hydrophobic residues (primarily Leu and Phe) in position 4 (**Fig. 4A,B and Fig. S8**).

We initially examined the impact of phosphorylation site residue substitutions LIMK1-induced phosphorylation and growth inhibition in yeast (**Fig. 5A**). In contrast to yeast expressing cofilin[WT] (with Ser3), those expressing cofilin[S3T] were unaffected by LIMK1 induction and grew similarly to yeast harboring cofilin[S3A]. Because our cofilin library did not include Tyr3 variants, we also examined growth of yeast expressing cofilin[S3Y]. While cofilin[S3Y] only partially rescued growth in the presence of DOX, growth was further inhibited by galactose-induced expression of LIMK1. Matching these observations, both cofilin[WT] and cofilin[S3Y] became phosphorylated upon LIMK1 induction (**Fig. 5B**) as judged by mobility shift on Phos-tag SDS-PAGE, while only a trace amount of cofilin[S3T] phosphorylation was observed. In similar experiments performed with yeast expressing the hyperactive LIMK1[CAT], cofilin[S3T] was still phosphorylated to only a small extent and fully supported growth (**Fig. S9**). Consistent with these results and a recent report[17], LIMK1 phosphorylation of cofilin was blocked by either S3A or S3T mutation, and was reduced but not eliminated when Ser3 was substituted with Tyr (**Fig. 5D**). LIMK2

was similarly unable to phosphorylate cofilin[S3T] (**Fig. S9D**), indicating that this strong phosphoacceptor residue preference is conserved across isoforms. Collectively, these results confirm that while LIMKs are "dual-specificity" kinases in the sense that they can phosphorylate both Ser and Tyr, they are almost entirely devoid of activity on Thr residues.

Cofilin/ADF proteins are members of a larger family of actin-binding proteins having an ADF fold. Theoretically other ADF domain proteins could be regulated by phosphorylation at sites analogous to Ser3. For example, the N-terminal region of twinfilin proteins, which consist of two ADF domains arranged in tandem, has a potential phosphoacceptor residue at position 5, analogous to cofilin Ser3 (**Fig. 5F**). Notably, this residue is Thr in mammalian twinfilins, but is a Ser in the single ortholog from budding yeast, which lack LIMKs. Furthermore, the LIMK docking region of cofilin centered on helix α5 is well conserved to the N-terminal ADF domain of twinfilin (**Fig. S10**).[16] We therefore wondered whether the identity of the phosphoacceptor residue might serve to insulate twinfilin from phosphoregulation by LIMKs. We thus examined if LIMK1 could phosphorylate mouse or budding yeast twinfilin *in vitro* and whether phosphorylation was influenced by the identity of the putative phosphoacceptor. We found that LIMK1 robustly phosphorylated budding yeast twinfilin at ~40% of its cofilin phosphorylation rate yet had barely detectable activity on mouse twinfilin-1 (**Fig. 5G**). As anticipated, substituting Ser5 of yeast twinfilin with Thr abolished phosphorylation, confirming Ser5 to be the site of LIMK1 phosphorylation and suggesting flexibility in the identity of that residue in an organism lacking LIMKs. However, LIMK1 did not substantially phosphorylate mouse twinfilin-1[T5S], indicating a Ser phosphoacceptor residue is necessary but not sufficient for

LIMK phosphorylation of ADF fold proteins. Collectively these results suggest that insulating twinfilin from inhibition by LIMK1 involves both the phosphoacceptor and additional structural features that prevent phosphorylation.

Identification of phosphoregulatory constraints on the cofilin N-terminal sequence

In our screen, LIMK1-resistant variants were dominated by those having a Thr phosphoacceptor residue. However, Ser3 sequences having Leu, Phe or Trp at position 4 also persisted despite LIMK1 expression, while Ser3 sequences depleted upon LIMK1 induction were enriched for Gly and other small residues at position 4 (**Fig. 4B**). We confirmed in growth assays on solid media that cofilin^{G4L} and cofilin^{G4F}, but not cofilin^{G4A}, protected yeast from LIMK1-induced growth inhibition (**Fig. 6A, Fig. S11**). The most straightforward explanation for these results would be if variants with bulky hydrophobic residues at position 4 were poor substrates of LIMK1. However, we found instead that cofilin^{G4L} and cofilin^{G4F} became highly phosphorylated upon expression of LIMK1 in yeast, as did cofilin^{G4A} (**Fig. 6B,C**). Furthermore, LIMK1 phosphorylated cofilin^{G4A}, cofilin^{G4L}, and cofilin^{G4F} at rates faster than cofilin^{WT} *in vitro*, while cofilin^{G4P} was not detectably phosphorylated (**Fig. 6D**). A comparison of steady-state kinetic parameters confirmed that cofilin^{G4F} (k_{cat} = 0.55 ± 0.05 s^{-1}, K_M = 12 ± 3 µM) was a more efficient LIMK1 substrate than cofilin^{WT} (k_{cat} = 0.20 ± 0.03 s^{-1}, K_M = 9 ± 3 µM) over the full range of substrate concentrations (**Fig. 6F,G**). The LIMK1 preference for cofilin^{G4F} over its wild-type substrate and the extent of cofilin^{G4F} phosphorylation *in vivo* suggested that the cofilin^{G4F} growth phenotype in yeast was not due to reduced phosphorylation.

As cofilin[G4F] is not impaired as a LIMK1 substrate, an alternative explanation for its ability to support growth in the presence of LIMK1 is that it alleviates the inhibitory effect of Ser3 phosphorylation. To investigate this possibility, we used cofilin variants harboring a phosphomimetic S3D mutation to model constitutive phosphorylation. Consistent with its inability to sever actin, cofilin[S3D] as previously shown[11] was unable to support yeast growth (**Fig. 7A**). In contrast, yeast expressing cofilin[S3D,G4F] grew at an equivalent rate to those expressing the cofilin[G4F] single mutant. When examined *in vitro,* cofilin[S3D,G4F] bound pyrene-actin similar to cofilin[G4F], effectively correcting the phospho-mimetic S3D substitution (**Fig. 7C**). Together, these results suggest that the presence of a large hydrophobic residue immediately downstream of the phosphoacceptor residue uncouples cofilin inhibition from phosphorylation. Accordingly, the absolute conservation of a Gly residue at position 4 may be due at least in part because it is required for phosphoregulation of actin binding and severing.

Discussion

Cofilin regulation of actin depolymerization is essential for normal cytoskeletal function. Much of the functional regulation of cofilin converges on its N-terminal site of phosphorylation, yet how this short four amino acid stretch simultaneously impacts its relationship with its regulators, the LIM kinases, and its effector, actin, have not been explored. We used a yeast-based dual-functionality screen of a cofilin N-sequence library and subsequent functional analysis to simultaneously determine the fitness of all possible combinations of N-terminal residues and to assess the importance of each of these amino acids. We identified specific features required for maintenance of cofilin core functionality, targeting of cofilin phosphorylation by its kinases, evasion of non-cognate kinase

phosphorylation, and phosphoregulation of its ability to bind and sever actin. These results demonstrate how extraordinarily tight control of multiple functional outputs can converge on a trivially short sequence of residues.

This system allowed us to simultaneously determine the fitness of all possible combinations of N-terminal residues. Previous mutagenesis library screens have similarly used yeast growth as a readout for protein function, binding interactions, or kinase activity[25-27]. Other approaches such as phage display or droplet-based screening could theoretically allow high-throughput analysis of actin binding, with a potential advantage of consistent levels of expression and a single readout[28]. However, it would be difficult to conduct large scale in vitro screens examining actin severing or phosphorylation by LIMK. Some potential advantages of our approach are that all interactions occur in a eukaryotic cellular environment, that selection requires actin severing (and not simply binding) activity, and that we are able to simultaneously screen for cofilin function and regulation by LIMK.

Given the high degree of conservation of the cofilin N-terminal sequence, we were surprised to find relative loose constraints on the sequence for maintaining sufficient activity to support growth. The deleterious effect of most substitutions was largely driven by selectivity for aliphatic and β-branched amino acids at the fifth position. This range of residues is reflected in various cofilin orthologs, including those from mammals (Val), *Arabidopsis* (Met)*,* maize (Leu), *Acanthamoeba* (Ile)*,* and starfish (Thr)[11,29,30]. In all three-dimensional structures of cofilin solved to date, the sidechain of this residue contacts the globular domain of cofilin, suggesting that it may stabilize the protein (**Fig. S12**). Indeed, cofilin^{V5E} aggregated readily during expression and purification. The suitability of variants

at other positions was not readily predicted from known structures. For example, despite allowing interaction with actin, the presence of bulky residues in place of Gly4 is incompatible with the actin-binding mode observed by cryo-EM and would thus require an alternative conformation (**Fig. S13)**. By contrast, a more conservative Ala substitution was deleterious. Substitutions to Ala2 were most tolerated overall and had little effect on cofilin severing activity. This residue is missing in *Acanthamoeba* actophorin and is thus dispensable for actin binding[31]. We conjecture that the conservation of small residues at this position may be important for maintaining N-terminal processing that would be dispensable in the context of our screen. We note that in our viability-based screen, variants promoting the fastest growth would not necessarily be those that bind actin with the highest affinity, and that a "hyperactive" cofilin could be deleterious for growth. However, when examined individually, mutations that imparted fitness defects consistently reduced the actin binding, and there was a general correlation between rank in the screen and affinity. Finally, we note that many variants supporting growth did appear to have fitness defects evident during competitive growth, but a large number were indistinguishable from the WT sequence.

Our finding that almost all cofilin variants with a Thr3 phosphoacceptor residue were resistant to LIMK1 expression and our subsequent biochemical analysis agree with a recent report that cofilin^{S3T} could not be phosphorylated by LIMK[17]. The few Thr3 cofilin sequences selectively depleted by LIMK1 are strongly enriched for Tyr residues at the adjacent position (~40% of sequences, **Fig. S14**). As LIMK1 can efficiently phosphorylate cofilin^{S3Y} it would seem likely that these variants are phosphorylated on Tyr4 rather than Thr3[17]. Ser/Thr kinase phosphoacceptor specificity is canonically determined by the

identity of the residue immediately downstream of the DFG motif in the kinase activation loop (DFG+1), which appears to promote a conformation that facilitates phosphate transfer.[32] However, due to the non-canonical nature of the interaction, cofilin Ser3 is too distant from the LIMK DFG+1 residue for it to impart substrate specificity.[16] Furthermore, the relative openness of the LIMK active site and the flexibility of the cofilin N-terminus suggest that steric clashes with the Thr3 sidechain are unlikely to preclude phosphorylation. Alternatively, LIMK and cofilin[S3T] may form a non-productive complex that sequesters the phosphoacceptor from the catalytic center. The majority of Ser/Thr kinases across species prefer Ser over Thr, potentially explaining why Ser3 is conserved even in organisms lacking LIMK.

In addition to optimizing a substrate for its cognate kinase, a phosphorylation site sequence may also be tuned to avoid the large number of other kinases present within the cell. As most kinases require extended interactions on either side of the phosphosite, the location of the cofilin Ser3 so close to the N-terminus may facilitate evasion of non-cognate kinases. By contrast, LIMK-cofilin binding does not require these canonical interactions as demonstrated by its ability to phosphorylate the N-terminal Ser1 of actophorin[30]. Notably, the extended N-terminal sequences of yeast and plant cofilin orthologs may facilitate a more canonical mode of interaction in organisms lacking LIMKs. By a similar token, because it lacks selective pressure to evade LIMKs, yeast twinfilin is an efficient LIMK substrate, which we could attribute to its Ser phosphoacceptor. Because LIMK-induced growth suppression appears entirely reversed in the context of a cofilin S3A mutation, twinfilin phosphorylation does not appear to confer a growth defect, but its potential phosphoregulation under other conditions warrants further investigation. Though

conservation of Thr at the N-terminus of mammalian twinfilin does not appear necessary to evade phosphorylation by LIMK, it may contribute to its insulation from other pathways.

Save for exclusion of Pro residues, LIMK1 appeared almost entirely promiscuous for the surrounding sequence the cofilin phosphorylation site, consistent with a prior report examining a small panel of mutants[33]. Indeed, LIMK-resistant sequences having bulky hydrophobic residues in place of Gly4 were phosphorylated faster by LIMK1. Instead, our observation that G4F substitution can "correct" the phosphomimetic S3D mutation in vitro and in cells suggest that Gly4 is critical for cofilin to be regulated by phosphorylation. How might this phosphoresistance occur? Molecular dynamic simulations indicate that the phospho-mimetic S3D substitution obstructs docking of the cofilin N-terminus to actin[12]. If the hydrophobic binding site for the N-terminal tail repels a charged phospho-Ser3, it is possible the sidechain of Leu or Phe recapitulates the N-terminal binding interaction. A distinct conformation would be required to accommodate the larger residues but could restore hydrophobic interactions despite Ser3 phosphorylation. Decreased severing may also be attributable to increased dynamics of the N-terminal region as seen in molecular dynamics simulations of the S3D mutant, and this phenomenon may require a flexible Gly4 residue. An alternative model has suggested that phospho-Ser3 participates in intramolecular interactions with one or more basic patches within the cofilin globular domain to sequester the N-terminus away from actin[34]. In this scenario, reducing flexibility of the N-terminal sequence by substituting Gly4 might preclude these intramolecular interactions, leaving the N-terminus free to interact with actin through the canonical binding mode. Further research will be required to distinguish between these models.

Phosphosite sequence conservation within disordered regions is generally thought to be driven by maintaining for signal fidelity in promoting recognition by kinases.[35,36] However, such sequences must also support core function, evade phosphorylation by other kinases, and maintain the capacity for phospho-regulation. In the specific case explored here, each aspect of cofilin functionality is mediated by distinct sequence features of its N-terminal sequence, which collectively restrain sequence variation across the entire tail (**Fig. 7D**). Overall, our studies provide a clearer understanding of specific determinants of cofilin regulation and how it interacts with and severs actin filaments.

Materials and Methods

Plasmids

The high-copy constitutive yeast expression vector (pRS423-GPD) for His_6-tagged human cofilin-1, the bacterial expression vector for N-terminally His_6-SUMO-tagged human cofilin-1 (in pEBDuet28a), and the galactose-inducible yeast expression vectors (pRS415-GAL1) for N-terminally FLAG-tagged LIMK1^{CAT} and full length LIMK1 were previously described.[16] The bacterial expression vector for N-terminally His_6-tagged human cofilin-1 was generated by Gibson assembly using pEBDuet28a-His_6-SUMO-cofilin-1 as a template with primers flanking the SUMO coding sequence. The high-copy yeast expression vector for yeast Cof1 was generated by shuttling the genomic DNA fragment from pRS316-*COF1* (produced by Mark Hochstrasser's laboratory)[37] into pRS423. Bacterial expression vectors for GST-tagged mouse twinfilin-1 and yeast twinfilin were a gift from the Bruce Goode laboratory.[38,39] Point mutations in all plasmids

were made by either QuikChange site directed mutagenesis or by Gibson assembly following amplification with mutagenic primers.

To generate the cofilin-1 N-terminal mutagenesis library, an oligonucleotide pool spanning the region was synthesized with phosphoramidate trimer mixtures encoding all 20 amino acids at the sites encoding Ala2, Gly4 and Val5, and an ACC/AGC mixture (encoding a mix of Ser and Thr) in place of Ser3. Oligonucleotides were PCR amplified and inserted into pRS423-GPD-His$_6$-cofilin-1 plasmid using the 5' BamHI restriction site upstream of the start codon and a 3' SalI restriction sites that was introduced by silent mutagenesis within the cofilin-1 coding sequence. Electrocompetent DH10β *E. coli* (Invitrogen ElectroMAX) were transformed with ligation products, providing at least 1000 transformants per library variant. Plasmid library DNA was recovered from the pool of transformed bacterial colonies, PCR amplified and sequenced on an Illumina NovaSeq instrument to confirm full representation of all variants.

Cofilin library screening

The *S. cerevisiae* TeTO$_7$-*COF1* strain[22] (Horizon Discovery) was transformed sequentially with pRS415-GAL1-FLAG-LIMK1 and the cofilin library plasmid pool and frozen in aliquots to conduct replicate screens. For each screen, transformed yeast were pooled and grown at 30 °C with shaking in liquid culture with selective media (SC-His-Leu) containing 2% glucose at a starting OD$_{600}$ of 0.1. After 4 - 5 population doublings, a portion was diluted to an OD$_{600}$ of 0.1 with SC-His-Leu media containing 2% raffinose and 10 mg/L DOX. After the culture reached an OD$_{600}$ of 1 - 1.5, a portion was reserved for

plasmid recovery (T_0 sample), and the remaining culture was split and diluted into SC-His-Leu liquid media containing 10 mg/L DOX and either 2% glucose or 2% raffinose/1% galactose to OD_{600} = 0.1. Cultures were subjected to three growth and dilution cycles in which they were grown to an OD_{600} of 1 - 2 and diluted back to an OD_{600} of 0.1, saving a portion for plasmid recovery at each step (T_1 – T_3 samples). After the screen was complete, plasmids were recovered from cell pellets using a QIAGEN Spin Miniprep kit. The N-terminal variable region of the plasmid pool was PCR amplified to attach sequencing adaptors and add barcodes and sequenced on an Illumina NovaSeq instrument.

The relative representation of each cofilin variant at each timepoint was calculated by dividing the number of reads that variant by the total read count. The enrichment or depletion of a cofilin-1 library sequence in the absence of LIMK1 induction was calculated as the ratio of its relative representation at T_0 and T_2 in glucose. Enrichment or depletion following LIMK1 induction was calculated from timepoints T_1 and T_3 in glucose vs. galactose. Sequences were ranked by the average log fold changes in representation across three replicate screens. Data were processed using Microsoft Excel and sequence logos were generated using pLogo.[40]

Immunoblotting

Yeast lysates were fractionated by either conventional SDS-PAGE or Mn^{2+} Phos-tag SDS-PAGE before transfer to polyvinylidene difluoride (PVDF) membranes. Membranes were blocked with 5% powdered milk in Tris-buffered saline solution with 0.05% Tween-20 (TBST) for 2 hours at room temperature and incubated with primary

antibodies diluted 1:2000 in TBST containing 5% milk overnight at 4 °C. Primary antibodies included: anti-Penta-His (QIAGEN, #34650), anti-FLAG M2 (Sigma-Aldrich, #F3165), and anti-Kss1 (Santa Cruz, #sc-6775-R). Membranes were washed with TBST before incubation with fluorophore-labeled secondary antibodies (1:20,000 dilution) in TBST containing 5% bovine serum albumin (BSA) for 1 hr. at room temperature. Membranes were washed with TBST and imaged with an Odyssey CLx imager (LI-COR Biosciences). Fluorescence signals were quantified using Image Studio software.

Yeast growth assays

To evaluate the ability of cofilin variants to support growth, overnight cultures of TeTO$_7$-*COF1* yeast transformed with the indicated expression vectors were diluted to OD$_{600}$ of 0.1 in selective media and grown in a shaking incubator at 30 °C until 3 - 4 additional doublings had occurred. Portions of each culture were reserved for immunoblotting, and a series of 5-fold dilutions (OD$_{600}$ range from 0.5 to 0.0008) were spotted onto SC-His agar plates with or without 10 mg/L DOX. Plates were incubated at 30 °C for 3 - 4 days before imaging. Lysates for immunoblotting were prepared by pelleting cells at 4696 g for 2 min and vortexing the cell pellet with glass beads in TCA extraction buffer (10 mM Tris, pH 8.0, 10% TCA, 25 mM NH$_4$OAc, 1 mM EDTA). Proteins were pelleted by centrifugation and resuspended in 100 mM Tris (pH 11.0) with 3% SDS before boiling for 5 min. Protein concentrations were determined by BCA assay, equal amounts were separated by SDS-PAGE and subjected to immunoblotting as described above.

The effect of LIMK1 expression on cofilin phosphorylation and yeast growth was evaluated using TeTO$_7$-*COF1* yeast co-transformed expression vectors for the indicated cofilin variants and pRS415-GAL1-FLAG-LIMK1, pRS415-GAL1-FLAG-LIMK1^{CAT} or empty pRS415-GAL1 vector as a control. Yeast were grown overnight in SC-His-Leu containing 2% raffinose to derepress the GAL promoter, diluted to an OD$_{600}$ of 0.1, and allowed to grow through 3 - 4 population doublings at 30 °C. A portion of the cells were spotted and grown as described above on SC-His-Leu agar plates containing 2% glucose, 2% glucose with 10 mg/L DOX, or 2% raffinose/1% galactose with 10 mg/L DOX. The remaining culture was split, and LIMK1 expression was either induced or repressed either galactose to 1% or glucose to 2%. After 12 hr, equal OD$_{600}$ units of each culture were pelleted at 4696 x g for 2 min and lysates prepared by TCA extraction as described above. Lysates were fractionated by both conventional SDS-PAGE and Mn^{2+} Phos-tag SDS-PAGE before immunoblotting. Phos-tag acrylamide gels were prepared with 12% acrylamide, 100 μM Mn^{2+}, and 50 μM Phos-tag Acrylamide AAL-107 provided by Fujifilm.

Protein purification

Human cofilin-1 variants were expressed in *E. coli* Rosetta(DE3) cells overnight at 16 °C after induction with 0.5 mM IPTG. Cells were pelleted, snap-frozen on dry ice/EtOH, thawed on ice, and resuspended in bacterial lysis buffer (20 mM Tris, pH 7.5, 140 mM NaCl, 1 mM DTT, 0.4% Igepal CA-630, 1 mM MgCl$_2$, 200 μg/ml lysozyme, 10 μg/ml pepstatin A, Roche cOmplete protease inhibitor cocktail, and 30 U/ml DNAse 1). Cells were lysed by sonication and pelleted by centrifugation (13,870 x g for 30 min. at 4 °C). The supernatant was extracted and incubated with TALON affinity resin while rotating at

4 °C for 1 - 2 hr. A gravity flow column was used to separate the TALON resin from the remaining cell lysate. The resin was washed once with 0.5% Igepal CA-630 in PBS and once in 10 mM imidazole, 20 mM Tris, pH 8.0, 140 mM NaCl. His$_6$-SUMO-tagged cofilin variants were washed an additional time with cofilin storage buffer (10 mM HEPES, 100 mM NaCl, 1 mM DTT, 10% glycerol) and eluted from the TALON resin by overnight digestion with SUMO protease at 4 °C. N-terminally His$_6$-tagged cofilin variants were eluted by imidazole wash (20 mM Tris, pH 7.5, 250 mM imidazole, 100 mM NaCl, 1 mM DTT) and dialyzed into cofilin storage buffer overnight at 4 °C. Cofilin variants used for actin binding and severing assays were further purified by size exclusion chromatography on a Superdex200 10/300 GL column in 25 mM Tris, pH 8.0, 0.5 mM DTT, 0.25 mM EGTA. Cofilin eluted as a single peak across multiple fractions, which were pooled and concentrated using Amicon Ultra centrifugal filter concentrators (3,000 Da cutoff). Protein concentration was determined by A$_{280}$ absorbance using a Thermo Scientific 2000 nanodrop instrument.

GST-tagged yeast and mouse twinfilin variants were expressed in *E. coli* and cells were lysed as described above. Clarified cell lysate was incubated with glutathione Sepharose resin for 2 hr at 4 °C. The resin was washed once with 0.5% Igepal CA-630 in PBS and once with cofilin storage buffer. Proteins were eluted by overnight digestion with GST-tagged HRV 3C protease at 4 °C. Twinfilin concentration was determined by comparison to BSA standards following SDS-PAGE and Coomassie-staining.

FLAG-tagged human LIMK1 was purified using a yeast expression system. Budding yeast strain TH_5610 transformed with pRS423-GPD-His$_6$-cofilin-1^{S3A} (to support growth of yeast during LIMK induction) and pRS415-GAL1-FLAG-LIMK1 were

grown in selective liquid media (SC-His-Leu) containing 2% raffinose at 30 °C with shaking to an OD_{600} of 2. LIMK1 expression was induced by the addition of galactose and nutrient-rich yeast extract peptone (YP) broth, bringing the final concentrations of media to 1xYP media and 1% galactose. Cells were induced for 12 hr. at 30 °C before cells were harvested, pelleted, washed once with sterile water, and snap-frozen. The cell pellet was thawed on ice and resuspended in yeast lysis buffer (50 mM HEPES, pH 7.4, 150 mM NaCl, 1 mM EDTA, 0.5% Triton X-100, 10% glycerol, 0.5 mM DTT, 10 µg/ml leupeptin, 2 µg/ml pepstatin A, 2.5 mM $NaPP_i$, 10 µg/ml aprotinin, 1 mM βGP and 1 mM Na_3VO_4). Cells were lysed by vortexing with glass beads at 4 °C. Cell lysates were pelleted, and the recovered supernatant was incubated with M2 anti-FLAG affinity resin in batch for 1 - 2 hr at 4 °C. The resin was washed twice with yeast lysis buffer and twice with FLAG wash buffer (50 mM HEPES, pH 7.4, 100 mM NaCl, 1 mM DTT, 0.01% Igepal CA-630, 10% glycerol, 1 mM βGP, and 100 µM Na_3VO_4) before the protein was eluted with 0.5 mg/mL 3x FLAG peptide (Sigma). LIMK1 concentration was determined by running alongside BSA standards on SDS-PAGE with Coomassie-staining.

To generate Ser3-phosphorylated cofilin, FLAG-LIMK1 catalytic domain was expressed in yeast and purified as previously described but not eluted from the resin. Cofilin^{WT} purified as above was incubated with resin-bound LIMK1 catalytic domain overnight at 4 °C in 50 mM Tris, pH 7.5, 100 mM NaCl, 5 mM $MgCl_2$, 5 mM $MnCl_2$, 1 mM DTT, and 1 mM ATP. The resin was then pelleted and the supernatant dialyzed overnight at 4 °C into 50 mM Tris, pH 6.8 with 1 mM DTT. Phosphorylated and unphosphorylated cofilin species were separated by cation-exchange chromatography using a MonoS 5/50 GL column. Cofilin phosphorylation was assessed by Mn^{2+} Phos-tag SDS-PAGE.

Actin filament binding assays

The binding affinity of cofilin variants for actin filaments was compared using *in vitro* pyrene quenching assays as previously reported[12,24,41]. Serial dilutions of purified cofilin variants were incubated with polymerized pyrene-labeled actin (1 µM, ~83% labeled) in 1X KMI buffer (20 mM imidazole, pH 7.0, 50 mM KCl, 2mM $MgCl_2$, 2 mM DTT, 200 µM ATP, 1 mM NaN_3) for 1 hr at 25 °C. Pyrene quenching was measured in a plate reader (excitation 366 nm, emission scan 409 nm). Data were normalized to the signal for unquenched pyrene-actin alone and fit to a model-independent sigmoidal 4 parameter dose-response curve using GraphPad Prism.

Actin severing assays

The severing activity of cofilin variants was assessed using TIRF microscopy as previously described[12]. Alexa-647 labeled actin filaments (5% labeled) were polymerized for 1 hour in 1X KMI buffer. Serial dilutions of purified cofilin variants were incubated with the polymerized actin for 1 hr before placement on poly-L-lysine coated glass slips and imaged directly by TIRF microscopy. The recorded images were skeletonized using ImageJ FIJI and analyzed with the Persistence Software ver. 4.2.3 to quantify the actin filament lengths.[42] Persistence Software is available to download for free at delacruzlab.yale.edu/persistence-software.

Radiolabel kinase assays

Purified cofilin or twinfilin (2 µM) was incubated with LIMK1 catalytic domain (2 nM) and ATP (12.5 µM with 0.1 µCi/µL of [γ-^{32}P] ATP) in reaction buffer (50 mM Tris, pH 7.5,

100 mM NaCl, 5 mM MgCl$_2$, 5 mM MnCl$_2$, and 1 mM DTT) at 30 °C. At 5 min and 10 min incubation aliquots were removed and quenched with 4x SDS-PAGE loading buffer. Samples were boiled 5 min and fractionated by 15% acrylamide SDS-PAGE. Gels were Coomassie-stained and dried before exposure to a phosphor screen. Phosphor imaging and quantification of radiolabel incorporation was conducted using a BioRad Molecular Imager FX Pro Plus. Phosphorylation rates were calculated as the change in phosphorylation across three timepoints (0, 5, and 10 min.) and normalized to the phosphorylation rate of wild-type cofilin-1. Radiolabel kinase assays comparing the phosphorylation of twinfilin variants by LIMK1 were conducted using the same parameters. Reaction kinetics for cofilin^{WT} and cofilin^{G4F} were conducted with the same reaction conditions as described previously; differing only in cofilin concentration. A set of seven serial dilutions of both cofilin variants was prepared using the cofilin storage buffer described previously. In the final reactions, cofilin concentrations ranged from 50 to 0.5 µM. A titration range of [γ-^{32}P] ATP (10 nCi, 5 nCi, and 2.5 nCi) was pipetted on filter paper and exposed to a phosphor screen concurrently with the dried Coomassie-stained gel. Quantification of radiolabel signal was performed using a BioRad Molecular Imager FX Pro Plus and absolute values were obtained by comparison to the standard curve of [γ-^{32}P] ATP. Reaction kinetics were calculated using Microsoft Excel and GraphPad Prism.

Acknowledgments

We thank Cameron Schmitz for technical assistance in generating cofilin mutants. This research was supported by National Institute of General Medical Sciences of the National Institutes of Health under awards R01 GM102262 (awarded to BET and TJB) and R35 GM136656 (awarded to EMDLC).

Table 1. Properties of cofilin variants chosen for individual evaluation. For growth on solid media: +++, equal growth in the presence or absence of DOX; ++, ~5-fold lower growth on DOX; +, some detectable growth on DOX; -, no growth on DOX. For actin binding: *, none detectable; **, not assayed due to protein aggregation. ND, not determined. Rank and Log_2 fold change values for all variants are provided in Dataset S2.

Cofilin Variant	Growth on plate	Actin binding $K_{0.5}$, µM	$K_{0.5}$ 95% confidence interval µM	Screen rank	Log_2 fold change
WT	+++	0.61	0.54 – 0.67	87	3.19
A2L	+++	0.53	0.47 – 0.57	182	3.05
A2D	+++	1.50	1.39 – 1.60	822	2.55
S3D	-	4.65	4.41 - 4.49	ND	ND
S3D,G4F	++	1.06	1.02 – 1.09		
G4A	++	3.15	3.03 – 3.26	4424	-0.45
G4L	+++	ND	ND	1260	2.33
G4P	+	ND	ND	5800	-2.41
G4F	++	1.00	0.88 – 1.09	2292	1.76
G4E	-	*	*	9618	-4.45
G4K	-	3.48	3.30 – 3.66	8153	-3.98
V5E	-	**	**	10647	-4.70
V5M	+++	0.48	0.05 - 0.11	262	2.95
A2L,V5M	+++	0.85	< 0.93	197	3.02

Figure 1. A dual-functionality screen of a cofilin-1 N-terminal mutagenesis library.
(A) Growth on solid media of TeTO$_7$-*COF1* yeast exogenously expressing the indicated cofilin in the presence or absence of doxycycline. Image representative of n = 2. (B) Yeast harboring the indicated expression plasmids were grown on either glucose or galactose to induce LIMK1 expression in the presence or absence of doxycycline. Image representative of n = 7. (C) Design of cofilin N-terminal combinatorial mutagenesis library. X indicates any of the 20 natural amino acids. (D) Schematic of the cofilin library dual-functionality screen. (E) The graphs show change in relative representation of the indicated cofilin variants over time during culture in either glucose (no LIMK1) or galactose (LIMK1 induction). Individual datapoints from three replicate screens performed independently are shown. Blue circles indicate timepoints used to identify functional sequences. Orange circles indicate timepoints used to identify sequences inhibited by LIMK1. Time zero is after de-repression of P_{GAL}-LIMK1 over the course of 1 - 2 population doublings in raffinose + DOX media.

Figure 2. Cofilin library screen results for yeast growth rescue. (A) Waterfall plot depicting enrichment or depletion of all 16,000 cofilin library sequences expressed in TeTO7-*COF1* yeast grown in doxycycline. Sequences are ordered left to right from highest to lowest enrichment. Data show the average from three independent screens. (B) Probability logo of cofilin N-terminal library sequences enriched by screen. Red lines

indicate the threshold for statistical significance (p = 0.05). (C) Heat map for cofilin N-terminal library sequences enriched by screen. Color scale indicates the mean enrichment score of the 800 (positions 2, 4 and 5) or 8000 (position 3) library sequences containing the indicated residue across 3 independent replicates. (D) Distribution of cofilin sequences by enrichment scores according to identity of residues at the indicated positions. Boxes indicate 2nd and 3rd quartile of distribution with median line. Bars indicate sequences in the 10 – 90 percentile range.

Figure 3. Impact of cofilin N-terminal mutations on yeast growth and actin binding.
(A,B) Growth assay comparing cofilin mutants containing substitutions of residues Ala2, Gly4, or Val5. Images representative of n = 3. (C-E) Pyrene quenching assays comparing cofilin^{WT} to the indicated cofilin mutants. Actin concentration pre-polymerization was 1 µM. Replicates for binding assays are as follows: cofilin^{WT} (n = 4), cofilin^{A2L} (n = 3), cofilin^{A2D} (n = 4), cofilin^{S3D} (n = 3), phospho-cofilin^{WT} (n = 2), cofilin^{G4A} (n = 3), cofilin^{G4F} (n = 2), cofilin^{G4E} (n = 2), cofilin^{G4K} (n = 3), cofilin^{V5M} (n = 3), and cofilin^{A2L,V5M} (n = 2).

Figure 4. Cofilin library screen results for yeast growth rescue with and without LIMK1. (A) Scatter plot depicting change in representation of Ser3 cofilin library sequences with and without LIMK1 expression. Data points show average enrichment in glucose and galactose across three replicates. (B) Logo of sequences supporting growth in yeast but are selectively inhibited by LIMK1 (lower right quadrant in panel A) on the background of all sequences that support yeast growth in the absence of LIMK1 (upper and lower right quadrants in panel A). (C) Distribution of cofilin sequences by enrichment scores with and without LIMK1 expression according to identity of the phosphoacceptor residue. Boxes indicate 2nd and 3rd quartiles, and bars indicate sequences with 10- and 90-percentiles of distribution with median line.

Organism	Protein	N-terminal Sequence	LIMK
H. sapiens, *M. musculus*	mCofilin-1	MA**S**GVAV	Yes
	mTwinfilin-1	MSHQ**T**GIQA	
S. cerevisiae	yCofilin	MSR**S**GVAV	No
	yTwinfilin	MSTQ**S**GIVA	

Figure 5. Effects of cofilin Ser3 substitutions on phosphorylation by LIMK1. (A) TeTO$_7$-*COF1* yeast expressing the indicated human cofilin-1 mutants were grown in the presence of doxycycline to repress endogenous Cof1 expression and/or galactose to induce LIMK1 expression. Images are representative of n = 3 independent experiments. (B) Immunoblots with the indicated antibodies of lysates corresponding to yeast plated in panel A. Samples in the cofilin blot were separated on Phos-tag SDS-PAGE to resolve phosphorylated (upper band) from unphosphorylated protein (lower band). Kss1 serves as a loading control. (C) Quantification of immunoblots measuring the fraction of total cofilin phosphorylated by LIMK1. Error bars show SD for n = 3 independent experiments. (D) *In vitro* radiolabel kinase assay comparing phosphorylation of purified cofilin (2 μM) by LIMK1 (2 nM) catalytic domain for 10 min at 30 °C. Catalytically dead LIMK1^{D460N} kinase domain serves as a negative control. (E) Quantification of phosphorylation rates of cofilin mutants normalized to cofilin^{WT}. Error bars show SD for n = 4 independent experiments. (F) Table comparing N-terminal sequences of mammalian and yeast isoforms of cofilin and twinfilin. (G) *In vitro* radiolabel kinase assay comparing phosphorylation of purified mouse and budding yeast isoforms of twinfilin (2 μM) by LIMK1 kinase domain (2 nM) for 10 min at 30 °C. LIMK1^{D460N} kinase domain serves as a negative control. (H) Quantification of phosphorylation rates normalized to cofilin^{WT}. Error bars show SD for n = 3 independent experiments.

Figure 6. Effects of cofilin Glycine 4 substitutions on LIMK1 substrate suitability.
(A) Yeast growth assay with the TeTO₇-*COF1* strain expressing the indicated cofilin variants with or without LIMK1 expression. Representative of n = 3 independent experiments. (B) Immunoblots with the indicated antibodies of lysates corresponding to yeast plated in panel A separated by standard or Phos-tag PAGE. (C) Quantification of immunoblots measuring the fraction of total cofilin phosphorylated by LIMK1. Error bars show SD for n = 3 independent experiments. (D) *In vitro* radiolabel kinase assay comparing phosphorylation of purified 2 µM cofilin by 2 nM LIMK1 kinase domain for 10 min at 30 °C. Catalytically inactive LIMK1^{D460N} kinase domain serves as a negative control. (E) Quantification of phosphorylation rates of cofilin mutants normalized to cofilin^{WT}. Error bars show SD for n = 6 independent experiments. (F) *In vitro* radiolabel

kinase assay comparing phosphorylation of cofilin^{WT} and cofilin^{G4F} by 2 nM LIMK1 kinase domain across a range of concentrations. Reactions performed for 10 min at 30 °C. (G) Quantification of phosphorylation rates of cofilin mutants. Error bars show SD for n = 3 independent experiments. Kinetic parameters include SE for k_{cat} and K_m.

Figure 7. Cofilin G4F mutation counters the inhibitory effects of phosphomimetic S3D substitution. (A) Yeast growth assay with TeTO₇-*COF1* strain expressing the indicated cofilin variants in the presence or absence of doxycycline. (B) Immunoblots of cofilin protein levels corresponding to yeast plated in panel A. Kss1 serves as a loading control. Images are representative of n = 3 independent experiments. (C) Pyrene quenching assays comparing cofilin[WT] to cofilin variants with a phosphomimetic substitution or phosphorylation modification at residue Ser3. Actin concentration pre-polymerization was 1 µM. WT and G4F single mutant data are the same as shown in Fig 3 and are included here for comparison. Binding parameters are listed in Table 1. Replicates for binding assays are as follows: cofilin[WT] (n = 4), cofilin[S3D] (n = 3), phospho-cofilin[WT] (n = 2), cofilin[G4F] (n = 2), and cofilin[S3D,G4F] (n = 3). (D) Overlapping constraints promoting conservation of the cofilin phosphosite motif sequence.

P_{GAL4}-LIMK1^{CAT}	Cofilin	- DOX Glucose	+ DOX Glucose	+DOX Galactose
+	WT			
+	S3A			
-	WT			
-	-			

Figure S1. Growth inhibition of yeast by galactose-induced LIMK1^{CAT} expression. Yeast harboring the indicated expression plasmids were grown on either glucose or galactose to induce LIMK1^{CAT} expression in the presence or absence of doxycycline. Image representative of n = 7 independent experiments.

Figure S2. N-terminal His₆-tagged cofilin can sever actin and be phosphorylated by LIMK1. (A) TIRF microscopy Alexa647-labeled actin filaments by N-terminally His₆-tagged cofilin-1. Cofilin concentrations listed in upper right corner of each image. (B) Quantification of actin filament length at concentrations for His₆-tagged cofilin-1. (C) *In vitro* radiolabel kinase assay comparing phosphorylation of purified 2 µM tagged and un-tagged cofilin^{WT} and cofilin^{S3T} by 2 nM LIMK1 kinase domain for 10 min at 30 °C. Kinase inactive mutant (D460N) was assayed alongside. n = 2. (D) Quantification of phosphorylation rates normalized to untagged cofilin^{WT}.

Figure S3. Cofilin N-terminus combinatorial motifs identified by library screen. (A) Consensus sequence logo of the top 400 sequences enriched by the screen shows an enrichment for Gly and Leu residues at position 4. Consensus logos of all sequences within the top 400 that contain a Gly4 (B), Leu4 (C), Val5 (D), and Met5 (E). Red lines indicate a threshold for statistical significance (P=0.05). Letters large enough to extend past this line represent significant representation. (F) Sequence distribution of cofilin sequences containing an L-V, L-M, G-V, or G-M combination of residues. Box*es* indicate 1st and 3rd quartile of distribution. Bars indicate sequences with minimum and maximum enrichment. Bars indicate sequences with 10- and 90-percentiles of distribution.

Figure S4. Cofilin N-terminus X-T-X-T motif identified by library screen. Consensus logos of all sequences within the top 400 that contain a Thr5 (A) or Thr3 (B) residue. Red lines indicate a threshold for statistical significance (P=0.05). Letters large enough to extend past this line represent significant representation. (C) Sequence distribution of cofilin sequences containing an X-T-X-T motif or an X-S/T-G-V motif. Box*es* indicate 1[st] and 3[rd] quartile of distribution. Bars indicate sequences with 10- and 90-percentiles of distribution.

Figure S5. Cofilin variant protein levels within yeast. Immunoblots of cofilin protein levels in lysates corresponding to yeast plated in Fig 3A and C. Kss1 serves as a loading control. (B, D) Quantification of protein levels normalized to the Kss1 signal as a fraction of the normalized cofilin[WT] signal. Error bars show SD for n = 3 independent experiments.

Figure S6. *In vitro* **actin severing assays comparing cofilin mutants.** TIRF microscopy images of Alexa647-labeled actin filaments incubated with the indicated concentration of WT or mutant cofilin.

Figure S7. Quantification of i*n vitro* actin severing assays comparing cofilin mutants. Changes in average Alexa647-labeled actin filament length (μm) measured against changes in cofilin concentration (μM) quantified from TIRF microscopy images.

Figure S8. Distributions of Ser3 and Thr3 cofilin library sequences. Two-dimensional plot of Ser3 (A) and Thr3 (B) cofilin library sequences ±LIMK1. Cofilin sequences containing serine as the phosphoacceptor residue are selectively depleted by LIMK1 induced-expression. Each library sequence is plotted according to its average enrichment in glucose and galactose across three replicates.

Figure S9. Exacerbated effects LIMK1 catalytic domain on growth inhibition of cofilin Serine 3 substitutions. (A) Plate-based yeast growth assay demonstrates that cofilin[S3T] growth rescue is unaffected by LIMK1 expression. Representative of n = 3 independent experiments. (B) Immunoblots of lysates corresponding to yeast plated on A. Cofilin species are separated based on phosphorylation state by Phos-tag SDS-PAGE. Kss1 serves as a loading control. (C) Quantification of immunoblots measuring the fraction of total cofilin phosphorylated by LIMK1. Error bars show SD for n = 3 independent experiments. (D) Phosphorylation of the indicated forms of cofilin by LIMK2 catalytic domain.

Figure S10. Suitability of twinfilin as a LIMK1 substrate. (A) Diagram comparing ADF-homology domains of human cofilin-1 and twinfilin-1. (B) Conserved cofilin residues (orange) (PDBID:5YU8)[43] important for LIMK1 binding are also conserved in the twinfilin-1 N-terminal ADF-H domain (C) (PDBID:7CCC)[44].

Figure S11. Exacerbated effects of LIMK1 catalytic domain on growth inhibition in the context of Glycine 4 cofilin mutations. (A) Plate-based yeast growth assay demonstrates that cofilin^{G4L} and cofilin^{G4F} growth rescue is unaffected by LIMK1 expression. Representative of n = 3 independent experiments. (B) Immunoblots of lysates corresponding to yeast plated on A. Cofilin species are separated based on phosphorylation state by Phos-tag SDS-PAGE, with the slower migrating species corresponding to phosphorylated cofilin. (C) Quantification of immunoblots measuring the fraction of total cofilin phosphorylated by LIMK1. Error bars show SD for n = 3 independent experiments.

Figure S12. Comparison of N-terminal structures of cofilin orthologs. (A) Structure of human cofilin-1. PDB ID: 5HVK.[16] (B) Structure of *Arabidopsis* ADF. PDB ID: 1F7S.[29] (C) Structure of *Acanthamoeba* actophorin. PDB ID: 1AHQ.[45]

Figure S13. Potential structural rationale for insensitivity of cofilin[G4F] to phosphorylation. (A) Structure showing conformation of the cofilin[WT] N-terminus (cartoon and stick representation, cyan) when bound to actin (surface representation, beige). Acetylated Ala2 was absent from the reported structure and has been modeled here using the building feature in PyMol. PDB ID: 5YU8 [43] (B) Model of cofilin[G4F] based on the previous structure produced by the mutagenesis and sculpting functions in PyMol suggests showing a potential alternative conformation that might reduce the inhibitory effect of Ser3 phosphorylation.

A

B

Figure S14. Distributions of X-T-Y-X cofilin library sequences. (A) Two-dimensional plot of cofilin library sequences containing a X-T-Y-X motif (blue) compared to all sequences contain a Thr residue at position 3 (orange) ±LIMK1. The majority of cofilin

sequences containing a threonine as the phosphoacceptor residue and a tyrosine residue at position 4 are selectively depleted by LIMK1 induced-expression. Each library sequence is plotted according to its average enrichment in glucose and galactose across three replicates. (B) Consensus sequence logo of cofilin sequences containing a Thr3 phosphoacceptor residue selectively depleted by LIMK1 induction when compared to a background of all Thr3 cofilin sequences that support yeast growth.

References

1. Pollard, T. D. Actin and Actin-Binding Proteins. *Cold Spring Harb Perspect Biol* **8**, (2016)
2. Mizuno, K. Signaling mechanisms and functional roles of cofilin phosphorylation and dephosphorylation. *Cell Signal* **25**, 457-469 (2013)
3. Maciver, S. K., and Hussey, P. J. The ADF/cofilin family: actin-remodeling proteins. *Genome Biol* **3**, reviews3007 (2002)
4. Kanellos, G., and Frame, M. C. Cellular functions of the ADF/cofilin family at a glance. *J Cell Sci* **129**, 3211-3218 (2016)
5. Maciver, S. K., and Weeds, A. G. Actophorin preferentially binds monomeric ADP-actin over ATP-bound actin: consequences for cell locomotion. *FEBS Lett* **347**, 251-256 (1994)
6. De La Cruz, E. M. How cofilin severs an actin filament. *Biophys Rev* **1**, 51-59 (2009)
7. Suarez, C., Roland, J., Boujemaa-Paterski, R., Kang, H., McCullough, B. R., Reymann, A. C., Guerin, C., Martiel, J. L., De la Cruz, E. M., and Blanchoin, L. Cofilin tunes the nucleotide state of actin filaments and severs at bare and decorated segment boundaries. *Curr Biol* **21**, 862-868 (2011)
8. Bibeau, J. P., Gray, S., and De La Cruz, E. M. Clusters of a Few Bound Cofilins Sever Actin Filaments. *J Mol Biol* **433**, 166833 (2021)
9. Oosterheert, W., Klink, B. U., Belyy, A., Pospich, S., and Raunser, S. Structural basis of actin filament assembly and aging. *Nature* **611**, 374-379 (2022)
10. Elam, W. A., Kang, H., and De la Cruz, E. M. Biophysics of actin filament severing by cofilin. *FEBS Lett* **587**, 1215-1219 (2013)
11. Moriyama, K., Iida, K., and Yahara, I. Phosphorylation of Ser-3 of cofilin regulates its essential function on actin. *Genes Cells* **1**, 73-86 (1996)
12. Elam, W. A., Cao, W., Kang, H., Huehn, A., Hocky, G. M., Prochniewicz, E., Schramm, A. C., Negron, K., Garcia, J., Bonello, T. T., Gunning, P. W., Thomas, D. D., Voth, G. A., Sindelar, C. V., and De La Cruz, E. M. Phosphomimetic S3D cofilin binds but only weakly severs actin filaments. *J Biol Chem* **292**, 19565-19579 (2017)
13. Lappalainen, P., Fedorov, E. V., Fedorov, A. A., Almo, S. C., and Drubin, D. G. Essential functions and actin-binding surfaces of yeast cofilin revealed by systematic mutagenesis. *EMBO J* **16**, 5520-5530 (1997)
14. Nagaoka, R., Abe, H., and Obinata, T. Site-directed mutagenesis of the phosphorylation site of cofilin: its role in cofilin-actin interaction and cytoplasmic localization. *Cell Motil Cytoskeleton* **35**, 200-209 (1996)
15. Huehn, A. R., Bibeau, J. P., Schramm, A. C., Cao, W., De La Cruz, E. M., and Sindelar, C. V. Structures of cofilin-induced structural changes reveal local and asymmetric perturbations of actin filaments. *Proc Natl Acad Sci U S A* **117**, 1478-1484 (2020)
16. Hamill, S., Lou, H. J., Turk, B. E., and Boggon, T. J. Structural Basis for Noncanonical Substrate Recognition of Cofilin/ADF Proteins by LIM Kinases. *Mol. Cell* **62**, 397-408 (2016)
17. Salah, E., Chatterjee, D., Beltrami, A., Tumber, A., Preuss, F., Canning, P., Chaikuad, A., Knaus, P., Knapp, S., Bullock, A. N., and Mathea, S. Lessons from

LIMK1 enzymology and their impact on inhibitor design. *Biochem J* **476**, 3197-3209 (2019)

18. Mezna, M., Wong, A. C., Ainger, M., Scott, R. W., Hammonds, T., and Olson, M. F. Development of a high-throughput screening method for LIM kinase 1 using a luciferase-based assay of ATP consumption. *J Biomol Screen* **17**, 460-468 (2012)

19. Lappalainen, P., and Drubin, D. G. Cofilin promotes rapid actin filament turnover in vivo. *Nature* **388**, 78-82 (1997)

20. Nakano, K., Satoh, K., Morimatsu, A., Ohnuma, M., and Mabuchi, I. Interactions among a fimbrin, a capping protein, and an actin-depolymerizing factor in organization of the fission yeast actin cytoskeleton. *Mol Biol Cell* **12**, 3515-3526 (2001)

21. Iida, K., Moriyama, K., Matsumoto, S., Kawasaki, H., Nishida, E., and Yahara, I. Isolation of a yeast essential gene, COF1, that encodes a homologue of mammalian cofilin, a low-M(r) actin-binding and depolymerizing protein. *Gene* **124**, 115-120 (1993)

22. Mnaimneh, S., Davierwala, A. P., Haynes, J., Moffat, J., Peng, W. T., Zhang, W., Yang, X., Pootoolal, J., Chua, G., Lopez, A., Trochesset, M., Morse, D., Krogan, N. J., Hiley, S. L., Li, Z., Morris, Q., Grigull, J., Mitsakakis, N., Roberts, C. J., Greenblatt, J. F., Boone, C., Kaiser, C. A., Andrews, B. J., and Hughes, T. R. Exploration of essential gene functions via titratable promoter alleles. *Cell* **118**, 31-44 (2004)

23. Moriyama, K., and Yahara, I. The actin-severing activity of cofilin is exerted by the interplay of three distinct sites on cofilin and essential for cell viability. *Biochem J* **365**, 147-155 (2002)

24. De La Cruz, E. M. Cofilin binding to muscle and non-muscle actin filaments: isoform-dependent cooperative interactions. *J Mol Biol* **346**, 557-564 (2005)

25. Ahler, E., Register, A. C., Chakraborty, S., Fang, L., Dieter, E. M., Sitko, K. A., Vidadala, R. S. R., Trevillian, B. M., Golkowski, M., Gelman, H., Stephany, J. J., Rubin, A. F., Merritt, E. A., Fowler, D. M., and Maly, D. J. A Combined Approach Reveals a Regulatory Mechanism Coupling Src's Kinase Activity, Localization, and Phosphotransferase-Independent Functions. *Mol Cell* **74**, 393-408 e320 (2019)

26. Shi, G., Song, C., Torres Robles, J., Salichos, L., Lou, H. J., Lam, T. T., Gerstein, M., and Turk, B. E. Proteome-wide screening for mitogen-activated protein kinase docking motifs and interactors. *Sci. Signal.* **16**, eabm5518 (2023)

27. Hietpas, R., Roscoe, B., Jiang, L., and Bolon, D. N. Fitness analyses of all possible point mutations for regions of genes in yeast. *Nat Protoc* **7**, 1382-1396 (2012)

28. Rossenu, S., Leyman, S., Dewitte, D., Peelaers, D., Jonckheere, V., Van Troys, M., Vandekerckhove, J., and Ampe, C. A phage display-based method for determination of relative affinities of mutants. Application of the actin-binding motifs in thymosin beta 4 and the villin headpiece. *J Biol Chem* **278**, 16642-16650 (2003)

29. Bowman, G. D., Nodelman, I. M., Hong, Y., Chua, N. H., Lindberg, U., and Schutt, C. E. A comparative structural analysis of the ADF/cofilin family. *Proteins* **41**, 374-384 (2000)

30. Blanchoin, L., Robinson, R. C., Choe, S., and Pollard, T. D. Phosphorylation of Acanthamoeba actophorin (ADF/cofilin) blocks interaction with actin without a change in atomic structure. *J Mol Biol* **295**, 203-211 (2000)

31. Quirk, S., Maciver, S. K., Ampe, C., Doberstein, S. K., Kaiser, D. A., VanDamme, J., Vandekerckhove, J. S., and Pollard, T. D. Primary structure of and studies on Acanthamoeba actophorin. *Biochemistry* **32**, 8525-8533 (1993)

32. Chen, C., Ha, B. H., Thevenin, A. F., Lou, H. J., Zhang, R., Yip, K. Y., Peterson, J. R., Gerstein, M., Kim, P. M., Filippakopoulos, P., Knapp, S., Boggon, T. J., and Turk, B. E. Identification of a major determinant for serine-threonine kinase phosphoacceptor specificity. *Mol. Cell* **53**, 140-147 (2014)

33. Amano, T., Tanabe, K., Eto, T., Narumiya, S., and Mizuno, K. LIM-kinase 2 induces formation of stress fibres, focal adhesions and membrane blebs, dependent on its activation by Rho-associated kinase-catalysed phosphorylation at threonine-505. *Biochem J* **354**, 149-159 (2001)

34. Narita, A. ADF/cofilin regulation from a structural viewpoint. *J Muscle Res Cell Motil* **41**, 141-151 (2020)

35. Miller, C. J., and Turk, B. E. Homing in: Mechanisms of Substrate Targeting by Protein Kinases. *Trends Biochem Sci* **43**, 380-394 (2018)

36. Johnson, J. L., Yaron, T. M., Huntsman, E. M., Kerelsky, A., Song, J., Regev, A., Lin, T. Y., Liberatore, K., Cizin, D. M., Cohen, B. M., Vasan, N., Ma, Y., Krismer, K., Robles, J. T., van de Kooij, B., van Vlimmeren, A. E., Andree-Busch, N., Kaufer, N. F., Dorovkov, M. V., Ryazanov, A. G., Takagi, Y., Kastenhuber, E. R., Goncalves, M. D., Hopkins, B. D., Elemento, O., Taatjes, D. J., Maucuer, A., Yamashita, A., Degterev, A., Uduman, M., Lu, J., Landry, S. D., Zhang, B., Cossentino, I., Linding, R., Blenis, J., Hornbeck, P. V., Turk, B. E., Yaffe, M. B., and Cantley, L. C. An atlas of substrate specificities for the human serine/threonine kinome. *Nature* **613**, 759-766 (2023)

37. Kang, H., Bradley, M. J., Cao, W., Zhou, K., Grintsevich, E. E., Michelot, A., Sindelar, C. V., Hochstrasser, M., and De La Cruz, E. M. Site-specific cation release drives actin filament severing by vertebrate cofilin. *Proc Natl Acad Sci U S A* **111**, 17821-17826 (2014)

38. Johnston, A. B., Collins, A., and Goode, B. L. High-speed depolymerization at actin filament ends jointly catalysed by Twinfilin and Srv2/CAP. *Nat Cell Biol* **17**, 1504-1511 (2015)

39. Hilton, D. M., Aguilar, R. M., Johnston, A. B., and Goode, B. L. Species-Specific Functions of Twinfilin in Actin Filament Depolymerization. *J Mol Biol* **430**, 3323-3336 (2018)

40. O'Shea, J. P., Chou, M. F., Quader, S. A., Ryan, J. K., Church, G. M., and Schwartz, D. pLogo: a probabilistic approach to visualizing sequence motifs. *Nat. Methods* **10**, 1211-1212 (2013)

41. Cao, W., Goodarzi, J. P., and De La Cruz, E. M. Energetics and kinetics of cooperative cofilin-actin filament interactions. *J Mol Biol* **361**, 257-267 (2006)

42. Graham, J. S., McCullough, B. R., Kang, H., Elam, W. A., Cao, W., and De La Cruz, E. M. Multi-platform compatible software for analysis of polymer bending mechanics. *PLoS One* **9**, e94766 (2014)

43. Tanaka, K., Takeda, S., Mitsuoka, K., Oda, T., Kimura-Sakiyama, C., Maeda, Y., and Narita, A. Structural basis for cofilin binding and actin filament disassembly. *Nat Commun* **9**, 1860 (2018)

44. Mwangangi, D. M., Manser, E., and Robinson, R. C. The structure of the actin filament uncapping complex mediated by twinfilin. *Sci Adv* **7**, (2021)
45. Leonard, S. A., Gittis, A. G., Petrella, E. C., Pollard, T. D., and Lattman, E. E. Crystal structure of the actin-binding protein actophorin from Acanthamoeba. *Nat Struct Biol* **4**, 369-373 (1997)