Title: Genome size evolution and phenotypic correlates in the poison frog family Dendrobatidae

Abstract

Adaptive and neutral processes have produced a spectrum of genome sizes across organisms.
Amphibians in particular possess a wide range in C-values, from <1 to over 125 pg. However,
the genome size of most amphibians is unknown, and no single family has been comprehensively
assessed. We provide new estimates for 32 poison frog species representing the major lineages
within Dendrobatidae using Feulgen staining of museum specimens and flow cytometry of fresh
tissue. We show that genome size in Dendrobatidae has likely evolved gradually, with potential
evolutionary rate shifts in the genera *Phyllobates* and *Hyloxalus*, which respectively possess
species with the largest (13.0 pg) and second smallest (2.6 pg) genomes in the family.
Phylogenetically controlled regression analyses indicate that genome size is positively correlated
with snout-vent-length, oocyte number, and clutch size, but negatively correlated with active
metabolic rate and metabolic scope. While body size and metabolic rate are also correlates of
toxicity, we found no relationship between genome size and evolution of chemical defense
within Dendrobatidae. Genome size evolution in Dendrobatidae provides insight into the
processes shaping genome size evolution over short timescales and establishes a novel system in
which to study the mechanistic links between genome size and organismal physiology.

Keywords: transposable elements, metabolic rate, amphibians, phenotypic evolution

Resumen

Los procesos adaptativos y neutrales han producido un espectro de tamaños de genoma a través
de la diversidad biológica. Los anfibios en particular poseen una amplia gama de valores C,
desde <1 pg hasta más de 125 pg. Sin embargo, se desconoce el tamaño del genoma de la
mayoría de especies de anfibio, y no se ha evaluado exhaustivamente su evolución en detalle
dentro de ninguna familia. Proporcionamos estimados del tamaño del genoma para 32 especies
de ranas venenosas que representan los principales linajes de la familia Dendrobatidae utilizando
la tinción de Feulgen con especímenes de museo y citometría de flujo con tejido fresco.
Mostramos que el tamaño del genoma en Dendrobatidae probablemente evolucione bajo el fecto
de selección purificadora, potencialmente con regímenes de selección distintos al ancestral
géneros *Phyllobates* e *Hyloxalus*, que respectivamente poseen especies con el genoma más
grande (13,0 pg) y el segundo más pequeño (2,6 pg) de la familia. Análisis de regresión con
control filogenético indican que el tamaño del genoma se correlaciona positivamente con la
longitud hocico-cloaca, el número de oocitos y el tamaño de la nidada, y negativamente con la
tasa metabólica activa y el alcance metabólico. Aunque el tamaño corporal y la tasa metabólica
se correlacionan con la toxicidad, no encontramos relación entre el tamaño del genoma y la
evolución de la defensa química dentro de Dendrobatidae. Nuestra evaluación del tamaño del
genoma en Dendrobatidae proporciona información sobre los procesos que dirigen la evolución
del tamaño del genoma en escalas de tiempo cortas y un nuevo sistema que facilitará las
investigaciones de los vínculos mecánicos entre el tamaño del genoma y la fisiología del
organismo.

Palabras clave: Elementos transponibles, tasa metabólica, anfibios, evolución fenotípica

Introduction

Genomes can expand or contract through nonadaptive and adaptive mechanisms. In eukaryotes, genomes expand primarily through the self-replication of transposable elements (TEs), which generates long stretches of repetitive DNA (Kidwell 2002; Sun et al. 2012). Genomic TE activity largely explains why genome size does not scale with gene number in eukaryotes (the C-value paradox; Cavalier-Smith 1978). Genome expansion may be deleterious because accumulating introns and repetitive elements could increase mutational load and mutational hazard (Lynch et al. 2006), and set lower limits to traits like cell size and the rate of cell replication (Gregory 2002). Under this view, genome expansion is a function of genetic drift, and genome expansion is likely in populations where purifying selection is weak and genetic drift is strong (i.e., small effective population size Ne; Lynch and Conery 2003). Notably, however, some classical studies of neutral genome size evolution included data spanning prokaryotes and eukaryotes without phylogenetic correction and may have overestimated the role of neutral evolution in generating variation in genome size (Whitney and Garland 2010; Smith 2016).

Several selective hypotheses for the evolution of genome size exist based on observed correlations between genome size and fitness-related traits. For example, in some species genome size correlates positively with cell size and body size and negatively with developmental rate, developmental complexity (extent of tissue specialization), and metabolic rate (Cavalier-Smith 1982; Jockusch 1997; Kozłowski et al. 2003; Womack et al. 2019). Ecological or physiological constraints on any of these traits could indirectly impose selection on genome size. For instance, miniaturization is associated with genome size reduction, which results in smaller cells and makes achieving the minimum number of cells required to build functional organs possible (Levy and Heald 2015; Decena-Segarra et al. 2020). On the other extreme, large genomes may require physiological accommodations, such as enucleation of red blood cells in plethodontid salamanders with especially large genomes (Mueller et al. 2008). However, the direction and magnitude of physiological correlates with genome size vary between taxonomic groups. For instance, metabolic rate correlates negatively with genome size in birds, but this correlation is only weakly supported in mammals and amphibians (Licht and Lowcock 1991; Vinogradov 1995; Gregory 2002). Thus, the adaptive dynamics underlying genome size evolution may be lineage-specific and/or influenced by multiple independent traits, and therefore difficult to discern without denser taxon sampling at lower taxonomic levels.

Amphibians in particular are an ideal group for testing evolutionary hypotheses of genome size evolution, as they vary dramatically in genome size (Liedtke et al. 2018) and several of its potential life history correlates, including body size, cell size, ontogeny, reproductive ecology, and metabolic rates (Gomes et al. 2004; Elinson and del Pino 2012; Levy and Heald 2015; Liedtke et al. 2022; Johnson et al. 2023). Within amphibians, three clades of the family Dendrobatidae (*sensu* Santos et al. 2009) have independently evolved bright coloration and the ability to sequester dietary toxins, along with other correlated changes, together known as the aposematic syndrome (Santos et al. 2003; Vences et al. 2003; Santos and Cannatella 2011). Chemical defense evolution has been associated with higher aerobic capacity (Santos and Cannatella 2011) and substantial variation in body size (Silverstone 1975, 1976; Santos and Cannatella 2011), which, in some chemically defended species, positively correlates with defensive alkaloid quantity (Saporito et al. 2010). Furthermore, *HSP90*, a heat-shock protein that could be involved in suppressing transposable elements in multiple animal lineages, such as *Drosophila* (Specchia et al. 2010; Gangaraju et al. 2011), rodents (Olivieri et al. 2012; Xiol et al.

2012), and nematodes (Ryan et al. 2016), was shown to bind to alkaloids (Caty et al. 2019).
Thus, natural selection for chemical defenses could indirectly impact genome size by acting on
body size, metabolic rate, or *HSP90* function. Because the effects of body size and metabolic rate
on genome size are thought to operate in competing directions, the specific effects that chemical
defense would have on genome size are unclear. Here, we leverage the variation of a suite of
traits in Dendrobatidae (Santos et al. 2003) to investigate the patterns and possible drivers of
genome size evolution. Taking advantage of the diversity of preserved specimens available in
natural history collections, we generated genome size estimates for 32 species, and used
phylogenetic comparative methods and publicly available phenotypic data to inquire into the
drivers of genome size evolution in dendrobatids.

Methods

Specimen identification

Species identities of museum specimens and data in the Animal Genome Size Database (AGSD)
were checked using locality information and photographs or specimens when available (Table
S1; see supplementary materials for full details).

Cell extraction, staining, and imaging

We extracted red blood cells from formalin-fixed specimens stored in ethanol in the Museum of
Vertebrate Zoology (MVZ). Our general approach follows Hardie et al. (2002), with
modifications by Womack et al. (2019), who specifically analyzed formalin-fixed amphibian
specimens. Cells were stained and imaged as detailed below in two separate batches. Batch one
included 22 dendrobatid species and 6 non-dendrobatid species with published genome sizes;
batch two included 10 dendrobatid species and 12 non-dendrobatid species (Table S2). We drew
blood cells 4 times from 1-2 individuals per species, deposited cells arbitrarily across a set of
microscope slides to prevent batch bias and allowed the cell extracts to air dry for 30 minutes.
We then used Feulgen staining as detailed in Hardie et al. (2002). In brief, we fixed cells in 85
methanol:10 formalin:5 acetic acid for 24 hours, rinsed cells under running water for 10-minutes,
hydrolyzed cells in 5.0 N HCl for 2 hours, briefly submerged cells in 0.1 N HCl, then stained
cells for 2 hours with Schiff's reagent (Fisher Scientific 632-99-5). Next, we rinsed cells with
bisulfite solution 3 times for 5 minutes, followed by a 10-minute wash under running water and 3
2-minute rinses in distilled water. We air-dried slides and mounted cover glass on slides using
Permount (Fisher SP15500).

We imaged cells using a SeBa camera (SEBACAM5C) mounted to an Laxco LMC-2000
microscope with a 100x immersion oil lens. We white-balanced each slide individually using
SeBa imaging software SebaView v3.7 prior to imaging.

Genome and cell size estimation

Using ImageJ v1.53c we calculated the integrated optical density (IOD) from cell images, which
measures the amount of stained material (i.e., DNA), in 6-66 nuclei per species (average = 19.93
nuclei, SD = 12.18). To convert IOD values to genome size estimates, we generated IOD
measurements for 15 species with published genome sizes and 2 species for which we estimated
genome sizes using flow cytometry (Tables S1 and S2; see below for methods), and used them to
generate a standard curve. When possible we selected species with multiple published genome
size measurements for use in the standard curve. We obtained the median IOD across nuclei for

each specimen, as IOD scores exhibited left or right-skewed distributions in some samples. For
species with more than one sampled specimen we obtained the median of specimen medians.We
built our standard curve as a linear mixed-effects model with genome size as the dependent
variable, IOD as a fixed-effect predictor, and sampling batch as a random-effect predictor, using
the lmer() function in the *lme4* R package (Bates et al. 2015).

The use of cells from formalin-fixed specimens for genome size estimation based on Feulgen
staining can result in artifacts because Schiff's reagent binds aldehydes (including
formaldehyde), and preservation and long-term storage can result in cell damage (Greilhuber and
Temsch 2001; Hardie et al. 2002). This being said, multiple studies have quantified genome size
and other morphonuclear parameters using Feulgen-stained nuclei obtained from formalin-fixed
specimens, both from natural history (e.g., Itgen et al. 2019, 2022; Womack et al. 2019; Wang et
al. 2021), and hospital archival collections (e.g., Dorman et al. 1990; Salmon et al. 1992; Fang et
al. 2004), with accuracy often similar to other methods including flow cytometry. At least in
some cases, such measurements have not been found to be particularly susceptible to variation in
the formalin fixation process and posterior storage conditions (Schimmelpenning et al. 1990),
although more general tests of these remain scarce. With this in mind, Feulgen-based genome
size estimates from formalin-fixed specimens can be considered reliable, as long as the quality of
such measurements is first evaluated, for instance by the inclusion of a set of samples whose
parameters of interest are known (Greilhuber and Temsch 2001; Greilhuber 2008). In the case of
amphibians, studies using Feulgen staining on formalin-fixed cells have obtained tight-fitting
correlations between IOD and genome size determined using flow cytometry (e.g., Itgen et al.
2019; Womack et al. 2019), $R^2 = 0.91$-0.93), close to those from studies using alcohol-preserved
(e.g. Liedtke et al.2018, $R^2 = 0.99$) or fresh (e.g., Decena-Segarra et al. 2020)[1]) tissues.

We ensured the reliability of our estimates in three complementary ways: First, we removed
specimens of *Pseudacris nigrita*, *Oophaga pumilio*, and *Phyllobates terribilis* from the
calibration curve because cells from these specimens showed clear morphological anomalies.
Second, we evaluated the goodness of fit of our calibration curve using Nakagawa and
Schielzeth's (2013) marginal R^2, which quantifies the variance explained by the fixed effects of a
mixed-effects model, using the r.squaredGLMM()function of the *MuMIn* R package (Bartoń
2024), and generated confidence intervals around our calibration curve parameters via
bootstrapping to evaluate its precision and susceptibility to outlier specimens. Finally, we
included seven species in both staining batches to assess repeatability in genome size estimates
between batches.

To estimate the cell size of each species we manually traced the cell membrane of 2-22 cells per
species (average = 12.30 cells, SD = 5.07) in ImageJ and calculated the area within each cell.
Cell sizes were averaged by species for subsequent analysis.

Flow cytometry

We used flow cytometry to generate genome size estimates for four species of the genus
Phyllobates: *P. terribilis, P. bicolor, P, aurotaenia*, and *P. vittatus*. All frog-rearing protocols
were approved by the University of Michigan's Institutional Animal Care and Use Committee
(protocol #PRO00010325). Captive-bred specimens were euthanized with an overdose of topical

[1] We could not find an R^2 value reported in this study, but the authors report estimating genome size within 10% or
20% of published values for most of their standard species.

171   benzocaine, and muscle samples were immediately dissected and flash-frozen on dry ice, and
172   kept frozen until further use. See above for animal care and use protocol details. Genome size
173   was estimated as described by Johnston et al. (2019) and details of flow cytometry methods are
174   located in supplementary methods.

Phylogenetic inference

To investigate the evolutionary patterns and correlates of genome size from a phylogenetic
perspective, we generated a time-calibrated phylogeny of our study species based on the
relationships inferred by Grant et al. (2017) and Vacher et al. (2017), using publicly available
DNA sequences from six loci (one mitochondrial, five nuclear; Table S3). We pruned the Grant
et al. (2017) tree to include only our target species using *ape* (in R v5.3; Paradis and Schliep
2019). *Anomaloglossus surinamensis* was not included in the Grant et al. (2017) phylogeny, so
we placed it as the sister species to *Anomaloglossus stepheni* following Vacher et al. (2017).
Although *Aromobates tokuko* was recently shown to have a remarkably small genome (0.8 pg;
Liedtke et al. 2018), there are no DNA sequences available for this species, so we did not include
it in our analyses. Because dendrobatid systematics remain understudied, and there is evidence
for both cryptic species and species complexes (e.g., Tarvin et al. 2017; Guillory et al. 2019;
Márquez et al. 2020), we used tree tips and sequences from the closest available locality to the
collection site of the specimens used to estimate genome size (details in Table S3).

We aligned sequences for each locus (Table S3) using MUSCLE v3.8.1551 (Edgar 2004), and
chose a partitioning strategy and models of sequence evolution using PartitionFinder2 (Lanfear
et al. 2017). The final (concatenated) matrix contained 5,493 bp (2,513 bp from mtDNA; 2,980
bp across 5 nuclear loci ranging between 316 bp to1,200 bp), with 35% gaps and 1204
parsimony-informative sites. We then ran a concatenated analysis on BEAST 2.6.5 (Bouckaert et
al. 2019), setting the Grant et al. (2017) topology as a hard constraint, and using three normally-
distributed calibration priors based on previous studies: The root of Dendrobatidae (i.e. the root
of the tree) was set to $N(\mu = 38.534, \sigma = 5.151)$ following Santos et al. (2009), and the nodes
corresponding to Dendrobatinae and Colostethinae (*sensu* Grant et al. 2006) were set to
$N(\mu=26.655, \sigma=4.791)$ and $N(\mu=19.392, \sigma=3.793)$, respectively, based on Guillory et al. (2019).

We used empirical base frequencies and a calibrated Yule tree prior (Heled and Drummond
2012), and set the Yule birth rate and clock rate priors to gamma(0.01, 50), and the GTR relative
rate priors to *gamma*(2, 0.5) for transitions and *gamma*(2, 0.25) for transversions. All other priors
were left at the default. Independent uncorrelated log-normal clocks were set for all partitions,
and a preliminary MCMC was run for 5 million steps, sampling every 5,000. We recovered
posterior distributions of the parameters describing clock rate variation among branches
(*ucldStdev*) that leaned heavily towards zero in four of the five nuclear loci, suggesting very little
rate variation at these genes, so they were modeled under a strict clock in further runs. To obtain
a final tree, we ran two independent MCMC samplers for 20 million steps, sampling every 5,000,
and merged their results after discarding the initial 10% of each one as burnin. Convergence and
mixing within and between runs were assessed based on effective sample sizes (ESS) estimates
and trace plots generated in Tracer v1.7.1 (Rambaut et al. 2018). A maximum clade credibility
(MCC) tree with common ancestor node heights was produced in TreeAnnotator (distributed
with BEAST) and used for all downstream analyses. The xml file used as input for BEAST2,
which includes alignments and software parameters, is available on the Dryad repository for this
paper.

Patterns of genome size evolution

We explored the patterns of genome size evolution along the dendrobatid phylogeny in several complementary ways. First, we used BAMM (Rabosky 2014) to estimate its rate of evolution across the tree, and to assess the degree to which discrete rate shifts at specific branches can explain the extant variation in dendrobatid genome size. We used the R package BAMMtools (Rabosky 2014) to generate priors appropriate for the temporal and taxonomic scale of our tree and phenotype, and ran BAMM for 10,000,000 steps, sampling every 5,000, for a total of 20,000 samples from the posterior, out of which we discarded 10% as burnin. From the post-burnin BAMM samples, we estimated the posterior distribution for the number of rate shifts along the tree, as well as the frequency of different rate shift configurations and their associated rates of phenotypic evolution using BAMMtools.

Next, we examined the evidence for discrete (i.e., saltational) evolutionary regime shifts in the evolution of genome size using the *l1ou* R package (Khabbazian et al. 2016) to fit and rank all possible single and multi-optimum Ornstein-Uhlenbeck (OU) models of phenotypic evolution (Hansen 1997) for our tree. Briefly, these models allow different subtrees to have distinct evolutionary "optima," which we can use to gauge whether specific branches of the tree have experienced rapid jumps in genome size. We used the default lasso approach implemented in *l1ou*, and model fit was assessed using the phylogenetically corrected Bayesian information criterion (pBIC; Khabbazian et al. 2016). To quantify the strength of support for each model, we calculated the pBIC weight (w_i for model i) as

$$w_i = \frac{\exp(-0.5\Delta_{pBIC_i})}{\sum_{k=1}^{K} \exp(-0.5\Delta_{pBIC_k})}$$

Where Δ_{pBICi} is the difference between a given model (model *i*) and the best model (i.e., the one with the smallest pBIC), and *K* is the total number of models.

Finally, we evaluated the fit of different continuous trait evolution models to our data, in order to gain further insight into the processes driving the evolution of genome size in dendrobatids. The analyses above suggested there may have been saltational events in the evolution of genome size (see Results section), so we first selected a set of four single and multi-regime model configurations, and fit Brownian Motion (Felsenstein 1973), and OU models based on these regimes. In addition, we fit Early Burst (Harmon et al. 2010) and White Noise models across the whole tree. BM models were fit using the brownie.lite() function in *phytools* (Revell 2024), OU models were fit using the fit_OU() function in *l1ou*, and the EB and White Noise models were fit using the fitContinuous() function of the *geiger* package (Pennell et al. 2014). Model fit was evaluated using the corrected Akaike Information Criterion (AICc), since pBIC has not been implemented beyond OU models. To ensure compatibility between model-fitting algorithms, we also fit the single-regime BM and OU models with fitContinuous(), which obtained identical likelihoods and parameter estimates to the functions mentioned above.

Correlates of genome size

To assess phenotypic correlates of genome size, we obtained data for several physiological, ecological, and life history traits from compilations available in the literature (Santos and Cannatella 2011; Santos 2012; Carvajal-Castro et al. 2021; full list of traits available in Table 1).

In addition, we obtained elevation from the collection localities of the specific museum
specimens used for genome size estimation when available, or the literature otherwise. We used
snout-vent length (SVL) as a proxy for body size, which we measured directly on adult
specimens or their photographs, except for the four *Phyllobates* species used in flow cytometry,
for which SVL values were pulled from the literature (Márquez et al. 2020), as study specimens
were juveniles. Finally, we scored each species as able or unable to sequester alkaloids based on
data compiled by Santos, Tarvin, and O'Connell (2016). All trait data are available in Table S4;
note that up to 47% of species are missing data for some traits, so the results of our correlations
should be taken as preliminary evidence supporting hypotheses that require further validation.

To assess trait correlations, we fit univariate phylogenetic-generalized-least-squares (PGLS;
Grafen 1989; Martins and Hansen 1997) linear regressions using the phylolm()function of the
phylolm R package (Ho and Ané 2014). Covariance between traits due to phylogeny was
modeled under the OU model with an estimated ancestral state (OUfixedRoot model). We added
mass as a covariate in models involving metabolic rates (data were obtained from Santos 2012).
To visualize relationships between traits as well as their phylogenetic distribution we generated
phylomorphospace plots (Sidlauskas 2008) using *phytools*. Because some species of *Phyllobates*
were outliers in terms of genome size and some of the tested predictors (see Results), and may
therefore generate overestimates of phenotypic correlations, we ran our linear regressions both
including and excluding this genus from the dataset.

Results

Genome size estimation

Our calibration curve displayed a very close fit between IOD and published genome size for all
species in both staining batches (marginal $R^2 = 0.94$), with narrow conference intervals around
the slope and intercept (Fig. 1A, B), and highly consistent estimates between batches (Fig. 1C).
In light of these results, our Feulgen-based estimates using formalin-fixed cells can be
considered good proxies for genome size to evaluate the general patterns and correlates of this
trait across the range of dendrobatid genome sizes. We note, however, that for more precise
purposes, such as validating the completeness of *de novo* genome assemblies, or for intraspecific
analyses, the Feulgen-based genome size estimates presented here (and in other similar studies)
should be verified using additional methods, such as flow cytometry. Based on our IOD curve
and flow cytometry workflow, we estimated the genome size of 32 species with previously
unknown genome sizes (Table S5). Our estimated genome sizes (C-value) in Dendrobatidae
ranged from 2.6 pg (*Hyloxalus infraguttatus*) to 13.0 pg (*Phyllobates bicolor*; Fig. 2).

Figure 1. Standard curves used to convert IOD values to genome sizes. Panels A and B represent the curves inferred
for each independent imaging batch. Solid lines represent the slope and intercept for each specific batch and broken
lines 95% confidence intervals. Panel C shows the repeatability of our estimates by plotting genome size estimates
for the seven species that were imaged in both batches. The diagonal represents the identity line.

Figure 2. Genome size variation across the dendrobatid phylogeny. The position of *A. tokuko* is uncertain. Numbers
to the right of the dot plot represent estimated or measured genome sizes for each species in pg. MYA: Million
Years Ago.

Patterns of genome size evolution

The rate of genome size evolution was relatively similar across the dendrobatid tree and stable through time, with faster rates concentrated in branches containing the five species of *Phyllobates* and *Hyloxalus elachyhistus* and *H. infraguttatus* (Fig. 3AB). Rate estimates from BAMM were generally concordant with those estimated under Brownian Motion (the BAMM phenotypic model assumes Brownian evolution; Rabosky et al. 2013), suggesting these estimates are accurate despite the number of species analyzed (Fig. S1). BAMM and *l1ou* both provided some evidence for discrete rate shifts. Specifically, the posterior distribution for the number of rate shifts recovered by BAMM peaked at 1-2 shifts (Fig. 3C), and the top models in all analyses contained scenarios with 2-3 evolutionary regimes that were better or comparable with single-regime models, although single-regime scenarios were also well supported (Fig. S2).

Based on the above evidence for multiple regimes of genome size evolution in different branches of the dendrobatid tree, we conducted a model-fitting exercise where we compared ten different single and multi-regime models of trait evolution, evolving under BM, OU, and EB dynamics (Fig. 3). The two best-fitting models, which together accounted for 74% of AICc weight were both multi-regime OU models with either *Phyllobates* departing from the background (i.e., ancestral) regime (wAICc = 0.498), or *Phyllobates* and *H. infraguttatus* each having regimes separate from the background (wAICc = 0.241). None of the remaining models carried AICc weights above 0.1 (Fig. 4, Table 1).

Figure 3. Rates of genome size evolution in dendrobatid frogs. A) Instantaneous rates of phenotypic evolution along branches of the dendrobatid phylogeny and B) across the phylogeny over time. C) Posterior distribution of the number of discrete rate shifts recovered by BAMM. The color ramp in panel A was generated using the Jenks method in the plot.bammdata() function of BAMMtools. Panel B is based on 100 time slices, the central line depicts the median posterior rate, and the surrounding polygon the 95% highest posterior density.

Figure 4. Models of continuous trait evolution tested for genome size in dendrobatids, along with their AICc
weights. Branches evolving under the background evolutionary regime are colored gray, and subtrees evolving
under distinct regimes are colored blue (*Phyllobates*) or magenta (*H. infraguttatus* and *H. elachyhistus*). The AICc
for each regime configuration under different model types (i.e., BM, OU, etc.) are presented above each tree, and the
two best-fitting models are highlighted in bold. All ten models were compared simultaneously. wAICc: Corrected
Akaike weight; BM: Brownian Motion; EB: Early Burst; OU: Ornstein–Uhlenbeck; WN: White Noise.

Correlates of genome size

Six of the 15 tested traits were correlated with genome size (Table 2). Body size, oocyte number,
clutch size, and cell area showed positive correlations with genome size. Active metabolic rate
(AMR; all metabolic variables measured in VO_2 ml*h^{-1}) and aerobic scope were negatively
correlated with genome size, after controlling for body mass and phylogeny, while basal
metabolic rate (BMR) was not. Because aerobic scope measures the difference between active
and basal metabolic rates, this suggests that the correlation between scope and genome size is
driven mainly by the active metabolic rate. Excluding *Phyllobates* from the models resulted in
lower effect sizes and weaker statistical significance, as expected by the removal of 2-5 data
points near the edges of trait distributions. Yet, the magnitude and direction of the correlations
were qualitatively similar, indicating that the obtained correlations are not entirely driven by
species in this genus. Regression parameters and associated statistics for the 15 predictors,
including and excluding *Phyllobates*, are given in Table 2, and Figure 5 shows the relationships
between genome size and the six correlated predictors.

Figure 5. Phylomorphospace plots of the relationships between genome size and the six predictors that showed significant correlations. Large dots represent extant species, connected by lines that depict phylogeny. Small dots indicate bivariate ancestral state reconstructions. Species in the genus *Phyllobates* are colored yellow and all others black. For the two metabolic variables (aerobic scope and AMR) we plot the linear residuals with mass instead of the raw values to provide a visualization that controls for body mass. For model-fitting, mass was used as a covariate instead.

Discussion

Genome size evolution in Dendrobatidae

Dendrobatid genome sizes vary across the full range characterized in Anura, with a minimum C-value of 0.8 pg (*Aromobates tokuko*) and a maximum value of 13.0 pg (*Phyllobates bicolor*). The greater than 10-fold variation in genome size among dendrobatids (with N = 35 species) is broader than what has been observed in the few other similarly well-sampled amphibian families, such as plethodontid salamanders with a 5.2-fold range (13.90 – 71.65 pg; N = 95 species), hylid tree frogs with a 4.3-fold range (1.54 – 6.61 pg; N = 41 species), or bufonid toads with a 2.5-fold range (2.91 – 7.31 pg; N = 31 species) (Gregory 2022). Given the lack of comprehensive phylogenetic sampling in these and other families, we expect that additional cases of broad variation in genome size at lower taxonomic scales among amphibians will be described in the future.

The apparent occupation by dendrobatids of the entire range of genome sizes covered by frogs in
just under 40 million years could be an artifact of the general lack of data for anuran genome
sizes, as noted above. Alternatively, it may be explained by their much faster rate of genome size
evolution (Fig. 2B), compared to the average rates across all frogs and other amphibian orders,
which are over one order of magnitude lower (Liedtke et al. 2018). Previous work suggests that,
across amphibians, genome size has evolved in a mostly gradual manner, with rare but
pronounced saltational genome size shifts (Liedtke et al. 2018). Our results, which focused on a
shallower phylogenetic scale, point to a somewhat similar scenario, with key differences.
Phylogenetic comparative analyses provided good support for one or two discrete regime shifts
in the evolution of genome size, although single-regime models were also supported in some
cases (i.e., BAMM, *l1ou* under the pBIC), raising the possibility that the signature for discrete
shifts come from clades at the extremes of the distribution of a quickly-evolving trait.
Considering the number of species analyzed is on the low end for some of these analyses, we
refrain from drawing definitive conclusions in this regard.

Irrespective of the number of independent evolutionary regimes, OU models were
unambiguously supported over all others tested, suggesting that the diversification of genome
sizes across Dendrobatidae has likely been marked by the existence of relatively stable
"attractors", such as adaptive optima, towards which lineages are drawn along their evolutionary
trajectories. Under this framework, a regime shift constitutes a shift in the adaptive optimum,
which results in evolution towards the new optimum due to positive selection (i.e., adaptation),
followed by stabilizing selection around it. This contrasts with the patterns obtained across
amphibian species, where lineages were found to gradually explore phenotypic space in a
Brownian fashion, occasionally experience rapid jumps in genome size, and then continue their
Brownian walk, suggesting that within Dendrobatidae genome size may have experienced
stronger stabilizing selection than the general trend across Amphibia. Whether this is due to
dataset characteristics, such as the temporal scales involved (~40 vs. 350 MY), the degree of
phylogenetic sampling (34 species within a family vs. 464 species across three orders) or the
range of genome sizes investigated (0.8–13.0 pg vs. 0.8–121.0 pg), or it actually reflects
dendrobatids having distinct evolutionary dynamics from the majority of other amphibian
lineages remains to be clarified. Phylogenetic work with finer taxon sampling, as well as
mechanistic studies on the effects of genome size change in different taxa would be especially
insightful.

Phenotypic correlates of genome size

We identified a significant correlation between genome size and six fitness-related traits: active
metabolic rate (AMR), metabolic scope, snout-vent length (SVL), cell size, number of oocytes,
and clutch size. Correlations with metabolic scope and the number of oocytes are not significant
when we exclude the 5 *Phyllobates* lineages, yet qualitatively similar trends persist, and p-value
changes are small (metabolic scope: p = 0.0496 vs. 0.054; number of oocytes: 0.0498 vs. 0.071).
These changes are likely a combination of lower power (e.g., 18 vs 14 samples for oocyte
number) and the removal of some of the extreme values that *Phyllobates* species represent in the
dataset. In any case, similar traits to those in question, namely AMR and clutch size, remained
correlated to genome size regardless of the species involved, so our conclusions are robust to
these small statistical variations.

Amongst these correlates, all but AMR and scope were positively correlated with genome size.
AMR and scope are thought to increase with origins of chemical defense (Santos and Cannatella
2011), while body size (SVL) may be of particular significance in poison frogs because it
positively correlates with toxicity, bright coloration, and mating call complexity (Saporito et al.
2010; Santos and Cannatella 2011; Santos et al. 2014; Márquez et al. 2020). Despite our finding
that three traits of the aposematic syndrome correlate with genome size, we found no relationship
between alkaloid sequestration itself and genome size. AMR, scope, and SVL were correlated
with genome size across our entire dataset of toxic and non-toxic species, even when toxicity is
included as a covariate, further suggesting that the statistical relationship between these traits and
genome size is unrelated to chemical defense.

The mechanistic link between genome size and metabolic rate is unclear, although it has been
suggested to be mediated through cell size (Cavalier-Smith 1982; Gregory 2001), which we also
found to be positively correlated with genome size. Larger cells divide less frequently and have a
lower surface-area-to-volume ratio and thus a slower rate of nutrient and gas exchange, both of
which should decrease metabolic rate (Gregory 2001). However, this pattern is not consistent
across taxa. A negative correlation between genome size and basal metabolic rate (BMR) has
been observed multiple times in birds and in mammals, although with weak support (Vinogradov
1995; Gregory 2002; Kozłowski et al. 2003). A negative correlation between genome size and
BMR has been observed across all of Lissamphibia, but not when Anura and Caudata are
considered independently (Gregory 2003). In addition, AMR and genome size are negatively
correlated in some salamanders, but only at stress-inducing temperatures (Licht and Lowcock
1991), and ploidy level and AMR are negatively correlated in tadpoles but not froglets of the
hybridogenetic frog *Pelophylax esculentus* (Hermaniuk et al. 2017). Finally, genome size does
not appear to be associated with evolutionary saltations in BMR across extant vertebrates
(Gardner et al. 2020), although there are exceptions in certain mammalian lineages (Uyeda et al.
2017). While AMR was not considered in some of these studies, potentially due to lack of data
availability, our observation that AMR and metabolic scope are negatively correlated with
genome size (controlling for mass) is fairly novel amongst amphibians. Dendrobatids exhibit
highly specialized life history and physiology, including significant investment in parental care,
coloration, toxin sequestration, and diet, all of which could potentially impact variation in
metabolic rate. Additionally, AMR (but not BMR) is positively correlated with molecular
evolution rate in Dendrobatidae (Santos 2012). Thus, studies of Dendrobatidae may present an
opportunity to clarify the mechanistic link between genome size and metabolic rate.

Similarly to metabolic rate, the positive correlation between body size and genome size is
thought to be mediated through cell size, and, additionally, through cell number (Arendt 2007;
Hessen et al. 2013). While the mechanism underlying the positive correlation between genome
size and cell size remains unclear, it may be driven by variation in transcriptional output or
stoichiometric relationships between DNA amount and cytoplasmic protein levels (Mueller
2015). Because cell size is positively correlated with SVL and genome size in our data set,
selection on cell size or related attributes may mediate the positive correlation between body size
and genome size in poison frogs. In smaller species that have relatively large genomes for their
size compared to other vertebrates (e.g., *Ranitomeya*: 4.4 – 6.8 pg: Stuckert et al. 2021), cell size
could limit cell number and constrain morphological complexity (i.e., biological size), as has
been seen in other amphibians (Levy and Heald 2015; Womack et al. 2019; Decena-Segarra et
al. 2020). One potential consequence of miniaturization when genome size is not concomitantly

reduced is the reduced complexity of the visual nervous system and hindering of vision-based behaviors such as active foraging, as has been observed in salamanders that have very large genomes (>30 pg; Matsushima et al. 1989). Given that some dendrobatids are diet specialists that actively forage for ants and mites, and that this behavior is associated with chemical defense, the loss of active foraging could be of particular consequence in Dendrobatidae (Toft 1995; Santos and Cannatella 2011; Tarvin et al. 2024). Furthermore, visual communication is thought to play an important role in several aspects of poison frog biology (e.g., Summers et al. 1999; Pröhl 2005; Yang et al. 2019). Thus, natural selection for the maintenance of body structures used in vision could play a role in genome size reduction in some dendrobatid species. We note, however, that the genera with the smallest genomes in our dataset, *Hyloxalus* and *Aromobates*, are neither miniaturized, brightly colored, nor outliers with respect to AMR (a correlate of active foraging; Santos and Cannatella 2011), thus we do not find strong support for this dynamic amongst the lineages we sampled.

Finally, we note that differences in genome size among poison frog species may be associated with differences in the activity of TEs, which appear to be currently proliferating in the genomes of at least two species that we examined (Rogers et al. 2018; Dittrich et al. 2024). Large genomes are thought to be hazardous in terms of fitness because of their larger mutation target, slower replication rates, and impacts on biological size. However, our understanding of the dynamics of genome size evolution are built on a small portion of living species, and it is plausible that amphibians, given their large genome sizes, may operate under different rules. For example, genome size is expected to increase in small populations where drift has a greater impact than selection on the fate of genetic variation. However, range size, a correlate of population size, is unrelated to genome size in salamanders (Rios-Carlos et al. 2024). In some salamanders, genome size is unrelated to TE deletion rate or genome-wide selection efficiency, which are both expected to decrease with larger genomes under current theory (Rios-Carlos et al. 2024; Wang et al. 2024). Differences in recombination rates (Rios-Carlos et al. 2024), DNA loss rates (Sun and Mueller 2014; Wang et al. 2024), and TE activity may help further explain why so many amphibians, including some poison frogs, have large genomes.

Limitations

The wealth of formalin-fixed specimens in natural history collections around the world represent one of the richest and most complete resources to study phenotypic diversity — including genome size — in natural populations. However, chemicals used during specimen preservation such as formaldehyde and ethanol can affect nuclei compaction and Schiff's reagent stain intensity. Museum specimens are usually fixed using a 4% solution of formaldehyde then stored long-term in 70% ethanol. Although formalin- and alcohol-fixation is also part of the Feulgen staining process, an empirical comparison of genome size estimates using Feulgen staining of fixed versus fresh tissues would help establish possible impacts of fixation and storage. Until these impacts are known, we recommend that, when precise estimates are required, genome size estimates from formalin-fixed tissues should be confirmed using fresh tissues and/or additional biological replicates. Despite potential caveats of using formalin-fixed tissues, we believe that any introduced errors are unlikely to overturn the general evolutionary patterns we identified in this study.

Conclusion

The drivers of genome size evolution in eukaryotes remain a field of active research in evolutionary biology. Our results show that in dendrobatid poison frogs, genome size has evolved at much faster rates than in other amphibians — to the point of encompassing the entire range of genome sizes observed across anurans to date. Phylogenetic analyses suggest that this rapid evolution has proceeded primarily in a gradual manner and at a constant rate, constrained by purifying selection, and possibly with rare but appreciable events of punctuated evolution, although evidence for the latter is not definitive. Correlations between genome size and life history traits suggest that cell size may mediate changes in genome size within the group. Mechanisms underlying the relationship between cell size, metabolic rate, and genome size are not well understood, yet wide variation in each of these traits in Dendrobatidae could make the group suitable to provide mechanistic insight with future studies. Together these patterns suggest roles both for adaptive and neutral processes in shaping genome size evolution in Dendrobatidae. Our study contributes to understanding the multidimensional drivers of genome size, and sets the stage for future investigation into mechanisms driving genome size evolution in vertebrates.

Data Accessibility

Raw data files, R scripts and detailed metadata are available for download from the Dryad Digital Repository http://doi.org/10.5061/dryad.0cfxpnw7d.

Table Legends

Table 1. Phylogenetically-aware linear regression coefficients of genome size against 15 morphological, metabolic, and life history traits, including or excluding species in the genus *Phyllobates*. Body mass was included as a covariate in models involving metabolic rates.

Table 2. Fit of various models of continuous trait evolution to genome size data in Dendrobatidae. Models are ranked based on AICc weight.

References

Arendt, J. 2007. Ecological correlates of body size in relation to cell size and cell number: patterns in flies, fish, fruits and foliage. Biol. Rev. Camb. Philos. Soc. 82:241–256. Wiley.

Bartoń, K. 2024. MuMIn: Multi-Model Inference. R package version 1.48.4. https://CRAN.R-project.org/package=MuMIn.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models Usinglme4. J. Stat. Softw. 67. Foundation for Open Access Statistic.

Bouckaert, R., T. G. Vaughan, J. Barido-Sottani, S. Duchêne, M. Fourment, A. Gavryushkina, J. Heled,

G. Jones, D. Kühnert, N. De Maio, M. Matschiner, F. K. Mendes, N. F. Müller, H. A. Ogilvie, L.

du Plessis, A. Popinga, A. Rambaut, D. Rasmussen, I. Siveroni, M. A. Suchard, C.-H. Wu, D.

Xie, C. Zhang, T. Stadler, and A. J. Drummond. 2019. BEAST 2.5: An advanced software

platform for Bayesian evolutionary analysis. PLoS Comput. Biol. 15:e1006650.

Carvajal-Castro, J. D., F. Vargas-Salinas, S. Casas-Cardona, B. Rojas, and J. C. Santos. 2021.

Aposematism facilitates the diversification of parental care strategies in poison frogs. Sci. Rep.

11:19047.

Caty, S. N., A. Alvarez-Buylla, G. D. Byrd, C. Vidoudez, A. B. Roland, E. E. Tapia, B. Budnik, S. A.

Trauger, L. A. Coloma, and L. A. O. Connell. 2019. Molecular physiology of chemical defenses

in a poison frog. J. Exp. Biol. 222:jeb204149.

Cavalier-Smith, T. 1978. Nuclear volume control by nucleoskeletal DNA, selection for cell volume and

cell growth rate, and the solution of the DNA C-value paradox. J. Cell Sci. 34:247–278. The

Company of Biologists.

Cavalier-Smith, T. 1982. Skeletal DNA and the evolution of genome size. Annu. Rev. Biophys. Bioeng.

11:273–302. Annual Reviews.

Decena-Segarra, L. P., L. Bizjak-Mali, A. Kladnik, S. K. Sessions, and S. M. Rovito. 2020.

Miniaturization, genome size, and biological size in a diverse clade of salamanders. Am. Nat.

196:634–648.

Dittrich, C., F. Hoelzl, S. Smith, C. A. Fouilloux, D. J. Parker, L. A. O'Connell, L. S. Knowles, M.

Hughes, A. Fewings, R. Morgan, B. Rojas, and A. A. Comeault. 2024. Genome assembly of the

dyeing poison frog provides insights into the dynamics of transposable element and genome-size

evolution. Genome Biol. Evol. 16:evae109. Oxford University Press (OUP).

Dorman, A., D. Graham, B. Curran, K. Henry, and M. Leader. 1990. Ploidy of smooth muscle tumours:

retrospective image analysis study of formalin fixed, paraffin wax embedded tissue. J. Clin.

Pathol. 43:465–468. BMJ.

Edgar, R. C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

559   Nucleic Acids Res. 32:1792–1797.

560 Elinson, R. P., and E. M. del Pino. 2012. Developmental diversity of amphibians. Wiley Interdiscip. Rev.

561   Dev. Biol. 1:345–369. Wiley.

562 Fang, M., E. Lew, M. Klein, T. Sebo, Y. Su, and R. Goyal. 2004. DNA abnormalities as marker of risk

563   for progression of Barrett's esophagus to adenocarcinoma: image cytometric DNA analysis in

564   formalin-fixed tissues. Am. J. Gastroenterol. 99:1887–1894. Ovid Technologies (Wolters Kluwer

565   Health).

566 Felsenstein, J. 1973. Maximum-likelihood estimation of evolutionary trees from continuous characters.

567   Am. J. Hum. Genet. 25:471–492.

568 Gangaraju, V. K., H. Yin, M. M. Weiner, J. Wang, X. A. Huang, and H. Lin. 2011. Drosophila Piwi

569   functions in Hsp90-mediated suppression of phenotypic variation. Nat. Genet. 43:153–158.

570   Springer Science and Business Media LLC.

571 Gardner, J. D., M. Laurin, and C. L. Organ. 2020. The relationship between genome size and metabolic

572   rate in extant vertebrates. Philos. Trans. R. Soc. Lond. B Biol. Sci. 375:20190146. The Royal

573   Society.

574 Gomes, F. R., J. G. Chauí-Berlinck, J. E. P. W. Bicudo, and C. A. Navas. 2004. Intraspecific relationships

575   between resting and activity metabolism in anuran amphibians: influence of ecology and

576   behavior. Physiol. Biochem. Zool. 77:197–208. University of Chicago Press.

577 Grafen, A. 1989. The phylogenetic regression. Philos. Trans. R. Soc. Lond. B Biol. Sci. 326:119–157.

578   The Royal Society.

579 Grant, T., D. R. Frost, J. P. Caldwell, R. Gagliardo, C. F. B. Haddad, P. J. R. Kok, D. B. Means, B. P.

580   Noonan, W. E. Schargel, and W. C. Wheeler. 2006. Phylogenetic systematics of dart-poison frogs

581   and their relatives (Amphibia: Athesphatanura: Dendrobatidae). Bull. Am. Mus. Nat. Hist. 299:1–

582   262.

583 Grant, T., M. Rada, M. Anganoy-Criollo, A. Batista, P. H. Dias, A. M. Jeckel, D. J. Machado, and J. V.

584   Rueda-Almonacid. 2017. Phylogenetic systematics of dart-poison frogs and their relatives

revisited (Anura: Dendrobatoidea). South Am. J. Herpetol. 12:S1–S90.

Gregory, T. R. 2022. Animal Genome Size Database.

Gregory, T. R. 2001. Coincidence, coevolution, or causation? DNA content, cellsize, and the C☐value

enigma. Biol. Rev. Camb. Philos. Soc. 76:65–101. Wiley.

Gregory, T. R. 2002. Genome size and developmental complexity. Genetica 115:131–146.

Gregory, T. R. 2003. Variation across amphibian species in the size of the nuclear genome supports a

pluralistic, hierarchical approach to the C-value enigma. Biol. J. Linn. Soc. Lond. 79:329–339.

Oxford University Press (OUP).

Greilhuber, J. 2008. Cytochemistry and C-values: the less-well-known world of nuclear DNA amounts.

Ann. Bot. 101:791–804. Oxford University Press (OUP).

Greilhuber, J., and E. M. Temsch. 2001. Feulgen densitometry: some observations relevant to best

practice in quantitative nuclear DNA content determination. Acta Bot. Croat. 60:285–298.

Guillory, W. X., M. R. Muell, K. Summers, and J. L. Brown. 2019. Phylogenomic reconstruction of the

Neotropical poison frogs (Dendrobatidae) and their conservation. Diversity 11.

Hansen, T. F. 1997. Stabilizing Selection and the Comparative Analysis of Adaptation. Evolution

51:1341.

Hardie, D. C., T. R. Gregory, and P. D. N. Hebert. 2002. From pixels to picograms: a beginners' guide to

genome quantification by Feulgen image analysis densitometry. J. Histochem. Cytochem.

50:735–749.

Harmon, L. J., J. B. Losos, T. Jonathan Davies, R. G. Gillespie, J. L. Gittleman, W. Bryan Jennings, K. H.

Kozak, M. A. McPeek, F. Moreno-Roark, T. J. Near, A. Purvis, R. E. Ricklefs, D. Schluter, J. A.

Schulte Ii, O. Seehausen, B. L. Sidlauskas, O. Torres-Carvajal, J. T. Weir, and A. Ø. Mooers.

2010. Early bursts of body size and shape evolution are rare in comparative data. Evolution

64:2385–2396. Wiley.

Heled, J., and A. J. Drummond. 2012. Calibrated tree priors for relaxed phylogenetics and divergence

time estimation. Syst. Biol. 61:138–149. Oxford University Press (OUP).

Hermaniuk, A., M. Rybacki, and J. R. E. Taylor. 2017. Metabolic rate of diploid and triploid edible frog

Pelophylax esculentus correlates inversely with cell size in tadpoles but not in frogs. Physiol.

Biochem. Zool. 90:230–239. Physiol Biochem Zool.

Hessen, D. O., M. Daufresne, and H. P. Leinaas. 2013. Temperature-size relations from the cellular-

genomic perspective. Biol. Rev. Camb. Philos. Soc. 88:476–489. Wiley.

Ho, L. si T., and C. Ané. 2014. A linear-time algorithm for Gaussian and non-Gaussian trait evolution

models. Syst. Biol. 63:397–408.

Itgen, M. W., G. R. Natalie, D. S. Siegel, S. K. Sessions, and R. L. Mueller. 2022. Genome size drives

morphological evolution in organ-specific ways. Evolution 76:1453–1468. Wiley.

Itgen, M. W., P. Prša, R. Janža, L. Skubic, J. H. Townsend, A. Kladnik, L. B. Mali, and S. K. Sessions.

2019. Genome size diversification in central American bolitoglossine salamanders (Caudata;

Plethodontidae). Copeia 107:560. American Society of Ichthyologists and Herpetologists (ASIH).

Jockusch, E. L. 1997. An evolutionary correlate of genome size change in plethodontid salamanders.

Proc. Biol. Sci. 264:597–604. The Royal Society.

Johnson, J. V., C. Finn, J. Guirguis, L. E. B. Goodyear, L. P. Harvey, R. Magee, S. Ron, and D.

Pincheira-Donoso. 2023. What drives the evolution of body size in ectotherms? A global analysis

across the amphibian tree of life. Glob. Ecol. Biogeogr. 32:1311–1322. Wiley.

Johnston, J. S., A. Bernardini, and C. E. Hjelmen. 2019. Genome size estimation and quantitative

cytogenetics in insects. Methods Mol. Biol. 1858:15–26.

Khabbazian, M., R. Kriebel, K. Rohe, and C. Ané. 2016. Fast and accurate detection of evolutionary

shifts in Ornstein–Uhlenbeck models. Methods Ecol. Evol. 7:811–824. Wiley.

Kidwell, M. G. 2002. Transposable elements and the evolution of genome size in eukaryotes. Genetica

115:49–63.

Kozłowski, J., M. Konarzewski, and A. T. Gawelczyk. 2003. Cell size as a link between noncoding DNA

and metabolic rate scaling. Proc. Natl. Acad. Sci. U. S. A. 100:14080–14085. Proceedings of the

National Academy of Sciences.

Lanfear, R., P. B. Frandsen, A. M. Wright, T. Senfeld, and B. Calcott. 2017. PartitionFinder 2: New

methods for selecting partitioned models of evolution for molecular and morphological

phylogenetic analyses. Mol. Biol. Evol. 34:772–773.

Levy, D. L., and R. Heald. 2015. Biological scaling problems and solutions in amphibians. Cold Spring

Harb. Perspect. Biol. 8:a019166. Cold Spring Harbor Laboratory.

Licht, L. E., and L. A. Lowcock. 1991. Genome size and metabolic rate in salamanders. Comparative

Biochemistry and Physiology Part B: Comparative Biochemistry 100:83–92.

Liedtke, H. C., D. J. Gower, M. Wilkinson, and I. Gomez-Mestre. 2018. Macroevolutionary shift in the

size of amphibian genomes and the role of life history and climate. Nat Ecol Evol 2:1792–1799.

Liedtke, H. C., J. J. Wiens, and I. Gomez-Mestre. 2022. The evolution of reproductive modes and life

cycles in amphibians. Nat. Commun. 13:7039. Springer Science and Business Media LLC.

Lynch, M., and J. S. Conery. 2003. The origins of genome complexity. Science 302:1401–1404.

American Association for the Advancement of Science (AAAS).

Lynch, M., B. Koskella, and S. Schaack. 2006. Mutation pressure and the evolution of organelle genomic

architecture. Science 311:1727–1730. American Association for the Advancement of Science

(AAAS).

Márquez, R., T. P. Linderoth, D. Mejía-Vargas, R. Nielsen, A. Amézquita, and M. R. Kronforst. 2020.

Divergence, gene flow, and the origin of leapfrog geographic distributions: The history of colour

pattern variation in Phyllobates poison-dart frogs. Mol. Ecol. 29:3702–3719. Wiley.

Martins, E. P., and T. F. Hansen. 1997. Phylogenies and the comparative method: A general approach to

incorporating phylogenetic information into the analysis of interspecific data. Am. Nat. 149:646–

667. University of Chicago Press.

Matsushima, T., U. Dicke, D. B. Wake, K. C. Nishikawa, and G. Roth. 1989. Mechanics and

neuromorphology of feeding in amphibians. Neth. J. Zool. 40:115–135. Brill.

Mueller, R. L. 2015. Genome biology and the evolution of cell-size diversity. Cold Spring Harb. Perspect.

Biol. 7:a019125. Cold Spring Harbor Laboratory.

Mueller, R. L., T. R. Gregory, S. M. Gregory, A. Hsieh, and J. L. Boore. 2008. Genome size, cell size,

and the evolution of enucleated erythrocytes in attenuate salamanders. Zoology (Jena) 111:218–

230. Elsevier BV.

Nakagawa, S., and H. Schielzeth. 2013. A general and simple method for obtainingR2from generalized

linear mixed-effects models. Methods Ecol. Evol. 4:133–142. Wiley.

Olivieri, D., K.-A. Senti, S. Subramanian, R. Sachidanandam, and J. Brennecke. 2012. The cochaperone

shutdown defines a group of biogenesis factors essential for all piRNA populations in Drosophila.

Mol. Cell 47:954–969. Elsevier BV.

Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R.

Pennell, M. W., J. M. Eastman, G. J. Slater, J. W. Brown, J. C. Uyeda, R. G. FitzJohn, M. E. Alfaro, and

L. J. Harmon. 2014. geiger v2.0: an expanded suite of methods for fitting macroevolutionary

models to phylogenetic trees. Bioinformatics 30:2216–2218.

Pröhl, H. 2005. Territorial behavior in dendrobatid frogs. Journal of H 39:354–365.

Rabosky, D. L. 2014. Automatic detection of key innovations, rate shifts, and diversity-dependence on

phylogenetic trees. PLoS One 9:e89543.

Rabosky, D. L., F. Santini, J. Eastman, S. A. Smith, B. Sidlauskas, J. Chang, and M. E. Alfaro. 2013.

Rates of speciation and morphological evolution are correlated across the largest vertebrate

radiation. Nat. Commun. 4:1958.

Rambaut, A., A. J. Drummond, D. Xie, G. Baele, and M. A. Suchard. 2018. Posterior summarization in

Bayesian phylogenetics using tracer 1.7. Syst. Biol. 67:901–904.

Revell, L. J. 2024. phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and

other things). PeerJ 12:e16505. PeerJ Inc.

Rios-Carlos, H., M. G. Segovia-Ramírez, M. K. Fujita, and S. M. Rovito. 2024. Genomic gigantism is not

associated with reduced selection efficiency in neotropical salamanders. J. Mol. Evol. 92:371–

380. Springer Science and Business Media LLC.

Rogers, R. L., L. Zhou, C. Chu, R. Márquez, A. Corl, T. Linderoth, L. Freeborn, M. D. MacManes, Z.

Xiong, J. Zheng, C. Guo, X. Xun, M. R. Kronforst, K. Summers, Y. Wu, H. Yang, C. L.

Richards-Zawacki, G. Zhang, and R. Nielsen. 2018. Genomic takeover by transposable elements

in the strawberry poison frog. Mol. Biol. Evol. 35:2913–2927.

Ryan, C. P., J. C. Brownlie, and S. Whyard. 2016. Hsp90 and physiological stress are linked to

autonomous transposon mobility and heritable genetic change in nematodes. Genome biology and

evolution 8:3794–3805.

Salmon, I., P. Gasperin, J. L. Pasteels, R. Heimann, and R. Kiss. 1992. Relationship between

histopathologic typing and morphonuclear assessments of 238 thyroid lesions. Digital cell image

analysis performed on Feulgen-stained nuclei from formalin-fixed, paraffin-embedded materials.

Am. J. Clin. Pathol. 97:776–786. Oxford University Press (OUP).

Santos, J. C. 2012. Fast molecular evolution associated with high active metabolic rates in poison frogs.

Mol. Biol. Evol. 29:2001–2018.

Santos, J. C., M. Baquero, C. Barrio-Amorós, L. A. Coloma, L. K. Erdtmann, A. P. Lima, and D. C.

Cannatella. 2014. Aposematism increases acoustic diversification and speciation in poison frogs.

Proc. R. Soc. Lond. B Biol. Sci. 281:20141761.

Santos, J. C., and D. C. Cannatella. 2011. Phenotypic integration emerges from aposematism and scale in

poison frogs. Proc. Natl. Acad. Sci. U. S. A. 108:6175–6180.

Santos, J. C., L. A. Coloma, and D. C. Cannatella. 2003. Multiple, recurring origins of aposematism and

diet specialization in poison frogs. Proc. Natl. Acad. Sci. U. S. A. 100:12792–12797.

Santos, J. C., L. A. Coloma, K. Summers, J. P. Caldwell, R. Ree, and D. C. Cannatella. 2009. Amazonian

amphibian diversity is primarily derived from late Miocene Andean lineages. PLoS Biol. 7:0448–

0461.

Santos, J. C., R. D. Tarvin, and L. A. O'Connell. 2016. A review of chemical defense in poison frogs

(Dendrobatidae): ecology, pharmacokinetics, and autoresistance. Pp. 305–337 *in* B. A. Schulte, T.

E. Goodwin, and M. H. Ferkin, eds. Chemical Signals in Vertebrates 13. Springer International

Publishing, Switzerland.

Saporito, R. A., M. Isola, V. C. Maccachero, K. Condon, and M. A. Donnelly. 2010. Ontogenetic scaling

of poison glands in a dendrobatid poison frog. J. Zool. 282:238–245.

Schimmelpenning, H., U. G. Falkmer, K. Hamper, G. Seifert, and G. U. Auer. 1990. Variations in

Feulgen stainability of epithelial parenchymal cells extracted from paraffin-embedded salivary

gland specimens. Cytometry 11:475–480. Wiley.

Sidlauskas, B. 2008. Continuous and arrested morphological diversification in sister clades of

characiform fishes: a phylomorphospace approach. Evolution 62:3135–3156.

Silverstone, P. A. 1975. A revision of the poison-arrow frogs of the genus *Dendrobates* Wagler. Natural

History Museum of Los Angeles County 21.

Silverstone, P. A. 1976. A revision of the poison-arrow frogs of the genus *Phyllobates* Bibron in Sagra

(Family Dendrobatidae). Natural History Museum of Los Angeles County 27:1–53.

Smith, D. R. 2016. The mutational hazard hypothesis of organelle genome evolution: 10 years on. Mol.

Ecol. 25:3769–3775.

Specchia, V., L. Piacentini, P. Tritto, L. Fanti, R. D'Alessandro, G. Palumbo, S. Pimpinelli, and M. P.

Bozzetti. 2010. Hsp90 prevents phenotypic variation by suppressing the mutagenic activity of

transposons. Nature 463:662–665. Springer Science and Business Media LLC.

Stuckert, A. M. M., M. Chouteau, M. McClure, T. M. LaPolice, T. Linderoth, R. Nielsen, K. Summers,

and M. D. MacManes. 2021. The genomics of mimicry: Gene expression throughout

development provides insights into convergent and divergent phenotypes in a Müllerian mimicry

system. Mol. Ecol. 30:4039–4061. Wiley.

Summers, K., R. Symula, M. Clough, and T. Cronin. 1999. Visual mate choice in poison frogs. Proc.

Biol. Sci. 266:2141–2145.

Sun, C., and R. L. Mueller. 2014. Hellbender genome sequences shed light on genomic expansion at the

base of crown salamanders. Genome Biol. Evol. 6:1818–1829. Oxford University Press (OUP).

Sun, C., D. B. Shepard, R. A. Chong, J. López Arriaza, K. Hall, T. A. Castoe, C. Feschotte, D. D.

Pollock, and R. L. Mueller. 2012. LTR retrotransposons contribute to genomic gigantism in

plethodontid salamanders. Genome Biol. Evol. 4:168–183. Oxford University Press (OUP).

Tarvin, R. D., J. L. Coleman, D. A. Donoso, M. Betancourth-Cundar, K. López-Hervas, K. S. Gleason, J.

R. Sanders, J. M. Smith, S. R. Ron, J. C. Santos, B. E. Sedio, D. C. Cannatella, and R. Fitch.

2024. Passive accumulation of alkaloids in non-toxic frogs challenges paradigms of the origins of

acquired chemical defenses.

Tarvin, R. D., E. A. Powell, J. C. Santos, S. R. Ron, and D. C. Cannatella. 2017. The birth of

aposematism: high phenotypic divergence and low genetic diversity in a young clade of poison

frogs. Mol. Phylogenet. Evol. 109:283–295. Elsevier Inc.

Toft, C. A. 1995. Evolution of diet specialization in poison-dart frogs (Dendrobatidae). Herpetologica

51:202–216.

Uyeda, J. C., M. W. Pennell, E. T. Miller, R. Maia, and C. R. McClain. 2017. The evolution of energetic

scaling across the vertebrate tree of life. Am. Nat. 190:185–199.

Vacher, J.-P., P. J. R. Kok, M. T. Rodrigues, J. D. Lima, A. Lorenzini, Q. Martinez, M. Fallet, E. A.

Courtois, M. Blanc, P. Gaucher, M. Dewynter, R. Jairam, P. Ouboter, C. Thébaud, and A.

Fouquet. 2017. Cryptic diversity in Amazonian frogs: Integrative taxonomy of the genus

Anomaloglossus (Amphibia: Anura: Aromobatidae) reveals a unique case of diversification

within the Guiana Shield. Mol. Phylogenet. Evol. 112:158–173.

Vences, M., J. Kosuch, R. Boistel, C. F. B. Haddad, E. La Marca, S. Lötters, and M. Veith. 2003.

Convergent evolution of aposematic coloration in Neotropical poison frogs: a molecular

phylogenetic perspective. Org. Divers. Evol. 3:215–226.

Vinogradov, A. E. 1995. Nucleotypic effect in homeotherms: Body-mass-corrected basal metabolic rate

of mammals is related to genome size. Evolution 49:1249–1259. Wiley.

Wang, J., M. W. Itgen, H. Wang, Y. Gong, J. Jiang, J. Li, C. Sun, S. K. Sessions, and R. L. Mueller.

2021. Gigantic genomes provide empirical tests of transposable element dynamics models.

Genomics Proteomics Bioinformatics 19:123–139. Oxford University Press (OUP).

Wang, J., G. Zhang, C. Sun, L. Chang, Y. Wang, X. Yang, G. Chen, M. W. Itgen, A. Haley, J. Tang, and

768   R. L. Mueller. 2024. Size evolution of gigantic genomes suggests stochastic outcomes of

769   transposable element/host silencing interactions.

770   Whitney, K. D., and T. Garland Jr. 2010. Did genetic drift drive increases in genome complexity? PLoS

771   Genet. 6:e1001080. Public Library of Science (PLoS).

772   Womack, M. C., M. J. Metz, and K. L. Hoke. 2019. Larger genomes linked to slower development and

773   loss of late-developing traits. Am. Nat. 194:854–864. University of Chicago Press.

774   Xiol, J., E. Cora, R. Koglgruber, S. Chuma, S. Subramanian, M. Hosokawa, M. Reuter, Z. Yang, P.

775   Berninger, A. Palencia, V. Benes, J. Penninger, R. Sachidanandam, and R. S. Pillai. 2012. A role

776   for Fkbp6 and the chaperone machinery in piRNA amplification and transposon silencing. Mol.

777   Cell 47:970–979. Elsevier BV.

778   Yang, Y., M. R. Servedio, and C. L. Richards-Zawacki. 2019. Imprinting sets the stage for speciation.

779   Nature 574:99–102. Springer US.

780