Assessing above and belowground recovery from ammonium sulfate addition and wildfire in a lowland heath: mycorrhizal fungi as potential indicators

Running head: Mycorrhizas as restoration ecology assessment tools

Authors: Jill Kowal[1], Raquel Pino-Bodas[1,2], Elena Arrigoni[1], Guillaume Delhaye[1], Laura M. Suz[1], Jeffrey G. Duckett[4], Martin I. Bidartondo[1,3] and Silvia Pressel[4]

1- Royal Botanic Gardens, Kew, England

2- Universidad Rey Juan Carlos, Móstoles, Spain

3- Imperial College London, England

4- Natural History Museum, London, England

Corresponding Author: Dr. Jill Kowal, j.kowal@kew.org

Author Contributions

JK, RP, conceived the ideas and designed methodology; all authors but GD collected field and laboratory data; all authors analysed the data; JK led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

ABSTRACT:

Atmospheric pollution containing soil-nitrifying ammonium sulphate is affecting semi-natural ecosystems worldwide. Long-term additions of ammonium sulphate on nitrogen (N)-limited habitats such as heathlands increase climate stress affecting recovery from wildfires. Yet although heathland vegetation largely depends on ericoid mycorrhizal fungi (ErM) to access soil N, we lack a detailed understanding of how prolonged exposure to ammonium sulphate may alter ErM community composition and host

plants' reliance on fungal partners following wildfire and substantial reductions in ammonium sulphate pollution. Effects on ecosystem processes, particularly carbon stores, also remain uncertain. Ammonium sulphate additions occurred bi-weekly for five years after a 2006 wildfire burnt a UK heathland. Ten years after the treatments ceased (2021), we measured vegetation structure, lichen and lichen photobiont composition, soil characteristics, ErM colonisation, ErM diversity in roots and soil, and assessed their potential as recovery indicators. We found heather height and density, and moss groundcover, were greater in N-enriched plots. Lichen community indices showed significant treatment effects without photobiont differences. Soil pH and Mg, and the proportion of putative ErM fungi in soil were significantly lower in treated plots while soil cation exchange capacity was significantly higher. Increases in soil pH were positively correlated to soil ErM abundance. Soil carbon stock measures were variable and negatively related with soil ErM. Our results indicate that atmospheric pollution following fire can have significant lingering effects and mycorrhizal fungi diversity are a novel and effective ecological tool to assess ecosystem recovery on heathlands.

IMPLICATIONS FOR PRACTICE:

- Recovery of heathlands from wildfire and atmospheric pollution may require decadal scales. Conventional restoration assessment tools lack critical understanding of belowground soil and mycorrhizal fungi interactions and nutritional feedback loops with aboveground hosts.

- Given that ericoid mycorrhizal fungi (ErM) communities are key nutrient regulators of resilient heathlands, we recommend baseline measurements of ErM diversity and abundance in both soil and dominant plant roots be added to the recovery assessment toolkit prior to commencing restoration or management plans.

- What defines ErM recovery, and disentangling effects of ammonium sulphate pollution and fire, remain open questions which only long-term field experiments across pollution gradients will address.

RESEARCH ARTICLE

Introduction

Plants depend largely on their mycorrhizal fungal symbionts to access soil nitrogen (N) and phosphorus (P) particularly in nutrient-poor heathlands (Read 1996; Smith & Read 2008). Questions relating to the impact of atmospheric pollution on mycorrhizal fungal communities and their function in N- and P-limited systems such as heathlands are becoming more pressing as the ecosystem services that heathlands provide, especially regarding carbon (C) sequestration, come into focus. Particularly critical are the contributions of the dominant *Calluna vulgaris* roots and associated ericoid mycorrhizal fungi (ErM) to belowground C (Clemmensen et al. 2013) and how these may be affected by N deposition (Van Geel et. al. 2020), given the key role of ErM in plant N acquisition (Leake et al. 1990; van der Heijden et al. 2015). Nitrogen-limited heathlands across Europe are particularly vulnerable to atmospheric N deposition (Southon et al. 2013; Van Geel et al. 2020) which threatens their ecosystem function through loss of vegetation biodiversity aboveground (Tang et al. 2018; van Paassen et al. 2020) and soil changes affecting microbial communities belowground (Treseder 2008; Zhou et al. 2017; Kowal et al. 2022). Long-term N addition has been shown to alter root-associated fungal community composition, increase ErM root colonisation (Kiheri et al. 2020) or alleviate plant (heather) N deficits, thereby reducing reliance on ErM (Vesala et al. 2021). *Calluna vulgaris* age also plays a role in ecosystem function (Field et al. 2017).

European dry heathlands are reservoirs of biodiversity and provide essential ecosystem services such as C sequestration (Friggens et al. 2020; Gregg et al. 2021),

climate regulation and flood mitigation (Alonso et al. 2012). Currently, heathlands are threatened by anthropogenic pressures including encroachment by trees and agriculture plus soil acidification and eutrophication from transboundary pollutants, especially ammonium sulphate, a common N fertilizer and key component of particulate matter (PM) (EMEP 2019; Chen et al. 2018). Dry heathlands are dominated by dwarf ericaceous shrubs which associate with ErM and support diverse lichen and bryophyte communities. Whether heathlands and their belowground ErM can recover following substantial reductions in ammonium sulphate is difficult to assess on a field-scale.

Appreciation of the diversity of putative ErM colonising Ericaceae roots has expanded recently (Bougoure et al., 2007; Smith & Read 2008, Fehrer et al. 2019) to include many ascomycetes in the order Helotiales (Walker et al., 2011) and some Basidiomycota (Selosse et al. 2007; Vohnick et al. 2012). It is common for multiple ErM species to co-colonise a single host plant (Gorzelak et al. 2012) and their functionality, when combined, might become enhanced leading to increased resilience to environmental change (Perotto et al. 2002). Van Geel et al. (2020) identified 239 putative ErM operational taxonomic units (OTUs) associating with heather roots alone across 21 dry heathlands. Another recent study of ErM diversity in Norwegian heathlands found several new ErM OTUs within both Ascomycota and Basidiomycota associating with heather roots (Blaalid & Davey 2022). Further ErM diversity is being uncovered using different DNA sequence clustering methods, such as amplicon sequence variants (ASVs), which can now be linked to species hypotheses (SH) (Kõljalg et al. 2020; Põlme et al. 2020; Abarenkov et al. 2021).

Lichens are sensitive to air quality and have long been used as bioindicators (Nash 1996; Nimis et al. 2002; Giordani 2007). However, the effects of N deposition on terricolous lichen communities, typical of heathlands, have been largely ignored with only a few studies to date showing that N deposition negatively affected some heathland lichen species (Pilkington et al. 2007; Stevens et al. 2012; Edmondson et al. 2013). This represents a considerable knowledge gap since terricolous lichens provide important ecosystem services including soil stabilization, soil moisture retention and facilitation of seed germination (Asplund & Wardle 2017). Moreover, few studies have investigated the recovery of lichen communities once disturbance ceased (Edmondson et al. 2013; Benvenutto-Vargas & Ochoa-Hueso 2020). While variation in the composition of lichen communities under atmospheric pollution and different climatic gradients is well documented (Gadsdon et al. 2010), little is known about potential changes in the interactions between mycobiont and photobiont; this is surprising since adaptation of lichens to different environmental conditions is highly dependent on their photobiont (Peksa & Škaloud 2011). The ability of mycobionts to establish associations with several lineages of photobionts is considered an adaptive strategy to tolerate a wider range of environmental conditions (Fernandez-Mendoza et al. 2011; Jüriado et al. 2019). Recent studies show that photobionts switch along environmental gradients (Dal Grande et al. 2018); however, the effect of N deposition on mycobiont-photobiont interactions has not been investigated.

In this study we examined a subset of the experimental plots used by Southon et al. (2012) to determine whether there are lingering effects on ecosystem function more than ten years since the last ammonium sulphate addition and wildfire (for a description of the Southon et al.'s experiments, see Background Data S1, Table S1). Specifically, we

asked whether differences between the N-treated and control plots were still detectable (1) aboveground in the structure or diversity of plant and lichen communities and associated photobionts, and (2) belowground in soil C stocks and mycorrhizal and soil fungal community composition. We propose that the slow growth of the dominant heather shrubs renders this habitat resistant to full recovery of vegetation structure and lichen community composition from ammonium sulphate addition. However, we expect recovery in the soil chemistry, soil fungi and mycorrhizal community composition given more than 10 years of precipitation and background pollution to diffuse treatment effects. Finally, we present baseline molecular data on the ErM fungal community to assess its potential as a heathland recovery metric.

Methods

Experimental design and site

The study was carried out in June 2021 at Thursley NNR in southern England (51°09'25" N 0°42'04" W). The site comprises H2 dry heathland (National Vegetation Classification (NVC) scheme, Rodwell 2006) and is designated a Special Area of Conservation (European Union's Habitats Directive, 92/43/EEC). The soil is a podzol derived from Lower Greensand with a thin (<1 cm) humus layer overlying a sandy mineral layer (Power et al., 1998). In 2021, the mean annual precipitation was 781 mm and mean annual temperature 11.9° C (minimum 6° C in January and maximum 20° C in August) (Fig. S1). Background N and sulphur (S) deposition levels were 14 kg ha^{-1} year^{-1} and 12.7 kg ha^{-1} year^{-1} in 2020 (www.worldweatheronline.com, Levy et al., 2020), higher than the N levels measured at the site (8 kg ha^{-1} year^{-1}) from 1998 and 2000 (Power & Barker 2003); S alone was not measured.

We studied three of the four blocks assessed in Southon et al. (2012) (Fig. S2a, b, c); each block contains four 4 m^2 plots, each with four 2 m^2 subplots. These three blocks were selected as they were the only blocks with the same number of treatments; two control plots and two treated plots which received bi-weekly additions of ammonium sulphate at a rate of 30 kg^{-1} ha (dissolved and distributed using a fine nozzle spray) during 1998 through 2010. This rate approximated levels reported in Europe, e.g., the Netherlands, during this period (Bobbink et al. 2013).

Vegetation and lichen survey

We carried out a vegetation and lichen survey of the three blocks. For each subplot we measured vegetation density, abundance and height, and recorded plant and lichen species composition. Following the lichen survey, three specimens of each lichen species were collected to confirm the identifications and carry out DNA identification of photobionts in the laboratory (Methods S1). Nomenclature for bryophytes follows Blockeel et al. (2021) and for lichens Burgaz et al. (2020) and Pino-Bodas et al. (2021).

Soil sampling

To assess soil characteristics and chemistry, we collected two soil samples per plot to a depth of 15 cm. Each sample comprised three subsamples which combined gave at least 500 g dry weight (DW). Soil samples were stored at 4°C and processed by NRM Laboratories (Bracknell, England, Methods S2). We measured the following: water content, bulk density, dry matter, pH, total C and N concentrations (% g/100g, on a dry matter basis); N, P, potassium (K), magnesium (Mg), nitrate (NO_3), ammonium (NH_4^+), sodium (Na), calcium (Ca), all (mg/kg); cation exchange capacity (CEC) and C stock (t/ha).

Approximately 50 g of fresh soil was removed from each soil sample (taking care to remove at least 15 g from each subsample) and placed at -80°C for environmental DNA analysis (see below).

Plant leaves, roots and stems analyses

Calluna vulgaris leaf tissue (50 g DW) was collected from at least three randomly selected plants from each plot (n=12) to measure foliar N, P and C. Immediately after collection, samples were dried (overnight at 60°C) and processed by NRM Laboratories (Bracknell, England; see Methods S2).

For ErM fungal root colonisation and community assessments, three young plants of *C. vulgaris* were collected. For each of their roots, one half was stored in ethanol (70% v/v) at 4°C for ink staining to assess ErM root colonisation, the other half was stored in cetyltrimethylammonium bromide (CTAB) extraction buffer at -20°C for molecular identification. Ericoid mycorrhizal colonisation was scored as present only when typical ErM hyphal coils were visible within the cell walls of at least three root cells per root section (Kowal et al. 2020a, b).

To assess lateral stem seasonal growth, we selected five mature plants of *C. vulgaris* per plot and removed the largest stem at the plant's base. The stem cuttings were kept at 4°C for further processing and analysis. Stem cuttings were hand sectioned (approximately 0.25 mm thick) and annual growth ring widths were measured (x 3 per section) on both the widest and narrowest radial hemispheres (x40) under a Zeiss Axioscop 2 microscope equipped with an AxioCam MRc digital camera and auto-calibrated using Axiovision Microscope Software (Fig. S3). The results were tabulated and aligned to compare years of treatment and the years post-treatment.

Soil fungal community analysis

Total soil DNA was extracted in duplicate from 0.25 g of soil from each plot (12 plots, 24 extractions, DNeasy PowerSoil Pro Kit, Qiagen, Hilden, Germany). DNA extraction quality (260/280nm and 260/230nm ratios) was assessed using a NanoDrop2000 spectrophotometer (Thermo Fisher Scientific) and DNA concentration was quantified using a Quantus™ Fluorometer and QuantiFluor® dsDNA System (Promega, Madison, Wisconsin, United States). The DNA extractions were processed by Macrogen (Seoul, South Korea) for library preparation and Illumina sequencing with the primer set ITS86F and ITS4 (White et al. 1990; Turenne et al. 1999) targeting the ITS2 region.

Root fungal community analysis

To identify the ErM fungal communities colonising *C. vulgaris* roots, we pooled ten roots from each of the heather plants sampled per plot (30 roots) and carried out three DNA extractions (DNeasy PowerSoil Pro Kit, Qiagen, Hilden, Germany) for each plot (n = 12 extractions per block, 36 extractions overall) followed by amplification, cloning and sequencing of the complete ITS region (Methods S3). An ABI3730 genetic analyser (Applied Biosystems) was used for Sanger DNA sequencing, and forward and reverse sequences were assembled using Geneious v. 8.1.9. Partial SSU and LSU regions were trimmed using ITSx (Bengtsson-Palme et al. 2013) and the resulting full-length ITS sequences were clustered into Amplicon Sequence Variants (ASVs) in VSEARCH (Rognes et al., 2016). Amplicon Sequence Variants forming new Species Hypothesis (SHs) were excluded from further analyses as their taxonomic classification remains poorly resolved.

For both roots and soil, taxonomic classification was performed using the SH matching v2.0.0 tool on the PlutoF workbench (Abarenkov et al. 2010). Those assigned to the same SH were grouped and categorised by putative ecological function (mycorrhiza, ericoid associate or other) (Vohnik et al. 2012, 2016; Grelet et al. 2017; Van Geel et al. 2020). For statistical analysis, we only analysed taxa (OTUs) represented by > 100 reads. The overall proportion of ErM and non-ErM fungi was calculated as the total ErM and non-ErM sequences over the total number of sequences that passed the quality control in the SH matching analysis.

To compare our baseline analysis with a recent European heathland survey of root-derived ErM diversity, we grouped the OTUs reported in Van Geel et al. (2020) into SHs by filtering only OTUs derived from *C. vulgaris* roots in heathlands and assigning a putative ecological function to the SH taxonomy (Methods S3; see Methods S4 for the analysis of lichen photobiont DNA).

Statistical analyses

Statistical analyses were conducted in R (R core Team 2022, v. 4.1.2) unless otherwise stated. Soil characteristics, leaf chemistry, aboveground biomass measures, lichen richness, abundance and diversity (Shannon and inverse Simpson indexes) were compared between treatments using linear mixed effects models, using block and replicates as random variables (function brm in the *brms* package). Shannon and inverse Simpson indexes were calculated using BiodiversityR (Kindt & Kindt 2019). Wilcoxon rank sum tests were applied with continuity correction. The effect of the treatment on the composition of total fungal and ErM fungal root and soil communities was tested using a nested PERMANOVA (nperm = 999) with block as a random variable on a Bray-

Curtis dissimilarity matrix, using the *adonis2* function in the *vegan* package (Oksanen et al. 2013). The predicted values for both soil ErM richness and ErM colonisation rates were tested using soil covariates in simple linear regression models.

Effect of treatment on ErM colonisation presence or absence in heather roots was also tested using chi-square tests with Yates correction using Graphpad Prism (V 9.5 525). Correlation matrices were used to test ErM root colonisation and root and soil richness with soil chemical characteristics (Prism V 9.5 525).

To test significant differences between lichen community composition in control and N-addition subplots PERMANOVA (nperm = 999) was applied in *vegan*. The Bray-Curtis dissimilarity between communities was represented graphically by non-metric multidimensional scaling (NMDS) ordination. Indicspecies (De Cáceres & Jansen 2015) was used to identify lichen indicator species for each treatment, performed with *multpatt* (999 permutations).

Results

Vegetation and lichen structure, composition, abundance and diversity

The results of the vegetation and lichen surveys are summarized in Tables S2a, b, c. *Calluna vulgaris* dominated the shrub community and was present in every subplot. *Ulex europaeus and Erica tetralix* were present occasionally and with significantly less coverage with 112 and 24 grid points, respectively, compared with 2,429 *C. vulgaris* points. The ground layer consisted of either leaf litter, bryophytes (the mosses *Hypnum jutlandicum*, *Campylopous introflexus* and *Dicranum scoparium* and the leafy liverwort *Cephaloziella divaricata*) or a combination of the two along with 21 species of lichens,

all belonging to the genus *Cladonia*. The most frequent *Cladonia* species were *C. furcata*, *C. crispata* subsp. *cetrariiformis*, *C. portentosa* and *C. verticillata.*

The mean height and tallest points of the heather canopy were both significantly higher in the N-addition plots than the control plots (ANOVA, *P* value <0.05) but no treatment differences were detected in the vegetation density at either ground level or 20cm (Fig. 1).

Campylopus introflexus ground cover was significantly higher in the N-addition plots compared with the control plots (Table S3; Fig. 2D) but no treatment effects were seen in the other bryophyte species*.* Similarly, we found no differences in heather lateral stem growth between the two treatment plots when measuring growth rings in the immediate years following the fire (2006), during the years of further N-addition (2006-2011), or after N additions ceased (2012 to 2021 harvest) (Table S4). However, the fixed effect of year shows significant differences in growth patterns (*P* value: < 0.0001) with heather growth more robust in the first three years (average 0.36 mm/year) and slower in the last three years (0.16 mm/year). Notably, we observed a growth peak in the N-addition plots around 2014, two years after treatments ceased (Fig. S4).

Control plots had significantly higher abundance of lichens (all *Cladonia* spp., P-value=0.014), while the N-addition plots had significantly higher Simpson and Shannon diversity values (P-value = 0.0002, P-value = 0.005, respectively; Fig. 2; Table S5a). No significant differences in richness were found between treatments (P-value = 0.633).

PERMANOVA analysis showed significant differences in the lichen communities between treatments (R2 = 0.06, F = 3.0074, *P* value = 0.004, Fig. S5). The species

indicator analysis found significant differences for the presence and/or abundance of four species (Table S5b; Fig. S6). *Cladonia crispata (P* value = 0.024), *C. furcata* (*P* value = 0.002) and *C. gracilis* (*P* value = 0.018) were associated with the control plots while *C. fimbriata* (*P* value = 0.002) was associated with N-addition plots. Differences in presence and/or abundance in other *Cladonia* spp. (Table S6), and the lichen crust cover recorded in Southon et al. (2012) were no longer apparent.

Soil characteristics and leaf chemistry

Soil chemistry analyses revealed significant differences in several variables between treatments; soil pH and Mg were significantly lower in the ammonium sulphate-addition plots (*P* value < 0.01, < 0.05), while CEC was higher (*P* value < 0.01)(Table S7).

The Principal Component Analysis biplot indicates that soil pH, Mg and CEC explain most soil characteristic differences between control and N-addition plots and that Mg strongly correlates with pH (Fig. 3A). The group of soil characteristics are also clustered by control and N-addition, further supporting a difference due to treatment (Fig. 3B). Approximately 47% of the variance is explained in the first (31.3%) and second (16.2%) principal components.

Carbon stock in the control plot samples ranged from 13 to 40 C t/ha while the ammonium sulphate-addition plots ranged from 16 to 48 C t/ha but this was not significantly different. Foliar P, N and C were not significantly different between the treatment plots (Table S8).

Root ErM fungal colonisation, richness and diversity

ErM fungal colonisation

All *Calluna vulgaris* roots analysed showed some ErM colonisation. The analysis of treatment impact on proportion of individual roots colonised, by block and plot, was not significant (Fig. 4A). However, when considering the proportion of overall root cells colonized in all plots combined, the N-addition plots had significantly lower colonisation than control plots (Fig. 4B, Table S9).

Root fungal richness and diversity

A total of 352 fungal full ITS sequences out of the 368 sequences identified passed the chimera detection and quality control steps. These were clustered into 218 ASVs yielding 23 unique SHs. From these, 56 were assigned as putative ErM ASVs matching to seven unique ErM SHs (Table S10). Three of the seven unique ErM SHs were present in the roots and soil (this study) and sequences from Van Geel et al. (2020) analysed for comparison in this study.

Taxonomic classifications for the following order and families were assigned as: (1) ErM - Chaetothyriales, Helotiales, *Hyaloscypha/Pezoloma*, *Hymenoscyphus, Meliniomyces; Oidiodendron*, Sebacinales and Serendipitaceae (Sebacinaceae isolates however have not been resynthesized in culture, Allen et al. (2003)); and (2) ericoid associates - Mycenaceae and Hymenochaetales. Although their mycorrhizal status remains uncertain, some fungi are recognized as frequent associates of Ericaceae roots and are therefore differentiated herein as 'ericoid associates'.

The ErM fungi represented ~ 19% and 29% of sequences for the control and N-addition plots, respectively. The different taxonomic breakdown of putative ErM and non-ErM fungi can be seen in Fig. 5A, C.

Among the first five dominant SHs were the ericoid mycorrhizal genera *Hyaloscypha/Pezoloma* (SH1156094.08FU, 22 ASVs) and *Oidiodendron* (SH1185398.08FU, 11 ASVs), which represented 6.4%/7.7%, and 7%/12.1% of the relative abundance in control/N-addition plots, respectively (Table S9). Our SH re-analysis with the 239 putative ErM OTUs previously reported in *C. vulgaris* across 21 European heathlands (van Geel et al. 2020) yielded 82 unique ErM SHs (Table S11).

There were no significant differences in community composition between treatments when comparing all root fungi or the ErM only (Fig. 6A, B).

Soil fungal richness and diversity

There were significant differences in the overall soil fungi community composition between control and N-addition plots (Fig. 6C). Over 53% of all soil fungi reads were putative ErM which clustered into 1,209 OTUs (97% similarity) matching to 570 unique SHs. Of these, 40 were putative ErM SHs (Table S11). When comparing the soil ErM fungi only, the effect of treatment was not significant (Fig. 6D).

Ericoid mycorrhizal richness, soil chemistry covariates

The overall regression model combining soil pH and soil C stock predicts soil ErM SH richness (adjusted R^2 = 0.756, $F_{(2,9)}$ = 18.05, $p < 0.001$). Individually each variable significantly predicts soil ErM SH richness (b = 291,123, p = 0.002; b = -3101, p = 0.015, respectively).

Pearson r correlation coefficients between ErM root colonization and ErM soil SH with soil and leaf chemical parameters are reported in Fig. S7.

Diversity of photobionts associated with Cladonia

A total of 214 ITS rDNA photobiont sequences associated with the genus *Cladonia* were generated; all belonged to *Asterochloris italiana* (Fig. S8). Six haplotypes were identified and connected in a single network. One haplotype was restricted to N-addition plots, three to control plots and the other two haplotypes were present in both treatment plots (Fig. S9). The most abundant haplotype was present in 96.2% of all the specimens and equally represented on plots of both treatments. The genetic diversity of photobionts on control plots was higher than in N-addition plots (Table S12); however, the AMOVA showed that most photobiont variation was within subplots (Table S13).

Discussion

Our results provide novel insights on the long-term effects of ammonium sulphate addition to European lowland heathlands and highlight the hitherto unappreciated potential of mycorrhizal fungi as bioindicators of heathland recovery. We show that ammonium sulphate addition continues to exert an effect on heathland vegetation structure and ecosystem function more than ten years since treatments ceased and 15 years since a wildfire. Marked differences between treated and control plots are apparent in plant biomass and lichen community composition and diversity, soil characteristics and overall soil fungi community composition. As expected, heather height, both in terms of mean height and highest canopy points, continues to show a N effect, indicating that N-addition has a long-term effect on dominant heathland vegetation structure. However, because all the study heather plots were reaching life-cycle maturity when the plots burned, making new seedling recruitment unlikely (Schellenberg & Bergmeier 2022) it is possible that plant biomass responses and heather

regeneration may have been different if the heather stands had burned at an earlier plant life-history stage (Meyer-Grünefeldt et al. 2015). We expected to see comparable growth trends in the stem ring diameters given N additions induce heather canopy growth (Aerts 1989), instead we found that heather stem expansion was insensitive to treatment and that growth regulation was dominated by the terminal stems. Stem thickness and any effects from N addition may have already been reached before the 2006 fire.

In line with previous findings (Terry et al. 2004), our survey also revealed significant differences in lichen community composition more than ten years after ceasing N addition. Thus is likely due to slow growth of lichens (Hale 1959; Karlén & Black 2002) requiring decadal recovery periods.

When considering lichen photobiont diversity, although the number of haplotypes found in the control plots was higher than in the N-addition plots, the two most frequent haplotypes were equally present across treatments, all belonging to *Asterochloris italiana*, a photobiont commonly associated with *Cladonia* species in Europe (Pino-Bodas & Stenroos 2021) and able to tolerate pH values of around 4 (Škvorová al. 2022). Recent studies identified differences in soil chemistry, including pH, N and C content, as major factors driving photobiont switching (Škvorová et al. 2022). However, we found no clear evidence of photobiont switching between our treatment groups, perhaps indicating that treatment levels and soil pH differences were insufficient to elicit such a response.

Turning to bryophyte diversity, N additions are known to promote the invasive moss *Campylopus introflexus* (Sparrius & Kooijman 2013; Caporn et al. 2014; Sérgio et al. 2018), and indeed we recorded significantly higher ground cover of this species in the

ammonium sulphate-addition plots. This contrasts with Southon et al.'s (2012) results

showing significant differences in species abundance of the moss *Hypnum jutlandicum*

and the leafy liverwort *Cephaloziella divaricate* but not of *C. introflexus*. The most likely

explanation for this difference is the presence in their study, but not ours, of several

other competing pioneer moss species (e.g., *Ceratodon purpureus*, *Pohlia nutans*,

Polytrichum juniperinum). Post-fire bryophyte succession on a typical NVC H2 heathland

predominantly includes a mosaic of *C. introflexus, C. pyriformis, Ceratodon purpureus,*

Marchantia polymorpha ssp. *ruderalis, Pohlia nutans* and *Polytrichum* spp., progressing

to dominant *Campylopus* spp. and *Hypnum jutlundicum* by year 20 (Duckett et al. 2008;

Burch 2009; Pressel et al. 2021). This is consistent with our observations that only *C.*

introflexus and *H. jutlundicum* were present in the experimental plots. Southon et al.

(2012) noted that *Cephaloziella divaricata* was significantly more abundant in the

control plots. While we did observe *C. divaricata* in many of the plots, we excluded this

species from our analyses because it is miniscule and only detectable by close

examination with a hand lens, making it impractical in the context of this study. Given

the potential of leafy liverworts such as *Cephaloziella* as sources of mycorrhizal inocula

to improve the re-establishment of vascular plants following disturbance (Kowal et al.

2016), further studies are needed to determine the long-term effects of N-deposition

on this genus.

Surprisingly, observed aboveground differences in bryophyte abundance did not

correspond to differences in current soil nitrate or ammonium concentrations. It is likely

that given the sandy nature of heathland soils, ammonium and nitrate have been

leached, but lingering effects on soil pH remained, since we found significant differences

in soil pH values between our control and ammonium sulphate plots, in line with

previous reports (Kadyampakeni et al. 2018). The significant differences documented in this study in soil fungal community composition between treatments, but equivocally in mycorrhizal fungal colonisation of *Calluna* roots, further highlight that more experiments are needed to guide policy makers on which components and levels of pollution can lead to fundamental changes in mycorrhizas and C stocks. Differences in pH were the most important predictor of ErM abundance in soil. While N deposition has been linked to lower soil pH, further direct experimental measurements are needed to test this hypothesis in heathland. Southon et al. (2012) reported increased microbial biomass in N-addition plots possibly reflecting greater presence of bacteria and non-mycorrhizal fungi. In this light, our regression results predicting higher ErM abundance when the C stock is lower, are intriguing as they may indicate subtle feedback cycles not captured in our analysis, possibly related to the leaching properties of sandy dry heathland soil. A more plausible explanation is the inherent limitations imposed by the mature age of the canopy which reduces C sequestration potential (Field et al. 2017). Thus, the age of the heather host and relative contribution to soil C stock may yet be another important covariate not measurable in the current study design. Variation in soil C stock across plots may also be explained by uneven distribution of sandy soil and soil moisture (Li et al. 2022), which we found strongly and positively correlated with C stock.

Differences in soil pH can also directly lead to changes in community composition of bryophytes and terricolous lichens (Rai & Gupta 2022). However, all terricolous lichens recorded in our survey belonged to the generally heliophilous genus *Cladonia* (Ahti 2000). Indicating that the most important factor influencing differences in *Cladonia* species richness between the control and ammonium sulphate-addition plots

was the density of the heather canopy, as measured by Southon. The indicator species of the ammonium sulphate-addition plots was *C. fimbriata*, which was mainly present at the base of heather near ground level. This species is common in a wide diversity of forests (Burgaz et al. 2020) and has been associated with disturbed sites in the UK (Pino-Bodas et al. 2021). Indicator species in the control plots included *C. crispata*, a characteristic species of mature heathlands (Coppins & Shimwell 1971; Sanderson 2017) and *C. furcata*, a species growing in a wide range of habitats, also frequent in heathlands.

Our molecular analyses of heather root-derived fungal DNA detected 40 unique SHs, of which seven belonged to putative ErM fungi, compared to 82 SHs in heather across 21 dry European heathlands including nine in England (van Geel et al. 2020). Van Geel et al.'s sampling effort i.e., 10 roots from three plants (for each of the three 1 x 1 m plots per site) was similar to that in this study. Six out of the seven ErM fungi found in the roots and 24 out of the 40 ErM fungi detected in soil at Thursley were also identified in their study. Thus, the diversity of ErM hosted by heather at Thursley yielded only one new SH not previously detected at the regional scale.

Our results showing a weak positive relationship between ErM colonisation and P concentration in heather leaves reflect slower growth rates in the aging experimental heather stands alongside less heather regeneration (Calvo-Fernández, 2018). We also found that soil ammonium was negatively correlated to soil putative ErM taxa, but are unable to infer whether this is a consequence of the long-term ammonium sulphate addition experiment since we found no measurable link between soil N and ErM colonisation of *Calluna* roots. ErM colonisation may have recovered to pre-treatment colonisation levels or perhaps was never affected by the treatment. Nevertheless, our observed differences in soil pH and soil fungal community composition between

treatments however indicate that treatment effect is likely in this case and is consistent with Van Geel et al. (2020). It is worth noting that N deposition has been previously shown to increase P acquisition (and K) in heather (Rowe et al. 2008), thus it is possible that the weak positive correlation between ErM colonisation and foliar P is a lingering effect of past N additions and that our sample size was simply too small to detect a significant difference between treatments. That soil C stocks in both the Scottish and this study (at least in the top 15 cm of soil) showed no significant differences between N treated and control plots is probably attributable to considerable leaching in this habitat, as discussed before. This, as well as soil pH should be tested in further experiments as explanatory variables for ErM colonisation, diversity and functionality. It is also possible that cumulative historical background N deposition had already impacted the heathland belowground to a level whereby it is difficult to detect strong soil carbon or mycorrhizal effects, despite adding massive amounts of ammonium sulphate and then stopping. Thus, a belowground tipping point may have been crossed long ago whereby the control is largely indistinguishable from treatment.

Despite heathlands harbouring significant biomass belowground, there is a scarcity of experiments testing the effects of atmospheric pollution on these habitats belowground. The fact that differences in vegetation biomass, lichen composition, soil chemistry and mycorrhizal fungal community recorded here can be attributed to the historical ammonium sulphate applications is notable given that mean annual precipitation and temperatures at Thursley have fluctuated dramatically since 2009 (Fig. S1). This suggests that heathlands may be an ecosystem resilient to climate change but more sensitive to deposition of nutrients. Our study thus provides a baseline for

monitoring the above and belowground effects of both climate change and nutrient inputs on lowland heathlands.

Predicting interactions between atmospheric pollution, ErM community composition and C sequestration is particularly complex as ErM can be free-living saprotrophs, and their function is context-dependent and strongly related to ErM species diversity (Ward et al. 2022). . In summary, the feasibility of developing objective indices based on ErM abundance and community composition offers a promising new ecological indicator for comparing atmospheric pollution effects and recovery across heathlands in addition to aboveground lichen and vegetation structure and phenology.

References

Abarenkov K, Zirk A, Piirmann T, Pöhönen R, Ivanov F, Nilsson RH, Kõljalg U (2021) Full UNITE+INSD dataset for Fungi. Version 10.05.2021. UNITE Community

Abarenkov K, Tedersoo L, Nilsson RH, Vellak K, Saar I, Veldre V, Parmasto E, Prous M, Aan A, Ots M, Kurina O, Ostonen I, Jõgeva J, Halapuu S, Põodmaa K, Toots M, Truu J, Larsson K-H, Kõljalg U (2010) PlutoF—a web-based workbench for ecological and taxonomic research, with an online implementation for fungal ITS sequences. *Evolutionary Bioinformatics* 6:EBO-S6271

Aerts R (1989) Above-ground biomass and nutrient dynamics of *Calluna vulgaris* and *Molinia caerulea* in a dry heathland *Oikos* 56:31-38

Ahti, T (2000) Cladoniaceae. *Fl Neotrop Monogr* 78:1-363

Allen TR, Millar T, Berch SM, Berbee ML (2003) Culturing and direct DNA extraction find different fungi from the same ericoid mycorrhizal roots. *New Phytologist* 160:255-272

Alonso I, Weston K, Gregg R, Morecroft M (2012) Carbon storage by habitat – Review of the evidence of the impacts of management decisions and condition on carbon stores and sources. Natural England Research Reports, Number NERR043

Asplund J & Wardle DA (2017) How lichens impact on terrestrial community and ecosystem properties. *Biological Reviews* 92:1720-1738

Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, De Wit P, Sánchez-García M, Ebersberger I, de Sousa F, Amend A, Jumpponen A, Unterseher M, Kristiansson E, Abarenkov K, Bertrand YJK., Sanli K, Eriksson KM, Vik U, Veldre V, Nilsson RH (2013) Improved software detection and extraction of ITS1 and ITS2 from ribosomal

ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. *Methods in Ecology and Evolution* 4:914-919

Benvenutto-Vargas VP & Ochoa-Hueso R (2020) Effects of nitrogen deposition on the spatial pattern of biocrusts and soil microbial activity in a semi-arid Mediterranean shrubland. *Functional Ecology* 34:923-937

Blaalid R & Davey ML (2022) Habitat protection approaches facilitate conservation of overlooked fungal diversity – A case study from the Norwegian coastal heathland system. *Frontiers in Fungal Biology* 3: 886385

Blockeel T, Hodgetts NG, Pilkington SL, Pescott OL (2021) A Census Catalogue of British and Irish Bryophytes. British Bryological Society. ISBN: 978-0-9561310-4-1

Bougoure DS, Parkin PI, Cairney JW, Alexander IJ, Anderson IC (2007) Diversity of fungi in hair roots of Ericaceae varies along a vegetation gradient. *Molecular Ecology* 16:4624-36

Bobbink R, Bal D, van Dobben HF, Jansen A JM, Nijssen M, Siepel H, Schaminée JHJ, Smits N AC, de Vries W (2013) Part 2 The effects of nitrogen deposition on the structure and functioning of ecosystems. Available from: https://ec.europa.eu/environment/nature/natura2000/platform/documents/part-i-chapter_2_nov-2012_2013-09-10_en.pdf [accessed online December 2022]

Burgaz AR, Ahti T, Pino-Bodas R (2020) Mediterranean Cladoniaceae. Sociedad Española de Liquenología, Madrid. ISBN: 978-84-09-21610-9

Burch J (2009) The regeneration of bryophytes after the burning of dry heath and wet heath moorland on the North York Moors. Natural England Commissioned Report NECR011

Calvo-Fernández J, Taboada Á, Fichtner A, Härdtle W, Calvo L, Marcos E (2018) Time-and age-related effects of experimentally simulated nitrogen deposition on the functioning of montane heathland ecosystems. *Science of the Total Environment* 613:149-159

Cape JN, van der Eerden LJ, Sheppard LJ, Leith ID, Sutton MA (2009) Evidence for changing the critical level for ammonia. *Environmental Pollution* 157:1033-1037

Caporn SJ, Carroll JA, Dise NB, Payne RJ (2014) Impacts and indicators of nitrogen deposition in moorlands: Results from a national pollution gradient study. *Ecological indicators* 45:227-234

Chen S, Zhao Y, Zhang R (2018) Formation Mechanism of Atmospheric Ammonium Bisulfate: Hydrogen-Bond-Promoted Nearly Barrierless Reactions of SO_3 with NH_3 and H_2O. *Chemphyschem* 19:967-972

Clemmensen KE, Bahr A, Ovaskainen O, Dahlberg A, Ekblad A, Wallander H, Stenlid J, Finlay RD, Wardle DA, Lindahl BD (2013) Roots and associated fungi drive long-term carbon sequestration in boreal forest. *Science* 339:1615-1618

Coppins B & Shimwell D (1971) Cryptogam complement and biomass in dry *Calluna* heath of different ages. *OIKOS* 22:204-209

Dal Grande F, Rolshausen G, Divakar PK, Crespo A, Otte J, Schleuning M, Schmitt I (2018) Environment and host identity structure communities of green algal symbionts in lichens. *New Phytologist* 217:277-289

De Cáceres M & Jansen F (2015) Indicspecies: Relationship between species and groups of sites. R package version 1.7. 6

Duckett JG, Matcham HW, Pressel S (2008) Thursley Common National Nature Reserve: bryophyte recolonisation one year after the great fire of July 2006. *Field Bryology* 94:3-11

Edmondson J, Terribile E, Carroll JA, Price EAC, Caporn SJM (2013) The legacy of nitrogen pollution in heather moorlands: ecosystem response to simulated decline in nitrogen deposition over seven years. *Science of the Total Environment* 444:138-144

European Monitoring and Evaluation Programme (EMEP) updated 2019. https://www.emep.int/mscw/pollutants.html#acid [accessed 21 April 2022]

Field CD, Evans CD, Dise NB, Hall JR, Caporn SJM (2017) Long-term nitrogen deposition increases heathland carbon sequestration. *Science of the Total Environment* 592:426-435

Fehrer J, Réblová M, Bambasová V, Vohník M (2019) The root-symbiotic *Rhizoscyphus ericae* aggregate and *Hyaloscypha* (Leotiomycetes) are congeneric: Phylogenetic and experimental evidence. *Studies in Mycology* 92:195-225

Fernandez-Mendoza F, Domaschke S, Garcıa MA, Jordan P, Martın MP, Printzen C (2011) Population structure of mycobionts and photobionts of the widespread lichen *Cetraria aculeata*. *Molecular Ecology* 20:1208-1232

Friggens NL, Hester AJ, Mitchell RJ, Parker TC, Subke J-A, Wookey PA (2020) Tree planting in organic soils does not result in net carbon sequestration on decadal timescales. *Global Change Biology* 26:5178-5188

Gadsdon SR, Dagley JR, Wolseley PA, Power SA (2010) Relationships between lichen community composition and concentrations of NO_2 and NH_3. *Environmental Pollution* 158:2553-2560

Giordani P (2007) Is the diversity of epiphytic lichens a reliable indicator of air pollution? A case study from Italy. *Environmental Pollution* 146:317-323

Gregg R, Elias JL, Alonso I, Crosher IE, Muto P, Morecroft MD (2021) Carbon storage and sequestration by habitat: a review of the evidence (second edition) Natural England Research Report NERR094. Natural England, York

Grelet GA, Ba R, Goeke DF, Houliston GJ, Taylor AF, Durall DM (2017) A plant growth-promoting symbiosis between *Mycena galopus* and *Vaccinium corymbosum* seedlings. *Mycorrhiza* 27:831-839

Gorzelak MA, Hambleton S, Massicotte HB (2012) Community structure of ericoid mycorrhizas and root-associated fungi of *Vaccinium membranaceum* across an elevation gradient in the Canadian Rocky Mountains. *Fungal Ecology* 5:36-45

Hale JME (1959) Studies on lichen growth rate and succession. *Bulletin of the Torrey Botanical Club* 66:126-129

Jüriado I, Kaasalainen U, Jylhä M, Rikkinen J (2019) Relationships between mycobiont identity, photobiont specificity and ecological preferences in the lichen genus *Peltigera* (Ascomycota) in Estonia (northeastern Europe). *Fungal Ecology* 39:45-54

Kadyampakeni DM, Nkedi-Kizza P, LeivaJ A, Muwamba Fletcher E, Morgan KT (2018) Ammonium and nitrate transport during saturated and unsaturated water flow through sandy soils. *Journal of Plant Nutrition and Soil Science* 181:198-210

Karlén W & Black JL (2002) Estimates of lichen growth-rate in northern Sweden. *Geografiska Annaler: Series A, Physical Geography* 84:225-232

Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20:1160-1166

Kiheri H, Velmala S, Pennanen T, Timonen S, Sietiö O, Fritze H, Heinonsalo J, van Dijk N, Dise N, Larmola T (2020) Fungal colonisation patterns and enzymatic activities of peatland ericaceous plants following long-term nutrient addition. *Soil Biology and Biochemistry* 147:107833

Kindt R & Kindt MR (2019) Package 'BiodiversityR'. *Package for community ecology and suitability analysis* 2:11-12

Kosecka M, Guzow-Krzemińska B, Černajová I, Škaloud P, Jabłońska A, Kukwa M (2021) New lineages of photobionts in Bolivian lichens expand our knowledge on habitat preferences and distribution of *Asterochloris* algae. *Scientific Reports* 11:1-12

Kõljalg U, Nilsson HR, Schigel D, Tedersoo L, Larsson K-H, May TW, Taylor AFS, Jeppesen TS, Frøslev TG, Lindahl BD, Põldmaa K, Saar I, Suija A; Savchenko A, Yatsiuk I, Adojaan K, Ivanov F, Piirmann T, Pöhönen R, Zirk A, Abarenkov K (2020) The taxon hypothesis paradigm—On the unambiguous detection and communication of taxa. *Microorganisms* 8:1910

Kowal J, Pressel S, Duckett JG, Bidartondo, MI (2016) Liverworts to the rescue: an investigation of their efficacy as mycorrhizal inoculum for vascular plants. *Functional Ecology* 30:1014-1023

Kowal J, Arrigoni E, Serra J, Bidartondo M (2020ª) Prevalence and phenology of fine root endophyte colonization across populations of *Lycopodiella inundata*. *Mycorrhiza* 30:577-587.

Kowal J, Arrigoni E, Lane S (2020b). Acidified Blue Ink-staining Procedure for the Observation of Fungal Structures Inside Roots of Two Disparate Plant Lineages. *Bio-protocol* 10:21769

Kowal J, Arrigoni E, Jarvis S, Zappala S, Forbes E, Bidartondo MI, Suz LM (2022) Atmospheric pollution, soil nutrients and climate effects on Mucoromycota arbuscular mycorrhizal fungi. *Environmental Microbiology* 24:3390-3404

Leake JR, Shaw G, Read DJ (1990) The role of ericoid mycorrhizas in the ecology of Ericaceous plants. *Agriculture Ecosystems & Environment* 29:237-250

Levy PE, Smith RI, Dore AJ, Tang YS, Stedman JR (2020) Sulphur and nitrogen atmospheric Concentration Based Estimated Deposition (CBED) data for the UK 2014-2016. NERC Environmental Information Data Centre. (Dataset)

Li Q, Ambus PL, Michelsen A, Kapper Schmidt I, Beier C, Dietzen CA, Reinsch S, Frost Arndal M, Steenberg Larsen K (2022) Re-visiting soil carbon and nitrogen stocks in a temperate heathland seven years after the termination of free air CO_2 enrichment (FACE), *Geoderma* 428:116185.

Meyer-Grünefeldt M, Calvo L, Marcos E, von Oheimb G, Härdtle W (2015) Impacts of drought and nitrogen addition on *Calluna* heathlands differ with plant life-history stage. *Journal of Ecology* 103:1141-1152

Nash III TH (1996) Nitrogen, its metabolism and potential contribution to ecosystems. In: Lichen Biology. T.H. Nash III, ed. Cambridge University Press, Cambridge, pp. 136-153

Nimis PL, Scheidegger C, Wolseley PA (2002) Monitoring with Lichens. In: Nimis PL, Scheidegger C, Wolseley PA (eds) Monitoring Lichens. NATO Science Series, vol. 7. Springer, Dordrecht

Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara RB, Simpson GL, Solymos P, Stevens MHH, Wagner H (2013) Community ecology package. R package version 2.0. http://CRAN.Rproject.org/package=vegan

Peksa O & Škaloud P (2011) Do photobionts influence the ecology of lichens? A case study of environmental preferences in symbiotic green alga *Asterochloris* (Trebouxiophyceae). *Molecular Ecology* 20:3936-3948

Perotto S, Girlanda M, Martino E (2002) Ericoid mycorrhizal fungi: some new perspectives on old acquaintances. *Plant and Soil* 244:41-53

Pilkington MG, Caporn SJ, Carroll JA, Cresswell N, Lee J A, Emmett BA, Bagchi R (2007) Phosphorus supply influences heathland responses to atmospheric nitrogen deposition. *Environmental Pollution* 148:191-200

Pino-Bodas R & Stenroos S (2021) Global biodiversity patterns of the photobionts associated with the genus *Cladonia* (Lecanorales, Ascomycota). *Microbial Ecology* 82:173-187

Pino-Bodas R, Sanderson N, Cannon P, Aptroot A, Coppins B, Orange A, Simkin J (2021) Lecanorales: Cladoniaceae, including the genera *Cladonia*, *Pilophorus* and *Pycnothelia*. *Revisions of British and Irish Lichens* 19:1-45

Põlme S, Abarenkov K, Henrik Nilsson R, Lindahl BD, Clemmensen K, Kauserud H, Nguyen N Kjoller R, Bates ST, Baldrian P, Guldberg Frøslev T, Adojaan K, Vizzini A, Suija A, Pfister D, Baral HO, Järv H, Madrid H, Nordén J, Liu J-K *et al* (2020) FungalTraits: a user-friendly traits database of fungi and fungus-like stramenopiles. *Fungal Diversity* 105:1-16

Power SA, Ashmore MR, Cousins DA, Sheppard LJ (1998) Effects of nitrogen addition on the stress sensitivity of *Calluna vulgaris*. *New Phytologist* 138:663-67

Power SA & Barker CG (2003) Deposition measurements at Thursley Common heathland nature reserve. In: Empirical Critical Loads for Nitrogen: Proceedings of An Expert Workshop, Berne, Switzerland 11–13 November 2002 (eds Acherman B., Bobbink R.), pp. 259-261. Swiss Agency for the Environment, Forests and Landscape, Berne.

Pressel S, Kowal J, Duckett JG (2021) A tale of two fires: heathland bryophyte successions at Thursley, Surrey, long and short-term. *Field Bryology* 125:34-49

Rai H & Gupta RK (2022) Lichenized fungi, a primary bioindicator/biomonitor for bio-mitigation of excessive ambient air nitrogen deposition worldwide. In *Relationship Between Microbes and the Environment for Sustainable Ecosystem Services,* Vol. 1: 267-301. Elsevier

Read DJ (1996) The structure and function of the ericoid mycorrhizal root. *Annals of Botany* 77:365-374

Rodwell JS (2006) NVC Users' Handbook, JNCC, Peterborough, ISBN 978 1 86107 574 1.

Rognes T, Flouri T, Nichols B, Quince C, Mahé F (2016) VSEARCH: a versatile open-source tool for metagenomics. *PeerJ* 4:e2584

Rowe EC, Smart SM, Kennedy VH, Emmett BA, Evans CD (2008) N deposition increases the acquisition of phosphorus and potassium by heather *Calluna vulgaris*. *Environmental Pollution* 155: 201–207. https://doi.org/10.1016/j.envpol.2007.12.008.

Sanderson NA (2017) *The New Forest Heathland Lichen Survey 2011–2015*. A report by Botanical Survey & Assessment to Natural England, Forest Enterprise & The National Trust.

Schellenberg J & Bergmeier E (2022) The *Calluna* life cycle concept revisited: implications for heathland management. *Biodiversity and Conservation* 31:119-141

Selosse M-A, Setaro S, Glatard F, Richard F, Urcelay C & Weiß M (2007) Sebacinales are common mycorrhizal associates of Ericaceae. *New Phytologist* 174:864-878

Sérgio C, Garcia CA, Stow S, Martins A, Vieira C, Hespanhol H, Sim-Sim M (2018) How are anthropogenic pressures facilitating the invasion of *Campylopus introflexus* (Dicranaceae, Bryopsida) in mainland Portugal? *Cryptogamie Bryologie* 39:283-292

Škvorová Z, Černajová I, Steinová J, Peksa O, Moya P, Škaloud P (2022) Promiscuity in lichens follows clear rules: Partner switching in *Cladonia* is regulated by climatic factors and soil chemistry. *Frontiers in Microbiology* 12:781585

Smith SE & Read DJ (2008) Mycorrhizal symbiosis, 3rd edition. Academic, London, ISBN: 978-0-12-370526-6

Southon GE, Green ER, Jones AG, Barker CG, Power SA (2012) Long-term nitrogen additions increase likelihood of climate stress and affect recovery from wildfire in a lowland heath. *Global Change Biology* 18:2824-2837

Southon GE, Field C, Caporn SJM, Britton AJ, Power SA (2013) Nitrogen deposition reduces plant diversity and alters ecosystem functioning: Field-scale evidence from a nationwide survey of UK heathlands. *PLOS One* 8:e59031

Sparrius LB & Kooijman AM (2013) Nitrogen deposition and soil carbon content affect nitrogen mineralization during primary succession in acid inland drift sand vegetation. *Plant and Soil* 364:219-228

Stevens CJ, Smart SM, Henrys PA, Maskell LC, Crowe A, Simkin J, Cheffings CM, Whitfield C, Gowing DJG, Rowe EC, Dore AJ, Emmett BA (2012) Terricolous lichens as indicators of nitrogen deposition: evidence from national records. *Ecological Indicators* 20:196-203

Tang YS, Braban CF, Dragosits U, Dore AJ, Simmons I, van Dijk N, Poskitt J, Dos Santos Pereira, G, Keenan PO, Conolly, C, Vincent K, Smith RI, Heal MR, Sutton, MA (2018) Drivers for spatial, temporal and long-term trends in atmospheric ammonia and ammonium in the UK. *Atmospheric Chemistry and Physics* 18:705-733

Terry AC, Ashmore MR, Power SA, Allchin EA, Heil GW (2004) Modelling the impacts of atmospheric nitrogen deposition on *Calluna*-dominated ecosystems in the UK. *Journal of Applied Ecology* 41: 897-909

Treseder KK (2008) Nitrogen additions and microbial biomass: a meta-analysis of ecosystem studies. *Ecology Letters* 11:1111-1120

Turenne CY, Sanche SE, Hoban DJ, Karlowsky JA, Kabani AM (1999) Rapid identification of fungi by using the ITS2 genetic region and an automated fluorescent capillary electrophoresis system. *Journal of Clinical Microbiology* 37:1846-1851

van der Heijden MGA, Martin FM, Selosse M-A, Sanders IR (2015) Mycorrhizal ecology and evolution: the past, the present, and the future. *New Phytologist* 205:1406-1423

Van Geel M, Hans Jacquemyn H, Peeters G, van Acker K, Ceulemans T (2020) Diversity and community structure of ericoid mycorrhizal fungi in European bogs and heathlands across a gradient of nitrogen deposition *New Phytologist* 228:1640-1651

van Paassen JG, Britton AJ, Mitchell RJ, Street LE, Johnson D, Coupar A, Woodin SJ (2020) Legacy effects of nitrogen and phosphorus additions on vegetation and carbon stocks of upland heaths. *New Phytologist* 228:226-237

Vesala R, Kiheri H, Hobbie EA, van Dijk N, Dise N, Larmola T (2021) Atmospheric nitrogen enrichment changes nutrient stoichiometry and reduces fungal N supply to peatland ericoid mycorrhizal shrubs. *Science of the Total Environment* 794:148737

Vohník M, Sadowsky JJ, Kohout P, Lhotáková Z, Nestby R, Kolaříket M (2012) Novel root-fungus symbiosis in Ericaceae: Sheathed ericoid mycorrhiza formed by a hitherto undescribed basidiomycete with affinities to Trechisporales. *PLOS One* 7:e39524

Vohník M, Pánek M, Fehrer J, Selosse M-A (2016) Experimental evidence of ericoid mycorrhizal potential within Serendipitaceae (Sebacinales). *Mycorrhiza* 26:831-846

Walker JF, Aldrich-Wolfe L, Riffel A, Barbare H, Simpson NB, Trowbridge J, Jumpponen A (2011) Diverse Helotiales associated with the roots of three species of Arctic Ericaceae provide no evidence for host specificity. *New Phytologist* 191:515-527

Ward EB, Duguid MC, Kuebbing SE, Lendemer JC, Bradford MA (2022) The functional role of ericoid mycorrhizal plants and fungi on carbon and nitrogen dynamics in forests. *New Phytologist* 235:1701-1718

White T, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of ribosomal RNA genes for phylogenetics. Pages 315-322 In: Innis M, Gelfald D, Sninsky J, White T, editors. *PCR protocol. A Guide to Methods and Applications*. London: Academic Press

Zhou Z, Wang C, Zheng M, Jiang L, Luo Y (2017) Patterns and mechanisms of responses by soil microbial communities to nitrogen addition. *Soil Biology and Biochemistry* 115:433-441

Acknowledgements

Funding for this study came from Kew Science Pilot Grant scheme, the Joint Nature Conservation Committee and the Ecological Continuity Trust. R. P-B. was funded by the program 'Atracción de Talento Investigador', Comunidad de Madrid (Spain), 2020-T1/BIO-20503. We thank Emma Green and Georgina Southon for providing their time and raw data from previous long-term experiment studies at Thursley NNR. We are ever grateful to James Giles, the Thursley NNR site warden and Natural England who continue to support our science investigations by providing access to the site and Bliss Furtado, who helped analyse the heather roots for colonisation levels.

Data availability statement

Our soil and root-derived fungal sequences have been deposited in National Center for Biotechnology Information (NCBI) (accession numbers: pending) and one sequence per photobiont haplotype has been submitted to GenBank (accession numbers: OQ428652-OQ428657). All other raw data is in supplemental information.

Fig. 1 Comparison of aboveground biomass measures, between treatments (±SD). (A) mean height of vegetation; (B) tallest point within a subplot; (C) number of *Calluna vulgaris* points per quadrat (50 cm x 50 cm) counted at 20cm; and (D) *Campylopus introflexus* ground cover. Means ± SD.

Fig. 2 Treatment effects on lichen richness (A); abundance (B); diversity using Shannon

Index, H (C); and, diversity, using inverse Simpson index (D). Means ± SD.

Fig. 3 (A) Principal Component Analysis of soil characteristics; (B) Loading plots of soil

characteristics based on treatments.

Fig. 4 Root colonisation by treatment analysed (A) as mean percentage of individual roots colonized ($p > 0.05$); and (B) Chi-square of all *Calluna vulgaris* cells evaluated ($X^2(1, N = 50,918) = 11.2$, $p < .0001$). Significant *P* value differences are shown as: ***, < 0.001, **, < 0.01; * < 0.05, ns = not significant. Means ± SD.

Fig. 5 Krona diagrams showing all (A-B) and ErM-only (C-D) fungal taxa identified within *Calluna vulgaris* roots (A-C) and in soil (B-D). The ErM taxa are highlighted in green in charts A and B. Percentages represent the abundance of each taxon in relation to the total of all (A) and ErM only (C) fungal community in roots, and to the dominant soil community taxa (>10,000 reads) in soil (B and D).

Fig. 6 Non-metric multidimensional scaling (NMDS) and PERMANOVA analyses of root (A,B) and soil (C,D) fungal community differences between control and nitrogen (N)-addition treatments. A) All root fungi; B) Putative root-derived ericoid mycorrhizal fungi (ErM) only; C) All soil fungi; D) Putative soil ErM fungi only. All NMDS converge with stress < 0.2.

Supporting Material:

Assessing above and belowground recovery from ammonium sulfate addition and wildfire in a lowland heath: mycorrhizal fungi as potential indicators

Authors: Jill Kowal[1], Raquel Pino-Bodas[1,2], Elena Arrigoni[1], Guillaume Delhaye[1], Laura M. Suz[1], Martin I. Bidartondo[1,3], Jeffrey G. Duckett[4] and Silvia Pressel[4]

5- Royal Botanic Gardens, Kew, England
6- Universidad Rey Juan Carlos, Móstoles, Spain
7- Imperial College London, England
8- Natural History Museum, London, England

Supporting Tables

Table S1 Chronology of previous N-addition experiments at Thursley National Nature Reserve with fire history

Study Years	N+ rate	Organism variable tested: Results of N+	Authors
1976	FIRE		
1994-1996	7±7 and 15±4 kg N ha^{-1} yr^{-1})	1. *Calluna vulgaris* (heather) canopy: increased gaps; shoot water loss rate faster 2. heather beetles: adult weight increased leading to insect damage	Power et al. 1998. *New Phytol* 138.
1989-1996	0 or up to 15±4 kg N ha^{-1} yr^{-1}	Recovery after 7 years: 1. heather growth: persistent differences in growth and phenology; suppression of lichen abundance; stimulation of mosses 2. soil pH: recovered 3. soil microbial community size/activity	Power et al. 2006. *Glob Change Biol* 12.
1998-2006 FIRE 2010	0 or 30 kg N ha^{-1} yr^{-1}	Half the original plots (4 of 8 blocks) but double the N-addition: 1. drought injury: exacerbated 2. heather growth: persistent differences in growth and phenology (significantly more bud bursts than the controls over three consecutive years) 3. ongoing suppression of lichens and increased vigour in some bryophytes - the moss *Hypnum jutlandicum* and the liverwort *Cephaloziella divaricata*, but not the majority of mosses (*Campylopus introflexus*, *Ceratodon purpureus*, *Dicranum scoparium*, *Pohlia nutans* and *Polytrichum juniperinum* 4. soil pH reduction; C and N stocks increased (prior and post-fire) 5. soil microbial community size/activity elevated	Jones & Power 2012; Southon et al. 2012. *Glob Change Biol* 18.

Table S2a,b,c Summary of vegetation (a) density measures (b) and lichen species abundance (c) at Thursley National Nature Reserve, NVC H2 habitat (recorded July 2021).

(see excel SI)

Table S3 Comparison of aboveground mean biomass measures of *Calluna vulgaris*, *Campylopus introflexus* and *Hypnum jutlandicum* by treatment and block using two-way ANOVA contrasted with results measured by Southon et al. (2012). Mean values shown with standard deviation. *P* value significance = *<0.05; **<0.01; ***<0.001; ns, not significant; n/a, not applicable as not tested. N = 48.

| | treatments | | | factor *P* value | |
Biomass measures	Control	N-addition	subplot level n	Treatment	Southon
Height (mean cm ± SEM)					
C. vulgaris highest point	44.9±4.8	51.7±3.0	24	*	**
C. vulgaris mean height	40.5±4.4	47.3±2.1	24	*	**
Density of vegetation					
C. vulgaris at 20 cm	24.1±6.7	28.6±4.0	40 points	ns	**
C. vulgaris ground level	26.5±3.6	23.3±0.9	40 points	ns	n/a
C. introflexus	8.9±2.5	14±2.8	40 points	*	ns[1]
H. jutlandicum	1.9±1.9	6.4±3.2	40 points	ns	***
Stem rings					
Ring years (widest radius)	ns	ns	5 stems	ns	n/a
diameter	ns	ns	5 stems	ns	n/a
growth (mm) first 5 years	ns	ns	5 stems	ns	n/a

Table S4 Summary of growth ring measures.

(see Excel SI)

Table S5a Differences in lichen richness and diversity compared with the previous long-term experiment (LTE, Southon et al. 2012). Values are shown as mean ± standard deviation (n=48). Significant *P* value differences are shown as: ***, < 0.001; **, < 0.01; * < 0.05; ns, not significant.

	Treatment		P value	t value	LTE
	Control	N-addition			
Abundance	58.83± 22.30	46.42 ± 22.10	0.014*	-2.45	n/a
Richness	7.83 ± 2.80	8.13 ± 2.60	0.624	0.49	ns
Shannon index	1.27± 0.45	1.55 ± 0.40	0.005**	2.76	n/a
Simpson Index	2.83 ± 0.99	3.98 ± 1.64	0.000***	0.31	n/a

Table S5b Results of the indicator species analysis, indicating the lichen species associated with the different treatments, Control and N-addition.

Species	Indicator statistic	P values	Treatment	
			Control	N-addition
Cladonia fimbriata	0.446	**0.002**		X
Cladonia crispata	0.326	**0.024**	X	
Cladonia furcata	0.499	**0.002**	X	
Cladonia gracilis	0.335	**0.018**	X	

Table S6 Mean abundance of each species per treatment.

Species	Treatment	
	Control	N+
Cladonia cervicornis	0.542 ± 0.977	0.750 ± 1.511
Cladonia chlorophaea	0.167 ± 0.482	0.458 ± 1.021
Cladonia ciliata	0.083 ± 0.408	0.120 ± 0.4482
Cladonia coniocraea	0.000	0.042 ± 0.204
Cladonia crispata	10.708 ± 13.795	3.583 ± 5.641
Cladonia cryptochlorophaea	0.042 ± 0.204	0.042 ± 0.204
Cladonia diversa	1.458 ± 1.769	2.208 ± 3.050
Cladonia fimbriata	0.875 ± 1.154	6.250 ± 7.697
Cladonia floerkeana	1.083 ± 1.586	2.625 ± 4.576
Cladonia furcata	27.917 ± 14.108	14.667 ± 8.820
Cladonia glauca	0.958 ± 1.706	0.750 ± 1.294
Cladonia gracilis	0.750 ± 1.225	0.125 ± 0.338
Cladonia macilenta	0.833 ± 2.099	1.125 ± 1.650
Cladonia merochlorophaea	0.042 ± 0.204	0.000

Cladonia novochlorophaea	0.167 ± 0.482	0.458 ± 2.245
Cladonia polydacdactyla	0.000	0.167 ± 0.482
Cladonia portentosa	6.292 ± 6.504	4.0832 ± 4.138
Cladonia ramulosa	3.125 ± 3.314	2.667 ± 3.422
Cladonia squamosa	0.333 ± 1.090	0.042 ± 0.204
Cladonia subulata	0.208 ± 0.509	0.208 ± 0.509
Cladonia verticillata	3.250 ± 4.406	6.042 ± 5.797

Table S7 Comparison of soil characteristics between treatment groups using linear mixed effects models. CEC, cation exchange capacity. Values are shown as mean ± standard deviation (n=24). Significant *P* value differences are shown as: **, < 0.01; * < 0.05.

Soil characteristics	Control	N-addition	*P* value	t value	df
pH	4.70±0.04	4.49±0.05	0.00**	-4.07	10
H_2O (mg/l)	5.53±0.67	6.16±0.94	0.52	8.29	10
Total N (% w/w)	0.04+0.01	0.04±0.01	0.53	0.65	10
Total C (% w/w)	1.29±0.23	1.53±0.32	0.35	0.75	10
C:N ratio	38.24±1.54	36.97±1.91	0.52	-0.66	8
C stock (t/ha)	23.44±3.61	26.37±5.11	0.57	0.58	10
P (mg/l)	4.43±0.85	5.00±0.61	0.37	0.94	8
K (mg/l)	9.08±4.82	13.53±3.15	0.20	1.41	8
Mg (mg/l)	11.83±1.07	8.33±1.51	0.04*	-2.32	10
Nitrate (mg/kg)	0.05±0.00	0.06±0.00	0.23	1.25	20
Ammonium (mg/kg)	0.08±0.03	0.14±0.04	0.15	1.50	22
Dry matter (%)	90.78±1.90	88.46±2.40	0.27	-1.18	8
Soil density (g/l)	1217±34	1179±32.81	0.17	1.22	8
Na (mg/l)	5.31±0.39	6.09±0.52	0.18	1.49	8
Ca (mg/l)	31.30±9.31	28.36±13.17	0.83	-0.22	22
CEC (meq/100g)	10.46±0.13	10.97±0.16	0.00**	3.21	20

Table S8 Results of mean foliar N, P and C analyses comparing the different treatments, Control and N-addition, all not significant.

Total leaf N		Total leaf P	
Control	N addition	Control	N addition
1.03	1.02	0.08	0.08
1.07	1.07	0.09	0.09
1.06	1.09	0.08	0.07
0.96	0.97	0.08	0.07
1.23	1.14	0.1	0.09
1.14	1.21	0.09	0.08
P value	0.9749	P value	0.2073
t, df	t=0.03231, df=10	t, df	t=1.348, df=10

Total leaf C		Leaf C:N	
Control	N addition	Control	N addition
54.4	54.2	52.7	53.1
54	54.2	50.6	50.7
54.1	54	51.2	49.7
50.1	53.7	52.4	55.1
53	53	43.2	46.3
53.1	53.8	46.5	44.3
P value	0.3216	P value	0.852
t, df	t=1.043, df=10	t, df	t=0.1915, df=10

Table S9 Summary of ErM colonisation count data of all root cells and results of Chi-square analysis (with Yates correction).

Root cells		Control	N-adition	Total
Count data				
colonised		13,113	12,444	25,557
uncolonised		11,773	13,588	25,361
Total		24,886	26,032	50,918
Percentages				
colonised		52.69%	47.80%	
uncolonised		47.31%	52.20%	
	Chi-square degree of freedom		121.5,1	
z			11.02	
P value			<0.0001****	

Table S10 Fungal Species Hypotheses (SH) present in *Calluna vulgaris* roots, their taxonomy and putative function, total number of DNA sequences per SH and Amplicon Sequence Variants (ASVs) per SH, by treatment group (Control, N-addition). Relative abundances are calculated as the proportion of sequences in each SH. Green highlights represent ericoid mycorrhiza taxa.

Species Hypothesis (SH)	SH Taxonomy	Putative function	#sequences/SH			#ASVs/SH			Relative abundance (%)		Presence in soil (this study)	Presence in Van Geel et al., 2020
			Total	Control	N-addition	Total	Control	N-addition	Control	N-addition		
SH1156373.08FU	Sordariomycetes sp.	other	101	63	38	43	28	17	36.84	20.99	x	
SH1186518.08FU	Lecanorales sp.	lichen	53	24	29	19	10	12	14.04	16.02		
SH1185398.08FU	*Oidiodendron maius*	**ErM**	34	12	22	11	5	8	7.02	12.15		x
SH1156094.08FU	*Hyaloscypha finlandica*	**ErM**	25	11	14	22	11	12	6.43	7.73	**x**	**x**
SH1142376.08FU	Rhytismatales sp.	other	12	11	1	7	7	1	6.43	0.55	x	
SH1186366.08FU	Chaetothyriales sp.	**ErM**	11	6	5	9	6	4	3.51	2.76		x
SH1236001.08FU	*Phialocephala bamuru*	other	10	4	6	7	3	5	2.34	3.31		x
SH1155585.08FU	Helotiales sp.	**ErM**	7	1	6	7	1	6	0.58	3.31	**x**	**x**
SH2604623.08FU	Lecanoromycetes sp.	lichen	7	2	5	5	2	3	1.17	2.76		
SH2618221.08FU	Agaricomycetes sp.	other	5	5	0	5	5	0	2.92	0.00		
SH1172345.08FU	Serendipitaceae sp.	**ErM**	4	1	3	4	1	3	0.58	1.66	**x**	**x**
SH1156864.08FU	*Mollisia cinerea*	other	4	0	4	3	0	3	0.00	2.21		
SH1172852.08FU	Hymenochaetales sp.	other	3	0	3	3	0	3	0.00	1.66	x	
SH1170828.08FU	*Mycena romagnesiana*	ericoid_associate	2	2	0	2	2	0	1.17	0.00	x	
SH1186383.08FU	Chaetothyriales sp.	**ErM**	2	0	2	2	0	2	0.00	1.10		
SH1186231.08FU	*Saitozyma podzolica*	other	2	0	2	2	0	2	0.00	1.10		
SH1170824.08FU	*Mycena zephyrus*	ericoid_associate	1	1	0	1	1	0	0.58	0.00	x	
SH1171997.08FU	*Luellia recondita*	other	1	0	1	1	0	1	0.00	0.55		

SH1155587.08FU	Ascomycota sp.	other	1	0	1	1	0	1	0.00	0.55	x	
SH1160531.08FU	*Penicillium adametzii*	other	1	0	1	1	0	1	0.00	0.55	x	
SH1193918.08FU	Serendipitaceae sp.	**ErM**	1	1	0	1	1	0	0.58	0.00		x
SH1172222.08FU	Mycosymbioces sp.	other	1	1	0	1	1	0	0.58	0.00	**x**	**x**
SH1171998.08FU	*Luellia* sp.	other	1	0	1	1	0	1	0.00	0.55	x	
SH_unknown	na	na	63	26	37	60	25	35	15.20	20.44		
Total			**352**	**171**	**181**	**218**	**109**	**120**				
ErM (%)			**23.9**	**18.7**	**28.7**	**25.7**	**22.9**	**29.2**				
non-ErM (%)			**76.1**	**81.3**	**71.3**	**74.3**	**77.1**	**70.8**				

1 **Table S11** Summary of Operational Taxonomic Units (OTUs), Amplicon Sequence Variants
2 (ASVs) and Species Hypotheses (SHs) detected in Thursley root and soil samples compared
3 with our SH compilation of root-derived fungi from 21 European heathlands 'VG' (Van Geel
4 et al., 2020).

Fungi source	OTUs		ASVs		SHs	
	all fungi	ErM	all fungi	ErM	all fungi	ErM
Roots (this study, Thursley)	n/a	n/a	218	87	23	7
Soil (this study, Thursley)	1209	73	n/a	n/a	570	40
Roots (VG heathlands)	n/a	239	n/a	n/a	n/a	82

5
6

7 **Table S12** Summary of genetic variation of photobionts associated with *Cladonia* species on
8 control and N-addition plots.

	Control	N-addition
Number of sequences (N)	113	101
Number of haplotypes (H)	5	3
Haplotype diversity (Hd)	0.0868	0.059
Nucleotide diversity (π)	0.001	0.0003

9
10

11 **Table S13** Analysis of molecular variance (AMOVA) within subplot using treatments as
12 grouping factor for ITS rDNA haplotypes of *Asterochloris italiana* associated with *Cladonia*. SS,
13 sum of squared differences of each observation from the mean; F_{st} denotes population
14 differentiation due to genetic structure. ** P-value >0.01; ns = no significant, P-value >0.05.

	SS[1]	Variance components	% Variation	Fixation Index
Among treatments	0.024	-0.001	-3.03	F_{st} = 0.128**
Among subplots within treatments	1.147	0.005	13.23	F_{st} = 0.102**
Within subplots	6.641	0.033	89.80	F_{st} = -0.030 ns
Total	7.812	0.037		

Supporting Figures

a)

b)

Fig. S1 Mean annual temperature and precipitation (mm), Thursley National Nature Reserve,
Surrey (2009-2022, June only).

a)

28
29
30
31 **b)**

32
33

c)

Fig. S2a, b, c   (a) Original field experimental block design (Power et al., 1998, modified by
Southon et al., 2012). Red Xs indicates blocks removed by Southon. Each block (numbered 1
through 4, in bold) has either six or four 4x4 m plots which are then subdivided into four 2x2

m sub-plots. The management treatments indicated in the key (old recovery and recovery)
were not significant. The three blocks tested in this study are circled in red and enlarged in
(b). These three blocks each have two ammonium sulphate addition plots and two control.
Source: Emma Green notes. (c) Map of Thursley trial plots, circled in red (Source: Natural
England) with corresponding aerial view of the plots tested in this study (5,6,7) adapted from
Google Earth (downloaded October 2021).

51
52   **Fig. S3** Image showing cross-section of *Calluna vulgaris* stem with three measurements per
53   ring. Pith in the center.
54
55

Fig. S4 Comparison between control 'C' and N-addition 'N' treatments of annual ring growth at Thursley post fire of 2006. There is no significant difference in the height of the peak in 2014 between control and N-addition treatments (LMM: t = 0.44, df = 52, p = 0.66).

Fig S5. Nonmetric multidimensional scaling showing *Cladonia* spp. community based on the Bray-Curtis dissimilarity matrix comparing Control and N-addition (N+) plots. All convergent with stress < 0.2.

Fig S6. Box plots comparing abundance responses to treatments for four different *Cladonia* spp. a) *C. crispata (P* value = 0.024), b) *C. gracilis* (*P* value = 0.018), c) *C. furcata P* value = 0.002), and d) *C. fimbriata* (*P* value = 0.002); N+, n-addition plots. Means ± SD.

Fig. S7 Heatmap of Pearson r correlation coefficients between ErM root colonization and ErM soil richness (SH) with soil and leaf chemical parameters.

A. italiana

OTU14

Fig. S8 Phylogeny of *Asterochloris* based on ITS rDNA. Maximum Likelihood tree of RAxML. Bold branches were supported by ≥ 75% of bootstrap values. The clade including all the sequences of photobionts from the Thursley *Cladonia* spp. is marked in red.

Fig. S9 Statistical parsimony haplotype network of *Asterochloris* based on ITS. Haplotypes are coloured according to (a) treatment, Control and N+ (nitrogen addition) and (b) blocks 5,6 and 7. The size of the circles is proportional to the frequency of each haplotype. Small circles represent haplotypes not observed in the data.

Supporting Data S1. Field Experiment: Thursley National Nature Reserve, Surrey, UK (1998-2010)

In 1989, a long-term experiment was set up at Thursley National Nature Reserve, to mimic and test ammonium sulphate on heathland ecosystem function (Power et al. 2006; Jones & Power 2012; Southon et al. 2012). The results showed that ammonium sulphate addition, applied bi-weekly between 1998 and 2010, resulted in significant changes in aboveground vegetation structure and diversity (Table S1) and belowground nutrient cycling, specifically C and N stocks. In 2006 the study was interrupted by a severe wildfire (Duckett et al. 2008) which affected all experimental plots and completely eliminated the successional heathland structure i.e., pioneer, building, mature and degeneration (Gimingham 1972). Following this event, the effects of fire and N deposition interactions were studied for an additional six years (Southon et al. 2012). Despite the wildfire, which removes most of the available N from *C. vulgaris* heathlands (Pate & Dell 1984; Read 1996), Southon et al. (2012) found continuing effects of the pre-fire N treatments on post-fire vegetation regeneration among pioneer species (Table S1). The dominant heather shrubs no longer had significant differences in growth parameters, likely due to post-disturbance regeneration from older stems (Schellenberg & Bergmeier 2022). The Southon et al. (2012) study also revealed significantly higher soil C in the N-addition plots, which was attributed to increases in the microbial biomass, although it did not examine mycorrhizal colonisation, root-associated fungi or soil fungi community structure.

References:

Gimingham CH (1972) Ecology of Heathlands. London: Chapman and Hall Ltd. ISBN 13: 9780412104602

Jones AG & Power S (2012) Field-scale evaluation of effects of nitrogen deposition on the functioning of heathland ecosystems. *Journal of Ecology* 100:331-342

Pate JS & Dell B (1984) Economy of mineral nutrients in sandplain species. In: Pate JS, Beard JS, eds. Kwongan. Plant life of the sandplain. Nedlands, Australia: University of Western Australia Press, 227-252

Power SA, Ashmore MR, Cousins DA, Sheppard LJ (1998) Effects of nitrogen addition on the stress sensitivity of *Calluna vulgaris*. *New Phytologist* 138:663-67

Power SA, Green ER, Barker CG, Bell NB, Ashmore MR (2006) Ecosystem recovery, heathland response to a reduction in N deposition. *Global Change Biology* 12:1241-1252

Schellenberg J & Bergmeier E (2022) The *Calluna* life cycle concept revisited: implications for
heathland management. *Biodiversity and Conservation* 31:119-141

Southon GE, Green ER, Jones AG, Barker CG, Power SA (2012) Long-term nitrogen additions
increase likelihood of climate stress and affect recovery from wildfire in a lowland heath.
Global Change Biology 18:2824-2837

Methods S1. Vegetation and lichen survey

For vegetation, each subplot we measured vegetation density, abundance and height and
recorded plant species composition Two quadrats (50 cm^2) subdivided into 25 squares (10
cm^2) were randomly set within each subplot. To obtain an estimate of vegetation density (or
cover) and abundance, using a similar point-grid quadrat system as in the original study
(Southon et al. 2012), plant species data were recorded 20 times at two heights - at ground
level and at 20 cm, for a total of 40 vegetation points per subplot. To gauge the canopy height
differences between treatments two measures were taken: mean height derived from four
randomly selected points and the two highest canopy points within each of the quadrats. For
lichens, species present in both quadrats were recorded with the number of squares in which
each species occurred as a measure of abundance. For the algal layer, two mm^2 of the algal
layer from clean healthy thalli were selected for DNA extraction. Lichens were identified using
published morphological characters (Ahti & Stenroos 2013; Burgaz et al. 2020; Pino-Bodas et
al. 2021). Additionally, thin layer chromatography (TLC) was performed to identify secondary
metabolites and confirm morphological identifications following the protocol described by
Orange (2001).

Ahti T & Stenroos S (2013) Cladoniaceae. In: Ahti T, Stenroos S, Moberg R (eds) *Nordic Lichen*

Flora 5:1-117

Methods S2. **Soil chemical extraction methods**

Upon returning to the laboratory, visible rocks and roots were removed and soil samples were

maintained overnight at 3-4°C. The next day they were delivered in cool boxes to an external

laboratory for processing and analyses (NRM Laboratories Services, Bracknell England). Below

are the abridged methods employed by NRM Laboratories.

Determination of soil bulk density

The total weight of the sample was recorded after drying overnight at < 30°C. This information

was used to calculate the soil bulk density on a dry matter basis for the set volume. A scoop

of set volume is used to take a sample of the dried and ground soil (after passing through a

2mm sieve). This is then levelled off before the weight is recorded.

Soil Density (g/l) = Weight of soil (g)/Volume of ring (l)

Reference: Soil Science – Methods and Applications. DL Rowell. ISBN 0-582-08784-8.

Determination of Electrical Conductivity (EC)

Matrix: Soil samples were air-dried at a temperature <30°C and sieved to pass a 2mm screen,

excluding stones and fibrous material and roots.

Soluble salts, other than calcium sulphate, are extracted from soil with saturated calcium

sulphate solution. The specific conductivity of the extract at 20°C is recorded as soil

conductivity. Results are expressed as uS/cm at 20°C.

Reference: The Analysis of Agricultural Materials, MAFF Reference Book RB427, ISBN 0 11
242762 6.

Determination of soil pH

Matrix: Soil samples were air-dried at a temperature <30°C and sieved to pass a 2mm screen,

excluding stones and fibrous material and roots.

A suspension of 5 grams of soil with 25 ml of water was shaken on an orbital shaker for 15

min and rested for 45 min prior to pH analysis using a Sentek pH electrode.

Reference method: ISO 10390.

Determination of total Carbon and Nitrogen

Matrix: Soil samples were dried and ground to pass through a 0.5mm screen.

Total Carbon and Total Nitrogen are determined by combustion analyser. Results are
expressed as percentage in soil.

Reference: AOAC Official Methods of Analysis (1990) Method 949.12. ISO 10694 & 13878.

Determination of plant available Phosphorus using Melich III extractable elements in soil
(for soil pH <6.0)

Matrix: Soil samples were air-dried at a temperature <30°C and sieved to pass a 2mm screen,
excluding stones and fibrous material and roots.

The sample is shaken with Mehlich III extracting solution which dissolves the 'active'
compounds. The extracted compounds are determined by Inductively Coupled Plasma Optical
Emission Spectroscopy. This extractant is composed of 0.2M acetic acid; 0.25M ammonium
nitrate; 0.015M ammonium fluoride; 0.013M nitric acid; and 0.001M EDTA. Exchangeable K,
Ca, Mg, and Na are extracted by the action of ammonium nitrate and nitric acid.
Micronutrients are extracted by NH4 and the chelating agent EDTA. Acetic acid and
ammonium fluoride are present for the extraction of phosphorus. Results are reported as
mg/l in soil on a dry matter basis.

Reference: Mehlich, A. 1984. Mehlich III soil test extractant: A modification of the Mehlich 2
extractant. Commun. Soil Sci. Plant Anal. 15:1409-1416.

Determination* of *mineral nitrogen in soil (available N)

Background: The protocol assumes soil available N includes nitrate-N, nitrite-N and
ammonium-N that can be extracted by 2M Potassium Chloride.

Matrix: The soil samples were maintained between 2-8°C to restrict nitrogen mineralization
which could lead to an overestimation of available N.

The soil sample is chopped and mixed to obtain a homogenous sample, removing stones and
roots. A portion is then shaken with 2M KCl to extract the mineral-N fractions and a dry matter
determination carried out. Once in solution the Nitrate-N, Nitrite-N and Ammonium-N can be
measured colourimetrically. Results are reported as mg/l in soil on a dry matter basis.

References:

Soil Sampling and Methods of Analysis. Canadian Society of Soil Science, Ed Martin R Carter.
ISBN 0-87371-861-5.
The Analysis of Agricultural Materials, MAFF Reference Book RB427, ISBN 0 11 242762 6.
Reference method ISO 10694 & 13878.

Determination extractable potassium, calcium, sodium and magnesium in soil

Matrix: Soil samples were air-dried at a temperature <30°C and sieved to pass a 2mm screen,

excluding stones and fibrous material and roots.

The available potassium and magnesium are extracted from the soil by shaking with 1M

ammonium nitrate at 20°C for 30 minutes. After filtration, the concentration of potassium

and magnesium in the extract is determined by atomic absorption spectrometry. The

instrument is calibrated using commercial potassium and magnesium standards traceable to

the SI unit.

References:

The Analysis of Agricultural Materials, DEFRA Reference Book RB427 ISBN 0 11 242762 6.

Fertiliser Recommendations for Agricultural and Horticultural Crops, DEFRA Reference Book
RB209 9th edition.

Determination* of *available sulphate

Matrix: Soil samples were air-dried at a temperature <30°C and sieved to pass a 2mm screen,

excluding stones, fibrous material and roots.

Available sulphate is extracted from the soil using a phosphate buffer extracting solution (1:2)

and the filtered extract of the sample is analysed by Inductively Coupled Plasma Emission

Spectroscopy.

Reference:

Soil Science - Methods and Applications, D L Rowell, 1995, Longman Scientific and Technical,
ISBN 0 582 087848.

Determination* of *available manganese

Matrix: Soil samples were air-dried at a temperature <30°C and sieved to pass a 2mm screen,

excluding stones, fibrous material and roots.

The available element manganese is extracted from the soil at 20°C with DTPA solution (1:2)

adjusted to a steady pH of 7.3.

Reference: Developed by Lindsay and Norvell: Soltanpour, P N, and Schwab, A P. 1977. Comm.
Soil Sci. Plant Anal. 8:195-207. Handbook on Reference Methods for Soil Analysis - 1992, ISBN
09627606 1 7.
237
238

Methods S3. **Details of molecular testing and bioinformatics analyses on root-associated and soil fungal communities**

DNA was extracted in duplicate from 250 mg of soil (fresh weight) for each plot, using the DNeasy PowerSoil Pro kit (Qiagen, Hilden, Germany) according to manufacturer's instructions. DNA extracts were quantified using a Quantus™ Fluorometer with QuantiFluor® Dye System for dsDNA (Promega, Madison, Wisconsin, United States) and purity-checked (260/280nm and 260/230nm ratios) via NanoDrop spectrophotometer (Thermo Scientific, Waltham, Massachusetts, United States). DNA extracts were sent to Macrogen (Macrogen Genome Center, Seoul, Republic of Korea) for downstream processing and bioinformatic analysis. PCR amplification of the ITS2 region was performed using the primers ITS1F/ITS86F (White et al., 1990; Turenne et al., 2000), and amplicon libraries were sequenced on an Illumina MiSeq (Illumina, San Diego, California, United States). Reads pre-processing and clustering were carried out by Macrogen Representative sequences from each OTU were assigned to a Species Hypothesis (SH) in house, using the SH matching v2.0.0 tool on PlutoF. The complete ITS region (ITS1, 5.8S and ITS2) was amplified using the fungal-specific primers ITS1F and ITS4 (White et al., 1990; Gardes and Bruns, 1993). PCR reactions were carried out in a final volume of 10 µl, consisting of 5 µl of JumpStart Taq ReadyMix (Sigma-Aldrich, St. Louis, Missouri, United States), 0.05 µl of each primer (100 µM), 2.4 µl of PCR-grade water and 2.5 µl of template. Cycling conditions were as follows: 94°C for 2 min, 35 amplification cycles (94°C for 30 s, 53°C for 45 s, 72°C for 1:30 min) and final extension at 72°C for 7 min. Considering that plant roots are colonised by multiple fungal species, PCR fragments were ligated into the pCR®4-TOPO® vector and cloned into One Shot® TOP10 competent *Escherichia coli* cells, following the manufacturer's protocol for Kanamycin selection with modifications. Eight to 12 colonies per sample were directly amplified using ITS1F/ITS4 primers into 15 µl of PCR mix consisting of: 7.5 µl of JumpStart Taq ReadyMix, 0.075 µl of each primer (100 µM) and 7.35 µl of PCR-grade water. Cycling conditions were as follows: 94°C for 7 min, 25 amplification cycles (94°C for 30 s, 53°C for 45 s, 72°C for 1 min with the addition of 5 s per cycle during the extension phase), and final extension at 72°C for 7 min. Prior to sequencing, PCR products were purified using a 40% v/v dilution in PCR-grade water of a Exonuclease I : FastAP solution (ratio 1:2) (Thermo Fisher Scientific, Waltham, Massachusetts, United States). Purified PCR products were prepared for sequencing with BigDye v. 3.1 (Applied Biosystems, Waltham, Massachusetts, United States) and primers ITS1F and ITS4.

<u>Soil fungal communities</u>

DNA was extracted in duplicate from 250 mg of soil (fresh weight) for each plot, using the

DNeasy PowerSoil Pro kit (Qiagen, Hilden, Germany) according to manufacturer's

instructions. DNA extracts were quantified using Quantus^TM Fluorometer with QuantiFluor®

Dye System for dsDNA (Promega, Madison, Wisconsin, United States) and purity-checked

(260/280nm and 260/230nm ratios) via NanoDrop spectrophotometer (Thermo Scientific,

Waltham, Massachusetts, United States). DNA extracts were sent to Macrogen (Macrogen

Genome Center, Seoul, Republic of Korea) for downstream processing and bioinformatic

analysis. PCR amplification of the ITS2 region was performed using the primers ITS1F/ITS86F

(White et al., 1990; Turenne et al., 1999), and amplicon libraries were sequenced on an

Illumina MiSeq (Illumina, San Diego, California, United States). Reads pre-processing and

clustering were carried out by Macrogen. Representative sequences from each OTU were

assigned to a Species Hypothesis (SH) in house, using the SH matching v2.0.0 tool on PlutoF.

References

Gardes M & Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes-
application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2:113-118.

Turenne CY, Sanche SE, Hoban DJ, Karlowsky JA, Kabani AM (1999) Rapid identification of
fungi by using the ITS2 genetic region and an automated fluorescent capillary
electrophoresis system. *Journal of Clinical Microbiology* 37:1846-1851.

White T, Bruns T, Lee S & Taylor J (1990) Amplification and direct sequencing of ribosomal RNA
genes for phylogenetics. In: Innis M, Gelfald D, Sninsky J, White T, editors. *PCR protocol. A Guide*
to Methods and Applications. London: Academic Press. p. 315-322.

Methods S4. **Lichen photobiont analysis**

The lichen photobiont DNA was extracted using Chelex-100 resin (BioRad) following Ferencová et al. (2017). We analysed three specimens for each of the four lichen species found to be consistently present at each subplot. The ITS rDNA region was amplified with 7.5 µl of Green Taq Mastermix (Promega, Madison, WI), 0.7 µl of each primer at 10 µM concentration, and 1 µl of DNA (Pino-Bodas & Stenroos 2021). To determine to which *Asterochloris* lineage the photobionts sequenced belonged, the new sequences were included in the ITS rDNA alignment of Pino-Bodas & Stenroos (2021), with additional sequences from Kosecka et al. (2021), including the new lineages found by the authors. The alignment was made in MAFFT using default parameters (Katoh et al. 2019). The phylogeny was estimated with Maximum likelihood, implemented in RAxML 7.0.3 (Stamatakis et al. 2005) with GTRGAMMA model and 500 pseudoreplicates of rapid bootstrap. *Vulcanochloris canariensis* was used as outgroup. Because the taxonomy of the photobiont is still poorly understood (Muggia et al. 2018), most recent studies have used photobiont haplotypes to study the diversity and to infer the mycobiont-photobiont interaction (Pérez-Ortega et al. 2012; Cao et al. 2015; Garrido-Benavent et al. 2020, 2022). The number of photobiont haplotypes and genetic diversity in each plot was analysed with DnaSP (Librado & Rozas 2009). Haplotype networks under statistical parsimony were constructed in PopART 1.7 (Leigh & Bryant 2015) selecting the TCS method (Clement et al. 2000). Gaps were considered as missing data. Analysis of molecular variance (AMOVA) was conducted to determine the contribution of treatment in the genetic differentiation of photobionts, using ARLEQUIN V3.5 (Excoffier & Lischer 2010).

References:

Cao S, Zhang F, Liu C, Hao Z, Tian Y, Zhu L, Zhou Q (2015) Distribution patterns of haplotypes for symbionts from *Umbilicaria esculenta* and *U. muehlenbergii* reflect the importance of reproductive strategy in shaping population genetic structure. *BMC microbiology* 15:1-12

Clement M, Posada DC, Crandall KA (2000) TCS: a computer program to estimate gene genealogies. *Molecular Ecology* 9:1657-1659

Excoffier L & Lischer HE (2010) Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* 10:564-567

Ferencová Z, Rico VJ, Hawksworth DL (2017) Extraction of DNA from lichen-forming and lichenicolous fungi: a low-cost fast protocol using Chelex. *The Lichenologist* 49:521-525

Garrido-Benavent I, Pérez-Ortega S, de los Ríos A, Fernández-Mendoza F (2020)
Amphitropical variation of the algal partners of *Pseudephebe* (Parmeliaceae, lichenized fungi).
Symbiosis 82:35-48

Garrido-Benavent I, Molins A, Barreno E (2022) Genetic variation in the symbiont partners in
the endangered macrolichen *Seirophora villosa* (Teloschistaceae: Ascomycota). *Botanical*
Journal of the Linnean Society 199:816-829

Leigh JW & Bryant D (2015) POPART: full-feature software for haplotype network
construction. *Methods in Ecology and Evolution* 6:1110–1116

Librado P & Rozas J (2009) DnaSP v5: a software for comprehensive analysis of DNA
polymorphism data. *Bioinformatics* 25:1451-1452

Muggia L, Leavitt S, Barreno E (2018) The hidden diversity of lichenized Trebouxiophyceae
(Chlorophyta). *Phycologia* 57:503-524

Pérez-Ortega S, Ortiz-Álvarez R, Green TGA, de Los Ríos A (2012) Lichen myco- and
photobiont diversity and their relationships at the edge of life (McMurdo Dry Valleys,
Antarctica). *FEMS Microbiology Ecology* 82:429-448

Pino-Bodas R & Stenroos S (2021) Global biodiversity patterns of the photobionts associated
with the genus *Cladonia* (Lecanorales, Ascomycota). *Microbial Ecology* 82:173-187

Stamatakis A, Ludwig T, Meier H (2005) RAxML-III: a fast program for maximum likelihood-
based inference of large phylogenetic trees. *Bioinformatics* 21:456-463