A Serum- and Feeder-Free System to Generate CD4 and Regulatory T Cells from Human iPSCs

Authors

Helen Fong[1], Matthew Mendel[1], John Jascur[1], Laeya Najmi[1], Ken Kim[1], Garrett Lew[1], Swetha Garimalla[1], Suruchi Schock[1], Jing Hu[1], Andres Villegas[1], Anthony Conway[1,2], Jason Fontenot[1], Simona Zompi[1]

Affiliation

[1] Sangamo Therapeutics, 7000 Marina Blvd, Brisbane, CA 94005; 501 Canal Blvd. Richmond, CA 94804

[2] Current affiliation: Replay, 5555 Oberlin Drive, Unit 120, San Diego, CA 92121

Corresponding authors

Helen Fong: hfong82@gmail.com

Simona Zompì: simona.zompi@gmail.com

Abstract

iPSCs can serve as a renewable source of a consistent edited cell product, overcoming limitations of primary cells. While feeder-free generation of clinical grade iPSC-derived CD8 T cells has been achieved, differentiation of iPSC-derived CD4sp and regulatory T cells requires mouse stromal cells in an artificial thymic organoid. Here we report a serum- and feeder-free differentiation process suitable for large-scale production. Using an optimized concentration of PMA/Ionomycin, we generated iPSC-CD4sp T cells at high efficiency and converted them to Tregs using TGFβ and ATRA. Using zinc finger nucleases, we demonstrated high non-viral, targeted integration of an HLA-A2 CAR in iPSCs. iPSC-Tregs +/- HLA-A2 CAR phenotypically, transcriptionally and functionally resemble primary Tregs and suppress T cell proliferation *in vitro*. Our work is the first to demonstrate an iPSC-based platform amenable to manufacturing CD4 T cells to complement iPSC-CD8 oncology products and functional iPSC-Tregs to deliver Treg cell therapies at scale.

Introduction

Engineered autologous T cell therapies have shown the ability to transform the oncology field, yet producing these products at scale for large numbers of patients has proven difficult. The ability to derive hypoimmunogenic engineered T cells from self-renewing human induced pluripotent stem cells (iPSCs) could dramatically increase access to these therapies [1]. The advantages of using iPSCs over primary cells include the capacity to easily multiplex edit the starting cell source, access to a self-renewable and well-characterized product, and standardization of the product through the development of a master cell bank (MCB). However, while using iPSCs as source material can have numerous benefits in terms of standardization and scale-up, differentiation of T cells from iPSCs has been challenging.

Access to a self-renewable, potentially unlimited source of cells is particularly attractive for cell types that are difficult to isolate or present in small numbers, such as regulatory T cells (Tregs). In recent years there has been much interest in the use of Tregs to treat a wide variety of diseases, including autoimmune disorders, as well as in organ transplantation [2]. As of 2022, there were more than 30 active and completed clinical trials testing the safety and efficacy of Treg cell therapies [3]. Most of the studies have focused on *in vitro*-expanded polyclonal Tregs showing limited clinical efficacy. Promising pre-clinical results testing antigen-specific CAR Tregs [4] led to the first CAR Treg clinical trial authorization (NCT04817774) for kidney transplant patients, using *ex vivo,* autologous, naïve Tregs engineered with a CAR construct recognizing HLA-A2, to investigate the induction of tolerance in HLA-A2 negative patients receiving an HLA-A2 positive renal graft.

Currently, sourcing $CD4^+$ Teff and Tregs for cell therapies is invasive, time-consuming, and yields small numbers of cells, particularly of Tregs which comprise only 5-7% of the circulating CD4 T cell population [5]. Furthermore, $CD4^+$ Teff and Tregs obtained from these sources are polyclonal

in nature and can introduce variability and non-specificity. While we have been successful in

introducing a CAR construct in primary Tregs to mediate a specific immunosuppressive response

[4], the need to obtain antigen specific CD4[+] Teff and Tregs more efficiently and in large numbers

remains, especially genetically engineered Tregs.

Numerous studies have shown the ability to generate large quantities of T cells from iPSCs,

focusing primarily on the differentiation to cytotoxic T lymphocytes (CTLs) [6-8]. Improvements have

recently been made in the differentiation processes to generate iPSC-derived CAR CD8sp T cells

that more closely resemble primary CD8$\alpha\beta^+$ T cells rather than innate $\gamma\delta$ T cells, by knocking out

TRAC and delaying and attenuating CAR expression [9]. The artificial thymic organoid (ATO)

system has also proven to be successful in generating functionally mature CD3$^+$CD8$\alpha\beta^+$ T cells

that produce IFN-γ, TNF-α, and IL-2 and proliferate upon stimulation [8].

Interestingly, the ATO system also produces a small population of CD3$^+$CD4$^+$ T cells with a

conventional phenotype and diverse TCR repertoire [8, 10] that can be converted to FOXP3$^+$ Tregs

(Yano H. et al., ASH 2022). However, a scalable method to generate clinically relevant numbers

of CD4$^+$ T cells and Tregs from iPSCs has yet to be demonstrated. This need has become more

compelling in recent years as CD4$^+$ T effectors (Teffs) cells are known to enhance CD8$^+$ cytotoxic

T lymphocyte (CTLs) responses [11] and have been shown to be therapeutically useful in CAR-T

cell therapies increasing product potency [12, 13] and showing long-term functional persistency in

clinical trials [14].

Overall, current methods to generate CD8 single positive (CD8sp) T cells, CD4 single positive

(CD4sp) T cells and Tregs from iPSCs utilize serum or mouse stromal cells at some stage of the

differentiation process making it difficult for use in clinical development. In addition, the use of an

ATO system is challenging to scale-up for manufacturing as it requires the maintenance of a 3D

organoid system and involves dissociation and extraction of the cells of interest from a mixed

population of mouse and human cells. Here we show for the first time the ability to efficiently

generate both CD4sp T cells and Tregs from iPSCs in completely serum- and feeder-free

conditions. We identified a potent cell stimulator to convert $CD4^+CD8^+$ double positive T (DP T)

cells to CD4sp T cells at high efficiency that can then be converted to $FOXP3^+$ Tregs using TGFβ,

all-trans retinoic acid (ATRA), and IL-2 with CD3/CD28/CD2 activation. We show that the iPSC-

CD4sp T cells have an effector memory (T_{EM}) phenotype, are transcriptionally similar to primary

CD4 T conventional (Tconv) cells and can be activated upon stimulation. The iPSC-Tregs possess

typical Treg characteristics, are transcriptionally more similar to primary Tregs than primary Tconv

and suppress primary CD4 Tconv proliferation. In addition, iPSCs engineered to express an HLA-

A2 CAR and a fully functional TCR under the control of the *TRAC* promoter could effectively

differentiate to CD4sp T cells and then Tregs that also suppress Tconv proliferation *via* activation

through the TCR or HLA-A2 CAR. Finally, we also show that the iPSC-Tregs are amenable to

cryopreservation and retain function post-thaw. This novel method provides a viable path forward

towards the development of a scalable iPSC-derived Treg cell therapy and the development of

iPSC-derived CD4sp T cells which could enhance the iPSC-derived CD8sp T cell products

currently in the clinic for oncology applications [12, 14, 15].

Results

Generation of CD34[+] hematopoietic stem and progenitor cells (HSPCs) and T cell progenitors from iPSCs

iPSC clones were derived from sorted CD4[+]CD25[+] cells from a healthy human donor. Derivation of iPSCs from T cells has been previously documented with resulting T cells differentiated from reprogrammed iPSCs possessing antigen specificity due to retention of a rearranged TCR [7, 16, 17]. Additionally, evidence suggests improved T cell lineage differentiation when iPSCs originate from mature T cells due to removal of the barrier to TCR rearrangement. [7, 18]. CD4[+]CD25[+] T cells were isolated and FACS-purified from healthy human leukapheresis blood and reprogrammed by Sendai virus carrying vectors encoding for the four Yamanaka factors: OCT-4, SOX2, KLF4, and C-MYC. Multiple clones were generated and characterized by immunocytochemistry to confirm expression of OCT4, SOX2, TRA-1-81, and SSEA-4 **(Supp. Figure 1A)**. In addition, G-band karyotyping confirmed the absence of any chromosomal aberrations **(Supp. Figure 1B)**.

To facilitate future translation from discovery to process development we utilized serum-free, commercially available reagents from the iPSC stage to the DP T stage. As a first step, a modified version of the spin embryoid body (EB) method was used to differentiate the iPSCs into early stages of hematopoiesis from days 0 to 8, followed by expansion of the hematopoietic stem and progenitor cells from days 8 to 14 [19, 20] **(Figure 1A)**. The EBs increased in size over time and by day 14, numerous free-floating HSPCs emerged around the periphery expressing both CD34 and CD45 (65%±13.5%, n=3) **(Figure 1B and 1C)**. The approximate yield of HSPCs per input iPSC was 16.5±9.3, n=3 **(Figure 2)**.

For commitment and differentiation to the T cell lineage, interaction between blood cell progenitors and a Notch ligand is required [21-23]. To differentiate the iPSC-HSPCs to lymphoid progenitors, the entire population of free-floating HSPCs was separated from the EBs, collected, and seeded onto

plates coated with lymphoid differentiation coating material (LDCM), a commercially available

matrix containing a Notch ligand **(Figure 1D)**. The HSPCs were then further differentiated to

lymphoid progenitor cells that were CD5 and CD7 DP by day 28 (**Figure 1C**, 67.4±0.1%, n=6)

and subsequently to CD4 and CD8 DP T cells by day 42 (**Figure 1C**, DP T cells, 86.3±0.1%, n=5)

using a commercially available iPSC-HSPC to T cell differentiation system **(Figure 1D)**. The

approximate yield of lymphoid progenitor cells and DP T cells was 2.3e3±1.4e3 per input iPSC

and 2.6e4±1.6e4 per input iPSC, respectively **(Figure 2)**. We tested the use of CXCL12 (SDF1a)

and a p38 inhibitor (SB203580) to enhance the number of DP T cells [17] and found approximately

a 1.8-fold increase in DP T cells **(Supp. Figure 2)**.

Generation of CD4-sp T cells and FOXP3$^+$ Tregs from iPSC-CD4$^+$CD8$^+$ DP T cells

During T cell development in the thymus, TCR signaling is required in DP thymocytes to induce

positive selection and subsequent generation of either mature CD4$^+$ or CD8$^+$ T cells [24]. In addition

to differentiation into T cells, the interaction between the TCR and MHC molecules in the thymus

also rescues thymocytes from programmed cell death [25]. Alternatively, it has been shown that DP

thymocytes can undergo positive selection and maturation into CD4$^+$ T cells without engagement

of the TCR if provided with a strong, persistent signal [24, 26, 27]. We tested the ability of the iPSC-

derived DP T cells (iPSC-DP T) to convert to CD4sp T cells by exposure to three different doses

(low, medium, high) of the potent cell stimulator PMA/Ionomycin (PMA/I) **(Figure 3A).** A change

in phenotype was observed by 24 hours when the iPSC-DP T cells appeared to be primarily

CD8sp initially but shifted dramatically to become CD4sp 4 days after treatment. After 6 days,

more than 50% of the cells were CD4sp at the lowest concentration of PMA/I **(Figure 3B)**. Medium

to high concentrations of PMA/I were also effective in converting iPSC-DP T cells to CD4sp T

cells albeit at lower efficiency. The cells remained DP in the absence of PMA/I. Commitment to

the CD4 lineage is typically specified by the zinc-finger transcription factor and master regulator

ThPOK [28]. At the lowest concentration of PMA/I, ThPOK could be observed in the

$CD3^+TCR\alpha\beta^+CD4^+$ population of developing T cells **(Figure 3C)**. Without potent PMA/I

stimulation, the DP T cells did not progress towards the CD4 lineage and the minor population of

$CD4^+$ cells did not express ThPOK. The yield of CD4sp cells from PMA/I treated iPSC-DP T cells

was approximately 4.3e3±2.3e3/input iPSCs **(Figure 2)**. The generated iPSC-CD4sp cells

displayed a mostly T_{EM} phenotype ($CD45RO^+CD62L^-$) and highly expressed the co-receptor CD28

similar to primary CD4sp (**Figure 3D**).

After successfully generating bona fide CD4sp T cells from iPSCs, we next interrogated whether

a regulatory phenotype could be induced to produce Tregs. Previous studies have shown the

ability to generate $FOXP3^+$ Tregs from $CD4^+CD25^-$ T cells, termed induced Tregs (iTregs), in the

presence of $TGF\beta$ and IL-2 [29, 30 31]. To determine if the iPSC-CD4sp T cells could also be induced

to express FOXP3, cells were treated with a commercially available cocktail of $TGF\beta$, all-trans

retinoic acid (ATRA), and IL-2 and supplemented with a soluble T cell activator composed of anti-

human CD3, CD28, and CD2 monospecific antibody complexes on day 50 of differentiation

(Figure 3A). Seven days post-treatment (day 57) with the $TGF\beta$ cocktail and soluble activator, an

increase in frequency of DP and CD8sp T cells was observed and, while reduced, the culture

retained a sizeable CD4sp T cell population **(Figure 3E)**. When gating on the CD4sp T cell

population the cells expressed CD25, low levels of CD127 and high levels of FOXP3 similar to

primary Tregs. Approximately 26% (+/-9.2%; n=6) of the total live population displayed a ≥90%

$CD4spCD25CD127lowFOXP3^+$ phenotype. In addition, the CD4sp T cell population maintained

expression of ThPOK while co-expressing FOXP3 **(Figure 3E)**. We observed an average yield of

5.9e3±4.4e3 (n=3) iPSC-Treg/input iPSC **(Figure 2)**.

iPSC-CD4sp T cells and iPSC-Tregs display a molecular signature similar to that of

primary CD4sp and Tregs respectively

As the populations of iPSC-CD4sp cells and iPSC-Tregs are heterogenous post-PMA/I and post-

TGFβ and ATRA treatment, we sought to isolate the CD4sp population at day 50 and the target

Treg population of CD4spCD25$^+$CD127$^-$FOXP3$^+$ iPSC-Tregs at day 57 by FACS to further

characterize them by single-cell RNA sequencing (scRNA-seq). For iPSC-Tregs, a population of

target cells that is >90% FOXP3 was obtained by gating on a "CD25 high" population from the

total population of live, single CD4sp T cells **(Supp Figure 3)**.

To determine the similarity between the iPSC-CD4sp to primary CD4 T cells and the iPSC-Tregs

to primary Tregs, their transcriptomes were compared. Four cell samples were analyzed: (1)

sorted iPSC-Tregs; (2) primary Tregs; (3) induced Tregs (generated from primary naïve CD4 T

cells); (4) iPSC-CD4sp; and (5) primary CD4 Tconvs. A global analysis of differentially expressed

genes by hierarchical clustering revealed that iPSC-CD4sp T cells were similar in their

transcriptional profile to primary Tconv while primary Tregs were similar to induced Tregs. The

iPSC-Tregs however, were distinct in their gene expression profile **(Figure 4A).** We sought a more

refined analysis and examined the expression of a core Treg gene signature in all samples [32-34]

(Figure 4B). We found that the iPSC-Tregs were similar to primary Tregs and induced Tregs with

some differences in gene expression. The iPSC-Tregs had upregulated expression of *FOXP3 and*

CTLA4 and downregulated expression of *TNFRSF1B*. Within the core Treg gene signature, iPSC-

CD4 T cells were less similar to primary Tregs and more similar to primary Tconv. Both primary-

and iPSC-CD4sp T cells upregulated *ANK3* expression and downregulated *TNFRSF1B*

expression compared to all other populations. However, the iPSC-CD4sp T cells expressed

several genes at a much higher level than their primary counterpart (*VAV3, PTPRK, IKZF2,*

HPGD, RBMS3).

Characterization and functional assessment of iPSC-CD4sp T cells and iPSC-Tregs

Further characterization and functional assessment were performed on sorted iPSC-CD4sp T

cells and compared to their primary Tconv counterpart. iPSC-CD4sp T cells were activated with

CD3/CD28/CD2 soluble activator. After activation, cells increased their expression of activation

markers CD25 and PD-1 at 24 hours **(Supp. Figure 4A and Supp. Figure 4B)** and at 48 hours

(Figure 5A and Figure 5B) while decreasing expression of CD28 similar to Tconv **(Supp. Figure**

4B and Figure 5B). While CD69 MFI increased after activation, the frequency of cells expressing

CD69 did not change. Activated iPSC-CD4sp T cells also secreted pro-inflammatory cytokines IL-

2 and TNF-α similar to activated primary Tconv but did not secrete IFN-γ. Similar levels of IL-4

and IL-10 were secreted by iPSC-CD4sp T cells compared to primary Tconv after stimulation

(Figure 5C).

We then characterized and examined the *in vitro* function of sorted iPSC-Tregs.

Immunophenotying of cells post-sort showed a typical Treg phenotype with an average of

83.4±6.3% FOXP3 expression (n=5) and average 91.6±1.6% Helios expression (n=2) from total

live cells.   Demethylation of the evolutionarily conserved non-coding element FOXP3 Treg-

specific demethylated region (FOXP3 TSDR) is required for stable FOXP3 expression in Tregs [35].

It is typically methylated in Tconv, induced Tregs and other non-Treg blood cells [36]. We examined

the methylation state of FOXP3 TSDR by high resolution melt quantitative polymerase chain

reaction (HRM-qPCR) in the sorted iPSC-Tregs. The iPSC-Tregs demonstrated >90%

demethylation similar to the primary Treg positive control. In contrast, primary CD4$^+$CD25$^-$ Tconv

as well iPSC-CD4sp T cells were highly methylated at FOXP3 TSDR **(Figure 5D)**. The sorted

iPSC-Tregs could be activated via their TCR using beads coupled to anti-CD3 and anti-CD28

antibodies (Dynabeads). After activation, the iPSC-Tregs increased the expression of early

activation marker CD69 on average of 3-fold with no change to their Treg phenotype **(Figure 5E-**

G).

Tregs typically exert their suppressive function through various mechanisms including

suppression by cytolysis [37, 38]. Cellular toxicity was determined by co-culturing activated and

unactivated iPSC-Tregs with primary CD4 Tconv target cells. Compared to activated cytotoxic

primary CD8 T cells, activated iPSC-Tregs had low cytolytic activity similar to activated primary

Tregs as measured by a LDH release assay **(Figure 6A).**

Additional suppressive mechanisms include secretion of anti-inflammatory cytokines like IL-10 [37].

IL-10 is produced by a variety of cells including $CD4^+$ Th2 cells, Tregs and IL-10-induced $CD4^+$ T

reg cells (Tr1 cells) [39]. Interestingly, iPSC-Tregs express higher, but variable, amounts of IL-10 as

compared to primary Tregs. Importantly, the iPSC-Tregs secreted low levels of pro-inflammatory

cytokines IL-2, IFN-γ, and TNF-α before and after activation, similar to primary Tregs **(Figure 6B)**.

The cardinal feature of Tregs is their ability to suppress the activation and proliferation of Tconv

or Teff cells by a cell-to-cell contact mechanism [37]. To determine the suppressive function of

activated iPSC-Tregs, the cells were co-cultured with activated primary Tconv and their

proliferation was measured by flow cytometric analysis. Activated iPSC-Tregs effectively

suppressed Tconv proliferation at a 1:1 ratio of Tregs:Tconv similar to activated primary Tregs

(Figure 6C). Without activation, iPSC-Tregs had significantly lower levels of suppression at 1:1

to 1:4 Tregs: Tconv ratios. An iPSC-Treg dose-dependent effect on Tconv suppression was also

observed similar to primary Tregs. As expected, when iPSC-CD4sp T cells were tested for

suppressive function, they were unable to prevent Tconv proliferation even at the highest ratio of

1:1 CD4sp T cells:Tconv **(Figure 6C)**. This confirms that the suppressive function is specific to

iPSC-Tregs and appears only after conversion of iPSC-CD4sp T cells to $FOXP3^+$ Tregs and after

cell activation.

Generation and functional assessment of HLA-A2 CAR^+ iPSC-Tregs

We next investigated the function of iPSC-Tregs equipped with a chimeric antigen receptor (CAR).

To this end, we genetically engineered iPSCs to express an HLA-A2 CAR using zinc-finger

nucleases (ZFNs) targeting exon 2 of the *TRAC* gene.

We sought to retain expression of the *TRAC* gene and utilized a donor sequence comprised of

TRAC cDNA encompassing the *TRAC* exonic sequence downstream of the integration site. This

donor sequence is placed upstream and in-frame with a T2A coding sequence such that the

engineered locus retains expression of the TCRα chain while expressing an HLA-A2 CAR under

the control of the *TRAC* promoter **(Figure 7A)**. Through optimized electroporation of the iPSCs

with TRAC ZFN mRNA and donor DNA we achieved a >80% targeted integration without selection

as assessed by sequencing of the bulk edited population.

The resulting engineered iPSCs were cloned, expanded, and sequenced to obtain a biallelically

modified iPSC clone with the HLA-A2 CAR inserted at the target site. Sequencing analysis

confirmed the precise and in-frame insertion of a single copy of donor DNA (HLA-A2-CAR),

without any detected mutations. The CAR-engineered iPSCs also displayed a normal pluripotent

phenotype and were karyotypically normal **(Supp. Figure 1)**. To generate HLA-A2 CAR[+] iPSC-

Tregs (CAR iPSC-Tregs) the iPSCs were differentiated to HSPCs, lymphoid progenitor cells, and

DP T cells using the same differentiation process utilized for unedited iPSCs as described above

(Supp. Figure 5A). HLA-A2 CAR iPSCs could also generate CD4sp T cells and FOXP3[+] Tregs

(Supp. Figure 5B, 5C). Of note, we observed a 2.2-fold lower yield of Tregs (n=3, ±0.2 fold) from

the HLA-A2 CAR iPSCs compared to Tregs generated from unedited. By flow cytometric analysis,

we found that ~90% (90.1%±1.8%, n=4) of the DP T cells were already expressing the CAR by

day 42 and retained high CAR expression at both the CD4sp T cell (99%±0.2%, n=4) and Treg

stages (99.85%±0.1%, n=4). The intensity of CAR expression increased as the cells differentiated

as measured by MFI **(Supp. Figure 5D)**.

Similar to unedited iPSC-Tregs, HLA-A2 CAR⁺ iPSC-Tregs could also be sorted. Four days post-

sort the expression levels of the CAR remained stable (81.3±12.4% of TCRαβ⁺ cells and

MFI=9814.3±1143.3, n=3) and the cells maintained a high level of demethylation at FOXP3 TSDR

(88%±10.3%; n=3).

Further characterization showed that the CAR iPSC-Tregs could be activated through both the

TCR and the HLA-A2 CAR demonstrating the engineering strategy successfully retained TRAC

expression while integrating a functional CAR. Activation of the CAR iPSC-Tregs using either

CD3/CD28 beads (Dynabeads) or HLA-A2 dextramer upregulated CD69 expression compared to

unactivated or dextramer control activation **(Supp. Figure 6A, 6B)**. TCR- and CAR-mediated

activation induced a very low level of pro-inflammatory cytokines while inducing higher levels of

IL-10 **(Supp. Figure 6C)** similar to levels of cytokines produced by unedited iPSC-treg upon TCR

activation **(Figure 6B).** The CAR iPSC-Tregs significantly suppress Tconv proliferation at similar

levels to wild-type iPSC-Tregs when activated either through the TCR or the HLA-A2 CAR, while

unactivated or control dextramer activated CAR iPSC-Tregs only displayed a basal level of

suppression **(Figure 7B).**

Characterization and functional assessment of cryopreserved CAR iPSC-Tregs

A requirement for a viable CAR iPSC-Treg therapy is the ability to efficiently cryopreserve and

store the cells for future use without loss of stability and function. Towards that goal, we

cryopreserved iPSC-Tregs post-sort and then thawed, characterized, and tested their suppressive

function. Approximately 80% of cells were recovered post-thaw (avg=82.5%, n=2) with a viability

comparable to pre-cryopreservation (avg=74%, n=2). The CAR iPSC-Tregs maintained their

phenotype post-thaw **(Supp. Figure 7A)**. The cells were also highly demethylated at FOXP3

TSDR (95.5±0.5%, n=2) and displayed an average 4-fold increase in CD69 after activation *via*

the HLA-A2 CAR **(Supp. Figure 7B).** Finally, thawed CAR iPSC-Tregs were able to suppress

Tconv proliferation when activated *via* the TCR or CAR **(Figure 7C).**

Discussion

We report here for the first time a serum- and feeder-free method to produce CD4sp T cells and functional Tregs with and without a CAR from iPSCs. While the generation of CD4sp T cells from iPSCs has been reported [8, 10], it currently requires the use of murine stromal cells, animal serum (fetal bovine serum, FBS) and ATOs. The use of xenogenic feeders, animal serum and 3D organoids is not ideal for the development of a clinical-grade therapeutic product. This is due to issues with safety, scalability and serum batch-to-batch heterogeneity as well as difficulty dissociating cells from 3D organoids. In addition, automation of 3D organoid generation to produce large-scale reproducible organoids is a nascent field and will require further optimization to minimize variability across differentiation batches [40]. The method proposed here circumvents many of these issues and increases its amenability for manufacturing cell therapies at scale.

The reprogramming of T cells to pluripotency and subsequent differentiation to mature T cells has been well documented particularly towards the generation of $CD8\alpha\beta^+$ CTLs [41, 42, 43]. Using T cell-derived iPSCs may aid in more efficiently generating iPSC-T cells as the early stages of T cell development that are characterized by failed TCR rearrangement and cell death are bypassed [44]. While antigen specificity is unknown, reprogramming from a T cell clone possessing a single TCR [7, 16, 45] or introducing a CAR or transgenic TCR at the iPSC or T cell stage [6, 9, 17, 46-48] can help confer specificity. Based on these findings, we sought to utilize a similar strategy to develop CD4sp T cells and Tregs from iPSCs. We were able to successfully obtain iPSCs from mature CD4sp T cells that expressed classic pluripotency markers and appeared as compact colonies with well-defined borders. The iPSCs were amenable to genome engineering for the targeted integration of an HLA-A2 CAR. Both the wild-type and engineered iPSC lines were karyotypically normal **(Supp. Figure 1)**. We did not perform deep characterization of off-targets in edited iPSCs at this research-grade stage. Most importantly, an advantage of using iPSCs over primary cells is

the ability to select through whole-genome sequencing (WGS) well-characterized clones with no

specific off-targets and expand them to generate a MCB for future clinical use.

Methods to differentiate iPSCs into HSPCs and progenitor T cells are readily available, but many

require the combination of multiple individual reagents from various sources which may contribute

to increased variability in differentiation. We found that utilizing commercially available reagents

specifically designed to generate HSPCs and T cell progenitors from iPSCs streamlined the

differentiation process while satisfying the serum- and feeder-free requirements. We utilized a

modified EB-based method to differentiate iPSCs and employed commercially available serum-

free media for the generation of $CD34^+CD45^+$ HSPCs capable of differentiation to lymphoid cells.

It is worth noting that while the process yielded a mixture of $CD34^+$ and $CD34^-$ cells, no enrichment

or purification for $CD34^+CD45^+$ target cells was required for further differentiation. Indeed, we

found that by separation of the free-floating progenitor cells from the main EB we were able to

generate lymphoid progenitor cells that co-expressed CD5 and CD7 marking their T cell

commitment [49, 50]. Continued differentiation of the lymphoid progenitor cells resulted in CD4 and

CD8 DP T cells again, without enrichment of the starting population.

Thymic DP T cells express both CD4 and CD8 co-receptors that promote TCR engagement of

MHC ligands to enhance signal transduction [24]. Through various strengths of binding affinity to

MHC ligands, the developing T cells will undergo positive selection to survive or cell death by

negative selection [51]. The step-by-step natural process followed by DP T cells after positive

selection to commit to either the CD4sp or CD8sp lineage is unclear. [24]. It has been suggested

that a persistent TCR signal is required to make the CD4 choice and block re-expression of the

CD8 co-receptor. To mimic the signal required for positive selection and commitment to the CD4

lineage at the DP T cell stage, we utilized a strategy previously shown to induce CD4 commitment

in murine DP T thymocytes without TCR engagement [26]. The addition of PMA/I activates protein

kinase c and elevates calcium levels, respectively and in a similar manner to TCR-mediated

activation with signals from co-stimulator molecules like CD4, CD8 and LFA-1 [26, 52]. At levels

between 0.016 uM to 0.16 uM of PMA and 0.13 uM of ionomycin, TRAC KO murine DP T

thymocytes are positively selected to CD4 T cells. Higher concentrations of PMA/I (1.6 uM and

1.3 uM, respectively) were inefficient in generating CD4 T cells. These findings are consistent with

our method in that the lower concentrations we used successfully converted cells to the CD4

lineage while higher concentrations of PMA/I were less efficient. Importantly, after conversion to

CD4sp T cells, ThPOK expression was observed further confirming its lineage commitment. The

kinetics of co-receptor expression in our system is different from what has been previously

published in mouse-derived thymocytes [26, 52, 53] and could be due to species-specific differences

between mouse and human thymocyte development [54], differences in the systems used (iPSC-

derived vs. thymus-derivates DP T cells) or related to the transient expansion of a highly

proliferative, immature intermediate (CD4$^-$ CD8$^+$ Mki67$^+$) [54]. Further characterization of the

intermediate stages will shed light on the mechanisms of PMA/I in our system.

Initial characterization of the iPSC-CD4sp T cells shows that they have a predominantly T_{EM}

phenotype, can be activated and secrete pro-inflammatory cytokines (IL-2 and TNF-α) upon

stimulation. However, we did not detect IFN-y from the iPSC-CD4sp T cells but observed IL-10

and IL-4 cytokines at similar levels to primary Tconv. The lack of IFN-y could suggest that the

iPSC-CDsp T cells are more similar to Th2 cells which are characterized by their secretion of IL-

4 and IL-10 and lack of IFN-y production when compared to Th1 cells [55] [56] (Yano et al. ASH 2019).

Infusion of a defined CD4:CD8 CAR T cell ratio allows for a potentially more potent and

predictable product [12]. While some studies demonstrate that naive CD4sp CAR-T in combination

with CD8sp CAR T cells confer higher antitumor responses than T_{EM} CD4sp in *in vivo* mouse

models [13], others show potent antitumor activity using a mixture of naive and effector like CD4sp

CAR T cells in humans [12]. As the field matures to identify the characteristics of an ideal CD4sp

product, further optimization of our protocol can fine tune the final phenotype of our iPSC-CD4sp.

In addition to maintaining a serum- and feeder-free differentiation system, the use of PMA/I to

generate iPSC-CD4sp T cells is an especially critical finding as providing a positive selection

signal towards the CD4 lineage is the first step to generating Tregs. After maturation, the iPSC-

CD4sp T cells were treated with TGFβ, ATRA, IL-2 and a soluble CD3/CD28/CD2 activator to

convert them to a FOXP3⁺ Treg phenotype. The generation of Tregs from naïve CD4 T cells using

TGFβ is well-documented [31]. While TGFβ-induced iTregs express FOXP3, they are only partially

demethylated at FOXP3 TSDR and are unstable in their expression of FOXP3. In addition,

polyclonal iTregs do not always display a regulatory phenotype, are not always suppressive and

can produce high levels of inflammatory cytokines [57]. In contrast, the iPSC-Tregs generated within

our system were highly demethylated at FOXP3 TSDR, suppressed Tconv proliferation and did

not produce inflammatory cytokines despite utilizing TGFβ towards its conversion. Transcriptomic

analysis by scRNA-sequencing further confirmed the identity and similarity of the iPSC Tregs to

primary Tregs. The iPSC-Tregs expressed typical markers of their primary natural Treg

counterpart, namely highly expressing *FOXP3*, *IKZF2* and *IL2RA* in the absence of activation.

The iPSC-Tregs also expressed a previously published core Treg signature [32-34] similar to that of

primary Tregs and less similar to iTregs and Tconv. Differential expression of genes between

iPSC-Treg and primary Treg, such as *IL7R* and *ID2*, could be due to culture conditions or to

differences between cells that differentiated *in vitro* as compared to *ex vivo* adult cells. Finally,

since scRNAseq utilizes relative expression levels, measuring the absolute expression of these

markers by bulk RNAseq may provide better quantitative results to distinguish these populations.

Previous studies have shown that iPSCs derived from T cells express TCRαβ prematurely which

interferes with DP T cell formation and can be overcome by supplying the cells with a strong Notch

signal [9]. Constitutive expression of CAR with simultaneous expression of the endogenous TCR,

however, offsets the effects of a stronger Notch signal. In this case, both elimination of TRAC

expression as well as attenuation of the CAR signal was required to restore DP T cell generation

[9]. These are important considerations for the generation of a scalable cell therapy product using

engineered iPSCs. While we were able to generate DP T cells from both wild-type and engineered

iPSCs, the exact Notch ligand in the LDCM reagent is unknown making the need to test a known

Notch ligand necessary for improving DP T cell yields. In addition, we observed a lower yield of

HLA-A2 CAR[+] iPSCs-Tregs as compared to Tregs generated from unedited iPSCs, supporting

the idea that premature TCR expression and CAR signaling influences DP T cell yields and

subsequent Treg generation. Defining the Notch ligand, optimizing CAR signaling as well as

applying compound-based measures to improve DP T cell yield from iPSCs can aid in the

generation of a more robust and reproducible differentiation protocol [9].

Cryopreservation is key to the development and delivery of successful cell therapy products. We

found that our CAR iPSC-Tregs were amenable to cryopreservation and showed similar viability

pre- and post-cryopreservation with approximately 80% recovery post-thaw. The phenotype of the

cells was also maintained pre- and post-thaw with appreciable upregulation of CD69 post-

activation via the CAR or TCR. Importantly, the cryopreserved iPSC-Tregs suppress Tconv

proliferation in vitro confirming that the cells maintain functionality **(Figure 7)**.

Further characterization of the iPSC-Tregs and iPSC-CD4sp T cells is still required, including

evaluation of its ability to expand to clinically relevant numbers and its function in a GvHD

mouse model. While additional data are essential to develop a viable cell therapy, we

demonstrated here a method to generate functional and stable iPSC-derived Tregs and CD4sp

T cells that removes the inconsistencies and safety issues caused by using xenogenic

materials. iPSCs can serve as an endless supply of CD4sp T cells and Tregs with and without a

CAR for therapeutic applications and our study supports further development of this process for

clinical use in oncology and autoimmune diseases.

Methods

Generation of human induced pluripotent stem cell lines. Previously cryopreserved CD4 T cells were used for reprogramming to pluripotency. CD4 T cells were isolated from mobilized leukapheresis material from a healthy male donor (AllCells, LLC) with informed written consent. The raw material was washed several times to remove platelets, labeled with CD4 microbeads (Miltenyi Biotec) and then isolated using the Miltenyi CliniMACS system. Isolated CD4 T cells were analyzed for purity prior to cryopreservation. To prepare for reprogramming, the CD4 T cells were thawed and cultured overnight in RPMI + 10% human AB serum (huABS, Valley Biomedical) and activated 1:1 with CD3/CD28 Dynabeads in 100 U/mL of IL-2 (Thermofisher). The next day, the CD4 T cells were debeaded and sorted for CD4$^+$CD25$^+$ cells and recovered overnight in RPMI + 10% huABS with 100 U/mL of IL-2. One day post-sort, the sorted CD4 T cells were reprogrammed with Sendai virus carrying the four transcription factors OCT, SOX2, KLF4, and C-MYC as previously reported using Cytotune iPS-2.0 Sendai Reprogramming Kit (Thermofisher; Fusaki Proc Japan Academy 2008). After transduction, cells were recovered in RPMI + 10% huABS for 24 hours and transferred to ReproTeSR media (StemCell Technologies) on Matrigel-coated plates (Corning). Cells were fed daily or every day until the appearance of colonies by day 14. Colonies were examined for appropriate morphology then manually selected for further expansion and transition to mTeSR1 media (StemCell Technologies)

Characterization of human induced pluripotent cells lines. Immunocytochemical analysis for pluripotency of iPSC lines was performed by washing cells in 1X PBS (Thermofisher), fixed in 4% paraformaldehyde for 20 minutes at 4°C, and blocked in 10% normal goat or rabbit serum with 0.5% Triton X-100 (Sigma) in PBS. Immunocytochemical analysis was performed using primary antibodies and corresponding AlexaFluor-conjugated secondary antibodies (Thermofisher). All antibodies were diluted and used according to manufacturer's instructions. Cell nuclei were

counterstained with DAPI in some experiments [58]. Images were collected using an EVOS Cell

Imaging System (Thermofisher). Karyotype analyses were performed by Cell Line Genetics.

iPSC culture and clonal expansion. iPSCs were maintained on Matrigel (Corning) in mTeSR1

media according to manufacturer's instructions (StemCell Technologies). For passaging, iPSC

colonies were detached briefly with either Accutase and manually scraped or with Gentle Cell

Dissociation Reagent. Cells were routinely passaged at a 1:8-1:10 split ratio in the presence of Y-

27632 (ROCK inhibitor, Tocris) for the first day after passaging and removed for subsequent media

changes.

For clonal expansion of transfected iPSCs, a method of low-density passaging was used. After

electroporation, iPSCs were passaged at a low seeding density (1e3, 2.5e3, 5e3 cells per well of

a 6-well plate) such that a single iPSC would give rise to a single colony in a Matrigel-coated 6-

well plate. iPSC colonies were then manually picked and expanded individually. Tightly packed

colonies with clearly defined borders were then collected for genomic DNA isolation and

subsequent screening for targeted integration.

Differentiation of HSPCs from iPSCs. iPSCs were differentiated as previously described with

modifications [19, 20]. Embryoid bodies were generated from iPSCs in APEL2 media (StemCell

Technologies) and cultured in non-tissue culture or ultra-low attachment 96-well round bottom

plates in the presence of 10 ng/mL BMP4, 10 ng/mL VEGF, 50 ng/mL SCF, 10 ng/mL bFGF, and

10 uM Y-27632 on day 0 (all cytokines from Thermofisher). EBs were then cultured in APEL2

media in the presence of 20 ng/mL VEGF, 10 ng/mL bFGF, 100 ng/mL SCF, 20 ng/mL FLT-3L, 20

ng/mL TPO and 40 ng/mL IL-3 from day 1 to day 6 with a media refresh every 2 days (all cytokines

from Thermofisher). On day 8, EBs were switched to complete StemPro-34 media (Thermofisher)

and supplemented with 100 ng/mL SCF, 20 ng/mL FLT3-L, 20 ng/mL TPO, and 40 ng/mL IL-3 in

order to expand developing HSPCs. Media was refreshed every 2 days until day 14. The EBs

were then collected and passed through a 70 uM cell strainer. The resulting cell suspension

containing the HSPCs was then used for downstream differentiation.

Differentiation of T cell progenitors from iPSC-HSPCs. StemDiff T cell kit (StemCell

Technologies) was used to differentiate HSPCs to lymphoid progenitor cells and to DP T cells.

Manufacturer's instructions were followed unless otherwise indicated. HSPCs were seeded at

50,000 cells/well of a 6-well plate coated with lymphocyte differentiation coating material (LDCM)

and cultured in StemSpan Lymphoid Progenitor Expansion Media to differentiate the HSPCs to

CD5+CD7+ lymphoid progenitor cells. After 14 days of culture (differentiation day 28), resulting

iPSC-lymphoid progenitor cells were harvested and seeded at 0.5e6/well of a 6-well plate with a

fresh coating of LDCM. The cells were cultured in StemSpan T Cell Progenitor Maturation Media

(maturation media) for differentiation to CD4+CD8+ DP T cells. After 14 days of culture

(differentiation day 42), the DP T cells were harvested for further differentiation.

Generation of CD4-single positive T cells and Tregs from iPSC-DP T cells. Induction of DP

T cells to CD4sp T cells was performed on LDCM in maturation media. On differentiation day 42,

a cell stimulation cocktail composed of phorbol 12-myristate 13-acetate and ionomycin was added

to the cultures at three different concentrations from the 500X stock concentration: low (0.01 uM

PMA, 0.16 uM I), medium (0.02 PMA uM, 0.335 uM I), and high (0.0405 uM PMA, 0.67 uM I). A

full media change was performed 24 hours later by harvesting the entire cell suspension, replating

into fresh maturation media and seeding at a 1:1 ratio onto new LDCM coated 6-well plates. The

cells were then fed every 3-4 days for an additional week until day 50. On day 50, the cells were

then treated with 20 uL/mL of Immunocult Human Treg Differentiation Supplement (StemCell

Technologies) containing recombinant human TGF-B1 and all-trans retinoic acid, 12.5 uL/mL of

Immunocult Human CD3/CD28/CD2 T cell activator (StemCell Technologies) and 10 ng/mL of

CTS IL-2 (Thermofisher). Media in the wells for conversion to Tregs was composed of a 50%

mixture of complete CTS OpTmizer and maturation media. A half media change was performed

on day 54 with a 50% mixture of CTS OpTmizer and maturation media supplemented with 10

ng/mL of IL-2.

Flow cytometry, sorting and antibodies. Flow cytometry was performed in FACS buffer

composed of PBS, 0.5% BSA, and 1 mM EDTA. For extracellular staining, antibodies were added

to cells according to manufacturer's instructions, incubated at 4°C for 20 minutes, and washed

before analysis. For intracellular staining, cells were first stained with extracellular markers and

fixed and permeabilized using the eBioscience FOXP3/Transcription Factor Staining Buffer set

(Thermofisher). Antibodies were then added according to manufacturer's instructions. Attune NxT

flow cytometer (Thermofisher) and Sony Sorter SH800 was used for flow cytometric analysis and

sorting, respectively. Antibodies (clones in parentheses) used for surface and intracellular staining

of the following molecules were obtained from Biolegend: CD45 (HI30), FOXP3 (206D), ThPOK

(11H11A14), TCRab (IP26), Helios (22F6), and CD127 (A019D5). Antibodies against CD34 (581)

and CD3 (UCHT1) were obtained from BD Biosciences. Antibodies against CD4 (OKT-4), CD4

(VIT4), CD8 (SK1), FOXP3 (PCH101), CD69 (REA824), CD25 (2A3), CD28 (S20013F), CD45RA

(HI100), CD45RO (UCHL1), CD27 (O323), CD62L (DREG-56), and PD-1 (EH12.1) were obtained

from Thermofisher. CD5 (UCHT1) and CD7 (CD7-687) antibodies were obtained from Miltenyi.

Antibodies to SOX2, OCT4, SSEA-4 were obtained from R&D Systems. TRA-1–81 was obtained

from Santa Cruz Biotechnology. Alexa-conjugated secondary antibodies and DAPI were obtained

from Thermofisher.

scRNA-sequencing RNA extraction, library generation, and sequencing analysis. RNA from

iPSC-Tregs, iPSC-CD4sp, primary Tregs, induced Tregs and primary Tconv was extracted using

an RNA extraction kit (Qiagen) following differentiation (iPSC-Tregs and induced Tregs) or

isolation and culture (primary Tregs and Tconv).

Single cell RNA-sequencing libraries were sequenced on a NextSeq 500. Samples had an

average depth of 18636 reads per cell. Read quality control, UMI counting, barcode counting, and

alignment to the GRCh38 GENCODE v32/Ensembl 98 reference genome from 10x Genomics

were performed using the "cellranger 6.0.2" pipeline. All downstream analysis were performed

using the Seurat packages in R for single-cell RNAseq analysis.

All samples were filtered on the following measures: number of unique genes detected, total

number of molecules detected, and level of expression of mitochondrial reads before integrating

the datasets. We removed cells in which greater than 20% mitochondrial reads, less than 200

feature counts, or greater than 6500 feature counts were detected.

The data was then normalized using the Seurat normalization function which divides the feature

counts by total counts for each cell, then multiplies the resulting fraction by a scale factor of

1.00e+4, and, lastly, log-normalizes this value.

Dimensionality reduction was performed using Seurat packages and utilizing principal

components 1 through 50 in the reduction. Clusters were determined at a resolution of 0.3. The

identity of each cluster was validated by evaluation of gene expression within each cluster of cell-

type specific markers for the cell-types of interest.

Assessment of iPSC-CD4sp T cell and iPSC-Treg Function. For assessment of Tconv or iPSC-

CD4sp T cell phenotype and activation, cells were sorted and cultured in X-VIVO15 (Lonza) +

10% huABS (Valley Biomedical). Cells were either left unactivated or activated with Immunocult

Human CD3/CD28/CD2 T cell activator per manufacturer's instructions (StemCell Technologies)

for 24 or 48hrs. Cells were then stained with indicated antibodies and run on flow cytometry as

previously described. For assessment of cytokine secretion, unactivated or activated iPSC-

CD4sp T cells were seeded at 5e4 cells/well in a round-bottom 96 well plate in 200 uL of media.

The supernatant was collected 24 hours later.

For assessment of iPSC-Treg function,cells were sorted on day 57 and cultured in complete

OpTmizer media supplemented with 1000U IL-2/mL (Thermofisher) for three days prior to

functional analysis. iPSC-Tregs were then activated with CD3/CD28 Dynabeads (Thermofisher)

at a ratio of 1:1 in media composed of X-VIVO15 (Lonza) + 10% huABS (Valley Biomedical) +

Penicillin / Streptomycin solution (pen/strep). For analysis of cytokine production, 5e4 activated

and unactivated cells were seeded in a round-bottom 96-well plate in 200 uL of media. One day

post-activation, supernatant was collected and analyzed for cytokines using human cytokine

assay X (Meso Scale Discovery). For assessment of Tconv suppression, activated and

unactivated iPSC-Tregs were co-cultured with activated (2:1 beads:cells) Tconv labeled with Cell

Trace Violet (Thermofisher) in X-VIVO15 media supplemented with pen/strep. iPSC-Tregs were

initially plated at 5e4/well in a 96-well round-bottom plate and serially diluted by a factor of 2 down

to 781.25 cells/well. 2.5e4 activated, CTV-labeled Tconv were then added to each serially diluted

well of iPSC-Tregs. Baseline Tconv proliferation was determined by culturing 2.5e4 Tconv in the

absence of iPSC-Tregs. Three days after co-culture, cells were collected, labeled for viability and

assessed for Tconv proliferation via Cell Trace Violet dye by flow cytometric analysis. For

assessment of iPSC-Treg cytolytic activity, activated and unactivated iPSC-Tregs were cocultured

with activated Tconv. 5e4 iPSC-Tregs were seeded with 5e4 primary Tconv and cultured for 3

days in X-VIVO15 media supplemented with pen/strep. The supernatant was then collected and

analyzed using the CyQuant LDH Cytotoxicity Kit (Thermofisher) following the manufacturer's

instructions. Samples were read on a SpectraMax microplate reader (Molecular Devices). Primary

Treg, Tconv and CD8 controls were cultured for 7-9 days prior to activation and assessment of

function.

FOXP3 TSDR demethylation assay. The Treg-specific demethylated region (TSDR)

hypomethylation status of FOXP3 gene was measured using a post-PCR High Resolution Melt

(HRM) analysis as previously described [4]. Briefly, genomic DNA was extracted from cell pellets

containing at least 1e6 cells/pellet using the DNeasy Blood and tissue kit (Qiagen), followed by

bisulfite conversion using the Epitech Fast Bisulfite Conversion Kit (Qiagen). The FOXP3 TSDR

regions were then amplified from the converted sample gDNAs by qPCR using specific primers

(forward: TTGGGTTAAGTTTGTTGTAGGATAG, reverse: ATCTAAACCCTATTATCACAACCCC,

Sigma Aldrich, USA) and Precision Melt Supermix (Biorad). The melting curves of amplified

product DNA fragments were quantified against the methylation standard curve generated using

commercial bisulfite converted human genomic DNAs (Qiagen) and expressed as a mean

methylation percentage.

Cytokine MSD methods. The concentrations of cytokines were measured with MSD U-Plex kits

(#K15067L-1, MSD, USA) following the manufacturer's protocol. Briefly, biotinylated capture

antibodies were incubated with specific linkers and then the mixture of linker coupled antibodies

was added to MSD U-Plex plates. After 1 hour incubation and wash step, standard curves and

samples were added to the plates. The plates were washed again after the sample incubation. A

mixture of SULFO-Tag conjugated detection antibodies was added to the wells and incubated.

After the final wash step, MSD GOLD Read Buffer B was added to the plate followed by reading

the electrochemiluminescence signals in a MESO SECTOR S 600 instrument.

ZFN and donor DNA constructs. The zinc finger nuclease (ZFN) pair targets the *TRAC* locus,

encoding for the alpha chain of the TCR, at exon 2. The ZFP backbone and FokI variants were

generated by reassembling zinc finger arrays from modules as previously described [59] and cloning

the resulting zinc finger array into a vector containing the desired FokI variant. All constructs were

verified by Sanger sequencing with two sequencing primers to yield overlapping sequence reads

of the entire ZFN coding sequence [60]. mRNA was transcribed from a PCR template using the 5×m

Message mMachine T7 ULTRA kit (Invitrogen) according to the manufacturer's instructions as

previously reported [60].

We designed and cloned the donor DNA sequence into a customized AAV vector containing AAV

inverted terminal repeats (ITRs). An AAV vector was initially selected to use AAV as DNA delivery

system but we ultimately optimized delivery using single strand DNA. The pAAV-partial-TRAC-

HLA-A2 CAR contains 110bp of genomic partial TRAC sequence (amplified by PCR) flanking the

TRAC homologous donor templates in frame with the second exon of TRAC, followed by a self-

cleaving T2A peptide, the HLA-A2 CAR and a bGHpA sequence.

Transfection of TRAC-ZFN mRNA and donor DNA. All transfections were performed on the

Maxcyte ATx (model GT) electroporation system. Approximately 3e6 iPSCs were used for

electroporation in an OC-100 processing assembly (Maxcyte). iPSCs were treated with Accutase

until cells were fully detached (~3min). After collection, cells were pelleted and washed once with

Maxcyte electroporation buffer. The iPSCs were then resuspended in Maxcyte electroporation

buffer at a concentration of 30e6 cells/mL and mixed with ZFN mRNA (10 ng/ul) and 8 ug of donor

DNA. The cell and nucleic acid mixture was dispensed into the OC-100 assembly and immediately

placed into the Maxcyte instrument using Optimization program 9. iPSCs were then recovered for

20 minutes at 37°C in the assembly and then transferred to Matrigel-coated plates in mTeSR1

media. Y-27632 was added to the media for 2-3 days post-transduction until confluent and then

passaged for clonal expansion.

Analysis of targeted integration in iPSCs. iPSCs were assessed quantitatively for on-target

insertions/deletions and targeted integration through NGS sequencing on the Illumina platform.

Genomic DNA from clonally expanded iPSCs was first amplified, followed by nested PCR

amplification to prepare ~200 bp amplicons for paired-end deep sequencing on the Illumina MiSeq

sequencer.

iPSC-Treg Cryopreservation. Post-sort CAR iPSC-Tregs were recovered in complete OpTmizer

supplemented with 1000U IL-2/mL at a concentration of 1e6/mL. 24 hours after recovery, CAR

iPS-Tregs were harvested, counted and cryopreserved in CryoStor CS10 (Biolife Solutions) at a

concentration of 4-5e6/mL. Cryopreserved cells were placed in an alcohol-free freezing container

(CoolCell, Corning) and placed at -80C overnight before transferring to LN2 for long-term storage.

References

1. Themeli, M., Riviere, I. & Sadelain, M. New cell sources for T cell engineering and adoptive immunotherapy. *Cell Stem Cell* **16**, 357-366 (2015).

2. Dominguez-Villar, M. & Hafler, D.A. Regulatory T cells in autoimmune disease. *Nat Immunol* **19**, 665-673 (2018).

3. Sanders, J.M., Jeyamogan, S., Mathew, J.M. & Leventhal, J.R. Foxp3+ regulatory T cell therapy for tolerance in autoimmunity and solid organ transplantation. *Front Immunol* **13**, 1055466 (2022).

4. Proics, E. et al. Preclinical assessment of antigen-specific chimeric antigen receptor regulatory T cells for use in solid organ transplantation. *Gene Ther* **30**, 309-322 (2023).

5. Raffin, C., Vo, L.T. & Bluestone, J.A. T(reg) cell-based therapies: challenges and perspectives. *Nat Rev Immunol* **20**, 158-172 (2020).

6. Themeli, M. et al. Generation of tumor-targeted human T lymphocytes from induced pluripotent stem cells for cancer therapy. *Nat Biotechnol* **31**, 928-933 (2013).

7. Nishimura, T. et al. Generation of rejuvenated antigen-specific T cells by reprogramming to pluripotency and redifferentiation. *Cell Stem Cell* **12**, 114-126 (2013).

8. Montel-Hagen, A. et al. Organoid-Induced Differentiation of Conventional T Cells from Human Pluripotent Stem Cells. *Cell Stem Cell* **24**, 376-389 e378 (2019).

9. van der Stegen, S.J.C. et al. Generation of T-cell-receptor-negative CD8ab-positive CAR T cells from T-cell-derived induced pluripotent stem cells. *Nature biomedical engineering* (2022).

10. Wang, Z. et al. 3D-organoid culture supports differentiation of human CAR(+) iPSCs into

highly functional CAR T cells. *Cell Stem Cell* **29**, 515-527 e518 (2022).

11. Borst, J., Ahrends, T., Babala, N., Melief, C.J.M. & Kastenmüller, K. CD4+ T cell help in

cancer immunology and immunotherapy. *Nature Reviews Immunology* **18**, 635-647

(2018).

12. Turtle, C.J. et al. CD19 CAR-T cells of defined CD4+:CD8+ composition in adult B cell

ALL patients. *J Clin Invest* **126**, 2123-2138 (2016).

13. Sommermeyer, D. et al. Chimeric antigen receptor-modified T cells derived from defined

CD8+ and CD4+ subsets confer superior antitumor reactivity in vivo. *Leukemia* **30**, 492-

500 (2016).

14. Melenhorst, J.J. et al. Decade-long leukaemia remissions with persistence of CD4+ CAR

T cells. *Nature* **602**, 503-509 (2022).

15. Tay, R.E., Richardson, E.K. & Toh, H.C. Revisiting the role of CD4(+) T cells in cancer

immunotherapy-new insights into old paradigms. *Cancer Gene Ther* **28**, 5-17 (2021).

16. Vizcardo, R. et al. Regeneration of human tumor antigen-specific T cells from iPSCs

derived from mature CD8(+) T cells. *Cell Stem Cell* **12**, 31-36 (2013).

17. Iriguchi, S. et al. A clinically applicable and scalable method to regenerate T-cells from

iPSCs for off-the-shelf T-cell immunotherapy. *Nature communications* **12**, 430 (2021).

18. Chang, C.W., Lai, Y.S., Lamb, L.S., Jr. & Townes, T.M. Broad T-cell receptor repertoire in

T-lymphocytes derived from human induced pluripotent stem cells. *PLoS One* **9**, e97335

(2014).

19.   Ng, E.S., Davis, R., Stanley, E.G. & Elefanty, A.G. A protocol describing the use of a

recombinant protein-based, animal product-free medium (APEL) for human embryonic

stem cell differentiation as spin embryoid bodies. *Nat Protoc* **3**, 768-776 (2008).

20.   Kotini, A.G. et al. Stage-Specific Human Induced Pluripotent Stem Cells Map the

Progression of Myeloid Transformation to Transplantable Leukemia. *Cell Stem Cell* **20**,

315-328 e317 (2017).

21.   Schmitt, T.M., Ciofani, M., Petrie, H.T. & Zuniga-Pflucker, J.C. Maintenance of T cell

specification and differentiation requires recurrent notch receptor-ligand interactions. *J

Exp Med* **200**, 469-479 (2004).

22.   Trotman-Grant, A.C. et al. DL4-mubeads induce T cell lineage differentiation from stem

cells in a stromal cell-free system. *Nat Commun* **12**, 5023 (2021).

23.   Mohtashami, M., Brauer, P.M. & Zuniga-Pflucker, J.C. Induction of Human T Cell

Development In Vitro with OP9-DL4-7FS Cells Expressing Human Cytokines. *Methods

Mol Biol* **2580**, 249-260 (2023).

24.   Singer, A., Adoro, S. & Park, J.H. Lineage fate and intense debate: myths, models and

mechanisms of CD4- versus CD8-lineage choice. *Nat Rev Immunol* **8**, 788-801 (2008).

25.   Jameson, S.C., Hogquist, K.A. & Bevan, M.J. Positive selection of thymocytes. *Annu

Rev Immunol* **13**, 93-126 (1995).

26.   Ohoka, Y. et al. In vitro differentiation and commitment of CD4+ CD8+ thymocytes to the

CD4 lineage, without TCR engagement. *Int Immunol* **8**, 297-306 (1996).

27.   Takahama, Y., Suzuki, H., Katz, K.S., Grusby, M.J. & Singer, A. Positive selection of

CD4+ T cells by TCR ligation without aggregation even in the absence of MHC. *Nature

371, 67-70 (1994).

28.   Park, J.H. et al. Signaling by intrathymic cytokines, not T cell antigen receptors, specifies

CD8 lineage choice and promotes the differentiation of cytotoxic-lineage T cells. *Nat*

Immunol **11**, 257-264 (2010).

29.   Fu, S. et al. TGF-beta induces Foxp3 + T-regulatory cells from CD4 + CD25 -

precursors. *Am J Transplant* **4**, 1614-1627 (2004).

30.   Chen, W. et al. Conversion of peripheral CD4+CD25- naive T cells to CD4+CD25+

regulatory T cells by TGF-beta induction of transcription factor Foxp3. *J Exp Med* **198**,

1875-1886 (2003).

31.   Mitsuhiro, K., Hiroko, N., Masahiro, O., Qianjin, L. & Akihiko, Y. Induced Regulatory T

Cells: Their Development, Stability, and Applications. *Trends in Immunology* **37**, 803-811

(2016).

32.   Pesenacker, A.M. et al. Treg gene signatures predict and measure type 1 diabetes

trajectory. *JCI Insight* **4** (2019).

33.   Pesenacker, A.M. et al. A Regulatory T-Cell Gene Signature Is a Specific and Sensitive

Biomarker to Identify Children With New-Onset Type 1 Diabetes. *Diabetes* **65**, 1031-

1039 (2016).

34.   Höllbacher, B. et al. Transcriptomic Profiling of Human Effector and Regulatory T Cell

Subsets Identifies Predictive Population Signatures. *ImmunoHorizons* **4**, 585-596 (2020).

35.   Huehn, J., Polansky, J.K. & Hamann, A. Epigenetic control of FOXP3 expression: the

key to a stable regulatory T-cell lineage? *Nat Rev Immunol* **9**, 83-89 (2009).

36.   Floess, S. et al. Epigenetic control of the foxp3 locus in regulatory T cells. *PLoS Biol* **5**,

e38 (2007).

37. Dikiy, S. & Rudensky, A.Y. Principles of regulatory T cell function. *Immunity* **56**, 240-255

(2023).

38. Vignali, D.A., Collison, L.W. & Workman, C.J. How regulatory T cells work. *Nat Rev*

Immunol **8**, 523-532 (2008).

39. Moore, K.W., de Waal Malefyt, R., Coffman, R.L. & O'Garra, A. Interleukin-10 and the

interleukin-10 receptor. *Annu Rev Immunol* **19**, 683-765 (2001).

40. Louey, A., Hernandez, D., Pebay, A. & Daniszewski, M. Automation of Organoid

Cultures: Current Protocols and Applications. *SLAS Discov* **26**, 1138-1147 (2021).

41. Seki, T. et al. Generation of induced pluripotent stem cells from human terminally

differentiated circulating T cells. *Cell Stem Cell* **7**, 11-14 (2010).

42. Loh, Y.H. et al. Reprogramming of T cells from human peripheral blood. *Cell Stem Cell* **7**,

15-19 (2010).

43. Brown, M.E. et al. Derivation of induced pluripotent stem cells from human peripheral

blood T lymphocytes. *PLoS One* **5**, e11373 (2010).

44. Hernandez, J.B., Newton, R.H. & Walsh, C.M. Life and death in the thymus--cell death

signaling during T cell development. *Curr Opin Cell Biol* **22**, 865-871 (2010).

45. Nagano, S. et al. High Frequency Production of T Cell-Derived iPSC Clones Capable of

Generating Potent Cytotoxic T Cells. *Mol Ther Methods Clin Dev* **16**, 126-135 (2020).

46. Kawamoto, H., Masuda, K. & Nagano, S. Regeneration of antigen-specific T cells by

using induced pluripotent stem cell (iPSC) technology. *Int Immunol* **33**, 827-833 (2021).

47.   Maeda, T. et al. Regeneration of Tumor-Antigen-Specific Cytotoxic T Lymphocytes from
iPSCs Transduced with Exogenous TCR Genes. *Mol Ther Methods Clin Dev* **19**, 250-
260 (2020).

48.   Li, S. et al. Strength of CAR signaling determines T cell versus ILC differentiation from
pluripotent stem cells. *Cell Rep* **42**, 112241 (2023).

49.   Awong, G. et al. Human proT-cells generated in vitro facilitate hematopoietic stem cell-
derived T-lymphopoiesis in vivo and restore thymic architecture. *Blood* **122**, 4210-4219
(2013).

50.   Edgar, J.M., Michaels, Y.S. & Zandstra, P.W. Multi-objective optimization reveals time-
and dose-dependent inflammatory cytokine-mediated regulation of human stem cell
derived T-cell development. *NPJ Regen Med* **7**, 11 (2022).

51.   Klein, L., Robey, E.A. & Hsieh, C.S. Central CD4(+) T cell tolerance: deletion versus
regulatory T cell differentiation. *Nat Rev Immunol* **19**, 7-18 (2019).

52.   Takahama, Y. & Nakauchi, H. Phorbol ester and calcium ionophore can replace TCR
signals that induce positive selection of CD4 T cells. *The Journal of Immunology* **157**,
1508-1513 (1996).

53.   Karimi, M.M. et al. The order and logic of CD4 versus CD8 lineage choice and
differentiation in mouse thymus. *Nat Commun* **12**, 99 (2021).

54.   Li, Y. et al. Development of double-positive thymocytes at single-cell resolution. *Genome*
Med **13**, 49 (2021).

55.   Theofilopoulos, A.N., Koundouris, S., Kono, D.H. & Lawson, B.R. The role of IFN-
gamma in systemic lupus erythematosus: a challenge to the Th1/Th2 paradigm in
autoimmunity. *Arthritis Res* **3**, 136-141 (2001).

56. Butcher, M.J. & Zhu, J. Recent advances in understanding the Th1/Th2 effector choice.
Fac Rev **10**, 30 (2021).

57. Tran, D.Q., Ramsey, H. & Shevach, E.M. Induction of FOXP3 expression in naive human
CD4+FOXP3 T cells by T-cell receptor stimulation is transforming growth factor-beta
dependent but does not confer a regulatory phenotype. *Blood* **110**, 2983-2990 (2007).

58. Fong, H. et al. Genetic correction of tauopathy phenotypes in neurons derived from
human induced pluripotent stem cells. *Stem Cell Reports* **1**, 226-234 (2013).

59. Paschon, D.E. et al. Diversifying the structure of zinc finger nucleases for high-precision
genome editing. *Nat Commun* **10**, 1133 (2019).

60. Miller, J.C. et al. Enhancing gene editing specificity by attenuating DNA cleavage
kinetics. *Nat Biotechnol* **37**, 945-952 (2019).

Acknowledgements. We thank Prof. Maria Themeli for critical review of the manuscript and for
giving critical advice for our research work. We also thank Prof. Megan Levings for critical advice
on the transcriptomics analysis; Dr. Andreas Reik, Dr. Lei Zhang and Dr. George Kwong for
technical assistance; Dr. Julie Gertner-Dardenne, Dr. Maurus de la Rosa, Dr. David Fenard, Dr.
Tobias Abel and Dr. Virginie Fabre for scientific discussions; Mrs. Cayla McEwen and Mrs. Jean
Alladin for administrative support.

Patents. Part of this research work is described in the following patents:

US Patent number 11512287: "Targeted disruption of T cell and/or HLA receptors"

Application PCT/US2021/057152 (Filed Oct 28, 2021): "Generation of CD4+ Effector and

Regulatory T cells from Human Pluripotent Stem Cells"

Application PCT/US2020/059730 (Filed Sep 11, 2020): "Generation of Engineered Regulatory T

cells"

Author contributions. H.F., M.M., L.N., G.L., S.G., A.C. and S.Z. designed the study; H.F., M.M.,

J.J., L.N., K.K., S.S., J.H. and A.V. performed the experiments; H.F., M.M., J.J., L.N., K.K., G.L.,

S.G., S.S., J.H. and A.V. analyzed the data; H.F., M.M., L.N., S.S., A.C., J.F. and S.Z. interpreted

the data; J.F. provided advice for the experiments; A.C. and S.Z. supervised the study; H.F. and

S.Z. wrote the manuscript

Competing interests. All authors are current or former Sangamo Therapeutics employees and

shareholders.

Correspondence and requests for materials should be addressed to H.F. and S.Z.

MAIN FIGURES

FIGURE 1. Differentiation of iPSCs to HSPCs, lymphoid progenitors and DP T cells in serum- and feeder-free conditions.

Figure 1. Differentiation of iPSCs to hematopoietic stem and progenitor cells (HSPCs), lymphoid progenitors and double positive (DP) T cells in serum- and feeder-free conditions.

A. Schematic for differentiation of iPSCs to HPSCs.
B. Embryoid bodies (EBs) at day 1 and 14 of differentiation. HSPCs emerged from the EB by day 14 and appeared as loosely packed cells (inset).
C. Representative flow cytometric analyses of HSPCs, lymphoid progenitor cells, and DP T cells from iPSCs. Cells were analyzed for phenotype over a 42-day period at the indicated timepoints (n=3).
D. Schematic for differentiation of iPSC-HSPCs to lymphoid progenitor cells and DP T cells.

FIGURE 2. Cell Yield per Input iPSC.

Figure 2. Yield of iPSC-derived hematopoietic stem and progenitor cells (HSPCs), lymphoid progenitors, double positive (DP) T cells, CD4sp T cells, and Tregs. Cell yield per input iPSC at various stages of differentiation (n=3).

FIGURE 3. Differentiation of iPSCs to CD4sp T cells and Tregs in serum- and feeder-free conditions.

D

Q1 (CD62L+CD45RO-)
Q2 (CD62L+CD45RO+)
Q3 (CD62L-CD45RO+)
Q4 (CD62L-CD45RO-)

E

CD4sp
CD8sp
DP
DN

Figure 3. Differentiation of iPSCs to CD4sp T cells and Tregs in serum- and feeder-free conditions.
   A. Schematic for differentiation of iPSCs to CD4sp T cells and Tregs.
   B. Representative kinetic analysis of developing CD4sp T cells after Phorbol 12-myristate 13-acetate and Ionomycin (PMA/I) addition. Phenotype was assessed by the expression levels of CD4 and CD8 by flow cytometry.
   C. Representative flow plots of iPSC-CD4sp T cells 8 days after PMA/I addition (day 50 of differentiation) to assess for CD4 commitment. Phenotype was assessed by the expression levels of CD3, TCR$\alpha\beta$, CD4, CD8, FOXP3 and ThPOK in cells with and without PMA/I treatment (n=6).
   D. Representative flow plots and expression of CD4 markers in unactivated T conventional (Tconv)–and sorted unactivated iPSC-CD4sp T cells. Phenotype was assessed by flow cytometric analysis (n=3) in Tconv and iPSC-CD4sp at day 50 of differentiation for the expression of CD4, CD62L, CD45RO and CD28. CD28 FMO control (gray) and CD28 expression (red).
   E. Frequency of cell subpopulations from all live cells at day 50 and day 57 of differentiation (top) and representative flow cytometric plots of iPSC-Tregs 7 days after TGFβ+ATRA addition (day 57 of differentiation) (bottom). CD4sp, CD8sp, double positive (DP), and double negative (DN) populations were assessed by flow cytometric analysis of CD4 and CD8 expression (n=6).

Figure 4. Single cell gene expression analysis of iPSC-Tregs.

A

Primary Tconv

iPSC-CD4sp

Induced Treg

Primary Treg

iPSC-Treg

B

Figure 4. Single cell gene expression analysis of iPSC-Tregs.

A. Hierarchical clustering of iPSC-Tregs, iPSC-CD4sp Tcells, cultured primary T conventional (Tconv), cultured primary Treg and induced Tregs based on differentially expressed genes (DEGs). DEGs with fold change <2 and p value<0.05, were subjected to CLC analysis to perform clustering.

B. scRNA-seq heatmap showing hierarchical clustering of a core Treg gene signature in iPSC-Tregs, iPSC-CD4sp, cultured primary Tregs, induced Tregs, and cultured primary Tconv.

FIGURE 5. Characterization of sorted and activated iPSC-CD4sp and iPSC-Tregs.

A

B

Figure 5. Characterization of sorted and activated iPSC-Tregs.

A. Representative flow plots and expression of CD25 in sorted unactivated and activated iPSC-CD4sp T cells or T conventional (Tconv) cells (n=3) by flow cytometry after 48 hours. CD25 Median fluorescence intensity (MFI) was calculated (right).

B. Expression of activation markers (CD69, PD-1 and CD28) was assessed by flow cytometry after 48 hours of activation. MFI was calculated (right).

C. Secretion of cytokines from sorted unactivated and activated iPSC-Tregs 24 hours after activation (n=3, iPSC-CD4sp; n=3, primary Tconv).

D. Assessment of FOXP3 TSDR demethylation in iPSC-Tregs, primary Tregs, and primary Tconv. Post-sort iPSC-Tregs were collected at day 57 for analysis and compared to primary controls (n=4, iPSC-Tregs; n=3, primary Tregs; n=3, primary Tconv).

E. Expression of Treg markers in sorted unactivated and activated iPSC-Tregs. Phenotype was assessed by flow cytometric analysis (n=6, unactivated and activated). ***p<0.001

F. Expression of CD69 in sorted unactivated and activated iPSC-Tregs. The expression of CD69 was assessed in both total live cell populations and target iPSC-Tregs by flow cytometry (left). Fold change in CD69 expression was assessed in activated total live and target iPSC-Tregs compared to unactivated total live cells and iPSC-Tregs (right, n=6, unactivated and activated).

G. Phenotype of unactivated and activated iPSC-Tregs. Treg markers were assessed by flow cytometric analysis (n=6).

Figure 6. Functional analysis of iPSC-Tregs.

Figure 6. Functional analysis of iPSC-Tregs.
- A. Cytotoxic activity of sorted unactivated and activated iPSC-Tregs as measured by an LDH release assay. iPSC-Tregs, primary Tregs, and primary CD8 T cells were cocultured with T conventional (Tconv) cells and LDH was measured after 3 days (n=3, iPSC-Tregs; n=3, primary Tregs; n=2, primary CD8 T cells).
- B. Secretion of cytokines from sorted unactivated and activated iPSC-Tregs 24 hours after activation (n=4, iPSC-Tregs; n=3, primary Tregs; n=3, primary Tconv). ****p<0.0001; **p=0.0011.
- C. Suppression of Tconv proliferation by unactivated and activated iPSC-Tregs, iPSC-CD4sp T cells and primary Tregs. iPSC-Tregs, primary Tregs, and iPSC-CD4sp T cells were cocultured with activated CTV labeled Tconv and CTV dilution was measured by flow cytometry after 3 days (n=6, iPSC-Tregs; n=4, primary Tregs; n=3, iPSC-CD4sp). ***p=0.0006; **p<0.001; *p<0.05.

FIGURE 7. Generation and Characterization of HLA-A2 CAR iPSC-Tregs

Figure 7. Generation and characterization of HLA-A2 CAR iPSC-Tregs.
   A. Editing strategy for generation of HLA-A2 CAR iPSCs. A donor construct showing the TCR coding sequence and HLA-A2 CAR sequence flanked by homology arms targets exon 2 of the endogenous T Cell Receptor Alpha Constant (TRAC) locus.
   B. Suppression of T conventional (Tconv) cell proliferation by HLA-A2 CAR+ iPSC-Tregs activated *via* the TCR or CAR. HLA-A2 CAR+ iPSC-Tregs were cocultured with activated CTV labeled Tconv and CTV dilution was measured by flow cytometry after 3 days (n=4). *p<0.05
   C. Suppression of Tconv proliferation by cryopreserved HLA-A2 CAR+ iPSC-Tregs activated via the TCR or CAR. HLA-A2 CAR+ iPSC-Tregs were cocultured with activated CTV labeled Tconv and CTV dilution was measured by flow cytometry after 3 days (n=2).

SUPPLEMENTARY FIGURES

SUPPLEMENTAL FIGURE 1. Characterization of iPSCs

Supp. Figure 1. Characterization of iPSCs.
- A. Representative images of T cell-derived iPSCs and HLA-A2 CAR edited iPSCs. iPSCs were assessed for morphology by phase contrast microscopy and expression of OCT4, SOX2, SSEA-4, and TRA-1-81 by immunocytochemistry.
- B. G-band karyotyping analysis confirming normal 46 XY karyotype in T cell-derived iPSCs and HLA-A2 CAR-edited iPSCs.

SUPPLEMENTARY FIGURE 2. Double positive (DP) T Cell Yield from Unedited and HLA-A2 CAR iPSCs with CXCL12 and p38i Treatment.

Supp. Figure 2. Double positive (DP) T cell yield from Unedited and HLA-A2 CAR iPSCs with CXCL12 and p38i treatment.
Yield of DP T cells from unedited and CAR edited iPSCs after 42 days of differentiation. CXCL12 and p38i were added at day 28 of differentiation (n=4, unedited; n=4, HLA-A2 CAR).

SUPPLEMENTAL FIGURE 3. Sorting Strategy for iPSC-CD4sp and for iPSC-Tregs

Supp. Figure 3. Sorting strategy for iPSC-CD4sp T cells and iPSC-Tregs.

Supp. Figure 4. Characterization of sorted and activated iPSC-CD4sp.

HF 24hr Tcon:MFI CD25 from Live CD4sp CD25

Supp. Figure 4. iPSC-CD4sp T cell characterization activation markers at 24 hours post-stimulation
   A. Representative flow plots and expression of CD25 in sorted unactivated and activated iPSC-CD4sp T cells or T conventional (Tconv) cells (n=3) by flow cytometry. CD25 Median fluorescence intensity (MFI) was calculated (right).
   B. Expression of activation markers (CD69, PD-1 and CD28) was assessed by flow cytometry after 24 hours of activation. MFI was calculated (right).

SUPPLEMENTARY FIGURE 5. Characterization of HLA-A2 CAR+ iPSC-Tregs pre-sort

Supp. Figure 5. Characterization of HLA-A2 CAR+ iPSC-Tregs

A. Representative flow cytometric plots of hematopoietic stem and progenitor cells (HSPCs), lymphoid progenitor cells, and double positive (DP) T cells from HLA-A2 CAR edited iPSCs (n=3).

B. Representative flow plots of iPSC-CD4sp T cells 8 days after PMA/I addition (day 50 of differentiation). Phenotype was assessed by the expression levels of CD3, TCRαβ, CD4, CD8, FOXP3 and ThPOK in cells with and without PMA/I treatment (n=3).

C. Representative flow plots of iPSC-Tregs 7 days after TGFβ+ATRA addition (differentiation day 50). CD4sp phenotype was assessed by the expression levels of CD3, TCRαβ, CD4, CD8, FOXP3 and ThPOK (n=3). Treg phenotype was assessed by the expression levels of CD4, CD8, CD25, CD127, and FOXP3 (n=3).

D. MFI of CAR expression as measured at differentiation days 42, 50, and 57.

SUPPLEMENTARY FIGURE 6. Characterization of sorted HLA-A2 CAR+ iPSC-Tregs

Supp. Figure 6. Functional characterization of HLA-A2 CAR+ iPSC-Tregs

A. Expression of Treg markers in sorted unactivated or control dextramer activated HLA-A2 CAR+ iPSC-Tregs and HLA-A2 CAR+ iPSC-Tregs activated via the TCR (CD3/CD28) or HLA-A2 CAR (HLA-A2 dextramer). Phenotype was assessed by flow cytometric analysis (n=4).

B. Expression of CD69 in sorted unactivated or control dextramer activated HLA-A2 CAR+ iPSC-Tregs and HLA-A2 CAR+ iPSC-Tregs activated via the TCR or HLA-A2 CAR. The expression of CD69 was assessed in both total live cell populations and target iPSC-Tregs by flow cytometry (left). Fold change in CD69 expression was assessed in activated total live and target iPSC-Tregs compared to unactivated total live cells and target iPSC-Tregs (right, n=4). *p<0.05.

C. Secretion of cytokines from sorted unactivated or control dextramer activated HLA-A2 CAR+ iPSC-Tregs and HLA-A2 CAR+ iPSC-Tregs activated via the TCR or HLA-A2 CAR (n=4, iPSC-Tregs).

SUPPLEMENTARY FIGURE 7. Characterization of Cryopreserved HLA-A2 CAR+ iPSC-Tregs

Supp. Figure 7. Characterization of Cryopreserved HLA-A2 CAR+ iPSC-Tregs
  A. Expression of Treg markers in cryopreserved unactivated HLA-A2 CAR+ iPSC-Tregs. Phenotype was assessed by flow cytometric analysis (n=2).
  B. Fold increase in CD69 expression was assessed in activated total live and target iPSC-Tregs compared to unactivated total live cells and iPSC-Tregs (n=2).