1

2 **Title:** Initial and corrective submovement encoding differences within primary motor cortex during
3 precision reaching
4

5 **Authors:**
6 Schwartze, Kevin[12]; Lee, Wei-Hsien[13]; Rouse, Adam G. [1234]
7

8 **Affiliations:**
9 [1]University of Kansas Medical Center, Department of Neurosurgery
10 [2]University of Kansas Medical Center, Department of Cell Biology and Physiology
11 [3]University of Kansas, Bioengineering Program
12 [4]University of Kansas, Department of Electrical Engineering and Computer Science
13

14 **Corresponding Author:**
15 Dr. Adam G. Rouse, MD, PhD
16 University of Kansas Medical Center
17 Department of Neurosurgery
18 arouse@kumc.edu

Abstract

Precision reaching tasks often require corrective submovements for successful completion. Most studies of reaching have focused on single initial movements, and the cortical encoding model was implied to be the same for all submovements. However, corrective submovements may show different encoding patterns from the initial submovement with distinct patterns of activation across the population. Two rhesus macaques performed a precision center-out-task with small targets. Neural activity from single units in primary motor cortex and associated behavioral data were recorded to evaluate movement characteristics. Neural population data and individual neuronal firing rates identified with a peak finding algorithm to identify peaks in hand speed were examined for encoding differences between initial and corrective submovements. Individual neurons were fitted with a regression model that included the reach vector, position, and speed to predict firing rate. For both initial and corrective submovements, the largest effect remained movement direction. We observed a large subset changed their preferred direction greater than 45º between initial and corrective submovements. Neuronal depth of modulation also showed considerable variation when adjusted for movement speed. By utilizing principal component analysis, neural trajectories of initial and corrective submovements progressed through different neural subspaces. These findings all suggest that different neural encoding patterns exist for initial and corrective submovements within the cortex. We hypothesize that this variation in how neurons change to encode small, corrective submovements might allow for a larger portion of the neural space being used to encode a greater range of movements with varying amplitudes and levels of precision.

Introduction

Our brain allows us to execute movement over a large dynamic range. While large, ballistic movements typically need less fine control, precision movements require it (Abrams et al. 1990; Craik 1948; Elliott et al. 2010; Sainburg et al. 1999; Woodworth 1899). One reaching movement can be broken down into 2 types of submovements: an initial submovement or any following corrective submovements (Milner and Ijaz 1990; Novak et al. 2002; Port et al. 1997; Pratt et al. 1994). Reaching movements have bell-shaped velocity profiles (Flash and Hogan 1985; Georgopoulos et al. 1981; Morasso 1981; Soechting 1984), a feature seen in both the initial and corrective submovement components (Hatsopoulos et al. 2007). However, by their nature, corrective submovements have a significantly smaller magnitude than the initial submovement.

The firing rates of individual neurons in the motor cortex signal information about upcoming movements. Populations of individual neurons within the motor cortex represent several reaching characteristics of action selection and execution (Kalaska et al. 1997), including force generation (Evarts 1968) and preferred direction (Fu et al. 1993; Georgopoulos et al. 1982, 1986). Technical advances have allowed for recording from more neurons to sample and build accurate predictions of upcoming movement features from specific neuronal firing within the cortex, (Kalaska and Crammond 1992; Moran and Schwartz 1999; Nicolelis et al. 1997). Models can describe how neural activity depends on various movement conditions (Moran and Schwartz 1999; Wang et al. 2007). Condition dependent activity is an increase of neural firing represented by movement specific parameters such as direction, speed, and spatial location (Georgopoulos et al. 1986; Moran and Schwartz 1999; Rouse, Adam G., Schieber 2017; Rouse and Schieber 2016; Wang et al. 2007).

Most experiments of neural encoding for reaching have only examined instructed movements that
were generally large, overlearned, and stereotyped to allow for trial averaging. For some analyses and
applications such as brain-computer interfaces (BCIs), identical models of encoding have been assumed
to fit both initial and corrective submovements. However, we hypothesize that initial and corrective
submovements are encoded differently (Evarts and Fromm 1977; Fu et al. 1995; Milner 1992; Scott
2004), allowing for a greater diversity of successive submovements. Understanding the fundamental
differences between submovement planning, execution, and implementation is vitally important for real-
world applications such as brain-computer interface algorithms (Umair et al. 2017). Current BCI
prostheses are capable of performing basic functions; however, users struggle to complete precision tasks
with the devices (Mridha et al. 2021). Recent BCI algorithms have started incorporating increasingly
complex decoding systems (Inoue et al. 2018; Karpowicz et al. 2022; Ma et al. 2022), but more
information about the underlying organization in the brain of corrective movements in healthy reaching
will help drive further development.

As researchers record larger populations of motor cortex neurons during reaching movements, the
results show previously indiscernible elegant patterns of neural activity (Churchland et al. 2006;
Churchland and Shenoy 2007; Maynard et al. 1999). These neural firing patterns can be temporally
mapped in multidimensional neural subspaces as (Churchland et al. 2012; Kaufman and Churchland
2016; Rouse and Schieber 2018) using dimensionality-reduction tools, such as principal components
analysis (PCA). Comparison of neural trajectories between initial and corrective submovements is thus far
unquantified and would provide important information about cortical encoding patterns during precision
reaching.

In the current study we utilized a custom precision center-out task to encourage multi-phasic
submovement corrections. We previously showed that condition-invariant neural activity reliably defines
the timing of both the initial and corrective submovements (Rouse et al. 2022). We now ask how neural
firing rates change with the specific reach conditions of velocity, position, and speed during initial and
corrective submovements. Kinematic behaviors were aligned with primary motor cortex neural recordings
to evaluate encoding pattern differences between initial and corrective submovements. We utilized a well-
established cortical tuning model (Wang et al. 2007) to evaluate encoding differences between
submovement types as well as modeled the observed trajectories of the observed neural tuning to measure
shared encoding subspaces.

Methods

Macaque Model

Two adult, male rhesus macaques (Macaca mulatta), P and Q, were used in the research project. All
standard procedures for the care and use of these nonhuman primates followed the Guide for the Care and
Use of Laboratory Animals and received approval from the University Committee on Animal Resources
at the University of Rochester Medical Center, Rochester, NY.

Experimental Design

Subjects P and Q were trained to perform a precision center-out-task using a custom interface equipped
with a joystick and monitor (Figure 1A). The joystick was 18 cm and had the centering spring removed,

allowing it to move freely with little resistance. The precision task required the subject to single handedly
manipulate the joystick moving the displayed cursor from the center into an indicated goal area. The goal
areas were spaced equidistant from each other in 45° increments, and designed to be narrow and deep,
wide and shallow, or wide and deep (Figure 1B).

Our precision center-out-task utilized small targets and a required hold time to elicit not only initial
center-out reaches but also corrective movements. After reaching the goal area, the animals were required
to hold for 500-600 ms. If the cursor left the goal area within the hold interval, we allowed them to move
and re-enter the goal as many times as required to complete a successful hold. Animals were provided a
liquid reward upon successful trial completion. Additional animal and task details are described in Rouse
et al. 2022.

Figure 1

Figure 1. Precision center-out-task design. A) Rhesus macaques manipulated a joystick controlling the
cursor on a digital screen. The cursor was required to move from the center starting area to the indicated
peripheral target zone. B) Peripheral target zone layouts designed to encourage corrective submovements
during precision reaching. Target zones are wide and deep (left), narrow and deep (middle), or wide and
shallow (right).

Neural Recordings

Both subjects P and Q were surgically implanted with chronic multielectrode arrays to detect
individual neuronal activity. Floating microelectrode arrays (MicroProbes for Life Science) were
implanted in primary motor cortex (M1) in each monkey, using methods described in detail previously
(Mollazadeh et al. 2011; Rouse et al. 2022; Rouse and Schieber 2016). For monkey P, recordings were
collected from six 16-channel arrays implanted in M1. For monkey Q, two 32-channel arrays and one 16-
channel array in M1 were used. Channels with spiking activity were collected with Plexon MAP (Plexon)
and Cerebus (Blackrock Microsystems, LLC.), thresholded manually online, and sorted off-line with a
custom, semi-automated algorithm.

Behavior Analysis

Our previous work found that most movements in our precision center-out-task are well defined
as distinct submovements with bell-shaped speed profiles separated by low-speed intervals. Here, we
again identified with initial and corrective submovements by identifying cursor speed peaks (Figure 2)
using a peak finding algorithm as previously described in Rouse 2022. Briefly, speed peaks were defined
as having a i) speed >250 pixels/s, ii) no larger peak within ±200-ms, and iii) having a peak prominence
of at least 50% of its two adjacent speed troughs. All speed-peak defined submovements from leaving the
center until completion of a successful hold were identified. The initial submovement was defined as the
large initial reach that moved at least halfway to the goal area and all subsequent submovements were
defined as corrective. Rare small amplitude submovements before the initial submovements as well as
submovements made entirely within the goal area were discarded from analysis.

Movement Vectors

All submovements, both initial and corrective, were approximated as straight-line vectors using
the beginning and end cursor locations (Figure 2C) to simplify the analysis of the relationship between
neural activity and submovement kinematics. Each vector is comprised of a starting position, the
movement direction, and distance travelled within the 200 ms time window from 100 ms before peak
speed until 100 ms after. All initial submovements had a similar starting point in the center region, as
expected during a center-out task. Most corrective submovements started in the periphery of the
workspace near the goal areas.

Figure 2. Submovement identification and vector approximations. A) Movements performed on the wide
and shallow peripheral target. Initial submovements are traced in blue and corrective submovements are
in red with speed peak occurrence noted with a circle. B) Speed profiles for the submovements in (2A)
with initial speed peaks denoted with a blue square and corrective speed peaks noted with a red square.
The dotted line shows the 250pixel/s threshold for submovement designation. C) Initial (blue) and
corrective (red) submovements approximated as single line vectors. Submovement starting location
indicated with a circle.

Multivariate Regression

Multivariate linear regression (Figure 3) was performed on initial and corrective submovements to analyze correlations of the kinematics to neural firing rates. We used the established cortical activity model proposed published by Wang et al. (2007). This model included 3 features: position, velocity, and speed defined by the straight-line vector. Position for each submovement was defined as the starting movement position 100 ms before peak speed. Velocity was the submovement vector from movement onset (100 ms before) to offset (100 ms after). Speed was the magnitude of the submovement vector.

The regression model uses the following equation:

$$FR(t - \tau) = b_0 + b_x x + b_y y + b_{\dot{x}} \dot{x} + b_{\dot{y}} \dot{y} + b_s S$$

Here x and y are the starting position such that x=0 and y=0 defines the center. The velocity vectors are defined by $\dot{x}$ and $\dot{y}$. The speed, S, was the magnitude of the velocity vector.

Firing rates were calculated with a Gaussian smoothing window (σ=30 ms) and sampled in 10ms bins. A sliding time lag (τ) from 300 ms before until 100 ms after peak movement speed was used to identify and select the highest total R^2 for each spiking unit for initial and corrective submovements separately. Once this single time lag was identified, the spiking units' relation to kinematic features was examined.

We were primarily interested in how the velocity tuning changed between initial and corrective submovements after correcting for differences in starting position and speed. A neuron's preferred reach direction (Θ_{Pref}) is the direction with maximal firing rate and can be defined as:

$$\theta_{Pref} = atan2(b_{\dot{y}}, b_{\dot{x}})$$

Using the 2-argument arctangent of the x and y velocity coefficients, $b_{\dot{x}}$ and $b_{\dot{y}}$. A neuron's depth of modulation (DOM) measures how much the firing rate changes between reaches in the preferred and anti-preferred direction.

$$\text{DOM} = \left\| b_{\dot{x}}, b_{\dot{y}} \right\|$$

It is the combined magnitude of the x and y velocity coefficients, $b_{\dot{x}}$ and $b_{\dot{y}}$. For examining the relationship between firing rate and reach direction, we display the marginal firing rates corrected for speed and starting position when firing rates as a function of reach direction are shown.

To estimate the confidence interval (C.I.) for the preferred direction estimate, bootstrapping by drawing with replacement over 10,000 iterations from the spiking unit population deemed to be significantly tuned. The 95 percent C.I. was then taken to be the range between the 2.5 and 92.5th percentile of the bootstrapped samples.

Neural Population Model

A neural population model was built using the cortical tuning parameters identified with our multivariate regression to further explore the differences in neural tuning between initial and corrective submovements. Since the corrective submovements were idiosyncratic due to motor error and not repeated to allow for trial averaging, we instead simulated the neural activity for initial and corrective submovements to each of the 8 center-out directions. These simulated movements were assumed to be all the same magnitude as the average speed of either initial or corrective submovements. The linear tuning

parameters for each spiking unit as a function of time were then used to generate simulated neural
trajectories to the 8 different targets.

Principal component analysis was used to identify the 2-dimensional neural subspace with the
most variance at the timepoint 150 ms before peak speed for both initial and corrective submovements.
These subspaces were calculated for both initial and corrective neural activity and the opposite neural
activity was also projected into the separately identified subspaces. The percentage of total variance
accounted for simulated activity by these subspaces for all activity from 300 ms before until 100 ms after
peak speed was calculated. Additionally, the total overlap of simulated neural activity (Elsayed et al.
2016; Rouse and Schieber 2018) was calculated using the formula:

209
$$Overlap \; = \; \frac{tr(\Sigma_{Init}\Sigma_{Cor})}{\|\Sigma_{Init}\|_F \cdot \|\Sigma_{Cor}\|_F}$$

210 The 2 covariance matrices for initial and corrective neural activity ($\Sigma_{Init}, \Sigma_{Cor}$) are compared by using the
211 trace of the multiplied covariance matrices to calculate the overlap. This result is normalized by the
212 Frobenius norm ($\|...\|_F$) of each covariance matrix to obtain a metric between 0 (completely orthogonal
213 neural activity) and 1 (complete alignment).

Results

Multivariate Regression

Subjects P and Q successfully completed 10964 and 8126 trials, respectively. Individual neurons
from each session were fit with a multivariate linear regression model separately for all initial and
corrective submovements. It included the 2-dimensional reach direction vector, hand position, and hand
speed to predict firing rate. Neurons were considered significantly tuned with an R^2 value of greater than
0.1 for either initial or corrective submovements. A total of 1293 significantly tuned neurons were
identified for subject P and 1185 for Subject Q. For initial submovements, the largest effect for almost all
units was velocity (Figure 3A,B). This was expected as there was little variability of speed or position for
the initial submovements as the starting hand position and distance to target remained approximately the
same for all trials. For corrective submovements analysis, the influence of the velocity vector remained
the largest factor; however, it decreased overall when compared to initial submovements (Figure 3C,D).
Significantly larger impacts by hand position and speed were both observed, with hand position having a
higher R^2 in both subjects. We examined the residual noise with the root-mean square error (RMSE), for
each unit after the fitting the three factors of velocity, position, and speed. A similar amount of deviation
for individual reaches was observed. Suggesting a relatively similar ability of our multivariate regression
to predict neural activity in both initial and corrective submovements. Albeit with a lower R^2 due to the
generally decreased modulation of firing rate for corrective movements.

Figure 3. Multivariate regression analysis. A,B) Partial R2 for firing rate explained for the velocity vector, starting hand position, and movement speed of all significantly tuned neurons using multivariate regression for initial and C,D) corrective submovements for monkeys P and Q. Partial pie charts represent the overall the regression elements in proportion to each other and the percent contribution to the total. E,F) Root-mean-square-error (RMSE) analysis of initial and corrective submovements. The dotted line shows equivalence for residual noise between initial and corrective. The direct correlation is plotted with the cyan line.

Preferred Direction

We next used the linear cortical tuning model for each unit to generate its cosine tuning curve as a function of reach direction. The direction with the largest increase of firing rate is defined as the preferred direction with the direction 180° opposite being the anti-preferred direction. We analyzed the preferred direction of each spiking unit for both initial and corrective submovements separately. If both

submovements are encoded identically, the modeled preferred direction would be the same. For both
subjects, the majority of neurons maintained approximately the same directional tuning, with less than a
45° change in preferred direction between submovement types (Figure 4B,C); however, a significant
number of neurons demonstrated a change in preferred direction to an entirely different goal target
location. Subject P had 42.5% (39.0-46.0%, 95% C.I.) of significantly tuned neurons change their
preferred direction at least 45° between submovement types (Figure 4B). Subject Q had a similar
representation of 40.0% (36.1-43.7%, 95% C.I.) of neurons changing preferred direction (Figure 4C).

Figure 4. Preferred direction analysis for all significantly (R^2>0.1) tuned M1 neurons. A) The cosine
tuning curve for an example neuron with both initial (blue) and corrective (red) submovements. The
preferred (max firing rate per direction) directions are marked with circles. Note, the firing rate may be
slightly negative for some submovements due to the correction made for position and speed to show the
relationship to reach direction. B) The number of neurons with a ≥45° change of preferred direction
encoding between initial and corrective submovements for Subject P and C) Subject Q.

Depth of Modulation

The depth of modulation is the firing rate change between the preferred and anti-preferred
directions (Figure 5A). As expected for most spiking units, a larger depth of modulation was observed
during initial submovements than during corrective submovements (5B,C). For a single linear encoding
model, proportional increases in modulation would be expected to accompany increases in movement
speed. However, when the depth of modulation is scaled linearly for movement speed, the changes in
firing rate showed considerable variation (Figure 5D,E). Some neurons were primarily active only during
initial submovements, other neurons exhibited significantly increased gain in firing rate for the small
corrective submovements, and some neurons exhibited the changes along the 45° diagonal consistent with
a fixed linear relationship with speed. No clear relationships were observed suggesting different patterns
of neural activity at the individual neuron level for the two submovement types.

Figure 5. Depth of modulation analysis for tuned M1 neurons. A) The cosine tuning curves of an example neuron for initial (blue) and corrective (red) submovements. The depth of modulation is the difference in firing rate between the preferred and anti-preferred directions for each submovement type (bold blue and red vertical lines). Note, the firing rate may be slightly negative for some submovements due to the correction made for position and speed to show the relationship to reach direction. B) Direct comparison of all individual neurons' depth of modulation analysis for all $R^2>0.1$ units for initial and corrective submovements for monkey P and C) monkey Q. The dotted line shows equivalence for panels A,B and 2x, 1x, and 0.5x corrective to initial firing rate ratio for D,E . D) Velocity adjusted depth of modulation comparison for all $R^2>0.1$ units for monkey P and E) monkey Q. F) Velocity adjusted spiking units separated by activity dedicated to primarily initial or corrective submovements for monkey P and G) monkey Q.

Neural population model

To further explore the tuning observed across the neural population, we built a simulated neural population with the tuning parameters observed. Using each spiking unit's tuning coefficients for initial and corrective submovements, we generated simulated neural activity for an entire population as if the subject performed the same amplitude center-out reaches in 8 directions. One set of firing rates simulated the tuning observed for initial and one set for corrective submovements. We next used principal component analysis (PCA) to separately identify and display the neural subspaces with the most initial and corrective variance. We then plotted the neural trajectories of the modeled firing rates for both initial and corrective submovements in these separate subspaces. In the initial subspace, larger magnitude neural activity occurred for the initial neural data than the corrective, as expected (Fig. 6). In the separate corrective subspace, however, larger firing rate changes occurred in these dimensions for the corrective neural data. Suggesting better representation of these corrective, smaller amplitude movements in different neural dimensions. The shifts in tuning indicate neural dimensions with increased neural activity depending on execution of an initial or corrective submovement. We hypothesize that different neural dimensions—as created by shifts in preferred direction and depths of modulation for a subset of neurons—allow for more accurate encoding of small precise movements than if the same, global neural subspace signaled both initial and corrective movements.

Initial Subspace, a.u.

Corrective Subspace, a.u.

Figure 6. Modeled firing rates using tuning parameters observed during initial and corrective submovements. Two different neural subspaces maximized variance for initial (top) and corrective (bottom) submovements. Left) The neural activity for simulated reaches with tuning observed for initial submovements. Middle) The neural activity for simulated reaches with tuning observed for corrective submovements. Right) Initial (blue) and corrective (red) neural tuning shown in the initial and corrective subspaces. The colors (left and middle) represent reaches to the 8 center-out targets. The dots indicate neural activity 150 ms before peak speed. The shifts in tuning indicate neural dimensions with increased neural activity depending on execution of an initial or corrective submovement. Percentages indicate how much of the total neural activity is accounted for in the given two-dimensional subspace. For monkey Q (neural trajectory plot not shown), the two dimensional subspace maximized for initial and corrective neural activity accounted for 52.8% and 47.1% of total neural activity, respectively. The initial subspace accounted for 33.1% of corrective and the corrective subspace also accounted for 33.1% of initial. The total overlap in simulated neural activity between initial and corrective submovements was 0.31for monkey P and 0.45 for monkey Q.

Discussion

Utilizing a custom precision center-out task, we quantified differences of several neural tuning parameters between initial and corrective submovements within the primary motor cortex. While differences are not unexpected, quantifying these differences is an important step in describing neural tuning to not only instructed movements but also subsequent corrective movements and could be a key foundation for BCI encoders to build upon. Our analysis primarily focused on individual neuron's preferred direction and depth of modulation.

Shifts in Neural Tuning

Directional tuning of neurons in the primary motor cortex is well established (Fu et al. 1993; Georgopoulos et al. 1982, 1986). We have demonstrated that a significant portion of individual neurons in primary motor cortex change their preferred direction when encoding for corrective versus initial submovements. We observed approximately 40% of spiking units were classified to have a >45° shift in preferred direction, an entire target area shift in the task protocol.

It has been well documented that larger and faster movements are typically elicited by increased neural descending signals (Desmedt and Godaux 1977; Farina et al. 2002; Freund 1983; Milner-Brown et al. 1973). As expected, our results showed an overall higher firing rate towards initial submovements, as would be expected due to their innately larger magnitude; however, when we adjusted for movement velocity, a range of relationships between speed and individual cells became apparent. Some neurons fired almost exclusively during initial submovements, and others exhibited a much larger response for corrective submovements than expected for a linear speed relationship. Rather than uniform increases in firing rate, our results align with sequential recruitment of some new cortical neurons with increasing speeds similar to the Henneman's size principal for motoneurons (1965). Additionally, a small fraction of units are even more active for smaller movements than larger (Evarts et al. 1983), highlighting that neural tuning to speed is far from consistent across the neural population.

Different Neural Subspaces

The observed shifts and modulation in neural tuning lead us to propose that different neural subspaces are utilized within the motor cortex to encode the initial and corrective submovements. These different neural subspaces can be thought of as different combinations or patterns of neurons selectively activated for movements in the same direction but with different amplitudes or task demands. The many neurons with shared preferred directions between initial and corrective submovements likely are due to the same agonist and antagonist muscles being responsible for reaching in each direction. However, for different magnitude movements, the amount of co-contraction and balance between agonist/antagonist muscle may shift along with changes in fixator muscle demands (Griffin et al. 2015).

Beyond the muscle activation differences, different subspaces potentially provide an elegant means to adjust the signal gain between cortical firing rates and muscle activation levels (Elsayed et al. 2016). The larger modulation in firing rate within a separate neural subspace mapped to small amplitude movements could provide more accurate encoding of small, precise movements compared to a system that uses only a single subspace with smaller changes in firing rate. If we assume cortical neurons and corresponding subspaces may be constrained by similar levels of noise, the larger signal mapping to smaller motor output would have a better signal-to-noise ratio. A subspace best suited for precision movement execution while other subspaces may be better for faster movements or other functions. Thus, expanding the motor capabilities without increasing the number of neurons required. Further studies are needed to better define and verify these diverse and dynamic encoding patterns undefinable by a single cortical tuning model.

The relationship between movement amplitude and target size in Fitt's Law is well established (Fitts 1954, 1964; MacKenzie and Buxton 1992; Monk and Jagacinski 1985; Mottet et al. 1994; Murata 1999; Takeda et al. 2019); however, the neural mechanisms that underlie Fitt's Law remain elusive. A

single, linear cortical model is unable to reproduce the motor control necessary for increased precision with smaller movements. Varying neural dynamics allowed by multiple neural subspaces might help optimize movement parameters (direction, speed, etc.) to accomplish a given task. Additionally, the increased sensory integration with movement execution for a precision task are likely to alter the underlying neural mechanisms encoding movement. Better defining these neural states may lead to better descriptions of the dynamic response required for precision movements that lead to Fitt's Law.

Limitations

By relying on user-generated errors, we were not able to systematically vary which targets required corrective submovements. Additionally, with the center out task, there was greater variety of starting position for corrective submovements than for initial. This is represented in the multivariate regression analysis (Figure 3) with the larger position effect on firing rates for corrective submovements. Despite this added influence, the velocity vector remained the primary factor for neural firing rates for both types of submovements lending credence to velocity and not position being the dominantly encoded motor cortex feature.

Noise in the model fits could explain part of the observed shift in neural tuning. In a direct comparison of root-mean-square-error (RMSE) of neural firing rates between initial and corrective submovements, we saw similar noise variance for predictions of both types of submovements (Figure 3E, F). Thus, it appears the model error due to noise was similar for both the initial and corrective data and strengthens our conclusions that true changes in preferred direction and depth of modulation are occurring for initial and corrective submovements (Stein et al. 2005).

Further experiments with all combinations of large and small movements that are both instructed and corrective are needed to more completely quantify these shifts in tuning. In future studies tasks that produce small initial submovements as well as small natural corrective submovements should be explored. Instructed corrective submovements induced by a change of target location or external force affecting arm trajectory should also be explored.

Conclusion

In summary, we have quantified the shifts in neuronal tuning within the primary motor cortex for initial versus corrective submovements using a precision reaching center-out-task. These shifts resulted in poor description of both submovement types by a single fixed linear model. We have highlighted how these shifts when viewed from a neural space perspective give evidence of different neural subspaces being utilized for generation of initial and corrective submovements.

Acknowledgements

We would like to thank Dr. Marc Schieber and his laboratory at the University of Rochester Medical Center for assistance with data collection.

Funding Support

This research was supported by funding from National Institute of Neurological Disease and Stroke grant R00-NS-101127, the Frank and Evangeline Thompson Opportunity Fund, and the Steve Palermo Fund.

Abrams RA, **Meyer DE**, **Kornblum S**. Eye-Hand Coordination: Oculomotor Control in Rapid Aimed Limb Movements. *J Exp Psychol Hum Percept Perform* 16: 248–267, 1990.

Churchland MM, **Cunningham JP**, **Kaufman MT**, **Foster JD**, **Nuyujukian P**, **Ryu SI**, **Shenoy K V.**, **Shenoy K V.** Neural population dynamics during reaching. *Nature* 487: 51–56, 2012.

Churchland MM, **Shenoy K V.** Temporal complexity and heterogeneity of single-neuron activity in premotor and motor cortex. *J Neurophysiol* 97: 4235–4257, 2007.

Churchland MM, **Yu BM**, **Ryu SI**, **Santhanam G**, **Shenoy K V.** Neural variability in premotor cortex provides a signature of motor preparation. *J Neurosci* 26: 3697–3712, 2006.

Craik. CRAIK, K. J. W., Theory of the human operator in control systems. I. The operator as an engineering system , British Journal of Psychology, 38 (1947/1948) p.56. *Br J Psychol* 38: 56–61, 1948.

Desmedt JE, **Godaux E**. Ballistic contractions in man: characteristic recruitment pattern of single motor units of the tibialis anterior muscle. 264: 673–693, 1977.

Elliott D, **Hansen S**, **Grierson LEM**, **Lyons J**, **Bennett SJ**, **Hayes SJ**. Goal-Directed Aiming: Two Components but Multiple Processes. *Psychol Bull* 136: 1023–1044, 2010.

Elsayed GF, **Lara AH**, **Kaufman MT**, **Churchland MM**, **Cunningham JP**. Reorganization between preparatory and movement population responses in motor cortex. *Nat Commun* 7: 1–15, 2016.

Evarts E V. Relation of pyramidal tract activity to force exerted during voluntary movement. *J Neurophysiol* 31: 14–27, 1968.

Evarts E V., **Fromm C**. Sensory responses in motor cortex neurons during precise motor control. *Neurosci Lett* 5: 267–272, 1977.

Evarts E V, **Fromm C**, **Kroller J**, **Jennings VAA**, **Kröller J**, **Jennings VAA**. Motor Cortex control of finely graded forces. *J Neurophysiol* 49: 1199–1215, 1983.

Farina D, **Fosciand M**, **Merletti R**. Motor unit recruitment strategies investigated by surface EMG variables. *J Appl Physiol* 92: 235–247, 2002.

Fitts P. The Information Capacity of the Human Motor System in Controlling the Amplitude of Movement [Online]. *J Exp Psychol* 47: 381–391, 1954http://www2.psychology.uiowa.edu/faculty/mordkoff/InfoProc/pdfs/Fitts 1954.pdf.

Fitts P. Information capacity of discrete motor responses. *J Exp Psychol* 47: 381–391, 1964.

Flash T, **Hogan N**. The coordination of arm movements: An experimentally confirmed

mathematical model. *J Neurosci* 5: 1688–1703, 1985.

Freund HJ. Motor unit and muscle activity in voluntary motor control. *Physiol Rev* 63: 387–436, 1983.

Fu QG, **Flament D**, **Coltz JD**, **Ebner TJ**. Temporal encoding of movement kinematics in the discharge of primate primary motor and premotor neurons. *J Neurophysiol* 73: 836–854, 1995.

Fu QG, **Suarez JI**, **Ebner TJ**. Neuronal specification of direction and distance during reaching movements in the superior precentral premotor area and primary motor cortex of monkeys. *J Neurophysiol* 70: 2097–2116, 1993.

Georgopoulos AP, **Kalaska JF**, **Caminiti R**, **Massey JT**. On the relations between the direction of two-dimensional arm movements and cell discharge in primate motor cortex. *J Neurosci* 2: 1527–37, 1982.

Georgopoulos AP, **Kalaska JF**, **Massey JT**. Spatial trajectories and reaction times of aimed movements: Effects of practice, uncertainty, and change in target location. *J Neurophysiol* 46: 725–743, 1981.

Georgopoulos AP, **Schwartz AB**, **Kettner RE**. Neuronal Population Coding of Movement Direction. *Science (80-)* 233: 1416–1419, 1986.

Griffin DM, **Hoffman DS**, **Strick PL**. Corticomotoneuronal cells are "functionally tuned." *Science (80-)* 350: 667–670, 2015.

Hatsopoulos NG, **Xu Q**, **Amit Y**. Encoding of movement fragments in the motor cortex. *J Neurosci* 27: 5105–14, 2007.

Henneman E, **Somjen G**, **Carpenter D**. Functional significance of cell size in spinal motoneurons. *J Neurophysiol* 28: 560–580, 1965.

Inoue Y, **Mao H**, **Suway SB**, **Orellana J**, **Schwartz AB**. Decoding arm speed during reaching. *Nat Commun* 9: 1–14, 2018.

Kalaska JF, **Crammond DJ**. Cerebral cortical mechanisms of reaching movements. *Science (80-)* 255: 1517–1523, 1992.

Kalaska JF, **Scott SH**, **Cizek P**, **Sergio LE**. Control of reaching movements. *Curr Opin Neurobiol* 7: 849–859, 1997.

Karpowicz BM, **Ali YH**, **Wimalasena LN**, **Sedler AR**, **Keshtkaran MR**, **Bodkin K**, **Ma X**, **Miller LE**, **Pandarinath C**. Stabilizing brain-computer interfaces through alignment of latent dynamics. *bioRxiv* 2022.04.06.487388, 2022.

Kaufman MT, **Churchland AK**. Many paths from state to state. *Nat Neurosci* 19: 1541–1542, 2016.

Ma X, **Rizzoglio F**, **Perreault EJ**, **Miller LE**, **Kennedy A**. Using adversarial networks to extend brain computer interface decoding accuracy over time. *bioRxiv* 2022.08.26.504777, 2022.

MacKenzie IS, **Buxton W**. Extending Fitts' law to two-dimensional tasks. *Conf Hum Factors Comput Syst - Proc* 219–226, 1992.

Maynard EM, **Hatsopoulos NG**, **Ojakangas CL**, **Acuna BD**, **Sanes JN**, **Normann RA**, **Donoghue JP**. Neuronal interactions improve cortical population coding of movement direction. *J Neurosci* 19: 8083–8093, 1999.

Milner-Brown HS, **Stein RB**, **Yemm R**. Changes in firing rate of human motor units during linearly changing voluntary contractions. *J Physiol* 230: 371–390, 1973.

Milner TE. A model for the generation of movements requiring endpoint precision. *Neuroscience* 49: 487–496, 1992.

Milner TE, **Ijaz MM**. The effect of accuracy constraints on three-dimensional movement kinematics. *Neuroscience* 35: 365–374, 1990.

Mollazadeh M, **Aggarwal V**, **Davidson AG**, **Law AJ**, **Thakor N V.**, **Schieber MH**. Spatiotemporal variation of multiple neurophysiological signals in the primary motor cortex during dexterous reach-to-Grasp movements. *J Neurosci* 31: 15531–15543, 2011.

Monk D., **Jagacinski R.** Fitts' law in two dimensions with hand and head movements [Online]. *J. Mot. Behav.* : 77–95, 1985.http://e.guigon.free.fr/rsc/article/JagacinskiMonk85.pdf.

Moran DW, **Schwartz AB**. Motor Cortical Representation of Speed and Direction During Reaching. *J Neurophysiol* 82: 2676–2692, 1999.

Morasso P. Spatial Control of Arm Movements. *Exp Brain Res* 223–227, 1981.

Mottet D, **Bootsma RJ**, **Guiard Y**, **Laurent M**. Fitts' law in two-dimensional task space. *Exp Brain Res* 100: 144–148, 1994.

Mridha MF, **Das SC**, **Kabir MM**, **Lima AA**, **Islam MR**, **Watanobe Y**. Brain-computer interface: advancement and challenges. *Sensors* 21: 1–46, 2021.

Murata A. International Journal of Human- Extending Effective Target Width in Fitts ' Law to a Two-Dimensional Pointing Task Extending Effective Target Width in Fitts ' Law to a Two-Dimensional Pointing Task. *Int J* 11: 137–152, 1999.

Nicolelis MAL, **Ghazanfar AA**, **Faggin BM**, **Votaw S**, **Oliveira LMO**. Reconstructing the engram: Simultaneous, multisite, many single neuron recordings. *Neuron* 18: 529–537, 1997.

Novak K, **Miller L**, **Houk J**. The use of overlapping submovements in the control of rapid hand movements. *Exp Brain Res* 144: 351–364, 2002.

Port NL, **Lee D**, **Dassonville P**, **Georgopoulos AP**. Manual interception of moving targets. I. Performance and movement initiation. *Exp Brain Res* 116: 406–420, 1997.

Pratt J, **Chasteen AL**, **Abrams RA**. Rapid aimed limb movements: Age differences and practice effects in component submovements. *Psychol Aging* 9: 325–334, 1994.

Rouse, Adam G., Schieber MH. Condition-Dependent Neural Dimensions Progressively Shift during Reach to Grasp. *Physiol Behav* 176: 139–148, 2017.

Rouse AG, **Schieber MH**. Spatiotemporal distribution of location and object effects in primary motor cortex neurons during reach-to-grasp. *J Neurosci* 36: 10640–10653, 2016.

Rouse AG, **Schieber MH**. Condition-dependent neural dimensions progressively shift during
reach to grasp. *Cell Rep* 25: 3158–3168, 2018.

Rouse AG, **Schieber MH**, **Sarma S V.** Cyclic, Condition-Independent Activity in Primary
Motor Cortex Predicts Corrective Movement Behavior. *eNeuro* 9: 1–15, 2022.

Sainburg RL, **Ghez C**, **Kalakanis D**. Intersegmental dynamics are controlled by sequential
anticipatory, error correction, and postural mechanisms. *J Neurophysiol* 81: 1045–1056, 1999.

Scott SH. Optimal feedback control and the neural basis of volitional motor control. *Nat Rev
Neurosci* 5: 532–544, 2004.

Soechting JF. Effect of target size on spatial and temporal characteristics of a pointing
movement in man. *Exp Brain Res* 54: 121–132, 1984.

Stein RB, **Gossen ER**, **Jones KE**. Neuronal variability: Noise or part of the signal? *Nat Rev
Neurosci* 6: 389–397, 2005.

Takeda M, **Sato T**, **Saito H**, **Iwasaki H**, **Nambu I**, **Wada Y**. Explanation of Fitts' law in
Reaching Movement based on Human Arm Dynamics. *Sci Rep* 9: 1–12, 2019.

Umair A, **Ashfaq U**, **Khan MG**. Recent trends, applications, and challenges of brain-computer
interfacing (BCI). *Int J Intell Syst Appl* 9: 58–65, 2017.

Wang W, **Chan SS**, **Heldman DA**, **Moran DW**. Motor cortical representation of position and
velocity during reaching. *J Neurophysiol* 97: 4258–70, 2007.

Woodworth RS. The accuracy of voluntary movement. *J Nerv Ment Dis* 26: 743–752, 1899.

556