A Compound that Inhibits Glycolysis in Prostate Cancer Controls Growth of Advanced Prostate Cancer

Takuma Uo[1,][*], Kayode K. Ojo[2†], Cynthia C. Sprenger[1†], Kathryn Soriano Epilepsia[1], B. Gayani K. Perera[3], Mamatha Damodarasamy[1], Shihua Sun[1], Soojin Kim[1], Hannah H. Hogan[1], Matthew A. Hulverson[2], Ryan Choi[2], Grant R. Whitman[2], Lynn K. Barrett[2], Samantha A. Michaels[2], Linda H. Xu[1], Vicky L. Sun[4], Samuel L.M. Arnold[1,4], Haley J. Pang[1], Matthew M. Nguyen[1], Anna-Lena B.G. Vigil[5], Varun Kamat[6], Lucas B. Sullivan[5, 7], Ian R. Sweet[6], Ram Vidadala[3], Dustin J. Maly[3], Wesley C. Van Voorhis[2,][*], Stephen R. Plymate[1, 8][*]

Affiliations:

[1]Department of Medicine, Division of Gerontology and Geriatric Medicine, University of Washington; Seattle, Washington 98109, USA.

[2]Department of Medicine, Division of Allergy and Infectious Disease, Center for Emerging and Reemerging Infectious Disease (CERID), University of Washington; Seattle, Washington 98109, USA.

[3]Department of Chemistry, University of Washington; Seattle, Washington 98195, USA.

[4]Department of Pharmaceutics, University of Washington; Seattle, Washington 98195, USA.

[5]Human Biology Division, Fred Hutchinson Cancer Center; Seattle, Washington 98109, USA.

[6]Department of Medicine, Division of Metabolism, Endocrinology and Nutrition, Diabetes Center, University of Washington; Seattle, Washington 98109, USA.

[7]Department of Biochemistry, University of Washington; Seattle, Washington 98195, USA.

[8]Geriatrics Research Education and Clinical Center, VA Puget Sound Health Care System; Seattle, Washington 98108, USA.

* Corresponding Authors   † These authors contributed equally to this paper

Running title: Novel antiglycolytic agents for prostate cancer therapy

Abbreviations: AA: antimycin A, ACC: acetyl-CoA Carboxylase, ACN: acetonitrile,

adeno-PCa: adenocarcinoma prostate cancer, ADME: adsorption-distribution-

metabolism-excretion, AMPK: AMP-activated protein kinase, AR: androgen receptor,

ARSi: androgen receptor signaling inhibitors, AUC: area under the curve, BKI: bumped

kinase inhibitor, BKIDCs: bumped kinase inhibitor (BKI) derived compounds, BSA:

bovine serum albumin, CAR: cell-associated radioactivity, Cl: clearance, CMC:

Chemistry Manufacturing and Controls, CRPC: castration-resistant prostate cancer,

CYP: cytochrome P450, CytB: cytochalasin B, DAPI: 4,6-diamidino-2-phenylindole,

DAB: 3,3'-diaminobenzidine, 2-DG: 2-deoxyglucose, 2-DG-6-P: 2-DG-6-phosphate,

DHAP: dihydroxyacetone phosphate, DPBS: Dulbecco's phosphate-buffered saline,

ECAR: extracellular acidification rate, FBS: fetal bovine serum, FCCP: carbonyl cyanide

4-(trifluoromethoxy)phenylhydrazone, FDA: Food and Drug Administration, FDG:

fluorodoxyglucose, GLP: Good Laboratory Practice, G6P: glucose-6-phosphate, hERG:

human ether-a-go-go related gene, HK: hexokinase, hr: human recombinant, IC_{50}: 50%

inhibition values, IND: Investigative New Drug, IRS-1: insulin receptor substrate-1, ITS:

Insulin-Transferrin-Selenium, KO: knock-out, KRB: Krebs-Ringer bicarbonate solution,

LC-MS: liquid chromatography coupled with mass spectrometry, MTS: 3-(4,5-

dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium,

inner salt, NE: neuroendocrine, OCR: oxygen consumption rate, Oligo: oligomycin 3-

OMG: 3-O-methyl D-glucose, OXPHOS: oxidative phosphorylation, OXPHOSi:

OXPHOS inhibitors, P_{app}: apparent permeability coefficient, PARP: poly ADP ribose

polymerase, PCa: prostate cancer, PDL: poly-D-lysine, PDX: patient-derived xenograft,

PEP: phosphoenolpyruvate, PG: phosphoglycerate, PK: pharmacokinetics, PKM2:

pyruvate kinase M2, PP: pyrazolopyrimidine, PrP: pyrrolopyrimidine, RLU: relative

luminescence units, RPPA: reverse-phase protein arrays, RPS6: ribosomal S6 protein,

SCID: severe combined immunodeficient, SD: standard deviation, SEM: standard error

of mean, SO: sodium oxamate, UPLC-HRMS: ultra-high-performance liquid

chromatography-high resolution mass spectrometry, V: volume of distribution

Corresponding authors: Wesley C. Van Voorhis, Department of Medicine, University

of Washington, MS 358061, 750 Republican St., Rm E-606, Seattle WA USA 98109-

4766, E-mail: WVanVoorhis@medicine.washington.edu; Stephen R. Plymate, University

of Washington School of Medicine, Box 358055, 850 Republican St., Seattle WA USA

98109-4766, E-mail: splymate@uw.edu; Takuma Uo, University of Washington, 850

Republican St. Rm S-325, Seattle WA USA 98109-4766, E-mail: tuo@uw.edu

Conflict of interest disclosure statement: SRP is the President, WCVV is the Chief

Executive Officer, DJM is the Chief Scientific Officer, and KKO is the Secretary of

ProsTech Inc. ProsTech Inc. was created to commercialize therapies for prostate

cancer, but as yet, has no commercial claims on BKIDCs or any other potential

commercial products. Other authors declare that they have no competing interests. The

patent numbers related to this article are as follows. Use Patent in Cancer Field: US

10350211 issued, Composition of Matter Patents: US 10632122 and US 10307425

both issued; and US patent application 17/758,577 pending. None of these four are

sublicensed for cancer or producing royalties.

ABSTRACT

Purpose: Metastatic castration-resistant prostate cancer remains incurable regardless of

recent therapeutic advances. Prostate cancer tumors display highly glycolytic

phenotypes as the cancer progresses. Non-specific inhibitors of glycolysis have not

been utilized successfully for chemotherapy, because of their penchant to cause

systemic toxicity. This study reports the preclinical activity, safety, and pharmacokinetics

of a novel small molecule preclinical candidate, BKIDC-1553, with antiglycolytic activity.

Experimental design: We tested a large battery of prostate cancer cell lines for inhibition

of cell proliferation, in vitro. Cell cycle, metabolic and enzymatic assays were used to

demonstrate their mechanism of action. A human PDX model implanted in mice and a

human organoid were studied for sensitivity to our BKIDC preclinical candidate. A

battery of pharmacokinetic experiments, absorption, distribution, metabolism, and

excretion experiments, and in vitro and in vivo toxicology experiments were carried out

to assess readiness for clinical trials.

Results: We demonstrate a new class of small molecule inhibitors where antiglycolytic

activity in prostate cancer cell lines is mediated through inhibition of hexokinase 2.

These compounds display selective growth inhibition across multiple prostate cancer

models. We describe a lead BKIDC-1553 that demonstrates promising activity in a

preclinical xenograft model of advanced prostate cancer, equivalent to that of

enzalutamide. BKIDC-1553 demonstrates safety and pharmacologic properties

consistent with a compound that can be taken into human studies with expectations of a

good safety margin and predicted dosing for efficacy. Conclusion: This work supports

testing BKIDC-1553 and its derivatives in clinical trials for patients with advanced

prostate cancer.

INTRODUCTION

Prostate cancer (PCa) is responsible for the deaths of more than 375,000 men worldwide and the second most common solid tumor in men(1). Although primary therapies such as surgery or radiotherapy appear to cure many men diagnosed with PCa, the disease becomes incurable once becomes metastatic(2, 3).

Treatment modalities for PCa have expanded significantly over the past decade with development of potent androgen receptor signaling inhibitors (ARSi) (e.g., abiraterone, enzalutamide)(2, 4). Additionally, tumors with loss of function mutations in homologous recombination repair pathway genes (e.g., BRCA1/2) define a subgroup of patients who can benefit from treatment with poly ADP ribose polymerase (PARP) inhibitors and possibly other DNA damaging therapies such as carboplatin(5). More recently, radionuclide therapies such as Radium 223 and PSMA targeting with Lutetium 177 have provided new avenues of therapy(6, 7). Although these therapies have been demonstrated to improve overall survival, the improvement is incremental, and resistance develops in the majority of cases(2, 8). One avenue of therapy for castration-resistant prostate cancer (CRPC) that has yet to realize its full potential involves the development of agents designed to selectively inhibit tumor metabolism(9, 10). To date, targeting tumor metabolism, particularly glycolysis or mitochondrial metabolism, has been stymied due to excessive toxicity and/or lack of efficacy(11).

Changes to cell metabolism is a hallmark of cancer that drives and supports oncogenesis(12-14). Both tumor genetics and tissue context define cancer dependence on specific metabolic pathways, with the tumor ultimately undergoing progression and being driven by combinations of oxidative phosphorylation, glycolysis, and potentially

fatty acid metabolism(15). One hundred years have passed since Otto Warburg's

pioneering work to report the dominance of aerobic glycolysis in tumors(16).

Nevertheless, non-specific antiglycolytic agents such as 2-deoxyglucose (2-DG), have

never been clinically translated as they not only affect tumor cells but also inhibit normal

proliferating cells(15, 17).

While glycolysis is not a significant driver in primary PCa, in more aggressive

lethal forms of the disease glycolysis is a major driver of tumor progression especially

since the tumor exists in a hypoxic environment(18-21). Challenging our ability to

develop drugs against this pathway are multiple enzymatic isoforms in the glycolytic

pathway that perform overlapping functions, complicating specific targeting(15).

Accordingly, flexibility and redundancy among metabolic genes could shape the

landscape of positive and negative impact on targeting tumor metabolism(9, 15). As far

as glycolysis is concerned, isoform specific contribution to tumor progression is

exemplified by pyruvate kinase M2 (PKM2) and hexokinase 2 (HK2), where activation

and inhibition have been shown to reduce tumor progression, respectively, despite

continued expression of corresponding isoforms such as PKM1 and HK1(22, 23). HK2

has been reported to be essential for PTEN/p53 deficient CRPC growth(24).

Previous protein kinase profiling of bumped kinase inhibitor (BKI) derived

compounds (BKIDCs) (BKIDC-1369, 1553, 1561, and 1649) revealed potential human

kinase targets, including Dual specificity mitogen-activated protein kinase kinase 2

(MEK2), human receptor-interacting serine/threonine-protein kinase 2 (RIPK2) and

protein kinase D (PKD1-3)(25). Given that other studies have shown adenocarcinoma

prostate cancer (adeno-PCa) cell lines are inhibited by pan-PKD inhibitors – irrespective

of AR status – we set out to screen for cell types that might be affected by BKIDC

compounds (26, 27). However, as we will show in this study, BKIDCs do not appear to

act against PCa growth by inhibiting protein kinases.

In this study, we report on the discovery of a novel series of BKIDCs, that display

selective inhibition of PCa growth and that are associated with a cessation of glycolysis

immediately on exposure to the compounds. Furthermore, we describe a lead

compound (referred to as BKIDC-1553) that demonstrates promising pharmacological

and safety properties and activity in a mouse preclinical model of advanced PCa.

MATERIALS AND METHODS

Data availability:

Derived data supporting the findings of this study are available upon request.

Information and requests for resources and reagents should be directed to and will be

fulfilled by lead contacts Stephen Plymate (splymate@uw.edu), Wesley Van Voorhis

(wesley@uw.edu), and Takuma Uo (tuo@uw.edu). Synthesis protocols of all unique,

non-FDA chemical compounds, synthesized for this publication are given below in the

section **Synthesis of PP and PrP based BKIDCs.**

Cell culture

LNCaP clone FGC (obtained in 2018. CRL-1740, RRID:CVCL_1379), C4-2 (obtained in

November 2019. CRL-3314, RRID:CVCL_4782), C4-2B (obtained in November 2019.

CRL-3315, RRID:CVCL_4784), VCaP (obtained in 2018. CRL-2876,

RRID:CVCL_2235), DU145 (obtained in 2018. HTB-81, RRID:CVCL_0105), PC3

(obtained in 2018. CRL-1435, RRID:CVCL_0035), HepG2 (obtained in May 2013, HB-

8065, RRID:CVCL_0027), HFF-1 (obtained in August 2020. SCRC-1041,

RRID:CVCL_3285), and lymphoblastoid WIL2NS (obtained in April 2011, CRL-8155,

RRID:CVCL_2761) were obtained from the American Type Culture Collection (ATCC).

293FT cell line was obtained in June 2016 from ThermoFisher Scientific (Invitrogen™,

Cat# R70007, RRID:CVCL_6911). LAPC4 (RRID:CVCL_4744) was obtained from Dr.

Charles Sawyers (Memorial Sloan Kettering Cancer Center, New York NY) through Dr.

Peter Nelson (Fred Hutchinson Cancer Center, FHCC) in October 2019. U2OS cells

(RRID:CVCL_0042) from Dr. Raymond Monnat (University of Washington) in October

2018, NCI-H660 in January 2020 (RRID:CVCL_1576) and LNCaP^{APIPC} from Dr. Peter

Nelson in November 2011(28), LNCaP95 (obtained in 2012. RRID:CVCL_ZC87) from

Dr. Alan Meeker (Johns Hopkins University), MSKCC EF1 derived from the organoid

line MSKCC-CaP4(29) from John K. Lee (FHCC) in January 2020. LTL331R cells

established from LTL331R tumor(29) were obtained from Dr. Yuzhuo Wang at

Vancouver Prostate Centre (Vancouver BC, Canada) in March 2023.

Cells were maintained in RPMI 1640 (Gibco™, Cat# 21-870-092) (LNCaP, C4-2, C4-

2B, DU145, PC3, MSKCC EF1, HepG2, WIL2NS, 293FT) supplemented with GlutaMax-

I (Gibco™, Cat# 35050061), DMEM (Gibco™, Cat# 10569-010) (VCaP, U2OS, HFF-1),

and IMDM (Iscove's Modified Dulbecco's Medium, Gibco™, Cat# 21056023) (LAPC4)

supplemented with 5% fetal bovine serum (FBS) (Gibco™, Cat# 16000-044), 100

units/mL penicillin, and 0.1 mg/mL streptomycin (Gibco™, Cat# 15140122). LNCaP95

cells were grown in phenol-red-free RPMI medium (Gibco™, Cat# 11835030)

supplemented with 5% charcoal stripped serum (Gibco™, Cat# 12676029). NCI-H660

was cultured in HITES medium (RPMI 1640 medium plus 5 µg /mL insulin, 10 µg/mL

transferrin, 30 nM sodium selenite (ITS (Insulin-Transferrin-Selenium), Corning, Cat#

25-800-CR), 10 nM hydrocortisone (Sigma Aldrich, Cat# H0135), 10 nM β-estradiol

(Sigma Aldrich, Cat# E2257)) with 10% FBS. The LTL331R cells were maintained in

RPMI1640 (ATCC modification: Gibco™, Cat# A1049101) containing 5% FBS, 1% ITS

(Gibco™, Cat# 41400-045), 10 nM hydrocortisone, 10 nM β-estradiol, 10 µM Y27632

(Tocris Bioscience, Cat# 1254), and 1% Cultrex Basement Membrane Extract,

PathClear (R&D Systems, Cat# 3432-010).

All cell lines were routinely tested for mycoplasma contamination with Universal

Mycoplasma Detection Kit (ATCC, Cat# 30-1012K ™) and used within 20 passages of

receipt. The latest tests relevant to this study were done in December 2018 (VCaP,

DU145, U2OS), January 2019 (293FT), May 2019 (LNCaP, LNCaP95, LNCaP^{APIPC},

PC3), November 2019 (C4-2, C4-2B, LAPC4), February 2020 (NCI-H660, MSKCC

EF1), September 2021 (HepG2), October 2021 (CRISPR-engineered LNCaP lines),

October 2023 (HFF-1, WIL2NS), December 2023 (LTL331R).

Cell authentication was carried out with ATCC FTA Sample Collection Kit for Human

Cell Authentication Service (ATCC 135-XV-20 ™). The latest test date was July 2023

for LNCaP (test specimen from cells collected in July 2021), LNCaP95 (July 2021),

U2OS (January 2019), PC3 (January 2019), VCaP (August 2019), DU145 (December

2018), and HepG2 (June 2020) and October 2023 for WIL2-NS (July 2020) and NCI-

H660 (February 2020).

Three cell lines were used for pharmacological characterization of BKIDC-1553. MDR1-

MDCKII (RRID:CVCL_IZ19) was used sed to assess transport of BKIDC-1553 in both

directions (apical to basolateral (A-B) and basolateral to apical (B-A)) across the cell

monolayer and monitored for key markers of MDR1 transporter expression.

Mycoplasma was monitored with MycoAlert® PLUS Mycoplasma Detection Kit (Lonza,

Cat# LT07-705) in April 2022 and the cell line was used in February 2023. The cell line

was used on passage 9 from thaw. Caco-2 cells at passage 19 (RRID:CVCL_0025)

were obtained from ATCC. Performance of the Caco-2 line in monolayer permeability

experiments was characterized by comparison of results for reference compounds to

published data. Cells were confirmed to be mycoplasma-free based on a luminescence-

based enzymatic assay (MycoAlert®, Lonza, Cat# LT07-118) conducted within 1 week

of the commencement and completion of the experiments. Cells used in the

experiments were at passage 37 after thawing stocks at passage 30.

Metabolomic flux assay

LNCaP95, LNCaP, and DU145 cells were plated at 500 K cells per well on 6-well plates

(Thermo Scientific™, Cat# 140675) coated with 10 µg/mL poly-D-lysine (PDL) (Sigma

Aldrich, Cat# A-003 M), overnight in media containing unlabeled 10 mM D-glucose and

2 mM L-glutamine. Cells were treated in triplicate with drug or vehicle for 30 min. Cells

were then quickly washed two times with RPMI media with L-glutamine and without

glucose (Gibco™, Cat# 11879020). Media was then changed to fresh RPMI with 10 mM

[U]^{13}C-glucose (Cambridge Isotope Laboratories, CLM 13961369) and unlabeled 2 mM

glutamine in the presence of different concentrations of BKIDC-1553-N-Me. DMSO

(Sigma-Aldrich, Cat# D5879) was used as vehicle control. Cells were then harvested 30

min after treatment with [U]^{13}C-glucose and inhibitors. Cells were then quickly washed

twice with ice cold blood bank saline and quenched with 500 μL HPLC-grade 80%

methanol (Fisher Chemical, Cat# A4561) containing 2.5 μM valine D8 extraction

standard. Samples were then centrifuged at 4°C at 17,000 g for 5 min. 400 μL of

supernatant was transferred to a new microfuge tube and centrifuged at 4°C at 17,000 g

for 5 min. 350 μL of supernatant was then transferred to a new microfuge tube and dried

down with a Labonco Centrivap Centrifugal Vacuum Concentrator at 4°C and placed at

-80°C until time of analysis. Samples were then re-constituted with 50 μL of 80% HPLC-

grade methanol, vortexed for 10 min at 4°C, and centrifuged at 17,000g for 5 min at

4°C. 40 μL of supernatant was transferred to a liquid chromatography-mass

spectrometry (LC-MS) vial for measurement by LC-MS. Metabolite quantitation was

performed using a Q Exactive HF-X Hybrid Quadrupole-Orbitrap Mass Spectrometer

equipped with an Ion Max API source and H-ESI II probe, coupled to a Vanquish Flex

Binary UHPLC system (Thermo Scientific). Mass calibrations were completed at a

minimum of every 5 days in both the positive and negative polarity modes using LTQ

Velos ESI Calibration Solution (Pierce). Samples were chromatographically separated

by injecting a sample volume of 1 μL into a SeQuant ZIC-pHILIC Polymeric column (150

x 2.1 mm 5 μM, EMD Millipore). The flow rate was set to 150 μL/min, autosampler

temperature set to 10°C, and column temperature set to 30°C. Mobile Phase A

consisted of 20 mM ammonium carbonate and 0.1% (v/v) ammonium hydroxide, while

Mobile Phase B consisted of 100% acetonitrile (ACN). The sample was gradient eluted

(%B) from the column as follows: 0-20 min: linear gradient from 85% to 20% B; 20-24

min: hold at 20% B; 24-24.5 min: linear gradient from 20% to 85% B; 24.5 min-end: hold

at 85% B until equilibrated with ten column volumes. Mobile Phase was directed into the

ion source with the following parameters: sheath gas = 45, auxiliary gas = 15, sweep

gas = 2, spray voltage = 2.9 kV in the negative mode or 3.5 kV in the positive mode,

capillary temperature = 300°C, RF level = 40%, auxiliary gas heater temperature =

325°C. Mass detection was conducted with a resolution of 240,000 in full scan mode,

with an AGC target of 3,000,000 and maximum injection time of 250 msec. Metabolites

were detected over mass range of 70-1050 m/z in full scan mode with polarity switching.

Quantitation of all metabolites was performed using Tracefinder 4.1 (Thermo Fisher

Scientific RRID:SCR_023045) referencing an in-house glycolytic metabolite standards

library using ≤ 5 ppm mass error.

Glucose uptake assay

Luciferase-based assay: LNCaP cells were plated at 240 K cells/well in PDL-coated

24-well plates (Corning, Cat# 353226). 24 h later, medium was changed to serum and

glucose-free medium containing BKIDC at 20 µM or cytochalasin B (CytB, C6762

Sigma-Aldrich, Cat# C6762) at 50 µM for 30 min prior to the experiment. Cells were

then rinsed and incubated media containing 2-DG and BKIDC or CytB for 10 min.

Glucose uptake was measured using the Glucose Uptake-Glo™ Assay (Promega, Cat #

J1341) according to the manufacturer's protocol. This luciferase-based assay measures

2-DG-6-phosphate (2-DG-6-P) which accumulates as a dead-end product after

hexokinase-mediated conversion of 2-DG. Luminescence signals were measured on

96-well plate (Corning, Cat# 3915) on Promega GloMax® plate reader

(RRID:SCR_015575). Either BKIDC-1553-N-Me or CytB was added after cellular

harvest, to demonstrate these compounds did not interfere with the assay to evaluate 2-

DG accumulation. The data were presented as mean relative luminescence units (RLU)

± standard deviation (SD) (n=3).

Radiometric assay: Uptake of ^{3}H-3-O-methylglucose (American Radiolabeled

Chemicals, St. Louis, MO) and ^{3}H-2-deoxyglucose (Perkin Elmer, Waltham, MA) was

measured as carried out previously(30, 31). Cells were incubated in 12 x 75 mm

polypropylene test tubes containing Krebs-Ringer bicarbonate solution (KRB) (with 5

mM bicarbonate and 0.1% BSA) at 37°C for 30 min in a shaking water bath. Cell

densities were 2 x10^6 cells per 200 µL for all cell types. Radiolabeled molecules (0.175

µCi) were added to the tubes using an Eppendorf Repeater pipet and the cells were

further incubated in the presence of a radiotracer for 30 min. (Experiments

demonstrated that between 10 and 30 min, uptake of tracer was linear). The specific

activities and the concentrations in the incubation media of the radiotracers were as

follows: [^{3}H]3-O- MG, 90 Ci/mmol, 24 nM; [^{3}H]2-deoxyglucose, 20 Ci/mmol, 119 nM.

Accumulation of radiolabel was determined by separating the cell-associated

radioactivity (CAR) from the free radioactivity by transferring the cell suspension to a 0.4

mL centrifuge tube (USA Scientific, Ocala, FL) containing a layer of oil consisting of

1:37.5 n-dodecane:bromo-dodecane (Sigma–Aldrich, St. Louis, MO) and spinning at

12,535 g for 8 s (Beckman E centrifuge, Beckman Coulter, Fullerton, CA). The tube was

placed briefly in liquid nitrogen, cut through the radioactive-free oil layer with a razor

blade, and the bottom portion of the tube containing the cell pellet was placed into a 7

mL glass scintillation vial. The ^{3}H labels were counted using a Beckman liquid

scintillation counter (Model LS6500) after adding 5 mL of Ecolume liquid scintillation

cocktail (MP Biomedicals, Cat# 0188247001) and vortexing the vials. Values of non-

specific association of radioactivity was determined by performing the assay in the

presence of 5 mM glucose at 4°C. The CAR data for each sample was normalized by

subtracting off the non-specific values, and then dividing by the cell number and the

dose. Quench correction for samples was not needed since counts were normalized to

the dose and quenching did not vary from sample to sample.

In vitro AMPK activity measurement with total cell lysates

LNCaP95 cells were grown on 24-well plate for 24 h and treated with BKIDC at 20 µM

for 1 h. AMPK activity in total cell lysates was measured using CycLex AMPK Kinase

Assay kit (MBL International Corporation, Cat# CY-1182) according to the

manufacturer's instructions. Briefly, total cell lysates were prepared in lysis buffer (20

mM Tris-HCl, pH 7.5, 250 mM NaCl, 10% glycerol, 0.5% NP-40, 1 mM DTT)

supplemented with Halt Protease Inhibitor Cocktail and Halt Phosphatase Cocktail

(Thermo Scientific™, Cat# 78440) and incubated for 30 min at 30°C in a pre-coated

plate with an AMPK substrate peptide corresponding to mouse insulin receptor

substrate-1 (IRS-1)(supplied as part of kit). AMPK activity was measured by monitoring

the phosphorylation of Ser-789 in IRS-1 using an anti-mouse phospho-Ser-789 IRS-1

monoclonal antibody (AS-4C4: supplied as part of kit) and peroxidase-coupled anti-

mouse IgG antibody. Conversion of the chromogenic substrate tetramethylbenzidine

was quantified by absorbance measurement at 450 nm in BioRad iMark™ Microplate

Absorbance Reader (RRID:SCR_023799).

Real-time cellular bioenergetic assay

We used Agilent XFe24 analyzer (RRID:SCR_019539) to measure extracellular

acidification rate (ECAR) and oxygen consumption rate (OCR) which reflect the rates of

glycolysis and oxidative phosphorylation, respectively. Cells were plated at 40 K

cells/well in Seahorse XF24 V7 PS Cell Culture Microplates (Agilent, Cat# 10777-004)

coated with 10 µg/mL PDL and incubated overnight. The culture medium was then

replaced by Seahorse XF RPMI medium (Agilent, Cat# 103576-100) and cells were pre-

incubated in a CO_2-free incubator at 37°C for 1 h prior to metabolic analysis. The effect

of BKIDC was evaluated by pre-treating cells with 20 µM BKIDC during 1h-pre-

incubation period with and without glucose prior to mitochondrial and glycolytic stress

tests, respectively. Mitochondrial stress test was carried out by sequentially injecting

oligomycin (Sigma, Cat# O4876), carbonyl cyanide 4-

(trifluoromethoxy)phenylhydrazone (FCCP: Sigma, Cat# C2920), and rotenone (Sigma,

Cat# R8875)/antimycin A (Sigma, Cat# A8674). Glycolytic stress test was done by

sequential injection of glucose (Gibco™, Cat# A2494001), oligomycin, and 2-deoxy-D-

glucose (2-DG: Sigma, Cat# D8375). The acute effect of BKIDC was assessed by

directly adding BKIDC through the injection port. The bioenergetic profiles were

analyzed from at least 3 biological replicates. Results were automatically generated and

analyzed by Seahorse Wave Desktop Software, which were exported as files formatted

for GraphPad Prism (RRID:SCR_002798).

Measurement of intracellular ATP levels

50 K cells were plated on 48-plate (Corning, Cat# 353078) and grown overnight. Cells

were treated with the indicated compounds for 2 h unless specified. Metabolic inhibitors

used include: oligomycin, antimycin A, FCCP, rotenone, 2-DG, and sodium oxamate

(Cayman Chemical, Cat# 565-73-1). After removing culture media, 80 µL of the

CellTiter-Glo 2.0 Cell Viability Assay kit (Promega, Cat# G9241) was added to the cells.

Intracellular ATP levels under each condition were assessed on 96-well plate (Corning,

Cat# 3915) on GloMax® plate reader.

CRISPR/Cas9-mediated genome editing.

We used a CRISPR/Cas9 approach to establish a stable knockout LNCaP cell lines

deficient of either HK1 or HK2(32). sgRNA protospacer of HK1 (HK1-1:

GGTGAGGTGGAAATGAGCCA; HK1-3: GGAGGGCAGCATCTTAACCA) and HK2

(HK2-1: GATGCGCCACATCGACATGG; HK2-2: TAAGCGGTTCCGCAAGGAGA) was

cloned into lentiCRISPR v2 construct (RRID:Addgene_52961) using a protocol available

online (Zhang_lab_LentiCRISPR_library_protocol.pdf). gRNA sequence (control:

GTATTACTGATATTGGTGGG) was used to prepare the lentiviral construct for a non-

targeting control(33). The oligonucleotides were commercially synthesized at Eurofins

Scientific. 293FT cells were seeded on 6-well plates coated with poly-D-lysine. The

following day, 293T cells were co-transfected with lentiCRISPR v2 construct and the

second-generation lentiviral packaging plasmids pMD2.G for VSV-G

(RRID:Addgene_12259) and pCMVR8.74 for GAG, POL, TAT, and REV

(RRID:Addgene_22036) using Lipofectamine 3000 reagent (Invitrogen™, Cat#

L3000015). Lentivirus-containing supernatant was collected at 12 and 24 h after

transfection, filtered through a 0.45-µm filter (Cytiva, Cat# WHA-9916-1304), and used

to infect LNCaP as a recipient cell line, overnight in the presence of 5 µg/mL polybrene

(Sigma, Cat# 107689). The infected cells were selected with 5 µg/mL of puromycin

(Gibco™, A1113803) for 3 days. Selected cells were immediately used as pool cell

populations to evaluate glycolytic capacities in relation to BKIDC. On the other hand, to

isolate morphologically single colonies, they were also plated at the density of 0.5

cell/well in 96-well plates in the conditioned media from native LNCaP line. Resulting

colonies were screened by Western blot analysis for loss of either HK1 or HK2

expression. After a second confirmatory Western blot analysis, the genomic junctions

flanking corresponding sgRNA binding sites were amplified with sequence-specific

primers by PCR using Platinum Taq DNA Polymerase and the amplicons were cloned

into pCR™2.1-TOPO™ plasmid with TOPO™ TA Cloning™ Kit (Invitrogen™, Cat#

450641) and Sanger-sequenced using the T7 promoter primer

(TAATACGACTCACTATAGGG) and M13 Reverse primer (CAGGAAACAGCTATGAC)

at GENEWIZ (South Plainfield, NJ). Particular isolates (HK11-9, HK13-5, HK21-5,

HK22-1) were chosen for further experiments. Pool cell population which was infected

with non-targeting control virus served as control cell lines for the experiments with both

clonal cell lines and pool cell populations.

Cell proliferation assay

Cell   proliferation   was   determined   by   MTS   [3-(4,5-dimethylthiazol-2-yl)-5-(3-

carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium,   inner   salt]   assay   using

CellTiter 96® AQueous Non-Radioactive Cell Proliferation Assay (MTS) kit (Promega,

Cat# G3582) at 72 h after BKIDCs were added for all cell lines, except for VCaP at 120

h. Each cell line was tested in 96 well plates at 5 K cells/well in the culture media. The

data shown are representative of at least two independent biological experiments,

normalized to the values obtained from cells grown in medium containing 0.1% DMSO

vehicle and presented as mean ± standard deviation (SD) (n=3-5).

CellTiter-Glo® 3D Cell Viability Assay

LTL331R cells were plated on 96-well plate (Thermo Scientific, Cat# 167425) at 15 K

cells/well. At 2 days after plating, cells were treated with either BKIDC-1553 (1.25-20

μM) or enzalutamide (1.25-20 μM) using DMSO as vehicle control. At 72 h after

treatment, cell proliferation was determined using CellTiter-Glo™ 3D Cell Viability Assay

(Promega, Cat# G9683.) on GloMax® plate reader. The data shown are representative

of at least two independent biological experiments, normalized to the values obtained

from cells grown in medium containing 0.1% DMSO vehicle and presented as mean ±

standard deviation (SD) (n=3-4).

Crystal Violet Assay

Cell proliferation of LNCaP95 and PC3 was assessed by crystal violet assay at 72 h

after BKIDCs were added. Each cell line was tested in 96 well plates (Thermo

Scientific™, Cat# 167425) at 5 K cells/well in the culture media. Cells were fixed with

5% glutaraldehyde (Sigma-Aldrich, Cat# G6257) for 5 min and stained with 0.5% crystal

violet (Sigma-Aldrich, Cat# C3886) solution in 20% methanol (VWR Chemicals BDH®,

Cat# BDH1135) for 30 min followed by rinsing with distilled water 3 times. After

completely air-drying the plate overnight, 50 μL of 10% acetic acid was added to each

well to dissolve the cell-bound crystal violet. 10 min later, absorption was determined at

590 nm in BioRad iMark™ Microplate Absorbance Reader (RRID:SCR_023799). The

data shown are representative of at least three independent biological experiments,

normalized to the values obtained from cells grown in medium containing 0.1% DMSO

vehicle and presented as mean ± standard deviation (SD) (n=3-5).

Cell Cycle Analysis

Analysis was performed as previously described(34). Cells treated with BKIDCs for 30,

48, or 72 h were collected and resuspended in a solution of 10 μg/mL 4,6-diamidino-2-

phenylindole (DAPI) and 0.1% Nonidet P-40 detergent in a Tris buffered saline. The

suspension was analyzed on the LSR II flow cytometer (RRID:SCR_002159, BD

biosciences), with ultraviolet or violet excitation and DAPI emission collected at >450

nm. DNA content and cell cycle were analyzed as previously described using the

software program MultiCycle(34)(Phoenix Flow Systems, San Diego, CA).

Western blot analysis

Cells were lysed in M-PER Mammalian Protein Extraction Reagent (Thermo

Scientific™, Cat# 78501) supplemented with Protease Inhibitor Cocktail and Halt

Phosphatase Cocktail (Thermo Scientific™, Cat# 78440). Protein measurement was

determined using Pierce™ BCA Protein Assay Kits BCA assay (Thermo Scientific™,

Cat# 23225) using Pierce™ Bovine Serum Albumin Standard Pre-Diluted Set (Thermo

Scientific™, Cat# 23208). Samples were run on 4–15% Mini-PROTEAN® TGX™

Precast Protein Gels (Bio-Rad, Cat# 4561084) and proteins transferred to Immun-Blot®

PVDF Membrane (Bio-Rad, Cat# 1620177) using Bio-Rad TransBlot Turbo transfer

system (RRID:SCR_023156). Tris-buffered saline (TBS: 20 mM Tris-HCl, pH 7.5,

containing 0.15 M NaCl) with 0.3% Tween 20 (Fisher Scientific, Cat# BP337-100)(TBS-

T) was used. Membranes were blocked with the blocking buffer (TBS-T with 5% nonfat

milk (LabScientific, Cat# M-0842). Blots were incubated with appropriate antibodies in

the blocking buffer. TBS-T was used for all washes. Chemiluminescence with

Immobilon Forte Western HRP substrate (Millipore Sigma, Cat# WBLUF0500) and

near-Infrared fluorescence imaging were performed with FluorChem R (ProteinSimple)

and Odyssey® Imaging System (Li-COR Biosciences), respectively. The following

antibodies were used for Western blots with the indicated dilution factors: Hexokinase I

(C35C4) Rabbit mAb (Cell Signaling Technology, Cat# 2024, RRID:AB_2116996) at

1:1,000, HK2 Monoclonal Antibody (3D3) (Invitrogen™, Cat# MA5-15679,

RRID:AB_10986812) at 1:1,000, PARP Antibody (Cell Signaling Technology, Cat#

9542, RRID:AB_2160739) at 1:2,000, Acetyl-CoA Carboxylase (C83B10) Rabbit mAb

(Cell Signaling Technology, Cat# 3676, RRID:AB_2219397) at 1:1,000, Phospho-

Acetyl-CoA Carboxylase (Ser79) (D7D11) Rabbit mAb (Cell Signaling Technology, Cat#

11818, RRID:AB_2687505) at 1:500, Rabbit Anti-GAPDH Monoclonal Antibody

Unconjugated, Clone 14C10 (Cell Signaling Technology, Cat# 2118, RRID:AB_561053)

at 1:5000, S6 Ribosomal Protein (54D2) Mouse mAb (Cell Signaling Technology, Cat#

2317, RRID:AB_2238583) at 1:2000, Phospho-S6 Ribosomal Protein (Ser235/236)

(D57.2.2E) XP Rabbit mAb (Cell Signaling Technology, Cat# 4858, RRID:AB_916156)

at 1:2000, Anti-β-Actin antibody-Mouse monoclonal (Sigma-Aldrich, Cat# A1978,

RRID:AB_476692) at 1:5,000. Secondary antibodies used for chemiluminescence

detection were HRP conjugated anti-mouse antibody (Cytiva, Cat# NA931) and anti-

rabbit antibody (Vector Laboratories, Cat# PI10001). Near-infrared detection was

carried out with IRDye® 680RD Donkey anti-Rabbit IgG (H + L) (LI-COR Biosciences,

Cat# P/N 925-68073) and RDye® 800CW Donkey anti-Mouse IgG (H + L) (LI-COR

Biosciences, Cat# P/N 925-32212). Western blot Images were processed with GIMP

(GNU Image Manipulation Program, Version 2.10).

In vivo BKIDC-1553 studies

Male severe combined immunodeficient (SCID) mice were used for all in vivo

experiments (Charles River Laboratories, strain code# 236). PC3 (100,000 cells) were

mixed 1:1 with Matrigel (Corning, Cat# 356234) and injected subcutaneously into rear

flanks of intact SCID mice (10 mice were enrolled per treatment group for PC3 study).

When tumors reached ~100 mm^3, either vehicle (7% Tween-80, 3% EtOH in

pharmaceutical grade PBS) or 20 mg/kg of BKIDC-1553 was given by oral gavage three

times weekly (Mon, Wed, Fri), for up to 4 weeks. A patient-derived PCa xenograft,

LuCaP 35 (RRID:CVCL_4853), was implanted subcutaneously into rear flanks of intact

SCID mice (The number of enrolled mice was: 8 for control, 9 for 1553, 7 for

enzalutamide). When tumors reached ~100 mm^3, mice were given vehicle (7% Tween-

80, 3% EtOH in pharmaceutical grade PBS), 20 mg/kg of BKIDC-1553, or 10 mg/kg

enzalutamide (Medivation, log# 083112-02, lot: BREC-0035-123T) by oral gavage three

times weekly (Mon, Wed, Fri) for up to 5 weeks. All mice were assigned to treatment or

control groups using a random number generator. ARRIVE guidelines were followed for

all animal experiments. All animal studies were approved by the University of

Washington Institutional Animal Care and Use Committee (protocol 2797-04).

Determining BKIDC-1553 concentrations in tumors

Each tumor was flushed with Dulbecco's phosphate-buffered saline (DPBS), weighed

and placed in storage at -80°C. On the day of homogenization, 0.9% NaCl was added to

each sample to achieve 100 mg/mL of tissue concentration. Samples were then

homogenized using a handheld motorized homogenizer (Thermo Fisher Scientific) on

ice, and 100 µL of each homogenate sample was placed in a 96-well plate. In addition,

tissues from vehicle control-treated mice were homogenized and placed in a 96-well

plate to be used as blanks or standards. Calibration curves were generated individually

for each type of tumor. An internal standard was added to each sample to a final

concentration of 20 nM. All homogenates were stored at -80°C. On the day of tissue

extraction, 10 µL of each sample were added to a 96-well plate that contained 90 µL of

80% ACN:20% water. The samples were mixed and centrifuged at 4,200 g for 20 min.

25 µL of the supernatants were placed in a 96-well plate that contained 475 µL of 80:20

ACN:water. Then, the samples were mixed and centrifuged at 4,200 g for 10 min.

Supernatant from each sample was placed in a new 96-well plate, sealed and analyzed

with an Acquity UPLC in tandem with a Xevo TQ-S micro mass spectrometer (Waters

Corp). Sample concentrations were determined using the internal standard normalized

calibration curves. To report tissue concentrations of BKIDC-1553, diluted homogenates

were adjusted post-integration on a weight:weight basis. Individual tumor

concentrations, means, and standard deviations are reported in **Supplementary Table**

5.

Pharmacology (PK), Absorption, Distribution, Metabolism, and Excretion (ADME),

and Toxicology Methods:

PK, ADME and Toxicology methods with BKIDC-1553 were carried out as described

previously where the compound was referred to as Compound **32** (35) and reanalysis of

PK data was carried out with Phoenix WinNonlin software (Certara, NJ). A human dose

prediction for BKIDC-1553 was generated using estimated human pharmacokinetic

parameters and BKIDC-1553 concentrations associated with efficacy in a preclinical

prostate cancer mouse model. BKIDC-1553 clearance (Cl) and volume of distribution

(V) in a 74 kg human were estimated by allometric scaling using pharmacokinetic

parameters observed with IV dosing in rats, dogs, and non-human primates (Table

S7)(36-38). Since the exponent for Cl using body weight was >1, the product of Cl x

brain weight was used(39). Based on the rapid and nearly complete absorption

observed in preclinical models, BKIDC-1553 was assumed to have an oral

bioavailability of 1.0 in humans. The BKIDC-1553 concentration associated with efficacy

in vivo was determined using a population pharmacokinetic modeling approach. Briefly,

observed BKIDC-1553 plasma concentrations in mice were fit with a 1-compartment

structural model and a proportional error model. BKIDC-1553 concentrations were

simulated based on post-hoc estimates of pharmacokinetic parameters (absorption rate

constant (ka), Cl, V) for each mouse (n = 3). The simulated concentrations were used to

estimate the maximum BKIDC-1553 plasma concentration (C_{max}) and area under the

curve (AUC) over the last day of dosing ($AUC_{264\text{-}288h}$) and the entire dosing period

($AUC_{0\text{-}288h}$).

PK experiments were performed as previously described(35) with the following

modifications. Male SCID mice were given an oral dose of 20 mg/kg BKIDC-1553 (in

7% Tween-80, 3% EtOH in pharmaceutical grade PBS vehicle) on Monday.

Wednesday, and Friday for two weeks for a total of 6 doses. Blood was sampled after

the 1st, 4th, and 6th doses, with a terminal bleed taken on the third Monday, 72 h after the

final dose. Blood samples at each dose were taken pre-dose and at 1 h, 2 h, 4 h, 6.5 h,

and 24 h post dose. Plasma protein determination was performed using the dialysis

membrane technique as previously described(40). Caco-2 permeability testing was

performed in Hanks balanced salt solution containing 20 mM HEPES using 0.3 cm^2

polycarbonate filter transwells seeded with 6 x 10^4 cells. On day 24 post-seeding, donor

solutions containing test compounds in DMSO (0.1% final) were added to confluent cell

monolayers and compound flux was assessed from the acceptor chamber at 5-6 time

points over 120 min. Acceptor chamber volume was replaced with blank buffer and

dilution corrected with each sample collection. Samples were stored frozen at -80°C

until analysis by LC-MS. Apparent permeability coefficient (P_{app}) calculations were

determined using the equation:

$$P_{app} = \frac{dQ}{dt} \times \frac{1}{C_{initial\,donor}} \times A$$

Where dQ/dt = apparent steady-state transport rate (µmol/s); A = monolayer surface

area; and $C_{initial\,donor}$ = concentration in donor at experiment initiation. Apparent efflux

ratios were calculated based on the P_{app} values in the A-B and B-A directions.

Additional permeability testing in MDR1-MDCKII cells (RRID:CVCL_IZ19) was

conducted by Eurofins CRO services (Eurofins Scientific, Luxembourg) using 10 µM

BKIDC-1553.   Hepatic microsome metabolism of BKIDC-1553 was performed with

cryopreserved liver microsomes from dogs, humans, and rats. Microsome incubations

were prepared by pre-incubating 1.0 mg/mL cryopreserved liver microsomes in

phosphate buffer (pH 7.4, 100 mM) for 5 min at 37□C. The reaction was started with the

addition of UDPGA (5 mM), NADPH (2 mM) and incubation was performed for 1 h. The

final concentration of test article was 10 µM with <0.5% organic. The sample was

quenched with an equal volume of acetonitrile, vortexed, centrifuged at 13,000 rpm for 5

min, the supernatant (170 µL) was transferred to 96 well plate and partially dried under

nitrogen to ~ 120 µL for ultra-high-performance liquid chromatography-high resolution

mass spectrometry (UPLC-HRMS, with UPLC instrument: Waters ACQUITY I-Class

system, and Mass Spectrometer instrument: Waters Xevo G2 XS) analysis. The positive

control was a separate incubation of verapamil using the same microsomes to confirm

the viability of the incubation. The negative controls were separate incubation of the test

article using the incubation media and same microsome incubation without NADPH and

UDPGA. The LC/MS full scan data was processed manually, assisted by Metabolynx, to

identify the metabolites. Product ion spectra were acquired by a separate UPLC-

HRMS/MS run to assign the structures of the metabolites by MS/MS data interpretation.

In vitro inhibition of Cytochrome P450 (CYP) enzymes was performed in five major

isoforms (CYPs 1A2, 2C9, 2C19, 2D6, and 3A4) in human liver microsomes using up to

20 µM BKIDC-1553 and included standard CYP specific inhibitors (furafylline,

sulfaphenazole, ticlopidine, quinidine, and ketoconazole) and substrates (phenacetin,

tolbutamide, *(S)*-mephenytoin, dextromethorphan, and midazolam/testosterone).

Various concentration of BKIDC-1553 and positive control inhibitors were co-incubated

at 37°C with each substrate in human liver microsomes (XenoTech LLC, Lenexa KS) for

10-40 min, with reactions initiated by the addition of an NADPH-regenerating system

and quenched by addition of ice-cold acetonitrile containing 0.15 µg/mL of diazepam

internal standard. Quenched samples were analyzed by LC-MS (Waters/Micromass

Xevo TQD triple-quadrupole) and concentrations determined relative to standards in

quenched microsomal matrix. Inhibitory activities were based on the reduction in

formation of the specific CYP-mediated metabolite relative to no-inhibitor controls and

reported as 50% inhibition values (IC_{50}). Additional CYP inhibition studies were

conducted by Eurofins CRO services on CYPs 2B6 and 2C8 with 10 µM BKIDC-1553

using Clopidogrel and Montelukast as inhibitors and bupropion and amodiaquine as

substrates for 2B6 and 2C8, respectively. The mutagenic toxicity of BKIDC-1553 was

assessed by measuring its ability to induce reverse mutations in Salmonella TA98

(frame shift) and TA100 (base-pair) strains in the presence and absence of a

metabolizing Aroclor 1254-induced rat S9 mix with the top concentration of 180 µg/well,

in the modified 24-well assay, which is equivalent to 5000 µg/plate in the standard plate

assay. A positive response is defined with a 2-fold increase in revertant colonies[41].

The genotoxicity of BKIDC-1553 was assessed in an in vitro micronucleus test carried

out with Chinese hamster V79 cells and determined by measuring the extent of

micronucleus formation with and without metabolizing Aroclor 1254-induced rat S9

mixture. BKIDC-1553 was tested at a top concentration of 500 µg/mL. The safety 44

screen assay was performed by Eurofins as described

(https://cdnmedia.eurofins.com/corporate-

eurofins/media/1069358/safetyscreen44_epdsfl420june16.pdf)(42).

Safety testing of BKIDC-1553 in mice, using oral gavage of BKIDC-1553 suspended in

a lipid-based vehicle (60% Phosal 53 MCT: 30% PEG-400: 10% ethanol), was

performed with 35 mg/kg, 70 mg/kg, and 100 mg/kg BKIDC-1553 once daily oral doses

for 7 days. Mice were monitored daily for signs of toxicity and glucose levels were

sampled from tail snip using an Accu-Chek Guide Me glucometer (Roche) at pre-dose,

2, and 4 h post dose time points. Blood samples were collected after the 3rd dose and

processed for LC-MS/MS (Waters Acquity Xevo TQ-S micro) analysis of BKIDC-1553

levels in plasma.

Non-GLP toxicology studies were carried out in dogs by StillMeadow Inc., (Sugarland,

Texas). In the first experiment, four groups of one male and one female >9-month-old

beagle dogs each were dosed orally by gavage tubes with either the vehicle or BKIDC-

1553 daily for 14 days. Group I was a Vehicle Control (80% Phosal 53 MCT, 15% PEG

400 and 5% Ethanol) at 1 mL/kg and Groups II, III and IV received 1.6 mg/kg, 5 mg/kg

and 15 mg/kg of BKIDC-1553 in the same vehicle at 1 mL/kg, respectively. Dogs were

monitored during the entire time of the experiment for changes in body weight and food

consumption and for acute signs of toxicity and general health. Heparinized plasma for

BKIDC-1553 quantitation was collected pre-dose, and 2, 4, 6, 24, and 48 h after the last

dose, and measured by LC-MS/MS. Dogs were monitored for blood glucose levels

(AlphaTRAK glucometer, Zoetis) prior to and 2 h after the first dose of the day, each day

of dosing. Blood and urine samples were also collected for serum chemistry,

hematology, and coagulation and urinalysis pre-dose and 24 h after the last dose.

These samples were analyzed by Antech Diagnostics. In the second non-GLP dog

toxicology experiment, a male and female >9-month-old beagle dog was first treated for

5 days with oral gavage twice a day (6 h apart) of the 4 mL/kg lipidic vehicle (80%

Phosal 53 MCT, 15% PEG 400 and 5% Ethanol), blood and urine were collected at the

24 hours after dosing, tested as above. After a 4-week washout period, the same dogs

were treated by oral gavage with BKIDC-1553 at 7.5 mg/mL in 4 mL/kg lipidic vehicle at

6 h intervals (30 mg/kg twice a day) for two days. Vomiting, decrease activity, and

salivation was noted in both animals on day two of BKIDC-1553 administrations, and

the next day, day 3, the BKIDC-1553 (7.5 mg/mL)-vehicle volume was reduced to 2

mL/kg administered twice daily at 6 h intervals (15 mg/kg twice a day) for an additional 3

days. Blood glucose was monitored by glucometer measurements before and 2 h after

administration of each dose. Plasma for PK measurements were collected as noted

above, after the last dose on the fifth day and the plasma BKIDC-1553 levels

determined via LC-MS/MS.

Rat cardiovascular (CV) safety studies were carried out at AbbVie Inc as previously

described(43).

Synthesis of PP and PrP based BKIDCs

1-(4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylpropan-2-ol (0.6 g, 1 eq)

was heated to 100 °C for 2 h in a microwave with a solution of 33% methylamine in

ethanol (5.7 mL, keeping the concentration of the pyrrolopyrimidine scaffold around 0.3

M in solution). The crude mixture was concentrated under reduced pressure and

purified by flash chromatography using 5% MeOH/DCM to obtain 1-(5-iodo-4-

(methylamino)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylpropan-2-ol with a 64% yield. [1]H

NMR (500 MHz, CDCl$_3$) δ 8.18 (s, 1H), 7.30 (s, 1H), 4.12 (s, 2H), 3.11 (s, 3H), 1.14 (s,

6H). Calcd for C$_{11}$H$_{15}$IN$_4$O 346.03, found (M+H)$^+$ 347.1.

A mixture of 4-chloro-3-iodo-1H-pyrazolo[3,4-d]pyrimidine (125 mg, 0.354 mmol) and

33% methyl amine in ethanol (1.2 mL) was heated to 100 °C for 2 h in a microwave

reactor. The crude mixture was concentrated under reduced pressure. Crude A was

obtained in 62% yield and was used in the next reaction without further purification. [1]H

NMR (500 MHz, MeOD) δ 8.16 (s, 1H), 5.39 (s, 1H), 3.04 (d, J = 9.8 Hz, 3H).

Compound A (60 mg, 0.22 mmol) was heated to 150 °C for 3 h with 2,2-dimethyloxirane

(40 uL, 2.0 eq) and K$_2$CO$_3$ (61 mg, 2.0 eq) in a mixture of 8.5:1.5 ACN/water [A=0.64M].

Crude mixture was concentrated under reduced pressure and purified by flash

chromatography using 5% MeOH/DCM to obtain compound B with a 57% yield.　[1]H

NMR (500 MHz, MeOD) δ 8.28 (s, 1H), 4.32 (s, 2H), 4.30 (s, 1H), 3.16 (s, 3H), 1.23 (s,

6H). Calcd for $C_{10}H_{14}IN_5O$ 347.02, found $(M+H)^+$ 348.3.

2-(6-cyclopropoxynaphthalen-2-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane was

synthesized according to previously published protocols(35).

676
677

General Suzuki coupling condition to obtain the final compounds: The appropriate

3-iodo pyrazolopyrimidine or 5-iodo pyrrolopyrimidine scaffold (1.0 eq), the desired

boronic acid or the boronate pinacole ester (1.5 eq), $PdCl_2.dppf.DCM$ (0.05 eq), K_3PO_4

(2.2 eq) were heated to 85 °C for 1 h in a microwave using a 3:1 mixture of

dioxane/water as the solvent. The crude mixture was concentrated and purified by

HPLC using ACN/water as the mobile phase to obtain the final compounds.

1553 (aka BKIDC-1553)

686   a

1-(4-amino-3-(6-cyclopropoxynaphthalen-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2-

methylpropan-2-ol was synthesized and characterized as previously described (35)

(identified as Compound **32**).

1553-N-Me

1-(3-(6-cyclopropoxynaphthalen-2-yl)-4-(methylamino)-1H-pyrazolo[3,4-d]pyrimidin-1-

yl)-2-methylpropan-2-ol: ^{1}H NMR (500 MHz, CDCl$_3$) δ 8.47 (s, 1H), 8.05 (s, 1H), 7.94 (d,

$J = 8.4$ Hz, 1H), 7.84 (d, $J = 9.1$ Hz, 1H), 7.73 (d, $J = 8.4$ Hz, 1H), 7.53 (s, 1H), 7.25 (m,

1H), 5.52 (q, $J = 4.8$ Hz, 1H), , 4.48 (s, 2H), 3.91 (m, 1H), 3.09 (d, $J = 4.8$ Hz, 3H), 1.26

(s, 6H), 0.90-0.88 (m, 4H). Calcd for C$_{23}$H$_{25}$N$_5$O$_2$ 403.2, found (M+H)$^+$ 404.4. HPLC

 purified product was determined to be ≥95 % pure under 220 nm and 254 nm detection

in analytical HPLC under both solvent systems.

1817

Conc HCl (3 mL) was added to 1-(6-ethoxynaphthalen-2-yl)-3-((1-methylpiperidin-4-

yl)methyl)imidazo[1,5-a]pyrazin-8-amine(35)(50 mg) and heated overnight at 85 C.

Crude mixture was diluted with aq. NaHCO$_3$ and the product was extracted using ethyl

acetate. The organic layer was dried over Na$_2$SO$_4$, concentrated, and purified by HPLC

using ACN/water as the mobile phase to obtain 1817. [1]HNMR(300 MHz, MeOD) δ 8.39

(s, 1H), 8.11 – 7.98 (m, 2H), 7.89 (dd, J = 8.5, 1.6 Hz, 1H), 7.55 (d, J = 8.5 Hz, 1H),

6.71 (d, J = 9.5 Hz, 1H), 4.46 (d, J = 6.7 Hz, 2H), 3.56-3.52 (m, 2H), 3.09 – 2.93 (m,

2H), 2.85 (s, 3H), 2.42-2.38 (s, 1H), 1.99-1.95 (m, 2H), 1.70-1.63 (m, 2H). Calcd for

C$_{21}$H$_{23}$N$_7$O 389.20, found (M+H)$^+$ 309.1. HPLC purified product was determined to be

≥99 % pure under 220 nm and 254 nm detection in analytical HPLC under both solvent

systems.

1826

Compound D (0.064 mmol, 1 eq) and sodium iodide (0.13 mmol, 2 eq) were dissolved

in DMF (0.3 mL). The mixture was cooled to 0 °C and NaH (0.13 mmol, 2 eq) was

added followed by compound C (0.096 mmol, 1.5 eq). The reaction mixture was allowed

to come to 70 °C and allowed to stir over night. The mixture was diluted with water and

the crude product was extracted into ethyl acetate. The organic layer was dried over

sodium sulfate and concentrated to obtain compound E (Calcd for $C_{40}H_{56}N_6O_9$ 764.41,

found $(M+Na)^+$ 787.6). Crude E was then stirred with a 30% TFA/DCM mixture (1.6 mL)

for 3 h at room temperature. The crude mixture was concentrated by rotary evaporation

and purified by flash chromatography using a 5% methanol/DCM supplemented with 2%

triethylamine as the mobile phase to obtain ~90% pure 1826 with a 32% yield. [1]H-

NMR(300 MHz, MeOD) δ 8.36 (s, 1H), 8.16 (s, 1H), 8.01 (d, J = 8.6 Hz, 1H), 7.94 (d, J

= 9.0 Hz, 1H), 7.81 (dd, J = 8.5, 1.6 Hz, 1H), 7.65 (d, J = 2.3 Hz, 1H), 7.27 (dd, J = 8.9,

2.4 Hz, 1H), 4.44 (s, 2H), 4.03 – 3.97 (m, 1H), 3.78 – 3.67 (m, 9H), 3.60 – 3.57 (m, 2H),

3.51 – 3.48 (m, 3H), 3.43-3.39 (m, 2H), 3.27-3.24 (m, 1H), 3.15-3.10 (m, 1H), 3.05 –

2.99 (m, 2H), 1.30 (s, 6H), 0.94 - 0.90 (m, 2H), 0.86 – 0.74 (m, 2H). Calcd for

$C_{30}H_{40}N_6O_5$ 564.3, found $(M+H)^+$ 565.6.

Immunohistochemistry (IHC)

Xenograft tumors were fixed with formalin and then paraffin embedded. The resulting

FFPE (formalin-fixed, paraffin-embedded) samples were sectioned at 10 μm, mounted

on slides, deparaffinized, rehydrated, heated up in Tris-based buffer (Vector

Laboratories, Cat# H-3301-250) in the pressure cooker to accomplish antigen

unmasking. Endogenous peroxidases were inactivated with 3% (v/v) H_2O_2 (Fisher

Scientific, Cat# H325) in Tris-buffered saline. The sections were then blocked in 2%

(v/v) normal goat serum (Invitrogen™, Cat# 10000C) at room temperature for 1 h before

application of primary antibody incubation at 1:500 dilution (Phospho-Acetyl-CoA

Carboxylase (Ser79) (D7D11) Rabbit mAb (Cell Signaling Technology, Cat# 11818,

RRID:AB_2687505)) overnight at 4°C. The 3,3′-diaminobenzidine (DAB) based-

chromogenic detection was performed using biotinylated secondary antibodies in

conjunction with VECTASTAIN® ELITE® ABC HRP Detection Kit (Vector Laboratories,

Cat# PK-6100) and ImmPACT® DAB Substrate Kit, Peroxidase (HRP) (Vector

Laboratories, Cat# SK-4105). DAB-stained slides were counterstained with hematoxylin,

dehydrated, and mounted with a coverslip. Appropriate negative controls were run in

parallel. For DAB-stained slides, at least three digital images per section were obtained

at 20× magnification with EVOS™ M7000 Imaging system (ThermoFisher Scientific)

with the same magnification and exposure. The images were opened in ImageJ

(RRID:SCR_003070, NIH image analysis freeware, http://imagej.nih.gov/ij/) and

converted to RGB stack. The threshold of the green channel was optimized until the

DAB-stained areas showed optimal contrast in the red channel. The same optimized

threshold was used for all the images and the intensity of DAB staining was expressed

as a percentage of stained area per a standard 20x field.

Kinobead profiling of 1553-N-Me using a 1:1 mixture of HCT116/HEK293T lysate

Competitive kinobead profiling of 1553-N-Me: Kinobeads were synthesized by

immobilizing five pan-kinase inhibitors on NHS activated sepharose (compounds 1

(33.3%), 2 (33.3%), 3 (8.3%), 4 (8.3%) and 7 916.7% from Golkowski et al (2017))(25),

using a slightly modified protocol(25). The synthesized beads were stored as a 50%

slurry in 20% ethanol until further use. A mixture (1:1) of HCT116 and HEK293T lysate

was generated with a final concentration of 2 mg/mL in mod. RIPA buffer (pH 7.8, 50

mM Tris, 150 mM NaCl, 10 mM NaF, 1% NP40, 0.25% sodium deoxycholate, 5%

glycerol, supplemented with Pierce Protease Inhibitor Tablets, 1 mM PMSF and

phosphatase inhibitor cocktail 2 and 3 from Sigma Aldrich). 1553-N-Me (20 µM) or

DMSO were added to the lysate mixture, followed by incubation for 30 min at 4 ºC. 150

µL of DMSO- or 1553-N-Me-treated lysate was then added to 5 µL of pre-washed

kinobeads. Samples were then rotated for 3 h at 4ºC. Following kinobead enrichment,

beads were washed with 500 µL of cold mod. RIPA lysis buffer (2x) followed by 500 µL

of chilled TBS (50 mM Tris, 150 mM NaCl, pH = 7.8) (3x) in order to remove detergent.

Each competitive profiling experiment was performed in quadruplicate. Following

washing, kinobeads were resuspended in 50 µL of denaturing buffer (6M guanidinium

chloride, 50 mM Tris containing 5 mM TCEP and 10 mM CAM, pH = 8.5) and heated to

95°C for 5 min. The bead slurry was the diluted with 50 µL of 100 mM TEAB

(triethylammonium bicarbonate buffer, pH 8.5). 0.8 µg of LysC (Wako) was then added

to the beads and the overall pH was adjusted to 8-9 with 1 N NaOH. A thermomixer

(Eppendorf) was then used to agitate the mixture at 37°C for 2 h at 1400 rpm. Further

dilution was performed by adding 100 µL of 100 mM TEAB, followed by the addition of

0.8 µg of sequencing grade trypsin (Pierce). Proteolytic digestion was performed for 14

h at 37°C at 800 rpm in the thermomixer. After overnight digestion, 200 µL of Buffer A

(5% ACN, 0.1% TFA) containing 1% formic acid was used to digest the samples and

the pH was adjusted to 2-3 with formic acid. Homemade stage-tips were prepared by

running 50 µL of Buffer B (80% ACN, 0.1% TFA, H_2O) through them followed by 50 µL

of Buffer A (5% ACN, 0.1% TFA, H_2O). The supernatants of spun down and kinobeads

were added to the stage-tips. Stage-tip loading was performed by letting the proteolytic

digest solutions pass through the stage-tips. Loaded stage-tips were then washed with

50 µL of Buffer A and eluted with 50 µL of Buffer B. Samples were dried with a

speedvac and the resuspended in Buffer A. Resuspended samples were then injected

onto an LC-MS. Samples were separated using a nanoAcquity UPLC instrument and a

homemade 10 cm capillary column packed with 5 µm 120 Å reverse-phase C18 beads

(ReproSil-Pur®, Maisch). Liquid chromatography was performed over 120 min using an

initial 20 min isocratic trapping of 3% Buffer B and a flow rate of 700 nL/min, followed by

a 100 min gradient of 35% Buffer B to an 80% Buffer B gradient at 350 nL/min. LC

Buffer A solvent was 0.1% Acetic acid and LC Buffer B was 99.9% ACN, 0.1% acetic

Acid. Data were collected using a Thermo Scientific Orbitrap Fusion Tribrid mass

spectrometer. Raw files were analyzed using MaxQuant, followed by Perseus by

filtering intensity values only identified by site, reverse, or potential contaminant. To

determine kinases that were significantly competed by treatment with 20 µM of 1553-N-

Me, a two-tailed two-sample t-test in Perseus with an FDR of 0.05 was applied. Kinases

were considered as being non-competed by an inhibitor if it had a Log2 ratio (Log2

Difference) <1.

Competitive profiling of 155-N-Me with immobilized 1826

1826 beads were synthesized using a previously published protocol with slight

modifications(25). In brief, 26.7 µL of a 20 mM DMSO stock of 1826 and 6 µL of triethyl

amine were added to 0.8 mL of 50% slurry of pre-washed NHS-sepharose beads. The

mixture was rotated at room temperature for 20 h in the dark. At the end of this time,

supernatant was removed, and the beads were washed thrice with 1:1 DMF/ethanol

solution. Then the beads were incubated with a mixture of 1:1 DMF/ethanol (528 µL),

1M aminoethanol dissolved in 20 mM acetic acid in 1:1 DMF/ethanol (272 µL) and 1 M

EDCI (120 µL) for 12 h in dark at room temperature for capping. The beads were then

washed 3x times with 1:1 DMF/ethanol, 2x with 0.5 M NaCl and finally with a 20%

ethanol solution. The beads were stored at 4°C until further use as a 50% slurry in 20%

ethanol. 1553-N-Me was profiled using 5 µL of the 1826 beads described above and a

1:1 mixture of HCT116/HEK293T lysate using the same protocol as described for

kinobead profiling of 1553-N-Me.

Measurement of in vitro AMPK activity

Measurement with recombinant proteins was assessed using recombinant human

AMPK (α1β1γ1 complex). Protein kinase activity of recombinant full-length AMPK

(A1/B1/G1) enzyme (SignalChem, Richmond, BC Canada) was determined using a

non-radioactive assay which measure changes in initial concentration of ATP via

luminescence. Enzyme kinase activity was measure in the presence of 45 µM peptide

substrate SAMStide (HMRSAMSGLHLVKRR) (SignalChem) in a buffered solution

containing 5 mM MOP (pH 7.5), 5 mM MgCl$_2$, 1 mM EGTA, 2.5 mM β-glycerol-

phosphate, 0.5 mM EDTA and 0.05 mM DTT. A series of AMPK (A1/B1/G1)

concentrations i.e., 40, 25, 15, and 10 ng were used to detect enzyme activity

enhancement. The reaction was initiated with addition of 1 µM ATP. Incubation of

enzyme reaction was at 37°C and 90 rpm agitation for 90 min. The assay was

terminated with the addition of Kinase-Glo$^®$ Luminescent Kinase Assays reagent

(Promega, Cat# V6711) and read in an EnVision Multilabel Plate Reader (Perkin Elmer,

USA). Compound A-769662 (Cayman Chemical, Cat# 11900) is a known positive

control that stimulates AMPK activity, Compound C (Cayman Chemical, Cat# 11967),

also known as dorsomorphin, is a known inhibitor of AMPK activity (both of these

compounds were used at 2 µM final concentrations), and DMSO was used as a no

compound control.

Activity measurement of hexokinase

The full-length human HK2 open reading frame (ORF) flanked by the linkers was

amplified by PCR using plasmid FLHKII-pGFPN3 (RRID:Addgene_21920) as a

template with primers

(*GGGTCCTGGTTCG*ATGATTGCCTCGCATCTGCTTGCCTACTTC and

*CTTGTTCGTGCTGTTTA*CTATCGCTGTCCAGCCTCACGGATGCGGC: Italic,

underlined, and double underlined text correspond to linker, initiation codon, and stop

codon, respectively.) The resulting PCR fragment was cloned into bacterial expression

plasmid AVA0421 by ligation-independent cloning(44, 45). The sequence integrity of the

cloned fragment (GenBank accession No.: NM_000189.5) was confirmed by Sanger

sequencing at GENEWIZ (South Plainfield, NJ). Human HK2 protein was expressed in

Escherichia coli BL21(DE3) using Studier auto-induction method at 20°C and 180

rpm(78), followed by cell lysis and purification using immobilized metal-affinity

chromatography (IMAC) in a Ni–NTA (Qiagen, Valencia, CA) column in binding buffer

composed of 20 mM HEPES pH 7.25, 500 mM NaCl, 5% glycerol, 30 mM imidazole,

0.5% CHAPS and 1 mM tris(2-carboxyethyl)phosphine (TCEP) as previously

described(79). This was followed by elution in the same buffer supplemented with 250

mM imidazole and size-exclusion chromatography in a 26/60 Superdex 75 size

exclusion column. SDS-PAGE analysis revealed that the purified HK2 enzyme was near

homogeneity.

For **Figure S13A** and **B**, hexokinase activity was measured using Hexokinase

Colorimetric Assay Kit (Sigma-Aldrich, Cat# MAK091) on 96-well plates according to the

manufacturer's instructions. 50 µL of sample containing either LNCaP crude extract or

recombinant HK2 protein (R&D Systems™, Cat# 8179HK020) was added to a well of

96-well plates. The reaction was initiated by addition of 50 µL reaction mixture

containing HK Assay Buffer, HK Enzyme mix, HK Developer, HK Coenzyme, HK

Substrate, and test compounds (BKIDC-1553-N-Me or 3-bromopyruvate (Sigma-Aldrich,

Cat# 16490) using DMSO as a vehicle control). Hexokinase activity was determined by

measuring a colorimetric product with absorbance at 450 nm proportional to the

enzymatic activity present. For **Figure S13C**, in-house recombinant human HK2 protein

was used to evaluate the effect of BKIDC on glucose phosphorylation activity. The non-

radioactive Kinase-Glo® Luminescent Kinase Assays (Promega, Cat # V6711) was

used to measure ATP consumption in the presence of its natural substrate glucose.

Kinase reactions were performed in 25 mM HEPES-KOH, pH 7.5 containing 150 mM

NaCl, 10 mM $MgCl_2$, and 10 mM $CaCl_2$. The reaction mixture contained 1 mM of D-

glucose substrate, 10 µM of BKIDC or 100 µM 3BP. The activity of HK2 was measured

in a total reaction volume of 25 µL using a serial dilution of the enzyme in the range

between 70 and 840 nM. All reactions were initiated by the addition of 10 µM ATP and

terminated after 120 min incubation at 37°C and 90 rpm shaking with addition of Kinase-

Glo® reagent. Change in initial ATP concentration was measured as bioluminescence

readout in counts per second on the Envision plate reader (PerkinElmer).

Statistical Analysis

Statistical analyses were conducted using GraphPad Prism (RRID:SCR_002798) v8

and are indicated within all figures. One-way ANOVA followed by Tukey's or Dunnett's

test for multiple comparison or 2-tailed unpaired t test was performed to determine

significance, which was defined as $p < 0.05$.

RESULTS

BKIDC effects in Prostate Cancer cell lines

Our lead compound, BKIDC-1553 has oral availability with a favorable safety and

pharmacokinetics profile (compound **32** in (35)). BKIDC-1553 was tested in the MTS (3-

(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay for proliferation

inhibition in a series of human PCa cell lines (LNCaP, LNCaP95, C4-2, C4-2B, LAPC4,

VCaP, PC3, DU145, LNCaP^{APIPC}(28), MSKCC EF1 and NCI-H660) and non-prostate

control lines (HepG2, U2OS, WIL2-NS and HFF) (**Figure 1A**). From the data we see

that androgen receptor (AR) positive lines LNCaP, LNCaP95, C4-2, C4-2B, LAPC4 and

VCaP as well as AR negative cell lines LNCaP^{APIPC} (LNCaP cell engineered to grow

without AR(28)) and the neuroendocrine (NE) lines MSKCC EF1 and NCI-H660

respond to BKIDC-1553. Thus, the responsiveness to BKIDC-1553 was found in both

AR positive and AR negative PCa cell lines. The concentration to inhibit cell proliferation

by 50% ranged from 2.5-20 µM in these cell lines; concentrations achievable in

experimental animals without toxicity. No response to BKIDC-1553 is seen in the AR-

negative and non-NE PCa lines (PC3 and DU145). We also saw no response to BKIDC-

1553 from non-PCa cell lines including a human bone cancer line (U2OS), human

foreskin fibroblasts (HFF), a lymphoblastoid (WIL2-NS) or hepatoma (HepG2) cell lines

(**Figure 1A**). We performed crystal violet growth assays three times with LNCaP95 and

PC3 cells in dose response to BKIDC-1553 (**Supplementary Figure S3**). They clearly

demonstrate inhibition of cell growth in the BKIDC sensitive cell line, LNCaP95 cells, but

not the BKIDC resistant line, PC3. These experiments show that the effects seen with

the MTS assay are due to cell proliferation and not due to metabolic responses. In

summary, BKIDC-1553 was active in suppressing the growth of most AR positive and

AR negative metastatic PCa cell lines but was not effective in suppressing the growth of

normal fibroblast cells or bone, lymphocyte, or hepatocyte derived cancer cell lines that

are tested in this paper.

We next evaluated if PKD was the target of BKIDCs. We first noted that unlike

previously reported PKD inhibitors(26, 27), BKIDC-1553 lacked activity against PC3, an

AR negative and NE negative cell line. To gain insight into the targets of BKIDC-1553,

we performed structure–activity relationship studies with small molecules based on

pyrrolopyrimidine (PrP) or pyrazolopyrimidine (PP) scaffolds (**Figure 1B**) with LNCaP95

and PC3 as a representative of responders and non-responders, respectively (**Table S1**

and S2). A series of compounds with variable R1 and R2 substituents still displayed

antiproliferative activity in LNCaP95 but not PC3 (**Figure 1C and Table S1 and S2**). We

created BKIDCs, represented by BKIDC-1553-N-Me that contain substituents (alkyl

groups on the exocyclic nitrogen of pyrimidine) that disrupt canonical hinge region

hydrogen bonding interactions in the ATP-binding site of protein kinases (**Figure 1B**).

The protein targets of BKIDC-1553-N-Me were determined with competitive proteomic

profiling experiments and no protein kinases, including PKD, MEK2, and RIPK2, were

identified as hits (**Figure S1**). Yet, BKIDC-1553-N-Me maintained its antiproliferative

activity against LNCaP95 (**Figure S2**). This demonstrated PKD. MEK2, and RIPK2 were

not BKIDC targets and suggested the antiproliferative actions of BKIDCs on PCa cells

acts via a mechanism not requiring protein kinase inhibition. Indeed, we confirmed that

BKIDC-1553-N-Me showed a similar antiproliferative spectrum across the cell lines

tested with BKIDC-1553 (**Figure S2**). In this screen, we also found a closely-related

molecule, BKIDC-1817 (**Figure 1B**: a 1H-pyrazolo[3,4-d]pyrimidin-4-amine

compound)(46), had no inhibitory activity in LNCaP95 or PC3 cells (**Figure S2**). BKIDC-

1817 served as a negative control in subsequent studies.

BKIDC effects on cell cycle and cell-signaling pathways

We carried out a series of experiments to characterize cell cycle effects, cell-

signaling pathways, and energy sources to investigate the cellular processes inhibited

by BKIDC-1553 and -1553-N-Me. Thirty-hour exposure to BKIDC-1553 and 1553-N-Me

commonly resulted in G1 cell cycle arrest of responders (LNCaP and LNCaP95), but not

a non-responder line (PC3) which had no enrichment of the G1 population (**Figure 2A,**

Figure S4A and S4C). In addition, our negative control BKIDC-1817 did not affect the

cell cycle of any cell line. Similar patterns of cell cycle profiles were observed at 48- and

72-h time points with no obvious appearances of the sub-G1 population in LNCaP95

(47)(**Figure S4B**), suggesting that cytostatic rather than cytopathic actions of BKIDCs

are responsible for their antiproliferative activities.

Since G1 but not G2/M arrest was dominantly associated with BKIDC-induced

changes in cell cycle profiles, we took advantage of reverse-phase protein arrays

(RPPA) to characterize robust biochemical alterations leading to G1 cell cycle arrest(48)

(**Figure 2B and 2C, Figure S5, Table S3**). This high-throughput antibody-based

targeted proteomics platform allowed us to observe a time- and dose-dependent

decrease in phosphorylated ribosomal protein S6 at serine 235/S236 (P-RPS6$^{S235/S236}$)

and serine 240/S244 (P-RPS6$^{S240/S244}$) in LNCaP95 but not PC3 cells as early as 4 h

after treatment with BKIDC-1553 and 1553-N-Me (**Figure 2C, Figure S6**). RPS6 is a

canonical downstream target of mTORC1 kinase which is negatively regulated by the

metabolic sensor AMP-activated protein kinase (AMPK)(49, 50). Indeed,

antiproliferative BKIDC-1553-N-Me, but not our negative control BKIDC-1817,

consistently activated AMPK in LNCaP95 cells as reflected by increased enzymatic

activity and increased phosphorylation of acetyl-CoA carboxylase (ACC) at Ser79 (the

major substrate for AMPK)(49) (**Figure 3A and 3B**).

We next performed an in vitro enzyme assay using purified human AMPKα1β1γ1

complex to examine whether BKIDC-1553 and 1553-N-Me could be direct activators of

AMPK. A well-established allosteric activator A-769662 caused 2-fold increase in AMPK

activity (51) while BKIDCs failed to do so, suggesting BKIDCs do not stimulate AMPK

activity directly (**Table S4**).

Since AMPK serves as a cellular sensor of ATP-ADP-AMP balance in cells, the

alteration in ATP metabolism could be upstream of pathways that affect ATP levels and

produce AMPK activation(49). We examined BKIDC's effects on ATP supply from two

major pathways, glycolysis in the cytosol and oxidative phosphorylation (OXPHOS) in

mitochondria. OXPHOS inhibition, which was achieved by oligomycin (inhibitor for F0

component of H^+-ATP-synthase), antimycin A (AA: inhibitor for complex III), rotenone

(rot: complex I inhibitor), or carbonyl cyanite-4 (trifluoromethoxy) phenylhydrazone

(FCCP: uncoupler)(52), cooperated with BKIDCs to deplete intracellular ATP (**Figure**

3C, Figure S7). This suggested that BKIDCs inhibit glycolysis-derived ATP production,

and indeed a similar effect was seen with the glycolysis inhibitor 2-deoxyglucose (2-

DG). Also, similar to 2-DG, BKIDC-1553 and -1553-N-Me cooperated with oligomycin to

exhibit cytopathic, apoptotic actions in LNCaP as judged by increased PARP cleavage

and cell detachment (47) (**Figure S8**).

BKIDC effects on glycolysis

To examine the effects of BKIDCs on glycolysis directly, we employed the

Seahorse assay to measure the extracellular acidification rate (ECAR), a proxy for

glycolysis rate in cells(53). ECAR was suppressed by BKIDCs in LNCaP and LNCaP95

cells but not in BKIDC non-responder cell lines, including DU145, PC3, and U2OS

(**Figure 3D and 3E, Figure S9**). Mitochondrial respiration as judged by oxygen

consumption rate (OCR) (54) was not affected by BKIDC-1553-N-Me in both responder

(LNCaP, LNCaP95) and non-responder cell lines (PC3, DU145)(**Figure S9**). The rapid

decrease in ECAR with BKIDCs suggests that BKIDCs have a direct antiglycolytic effect

(**Figure 3F**). Under the same conditions, BKIDCs induced a moderate compensatory

increase in the oxygen consumption rate (OCR) in LNCaP95(**Figure 3F**), suggesting

aerobic, mitochondrial oxygen consumption partially compensates for inhibition of

glycolysis.

In order to determine which glycolytic step is interrupted by BKIDCs, we

performed a metabolic flux assay with D-[U-^{13}C] glucose(55) (**Figure 4A and 4B**). We

exposed cells to labelled glucose with different concentrations of BKIDC-1553-N-Me

and DMSO as a solvent control for 30 min. We did not see significant changes in

glucose levels among cells lines and treatment. In LNCaP and LNCaP95 cells, we

clearly saw a decrease in the levels of 6 carbon sugar phosphate which includes

glucose 6-phosphate and fructose 6-phosphate. Consistently we observed marked

reduction in levels of downstream metabolites, including dihydroxyacetone phosphate

(DHAP), phosphoglycerate (PG), phosphoenolpyruvate (PEP), and pyruvate (**Figure**

4B). Importantly, the non-responder cell line DU145 did not display such metabolic

alterations. These experiments suggested that either glucose uptake or glucose

phosphorylation were affected by BKIDC exposure in susceptible PCa cells(52).

Supporting this, results from the Glucose Uptake-Glo Assay also were consistent with

BKIDC-1553-N-Me blocking glucose uptake and/or phosphorylation (**Figure S10A**).

To distinguish between glucose uptake and glucose phosphorylation steps, we

used two different radiolabeled glucose analogs, [^{3}H] 2-DG and [^{3}H] 3-O-methyl D-

glucose (3-OMG), which are taken up into cells in the same manner as glucose (**Figure**

4C and 4D). 2-DG undergoes phosphorylation to become trapped and accumulates in

the cells as a form of 2DG-6P. On the other hand, 3-OMG cannot be phosphorylated

and its intracellular levels reflect the cell's capability of glucose uptake alone(30). We

confirmed time-dependent accumulation of 2-DG signals in both LNCaP and DU145

(**Figure S10B**). This accumulation was inhibited by the presence of BKIDC-1553-N-Me

in LNCaP but not DU145 cells. BKIDC-1553-N-Me did not inhibit 3-OMG uptake in

LNCaP cells, suggesting that BKIDC inhibits the glucose phosphorylation step, not

glucose uptake, in BKIDC susceptible cells (**Figure 4D**).

Hexokinases carry out the phosphorylation of glucose and we wanted to learn if

BKIDCs were dependent on a specific isoform of hexokinase (HK). Among five isoforms

of hexokinases expressed in humans (HK1, HK2, HK3, GCK1/HK4, and HKDC1)(56),

HK1 and HK2 are reportedly predominant in LNCaP cells(57). We used a CRISPR-

Cas9 system to generate HK1 and HK2 knock-out (KO) LNCaP clones. We successfully

established several independent HK1 and HK2 KO clones, demonstrating that HK1 and

HK2 are dispensable, and deletion of each gene did not affect cell proliferation and

survival (**Figure 5 and Figure S11**). We then characterized their sensitivity to BKIDCs.

We evaluated intracellular ATP levels, in the presence of oligomycin (OXPHOSi), to

determine BKIDC's antiglycolytic activity in the HK1 and HK2 KO lines. Like the parental

LNCaP line, individual HK1 KO clones were sensitive to both BKIDC-1553 and 1553-N-

Me, demonstrating lower ATP levels in the presence of oligomycin. In sharp contrast,

HK2 KO clones displayed remarkable resistance to antiglycolytic action of BKIDCs, in

that ATP levels were not suppressed in the presence of oligomycin. Pooled KO lines

with different sgRNAs also displayed resistance to BKIDC only with HK2 KO pools, but

not with HK1 KO (**Figure S12**). This suggested that BKIDCs exert their antiglycolytic

activity through HK2, but not HK1, in susceptible PCa cell lines. Collectively, these

findings implicate HK2-mediated glucose phosphorylation as the primary target of

BKIDC. An unexpected finding was that both HK1 and HK2 KO clones retained

antiproliferative sensitivity to BKIDCs (**Figure S11**). This suggests that BKIDCs may

exert antiproliferative effects beyond effects on glycolysis. But from these experiments,

it is clear that BKIDCs require HK2, but not HK1, for their effects on ATP generation

from glycolysis.   We explored whether HK2 could be directly targeted by BKIDCs. We

were unable to demonstrate that BKIDCs inhibit human recombinant HK2 (hrHK2)

enzymatic activity, in phosphorylation of glucose to glucose-6-phosphate (**Figure S13**).

Similarly, HK2 expression does not determine whether a cell line is sensitive. BKIDC-

resistant cell lines PC3 and DU145 express HK2 and its expression levels are

comparable irrespective of susceptibility to BKIDC (**Figure S14**). This

suggests that BKIDCs do not act directly on HK2, but likely act on regulatory molecules

(pathway's) that are known to modulate HK2 function(58).

1057

BKIDC effects in preclinical models of prostate cancer

We next tested the therapeutic potential of BKIDC-1553 in in vivo preclinical PCa

models. The slow clearance of BKIDC-1553 from the systemic circulation of mice led to

exposures of over 2 µM for an extended period of up to 72 h post-dose, which allowed a

Monday, Wednesday, Friday dosing regimen (**Supplementary Figure S15**)**.** BKIDC-

1553, administered 20 mg/kg orally, 3 times/week, decreased tumor growth after 5

weeks in a human PCa patient-derived xenograft (PDX), LuCaP 35 (59) while PC3

tumors derived from adeno-PCa cell-line derived xenografts were resistant to BKIDC-

1553 (**Figure 6A**). Intratumoral concentrations of BKIDC-1553 48 h after the end of

dosing were similar in PC3 and LuCaP 35 tumors (about 8 µM, **Table S5**), ruling out

that observed drug resistance of PC3 is due to low exposure to BKIDC-1553.

Importantly, antitumor activities of BKIDC-1553 were comparable to those of standard

ARSi enzalutamide treatment in AR+ LuCaP 35 model. As demonstrated in the

responder line LNCaP95 cells in vitro (**Figure 3A**), we observed increased P-ACC

immunoreactivities in 1553-treated LuCaP 35 tumors but not enzalutamide-treated

tumors (**Figure 6B**). This implies 1553 also elicited metabolic rewiring in association

with AMPK activation in vivo. These observations are consistent with our in vitro studies

and may reflect the association of antiglycolytic action of BKIDC-1553 with antitumor

activity. These results demonstrate that BKIDC-1553 has good systemic exposure and

tissue penetration needed to achieve therapeutic levels. Furthermore, we tested

organoid cultures derived from neuroendocrine prostate cancer that is androgen

insensitive and resistant to AR-directed therapy. BKIDC-1553 elicited antiproliferative

activities in LTL331R organoids derived from treatment-induced neuroendocrine

prostate cancer xenograft model LTL331R(29) (**Figure 6C**). Taken together, these

preclinical studies suggest BKIDC-1553 could be developed as a safe and effective

therapy of advanced PCa.

1084

BKIDC-1553 is an advanced preclinical candidate

BKIDC-1553 has the properties of a suitable advanced preclinical candidate in

models of pharmacokinetics (PK), in vitro adsorption-distribution-metabolism-excretion

(ADME), and toxicology in vitro and in vivo.

PK: We previously reported BKIDC-1553 (compound 32) has excellent PK

parameters when it is administered in a single oral dose to mice, rats, calves, dogs and

monkeys (35). We have reanalyzed these data, summarized in **Supplementary Data**

Table 6, and demonstrate >65% oral bioavailability in three species and slow clearance

in all species. The exposure obtained in each species is in accordance with allometric

scaling, suggesting oral dosing will be successful in humans, and allometric scaling can

be used to estimate the human dose for efficacy. To establish the efficacious exposure,

we dosed SCID mice for 2 weeks with a dose that would allow for systemic exposures

equivalent to concentrations successful in inhibiting human PDX LuCaP 35 in vitro,

namely, 20 mg/kg BKIDC given orally on Monday, Wednesday, and Friday. Plasma

samples were collected to assess for BKIDC levels. The AUC_{0-288h} (12 days) was

estimated to be 3,190 µmol·h/L. The highest exposure over any 24 h period was

estimated to be 381.8 µmol·h/L with a C_{max} of 19.4 µM, and these values were used to

compare to the highest exposure over a 24 h period during toxicology experiments for

determination of a safety index. BKIDC-1553 is predicted to have a ~17 h half-life in

humans based on an estimated Cl of 1.1 mL/min/kg and V of 1.6 L/kg in humans. Using

the average BKIDC concentration (11.1 µM)   and C_{max} (19.4 µM) associated with

efficacy in the preclinical mouse prostate cancer model, the estimated efficacious oral

dose in a 74 kg man will be 500 mg per day.

ADME: We previously reported that BKIDC-1553 has moderate solubility and oral

absorption consistent with an oral preclinical candidate(35). BKIDC-1553 binds to

human plasma proteins at 97%, which correlates with approximately 3% free fraction in

plasma. Studies of permeability in Caco-2 cells and MDR1-MDCKII cells demonstrated

high permeability (Caco-2 $P_{app\ A-B}$: 61 ± 9.6 x 10^{-6} cm/s and $P_{app\ B-A}$: 35 ± 1.0 x 10^{-6} cm/s;

MDR1-MDCKII $P_{app\ A-B}$: 15.6 ± 0.3 x 10^{-6} cm/s and $P_{app\ B-A}$: 4.2 ± 0.6 x 10^{-6} cm/s) and no

apparent efflux pump activity (Caco-2 efflux ratio: 0.6 ± 0.1; MDR1-MDCKII efflux ratio:

0.27 ± 0.03). Hepatocyte metabolism with human, rats, and dog microsomes

demonstrated similar metabolites for all three species (**Figure S16**). Inhibition of

cytochrome P450 (CYP) was assessed in vitro, and 20 µM BKIDC-1553 had no effect

on CYPs 1A2, 2C9, 2C19, 2D6, or 3A4. In separate studies, 10 µM BKIDC-1553

resulted in <30% inhibition of CYP2B6, but did inhibit CYP2C8 by a mean of 56%.

These data indicate that there is low potential for drug-drug-interactions associated with

CYP 1A2, 2B6, 2C9, 2C19, 2D6, or 3A4. Follow up dose response studies are planned

in order to determine the IC_{50} for BKIDC-1553 for CYP2C8 metabolism and better

predict the possible effects on CYP2C8 drug metabolism before submission for

Investigative New Drug (IND) status with the U.S. Food and Drug Administration (FDA).

Toxicology: As noted above (**Figure 1**), BKIDC-1553 has little to no effect on

normal mammalian cells (HFF) in vitro. We previously reported BKIDC-1553 has no

discernable effect on the cardiac risk channel, hERG, and an 80 human protein kinase

screen revealed possible interaction only with PKD3 (IC_{50} of 0.12 µM), and MEK2 (IC_{50}

of 0.97 µM)(35). BKIDC-1553 was not genotoxic in a bacterial mutagenicity assay

(microAmes assay) or an in vitro micronucleus study. BKIDC-1553 was assessed at 10

µM for possible binding or inhibition to 44 receptors and enzymes in a Safety 44 Screen

(https://cdnmedia.eurofins.com/corporate-

eurofins/media/1069358/safetyscreen44_epdsfl420june16.pdf)(42). Of the 44 receptors

and enzymes, BKIDC-1553 had activity in excess of 66% of control for only one target,

Angiotensin Converting Enzyme (ACE), which was 100% inhibited at 10 µM. We

described previously that mice tolerated a single 500 mg/kg oral dose and 5 days of

daily dosing at 100 mg/kg using a polyetheleneglycol-ethanol-saline vehicle(35). We

conducted an additional toxicology experiment for 7 days in mice with a lipid-based

(Phosal 53 MCT) vehicle and found that mice tolerated 35 mg/kg/day and 70 mg/kg/day

without clinical signs, CBC or biochemical changes, or histologic abnormalities, but did

not tolerate 100 mg/kg/day, which led to clinically moribund animals after 3 days of

dosing. Investigation of these mice revealed plasma glucose levels averaging >300

mg/dL, which presumably resulted in hyperglycemia. No other signs or sources of

toxicity could be found in these animals. BKIDC-1553 blood exposures at this level were

>5-times the exposure necessary for therapeutic effects on PDX, and thus a 5X margin

of safety exists for therapy. Similar dose acceleration experiments were conducted in

Beagle dogs treated with 1.7, 5, or 15 mg/kg BKIDC-1553 for 14 days, and no clinical,

hematologic, or biochemical abnormalities were observed. However, the exposure

($AUC_{last\ 24\ hours}$) from the highest dose (15 mg/kg) was estimated to yield only 60% of

exposure estimated in therapeutic dosing in mice. We subsequently investigated a

dosing regimen of 30 mg/kg administered orally twice a day for 4 doses (2 days),

followed by 15 mg/kg administered twice a day in a lipidic vehicle for 6 doses (3 days),

which yielded exposures approximately equal to therapeutic exposures measured in

mice. In this experiment, after administration of 2 doses of 30 mg/kg twice a day to

dogs, two subsequent doses resulted in emesis, leading to the dose reduction to 15

mg/kg twice a day, noted above. Similar vomiting was not seen dogs treated with 5-day

twice a day administration of vehicle alone. This suggests that vomiting was not due to

the vehicle, and vomiting may limit dose escalation in dogs. Moreover, plasma glucose

levels, serum chemistry, hematology, and coagulation and urinalysis were

unremarkable. Advanced cardiac testing was performed in anesthetized rats with

accelerating IV infusions of BKIDC-1553 to monitor drug induced changes in mean

arterial blood pressure, heart rate, and cardiac contractility (**Table S7**). There were no

significant changes in the cardiovascular parameters at IV doses of 3 mg/kg and 10

mg/kg but at 30 mg/kg there was a 25% increase in cardiac contractility. Plasma

BKIDC-1553 levels at the 30 mg/kg were 55 µM, far surpassing the exposure necessary

for therapy, thus the compound has a good cardiovascular safety index in rats.

Altogether, in vivo assessments in mice, rats, and dogs demonstrate tolerability of

BKIDC-1553 at doses that are pharmacologically active indicating potential therapeutic

benefit to patients without drug-induced toxicity.

1170

DISCUSSION

Despite considerable efforts, precision oncology in advanced PCa has faced

challenges, including drug resistance and limited populations who benefit from the

related therapeutics(8). Metabolic reprogramming is observed in emergent CRPC after

initial therapies fail(60). In particular, emergent CRPC becomes highly glycolytic as

demonstrated by positive [18]F-fluorodoxyglucose ([18]F-FDG) labeling in metastatic CRPC

but not primary PCas(61-63). Strong avidity to FDG in advanced PCa suggests that

targeting glycolysis would have a major impact on PCa progression and would be

especially important in treating those tumors that currently are untreatable or have

limited potential treatments(64).

However, the therapeutic targeting of metabolic complexes or enzymes in

glycolysis is challenging, affecting not only tumor cells but also normal proliferating

cells. Multiple enzymatic isoforms can also perform overlapping functions, and

metabolic flexibility can divert processes away from inhibited pathways. Mammals

possess five HK isoenzymes, including HK1, HK2, HK3, HK4/GCK1, and HKDC1 with

overlapping and specific tissue and cellular distribution patterns to fix glucose as the

first step in glycolysis(56). Along with catalytic domains, HK1 and HK2 contain N-

terminal motifs to interact with VDAC1 for association with mitochondria and the

glucose-6-phosphate (G6P)-binding sties for feedback allosteric inhibition(56). HK2 is

required for oncogenic transformation and progression of some cancers, but HK2 is

minimally expressed in most adult tissues and its systemic deletion in adult mice does

not lead to overt side effects(22). Thus, HK2 garners much attention as an ideal and

selective target in cancer therapeutics(65, 66). Target-based screening has identified

HK2 selective inhibitors that competitively or allosterically inhibit catalytic functions or

promote dissociation from the mitochondria, though they are still in early phases of

clinical development(67).

As demonstrated here, using phenotypic screening and target deconvolution, we

identified a series of compounds that fulfil a major unmet need for development of

metabolism-based therapeutic interventions(15). It is clear from Seahorse ECAR

measurements that BKIDCs interrupt glycolysis almost immediately after exposure in

susceptible PCa cells. This is accompanied by a rapid reduction in glycolysis

metabolites. Experiments showed that 2-DG accumulation, but not 3-OMG

accumulation, was interrupted by BKIDCs. This demonstrated that hexokinase activity,

but not glucose importation, was targeted by BKIDCs. Finally, knockout of HK2, but not

HK1, abrogated the reduced ATP levels observed with BKIDCs combined with

oligomycin. This demonstrated that BKIDCs interrupt ATP generation by glycolysis, and

appear to do so in a HK2 dependent fashion. This makes sense, in that PCa is known to

activate HK2 expression and become dependent on it for glycolysis(24).

We explored whether HK2 could be directly targeted by BKIDCs. BKIDC-1553

did not inhibit recombinant-HK2 (**Figure S13**). In addition to this, HK2 expression does

not determine whether a cell line is sensitive. BKIDC-resistant cell lines PC3 and

DU145 express HK2 protein (**Figure S14**). This suggests that BKIDCs do not act

directly on HK2, but could act on the many molecules that are known to modulate HK2

function(58). We hypothesize the regulatory pathways of BKIDC-responsive prostate

cancers are different from non-BKIDC responsive PC3 and DU145 cells, and other cell

lines. This difference suppressing the HK2 function of susceptible PCa cells, leading to

the suppression of glycolysis immediately after exposure of BKIDCs (Seahorse

experiments, **Figure 3F**).

As noted, both HK2 and HK1 KO cell lines retained sensitivity to antiproliferative

effects of BKIDCs. Observed antiproliferative phenotypes may reflect

polypharmacological effects of BKIDCs. Indeed, it is still a matter of debate whether

HK2 inhibition is sufficient to elicit antitumor activity(24, 68-71). HK1 and HK2 are

dispensable in LNCaP cells as knockout of each gene did not affect cell proliferation

and survival (this study) (69). However, metabolic reprogramming of the HK1 or HK2

KO population may change the coupling of glycolysis to antiproliferative effects. The

lack of effect on proliferation may be due to BKIDC effects on other targets or may

reflect the plasticity of PCa, allowing proliferation after complete KO of HK2(72, 73).

BKIDCs elicit cell-cycle arrest and AMPK activation, presumably due to the

integrated stress response(74). Depending on the context of AMPK expression, e.g.

spatial location in a cell, AMPK activation can be associated with a proliferation

response(75). Indeed, AMPK may work together with HK2 inhibition in the BKIDCs'

effects on PCa. As we demonstrated here, BKIDCs synergize with OXPHOS inhibition

(oligomycin) to promote cell death. A similar effect to BKIDCs could be seen with the

non-specific glycolysis inhibitor, 2-DG. However, BKIDCs did not exhibit the toxicity to

multiple cell lines that 2-DG demonstrates, making BKIDCs a good candidate for clinical

therapy. HK2 inhibition reportedly renders therapeutic vulnerability to interference with

OXPHOS or glutamine metabolism(76-78). This warrants exploration of synthetic lethal

pairings for metabolism-targeted combined therapies to improve clinical results of

BKIDC actions.

BKIDC-1553-N-Me and BKIDC-1553 were evaluated as possible leads to

advance as preclinical candidates. BKIDC-1553-N-Me would be advantageous,

potentially, through its lack of demonstrated "off-target" activity on protein kinases

PKD3, RIPK2, and MEK2, and BKIDC-1553 has weak inhibitory activity towards these

kinases. However, BKIDC-1553-N-Me was found to be rapidly converted to BKIDC-

1553 after oral dosing in mice, and thus was not advantageous to BKIDC-1553 for oral

administration.

BKIDC-1553 has the properties of a suitable advanced preclinical candidate with

its ability to arrest prostate cancer xenograft growth in the SCID implantation model, as

well as its PK, in vitro ADME, and toxicology attributes in vitro and in vivo. In the SCID

xenograft LuCaP 35 model, 20 mg/kg of BKIDC-1553 delivered orally three times

weekly was efficacious; even after 8 weeks of therapy, no toxicity was observed in the

BKIDC-1553-treated animals, though vehicle controls with tumors needed to be

euthanized per animal care committee regulations. The PK parameters of BKIDC-1553

have been examined in mice, rats, dogs, and monkeys, and excellent oral bioavailability

has been found in all species tested. In vitro ADME testing supports that CYP

metabolism likely leads to clearance of BKIDC-1553, but consistent with its slow

clearance in multiple species by liver microsomes and hepatocytes, it is slowly cleared

in all species tested in vitro. Moreover, BKIDC-1553 is highly bound to plasma protein

(97% human plasma protein bound), which likely contributes to its slow clearance (only

unbound drug is subject to hepatic clearance) and its safety (less free drug to contribute

to safety issues in vivo). This slow clearance explains why such low doses (20 mg/kg),

intermittently orally administered (3 times per week) in mice are successful for therapy

of the PDX model. Comparative in vitro metabolism testing demonstrates that human,

rat, and dog hepatocytes metabolize BKIDC-1553 to form identical metabolites,

demonstrating that rats and dogs are excellent to study for final GLP-toxicology required

for application for an IND status for BKIDC-1553. In vitro ADME studies suggest low

probability of drug-drug interactions, perhaps with the exception of drugs that interact

with CYP2C8, but this needs confirmatory dose response studies to adequately assess

this risk. In vitro safety studies demonstrate lack of binding to hERG cardiac toxicity ion

channels at 10 µM; and 10 µM is far beyond the unbound free levels measured during

successful therapy. An 80 human kinase off-target screen revealed possible interaction

only with PKD3 (IC_{50} of 0.12 µM), and MEK2 (IC_{50} of 0.97 µM)(35). However, these

studies were carried out without ATP. ATP is a competitor for inhibition of BKIDC-1553.

Therefore, given the high intracellular concentration of ATP (mM range) and high

plasma protein binding of BKIDC-1553 (only free levels, ~3% of plasma levels, should

drive off-target effects), it is not likely that these protein kinases will be inhibited during

therapy. BKIDC-1553 gave no signals for genotoxicity (microAmes test or in vitro

micronucleus assays) or for interaction with a 44 off-target screen of receptors and

enzymes of toxicological concern, except for 100% inhibition of angiotensin converting

enzyme (ACE) at 10 µM. ACE inhibitors are generally considered safe, but follow on

dose response experiments will be carried out to further explore this result prior to IND

application. Finally, preliminary experiments in mice and dogs with short term, multiple

day oral administration experiments demonstrate safety at observed therapeutic levels.

Mouse experiments suggest a therapeutic window of about 5-fold, with hyperglycemia

being the limiting toxicity. Dog experiments suggest vomiting could be an issue at

exposures above therapeutic exposures. Both hyperglycemia and vomiting are toxicity

issues that can be monitored and managed easily in human clinical trials, and during

therapy after regulatory approval. A pre-IND assessment with the U.S. FDA suggests

that pending a final Good Laboratory Practice (GLP) Toxicology and Chemistry

Manufacturing and Controls (CMC) development study, BKIDC-1553 is on track for a

successful IND application. Thus, we conclude BKIDC-1553 is an outstanding

preclinical candidate for development against prostate cancer.

1293

In summary, we overcame the hurdle of targeting tumor metabolism by

developing a series of druggable small molecules, BKIDCs, that elicit antiproliferative

effects selectively in PCa models. BKIDCs rapidly shut down glycolytic energy

production, an effect that appears dependent on HK2 expression. BKIDC-1553 was

effective in vitro and in vivo in shutting down prostate cancer growth, and demonstrates

outstanding PK, ADME, and safety properties consistent with a late preclinical lead.

Only CMC/Formulation/Stability and GLP toxicology studies remain before an IND

status can be requested from the U.S. FDA, to advance this promising molecule to

clinical trials of advanced prostate cancer. Since this compound has a unique

mechanism of action, it will be a welcome addition to the chemotherapy of advanced

prostate cancer, which generally becomes resistant to the available androgen-directed

therapy, and may be an addition or alternative to rescue cytotoxic chemotherapy

regimens that are very toxic and fail to deliver lasting response.

1307

1308

References

1.	Wang L, Lu B, He M, Wang Y, Wang Z, Du L. Prostate Cancer Incidence and Mortality: Global Status and Temporal Trends in 89 Countries From 2000 to 2019. Front Public Health. 2022;10. doi: 10.3389/fpubh.2022.811044.

2.	Rebello RJ, Oing C, Knudsen KE, Loeb S, Johnson DC, Reiter RE, Gillessen S, Van der Kwast T, Bristow RG. Prostate cancer. Nat Rev Dis Primers. 2021;7(1):9. doi: 10.1038/s41572-020-00243-0.

3.	Sumanasuriya S, De Bono J. Treatment of Advanced Prostate Cancer-A Review of Current Therapies and Future Promise. Cold Spring Harb Perspect Med. 2018;8(6). doi: 10.1101/cshperspect.a030635.

4.	Watson PA, Arora VK, Sawyers CL. Emerging mechanisms of resistance to androgen receptor inhibitors in prostate cancer. Nat Rev Cancer. 2015;15(12):701-11. doi: 10.1038/nrc4016.

5.	Mateo J, Lord CJ, Serra V, Tutt A, Balmana J, Castroviejo-Bermejo M, Cruz C, Oaknin A, Kaye SB, de Bono JS. A decade of clinical development of PARP inhibitors in perspective. Ann Oncol. 2019;30(9):1437-47. doi: 10.1093/annonc/mdz192.

6.	Sartor O, de Bono J, Chi KN, Fizazi K, Herrmann K, Rahbar K, Tagawa ST, Nordquist LT, Vaishampayan N, El-Haddad G, Park CH, Beer TM, Armour A, Perez-Contreras WJ, DeSilvio M, Kpamegan E, Gericke G, Messmann RA, Morris MJ, Krause BJ, Investigators V. Lutetium-177-PSMA-617 for Metastatic Castration-Resistant Prostate Cancer. N Engl J Med. 2021;385(12):1091-103. doi: 10.1056/NEJMoa2107322.

7.   Parker C, Nilsson S, Heinrich D, Helle SI, O'Sullivan JM, Fossa SD, Chodacki A,

Wiechno P, Logue J, Seke M, Widmark A, Johannessen DC, Hoskin P, Bottomley D,

James ND, Solberg A, Syndikus I, Kliment J, Wedel S, Boehmer S, Dall'Oglio M,

Franzen L, Coleman R, Vogelzang NJ, O'Bryan-Tear CG, Staudacher K, Garcia-Vargas

J, Shan M, Bruland OS, Sartor O, Investigators A. Alpha emitter radium-223 and

survival in metastatic prostate cancer. N Engl J Med. 2013;369(3):213-23. doi:

10.1056/NEJMoa1213755.

8.   Mateo J, McKay R, Abida W, Aggarwal R, Alumkal J, Alva A, Feng F, Gao X,

Graff J, Hussain M, Karzai F, Montgomery B, Oh W, Patel V, Rathkopf D, Rettig M,

Schultz N, Smith M, Solit D, Sternberg C, Van Allen E, VanderWeele D, Vinson J, Soule

HR, Chinnaiyan A, Small E, Simons JW, Dahut W, Miyahira AK, Beltran H. Accelerating

precision medicine in metastatic prostate cancer. Nat Cancer. 2020;1(11):1041-53. doi:

10.1038/s43018-020-00141-0.

9.   Choi SYC, Ribeiro CF, Wang Y, Loda M, Plymate SR, Uo T. Druggable

Metabolic Vulnerabilities Are Exposed and Masked during Progression to Castration

Resistant Prostate Cancer. Biomolecules. 2022;12(11). doi: 10.3390/biom12111590.

10.   Zadra G, Ribeiro CF, Chetta P, Ho Y, Cacciatore S, Gao X, Syamala S, Bango

C, Photopoulos C, Huang Y, Tyekucheva S, Bastos DC, Tchaicha J, Lawney B, Uo T,

D'Anello L, Csibi A, Kalekar R, Larimer B, Ellis L, Butler LM, Morrissey C, McGovern K,

Palombella VJ, Kutok JL, Mahmood U, Bosari S, Adams J, Peluso S, Dehm SM,

Plymate SR, Loda M. Inhibition of de novo lipogenesis targets androgen receptor

signaling in castration-resistant prostate cancer. Proc Natl Acad Sci U S A.

2019;116(2):631-40. doi: 10.1073/pnas.1808834116.

11.   Zhang X, Dang CV. Time to hit pause on mitochondria-targeting cancer

therapies. Nat Med. 2023;29(1):29-30. doi: 10.1038/s41591-022-02129-y.

12.   Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. Cell.

2011;144(5):646-74. Epub 2011/03/08. doi: 10.1016/j.cell.2011.02.013.

13.   Faubert B, Solmonson A, DeBerardinis RJ. Metabolic reprogramming and cancer

progression. Science. 2020;368(6487). Epub 2020/04/11. doi:

10.1126/science.aaw5473.

14.   Vander Heiden MG, DeBerardinis RJ. Understanding the Intersections between

Metabolism and Cancer Biology. Cell. 2017;168(4):657-69. doi:

10.1016/j.cell.2016.12.039.

15.   Stine ZE, Schug ZT, Salvino JM, Dang CV. Targeting cancer metabolism in the

era of precision oncology. Nat Rev Drug Discov. 2022;21(2):141-62. doi:

10.1038/s41573-021-00339-6.

16.   Warburg O. Über den Stoffwechsel der Carcinomzelle. Naturwissenschaften.

1924;12(50):1131-7. doi: 10.1007/BF01504608.

17.   Landau BR, Laszlo J, Stengle J, Burk D. Certain metabolic and pharmacologic

effects in cancer patients given infusions of 2-deoxy-D-glucose. J Natl Cancer Inst.

1958;21(3):485-94. Epub 1958/09/01.

18.   Ahmad F, Cherukuri MK, Choyke PL. Metabolic reprogramming in prostate

cancer. Br J Cancer. 2021;125(9):1185-96. Epub 2021/07/16. doi: 10.1038/s41416-021-

01435-5.

19.   Uo T, Sprenger CC, Plymate SR. Androgen Receptor Signaling and Metabolic

and Cellular Plasticity During Progression to Castration Resistant Prostate Cancer.

Front Oncol. 2020;10:580617. doi: 10.3389/fonc.2020.580617.

20.   Cameron S, Deblois G, Hawley JR, Qamra A, Zhou S, Tonekaboni SAM,

Murison A, Van Vliet R, Liu J, Locasale JW, Lupien M. Chronic hypoxia favours

adoption to a castration-resistant cell state in prostate cancer. Oncogene.

2023;42(21):1693-703. Epub 2023/04/06. doi: 10.1038/s41388-023-02680-z.

21.   Beier AK, Puhr M, Stope MB, Thomas C, Erb HHH. Metabolic changes during

prostate cancer development and progression. J Cancer Res Clin Oncol.

2023;149(5):2259-70. Epub 2022/09/24. doi: 10.1007/s00432-022-04371-w.

22.   Patra KC, Wang Q, Bhaskar PT, Miller L, Wang Z, Wheaton W, Chandel N,

Laakso M, Muller WJ, Allen EL, Jha AK, Smolen GA, Clasquin MF, Robey B, Hay N.

Hexokinase 2 is required for tumor initiation and maintenance and its systemic deletion

is therapeutic in mouse models of cancer. Cancer Cell. 2013;24(2):213-28. Epub

2013/08/06. doi: 10.1016/j.ccr.2013.06.014.

23.   Davidson SM, Schmidt DR, Heyman JE, O'Brien JP, Liu AC, Israelsen WJ,

Dayton TL, Sehgal R, Bronson RT, Freinkman E, Mak HH, Fanelli GN, Malstrom S,

Bellinger G, Carracedo A, Pandolfi PP, Courtney KD, Jha A, DePinho RA, Horner JW,

Thomas CJ, Cantley LC, Loda M, Vander Heiden MG. Pyruvate Kinase M1 Suppresses

Development and Progression of Prostate Adenocarcinoma. Cancer Res.

2022;82(13):2403-16. Epub 2022/05/19. doi: 10.1158/0008-5472.CAN-21-2352.

24.   Wang L, Xiong H, Wu F, Zhang Y, Wang J, Zhao L, Guo X, Chang LJ, Zhang Y,

You MJ, Koochekpour S, Saleem M, Huang H, Lu J, Deng Y. Hexokinase 2-mediated

Warburg effect is required for PTEN- and p53-deficiency-driven prostate cancer growth.

Cell Rep. 2014;8(5):1461-74. Epub 2014/09/02. doi: 10.1016/j.celrep.2014.07.053.

25. Golkowski M, Vidadala RS, Lombard CK, Suh HW, Maly DJ, Ong SE. Kinobead

and Single-Shot LC-MS Profiling Identifies Selective PKD Inhibitors. J Proteome Res.

2017;16(3):1216-27. doi: 10.1021/acs.jproteome.6b00817.

26. Tandon M, Salamoun JM, Carder EJ, Farber E, Xu S, Deng F, Tang H, Wipf P,

Wang QJ. SD-208, a novel protein kinase D inhibitor, blocks prostate cancer cell

proliferation and tumor growth in vivo by inducing G2/M cell cycle arrest. PLoS One.

2015;10(3):e0119346. doi: 10.1371/journal.pone.0119346.

27. Tandon M, Johnson J, Li Z, Xu S, Wipf P, QJ W. New pyrazolopyrimidine

inhibitors of protein kinase D as potent anticancer agents for prostate cancer cells. PloS

ONE. 2013;8:e75601.

28. Bluemn EG, Coleman IM, Lucas JM, Coleman RT, Hernandez-Lopez S,

Tharakan R, Bianchi-Frias D, Dumpit RF, Kaipainen A, Corella AN, Yang YC, Nyquist

MD, Mostaghel E, Hsieh AC, Zhang X, Corey E, Brown LG, Nguyen HM, Pienta K,

Ittmann M, Schweizer M, True LD, Wise D, Rennie PS, Vessella RL, Morrissey C,

Nelson PS. Androgen Receptor Pathway-Independent Prostate Cancer Is Sustained

through FGF Signaling. Cancer Cell. 2017;32(4):474-89.e6. doi:

10.1016/j.ccell.2017.09.003.

29. Ci X, Hao J, Dong X, Xue H, Wu R, Choi SYC, Haegert AM, Collins CC, Liu X,

Lin D, Wang Y. Conditionally Reprogrammed Cells from Patient-Derived Xenograft to

Model Neuroendocrine Prostate Cancer Development. Cells. 2020;9(6). doi:

10.3390/cells9061398.

30.   Tanti JF, Cormont M, Gremeaux T, Le Marchand-Brustel Y. Assays of glucose

entry, glucose transporter amount, and translocation. Methods in molecular biology

(Clifton, NJ). 2001;155:157-65. doi: 10.1385/1-59259-231-7:157.

31.   Sweet IR, Cook DL, Lernmark A, Greenbaum CJ, Wallen AR, Marcum ES,

Stekhova SA, Krohn KA. Systematic screening of potential beta-cell imaging agents.

Biochem Biophys Res Commun. 2004;314(4):976-83. doi: 10.1016/j.bbrc.2003.12.182.

32.   Sanjana NE, Shalem O, Zhang F. Improved vectors and genome-wide libraries

for CRISPR screening. Nat Methods. 2014;11:783-4.

33.   Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, Smith I,

Tothova Z, Wilen C, Orchard R, Virgin HW, Listgarten J, Root DE. Optimized sgRNA

design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat

Biotechnol. 2016;34(2):184-91. doi: 10.1038/nbt.3437.

34.   Rabinovitch PS. DNA content histogram and cell-cycle analysis. Methods Cell

Biol. 1994;41:263-96. Epub 1994/01/01. doi: 10.1016/s0091-679x(08)61723-9.

35.   Vidadala RS, Rivas KL, Ojo KK, Hulverson MA, Zambriski JA, Bruzual I, Schultz

TL, Huang W, Zhang Z, Scheele S, DeRocher AE, Choi R, Barrett LK, Siddaramaiah

LK, Hol WG, Fan E, Merritt EA, Parsons M, Freiberg G, Marsh K, Kempf DJ, Carruthers

VB, Isoherranen N, Doggett JS, Van Voorhis WC, Maly DJ. Development of an Orally

Available and Central Nervous System (CNS) Penetrant Toxoplasma gondii Calcium-

Dependent Protein Kinase 1 (TgCDPK1) Inhibitor with Minimal Human Ether-a-go-go-

Related Gene (hERG) Activity for the Treatment of Toxoplasmosis. J Med Chem.

2016;59(13):6531-46. Epub 2016/06/17. doi: 10.1021/acs.jmedchem.6b00760.

36. Mordenti J. Man versus beast: pharmacokinetic scaling in mammals. J Pharm

Sci. 1986;75(11):1028-40. doi: 10.1002/jps.2600751104.

37. Weiss M, Sziegoleit W, Forster W. Dependence of pharmacokinetic parameters

on the body weight. Int J Clin Pharmacol Biopharm. 1977;15(12):572-5.

38. Mahmood I, Balian JD. Interspecies scaling: predicting clearance of drugs in

humans. Three different approaches. Xenobiotica. 1996;26(9):887-95. doi:

10.3109/00498259609052491.

39. Tang H, Mayersohn M. A mathematical description of the functionality of

correction factors used in allometry for predicting human drug clearance. Drug Metab

Dispos. 2005;33(9):1294-6. doi: 10.1124/dmd.105.004135.

40. Curran RE, Claxton CR, Hutchison L, Harradine PJ, Martin IJ, Littlewood P.

Control and measurement of plasma pH in equilibrium dialysis: influence on drug

plasma protein binding. Drug Metab Dispos. 2011;39(3):551-7. doi:

10.1124/dmd.110.036988.

41. Escobar PA, Kemper RA, Tarca J, Nicolette J, Kenyon M, Glowienke S, Sawant

SG, Christensen J, Johnson TE, McKnight C, Ward G, Galloway SM, Custer L, Gocke

E, O'Donovan MR, Braun K, Snyder RD, Mahadevan B. Bacterial mutagenicity

screening in the pharmaceutical industry. Mutat Res. 2013;752(2):99-118. Epub

2012/12/25. doi: 10.1016/j.mrrev.2012.12.002.

42. Bowes J, Brown AJ, Hamon J, Jarolimek W, Sridhar A, Waldron G, Whitebread

S. Reducing safety-related drug attrition: the use of in vitro pharmacological profiling.

Nat Rev Drug Discov. 2012;11(12):909-22. Epub 2012/12/01. doi: 10.1038/nrd3845.

43.   Segreti JA, Marsh KC, Polakowski JS, Fryer RM. Evoked Changes in

Cardiovascular Function in Rats by Infusion of Levosimendan, OR-1896

[(R)-N-(4-(4-Methyl-6-oxo-1,4,5,6-tetrahydropyridazin-3-

yl)phenyl)acetamide], OR-1855 [(R)-6-(4-Aminophenyl)-5-methyl-4,5-

dihydropyridazin-3(2H)-one], Dobutamine, and Milrinone: Comparative

Effects on Peripheral Resistance, Cardiac Output, dP/dt, Pulse Rate, and Blood

Pressure. Journal of Pharmacology and Experimental Therapeutics. 2008;325(1):331-

40. doi: 10.1124/jpet.107.132530.

44.   Choi R, Kelley A, Leibly D, Hewitt SN, Napuli A, Van Voorhis W. Immobilized

metal-affinity chromatography protein-recovery screening is predictive of

crystallographic structure success. Acta Crystallogr Sect F Struct Biol Cryst Commun.

2011;67(Pt 9):998-1005. doi: 10.1107/S1744309111017374.

45.   Ojo KK, Gillespie JR, Riechers AJ, Napuli AJ, Verlinde CL, Buckner FS, Gelb

MH, Domostoj MM, Wells SJ, Scheer A, Wells TN, Van Voorhis WC. Glycogen

synthase kinase 3 is a potential drug target for African trypanosomiasis therapy.

Antimicrob Agents Chemother. 2008;52(10):3710-7. doi: 10.1128/AAC.00364-08.

46.   Shrestha A, Ojo KK, Koston F, Ruttkowski B, Vidadala RSR, Dorr CS, Navaluna

ED, Whitman GR, Barrett KF, Barrett LK, Hulverson MA, Choi R, Michaels SA, Maly DJ,

Hemphill A, Van Voorhis WC, Joachim A. Bumped kinase inhibitor 1369 is effective

against Cystoisospora suis in vivo and in vitro. Int J Parasitol Drugs Drug Resist.

2019;10:9-19. doi: 10.1016/j.ijpddr.2019.03.004.

47. Wlodkowic D, Telford W, Skommer J, Darzynkiewicz Z. Apoptosis and beyond:

cytometry in studies of programmed cell death. Methods Cell Biol. 2011;103:55-98. doi:

10.1016/B978-0-12-385493-3.00004-8.

48. Gujral TS, Karp RL, Finski A, Chan M, Schwartz PE, MacBeath G, Sorger P.

Profiling phospho-signaling networks in breast cancer using reverse-phase protein

arrays. Oncogene. 2013;32(29):3470-6. Epub 2012/09/05. doi: 10.1038/onc.2012.378.

49. Garcia D, Shaw RJ. AMPK: Mechanisms of Cellular Energy Sensing and

Restoration of Metabolic Balance. Mol Cell. 2017;66(6):789-800. doi:

10.1016/j.molcel.2017.05.032.

50. Meyuhas O. Ribosomal Protein S6 Phosphorylation: Four Decades of Research.

Int Rev Cell Mol Biol. 2015;320:41-73. Epub 2015/11/29. doi:

10.1016/bs.ircmb.2015.07.006.

51. Cool B, Zinker B, Chiou W, Kifle L, Cao N, Perham M, Dickinson R, Adler A,

Gagne G, Iyengar R, Zhao G, Marsh K, Kym P, Jung P, Camp HS, Frevert E.

Identification and characterization of a small molecule AMPK activator that treats key

components of type 2 diabetes and the metabolic syndrome. Cell Metab. 2006;3(6):403-

16. Epub 2006/06/07. doi: 10.1016/j.cmet.2006.05.005.

52. Lehninger AL NDCM. Principles of Biochemistry, 2nd edn. textbook. 1993.

53. Mookerjee SA, Nicholls DG, Brand MD. Determining Maximum Glycolytic

Capacity Using Extracellular Flux Measurements. PLoS One. 2016;11(3):e0152016.

doi: 10.1371/journal.pone.0152016.

54.   Gerencser AA, Neilson A, Choi SW, Edman U, Yadava N, Oh RJ, Ferrick DA,

Nicholls DG, Brand MD. Quantitative microplate-based respirometry with correction for

oxygen diffusion. Analytical chemistry. 2009;81(16):6868-78. doi: 10.1021/ac900881z.

55.   Antoniewicz MR. A guide to (13)C metabolic flux analysis for the cancer biologist.

Exp Mol Med. 2018;50(4):19. Epub 2018/04/17. doi: 10.1038/s12276-018-0060-y.

56.   Farooq Z, Ismail H, Bhat SA, Layden BT, Khan MW. Aiding Cancer's "Sweet

Tooth": Role of Hexokinases in Metabolic Reprogramming. Life (Basel). 2023;13(4).

Epub 2023/04/28. doi: 10.3390/life13040946.

57.   Ghandi M, Huang FW, Jane-Valbuena J, Kryukov GV, Lo CC, McDonald ER,

3rd, Barretina J, Gelfand ET, Bielski CM, Li H, Hu K, Andreev-Drakhlin AY, Kim J, Hess

JM, Haas BJ, Aguet F, Weir BA, Rothberg MV, Paolella BR, Lawrence MS, Akbani R,

Lu Y, Tiv HL, Gokhale PC, de Weck A, Mansour AA, Oh C, Shih J, Hadi K, Rosen Y,

Bistline J, Venkatesan K, Reddy A, Sonkin D, Liu M, Lehar J, Korn JM, Porter DA,

Jones MD, Golji J, Caponigro G, Taylor JE, Dunning CM, Creech AL, Warren AC,

McFarland JM, Zamanighomi M, Kauffmann A, Stransky N, Imielinski M, Maruvka YE,

Cherniack AD, Tsherniak A, Vazquez F, Jaffe JD, Lane AA, Weinstock DM,

Johannessen CM, Morrissey MP, Stegmeier F, Schlegel R, Hahn WC, Getz G, Mills

GB, Boehm JS, Golub TR, Garraway LA, Sellers WR. Next-generation characterization

of the Cancer Cell Line Encyclopedia. Nature. 2019;569(7757):503-8. doi:

10.1038/s41586-019-1186-3.

58.   Ciscato F, Ferrone L, Masgras I, Laquatra C, Rasola A. Hexokinase 2 in Cancer:

A Prima Donna Playing Multiple Characters. Int J Mol Sci. 2021;22(9). doi:

10.3390/ijms22094716.

59.   Corey E, Quinn JE, Buhler KR, Nelson PS, Macoska JA, True LD, Vessella RL.

LuCaP 35: a new model of prostate cancer progression to androgen independence.

Prostate. 2003;55(4):239-46. doi: 10.1002/pros.10198.

60.   Chetta P, Zadra G. Metabolic reprogramming as an emerging mechanism of

resistance to endocrine therapies in prostate cancer. Cancer Drug Resist.

2021;4(1):143-62. doi: 10.20517/cdr.2020.54.

61.   Fox JJ, Gavane SC, Blanc-Autran E, Nehmeh S, Gonen M, Beattie B, Vargas

HA, Schoder H, Humm JL, Fine SW, Lewis JS, Solomon SB, Osborne JR, Veach D,

Sawyers CL, Weber WA, Scher HI, Morris MJ, Larson SM. Positron Emission

Tomography/Computed Tomography-Based Assessments of Androgen Receptor

Expression and Glycolytic Activity as a Prognostic Biomarker for Metastatic Castration-

Resistant Prostate Cancer. JAMA Oncol. 2018;4(2):217-24. doi:

10.1001/jamaoncol.2017.3588.

62.   Jadvar H. Is There Use for FDG-PET in Prostate Cancer? Semin Nucl Med.

2016;46(6):502-6. doi: 10.1053/j.semnuclmed.2016.07.004.

63.   Bakht MK, Lovnicki JM, Tubman J, Stringer KF, Chiaramonte J, Reynolds MR,

Derecichei I, Ferraiuolo RM, Fifield BA, Lubanska D, Oh SW, Cheon GJ, Kwak C,

Jeong CW, Kang KW, Trant JF, Morrissey C, Coleman IM, Wang Y, Ahmadzadehfar H,

Dong X, Porter LA. Differential Expression of Glucose Transporters and Hexokinases in

Prostate Cancer with a Neuroendocrine Gene Signature: A Mechanistic Perspective for

(18)F-FDG Imaging of PSMA-Suppressed Tumors. J Nucl Med. 2020;61(6):904-10. doi:

10.2967/jnumed.119.231068.

64.   Wibmer AG, Morris MJ, Gonen M, Zheng J, Hricak H, Larson S, Scher HI,

Vargas HA. Quantification of Metastatic Prostate Cancer Whole-Body Tumor Burden

with (18)F-FDG PET Parameters and Associations with Overall Survival After First-Line

Abiraterone or Enzalutamide: A Single-Center Retrospective Cohort Study. J Nucl Med.

2021;62(8):1050-6. doi: 10.2967/jnumed.120.256602.

65.   Shan W, Zhou Y, Tam KY. The development of small-molecule inhibitors

targeting hexokinase 2. Drug Discov Today. 2022;27(9):2574-85. doi:

10.1016/j.drudis.2022.05.017.

66.   Ciscato F, Filadi R, Masgras I, Pizzi M, Marin O, Damiano N, Pizzo P, Gori A,

Frezzato F, Chiara F, Trentin L, Bernardi P, Rasola A. Hexokinase 2 displacement from

mitochondria-associated membranes prompts Ca(2+) -dependent death of cancer cells.

EMBO Rep. 2020;21(7):e49117. Epub 2020/05/10. doi: 10.15252/embr.201949117.

67.   Shan W, Zhou Y, Tam KY. The development of small-molecule inhibitors

targeting hexokinase 2. Drug Discov Today. 2022. doi: 10.1016/j.drudis.2022.05.017.

68.   Xu S, Herschman HR. A Tumor Agnostic Therapeutic Strategy for Hexokinase 1-

Null/Hexokinase 2-Positive Cancers. Cancer Res. 2019;79(23):5907-14. Epub

2019/08/23. doi: 10.1158/0008-5472.CAN-19-1789.

69.   Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S,

Wilson CJ, Lehar J, Kryukov GV, Sonkin D, Reddy A, Liu M, Murray L, Berger MF,

Monahan JE, Morais P, Meltzer J, Korejwa A, Jane-Valbuena J, Mapa FA, Thibault J,

Bric-Furlong E, Raman P, Shipway A, Engels IH, Cheng J, Yu GK, Yu J, Aspesi P, Jr.,

de Silva M, Jagtap K, Jones MD, Wang L, Hatton C, Palescandolo E, Gupta S, Mahan

S, Sougnez C, Onofrio RC, Liefeld T, MacConaill L, Winckler W, Reich M, Li N, Mesirov

JP, Gabriel SB, Getz G, Ardlie K, Chan V, Myer VE, Weber BL, Porter J, Warmuth M,

Finan P, Harris JL, Meyerson M, Golub TR, Morrissey MP, Sellers WR, Schlegel R,

Garraway LA. The Cancer Cell Line Encyclopedia enables predictive modelling of

anticancer drug sensitivity. Nature. 2012;483(7391):603-7. doi: 10.1038/nature11003.

70.   Nogueira V, Patra KC, Hay N. Selective eradication of cancer displaying

hyperactive Akt by exploiting the metabolic consequences of Akt activation. Elife.

2018;7. Epub 2018/04/25. doi: 10.7554/eLife.32213.

71.   Xu S, Catapang A, Braas D, Stiles L, Doh HM, Lee JT, Graeber TG, Damoiseaux

R, Shirihai O, Herschman HR. A precision therapeutic strategy for hexokinase 1-null,

hexokinase 2-positive cancers. Cancer Metab. 2018;6:7. Epub 2018/07/11. doi:

10.1186/s40170-018-0181-8.

72.   Beltran H, Hruszkewycz A, Scher HI, Hildesheim J, Isaacs J, Yu EY, Kelly K, Lin

D, Dicker A, Arnold J, Hecht T, Wicha M, Sears R, Rowley D, White R, Gulley JL, Lee J,

Diaz Meco M, Small EJ, Shen M, Knudsen K, Goodrich DW, Lotan T, Zoubeidi A,

Sawyers CL, Rudin CM, Loda M, Thompson T, Rubin MA, Tawab-Amiri A, Dahut W,

Nelson PS. The Role of Lineage Plasticity in Prostate Cancer Therapy Resistance. Clin

Cancer Res. 2019;25(23):6916-24. doi: 10.1158/1078-0432.CCR-19-1423.

73.   Davies AH, Beltran H, Zoubeidi A. Cellular plasticity and the neuroendocrine

phenotype in prostate cancer. Nat Rev Urol. 2018;15(5):271-86. doi:

10.1038/nrurol.2018.22.

74.   Vara-Ciruelos D, Dandapani M, Hardie DG. AMP-Activated Protein Kinase:

Friend or Foe in Cancer? Annual Review of Cancer Biology. 2020;4(1):1-16. doi:

10.1146/annurev-cancerbio-030419-033619.

75.   Khan AS, Frigo DE. A spatiotemporal hypothesis for the regulation, role, and

targeting of AMPK in prostate cancer. Nat Rev Urol. 2017;14(3):164-80. doi:

10.1038/nrurol.2016.272.

76.   DeWaal D, Nogueira V, Terry AR, Patra KC, Jeon S-M, Guzman G, Au J, Long

CP, Antoniewicz MR, Hay N. Hexokinase-2 depletion inhibits glycolysis and induces

oxidative phosphorylation in hepatocellular carcinoma and sensitizes to metformin.

Nature Communications. 2018;9(1):446. doi: 10.1038/s41467-017-02733-4.

77.   Mendez-Lucas A, Lin W, Driscoll PC, Legrave N, Novellasdemunt L, Xie C,

Charles M, Wilson Z, Jones NP, Rayport S, Rodriguez-Justo M, Li V, MacRae JI, Hay

N, Chen X, Yuneva M. Identifying strategies to target the metabolic flexibility of tumours.

Nat Metab. 2020;2(4):335-50. doi: 10.1038/s42255-020-0195-8.

78.   Guo W, Kuang Y, Wu J, Wen D, Zhou A, Liao Y, Song H, Xu D, Wang T, Jing B,

Li K, Hu M, Ling J, Wang Q, Wu W. Hexokinase 2 Depletion Confers Sensitization to

Metformin and Inhibits Glycolysis in Lung Squamous Cell Carcinoma. Front Oncol.

2020;10:52. doi: 10.3389/fonc.2020.00052.

1613

1614

1615

Acknowledgements:

We thank Dr. Taran Gujral and Stella Shin (Fred Hutchinson Cancer Center) for

help with the RPPA data, Michael S. Dey PhD and D. Bruce Burlington PhD for

their consultation, Dr. Michael Schweizer for discussion and manuscript editing,

and Mika Munari for technical assistance. We are grateful to Prof. Susan

Charman and the team within the Centre for Drug Candidate Optimisation,

Monash Institute of Pharmaceutical Sciences, Monash University, Parkville,

Australia for testing the ADME properties of BKIDC-1553. We are grateful to

Medicines for Malaria Venture (MMV) for providing access to testing for ADME

for BKIDC-1553. Funding: Lopker Family Foundation in honor of Karl Lopker,

who died of prostate cancer in 2018; WE-REACH, supported by University of

Washington and National Institutes of Health grant 5U01HL152401; US

Department of Defense grant W81XWH-17-1-0323 (SRP, DM, WCVV); US

Department of Defense grant W81XWH-20-1-0146 (SRP); US Department of

Veterans Affairs Merit Review grant I01BX0033248 (SRP); US Department of

Veterans Affairs Service Directed Project number SDR-PLYMATE (SRP);

National Institutes of Health/National Cancer Institute grant CA255830 (TU) and

National Institutes of Health/National Cancer Institute grant P50CA097186

(KKO).

1635

1636

1637

 Figures

Figure 1. BKIDC effects on Prostate Cancer Cell lines and other Cell Lines.

(A) Cell proliferation was evaluated by MTS assay at 72 h after BKIDC-1553 was added

at the range between 0.63-20 μM, except for VCaP at 120 h (DMSO as a vehicle

control). The data shown are representative of at least two independent experiments,

normalized to the values obtained from cells grown in medium containing 0.1% DMSO

vehicle and presented as mean ± standard deviation (SD) (n=3-5). One-way ANOVA

followed by Dunnett's test for multiple comparison: **$p <0.01$. **(B)** Structures of

representative BKIDCs. Lead BKIDCs have two main scaffolds: Pyrrolopyrimidines

(PrP) and pyrazolopyrimidines (PP) are two main scaffolds of BKIDCs. *Toxoplasma*

gondii calcium dependent protein kinase 1 (*Tg*CDPK1) has the small gatekeeper

residue glycine and can hold BKIDC-1553 in the ATP binding pocket. BKIDC-1553-N-

Me, 1820, and GP228 have distinct R2 groups while commonly lacking the protein

kinase binding and thus inhibition activity. **(C)** Graphical presentation of screening

results of over 123 BKIDC compounds (**Table S1**) tested at 10 µM in LNCaP95 and

PC3 (**Table S2**). Each dot represents one compound and the data are shown as mean

± SD. Compounds were categorized into three groups depending on the degree of

inhibition of proliferation in LNCaP95 and PC3.

1656

Figure 2. Cytostatic effects of anti-proliferative BKIDCs. (A) The percentage of cells

at G1, S, and G2 phase of cell cycle at 30 h after treatment with the indicated BKIDCs.

Cell cycle analysis showing G1 cell cycle arrest in LNCaP and LNCaP95 cells

compared to PC3 when treated with BKIDC-1553 and 1553-N-Me. These results were

confirmed in 2 subsequent independent experiments (**Figure S4**) **(B)** Heat map of

antigen levels detected by RPPA. Cellular lysates of LNCaP95 treated with the

indicated BKIDC at 20 μM for 24 h were subjected to RPPA with 70 distinct antibodies

(Antibodies and the corresponding RPPA signal intensities relative to β-actin signals are

listed in **Table S3**). Note BKIDC-1553 and 1553-N-Me commonly affected only

abundance of ribosomal S6 protein (RPS6) phosphorylated at S235/S236 or

S240/S244, the first two bands seen with BKIDC-1553 and the only bands seen with

BKIDC-1553-N-Me. The response to BKIDC-1817 was comparable to the one to

DMSO. **(C)** Time course and dose dependent studies of BKIDCs in LNCaP95 and PC3.

Time- and dose-dependent decrease in phosphorylated RPS6 signals in RPPA in

LNCaP95 but not PC3 as early as at 4 h after treatment with BKIDC-1553 and 1553-N-

Me.

1674

Figure 3. BKIDCs are antiglycolytic agents. (A) AMPK activation in LNCaP95 treated

with BKIDC-1553-N-Me. AMPK activity in total LNCaP95 lysates was measured after 1

h exposure to BKIDC-1553-N-Me or 1817. One-way ANOVA followed by Tukey's test

for multiple comparison: $**p < 0.01$. **(B)** Phosphorylated acetyl-CoA carboxylase (P-

ACC) at Ser79 was increased after treatment with BKIDC-1553 but not 1817. Metabolic

inhibitors 2-deoxyglucose (2-DG) and oligomycin (oligo) served as positive controls for

AMPK activation to increase P-ACC. **(C)** Effects of BKIDCs on intracellular ATP levels

in LNCaP and LNCaP95 cells. Cells were treated with BKIDC-1553-N-Me, glycolysis

inhibitor 2-DG, and OXPHOS inhibitors (OXPHOSi) singly or in combination for 120 min.

The data shown are representative of at least two independent experiments, normalized

to the values obtained from cells grown in medium containing 0.1% DMSO vehicle and

presented as mean ± SD (n=2). Rot: rotenone, FCCP: carbonyl cyanite-4

(trifluoromethoxy) phenylhydrazone, AA: Antimycin A. One-way ANOVA followed by

Dunnett's test for multiple comparison (vs. Oligomycin alone). **(D)** Extracellular

acidification rates (ECAR) were determined in cultured cells by the Seahorse XFe24

analyzer. Cells were pre-treated with 20 μM BKIDC for 1 h prior to application to

Seahorse analyzer. Increase in ECAR by glucose injection (glycolytic rates) was

inhibited by BKIDC-1553-N-Me in LNCaP95 but not PC3. Maximum glycolytic capacity

was evaluated by oligomycin injection. 2-DG injection shuts down glycolysis as reflected

by return of ECAR to the basal levels. **(E)** The effect of BKIDC-1553-N-Me on glycolytic

rates. The data shown are normalized to the values obtained from cells pre-treated with

DMSO vehicle and presented as mean ± SD (n=3). One-way ANOVA followed by

Tukey's test for multiple comparison: $**p < 0.01$. **(F)** Cells were exposed to BKIDC at the

indicated time by injection at the final concentration of 10 μM. Decrease in ECAR was

observed in LNCaP95 but not PC3 immediately after injection of BKIDC-1553-N-Me.

There was a compensatory increase in oxygen consumption rates (OCR) which reflects

mitochondrial oxidative respiration.

1702

Figure 4. Glucose phosphorylation is the primary target for BKIDC as an

antiglycolytic agent.

(A) Schematic flowchart of glycolysis and the metabolites measured in this metabolic

flux assay with [13]C-labeled glucose with BKIDC-1553-N-Me this experiment (dark

boxes). **(B)** Metabolic flux assay with [13]C-labeled glucose. [13]C enrichment of glycolytic

intermediates was determined by liquid chromatography coupled with mass

spectrometry (LC-MS) in DU145, LNCaP, and LNCaP95 cells at 30 min after exposure

to [U-^{13}C]-glucose in the presence and absence of BKIDC-1553-N-Me. The X-axis is the

concentration of BKIDC-1553-N-Me, with 0-20 µM added to the indicated cell lines.

Black columns indicate where unlabeled carbon (^{12}C, thus M+0 [M=mass expected for

^{12}C]) is found and grey columns originate from the ^{13}C-labeled glucose (^{13}C, thus M+6

or M+3). In LNCaP and LNCaP95 cells, addition of BKIDC-1553-N-Me inhibited glucose

being metabolized to 3-carbon glycolytic intermediates, DHAP, PG, PEP, or pyruvate,

indicating a block in glycolysis, at or prior to the six-carbon to 3-carbon fructose-

bisphosphate aldolase step. Note that in BKIDC non-responsive DU145 cells there is no

BKIDC-1553-N-Me effect on hexose-6-phosphate and 3-carbon glycolytic intermediates

(DHAP, PEP, PG, or pyruvate). The results trended to indicate that BKIDC blocked

effects at hexose-6-phosphate levels. One-way ANOVA followed by Dunnett's test for

multiple comparison for M +3 and M + 6 isotopomers: *p <0.05, **p <0.01, ***p <0.001,

****p <0.0001. **(C)** To distinguish between glucose uptake and phosphorylation steps,

two different radiolabeled glucose analogs were used for radiometric assays in LNCaP

and DU145 cells. Both 2-DG and 3-O-methylglucose (3-OMG) are taken up into cells in

the same manner as glucose. Unlike 2-DG which is trapped by hexokinase (HK)-

mediated phosphorylation and accumulated in the cells as a form of 2-DG-6P, 3-OMG is

not phosphorylated, and its intracellular levels reflect the cells' capability of Glut1-

mediated glucose uptake itself. **(D)** 2-DG 2-[1,2-^{3}H(N)] accumulation was inhibited by

the presence of BKIDC-1553-N-Me at 20 µM in LNCaP but not DU145 cell lines. 3-O-

[methyl-^{3}H] D-glucose uptake was not affected by BKIDC-1553-N-Me in LNCaP. These

data support the conclusion that BKIDC block phosphorylation but not uptake of glucose

in BKIDC-susceptible cell line LNCaP cell line. Two-tailed t tests: ****p <0.0001

1733

(A)

(B)

1734

Figure 5. The effect of HK deficiency on antiglycolytic actions of BKIDC-1553-N-

Me. (A) CRISPR-Cas9 system was used to generate HK1 and HK2 knock-out (KO)

LNCaP clones and two independently derived KO clones for HK1 and HK2 were tested.

Intracellular ATP levels, in the presence of a mitochondrial respiration blocker

oligomycin (Oligo), were used as a proxy to monitor the activity of glycolysis to generate

ATP. 2-Deoxyglucose (2-DG) and sodium oxamate (SO) were employed to block

glycolysis at different steps. SO is an inhibitor of lactate dehydrogenase thus blocking

the final step of glycolysis conversion of pyruvate into lactate. Note HK2 KO clones

displayed remarkable resistance to decreasing ATP when co-treated with Oligo and

BKIDC-1553-N-Me while they were sensitive to 2-DG and SO. One-way ANOVA

followed by Dunnett's test for multiple comparison (vs. Oligomycin alone). **(B)** Western

blot validation of HK1 and HK2 deficiency in LNCaP KO clones. GAPDH served as a

loading control.

Figure 6. Preclinical anti-prostate cancer activity of BKIDC-1553

(A) Growth Curves of Prostate Cancer Xenografts Implanted in SCID Mice. The BKIDC-resistant human prostate cancer cell line PC3 and a human patient-derived xenograft (PDX) model (LuCaP 35) were injected subcutaneously in SCID mice. Mice were treated for 4-5 weeks with 20 mg/kg BKIDC-1553 po three times a week. The LuCaP 35 tumors responded to BKIDC-1553 to a similar degree as enzalutamide (Enza) and no decrease was seen in PC3 xenografts. The data are presented as mean ± standard error of mean (SEM) (n=7-10). One-way ANOVA followed by Tukey's test for multiple comparison (LuCaP35) and two-tailed t tests (PC3): *p <0.05, **p <0.01, ns: not significant. **(B)** Immunohistochemical analysis of P-ACC in LuCaP 35 tumors. Representative images of LuCaP 35 tumors stained with an anti-P-ACC antibody are shown (at 20x magnification). Box and whisker plots show scoring analysis of anti-PACC immunoreactivities in tumors in the specified treatment. One-way ANOVA followed by Dunnett's test for multiple comparison (vs. Control: *p <0.05). The number of animals used for IHC study was: 6 for control, 8 for 1553, 6 for enzalutamide. IHC score is defined in Materials and Methods. **(C)** Dose response of LTL331R line (patient-derived in vitro model of neuroendocrine prostate cancer) to BKIDC-1553 and enzalutamide using CellTiter-Glo® 3D Cell Viability Assay. The data shown are normalized to the values obtained from cells treated with DMSO vehicle and presented as mean ± SD (n=3). One-way ANOVA followed by Dunnett's test for multiple comparison (vs. Control).

(A)

LuCaP 35 — Tumor volume (mm³) vs Weeks; Control, 1553, Enza

PC3 — Tumor volume (mm³) vs Weeks; Control, 1553; ns

(B)

Control | 1553 | Enza

P-ACC

IHC score — Control, 1553, Enza; *

(C)

Relative Proliferation vs BKIDC-1553 (µM): 0, 1.25, 5, 10, 20; ***, ****

Relative Proliferation vs Enzalutamide (µM): 0, 1.25, 5, 10