Anti-manic effect of deep brain stimulation of the ventral tegmental area in an animal model of mania induced by methamphetamine

Running title: Antimanic effects of deep brain stimulation

Roger B. Varela, PhD[1]; Suelen L. Boschen, PhD[2,3]; Nathanael Yates, PhD[1]; Tristan Houghton, BSc[1]; Charles D. Blaha, PhD[2]; Kendall H. Lee, PhD[2]; Kevin E. Bennet, PhD[2,4]; Abbas Z. Kouzani, PhD[5]; Michael Berk, PhD[6]; João Quevedo[7,8,9,10]; Samira S. Valvassori, PhD[10]; Susannah J. Tye, PhD[1,11,12,13]

[1] Functional Neuromodulation and Novel Therapeutics Laboratory, Queensland Brain Institute, Asia Pacific Centre for Neuromodulation, The University of Queensland – Brisbane, QLD, Australia.

[2] Neural Engineering Laboratories, Department of Neurologic Surgery, Mayo Clinic – Rochester, MN, United States.

[3] Applied Computational Neurophysiology and Neuromodulation Laboratory, Department of Neurologic Surgery, Mayo Clinic – Rochester, MN, United States

[4] Division of Engineering, Mayo Clinic - Rochester, MN, United States

[5] Deakin University, School of Engineering, Geelong, VIC, 3216, Australia.

[6] Deakin University, IMPACT – the Institute for Mental and Physical Health and Clinical Translation, School of Medicine, Barwon Health, Geelong, Australia

[7] Center for Interventional Psychiatry, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth Houston), Houston, TX, USA.

[8] Center of Excellence on Mood Disorders, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, USA.

[9] Translational Psychiatry Program, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, USA.

[10] Translational Psychiatry Laboratory, Graduate Program in Health Sciences, University of Southern Santa Catarina (UNESC), Criciúma, SC, Brazil.

[11] Translational Neuroscience Laboratory, Department of Psychiatry and Psychology, Mayo Clinic – Rochester, MN, United States.

[12] Department of Psychiatry, University of Minnesota, Minneapolis, MN, United States.

[13] Department of Psychiatry and Behavioral Sciences, Emory University, Atlanta, GA, United States.

*Address for correspondence: S. J. Tye, PhD, Functional Neuromodulation and Novel Therapeutics Laboratory, Queensland Brain Institute, The University of Queensland – Brisbane, QLD, Australia. Phone: #61 7 344 32493. E-mail: s.tye@uq.edu.au.

Acknowledgments

Funding for this project was provided by the National Health and Medical Research Council (NHMRC; APP1160472) (Australia) *Coordenação de Aperfeiçoamento de Pessoal de Nível Superior* (CAPES) (Brazil). MB is supported by a NHMRC Senior Principal Research Fellowship and Leadership 3 Investigator grant (GNT1156072 and 2017131). RV is supported by CREDIT: The CRE for the Development of Innovative Therapies for Psychiatric Disorders (GNT1153607). The authors gratefully acknowledge the QBI Advanced Microscopy and Histology facilities for their support and assistance in this work.

Conflict of Interest Statement

RBV, NY, and SJT were supported the Asia Pacific Centre for Neuromodulation and Medtronic (Australia). MB was supported by a Senior Principal Research Fellowship (1156072). JQ received clinical research support from LivaNova; has speaker bureau membership with Myriad Neuroscience, Janssen Pharmaceuticals, and Abbvie; is stockholder at Instituto de Neurociencias Dr. Joao Quevedo; and receives copyrights from Artmed Editora, Artmed Panamericana, and Elsevier/Academic Press. The other authors have nothing to disclose.

Abstract

Treatment of refractory bipolar disorder (BD) is extremely challenging. Deep brain stimulation (DBS) holds promise as an effective treatment intervention. However, we still understand very little about the mechanisms of DBS and its application on BD. The present study aimed to investigate the behavioural and neurochemical effects of ventral tegmental area (VTA) DBS in an animal model of mania induced by methamphetamine (m-amph). Wistar rats were given 14 days of m-amph injections, in the last day animals were submitted to 20 minutes of VTA DBS in two different patterns: intermittent low frequency stimulation (LFS) or continuous high frequency stimulation (HFS). Immediately after DBS, manic-like behaviour and nucleus accumbens (NAc) phasic dopamine (DA) release were evaluated separately through open-field test and fast-scan cyclic voltammetry. Levels of NAc dopaminergic markers were evaluated by immunohistochemistry. M-amph induced hyperlocomotion in the animals and both DBS parameters reversed this alteration. M-amph increased DA reuptake time post-sham compared to baseline levels, and both LFS and HFS were able to block this alteration. LFS was also able to reduce phasic DA release when compared to baseline. LFS was able to increase dopamine transporter (DAT) expression in the NAc. These results demonstrate that both VTA LFS and HFS DBS exert anti-manic effects and modulation of DA dynamics in the NAc. More specifically the increase in DA reuptake driven by increased DAT expression may serve as a potential mechanism by which VTA DBS exerts its anti-manic effects.

Keywords: Bipolar Disorder; Mania; Depression; Methamphetamine; Deep Brain Stimulation; Dopamine, Psychiatry.

1 Introduction

Bipolar disorder (BD) is a major psychiatric disorder characterized by intermittent and interspersed episodes of mania/hypomania and depression. Although manic episodes are the clinical hallmark for this disorder, BD patients typically spend three times as long with depression as mania/hypomania (1). In fact, bipolar depression is considerably more persistent than non-bipolar depression (2). Lithium, the gold standard mood stabilizer, is used for treatment of bipolar depression (3). However, about two thirds of BD patients may not respond to mood stabilizer treatment and, refractory bipolar treatment is even more challenging due the potential risk of mood shift (4, 5).

The treatment of psychiatric disorders is challenging, as it involves dysfunctions of multiple brain regions and often presents overlapping symptoms and pathologies. Consequently, there is an increasing focus on how the dysfunction of specific brain circuits can lead to specific transdiagnostic psychiatric symptoms (6). One of the more important and studied brain circuits involved in BD is the mesoaccumbens dopaminergic pathway with dopamine (DA)-containing neuronal cells in the ventral tegmental area (VTA) that directly innervate the nucleus accumbens (NAc) (7, 8). Dysregulation of mesoaccumbens dopaminergic transmission in the NAc is often implicated in disorders of motivated behaviour, including BD (9).

For those mood disorders patients who do not respond to any available antidepressant treatments, deep brain stimulation (DBS) holds promise as an effective treatment intervention (10). Recently Drobisz and Damborská (11) compiled evidence reports demonstrating that stimulation of different brain regions can exert antidepressant effects in treatment-resistant depression patients. Although suggesting promising results, DBS for treatment-resistant depression carries many issues that need to be addressed. Very little is understood about the mechanisms of DBS for treatment of depression and moreover, the application of DBS on BD is even less described, as few clinical DBS studies have included BD patients (12). The main concern for the application of DBS on bipolar patients is the observation of treatment associated adverse events in isolated cases (e.g., changes in mood, induced hypomania state). However, the studies about its applicability are still controversial as noted in the current literature, highlighting the need for further studies on the effects of different patterns of stimulation in this condition (13).

It is known that the electrical stimulation parameters have an important role in the mechanism of action of DBS. High frequency stimulation (HFS) (130–185 Hz) induces neuronal depolarization and generates action potentials by opening local voltage-gated sodium channels (14, 15). However, HFS may have a more pronounced inhibitory effect than low frequency stimulation (LFS) under 100 Hz (16). This is thought to occur due to a 'depolarization block', where repeated stimulations inhibit sodium ion driven action potentials (17). Also, HFS could have different neural activity effects depending on targeted excitatory or inhibitory neurons (18). Alternatively, Friedman and colleagues (19) designed a programmed LFS based on the difference of VTA neuronal firing patterns in a genetic rat model of depression and control animals. This programmed LFS consists of bursts containing numerous spikes, mimicking and resetting the natural VTA firing configuration, thereby potentially alleviating depressive symptoms in rats (16, 20).

Animal models are important tools to investigate DBS therapeutic mechanisms and applications as they provide a means to control variables that are not easily possible in human studies. Under these conditions, new targets and patterns of stimulations can be tested to refine the functionality of this

interventive technique (11, 21). Methamphetamine (m-amph) administration in rats is an animal model of human mania with face, construct and predictive validities, since it can induce manic-like behaviours (including hyperactivity) and responds to mood stabilizers treatment (22). M-amph increases extracellular levels of DA through multiple mechanisms, mindful that dopamine is a critical element in bipolar pathophysiology (23) and the NAc is particularly affected due to its high concentration of DA terminals (24, 25).

There are several potential mechanisms through which DBS of the mesoaccumbens dopaminergic pathway could correct aberrant DA signaling in the NAc that is characteristic of mania. In line with the depolarization block hypothesis of DBS, inhibiting DA release from the VTA to the NAc should reduce the synaptic concentrations of DA in the NAc induced by m-amph. Alternatively, DBS applied to this region could also reset the natural firing configuration, modulating aberrant activity in the mesolimbic DA circuit, normalizing synaptic DA concentrations in the NAc and consequent abnormal behavior. Therefore, the present study aimed to investigate the effects of VTA DBS (both LFS and HFS) in an animal model of mania induced by m-amph to better describe the effects of DBS on mania states, and potential application for BD treatment.

2 Material and methods

2.1 Animals

The subjects were adult male Wistar rats (N=15-20/group, weighting 250–350 g) obtained from both the Mayo Clinic or The University of Queensland breeding colonies. The animals were housed two or five per cage (for behaviour and DA quantification, respectively), with food and water available *ad libitum* and maintained on a 12-h light/dark cycle (lights on at 6:00 a.m.) at a temperature of 22±1 °C. All experimental procedures were performed in accordance with, and with the approval of either the Mayo Clinic or The University of Queensland Institutional Animal Care and Use Committees, respectively. All experiments were performed at the same time during the day to mitigate circadian variations.

2.2 Experiment 1: Effects of VTA DBS on mania-like behaviour

After a week of acclimatization, rats received 14 days of either saline (NaCl 0.9%), or m-amph (1mg/kg) treatment via intraperitoneal (I.P.) injections (Sigma-Aldrich, USA). On day 10, animals were anesthetized with isoflurane (1–3%) for stereotactic surgery. A 1.5–2 cm incision was made, and a small hole was broached in the skull at the site corresponding to the target. Twisted bipolar stainless steel electrodes (10 mm long, 0.075 mm in diameter; Plastics One, Virginia, USA) were unilaterally implanted into the VTA (from bregma: antero-posterior (AP) -6.8mm, medial-lateral (ML) +0.6mm and, dorsal-ventral (DV) −8.6 mm) (26). Stimulating electrodes and four screws were mounted to the skull with dental cement. A control group was not subjected to stereotactic surgery and had no electrode implanted. Unilateral implantation was applied since a previous study reported that unilateral stimulation of this brain region was sufficient for clinical outcomes (27).

At the 15[th] day, animals received a single I.P. injection of m-amph or saline (challenge), 1 hour and 40 minutes after the injection the animals were submitted to 20 minutes of DBS in two different patterns: Intermittent LFS (Frequency: 10Hz; 2ms pulse width; 2 burst/sec; 300µA; monophasic) or continuous HFS (130Hz; 2ms pulse width; 200µA; monophasic) controlled through Iso-Flex/Master-8 system (AMPI©, Israel)

(Figure 1A) (n=12-16). Stimulation parameters were based on Gazit et al. (16), mimicking physiological pattern of VTA firing and Kouzani et al. (28), current clinical protocol. Sham animals were submitted to the same surgical procedure, had the electrode implanted and were attached to the stimulation device for the same period as the active DBS groups, but stimulation was turned off. Immediately after the DBS session, animals' locomotor activity and risk-taking behaviour were evaluated through the open-field test (OFT). For this test, animals were placed in an arena (50x50cm) surrounded by 50 cm high walls. A video camera on the top of the arena was connected to a computer for recording behavioural parameters. The animals were carefully put into the central area and then left to explore the arena for 20 min. Total distance travelled and the time spent in the central area of the arena were recorded through the EthoVision XT behavioral software (Noldus©, Netherlands). Experiment 1 timeline is depicted in Figure 1B. Timepoints for m-amph injection and OFT were based on the original animal model study considering pharmacokinetics of m-amph (22), timepoint and duration of DBS was based on Gazit et al. study (16).

Figure 1: (A) Intermittent low frequency stimulation (LFS): 10Hz stimulation divided in 2 bursts per second, each burst composed of 5 spikes with an inter-spike interval of 80ms and a pause of 180 ms after each burst, according to Gazit et al., 2015. High frequency stimulation (HFS): Continuous stimulation at 130Hz; (B) Experiment 1: Animals received daily intraperitoneal injections of methamphetamine (m-amph) or Saline (Sal) for 14 days. On the tenth day animals were submitted to stereotactic surgery for unilateral implantation of a deep brain stimulation (DBS) electrode in the ventral tegmental area. On the fifteenth day, animals received a single injection of m-amph or Sal and were submitted to 20 minutes of LFS or HFS DBS prior to open field test (OFT); (C) Experiment 2: Animals were submitted to 14 daily injections of m-amph or Sal. On the fifteenth day animals were submitted to fast-scan cyclic voltammetry surgery. After implantation of electrodes and optimization, animals received a single injection of m-amph or Sal and had five evoked dopamine (DA) responses recorded before 20 minutes DBS session. Five evoked DA responses were recorded post-DBS session and after m-amph challenge. After FSCV recordings animals were euthanised and brains were removed.

2.3 Experiment 2: Effects of DBS on VTA-evoked phasic DA release in the NAc

This experiment was designed to investigate the biological mechanisms underlying DBS effects in significant conditions observed in experiment 1. After a week of acclimatization, rats received 14 days of either saline or m-amph treatment. On the 15th day, animals were submitted to fast-scan cyclic voltammetry (FSCV) analysis for evaluation of VTA-evoked phasic DA release in the NAc. Prior to FSCV surgery, animals were anaesthetised with a 1.5 mg/kg intraperitoneal urethane injection. An incision was made in the midline of the skull to expose bregma. Using bregma as reference, a 1 mm diameter hole was drilled

into the skull at the site for the DA recording electrode. The target for the recording electrode was the NAc (AP +1.2, ML +1.5, DV –6.5-8.0 from bregma). The VTA stimulation electrode was implanted according to coordinates described above. A hole was also made for the reference electrode in the opposite hemisphere to the stimulating and recording electrodes. A wireless instantaneous neurotransmitter concentration sensing (WINCS) Harmoni system (Mayo Clinic, USA) was used to generate a voltage-current plot. The location of the recording and stimulating electrodes were optimised by applying stimulation of the VTA (DA-evoked stimulation: 120 pulses; 60Hz; 300-500µA) and recording via the NAc-implanted carbon fiber recording electrode when located at progressively ventral DV coordinates. The recording electrode was lowered to improve the oxidation/reduction signal strength. Adjustments were made until there was a strong signal. Once recording and stimulation sites were optimized, animals received an injection of saline or m-amph (according to pre-treatment).

75 minutes after m-amph/saline injection, five baseline VTA-evoked DA responses were recorded every 5 minutes, period named pre-DBS. Animals were then submitted to VTA DBS (HFS or LFS) or Sham (no stimulation) for 20 minutes (n=4-6). During this time, evoked DA responses were not recorded. After Sham or DBS stimulation, VTA-evoked DA responses were recorded every 5 minutes for 25 minutes, period named post-DBS. After post-DBS recordings, all the animals were submitted to an m-amph challenge (1mg/kg) and 5 VTA-evoked DA responses were recorded for confirmation of DA dynamics, period named post-challenge (Figure 1C). The DA peak was determined by averaging the one second time (ten scans) with highest concentration after stimulation. DA reuptake was calculated through T1/2 method which consists in the time for DA concentration to drop to half after peak. The DA recording protocol is demonstrated in the scheme below.

2.4 Euthanasia and brain dissection

All the animals were euthanized with I.P. overdose of pentobarbital (150 mg/kg body weight) after the behavior or FSCV protocols. Experiment 2 animals had their brains removed and frozen by placing them on dry ice, followed by storage at − 80 °C for posterior histological and immunohistochemical analysis.

2.5 Histological analysis

Frozen tissue sections (10µm) were cut with cryostat (Cryostar NX70, ThermoScientific©, USA), mounted on ÜberFrost Slides (InstrumeC©, Australia) and immediately fixed using 4% PFA for 10 min. VTA sections were submitted to Nissl staining through cresyl violet for location of the stimulating electrodes. For immunostaining of NAc sections, the tissue section was encircled by using the wax PAP-pen (Daido Sangyo©, Japan) and permeabilized with 0.1% Revealit antigen recovery solution (0.1M sodium citrate, citric acid added until pH reaches 6, 0.1M SDS) for 30 min at 40°C using a shaker incubator. Subsequently, sections were washed in PBS for 5 min, and blocking solution for 30 min (0.5 % BSA in 0.05 % saponin, 0.05% sodium azide in 0.1M PBS). Samples were then incubated with the primary antibodies for 4 nights at room temperature protected from light in a humidity chamber. On the fourth day, the sections were washed three times with PBS for 15 min each followed with blocking solution for 5 min. Thereafter, samples were incubated overnight with the secondary antibodies at room temperature. The next day, the sections were washed three times with PBS for 15 min and then coverslipped by using fluorescence mounting

medium (Dako©, Denmark). Both primary and secondary antibodies were diluted in blocking solution. The primary antibodies used for immunohistochemical analysis included: Mouse monoclonal antibody against Tyrosine Hydroxylase (ImmunoStar Cat# 22941, RRID:AB_572268), rabbit monoclonal antibody against NeuN (Cell Signaling Technology Cat# 24307, RRID:AB_2651140), rat monoclonal antibody against Dopamine Transporter (Millipore Cat# MAB369, RRID:AB_2190413). Goat anti-Mouse IgG Alexa Fluor 488 (Thermo Fisher Scientific Cat# A-11001, RRID:AB_2534069), Goat anti-Rat IgG Alexa Fluor 647 (Molecular Probes Cat# A-21247, RRID:AB_141778) and Donkey anti-Rabbit IgG Alexa Fluor 555 (Molecular Probes Cat# A-31572, RRID:AB_162543) were used as secondary antibodies. Protein expression in the NAc was analysed by semi-quantitative intensity analysis using ImageJ software.

2.6 Microscopy and image analysis

All tissue imaging was captured using Zeiss Axio Imager 2 Microscope (https://www.zeiss.com/microscopy/us/products/light-microscopes/axio-imager-2-for-biology.html, RRID: SCR_018876) using a single field of view from the structure of interest, captured using a 20x objective lens, using LED illumination consistent exposure and light settings for all sections belong to a group. The images were then analysed using semi-quantitative immunofluorescent intensity analysis using Fiji software (http://fiji.sc, RRID:SCR_002285) (29) threshold analysis. Identical threshold settings were used between images to ensure consistency. Because DAT is strongly positive in blood vessels, we used a custom written macro to remove the blood vessels from our analysis in these stains.

2.7 Statistical analysis

Differences among the experimental groups were determined by two-way analysis of variance (ANOVA) followed by Tukey's post hoc test for OFT. Intragroup differences were determined by one-way ANOVA followed by Tukey's post hoc test for FSCV recordings and immunohistochemical analysis. In all comparisons, statistical significance was set at $p<0.05$. Statistical analysis and graphs were performed using Graphpad Prism software, V9.0 (http://www.graphpad.com/, RRID:SCR_002798).

3. Results

3.1 Open-field test

As demonstrated in Figure 2, no alteration was observed among the saline pre-treated groups in the OFT. M-amph chronic administration increased locomotor activity when compared to the Saline+control group. Although electrode implantation in the VTA was able to reverse this m-amph-induced hyperlocomotion, LFS and HFS led to a more significant reduction (Figure 2A). Time spent in the central area of the arena was also assessed as a risk-taking behavior parameter. Although this parameter shows the same trend of locomotor activity, no statistical significance was observed (Figure 2B). Data from the two-way ANOVA for m-amph pre-treatment [Distance: $F_{(1.95)}=5.692$, $p=0.019$; Risk: $F_{(1.95)}=7.554$, $p=0.007$]; for DBS [Distance: $F_{(3.95)}=8.421$, $p<0.0001$; Risk: $F_{(3.95)}=1.768$, $p=0.1586$] and for interaction between m-amph vs DBS [Distance: $F_{(3.95)}=2.691$, $p=0.0506$; Risk: $F_{(3.95)}=0.7622$, $p=0.5180$].

Figure 2: Effects of low frequency stimulation (LFS) and high frequency stimulation (HFS) on travelled distance (A) and time spent in center (B) in the open-field test of rats submitted to the animal model of mania induced by methamphetamine (m-amph). Bars represent means and error bars represent standard error according to two-way ANOVA. *=$p<0.05$ compared to de Saline+Ctrl group; ##=$p<0.01$ compared to the m-amph+Ctrl group; ##=$p<0.001$ compared to the m-amph+Ctrl group according to two-way ANOVA followed by Tukey's post-hoc test.

3.2 VTA-evoked phasic DA release

As depicted in Figure 3, stimulating electrodes were implanted in the VTA region for all animals submitted to the FSCV, except for one. Since evoked DA responses were detected in all animals, the misplaced electrode location could have been affected by the brain section used for histological analysis.

Figure 3: Left: Schematic illustration of coronal rat brain slices at the level of the VTA region (A: interaural 2.16mm; B: interaural 2.88mm). Image modified from Paxinos & Watson "The Rat Brain in Stereotactic Coordinates (2007). Right: Whole brain mirrored image of VTA DBS animal that was sliced to the site of electrode implantation. Electrode tract mark is visible in this slice.

Results from FSCV recordings are shown in Figure 4. Figure 4A, 4F, 4K and 4P demonstrate the dynamics of DA concentrations in the NAc after VTA-induced phasic release (stimulation happened at 5 seconds). Figure 4B, 4G, 4L and 4Q represents the DA peak concentration reached after VTA-induced

phasic release and, Figure 4C, 4H, 4M and 4R represent DA reuptake time. Figures 4D, 4E, 4I, 4J, 4N, 4O, 4S and 4T show pre- and post-sham/DBS representative colour plots.

As expected for Saline+ Sham animals, there was no difference in peak DA concentrations or reuptake on post-sham compared to baseline. However, m-amph challenge increased the DA peak concentration and reuptake time in these animals (Figure 4A, 4B and 4C). Data from the one-way ANOVA for Sal+Sham group [Peak: F(2.85)=4.331, p=0.0162; Reuptake: F(2.82)= 5.502, p=0.0057]. Neither sham period nor m-amph challenge altered the DA peak concentrations when compared to baseline. In addition, DA reuptake time was increased post-sham and post-challenge (Figure 4D, 4E and 4F). Data from one-way ANOVA for m-amph+Sham group [Peak: F(2.57)= 2.341, p=0.1054; Reuptake: F(2.57)=8.743, p=0.0006]. HFS had no effect on DA peak concentrations, like sham stimulation. However, HFS was able to block the increase in reuptake time post-stimulation which was observed in the sham group. Reuptake time was increased after m-amph challenge (Figure 4G, 4H and 4I). Data from one-way ANOVA for m-amph+HFS group [Peak: F(2.57)= 3.825, p=0.0276; Reuptake: F(2.57)= 12.28, p< 0.0001]. In contrast, LFS decreased the DA peak concentrations, which returned to baseline levels after m-amph challenge. LFS was able to block the increased DA reuptake time not only in the post-stimulation period but also during the post-challenge period (Figure 4J, 4K and 4L). Data from one-way ANOVA for m-amph+LFS group [Peak: F(2.72)= 7.569, p= 0.0010; Reuptake: F(2.71)= 2.004, p= 0.1423].

Figure 4: Effects of low frequency stimulation (LFS) and high frequency stimulation (HFS) on nucleus accumbens phasic dopamine (DA) release (A, F, K and P), DA peak concentration (B, G, L and Q), and DA reuptake time (C, H, M and R) of rats submitted to the animal model of mania induced by methamphetamine (m-amph). Bars and lines represent the mean and error bars represent standard errors according to one-way ANOVA. *=p<0.05; ***=p<0.01 and ****=p<0.001 compared to the baseline according to two-way ANOVA

followed by Tukey's post-hoc test. Representative pre- and post-DBS/Sham color plots are represented in figures D, E, I, J, N, O, S and T.

3.3 Immunohistochemistry

Figure 5A demonstrates the expression of DAT, NeuN and tyrosine hydroxylase (TH) in the NAc core and shell, represented as arbitrary intensity units. Representative images from IHC are demonstrated in Figure 5B. LFS appeared to increase DAT expression in both, NAc core and shell. No difference was observed in any of the other groups. Data from the one-way ANOVA for NAc core $F_{(3.16)}=4.485$, $p=0.0138$ and shell $F_{(3.16)}=4.717$, $p=0.0152$.

4. Discussion

Chronic m-amph injection increased locomotor activity in the OFT when compared to the control group. Hyperlocomotion is considered one of the most objective manic-like behaviors assessed in animals, akin to states of increased activity, energy, or drive in humans (30). Administration of psychostimulants, such as amphetamines, are known for inducing manic-like behaviors, and are considered a reliable animal model of mania, being based on monoaminergic dysregulation which responds to mood-stabilizer treatments, supporting face, construct and predictive validities (22). M-amph treatment increases time spent in the center of the OFT arena and it is considered a parameter of risk-taking behavior and decreased anxiety in rodents, which may be translated to poor judgment associated with mania (31, 32). In the present study, m-amph treatment appeared to increase risk-taking behavior, however this increase was not statistically significant.

Figure 5: Effects of high frequency stimulation (HFS) and low frequency stimulation (LFS) on nucleus accumbens (NAc) DAT, NeuN and TH expression in rats submitted to the animal model of mania induced by methamphetamine (m-amph). **(A)** Protein immunofluorescence presented in arbitrary intensity units, bars represent means and errors bars represent standard errors according to one-way ANOVA. *=p<0.05 compared to the Saline+Sham; #=p<0.05 compared to the m-amph+Sham group according to Tukey's post-hoc test. **(B)** First column represents a sample image NAc region staining, second to fifth columns shown representative images detailing DAT (far red), NeuN (red) and TH (green) staining for each experimental group.

Electrode implantation alone was able to reduce hyperactivity induced by chronic m-amph, although it did not alter locomotion in saline-treated animals. The positive effects of sham DBS are described in several depression clinical trials, although the effects of DBS are more substantial (33, 34). Perez-Caballero and colleagues (35) described that sham-DBS antidepressant effect is related to increased expression of inflammatory mediators and activation of several short-term repair mechanisms in the region of electrode implantation which can last for up to 2 weeks after electrode implantation and is blocked by anti-inflammatory drugs. Although not described in mania models so far, activation of inflammatory mediators and repair mechanisms in the VTA might contribute to anti-manic-like effects observed in the sham group. Since electrode implantation did not change behavior in saline animals it is unlikely that reduced locomotion observed in Sham/DBS groups are due VTA lesioning-induced locomotion impairment in m-amph group.

Both LFS and HFS were able to completely reverse m-amph-induced hyperlocomotion. Studies describing the effects of DBS on m-amph-induced hyperlocomotion are scant. However, Perez et al. (36) demonstrated that hippocampal HFS DBS reversed hyperactivity induced by acute m-amph administration in sensitized animals in a model of schizophrenia. DBS-induced inhibition of the hippocampus indirectly decreases VTA activity and dopaminergic signalling in the NAc, normalizing the behavioral responses (36). Those results are consistent with the present study, suggesting that direct inhibition of VTA through LFS and HFS DBS normalize m-amph induced hyperlocomotion.

We further investigated the role of phasic DA release on the behavioral effects of VTA LFS and HFS DBS in a different set of animals. For the saline-treated group, phasic DA release and reuptake time did not change post-sham when compared to baseline. Although not related to the first experiment protocol, experiment 2 animals received a m-amph challenge after the post-sham period to confirm DA responses. As previously described by other authors, an acute m-amph injection increased both evoked DA release and reuptake time (37). M-amph increases DA release through different mechanisms such as disruption of the dopamine transporter (DAT) (38), and increased DA synthesis after sub-toxic doses (<2.5mg/kg) (39, 40).

Like the saline group, m-amph-treated animals did not show any significant difference between evoked DA post-sham to baseline. In fact, a trend for decreased evoked DA was noticed post-sham, however it can be explained by the time-point difference of these recordings in relation to the m-amph injection. Hedges and colleagues (41) described that evoked DA reaches a peak 5-10 min after m-amph injection, gradually decreasing before reaching baseline levels around 60 minutes after injection. Thus, the apparent decrease in the present study could be explained by the fact that baseline was recorded 75 minutes after m-amph injection, and the post-sham was recorded 105 minutes after m-amph injection. Interestingly, time to DA reuptake was increased after sham when compared to baseline in this group. To the best of our knowledge this is the first study showing late effects of a m-amph challenge on phasic DA reuptake in pre-sensitized animals. The mechanisms underlying this effect remain unknown, however a previous study demonstrated that high extracellular concentrations of DA after phasic bursts can mediate internalization of DATs. As a result, upon cessation of phasic bursts, extrasynaptic DA clearance is slower, thus accumulating in the extracellular space (42). This rebound mechanism could explain, at least in part, the late effect of m-amph on phasic DA dynamics observed after the sham stimulation period.

HFS and LFS had similar effects on phasic DA dynamics. Although only LFS was able to decrease phasic DA release when compared to baseline, both patterns of stimulation prevented the slower DA reuptake observed in the sham group. The effects of intermittent LFS on DA release remain unknown, however this programmed LFS was designed to mimic the VTA firing configuration of a control rat, being able to normalize BDNF levels and alleviate depressive-like behavior in an animal model of depression (16, 43). However, it is interesting to note that LFS was able to prevent the slow-reuptake induced by the last m-amph challenge, in addition to increased NAc DAT expression, suggesting that LFS effects on phasic DA may be driven by DAT upregulation. In this regard, Miller and colleagues (44) demonstrated that VTA dopaminergic network connectivity is regulatable via the DAT but not a D2 receptor dependent mechanism, suggesting a master regulator role of this transporter. There is a genetic report suggesting linkage disequilibrium between bipolar disorder and the dopamine transporter (45). A PET neuroimaging study showed lower caudate transporter availability in bipolar disorder than in controls (46). Zhao et al. (47) demonstrated that chronic HFS can increase DAT levels, playing an important role on dopaminergic system modulation in an animal model of Tourette syndrome. Interestingly, a case report also demonstrated that 1 year of NAc HFS was able to increase striatal levels of DAT in an m-amph use disorder patient (48). Together with literature data, the present study suggests that DAT upregulation may act to re-establish normal DA dynamics, playing a key role on the anti-manic effects of DBS.

It is important to note that some limitations of the present study should be addressed in future investigations. The OFT test was conducted immediately after the VTA DBS session. Simultaneous DBS stimulation and OFT performance may provide a better temporal resolution of the behavioral effects of LFS and HFS VTA DBS. Animals submitted to the experiment 2 design were under urethane anaesthesia during phasic DA release recordings, which could affect the comparison of behavioral and electrochemical analysis. FSCV analysis in awake animals (even during OFT) might better reflect DA dynamics and mania-like behavior. FSCV is an electrochemical technique developed to measure rapid changes in neurotransmitters levels, and since it uses the method of background subtraction it can only measure phasic DA release but not tonic levels (49). Evaluation of tonic DA levels (through micro dialysis or multiple-cyclic square wave voltammetry (50)), and DAT activity would help to better understand the role of DA dynamics underlying the potential anti-manic effects of LFS and HFS VTA DBS.

5. Conclusion

In summary, this study demonstrated that both LFS and HFS acute DBS of the VTA region exerts anti-manic effects in animals with m-amph induced mania. In addition, LFS and HFS modulated DA dynamics (more specifically DA reuptake) and DAT expression in the NAc, suggest a potential mechanism whereby VTA DBS may exert its anti-manic effects. The next step is to better understand the real-time effects of DBS in manic-like behavior, as well as other aspects of DA signalling. However, these results point to the potential application of neuromodulation approaches to model and clinically modulate manic-like behaviours.

References

1.   Baldessarini RJ, Salvatore P, Khalsa HM, Gebre-Medhin P, Imaz H, González-Pinto A, et al. Morbidity in 303 first-episode bipolar I disorder patients. Bipolar Disord. 2010;12(3):264-70.
2.   Merikangas KR, Jin R, He JP, Kessler RC, Lee S, Sampson NA, et al. Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative. Arch Gen Psychiatry. 2011;68(3):241-51.
3.   Gitlin MJ. Antidepressants in bipolar depression: an enduring controversy. Int J Bipolar Disord. 2018;6(1):25.
4.   Fountoulakis KN. Refractoriness in bipolar disorder: definitions and evidence-based treatment. CNS Neurosci Ther. 2012;18(3):227-37.
5.   Schoeyen HK, Kessler U, Andreassen OA, Auestad BH, Bergsholm P, Malt UF, et al. Treatment-resistant bipolar depression: a randomized controlled trial of electroconvulsive therapy versus algorithm-based pharmacological treatment. Am J Psychiatry. 2015;172(1):41-51.
6.   Ressler KJ, Mayberg HS. Targeting abnormal neural circuits in mood and anxiety disorders: from the laboratory to the clinic. Nat Neurosci. 2007;10(9):1116-24.
7.   Fulton S, Pissios P, Manchon RP, Stiles L, Frank L, Pothos EN, et al. Leptin regulation of the mesoaccumbens dopamine pathway. Neuron. 2006;51(6):811-22.
8.   Rodríguez-López C, Clascá F, Prensa L. The Mesoaccumbens Pathway: A Retrograde Labeling and Single-Cell Axon Tracing Analysis in the Mouse. Front Neuroanat. 2017;11:25.
9.   Coenen VA, Panksepp J, Hurwitz TA, Urbach H, Mädler B. Human medial forebrain bundle (MFB) and anterior thalamic radiation (ATR): imaging of two major subcortical pathways and the dynamic balance of opposite affects in understanding depression. J Neuropsychiatry Clin Neurosci. 2012;24(2):223-36.
10.   Holtzheimer PE, 3rd, Mayberg HS. Deep brain stimulation for treatment-resistant depression. Am J Psychiatry. 2010;167(12):1437-44.
11.   Drobisz D, Damborská A. Deep brain stimulation targets for treating depression. Behav Brain Res. 2019;359:266-73.
12.   Gippert SM, Switala C, Bewernick BH, Kayser S, Bräuer A, Coenen VA, et al. Deep brain stimulation for bipolar disorder-review and outlook. CNS Spectr. 2017;22(3):254-7.
13.   Mallet L, Mesnage V, Houeto JL, Pelissolo A, Yelnik J, Behar C, et al. Compulsions, Parkinson's disease, and stimulation. Lancet. 2002;360(9342):1302-4.
14.   Breit S, Schulz JB, Benabid AL. Deep brain stimulation. Cell Tissue Res. 2004;318(1):275-88.
15.   Galati S, Mazzone P, Fedele E, Pisani A, Peppe A, Pierantozzi M, et al. Biochemical and electrophysiological changes of substantia nigra pars reticulata driven by subthalamic stimulation in patients with Parkinson's disease. Eur J Neurosci. 2006;23(11):2923-8.
16.   Gazit T, Friedman A, Lax E, Samuel M, Zahut R, Katz M, et al. Programmed deep brain stimulation synchronizes VTA gamma band field potential and alleviates depressive-like behavior in rats. Neuropharmacology. 2015;91:135-41.
17.   Grace AA, Bunney BS, Moore H, Todd CL. Dopamine-cell depolarization block as a model for the therapeutic actions of antipsychotic drugs. Trends Neurosci. 1997;20(1):31-7.
18.   Aum DJ, Tierney TS. Deep brain stimulation: foundations and future trends. Front Biosci (Landmark Ed). 2018;23:162-82.
19.   Friedman A, Deri I, Friedman Y, Dremencov E, Goutkin S, Kravchinsky E, et al. Decoding of dopaminergic mesolimbic activity and depressive behavior. J Mol Neurosci. 2007;32(1):72-9.
20.   Friedman A, Frankel M, Flaumenhaft Y, Merenlender A, Pinhasov A, Feder Y, et al. Programmed acute electrical stimulation of ventral tegmental area alleviates depressive-like behavior. Neuropsychopharmacology. 2009;34(4):1057-66.
21.   Hamani C, Nóbrega JN. Deep brain stimulation in clinical trials and animal models of depression. Eur J Neurosci. 2010;32(7):1109-17.

22.   Feier G, Valvassori SS, Varela RB, Resende WR, Bavaresco DV, Morais MO, et al. Lithium and valproate modulate energy metabolism in an animal model of mania induced by methamphetamine. Pharmacol Biochem Behav. 2013;103(3):589-96.

23.   Berk M, Dodd S, Kauer-Sant'anna M, Malhi GS, Bourin M, Kapczinski F, et al. Dopamine dysregulation syndrome: implications for a dopamine hypothesis of bipolar disorder. Acta Psychiatr Scand Suppl. 2007(434):41-9.

24.   Gerasimov MR, Ashby CR, Jr., Gardner EL, Mills MJ, Brodie JD, Dewey SL. Gamma-vinyl GABA inhibits methamphetamine, heroin, or ethanol-induced increases in nucleus accumbens dopamine. Synapse. 1999;34(1):11-9.

25.   Kuhn DM, Angoa-Pérez M, Thomas DM. Nucleus accumbens invulnerability to methamphetamine neurotoxicity. Ilar j. 2011;52(3):352-65.

26.   Paxinos G, Watson C. The Rat Brain in Stereotaxic Coordinates. 7th ed. Oxford: Elsevier; 2006 2nd November 2006.

27.   Germano IM, Gracies JM, Weisz DJ, Tse W, Koller WC, Olanow CW. Unilateral stimulation of the subthalamic nucleus in Parkinson disease: a double-blind 12-month evaluation study. J Neurosurg. 2004;101(1):36-42.

28.   Kouzani AZ, Kale RP, Zarate-Garza PP, Berk M, Walder K, Tye SJ. Validation of a Portable Low-Power Deep Brain Stimulation Device Through Anxiolytic Effects in a Laboratory Rat Model. IEEE Trans Neural Syst Rehabil Eng. 2017;25(9):1365-74.

29.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform for biological-image analysis. Nature Methods. 2012;9(7):676-82.

30.   Sharma AN, Fries GR, Galvez JF, Valvassori SS, Soares JC, Carvalho AF, et al. Modeling mania in preclinical settings: A comprehensive review. Prog Neuropsychopharmacol Biol Psychiatry. 2016;66:22-34.

31.   Einat H. Different behaviors and different strains: potential new ways to model bipolar disorder. Neurosci Biobehav Rev. 2007;31(6):850-7.

32.   Menegas S, Dal-Pont GC, Cararo JH, Varela RB, Aguiar-Geraldo JM, Possamai-Della T, et al. Efficacy of folic acid as an adjunct to lithium therapy on manic-like behaviors, oxidative stress and inflammatory parameters in an animal model of mania. Metab Brain Dis. 2020;35(2):413-25.

33.   Holtzheimer PE, Husain MM, Lisanby SH, Taylor SF, Whitworth LA, McClintock S, et al. Subcallosal cingulate deep brain stimulation for treatment-resistant depression: a multisite, randomised, sham-controlled trial. Lancet Psychiatry. 2017;4(11):839-49.

34.   Coenen VA, Bewernick BH, Kayser S, Kilian H, Boström J, Greschus S, et al. Superolateral medial forebrain bundle deep brain stimulation in major depression: a gateway trial. Neuropsychopharmacology. 2019;44(7):1224-32.

35.   Perez-Caballero L, Pérez-Egea R, Romero-Grimaldi C, Puigdemont D, Molet J, Caso JR, et al. Early responses to deep brain stimulation in depression are modulated by anti-inflammatory drugs. Mol Psychiatry. 2014;19(5):607-14.

36.   Perez SM, Shah A, Asher A, Lodge DJ. Hippocampal deep brain stimulation reverses physiological and behavioural deficits in a rodent model of schizophrenia. Int J Neuropsychopharmacol. 2013;16(6):1331-9.

37.   Jang EY, Yang CH, Hedges DM, Kim SP, Lee JY, Ekins TG, et al. The role of reactive oxygen species in methamphetamine self-administration and dopamine release in the nucleus accumbens. Addict Biol. 2017;22(5):1304-15.

38.   Xie Z, Miller GM. A receptor mechanism for methamphetamine action in dopamine transporter regulation in brain. J Pharmacol Exp Ther. 2009;330(1):316-25.

39.   Takeichi T, Hori O, Hattori T, Kiryu K, Zuka M, Kitamura O. Pre-administration of low-dose methamphetamine enhances movement and neural activity after high-dose methamphetamine administration in the striatum. Neurosci Lett. 2019;703:119-24.

40.	Shepard JD, Chuang DT, Shaham Y, Morales M. Effect of methamphetamine self-administration on tyrosine hydroxylase and dopamine transporter levels in mesolimbic and nigrostriatal dopamine pathways of the rat. Psychopharmacology (Berl). 2006;185(4):505-13.

41.	Hedges DM, Obray JD, Yorgason JT, Jang EY, Weerasekara VK, Uys JD, et al. Methamphetamine Induces Dopamine Release in the Nucleus Accumbens Through a Sigma Receptor-Mediated Pathway. Neuropsychopharmacology. 2018;43(6):1405-14.

42.	Lohani S, Martig AK, Underhill SM, DeFrancesco A, Roberts MJ, Rinaman L, et al. Burst activation of dopamine neurons produces prolonged post-burst availability of actively released dopamine. Neuropsychopharmacology. 2018;43(10):2083-92.

43.	Friedman A, Lax E, Abraham L, Tischler H, Yadid G. Abnormality of VTA local field potential in an animal model of depression was restored by patterned DBS treatment. Eur Neuropsychopharmacol. 2012;22(1):64-71.

44.	Miller DR, Guenther DT, Maurer AP, Hansen CA, Zalesky A, Khoshbouei H. Dopamine Transporter Is a Master Regulator of Dopaminergic Neural Network Connectivity. J Neurosci. 2021;41(25):5453-70.

45.	Greenwood TA, Alexander M, Keck PE, McElroy S, Sadovnick AD, Remick RA, et al. Evidence for linkage disequilibrium between the dopamine transporter and bipolar disorder. Am J Med Genet. 2001;105(2):145-51.

46.	Anand A, Barkay G, Dzemidzic M, Albrecht D, Karne H, Zheng QH, et al. Striatal dopamine transporter availability in unmedicated bipolar disorder. Bipolar Disord. 2011;13(4):406-13.

47.	Zhao M, Wang X, Deng J, Guan Y, Zhou J, Li T, et al. Globus Pallidus Internus Electric High-Frequency Stimulation Modulates Dopaminergic Activity in the Striatum of a Rat Model of Tourette Syndrome. World Neurosurg. 2019;127:e881-e7.

48.	Zhang C, Wei H, Zhang Y, Du J, Liu W, Zhan S, et al. Increased dopamine transporter levels following nucleus accumbens deep brain stimulation in methamphetamine use disorder: A case report. Brain Stimul. 2019;12(4):1055-7.

49.	Nguyen MD, Venton BJ. Fast-scan Cyclic Voltammetry for the Characterization of Rapid Adenosine Release. Comput Struct Biotechnol J. 2015;13:47-54.

50.	Oh Y, Heien ML, Park C, Kang YM, Kim J, Boschen SL, et al. Tracking tonic dopamine levels in vivo using multiple cyclic square wave voltammetry. Biosens Bioelectron. 2018;121:174-82.