ROM1 is redundant to PRPH2 as a molecular building block of photoreceptor disc rims

Tylor R. Lewis[1], Mustafa S. Makia[2], Carson M. Castillo[1], Ying Hao[1], Muayyad R. Al-
Ubaidi[2,3], Nikolai P. Skiba[1], Shannon M. Conley[4], Vadim Y. Arshavsky[1,5]* and Muna I.
Naash[2,3]*

[1]Department of Ophthalmology, Duke University Medical Center, Durham, NC, USA, 27710

[2]Department of Biomedical Engineering, University of Houston, Houston, TX, USA, 77204

[3]College of Optometry, University of Houston, Houston, TX, USA, 77204

[4]Department of Cell Biology, University of Oklahoma Health Sciences Center, Oklahoma City,
OK, USA, 73104

[5]Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham,
NC, USA, 27710

*Correspondence:
Vadim Arshavsky
vadim.arshavsky@duke.edu

Muna Naash
mnaash@central.uh.edu

Abbreviated title: ROM1 and photoreceptor disc enclosure

Keywords: vision, retina, photoreceptor, outer segment, cilia, tetraspanin

Abstract

Visual signal transduction takes place within a stack of flattened membranous "discs" enclosed within the light-sensitive photoreceptor outer segment. The highly curved rims of these discs, formed in the process of disc enclosure, are fortified by large hetero-oligomeric complexes of two homologous tetraspanin proteins, PRPH2 (a.k.a. peripherin-2 or rds) and ROM1. While mutations in PRPH2 affect the formation of disc rims, the role of ROM1 remains poorly understood. In this study, we found that the knockout of ROM1 causes a compensatory increase in the disc content of PRPH2. Despite this increase, discs of ROM1 knockout mice displayed a delay in disc enclosure associated with a large diameter and lack of incisures in mature discs. Strikingly, further increasing the level of PRPH2 rescued these morphological defects. We next showed that disc rims are still formed in a knockin mouse in which the tetraspanin body of PRPH2 was replaced with that of ROM1. Together, these results demonstrate that, despite its contribution to the formation of disc rims, ROM1 can be replaced by an excess of PRPH2 for timely enclosure of newly forming discs and establishing normal outer segment structure.

Introduction

The light-sensitive outer segment organelle of vertebrate photoreceptor cells is a modified cilium containing a stack of disc-shaped membranes, or "discs". While initially formed as serial evaginations of the photoreceptor plasma membrane, mature discs are separated from this membrane and are fully enclosed inside the outer segment (Steinberg et al., 1980; Burgoyne et al., 2015; Ding et al., 2015; Volland et al., 2015) (reviewed in (Spencer et al., 2020)). Membrane enclosure involves the formation of highly curved, hairpin-shaped disc rims. Two homologous tetraspanin proteins, PRPH2 (also known as peripherin-2 or rds) and ROM1, localize specifically to disc rims (Connell and Molday, 1990; Travis et al., 1991; Bascom et al., 1992) (reviewed in (Stuck et al., 2016)). Within the disc rim, PRPH2 and ROM1 form homo- and heteromeric complexes (Bascom et al., 1992; Goldberg et al., 1995; Goldberg and Molday, 1996b; Goldberg and Molday, 1996a) that are assembled into larger oligomers connected by disulfide bonds (Goldberg et al., 1998; Loewen and Molday, 2000; Chakraborty et al., 2009; Zulliger et al., 2018). Three parallel interconnected chains of these oligomers are wrapped around the circumference of the mature disc and support its hairpin-shaped structure (Pöge et al., 2021).

Despite a high level of structural homology between PRPH2 and ROM1 (Li et al., 2003), the consequences of their mutations for photoreceptor health are quite different. Around 200 mutations of *PRPH2* are shown to cause a heterogeneous set of inherited retinal diseases, including retinitis pigmentosa, cone-rod dystrophy and macular dystrophies (Landrum et al., 2018; Peeters et al., 2021). Yet, mutations in *ROM1* typically cause digenic retinitis pigmentosa in conjunction with mutations in *PRPH2* (Kajiwara et al., 1994; Dryja et al., 1997), except for a handful of reports of mutations in *ROM1* in patients without accompanying *PRPH2* mutations (Bascom et al., 1995; Sakuma et al., 1995; Reig et al., 2000). With regard to animal models, the *Prph2* knockout mouse, commonly known as the *rds* mouse, has a complete failure of outer segment formation (Cohen, 1983; Jansen and Sanyal, 1984), while the *Rom1* knockout mouse forms outer segments with relatively minor structural abnormalities (Clarke et al., 2000). This discrepancy could be explained, at least in part, by the importance of the cytoplasmic C-terminus of PRPH2, which has been shown to retain membranes at the photoreceptor cilium, thereby allowing their remodeling into outer segment discs (Salinas et al., 2017). However, most of the

disease-associated PRPH2 mutations spare its C-terminus, so the complete explanation of this
discrepancy awaits further investigation of the role of ROM1.

In this study, we investigated the role of ROM1 in the formation of photoreceptor disc
rims and the process of disc enclosure. We first revisited the phenotype of the *Rom1* knockout
(*Rom1*[-/-]) mouse and made several novel observations. Using quantitative proteomics, we showed
that the knockout of ROM1 causes a compensatory increase in the relative disc content of
PRPH2. Despite the total tetraspanin content being essentially the same as in WT discs, *Rom1*[-/-]
discs displayed a delay in their maturation and an increase in outer segment diameter. In
addition, *Rom1*[-/-] discs lacked the indentations of their rims, known as incisures, present in WT
discs. These changes could be explained, at least in part, by an alteration in tetraspanin
oligomerization likely arising from a rearrangement of intramolecular disulfide bridge(s) in
PRPH2. Strikingly, these morphological phenotypes can be rescued by transgenic overexpression
of PRPH2, suggesting that ROM1 is not absolutely necessary for the formation of normal outer
segments. In another set of experiments, we investigated whether disc rims could be formed in a
knockin mouse in which the tetraspanin core of PRPH2 was replaced with that of ROM1. While
the knockin photoreceptors failed to form orderly disc stacks, outer segment membranes
preserved the ability to form hairpin-shaped rims. Together, these results demonstrate that,
despite its contribution to disc formation and intrinsic ability to support disc rim structure,
ROM1 can be replaced by an excess of PRPH2 for the formation of a normal outer segment.

Results

Loss of ROM1 causes a compensatory increase in the relative outer segment content of PRPH2

Three proteins, rhodopsin, PRPH2 and ROM1, comprise ~98% of the total transmembrane
protein material in normal discs (Skiba et al., 2023). Recently, it has been proposed that the
overall dimensions of a disc are determined by the molar ratio between the total tetraspanin
content (PRPH2 and ROM1) forming the disc rim and rhodopsin forming the disc surface (Lewis
et al., 2023). Therefore, interpreting the phenotype of the *Rom1*[-/-] mouse requires an
understanding of how the loss of ROM1 affects the outer segment content of PRPH2. This is
particularly important because discs deficient in PRPH2 display a compensatory increase in the
relative amount of ROM1 (Lewis et al., 2023), suggesting that the opposite (i.e., a compensatory
increase in the amount of PRPH2 when there is a deficiency in ROM1) may also be true.

To address this possibility, we employed a quantitative mass spectrometry approach
(Skiba et al., 2023) recently used to determine the molar ratio among PRPH2, ROM1 and
rhodopsin in other mutant photoreceptors (Lewis et al., 2023). In these experiments, the outer
segment content of PRPH2 was determined as a molar fraction of rhodopsin. Because the
packing density of rhodopsin in discs is unaffected by any changes in disc dimensions (Liang et
al., 2004), this parameter reflects the relative content of PRPH2 in each disc, independent of any
changes in outer segment length or volume. We found that the relative content of PRPH2 in
Rom1[-/-] outer segments increased to an ~1:12 molar ratio to rhodopsin from an ~1:18 ratio in WT
outer segments (**Table 1**; see raw data in **Supplementary Table 1**). Strikingly, this 1:12 ratio is
equal to the ratio between total tetraspanin (PRPH2 + ROM1) and rhodopsin in WT outer
segments. Therefore, the knockout of ROM1 leads to a compensatory increase in the relative disc
content of PRPH2.

Loss of ROM1 delays disc enclosure causing increased outer segment diameter and occasional disc overgrowth

The retinal phenotype of $Rom1^{-/-}$ mice was analyzed at postnatal day 30 (P30), as photoreceptor cell degeneration was just beginning at this age (**Figs. 1A,B**). Interestingly, light microscopy imaging was sufficient to identify that the outer segment layer of these mice appeared disorganized and even shortened (**Figs. 1A,C**). Please note that this and the three subsequent figures also include data obtained with a mouse overexpressing PRPH2 on the $Rom1^{-/-}$ background (*PRPH2 OE*/$Rom1^{-/-}$), which will be described below, to facilitate a side-by-side comparison of all three phenotypes. Using transmission electron microscopy (TEM), we observed a number of outer segment structural abnormalities (**Fig. 2** and **Supplementary Fig. 1**), consistent with those noted in the original report (Clarke et al., 2000). In general, outer segments appeared to be shorter and wider, with some of them displaying overgrown disc membranes not aligned in a stack (yellow arrows, **Fig 2B**). The degree of this overgrowth varied, with the extended membranes occasionally wrapping around the entire outer segment.

Because these ultrastructural defects are reminiscent of those from several PRPH2 mutant lines in which photoreceptor disc enclosure is affected (Lewis et al., 2021), we assessed the status of disc enclosure in $Rom1^{-/-}$ outer segments. We contrasted retinal tissue with tannic acid and uranyl acetate instead of the traditionally used osmium tetroxide (Ding et al., 2015). This technique yields a darker staining of newly forming, "open" discs exposed to the extracellular space than discs fully enclosed within the outer segment. This approach revealed that even outer segments lacking gross abnormalities in $Rom1^{-/-}$ mice have an increased number of open discs at their base (**Figs. 3A,B**). Whereas WT rods contained a relatively constant number of ~7 open discs (consistent with previous reports (Ding et al., 2015; Volland et al., 2015; Lewis et al., 2021)), $Rom1^{-/-}$ rods had an average of ~14 open discs. In addition, the number of open discs in $Rom1^{-/-}$ rods was much more variable than in WT rods, with up to ~30 open discs occasionally observed. Lastly, we quantified outer segment diameters of $Rom1^{-/-}$ rods and found that they were on average ~35% wider than WT rods (**Fig. 3C**), which is close to the ~43% increase previously reported in (Clarke et al., 2000).

These findings show that ROM1 contributes to the process of disc enclosure and that the compensatory increase in the disc content of PRPH2 does not fully replace ROM1 in this capacity. Accordingly, the increase in outer segment diameter may be explained by prolonged delivery of disc membrane material to each disc before it becomes fully enclosed. On occasion, an expansion of open discs becomes uncontrolled leading to formation of membranous whorls.

Another phenotype of $Rom1^{-/-}$ mice was revealed using TEM of tangentially sectioned retinas (**Fig. 4**). Normal mouse discs contain a single deep indentation of their rims, called an incisure (yellow arrowheads, **Fig. 4**). In contrast, no incisures were observed in $Rom1^{-/-}$ discs. Because the rims of both disc circumference and incisure are built from PRPH2/ROM1, the lack of incisures in $Rom1^{-/-}$ discs may be explained by the entire pool of remaining tetraspanin being deposited at the circumference of larger discs.

ROM1 regulates the supramolecular organization of PRPH2

We next sought to address why the compensatory increase in the disc content of PRPH2 in $Rom1^{-/-}$ rods is insufficient to support normal disc morphogenesis. In this context, it is important to consider that defects in PRPH2 oligomerization, without an accompanying decrease in PRPH2 levels, also lead to defects in disc enclosure (Lewis et al., 2021). Therefore, we reasoned that loss

of ROM1 may modulate the status of PRPH2 oligomerization. In fact, the idea that ROM1 may regulate the formation of high order PRPH2 oligomers has been put forward in a previous study (Loewen and Molday, 2000).

To address the possibility that PRPH2 oligomerization may be disrupted without ROM1, we first performed SDS-PAGE of retinal lysates from WT and *Rom1*[-/-] mice under non-reducing conditions (i.e., no DTT in samples). Under these conditions, PRPH2 is represented by two bands, one corresponding to ~35 kDa monomers and another to ~75 kDa dimers linked by a disulfide bond (**Fig. 5A**). Both bands are somewhat diffuse, likely reflecting the heterogeneous status of PRPH2 glycosylation. The majority of PRPH2 in WT retinas is represented by dimers, consistent with its origin from large, disulfide-bound oligomers.

Two differences were observed in *Rom1*[-/-] retinas. First, there was an ~50% increase in the fraction of PRPH2 that runs in a monomeric state (**Figs. 5A,B**), indicating that PRPH2 oligomerization may be negatively affected by loss of ROM1. Second, the monomeric form of PRPH2 was represented by a doublet, consistent with a previous observation (Stuck et al., 2015). This doublet was consolidated into a single band under reducing conditions in the presence of DTT (**Fig. 5C**), which suggests that the two bands in *Rom1*[-/-] retinas represent pools of PRPH2 molecules with different patterns of internal disulfide bonds. Apart from the C150 residue involved in intermolecular disulfide bonding of a PRPH2 molecule to another PRPH2 or ROM1, there are six other conserved cysteines in PRPH2 that are involved in intramolecular disulfide bonding (Goldberg et al., 1998). Therefore, it is possible that ROM1 contributes to PRPH2 assuming the intermolecular disulfide bond conformation optimal for tetraspanin oligomerization at the disc rim.

To further investigate the status of PRPH2 oligomerization in *Rom1*[-/-] mice, we utilized a technique typically used to describe PRPH2 and ROM1 complexes of various sizes. In this approach, complexes are extracted from disc membranes in the presence of an non-ionic detergent and separated by velocity sedimentation on a sucrose gradient under non-reducing conditions (Clarke et al., 2000; Goldberg et al., 2001; Ding et al., 2004; Chakraborty et al., 2008; Chakraborty et al., 2009; Stuck et al., 2014; Zulliger et al., 2018; Milstein et al., 2020; Lewis et al., 2021). We performed velocity sedimentation of lysates from both WT and *Rom1*[-/-] retinas followed by SDS-PAGE of gradient fractions under non-reducing conditions (**Figs. 5D,E**). The distribution of the PRPH2 monomeric form across fractions was unaffected by the ROM1 knockout. In both WT and *Rom1*[-/-] retinas, it was most abundant in fractions #6-8, shown to correspond to PRPH2 core complexes (Chakraborty et al., 2008). However, the loss of ROM1 shifted the distribution of the disulfide-bound PRPH2 dimers to smaller oligomeric forms, consistent with a previous report (Conley et al., 2019). Taken together, these data show that loss of ROM1 does indeed affect the status of PRPH2 oligomerization and suggest that abnormal PRPH2 oligomerization underlies the defects in disc enclosure in *Rom1*[-/-] mice.

Transgenic overexpression of PRPH2 can compensate for the loss of ROM1

We next investigated whether a further increase in the level of PRPH2 could improve the morphological defects observed in *Rom1*[-/-] outer segments. We employed a transgenic line that expresses PRPH2 in both rods and cones under control of the human IRBP promoter (Nour et al., 2004). While originally called NMP, we refer to this line as *PRPH2 OE* (PRPH2 overexpressor). In the current study, we used *PRPH2 OE* mice with a single copy of this transgene, which has been shown to express ~30% excess PRPH2 over WT levels (Nour et al., 2004). *PRPH2 OE*

mice were crossed with *Rom1*[-/-] mice and retinas were analyzed as for WT and *Rom1*[-/-] mice
(**Figs. 1-3**). Strikingly, the gross morphological defects observed in *Rom1*[-/-] mice were
significantly improved by transgenic overexpression of PRPH2 (**Fig. 2A**). The outer segment
length was somewhat restored (**Figs. 1A,C**), although the degree of this effect did not reach
statistical significance. Furthermore, PRPH2 overexpression rescued the disc enclosure defect of
Rom1[-/-] outer segments by restoring the number of open discs to the WT level (**Figs. 3A,B**). In
addition, *PRPH2 OE/Rom1*[-/-] outer segments had normal diameters (**Figs. 3A,C**) and their discs
contained incisures (**Fig. 4**). These data indicate that, whereas ROM1 contributes to disc
formation in WT rods, it can be functionally replaced by a sufficient excess of PRPH2.

ROM1 is able to form disc rims in the absence of PRPH2

While we have shown that ROM1 contributes to the formation of disc rims, it is unclear whether
it can do so in the absence of PRPH2. Because disc formation is completely abolished in the
absence of PRPH2, we could not address this question using PRPH2 knockout mice. Instead, we
utilized a knockin mouse model, called RRCT (Conley et al., 2019), in which the tetraspanin
core of PRPH2 was replaced with that of ROM1 (**Fig. 6A**). This chimeric protein contains the C-
terminus of PRPH2 required for retaining disc membranes at the outer segment base (Salinas et
al., 2017), while replacing the tetraspanin core required for its oligomerization and functioning in
disc rim formation (Salinas et al., 2017; Conley et al., 2019).

As previously reported (Conley et al., 2019), homozygous RRCT photoreceptors formed
rudimentary outer segments (**Fig. 6B**). The small number and size of these outer segment-like
structures is likely explained by a very low expression of the chimeric protein (Conley et al.,
2019), which may lead to a low efficiency of disc membrane retention. We now show that some
of these structures contain stacked disc membranes that are partially enclosed within the outer
segment plasma membrane. Importantly, these disc membranes have discernible rims with a
characteristic hairpin-like shape. This finding indicates that ROM1 alone is, in principle, able to
support the formation of disc rims in the absence of PRPH2.

In an additional set of experiments, we explored the oligomerization status of ROM1 and
the RRCT chimera in the absence of PRPH2. Unfortunately, the severe disruptions of outer
segments in homozygous RRCT mice precluded us from obtaining a sufficient amount of retinal
lysate to perform this analysis. Instead, we used an *in vitro* system in which we co-transfected
equal amounts of RRCT and ROM1 in COS7 cells (**Fig. 6C**). In these cells, RRCT and ROM1
co-localized in structures that were distinct from the endoplasmic reticulum, as evident from the
lack of co-localization with the endoplasmic reticulum marker, calreticulin, suggesting that they
were normally processed in the biosynthetic membranes. Velocity sedimentation of lysates from
these cells revealed that both RRCT and ROM1 sediment in fractions #6-8 (**Figs. 6D-E**), which
is the same as for PRPH2/ROM1 core complexes from WT photoreceptors. These data indicate
that heterologously expressed RRCT and ROM1 can form core tetraspanin complexes, even in
the absence of PRPH2, that are able to support disc rim formation.

Disc rims can be formed without disulfide bonds between tetraspanin molecules

In a final experiment, we investigated whether the formation of intermolecular disulfide bonds
between tetraspanins is dispensable for disc rim formation. We have previously shown that
disulfide bonds between PRPH2 molecules are not absolutely required for this function because
disc rims were formed in *Prph2*[C150S/C150S] knockin mice lacking the cysteine residue forming this
bond (Lewis et al., 2020). Yet ROM1 also forms intermolecular disulfide bonds (Chakraborty et

al., 2008), which raises the question of whether disc rim formation in $Prph2^{C150S/C150S}$ knockin mice was driven by disulfide-linked ROM1 molecules.

To address this question, we analyzed disc rim structure in $Rom1^{-/-}$ mice bearing the $Prph2^{C150S/C150S}$ mutation. Outer segments of these mice had morphological defects comparable to those of $Prph2^{C150S/C150S}$ mice (**Fig. 7A**). Nonetheless, orderly disc stacks were occasionally observed in these retinas. Higher magnification images of these stacks revealed the presence of disc rims which looked comparable to those in $Prph2^{C150S/C150S}$ retinas (**Fig. 7B**). This finding demonstrates that, whereas disulfide bonds connecting tetraspanin molecules are essential for maintaining the overall outer segment structure, these bonds are not absolutely required for the formation of disc rims.

In addition, staining with tannic acid/uranyl acetate revealed that the outer segments of $Prph2^{C150S/C150S}$/$Rom1^{-/-}$ mice contained both open (yellow arrows, **Fig. 7A**) and enclosed discs (yellow arrowheads, **Fig. 7A**). This indicates that disulfide bonds among PRPH2 and ROM1 molecules are also dispensable for disc enclosure.

Discussion

In this study, we analyzed the role of ROM1 in supporting the formation of outer segment discs. In addition to the previously described morphological defects of $Rom1^{-/-}$ outer segments, we showed that these outer segments experience a delay in disc enclosure. This delay leads to an increase in disc diameter, loss of incisures and occasional uncontrolled disc membrane outgrowth. Remarkably, these defects can be reversed by sufficient overexpression of PRPH2, indicating a functional redundancy between these two tetraspanin proteins.

The idea of redundancy between PRPH2 and ROM1 is supported by the finding that mutations in both genetically interact to modulate disease (Kajiwara et al., 1994; Dryja et al., 1997). While individual PRPH2 mutations have greater phenotypic consequences, this could relate to both the unique role of the C-terminal tail of PRPH2 in the initial stages of disc formation (Salinas et al., 2017) as well as the fact that there is twice as much PRPH2 as ROM1 in outer segments, at least in mice (Kedzierski et al., 1999; Loewen and Molday, 2000; Skiba et al., 2023).

While evidence that *ROM1* mutations alone may cause human visual pathology is limited (Bascom et al., 1995; Sakuma et al., 1995; Reig et al., 2000), photoreceptors of $Rom1^{-/-}$ mice eventually degenerate (Clarke et al., 2000). We propose that the delay in disc enclosure is the primary defect underlying this degeneration. Because discs remain open longer than normal, they may incorporate more than a normal amount of membranous material, which is continuously delivered to the outer segment. This can lead to disc membrane overgrowth and, eventually, the formation of highly dysmorphic membrane whorls. This progressive collapse of outer segment ultrastructure would eventually lead to cell death. Notably, this same progression of structural defects occurs in certain mice in which PRPH2 oligomerization is affected (Lewis et al., 2021).

But why is the phenotype of the heterozygous *rds* mouse containing ~50% of normal PRPH2 protein (Hawkins et al., 1985; Cheng et al., 1997) much more severe than the $Rom1^{-/-}$ mouse lacking 100% of ROM1? This difference is not intuitive because the molar ratio between PRPH2 and ROM1 is ~2:1 and, therefore, each mouse is expected to have ~2/3 of the normal tetraspanin content. However, there is a compensatory increase in outer segment tetraspanin in

each of these two mice. We now show that relative content of total tetraspanin in *Rom1*$^{-/-}$ discs is
essentially the same as in WT discs (~1:12 molar ratio to rhodopsin) due to a compensatory
increase in PRPH2. On the other hand, there is only a partial compensatory increase of ROM1 in
heterozygous *rds* mice (~1:15 molar ratio of PRPH2+ROM1 to rhodopsin) (Lewis et al., 2023).
The mechanism behind these increases remains to be elucidated, but regardless, the lesser
severity of the *Rom1*$^{-/-}$ phenotype could be explained, at least in part, by a larger degree of this
compensatory tetraspanin increase.

Yet, the complete restoration of the total tetraspanin content in *Rom1*$^{-/-}$ outer segments
does not fully rescue their morphological phenotype. This implies that, in normal photoreceptors,
ROM1 contributes some unique functional features to the properties of tetraspanin oligomers. In
fact, our data suggest that the loss of ROM1 alters both the intra- and intermolecular properties
of the remaining PRPH2. We found that ROM1 likely facilitates an optimal internal disulfide
bond arrangement of PRPH2 in addition to promoting the formation of larger oligomeric chains
of PRPH2. One possible explanation for the ability of ROM1 to potentially facilitate the optimal
disulfide bond arrangement of PRPH2 may relate to the intracellular trafficking of these two
proteins. Whereas the majority of PRPH2 is normally trafficked through the unconventional
secretory pathway from the endoplasmic reticulum that bypasses the Golgi (Tian et al., 2014;
Zulliger et al., 2015; Pearring et al., 2021), ROM1 was shown to direct a small fraction of
PRPH2 through the conventional secretory pathway (Conley et al., 2019). It is intriguing to
speculate that a shift in the intramolecular disulfide bonds of PRPH2 in the absence of ROM1
may arise from a complete loss of PRPH2 trafficking through the conventional pathway.
Nonetheless, the outer segment deficiencies arising from loss of ROM1 were nearly completely
overcome by a sufficient excess of PRPH2.

Of particular interest is our finding that discs of *Rom1*$^{-/-}$ mice lack incisures. A recent
study put forth a model in which the total length of the disc rim, including the incisure, is
determined by the molar ratio between the total tetraspanin protein and rhodopsin (Lewis et al.,
2023). The lack of incisures in *Rom1*$^{-/-}$ discs may not appear intuitive since the total tetraspanin
content in these discs is essentially the same as in WT discs. However, *Rom1*$^{-/-}$ discs also have an
increased diameter, which allows the total rim length of WT and *Rom1*$^{-/-}$ discs to be about the
same. The reason why such a tradeoff between the disc diameter and the presence/absence of an
incisure exists remains to be determined, although it is likely to relate to the kinetics of disc
enclosure.

In conclusion, our study shows that ROM1 makes distinct contributions to the formation
of disc rims and the process of disc enclosure. Nonetheless, a sufficient excess of PRPH2 is able
to compensate for the loss of ROM1, indicating that ROM1 is redundant to PRPH2 as a
molecular building block of photoreceptor disc rims. Given that excess PRPH2 can rescue
defects associated with loss of ROM1, it is exciting to speculate that PRPH2 overexpression can
prevent the long-term photoreceptor degeneration associated with either loss of ROM1 or
deficiencies arising from mutations in PRPH2 itself. It also remains to be seen whether the
opposite is also true, whereby an excess ROM1 may rescue certain defects associated with
mutations in PRPH2.

Materials and Methods

Mouse (*Mus musculus*) husbandry

Animal maintenance and experiments were approved by the local Institutional Animal Care and Use Committee (IACUC; University of Houston, TX, USA) and guidelines as stated by the Association for Research in Vision and Ophthalmology (Rockville, MD). The generation of the $Prph2^{C150S}$ mouse was previously described in (Stuck et al., 2014). The generation of the RRCT mouse was previously described in (Conley et al., 2019). The generation of the *PRPH2 OE* mouse was previously described in (Nour et al., 2004). The $Rom1^{-/-}$ mouse, described in (Clarke et al., 2000), was generously provided by Roderick R McInnes (McGill University). All mice were genotyped to ensure that they did not contain either the *rd8* (Mattapallil et al., 2012) or *rd1* (Pittler et al., 1993) mutations commonly found in inbred mouse strains. All mice were on a C57BL/6 genetic background bearing the RPE65 L450 variant and housed under a 12/12 hour diurnal light (~30 lux) cycle.

Transmission electron microscopy (TEM)

For all main figures, fixation and processing of mouse eyes for TEM was performed as described previously (Ding et al., 2015). In the afternoon, anesthetized mice were transcardially perfused with 2% paraformaldehyde, 2% glutaraldehyde and 0.05% calcium chloride in 50 mM MOPS (pH 7.4) resulting in exsanguination. Enucleated eyes were fixed for an additional 2 h in the same fixation solution at room temperature. Eyecups were dissected from fixed eyes, embedded in 2.5% low-melt agarose (Precisionary, Greenville, NC) and cut into 200 µm thick slices on a Vibratome (VT1200S; Leica, Buffalo Grove, IL). Agarose sections were stained with 1% tannic acid (Electron Microscopy Sciences, Hatfield, PA) and 1% uranyl acetate (Electron Microscopy Sciences), gradually dehydrated with ethanol and infiltrated and embedded in Spurr's resin (Electron Microscopy Sciences). 70 nm sections were cut, placed on copper grids and counterstained with 2% uranyl acetate and 3.5% lead citrate (19314; Ted Pella, Redding, CA). The samples were imaged on a JEM-1400 electron microscope (JEOL, Peabody, MA) at 60 kV with a digital camera (BioSprint; AMT, Woburn, MA). Image analysis and processing was performed with ImageJ. For Supplementary Figure 1, fixation, processing, and imaging of mouse eyes for TEM was performed as described previously (Stricker et al., 2005).

Light microscopy of histological sections

Plastic embedded blocks generated for TEM were sectioned through the optic nerve in 500 nm sections and stained with methylene blue for light microscopy as previously described (Lobanova et al., 2008). Images were taken with a confocal microscope (Eclipse 90i and A1 confocal scanner; Nikon) with a 60× objective (1.4 NA Plan Apochromat VC; Nikon) using Nikon NIS-Elements software. Image analysis and processing was performed with ImageJ.

Preparation of retinal lysates

Eyecups were dissected and immediately frozen using liquid nitrogen and stored at −80°C prior to processing. Lysates were prepared essentially as previously described (Stuck et al., 2014). In short, individual eyecups were lysed in 200 µl solubilization buffer (PBS, pH 7.0, containing 1% Triton X-100, 5 mM EDTA, 5 mg/ml NEM and protease inhibitors (Roche, Mannheim, Germany)). Samples were incubated on ice for 1 h prior to being centrifuged for 30 min at 20,000g at 4°C. The supernatant was collected and subjected to either Western blot analysis directly or velocity sedimentation.

Western blotting of retinal lysates

Protein concentration was assayed by using a colorimetric Bradford assay (Bio-Rad, Hercules,
CA, USA). Western blotting was performed essentially as previously described (Spencer et al.,
2016). In short, lysates were incubated with Laemmli sample buffer (50 mM Tris-HCl, 2% SDS,
10% glycerol and 1% Bromophenol Blue) with or without 100 mM DTT for reducing or non-
reducing blots respectively. Samples containing 10 µg of total protein were incubated at 90°C for
5 minutes and run on a 10-20% Tris-HCl gel. Gels were transferred onto PVDF membrane and
blocked with Intercept (PBS) Blocking Buffer (Li-Cor, Lincoln, Nebraska) with 0.25% Tween-
20. Blots were incubated overnight at 4°C with 1:1000 dilution of polyclonal rabbit anti-PRPH2
C-terminal antibody (Kedzierski et al., 1999). After primary antibody incubation, blots were
washed and incubated with 1:10000 dilution of donkey anti-rabbit DyLight 800 (Invitrogen,
Carlsbad, CA) for 2 h. All experiments were repeated at least three times. Blots were imaged
using Odyssey CLx imaging system (Li-Cor).

Velocity sedimentation using sucrose gradients

Sucrose gradients were prepared as previously described (Chakraborty et al., 2009). In short,
gradients of 5–20% sucrose were prepared by sequentially layering 0.5 ml each of 20, 15, 10 and
5% sucrose solutions in PBS with 0.1% Triton X-100 and 10 mM NEM and allowing them to sit
at room temperature for 1 h to equilibrate. Gradients were then chilled on ice for 30 minutes
prior to loading lysate and centrifugation at 40,000 rpm on a TLS-55 swinging bucket rotor
(Beckman Coulter, Brea, CA) for 16 hours at 4°C. The bottom of each tube was pierced with a
21G needle and 12 fractions of ~180 µl each were collected. Western blotting was performed as
described above using 15 µl samples for each fraction.

Cell culture experiments

COS7 cells were transfected with 7.5 µg of RRCT-FLAG and ROM1 mammalian expression
constructs that were previously described (Conley et al., 2010; Conley et al., 2019) using a
calcium phosphate transfection protocol as previously described (Conley et al., 2019). After 48
hours, cells were processed for immunofluorescent staining or velocity sedimentation as
described below. Experiments were repeated three independent times.

Immunofluorescent staining was performed as described in (Conley et al., 2010). After
blocking, coverslips were incubated overnight in primary antibodies to label RRCT, ROM1 and
ER membranes as follows: 1) 1:2 dilution of monoclonal mouse anti-RRCT (2E7; described in
(Zulliger et al., 2015), 2) 1:400 dilution of polyclonal rabbit anti-ROM1 (2Rom1; described in
(Ding et al., 2005)), and 3) 1:250 dilution of polyclonal chicken anti-calreticulin at 1:250
(ab2908; Abcam, Cambridge, UK). After washing, cells were incubated with 1:000 dilution of
donkey anti-mouse Alexa 568 (Invitrogen), donkey anti-rabbit Alexa 647 (Invitrogen) and goat
anti-chicken Alexa 488 (Invitrogen) secondary antibodies for 1 h. Cells were mounted in
ProlongGold with DAPI (ThermoFisher). Images were captured on a BX-62 spinning disk
confocal microscope equipped with an ORCA-ER camera (Olympus, Japan) and analyzed with
Slidebook™ 5.2 software (Intelligent Imaging Innovations, Denver, CO). Images from the
spinning disk confocal microscope were deconvolved using the nearest neighbors paradigm.
Images were captured with 100x/1.40 oil objectives, and exposure times and display settings
(brightness and contrast) for all images were normalized to a control section where primary
antibody was omitted during processing. No gamma adjustments were made to
immunofluorescent images.

For velocity sedimentation, scraped cells were lysed in the same solubilization buffer
used to prepare retinal extracts. Lysates were separated as described above (200 µg protein
lysate/gradient). Resulting gradient fractions were separated by SDS-PAGE under reducing
conditions and blots were probed with either 1:500 dilution of monoclonal mouse anti-RRCT
(2B7; described in (Conley et al., 2014)) or monoclonal mouse anti-ROM1 (2H5, described in
(Conley et al., 2014)). Blots were probed with goat anti-mouse and anti-rabbit HRP secondaries
(SeraCare, Milford, MA), imaged on a Bio-Rad ChemiDoc imager and analyzed in Image Lab
Software version 6.0.1 (Bio-Rad).

Quantitative mass spectrometry

A crude preparation of rod outer segments was obtained as described in (Lewis et al., 2023).
Dissected mouse retinas were vortexed in 8% OptiPrep in mouse Ringer's solution (containing
130 mM NaCl, 3.6 mM KCl, 2.4 mM $MgCl_2$, 1.2 mM $CaCl_2$, and 10 mM HEPES, pH 7.4) that
was adjusted to 314 mOsm. The preparation was briefly left on ice to allow the remaining retinal
tissue to settle. The supernatant was removed and centrifuged at 20,000 x g. Pelleted outer
segments were gently washed with mouse Ringer's solution before lysis with 2% SDS in PBS.
Protein concentration was measured with the Bio-Rad Protein Assay kit (Bio-Rad). Samples
containing 5-10 µg of protein were mixed with 0.25-0.5 µg BSA (used as an internal standard in
this analysis) and cleaved with 1 µg trypsin/LysC mix (Promega, Madison, WI) using the SP3
beads protocol described in (Hughes et al., 2014). The combined digest of outer segments and
BSA was mixed with the digest of a chimeric protein consisting of concatenated tryptic peptides
of outer segment proteins, including rhodopsin, PRPH2 and ROM1, which is described in (Skiba
et al., 2023). Mass spectrometry, data processing and data analysis were also performed as
described in (Skiba et al., 2023). For each genotype, a total of two biological replicates were
analyzed.

Experimental design and statistical analysis

For the quantification of the number of photoreceptor nuclei (**Fig. 1B**), photoreceptor nuclei
were counted in 100 µm boxes at 500 µm intervals from the optic nerve spanning 2000 µm in
each direction for three mice of each genotype, as previously described (Lobanova et al., 2018).
Two-way ANOVA was performed to determine statistical significance across genotype and
location.

For the quantification of the number of open discs (**Fig. 3B**), darkly-stained new discs at
the base of the rod outer segment were counted until the first lightly-stained enclosed disc. 3
retinas were collected of each genotype with the number of outer segments analyzed as follows:
WT1, 44; WT2, 40; WT3, 68; *Rom1*$^{-/-}$1, 87; *Rom1*$^{-/-}$2, 42; *Rom1*$^{-/-}$3, 44; *PRPH2 OE/Rom1*$^{-/-}$1, 48;
PRPH2 OE/Rom1$^{-/-}$2, 35; *PRPH2 OE/Rom1*$^{-/-}$3, 59. Data were plotted with samples separated,
while statistical analysis was performed on the averages within each retina (n=3 for each
genotype). One-way ANOVA with Tukey's multiple comparisons test was performed to
determine statistical significance across genotypes. For the quantification of outer segment
diameter (**Fig. 3C**), 3 retinas were collected of each genotype with the number of outer segments
analyzed as follows: WT1, 88; WT2, 97; WT3, 92; *Rom1*$^{-/-}$1, 83; *Rom1*$^{-/-}$2, 81; *Rom1*$^{-/-}$3, 85;
PRPH2 OE/Rom1$^{-/-}$1, 75; *PRPH2 OE/Rom1*$^{-/-}$2, 66; *PRPH2 OE/Rom1*$^{-/-}$3, 78. Data were plotted
with samples separated, while statistical analysis was performed on the averages within each

retina (n=3 for each genotype). One-way ANOVA with Tukey's multiple comparisons test was performed to determine statistical significance across genotypes.

Densitometric analysis of non-saturated bands from Western blots (**Figs. 5,6**) was performed using Image Lab software v4.1 (Bio-Rad) and ImageJ. For quantification of the PRPH2 monomer:dimer ratio (**Fig. 5B**), 3 samples were used for both WT and *Rom1*$^{-/-}$ mice. Unpaired t-test was performed to determine statistical significance of the PRPH2 monomer:dimer ratio between WT and *Rom1*$^{-/-}$ retinas). For the velocity sedimentation analysis of PRPH2 supramolecular organization (**Fig. 5E**), 11 samples were used for WT and 4 samples were used for *Rom1*$^{-/-}$ mice. Two-way ANOVA was performed to determine statistical significance across genotypes and fractions followed by Sidak's multiple comparisons post-hoc test to determine statistical significance between genotypes in each fraction for the disulfide-bound form. For the quantification of ROM1 and RRCT in each fraction of transfected COS7 cells (**Fig. 6E**), 3 samples were analyzed.

All experiments were performed with mice of randomized sex. Sample sizes were determined based on previous published experiments. Data were graphed with Prism 9 (GraphPad, San Diego, CA) with error bars depicting the S.E.M. Statistical analyses were performed using Prism 9 (GraphPad). Where noted, statistical values are depicted in graphs with asterisks as follows: $p < 0.05$, *; $p < 0.01$, **, $p < 0.001$, ***; $p < 0.0001$, ****.

Acknowledgements

This work was supported by the National Institutes of Health grants EY030451 (VYA), EY005722 (VYA), EY010609 (MIN and MRA), EY034671 (MIN), EY033763 (TRL), AG070915 (SMC) and an Unrestricted Award from Research to Prevent Blindness Inc. (Duke University).

Competing interests

No competing interests declared.

References

Bascom, R. A., Liu, L., Heckenlively, J. R., Stone, E. M. and McInnes, R. R. (1995). Mutation analysis of the ROM1 gene in retinitis pigmentosa. *Hum Mol Genet* **4**, 1895-902.

Bascom, R. A., Manara, S., Collins, L., Molday, R. S., Kalnins, V. I. and McInnes, R. R. (1992). Cloning of the cDNA for a novel photoreceptor membrane protein (rom-1) identifies a disk rim protein family implicated in human retinopathies. *Neuron* **8**, 1171-84.

Burgoyne, T., Meschede, I. P., Burden, J. J., Bailly, M., Seabra, M. C. and Futter, C. E. (2015). Rod disc renewal occurs by evagination of the ciliary plasma membrane that makes cadherin-based contacts with the inner segment. *Proc Natl Acad Sci U S A* **112**, 15922-7.

Chakraborty, D., Ding, X. Q., Conley, S. M., Fliesler, S. J. and Naash, M. I. (2009). Differential requirements for retinal degeneration slow intermolecular disulfide-linked oligomerization in rods versus cones. *Hum Mol Genet* **18**, 797-808.

Chakraborty, D., Ding, X. Q., Fliesler, S. J. and Naash, M. I. (2008). Outer segment oligomerization of Rds: evidence from mouse models and subcellular fractionation. *Biochemistry* **47**, 1144-56.

Cheng, T., Peachey, N. S., Li, S., Goto, Y., Cao, Y. and Naash, M. I. (1997). The effect of peripherin/rds haploinsufficiency on rod and cone photoreceptors. *J Neurosci* **17**, 8118-28.

Clarke, G., Goldberg, A. F., Vidgen, D., Collins, L., Ploder, L., Schwarz, L., Molday, L. L., Rossant, J., Szel, A., Molday, R. S. et al. (2000). Rom-1 is required for rod photoreceptor viability and the regulation of disk morphogenesis. *Nat Genet* **25**, 67-73.

Cohen, A. I. (1983). Some cytological and initial biochemical observations on photoreceptors in retinas of rds mice. *Invest Ophthalmol Vis Sci* **24**, 832-43.

Conley, S. M., Stricker, H. M. and Naash, M. I. (2010). Biochemical analysis of phenotypic diversity associated with mutations in codon 244 of the retinal degeneration slow gene. *Biochemistry* **49**, 905-11.

Conley, S. M., Stuck, M. W., Burnett, J. L., Chakraborty, D., Azadi, S., Fliesler, S. J. and Naash, M. I. (2014). Insights into the mechanisms of macular degeneration associated with the R172W mutation in RDS. *Hum Mol Genet* **23**, 3102-14.

Conley, S. M., Stuck, M. W., Watson, J. N., Zulliger, R., Burnett, J. L. and Naash, M. I. (2019). Prph2 initiates outer segment morphogenesis but maturation requires Prph2/Rom1 oligomerization. *Hum Mol Genet* **28**, 459-475.

Connell, G. J. and Molday, R. S. (1990). Molecular cloning, primary structure, and orientation of the vertebrate photoreceptor cell protein peripherin in the rod outer segment disk membrane. *Biochemistry* **29**, 4691-8.

Ding, J. D., Salinas, R. Y. and Arshavsky, V. Y. (2015). Discs of mammalian rod photoreceptors form through the membrane evagination mechanism. *J Cell Biol* **211**, 495-502.

Ding, X. Q., Nour, M., Ritter, L. M., Goldberg, A. F., Fliesler, S. J. and Naash, M. I. (2004). The R172W mutation in peripherin/rds causes a cone-rod dystrophy in transgenic mice. *Hum Mol Genet* **13**, 2075-87.

Ding, X. Q., Stricker, H. M. and Naash, M. I. (2005). Role of the second intradiscal loop of peripherin/rds in homo and hetero associations. *Biochemistry* **44**, 4897-904.

Dryja, T. P., Hahn, L. B., Kajiwara, K. and Berson, E. L. (1997). Dominant and digenic mutations in the peripherin/RDS and ROM1 genes in retinitis pigmentosa. *Invest Ophthalmol Vis Sci* **38**, 1972-82.

Goldberg, A. F., Fales, L. M., Hurley, J. B. and Khattree, N. (2001). Folding and
subunit assembly of photoreceptor peripherin/rds is mediated by determinants within the
extracellular/intradiskal EC2 domain: implications for heterogeneous molecular pathologies. *J*
Biol Chem **276**, 42700-6.
Goldberg, A. F., Loewen, C. J. and Molday, R. S. (1998). Cysteine residues of
photoreceptor peripherin/rds: role in subunit assembly and autosomal dominant retinitis
pigmentosa. *Biochemistry* **37**, 680-5.
Goldberg, A. F. and Molday, R. S. (1996a). Defective subunit assembly underlies a
digenic form of retinitis pigmentosa linked to mutations in peripherin/rds and rom-1. *Proc Natl*
Acad Sci U S A **93**, 13726-30.
Goldberg, A. F. and Molday, R. S. (1996b). Subunit composition of the peripherin/rds-
rom-1 disk rim complex from rod photoreceptors: hydrodynamic evidence for a tetrameric
quaternary structure. *Biochemistry* **35**, 6144-9.
Goldberg, A. F., Moritz, O. L. and Molday, R. S. (1995). Heterologous expression of
photoreceptor peripherin/rds and Rom-1 in COS-1 cells: assembly, interactions, and localization
of multisubunit complexes. *Biochemistry* **34**, 14213-9.
Hawkins, R. K., Jansen, H. G. and Sanyal, S. (1985). Development and degeneration
of retina in rds mutant mice: photoreceptor abnormalities in the heterozygotes. *Exp Eye Res* **41**,
701-20.
Hughes, C. S., Foehr, S., Garfield, D. A., Furlong, E. E., Steinmetz, L. M. and
Krijgsveld, J. (2014). Ultrasensitive proteome analysis using paramagnetic bead technology.
Mol Syst Biol **10**, 757.
Jansen, H. G. and Sanyal, S. (1984). Development and degeneration of retina in rds
mutant mice: electron microscopy. *J Comp Neurol* **224**, 71-84.
Kajiwara, K., Berson, E. L. and Dryja, T. P. (1994). Digenic retinitis pigmentosa due
to mutations at the unlinked peripherin/RDS and ROM1 loci. *Science* **264**, 1604-8.
Kedzierski, W., Weng, J. and Travis, G. H. (1999). Analysis of the rds/peripherin.rom1
complex in transgenic photoreceptors that express a chimeric protein. *J Biol Chem* **274**, 29181-7.
Landrum, M. J., Lee, J. M., Benson, M., Brown, G. R., Chao, C., Chitipiralla, S.,
Gu, B., Hart, J., Hoffman, D., Jang, W. et al. (2018). ClinVar: improving access to variant
interpretations and supporting evidence. *Nucleic Acids Res* **46**, D1062-d1067.
Lewis, T. R., Makia, M. S., Castillo, C. M., Al-Ubaidi, M. R., Naash, M. I. and
Arshavsky, V. Y. (2021). Photoreceptor Disc Enclosure Is Tightly Controlled by Peripherin-2
Oligomerization. *J Neurosci* **41**, 3588-3596.
Lewis, T. R., Makia, M. S., Kakakhel, M., Al-Ubaidi, M. R., Arshavsky, V. Y. and
Naash, M. I. (2020). Photoreceptor Disc Enclosure Occurs in the Absence of Normal
Peripherin-2/rds Oligomerization. *Front Cell Neurosci* **14**, 92.
Lewis, T. R., Phan, S., Castillo, C. M., Kim, K.-Y., Coppenrath, K., Thomas, W.,
Hao, Y., Skiba, N. P., Horb, M. E., Ellisman, M. H. et al. (2023). Photoreceptor disc incisures
form as an adaptive mechanism ensuring the completion of disc enclosure. *bioRxiv*,
2023.04.06.535932.
Li, C., Ding, X. Q., O'Brien, J., Al-Ubaidi, M. R. and Naash, M. I. (2003). Molecular
characterization of the skate peripherin/rds gene: relationship to its orthologues and paralogues.
Invest Ophthalmol Vis Sci **44**, 2433-41.

Liang, Y., Fotiadis, D., Maeda, T., Maeda, A., Modzelewska, A., Filipek, S.,
Saperstein, D. A., Engel, A. and Palczewski, K. (2004). Rhodopsin signaling and organization
in heterozygote rhodopsin knockout mice. *J Biol Chem* **279**, 48189-96.
Lobanova, E. S., Finkelstein, S., Herrmann, R., Chen, Y. M., Kessler, C., Michaud,
N. A., Trieu, L. H., Strissel, K. J., Burns, M. E. and Arshavsky, V. Y. (2008). Transducin
gamma-subunit sets expression levels of alpha- and beta-subunits and is crucial for rod viability.
J Neurosci **28**, 3510-20.
Lobanova, E. S., Finkelstein, S., Li, J., Travis, A. M., Hao, Y., Klingeborn, M.,
Skiba, N. P., Deshaies, R. J. and Arshavsky, V. Y. (2018). Increased proteasomal activity
supports photoreceptor survival in inherited retinal degeneration. *Nat Commun* **9**, 1738.
Loewen, C. J. and Molday, R. S. (2000). Disulfide-mediated oligomerization of
Peripherin/Rds and Rom-1 in photoreceptor disk membranes. Implications for photoreceptor
outer segment morphogenesis and degeneration. *J Biol Chem* **275**, 5370-8.
Mattapallil, M. J., Wawrousek, E. F., Chan, C. C., Zhao, H., Roychoudhury, J.,
Ferguson, T. A. and Caspi, R. R. (2012). The Rd8 mutation of the Crb1 gene is present in
vendor lines of C57BL/6N mice and embryonic stem cells, and confounds ocular induced mutant
phenotypes. *Invest Ophthalmol Vis Sci* **53**, 2921-7.
Milstein, M. L., Cavanaugh, B. L., Roussey, N. M., Volland, S., Williams, D. S. and
Goldberg, A. F. X. (2020). Multistep peripherin-2/rds self-assembly drives membrane curvature
for outer segment disk architecture and photoreceptor viability. *Proc Natl Acad Sci U S A* **117**,
4400-4410.
Nour, M., Ding, X. Q., Stricker, H., Fliesler, S. J. and Naash, M. I. (2004).
Modulating expression of peripherin/rds in transgenic mice: critical levels and the effect of
overexpression. *Invest Ophthalmol Vis Sci* **45**, 2514-21.
Pearring, J. N., Martínez-Márquez, J., Willer, J. R., Lieu, E. C., Salinas, R. Y. and
Arshavsky, V. Y. (2021). The GARP Domain of the Rod CNG Channel's β1-Subunit Contains
Distinct Sites for Outer Segment Targeting and Connecting to the Photoreceptor Disk Rim. *J*
Neurosci **41**, 3094-3104.
Peeters, M., Khan, M., Rooijakkers, A., Mulders, T., Haer-Wigman, L., Boon, C. J.
F., Klaver, C. C. W., van den Born, L. I., Hoyng, C. B., Cremers, F. P. M. et al. (2021).
PRPH2 mutation update: In silico assessment of 245 reported and 7 novel variants in patients
with retinal disease. *Hum Mutat*.
Pittler, S. J., Keeler, C. E., Sidman, R. L. and Baehr, W. (1993). PCR analysis of
DNA from 70-year-old sections of rodless retina demonstrates identity with the mouse rd defect.
Proc Natl Acad Sci U S A **90**, 9616-9.
Pöge, M., Mahamid, J., Imanishi, S. S., Plitzko, J. M., Palczewski, K. and
Baumeister, W. (2021). Determinants shaping the nanoscale architecture of the mouse rod outer
segment. *bioRxiv*, 2021.08.18.456753.
Reig, C., Martinez-Gimeno, M. and Carballo, M. (2000). A heterozygous novel
C253Y mutation in the highly conserved cysteine residues of ROM1 gene is the cause of retinitis
pigmentosa in a Spanish family? *Hum Mutat* **16**, 278.
Sakuma, H., Inana, G., Murakami, A., Yajima, T., Weleber, R. G., Murphey, W. H.,
Gass, J. D., Hotta, Y., Hayakawa, M., Fujiki, K. et al. (1995). A heterozygous putative null
mutation in ROM1 without a mutation in peripherin/RDS in a family with retinitis pigmentosa.
Genomics **27**, 384-6.

Salinas, R. Y., Pearring, J. N., Ding, J. D., Spencer, W. J., Hao, Y. and Arshavsky, V. Y. (2017). Photoreceptor discs form through peripherin-dependent suppression of ciliary ectosome release. *J Cell Biol* **216**, 1489-1499.

Skiba, N. P., Lewis, T. R., Spencer, W. J., Castillo, C. M., Shevchenko, A. and Arshavsky, V. Y. (2023). Absolute quantification of photoreceptor outer segment proteins. *bioRxiv*, 2023.01.19.524794.

Spencer, W. J., Lewis, T. R., Pearring, J. N. and Arshavsky, V. Y. (2020). Photoreceptor discs: built like ectosomes. *Trends Cell Biol* **30**, 904-915.

Spencer, W. J., Pearring, J. N., Salinas, R. Y., Loiselle, D. R., Skiba, N. P. and Arshavsky, V. Y. (2016). Progressive Rod-Cone Degeneration (PRCD) Protein Requires N-Terminal S-Acylation and Rhodopsin Binding for Photoreceptor Outer Segment Localization and Maintaining Intracellular Stability. *Biochemistry* **55**, 5028-37.

Steinberg, R. H., Fisher, S. K. and Anderson, D. H. (1980). Disc morphogenesis in vertebrate photoreceptors. *J Comp Neurol* **190**, 501-8.

Stricker, H. M., Ding, X. Q., Quiambao, A., Fliesler, S. J. and Naash, M. I. (2005). The Cys214-->Ser mutation in peripherin/rds causes a loss-of-function phenotype in transgenic mice. *Biochem J* **388**, 605-13.

Stuck, M. W., Conley, S. M. and Naash, M. I. (2014). The Y141C knockin mutation in RDS leads to complex phenotypes in the mouse. *Hum Mol Genet* **23**, 6260-74.

Stuck, M. W., Conley, S. M. and Naash, M. I. (2015). Retinal Degeneration Slow (RDS) Glycosylation Plays a Role in Cone Function and in the Regulation of RDS·ROM-1 Protein Complex Formation. *J Biol Chem* **290**, 27901-13.

Stuck, M. W., Conley, S. M. and Naash, M. I. (2016). PRPH2/RDS and ROM-1: Historical context, current views and future considerations. *Prog Retin Eye Res* **52**, 47-63.

Tian, G., Ropelewski, P., Nemet, I., Lee, R., Lodowski, K. H. and Imanishi, Y. (2014). An unconventional secretory pathway mediates the cilia targeting of peripherin/rds. *J Neurosci* **34**, 992-1006.

Travis, G. H., Sutcliffe, J. G. and Bok, D. (1991). The retinal degeneration slow (rds) gene product is a photoreceptor disc membrane-associated glycoprotein. *Neuron* **6**, 61-70.

Volland, S., Hughes, L. C., Kong, C., Burgess, B. L., Linberg, K. A., Luna, G., Zhou, Z. H., Fisher, S. K. and Williams, D. S. (2015). Three-dimensional organization of nascent rod outer segment disk membranes. *Proc Natl Acad Sci U S A* **112**, 14870-5.

Zulliger, R., Conley, S. M., Mwoyosvi, M. L., Al-Ubaidi, M. R. and Naash, M. I. (2018). Oligomerization of Prph2 and Rom1 is essential for photoreceptor outer segment formation. *Hum Mol Genet* **27**, 3507-3518.

Zulliger, R., Conley, S. M., Mwoyosvi, M. L., Stuck, M. W., Azadi, S. and Naash, M. I. (2015). SNAREs Interact with Retinal Degeneration Slow and Rod Outer Segment Membrane Protein-1 during Conventional and Unconventional Outer Segment Targeting. *PLoS One* **10**, e0138508.

660 **Table 1. Quantification of molar ratios between tetraspanins and rhodopsin in WT and**
661 ***Rom1$^{-/-}$* mice**

Protein molar ratios*	WT	*Rom1$^{-/-}$*
PRPH2:rhodopsin	1:18.1 ± 0.5	1:12.2 ± 1.3
ROM1:rhodopsin	1:36.3 ± 3.9	n/a
PRPH2:ROM1	2.0:1 ± 0.3	n/a
(PRPH2 + ROM1):rhodopsin	1:12.1 ± 0.2	1:12.2 ± 1.3

662 *Values are shown as mean ± s.d. n/a: not applicable. Two outer segment preparations from mice
663 of each genotype were analyzed.

Figure 1. Light microscopy of retinas from WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} mice

(**A**) Representative light microscopy images of WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} retinas analyzed at P30. OS: outer segment; ONL: outer nuclear layer. Scale bar: 20 µm. (**B**) Quantification of the number of photoreceptor nuclei in a 100 µm segment of the retina at 500 µm increments away from the optic nerve (ON). Three retinas were analyzed for each genotype. Two-way ANOVA revealed statistically significant differences in the nuclear counts across genotypes (p = 0.0002). Sidak's multiple comparisons post-hoc test revealed statistically significant differences between the total nuclear count between WT and *Rom1*^{-/-} retinas (p = 0.0001) and between *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} retinas (p = 0.0148), but not between WT and *PRPH2 OE/Rom1*^{-/-} retinas (p = 0.3564). (**C**) Quantification of the outer segment (OS) layer length at 500 µm increments away from the optic nerve. Three retinas were analyzed for each genotype. Two-way ANOVA revealed statistically significant differences in the OS layer lengths across genotypes (p < 0.0001). Sidak's multiple comparisons post-hoc test revealed statistically significant differences for the total nuclear count between WT and *Rom1*^{-/-} retinas (p < 0.0001) and between WT and *PRPH2 OE/Rom1*^{-/-} retinas (p = 0.0006), but not between *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} retinas (p = 0.2924).

Figure 2. Ultrastructural analysis of retinas and rod outer segments from WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} mice

(**A**) Representative low magnification TEM images of WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-}
retinas analyzed at P30. RPE: retinal pigment epithelium; OS: outer segment; IS: inner segment.
Scale bar: 10 µm. (**B**) Representative high magnification TEM images of *Rom1*^{-/-} outer segments.
Yellow arrows indicate outer segment structural defects that range from slightly overgrown open
discs to membranous whorls. Scale bar: 1 µm.

Figure 3. Analysis of disc enclosure in rod outer segments of WT, *Rom1⁻/⁻* and *PRPH2 OE/Rom1⁻/⁻* mice

(A) Representative high magnification TEM images of tannic acid/uranyl acetate-stained retinas of WT, *Rom1⁻/⁻* and *PRPH2 OE/Rom1⁻/⁻* mice analyzed at P30. This approach stains newly forming "open" discs more intensely than mature enclosed discs. Yellow arrows point to darkly stained, unenclosed discs; yellow arrowheads point to lightly stained, enclosed discs. Scale bar: 0.5 µm. (B) Quantification of the number of darkly stained open discs at the rod outer segment base. Each data point represents a single outer segment. For each genotype, three retinas were analyzed with at least 35 outer segments analyzed per retina. Data were plotted with samples separated, while statistical analysis was performed on the averages within each retina (n=3 for

each genotype). One-way ANOVA revealed statistically significant differences in the number of
open discs across genotypes (p < 0.0001). Tukey's multiple comparisons post-hoc test revealed
statistically significant differences in the number of open discs between WT and *Rom1*$^{-/-}$ (p <
0.0001) and *Rom1*$^{-/-}$ and *PRPH2 OE/Rom1*$^{-/-}$ (p < 0.0001) mice, but not between WT and *PRPH2*
OE/Rom1$^{-/-}$ mice (p = 0.2686). **(C)** Quantification of the outer segment (OS) diameter. For each
genotype, three retinas were analyzed with at least 66 outer segments analyzed per retina. One-
way ANOVA revealed statistically significant differences in the OS diameters across genotypes
(p = 0.0074). Tukey's multiple comparisons post-hoc test revealed statistically significant
differences in the OS diameters between WT and *Rom1*$^{-/-}$ (p = 0.0083) and *Rom1*$^{-/-}$ and *PRPH2*
OE/Rom1$^{-/-}$ (p = 0.0197) mice, but not between WT and *PRPH2 OE/Rom1*$^{-/-}$ mice (p = 0.7198).
Note that these diameters were measured in longitudinal sections, in which outer segment are not
always sectioned across their widest part; therefore, these values are likely under-representations
of the true OS diameters. However, this does not affect the comparison across genotypes.

Figure 4. Ultrastructural analysis of disc incisures in WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} mice

Representative TEM images of retinas, tangentially sectioned through the outer segment layer, from WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} mice analyzed at P30. Yellow arrowheads indicate incisures observed in WT and *PRPH2 OE/Rom1*^{-/-} discs, but not *Rom1*^{-/-} discs. Scale bar: 1 μm.

Figure 5. Loss of ROM1 alters the oligomerization status of PRPH2

(**A**) Western blot probed for PRPH2 after protein separation by SDS-PAGE under non-reducing conditions. Each sample contained 10 µg of lysate obtained from eyecups of WT or *Rom1*[-/-] mice at P30. Under these conditions, PRPH2 runs as disulfide-bound dimers (~75 kDa) and monomers (~37 kDa). The monomer band of PRPH2 runs as a doublet in *Rom1*[-/-] but not WT eyecups. (**B**) Quantification of the ratio between monomer and disulfide-bound (dimer) bands of PRPH2 was

performed using densitometry of three independent lysates. For the doublet of monomer bands in
Rom1[-/-] lysates, both bands were used for quantification. Unpaired t-test revealed a statistically
significant difference in the PRPH2 monomer:dimer ratio between WT and *Rom1*[-/-] retinas (p =
0.0071). **(C)** Western blot probed for PRPH2 after protein separation by SDS-PAGE under
reducing conditions. Each sample contained 10 µg of lysate obtained from eyecups of WT or
Rom1[-/-] mice. **(D)** Lysates obtained under non-reducing conditions from eyecups of WT and
Rom1[-/-] mice were subjected to velocity sedimentation on 5-20% sucrose gradients. Twelve
fractions were collected with fraction #1 corresponding to 20% sucrose and fraction #12 to 5%
sucrose. Proteins from each fraction were subjected to non-reducing SDS-PAGE and Western
blotting for PRPH2. The distribution of molecular mass standards across fraction, as determined
in (Chakraborty et al., 2008), is shown above the panels. **(E)** Quantification of both the
monomeric and disulfide-bound bands of PRPH2 in each fraction was performed using
densitometry of at least four independent samples and normalized to the total PRPH2 content
across all fractions. Two-way ANOVA revealed statistically significant differences in the
PRPH2 content across genotypes and fractions (p < 0.0001 for disulfide-bound, p = 0.9489 for
monomer). Sidak's multiple comparisons post-hoc test revealed statistically significant
differences between genotypes for the disulfide-bound form in fractions #1 (p = 0.0022), #4 (p =
0.0002) and #5 (p = 0.0061).

Figure 6. The tetraspanin core of ROM1 is sufficient for disc rim formation

(**A**) Cartoon schematic of the RRCT chimeric tetraspanin protein. In the RRCT mouse, the *Prph2* gene has a knockin mutation that replaces it with a DNA sequence encoding a chimeric protein consisting of the tetraspanin body of ROM1 while retaining the C-terminal tail of PRPH2 that is essential for disc formation. (**B**) Representative TEM images of homozygous RRCT mice analyzed at P30. The boxed inset (left) is shown at a higher magnification (right) that reveals the

presence of disc rims. Scale bars: 1 µm (left); 0.1 µm (right). **(C)** Immunofluorescent images of
COS7 cells co-transfected with RRCT-FLAG and ROM1 constructs. Cells were stained with
antibodies against RRCT (red), ROM1 (blue) and calreticulin (green) to label ER membranes.
Nuclei were counterstained with DAPI (grey). Scale bar: 10 µm. **(D)** Lysates obtained under
non-reducing conditions from COS7 cells co-transfected with RRCT-FLAG and ROM1
constructs were subjected to velocity sedimentation on 5-20% sucrose gradients. Twelve
fractions were collected with fraction #1 corresponding to 20% sucrose and fraction #12 to 5%
sucrose. Proteins from each fraction were subjected to reducing SDS-PAGE and Western
blotting for ROM1 and RRCT. **(E)** Quantification of ROM1 and RRCT in each fraction was
performed using densitometry of three independent lysates and normalized to the total content
across all fractions.

Figure 7. Disc rims can be formed in the absence of intermolecular disulfide bonds between PRPH2 and ROM1

(A) Representative TEM images of tannic acid/uranyl acetate-stained retinas of *Prph2$^{C150S/C150S}$* and compound *Prph2$^{C150S/C150S}$/Rom1$^{-/-}$* mice analyzed at P30. Photoreceptor outer segments of each mouse have severely perturbed outer segment structure. Yet, disc enclosure is not completely prevented as there are both darkly-stained, "open" discs (yellow arrows) and lightly-stained, "enclosed" discs (yellow arrowheads) in each genotype. The boxed region of each image depicts an area in which disc stacking appears relatively normal and permits the analysis of disc rim structure. Scale bar: 1 µm. **(B)** Higher magnification images of the regions in which disc stacking appears normal in both *Prph2$^{C150S/C150S}$* and *Prph2$^{C150S/C150S}$/Rom1$^{-/-}$* mice. Disc rims are formed in each genotype. Scale bar: 0.1 µm.

 Supplementary Material

Supplementary Figure 1. Ultrastructural analysis of WT and *Rom1*^{-/-} retinas

Representative low magnification TEM images of WT and *Rom1*^{-/-} retinas contrasted with the conventionally used osmium tetroxide. RPE: retinal pigment epithelium; OS: outer segment; IS: inner segment. Scale bar: 10 µm.

Supplementary Table 1. Quantification of molar ratios between tetraspanins and rhodopsin in WT and *Rom1*^{-/-} mice – raw data