Domain-general and language-specific contributions to speech production in L2: an fMRI study using functional localizers.

Agata Wolna[1*], Jakub Szewczyk[2], Michele Diaz[3], Aleksandra Domagalik[4],

Marcin Szwed[1], Zofia Wodniecka[1*]

[1] Institute of Psychology, Jagiellonian University, Kraków, Poland
[2] Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands; Donders Institute for Brain Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands
[3] Social, Life, and Engineering Sciences Imaging Center, the Pennsylvania State University, Pennsylvania, United States
[4] Centre for Brain Research, Jagiellonian University, Kraków, Poland

Corresponding authors:
zofia.wodniecka@uj.edu.pl
agata.wolna@doctoral.uj.edu.pl

Language and Bilingualism Lab
Instytut Psychologii UJ
ul. Ingardena 6
30-060 Kraków

Abstract

For bilinguals, speaking in a second language (L2) compared to the native language (L1) is usually more difficult. In this study we asked whether the difficulty in L2 production reflects increased demands imposed on domain-general or core language mechanisms. We compared the brain response to speech production in L1 and L2 within two functionally-defined networks in the brain: the Multiple Demand (MD) network and the Language network. We found that speech production in L2 was linked to a widespread increase of brain activity in the domain-general MD network. The Language network did not show a similarly robust differences in processing speech in the two languages, however, we found increased response to L2 production in the language-specific portion of the left IFG. To further explore our results, we have looked at domain-general and language-specific response within the brain structures postulated to form a BILINGUAL LANGUAGE CONTROL network. Within this network, we found a robust increase in response to L2 in the domain-general, but also in some language-specific voxels including in the left IFG. Our findings show that L2 production strongly engages domain-general mechanisms, but only locally affects language-specific processing. These results put constrain on the current model of bilingual language control.

Introduction

Bilinguals often struggle with speaking in their second language (L2) compared to their native language (L1), even when they have a good level of proficiency. This difficulty is reflected in increased naming times and decreased accuracy in tasks requiring participants to name pictures of simple objects in L1 and L2 [1–5]. Similarly, neuroimaging research using *EEG* [6,7] and *fMRI* [8–13] consistently shows that speech production in L2 is linked to increased engagement of neural resources.

In this study, we investigated the neural correlates of L1 and L2 speech production to gain a better understanding of the source of the difficulty accompanying speech production in L2. Specifically, we asked whether production in L2 entails increased engagement of domain-general control resources, the core language processes, or both. Even though brain mechanisms engaged in speech production in L1 and L2 have been studied before, we are still lacking a full understanding of the mechanisms that contribute to speech production in the non-native language. One of the central axes of this debate is whether the increased effort related to speaking in L2 can be traced back to increased engagement of domain-general or language-specific processes. However, addressing this problem is a non-trivial task as the assumptions of the theoretical models of bilingual speech production as well as the tools used to interpret the brain responses to L2 and L1 do not clearly disentangle the language-specific and domain-general mechanisms engaged in speech production. Speech production in both L2 and L1 needs to rely on language-specific processes (conceptual preparation, lexical access, phonological encoding, articulation), however differences in the brain response to naming pictures in L2 and L1 usually have been attributed to increased engagement of cognitive control, not core language mechanisms. This interpretation draws on the assumption of theoretical models of bilingual speech production that speaking in a weaker language requires increased engagement of bilingual language control mechanisms that modulate the activation of the two languages and adjust them to the task and contextual demands at hand [14–16]. The problem is that from this perspective, all differences in neural activity between speech production in L2 and L1 should be interpreted as increased engagement of the control mechanisms. Therefore, this approach can easily overlook differences in language-specific mechanisms supporting the processing of L1 and L2, as the theoretical framework of bilingual language production does not differentiate between the domain-general and language-specific components of the postulated control mechanisms. In the current paper, we explored the brain response to speech production in L1 and L2 using an analytical approach in which functional networks are defined independently of the main task. This approach precisely disentangles the domain-general and language-specific components of bilingual speech production.

Speech production in L2 as a difficulty for domain-general control mechanisms

Many studies have linked the difficulty related to speech production in L2 to increased engagement of the domain-general control mechanisms [15,17–20]. The cognitive processes orchestrated

by the bilingual language control network largely overlap with those attributed to the Multiple Demand (MD) network [21] in both their functional definitions and anatomical localization in the brain. The MD network is a domain-general fronto-parietal network that supports a wide range of executive control mechanisms [22], working memory [23], fluid intelligence and problem solving [24]. The MD network also engages in the execution of complex and non-automated tasks [23,25,26] such that its activation closely reflects the difficulty of a given task [23]. This network is characterized by broad domain-generality [27,28] and can be simultaneously engaged in many aspects of a given task, including sensory (visual, auditory and motor) demands and rule complexity[27]. It has been suggested that the role of the MD network is to integrate information that is relevant to a current cognitive operation [29,30]. Importantly, although the MD network is sensitive to integration across a broad array of tasks, it is not strongly engaged by basic language processes (e.g., sentence comprehension [23,31]) Since L2 is typically less automatized than L1, the sensitivity of the MD network to task automatization, makes it a suitable candidate for supporting the additional demands related to producing speech in L2. This possibility is further corroborated by the topographical overlap between the MD network and structures that form the bilingual language control network.

Speech production in L2 as a difficulty for core language mechanisms

In addition to increased task demands, the difficulty related to speaking in L2 may reflect a language-processing problem and entail increased demands for language-specific computations in the brain which are supported by the language network: a fronto-temporal neural network specifically tuned to process language [31,32]. The Language network supports representational and computational aspects of language processing. It is engaged in language comprehension [32,33] and production [34] across different modalities and tasks. It supports language processing at the sub-lexical [35], lexical, and syntactic levels [36–38], it is also sensitive to the difficulty of language processing [39,40] and responds to differences between comprehension in native and non-native languages in polyglots [41]. What is more, the language network was shown to be sensitive to lexical access demands [34], and lexical access difficulty related to speaking in the L2 compared to the native language [42]. As such, it is plausible to assume that speech processing in L2 may present a language processing difficulty that is handled by the Language network.

What do we know about the neural basis of speech production in L2?

Neuroimaging research has shown that production in L2 compared to L1 engages a wide network of bilateral cortical and subcortical brain structures including the left and right prefrontal cortex, dorsal anterior cingulate cortex and pre-supplementary motor area, left and right inferior parietal lobe, cerebellum, thalamus as well as basal ganglia [15,20]. Abutalebi and Green [15] proposed that these structures form the *bilingual language control* network that supports a variety of processes necessary for a bilingual speaker to efficiently use two languages. The bilingual language control

network has been identified predominantly based on studies using the language switching tasks (LST [8,9,17,43–47]; for metaanalyses see [13,48]). However, these findings may be difficult to interpret as the LST captures brain activations underlying speech production in L1 and L2 in a situation requiring the speakers to switch between languages very frequently (e.g., every few seconds), which is not typically experienced in everyday life. Such frequent switches may engage cognitive control to a much higher degree than language production in more naturalistic situations. Specifically, frequent switching between the two languages may lead to increased brain response that not only reflects the engagement of neural networks supporting the L1 and L2 but also the mechanisms that support execution of difficult, non-automated tasks [25,26]. Finally, picture naming in L1 is usually significantly slowed down in the LST, which is interpreted as a consequence of rebalancing activation levels of L1 and L2 (the *reverse dominance* phenomenon [49]). As such, comparison of brain responses to picture naming in L1 and L2 in the LST may reflect artificial task demands that may even reverse language dominance due to switching between languages, rather than mechanisms that support production in L1 and L2 in a less demanding, single language context.

Then, it is not surprising that studies comparing language production in L1 vs L2 in single-language context show a different pattern of brain activity than those identified using the LST [13]. This discrepancy in results is also consistent with predictions of the adaptive control hypothesis of bilingual language control [18] which assumes that cognitive control resources are flexibly adjusted to the communicative context and therefore they may be differentially engaged in different tasks. For these reasons, it is essential to examine the brain correlates of bilingual speech production in a single language context. So far, several studies have used this approach [9–12,33,44,50–52]. However, their results do not converge on a set of brain regions responding more strongly to speech production in L2 compared to L1 (see **Table 1** and **Figure 1**). While the brain regions identified in these studies mostly pertain to the bilingual language control network [15], there is little overlap in peaks of activation across the studies. What is more, because of a high individual variability in the precise topography of functional networks supporting high-level cognitive processes [53,54], it is difficult to say the activations observed in the previous experiments fall within the language or the MD networks (this is especially problematic for activations falling within brain regions where language and MD networks are adjacent to each other).

Table 1. Summary of studies looking at L2 > L1 contrasts in the single language context. The table includes all available studies that tested and reported a direct comparison between L2 and L1. It includes the characteristics of the participants as well as a list of structures that were significantly activated for the L2 > L1 contrast.

Study	N	L1	L2	AoA	Balance (proficiency)	Stronger response to L2 > L1
De Bleser et al., 2003	11	Flemish	French	10	unbalanced	left orbito-frontal gyrus, cerebellum

Hernandez et al., 2009	12	Spanish	English	<5	unbalanced (more proficient in L2 based on the Boston Naming Test)	right postcentral gyrus, superior temporal gyrus, hippocampus, insula and pre-SMA
Liu et al., 2010	24	Chinese	English	12	unbalanced	left IFG, bilateral SMA, left precentral gyrus, left lingual gyrus, left cuneus, bilateral putamen, bilateral globus pallidus, bilateral cerebella
Videsott et al., 2010*	20	Ladin	Italian	5	balanced	right MFG
Fu et al., 2017	21	Chinese	English	10	unbalanced	left IPL, left cerebellum
Rossi et al., 2018	25	English	Spanish	late	unbalanced	left lateral occipital cortex, right fusiform and right posterior inferior temporal gyrus, left caudate, right postcentral and supramarginal gyri, and left MFG
Wang et al., 2020	27	Chinese	English	7,27	unbalanced	left and right IFG, left and right ACC, left and right insula, left and right Fusiform gyrus, left calcarine gyrus, left cerebellum

* Videsott et al., 2010 tested speech production in multilinguals: participants were also proficient in German (L3) and English (L4). In this table we only report the results of the L2 > L1 comparison, as it fits best the scope of this summary.

Figure 1. Peak activations corresponding to the L2 > L1 contrast in previous studies looking at differences between languages in single-language context. Data was plotted using a 5mm radius around each of the peaks reported in the papers. Rossi et al., 2018 did not report the exact peak locations for L2 > L1 contrast, for this reason this study is not included in the figure.

Lateralization of mechanisms supporting speech production in L1 and L2

Another perspective at looking at the brain basis of bilingual speech production is to ask whether within the Language and MD networks L1 and L2 engage similar or different regions. In this context, an interesting yet unanswered question related to the brain basis of speech production in L1 and L2 is the degree of lateralization. It is known that language is lateralized to the left hemisphere in most people, but research suggest that bilinguals may have reduced lateralization, especially if they acquired the L2 early in life [55–58]. However, so far, we know very little about differences in lateralization of L1 and L2 processing. Here we ask two questions related to lateralization of bilingual speech production. First, whether within the Language network L1 and L2 show a similar degree of lateralization and second, whether any differences in lateralization of L1 and L2 processing reflect differences in lateralization of language-specific processes or mechanisms supporting the domain-general control.

Current study

        In the current study we asked whether the brain's solution to processing L2 rely on domain-general control processes or on language specific mechanisms. More specifically, we explored the following questions: Is speech production in L2 linked to increased engagement of domain-general control regions? Is speech production in L2 linked to increased engagement of language-processing regions? Is there a difference in lateralization supporting L1 and L2 production? Additionally, to link our results to the neurocognitive model of bilingual language control [15], we explored brain activations during L1 and L2 speech production within the language-and domain-general selective areas forming the bilingual language control network. To address these questions, we used a precision fMRI approach based on functional localization [32]. To evaluate the brain response to speech production in L2 and L1 within the Language and MD networks, we created a set of subject-specific functional ROIs within each network and estimated the response within these fROIs to the critical conditions: speech production in L2 and L1. Using a similar approach, we created language and domain-general specific fROIs within each brain structure of the bilingual language control network (selected based on [15]; for details see Methods) and estimated responses to speech production in the L2 and L1 within each of the fROIs. A key advantage of this approach is that it precisely identifies functional networks for each subject separately which allows for better modelling of the individual differences in cortical organization of the functional networks in the brain. It also provides a straightforward functional interpretation of the observed brain activity without referring to the function assigned to a particular brain region based on its anatomical label (*reverse inference* problem [59]). Finally, using functional localizers increases the power of statistical tests as the comparisons between L1 and L2 are computed in a narrow set of ROIs instead of thousands of voxels in the entire brain. The increase in statistical power may be particularly important because some of the discrepancies in previous studies' results may be due to low statistical power.

Results
Behavioural results

        The analysis of naming latencies revealed that participants were significantly faster in L1 compared to L2. We also found a significant effect of the *Session* showing that participants were overall slower in the 2nd session. Importantly, the effect of *Session* did not interact with the effect of *Language*. The analysis of accuracy yielded very similar results: participants were more accurate in their L1 than L2, they were also less accurate in the 2nd session and there was no interaction between the two effects. The results of the behavioural analyses are presented in **Table 2.**

Table 2. Behavioural performance in Picture Naming task. The table presents results of linear mixed models predicting naming latencies and accuracy of naming pictures.

	Naming latencies				
Predictors	*Estimates*	*std. Error*	*CI*	*t*-value	*p* – value

Intercept	7.02	0.01	6.99 7.05	518.52	**<0.001**
Language	0.12	0.01	0.10 0.15	8.72	**<0.001**
Session	0.05	0.01	0.02 0.07	3.56	**<0.001**
Language x Session	0.01	0.05	-0.08 0.11	0.29	0.773

Accuracy					
Predictors	*Odds Ratio*	*std. Error*	*CI*	*Statistic*	*p*
Intercept	12.83	2.30	9.04 18.22	14.26	**<0.001**
Condition	0.07	0.02	0.04 0.13	-8.39	**<0.001**
Session	0.52	0.11	0.34 0.79	-3.04	**0.002**
Condition x session	0.74	0.33	0.30 1.78	-0.68	0.498

Neuroimaging results: localizer approach

In the Language network we did not observe a significant effect of *Language* or *Hemisphere*, however, we found a significant interaction between *Language* and *Hemisphere*. Pairwise comparisons revealed that the brain response to speaking in L1 was stronger compared to L2 in the right hemisphere (b = 0.09, t = 2.11, p = .038) but not different from L2 in the left hemisphere (b = -0.04, t = -0.94, p = .350). In the MD network, we found a significant effect of *Language* showing stronger responses to L2 than L1 production and *Hemisphere* showing stronger responses in the left than right hemisphere ROIs. Interestingly, we also found an interaction between *Language* and *Hemisphere*. Pairwise comparisons revealed that the response to L2 was stronger than to L1 in the left hemisphere (b = -0.187, t = -5.138, p < .001) but there were no differences between languages the right hemisphere (b = -0.04, t = -0.994, p = .325). The results of the localizer analyses are presented in **Table 3 and Figure 2** (see Supplementary Materials for individual functional ROI (fROI) results).

Table 3. Estimated brain response in Language and MD Network. The table presents results of linear mixed models predicting the % BOLD signal change in the two networks.

Language network					
Predictors	*Estimates*	*std. Error*	*CI*	*t*-value	*p* – value
Intercept	0.56	0.11	0.34 0.78	4.95	<0.001
Language (L2 vs L1)	-0.02	0.03	-0.09 0.04	-0.70	0.486
Hemisphere (left vs right)	-0.13	0.21	-0.54 0.28	-0.62	0.537
Language x Hemisphere	-0.12	0.04	-0.21 -0.04	-2.84	0.005
MD network					
Predictors	*Estimates*	*std. Error*	*CI*	*Statistic*	*p*
Intercept	0.46	0.06	0.34 0.58	7.43	<0.001

Language	0.11	0.03	0.05	0.18	3.35	0.001
Hemisphere	-0.19	0.09	-0.37	-0.02	-2.13	0.033
Language x Hemisphere	-0.15	0.03	-0.21	-0.09	-5.13	<0.001

Figure 2. Brain response to speech production in L1 and L2 within the MD and Language networks. The plot presents estimates of % signal change predicted by the statistical models. Panels A) and C) present data for estimates to L1 and L2 within the MD and Language networks averaged by hemisphere. Individual data points in these panels correspond to predicted responses of individual subjects; and panels B) and D) represent means for naming in L1 and L2 for each ROI. Whiskers in these panels represent Confidence Intervals for post-hoc pairwise comparisons, and significant differences between conditions for a given ROI are marked with asterisks.

Neuroimaging results: bilingual language control network

The first step of the analysis of responses to L2 and L1 speech production in the language- and domain-general selective fROIs within the bilingual language control network, was to establish the functional selectivity of the fROIs to the language and MD localizer contrasts as well as the overlap between the two types of fROIs within each of the anatomically defined ROIs forming part of the bilingual language control network (henceforth: BLC network ROIs). In the second step, we used the

fROIs which have shown robust responses to the localizer contrasts and evaluated the estimates of brain activity corresponding to L2 and L1 speech production within these fROIs (for details of the analytical approach see Methods).

To assess the functional selectivity of language and MD fROIs within the bilingual language control network ROIs we have first evaluated responses to the localizer contrasts within both language and MD fROIs. Full results of these analyses are presented in the Supplementary Materials. Among the MD fROIs, we observed a robust response to the MD localizer contrasts (hard > easy visual working memory task) in all fROIs except for the left putamen (p = .123). The MD fROIs were also found to be highly selective to the domain-general task (i.e., we did not observe significant responses to the language localizer contrast in any of them). For the language fROIs, we observed significant response to the language localizer contrast (listening to intact > degraded speech) in the left angular gyrus, left IFG (pars triangularis and opercularis), left MFG, left pre-SMA and right IFG (pars triangularis and opercularis). What is more, the left MFG fROI was sensitive to not only the language localizer contrast but also to the MD contrast, which means it is not specifically-selective for language processing. We also found that the overlap between language- and domain-general selective fROIs (in the BLC network ROIs for which both types of fROIs were successfully identified) was very small – the average overlap at the voxel level was not larger than 10% (see Figure 3).

In the next step, the language and MD fROIs that showed a significant response to their localizer contrasts were used to evaluate the brain response to L2 and L1 speech production. Response to L2 was significantly stronger than to L1 in 4 language fROIs and 7 MD fROIs. In the language-selective fROIs we found significant effects in the left IFG (pars triangularis and opercularis), left MFG and left pre-SMA. In the MD-selective fROIs we found increased response to L2 compared to L1 in the left ACC and left pre-SMA, left IFG (pars opercularis and triangularis), left MFG, and left caudate. The results of this analysis are presented in Table 4 and Figure 3.

Table 4. Difference between estimated brain response to speech production in L2 and L1 within language and MD fROIs in the Bilingual Language Control network ROIs. The table presents results of linear mixed models predicting the % BOLD signal change corresponding to L2 > L1 difference in each fROI.

Bilingual Language Control network

Localizer	ROI	Estimates	std.Error	t-value	effect size	p-value uncorrected	fdr corrected
Language	LeftIFGtr	0,12	0,04	2,84	0,90	0,007	0,012*
	LeftIFGoper	0,16	0,05	3,29	1,04	0,002	0,007*
	LeftMFG	0,13	0,04	3,49	1,10	0,001	0,007*
	LeftpreSMC	0,12	0,04	3,09	0,98	0,004	0,008*
	RightAngularG	-0,12	0,05	-2,37	-0,75	0,022	0,031*
	RightIFGoper	0,00	0,04	-0,04	-0,01	0,964	0,964

	RightIFGtr	-0,06	0,05	-1,16	-0,37	0,255	0,297
	LeftIFGtr	0,19	0,05	3,63	1,15	0,001	0,002*
	LeftIFGoper	0,19	0,05	3,79	1,20	0,000	0,002*
	LeftMFG	0,21	0,05	4,52	1,43	0,000	0,001*
	LeftAngularG	-0,09	0,04	-2,14	-0,68	0,039	0,053
	LeftpreSMC	0,10	0,04	2,87	0,91	0,007	0,012*
MD	LeftACC	0,08	0,03	2,45	0,78	0,019	0,029*
	LeftThalamus	0,05	0,04	1,06	0,34	0,294	0,323
	LeftCaudate	0,20	0,05	4,12	1,30	0,000	0,001*
	RightIFGtr	-0,07	0,05	-1,41	-0,45	0,166	0,203
	RightIFGoper	-0,01	0,04	-0,34	-0,11	0,735	0,735
	RightAngularG	-0,15	0,05	-3,07	-0,96	0,004	0,008*

Figure 3. Brain response to speech production in L1 and L2 within the bilingual language control network ROIs. Each panel presents the estimates for speech production in L2 and L1 in the language-specific fROIs (red) and MD-specific fROIs

(blue) within a given BLC ROI. For ROIs in which we identified both language- and MD-selective parcels, we also show the % of the voxels overlapping between the two fROIs.

Discussion

In this paper we explored the brain basis of the difficulty accompanying speech production in L2 and L1. Although some previous research has attempted to explore this issue before, there are several strengths and novelties of our contribution. First, we used newer analytical tools – the precision fMRI method based on functional localizers [32,60]. More specifically, we tested the overlap of brain responses to speech production in L2 and L1 with two functional networks in the brain: the Language network, which supports core language processes [32] and the MD network, which supports a wide range of domain-general control mechanisms [21,28,32]. The second novelty and strength of our contribution is that, unlike most of the previous studies - we have focused on speech production in the single language context – i.e., we used a task that does not require participants to frequently switch between languages. As such, our findings allow making conclusions about mechanisms of pure L2 vs. L1 production, not confounded with the additional language switching requirement. Finally, our study is very well-powered, based on a sample considerably larger (n=41) than any of the previous experiments looking at the same research question.

Using functional localizers, we have shown that difficulty in speech production in L2 is linked to robust activation in areas delineated by the MD localizer, suggesting that L2 leads to a greater recruitment of domain-general resources compared to L1. The language network did not show similarly widespread and consistently stronger responses to L2, except for the left inferior frontal gyrus (IFG) fROI.

The increased engagement of the MD network in speech production in L2 shows that bilingual language control is achieved by domain-general mechanisms. As indicated earlier, many of the previous studies evaluating and corroborating the assumption of domain-generality of the bilingual language control used the language switching task, which requires frequent switching between languages. Here, we show that in a single language context, production in L2 also draws on domain-general resources. Specifically, compared to L1, speech production in L2 was linked to widespread engagement of the left MD network. This is a clear indication that the difficulty in non-native speech production is not driven by an isolated control mechanism constrained to a single specific process, but instead, it engages a complex neural network. Nonetheless, based on the results of the analysis within the MD network, it is difficult to tell *which* exact domain-general mechanisms are more engaged in the L2 production compared to L1.

Within the core Language network, speech production in L2 was not linked to a consistent, network-wide increase in activation compared to production in L1. This result is in line with findings showing substantial overlap between brain representations of L1 and L2 [61–65]. It suggests that

language-specific processes are largely shared between languages and the most fundamental difference between bilingual and monolingual language processing is not the "language processing" itself. Importantly, we found no differences in how L1 and L2 speech production engaged the left temporal cortices, which have been linked to lexical access [66]. A difficulty at this stage of production has oftentimes been proposed as a locus of differences between L1 and L2 production [8,10,67]. At the same time, even though the Language network did not differ in its overall response to speech production in both languages, we found increased response to L2 compared to L1 in the language-specific left IFG fROI.

The left IFG has long been considered one of the key brain regions supporting language processing (in particular the articulation, cf. Broca's area). Activity in this structure has been linked to post-lexical compositional processes, such as phonetic, encoding, syllabification, or even later processes related to articulatory planning [66,68–70]. In the light of debate on the role of the left IFG in the language and domain-general processes as well as its contribution to controlling the two languages of a bilingual, it is particularly interesting that the increased activation in response to L2 compared to L1 was found in both language-specific and domain-general fROIs within the posterior part of left IFG (pars opercularis). This specific area is hypothesised to play a role of an interface between language representations and articulatory codes executed by the motor cortices [71]. At the same time the left IFG has been suggested to play a crucial role in general selection among the competing representations [72,73]. While this hypothesis was first put forward based on studies focusing on language processing, the engagement of the left IFG in selection of representations has been shown to extend to non-linguistic domains [74]. Drawing on the selection hypothesis, the bilingual language control model proposes that the left IFG is a crucial node supporting the bilingual language control [15,75].

Importantly, the left IFG, has been shown to be a heterogeneous structure, encompassing specialised language-specific and domain-general subregions [75,76]. Our data confirm this dissociation: language-specific and domain-general fROIs showed a minimal overlap at a subject-level even if the two functional subsections were identified in the same anatomical masks in the BLC network analysis (see voxel overlap plots for the left IFG triangularis and opercularis on Figure 3). On the one hand, our results may indicate that the domain-general and language-specific subsections of the left IFG closely cooperate to select appropriate words for production. On the other hand, however, the Language and MD networks were shown to support very different types of processes [76], so it is plausible that the increase in the domain-general and language-specific sub-sections of the left IFG in response to production in L2 reflects two different types of mechanisms. Given the functional characterization of the left IFG discussed above, we would like to propose that the observed increase in activation of the language-specific sub-section of the left IFG in response to speech production in L2 compared to L1 reflects increased difficulty in phonological encoding and articulatory processing [1,4].

Our results also show that compared to L1 production, L2 production recruits the left- hemisphere to a greater extent in both networks. In the Language network, compared to L1, speech production in L2 engaged more resources in the left hemisphere but fewer resources in the right hemisphere. While the left hemisphere is crucial for formulating propositional speech, the right hemisphere supplements it by controlling prosodic and non-propositional components of language [77]. The right hemisphere is also more engaged in automated speech (e.g., reciting months [78]). As such, the stronger engagement of the right hemisphere in the native than non-native language production may be driven by higher automatization of L1 compared to L2, especially in bilinguals who learned the L2 later in life. In the MD network, when speaking in L2 compared to L1 we found stronger activity in the left hemisphere, with no such differences in the right hemisphere. Interestingly, the differences in lateralization were much more pronounced in the MD network than in the Language network. This result may indicate that the stronger lateralization of L2, does not necessarily reflect the lateralization of language processes themselves but rather the lateralization of domain-general control mechanisms supporting speech production in L2.

The results discussed above provide a characterization of bilingual speech production mechanisms in relation to two well-defined functional systems in the brain. However, they did not allow to directly link our results to the most influential neurocognitive model of bilingual language control (BLC)[15]. In order to do so, we ran an additional analysis using the functional localizer approach in which we looked at the response to speech production in L2 and L1 within the language- and domain-general selective fROIs within the postulated BLC network regions. We found domain-general selective fROIs in almost all nodes of the BLC network (except for the left putamen), but not all of them responded to the L2 > L1 contrast: significant effects were found only in the left IFG, left MFG, left pre-SMA/ACC, left thalamus and left caudate. In the remaining fROIs we either observed a de-activation in response to L2 (left and right angular gyrus) or no difference between languages (right IFG triangularis and opercularis). As such, although the analyses within the BLC network ROIs largely confirm the claims of the neurocognitive model of bilingual language control, they also highlight some important differences. We believe that the differences between the model's predictions and our findings (e.g., for instance, no differences observed between L2 and L1 in the right IFG) can be reconciled by the fact that the BLC model was largely based on studies using the language switching paradigm, whereas in our study, participants named pictures in a single-language context. Since the right IFG is a structure implicated in cue detection and inhibition of non-target responses [15,18] and inhibition is postulated to be one of the key mechanisms of bilingual language control [14,15,18], the absence of differences in this structure in our study suggests that during production in a single language context, inhibition is not necessarily engaged to the same degree as in contexts requiring frequent switches between languages. At the same time, it has been claimed that bilingual language control in speech production is achieved   by a re-configuration of the domain-general network: it

draws on the same structures as the ones engaged in domain-general tasks, however, it adjusts the connections and engagement of the particular nodes to the task at hand [20]. Our data partially confirm this claim: we did find increased activation to L2 compared to L1 in most of the domain-general fROIs. However, we have also identified language-specific fROIs within the BLC network that responded more strongly to L2 compared to L1. As discussed above, overall, this result suggests that both domain-general and language-specific mechanisms cooperate to select appropriate words for production. With methods relying on group averaging, this spatially fine-grained distinction into the two networks may be easily overlooked.

To conclude, our study provides a characterization of brain mechanisms supporting speech production in L2 and L1. We used a methodological approach that allowed us to precisely disentangle the contribution of domain-general and language-specific mechanisms in the process of bilingual speech production. We found that the difficulty in speech production in L2, compared to L1, was linked to increased engagement of a wide range of domain-general mechanisms in the MD network, many of which coincided with brain structures postulated to form the bilingual language control network. Importantly, the absence of additional activation of the right IFG suggests that if each language is used separately (i.e., without the immediate need to switch them), language inhibition may not be required. However, production in L2 was also linked to increased engagement of language-specific resources in the left IFG which may reflect increased difficulty in phonological encoding and articulatory processing. What is more, we found that in both networks the L2 engaged resources lateralized to the left hemisphere, which suggests that the lateralization of language processing in bilinguals may in fact reflect lateralization of the control mechanisms.

Method

Participants

This study reports the results of data collected from the same group of subjects as the one used by Wolna et al [79]. Forty-two Polish-English late bilinguals took part in the study. One participant was excluded due to the excessive head motion (>2mm) in the main fMRI task resulting in a final sample of forty-one participants (31 females, 10 males, mean age = 23.29, SD = 3.24). All participants were Polish native speakers who learned English as their second language. They declared intermediate or advanced proficiency in English (B2 or higher) and obtain at least 20/25 points in the General English Test by Cambridge Assessment (mean score = 21,32, SD = 1,94). Their proficiency in English was also assessed again using LexTale [80] (mean score = 71,58%, SD = 9,91%). Detailed information on the participants' proficiency, age of acquisition and daily use of the languages (based on a language questionnaire) can be found in Table 5.

All participants gave written consent for participation in the study. The study was approved by the Ethics Committee of Jagiellonian University Institute of Psychology concerning experimental studies with human subjects.

Study Design

Each participant completed a series of behavioural tasks to assess their language proficiency as well as a detailed language questionnaire. Details of these tasks are described in Wolna et al., 2023, and their results are summarized in Table 5. In the scanner, each subject completed a main task (picture naming in L2 and L1) and the Language and Multiple Demand network localizers. To keep the language context under control, the main task was split between two sessions: in one of them, subjects completed the picture naming task in L2 and in another, in L1. Several additional tasks were included in the experiment, however, they're beyond the scope of this paper.

Table 5. Language experience of participants. Information on self-rated proficiency, language learning and daily use of languages is given for all languages that participants declared to know. L1 always refers to Polish and L2 to English. L3 and L4 were a variety of different languages (incl. German, French, Italian, Spanish, Russian, Czech, Japanese, Korean, Norwegian, Latin and Esperanto). Not all the participants reported to know an L3 or L4.

Language	**L1** (Polish)	**L2** (English)	**L3** (various)	**L4** (various)
n	41	41	36	17
Self-rated proficiency (1-10)				
reading	9,72	4,34	2,76	1,79
listening	9,62	4,10	2,45	1,84
writing	9,26	3,81	2,21	1,48
speaking	9,49	3,89	2,48	1,68
accent	9,56	3,45	2,45	1,65
Language learning				
Age of acquisition onset of learning	0,00	6,44	14,34	15,24
Age of acquisition: onset of using	3,17	9,15	15,41	16,88
Years of formal education	14,72	11,56	4,03	4,18
Daily use of languages				
passive	63,26%	31,79%	4,03%	3,25%
active	81,31%	16,31%	2,06%	1,31%
Verbal Fluency (mean number of words produced)				
category	20,04	13,36	-	-
letter	12,64	10,62	-	-

Main task: picture naming. In the main task, participants were asked to overtly name pictures of objects. In order to in order to minimize effects of language switching, which were not of theoretical interest in this study, participants named pictures in L1 and L2 on two separate sessions (on different days). The order of sessions was counterbalanced between participants. In each session, participants

named one block of pictures corresponded to one functional run. Each block contained 55 coloured pictures representing objects (from the CLT database [81]). Pictures were not repeated between blocks. Each block started with a fixation cross ("+") displayed on a screen for 7000 ms. Subsequently, 55 pictures were presented following a rapid event-related design. Each picture was displayed on a white background for 2000 ms and was followed by a fixation cross jittered at an interstimulus interval ranging from 607 ms to 10144 ms (M = 3091 ms, SD = 2225 ms). The order of picture presentation and interstimulus intervals was optimized using optseq2 [82]. Finally, at the end of each block, a fixation cross was presented for 8000 ms. Overall, each block of the Picture Naming task lasted 295s.

Language localizer. Participants listened to fragments of Alice in Wonderland in L1 and distorted speech recordings in which it was impossible to understand what the speaker was saying. The task followed the design introduced by Scott et al. [33] in which the localizer contrast was based on the intact > distorted speech condition contrast. In each functional run, participants listened to 6 short passages of Alice in Wonderland in their native language (18s each) and 6 passages of distorted speech (18s each). Additionally, 4 fixation blocks (12s each) were included in each run. The total duration of each run was 264s. Each participant completed 2 runs of the task.

Multiple Demand localizer. Participants were asked to perform a spatial working memory task [23,31]. The design of this task followed the one used by Fedorenko et al. [31] in which the localizer contrast was based on the hard > easy condition contrast. In each trial, participants saw four 3 x 4 grids and their task was to memorize the locations of black fields in each grid. In the easy condition, they had to memorize one location per grid (four in total) and in the hard condition, they had to keep track of two locations per grid (eight in total). At the end of each trial, participants had to choose a grid with all the memorized locations, from among two alternatives presented on the screen. Each trial started with a fixation cross displayed on a screen for 500 ms and was followed by four grids, each displayed for 1000 ms. After that, the choice task appeared on the screen until a response was given (for a max of 3750 ms) followed by feedback (right or wrong) displayed for 250 ms and a fixation cross which was displayed for 3750 ms minus the reaction time in the choice task. In each of the conditions – easy and hard – participants completed 4 trials (34s in total). Each run contained five easy condition blocks and five hard condition blocks. Additionally, six fixation blocks (16s) were included in each run. The order of experimental and fixation blocks was counterbalanced between participants and runs (four counterbalance lists were created and each participant completed two of them). The total duration of one run was 438s.

MRI Data Acquisition

MRI data were acquired using a 3T scanner (Magnetom Skyra, Siemens) with a 64-channel head coil. High-resolution, whole-brain anatomical images were acquired using a T1-MPRAGE sequence (208 sagittal slices; voxel size 0.9×0.9×0.9 mm3; TR = 1800 ms, TE = 2.32 ms, flip angle = 8°). A gradient field map was acquired with a dual-echo gradient-echo sequence, matched spatially with fMRI scans

(TE1 = 4.92 ms, TE2 = 7.38 ms, TR = 503 ms, flip angle = 60°). Functional T2*-weighted images were acquired using a whole-brain echo-planar (EPI) pulse sequence (50 axial slices, 3x3x3 mm isotropic voxels; TR = 1400 ms; TE = 27 ms; flip angle = 70°; MB acceleration factor 2; in-plane GRAPPA acceleration factor 2 and phase encoding A>>P) using 2D multiband echo-planar imaging (EPI) sequences from the Center for Magnetic Resonance Research (CMRR), University of Minnesota [83].

Data preprocessing

All data were visually inspected for artifacts. The non-brain tissue was removed using the FSL Brain Extraction Tool [84]. We preprocessed the functional data, using FSL FEAT (version 6.0.0 [85]). The preprocessing included high-pass temporal filtering (100s), spatial smoothing (FWHM = 5 mm), distortion correction using fieldmap with FUGUE, co-registration, and normalization using FLIRT [86,87] and motion correction using MCFLIRT [86] with 6 rigid-body transformations. Functional images were first co-registered to their anatomical images and then registered to an MNI standard space (FSL's 2mm standard brain template).

Data analysis

Neuroimaging data used in these analyses is stored in an OpenNeuro repository (https://openneuro.org/datasets/ds004456). Data and code necessary to reproduce the analyses and plots reported in this paper is available in the OSF repository (https://osf.io/s85hk/?view_only=f37a4a291bfd49f690f97548db66e80c).

Behavioral data analysis

Behavioral naming data recorded in the scanner during the picture naming task were transcribed and analyzed for naming latencies and accuracy. All answers corresponding to the dominant response and its synonyms were considered correct. Trials with no responses, inaudible responses and names unrelated to a given picture were classified as errors. We fitted a linear mixed-effect model to the naming latency data and a generalized mixed-effect linear model to the accuracy data (using the lmer() and glmer() functions from the lme4 for package [88] respectively). For both models we used the same formula:

Naming latencies / Accuracy ~ Language + Session + Language x Session
(1 + Language + Session | Subject) +
(1 + Language + Session | Item)

Language corresponded to a predictor with two levels: *L1* and *L2* and *Session* corresponded to a predictor with two levels: *first* and *second session*. Prior to the analysis the categorical predictors were deviation coded (L1 = -0.5; L2 = 0.5 and first session = -0.5; second session = 0.5). The analysis was performed using the lmer() function from the lme4 package. For each analysis, we first fitted a

maximal model and then identified the best random effects' structure [89]. Post-hoc pairwise comparisons were run using the lsmeans() function from the lsmeans package.

Neuroimaging data analyses

In the 1st level statistical analysis, a double-gamma hemodynamic response function was used to model the BOLD signal corresponding to each event (trial in the main task and blocks in the localizer tasks). The estimates of motion obtained with MCFLIRT were included in the 1st level GLM as nuisance covariates. We performed two types of analyses of the picture naming data: the localizer-based analyses and the whole-brain analysis.

Localizer-based analyses. Two functional runs of each localizer task were combined in a second-level analysis for each participant using an FSL fixed-effect model. To ensure better comparability of our findings with previous studies, as group-level localizer masks we used sets of functional parcels established and validated by previous studies instead of creating the group-level parcels based on our data (twelve language parcels [32]; twenty MD parcels [23]). The group-level parcels used in this analysis are available in an OSF repository (https://osf.io/s85hk/?view_only=f37a4a291bfd49f690f97548db66e80c). These group-level partitions were intersected with subject-specific data yielding a set of subject-specific group-constraint ROIs. Finally, we selected the top 10% most responsive voxels within each parcel based on the *z*-value map for a given localizer contrast (Language: speech > degraded speech, MD: hard > easy visual working memory task) and we binarized the obtained ROIs. We then used the functionally-defined ROIs for the language and MD networks to extract parameter estimates corresponding to % signal change in the two critical conditions of the main task: naming pictures in L1 and L2. Parameter estimates were extracted from the 1st level analyses for each participant separately. This was done using an FSL's FEATquery tool (http://www.FMRIb.ox.ac.uk/fsl/feat5/featquery.html). Further analyses were performed on the extracted parameter estimates in R (version: 4.0.2). Following an approach that proposes to model activity within a functional network on the network level, instead of relying on separate analysis for each ROI we fitted one linear mixed-effect model for each functional network according to the following formula:

$$\% \text{ signal change} \sim \text{Language} + \text{Hemishpere} + \text{Language x Hemisphere} +$$
$$(1 + \text{Language} + \text{Hemisphere} \mid \text{Subject}) +$$
$$(1 + \text{Language} \mid \text{ROI})$$

In each model, the fixed effect of Language corresponded to a predictor with two levels: *L1* and *L2* and the effect of *Hemisphere* corresponded to a predictor with two levels: *left hemisphere* and *right hemisphere*. Prior to the analysis this categorical predictor was deviation coded (L1 = -0.5; L2 = 0.5 and left hemisphere = -0.5; right hemisphere = 0.5). The analysis was performed using the lmer()

function from the lme4 package [88]. For each functional network, we first fitted a maximal model and then identified the best random effects' structure [89].

Bilingual language control network analysis. To evaluate language and domain-general selective responses within the Bilingual Language Control network, we used a similar approach as in the localizer-based analyses. First, we selected a set of ROIs corresponding to the Bilingual Language Control network, as described by Abutalebi and Green [15]. These included the left inferior frontal gyrus (IFG) – pars opercularis and pars triangularis, left middle frontal gyrus (MFG), left anterior cingulate cortex, left pre-supplementary motor area (pre-SMA), left angular gyrus, left thalamus, putamen and caudate as well as right IFG – pars opercularis and triangularis and right angular gyrus. The ROIs were derived from the FSL Harvard-Oxford Cortical and Subcortical atlases. As the atlases provide probabilistic masks, the selected ROIs were thresholded at 30% and binarized prior to further analyses. The pre-SMA is not available as a separate mask within the Harvard-Oxford Cortical atlas, hence it was created by splitting the Supplementary Motor Cortex ROI into anterior and posterior parts along y=0. Final set of masks used for this analysis is available at OSF (https://osf.io/s85hk/?view_only=f37a4a291bfd49f690f97548db66e80c). To create subject-specific language and domain-general selective masks within the BLC ROIs, each of the anatomically-defined ROI was intersected with activation map for each subject for corresponding to the (i) language localizer contrast, i.e., intact > degraded speech; and (ii) the MD localizer contrast, i.e., hard vWM > easy vWM. Subsequently, we selected top 10% of the most responsive voxels for each contrast and we binarized the obtained maps to create subject-specific functional ROIs (fROIs) within eacho of the anatomically-defined BLC ROI. Following that, response to L1 and L2 speech production in a picture naming task were extracted from the language- and domain-general selective fROIs using the same approach as described in the localizer-based analyses.

To assess the voxel overlap between the two networks, individual subject parcels for language and domain-general fROIs within each BLC network ROI were overlaid in a common space and a number of voxels overlapping between the two masks was calculated. Following that, the proportion of the overlapping voxels in each BLC network ROI was calculated using the following formula:

Proportion of overlapping voxels = shared / (lang fROI + MD fROI – shared)

Where shared corresponds to the number of voxels shared between the two fROIs and lang fROI and MD fROI corresponds to the total number of voxels in fROI defined using the language and MD localizers.

References

1.	Broos, W. P. J., Duyck, W. & Hartsuiker, R. J. Are higher-level processes delayed in second language word production? Evidence from picture naming and phoneme monitoring. *Lang. Cogn. Neurosci.* **33**, 1219–1234 (2018).

2.	Casado, A., Szewczyk, J., Wolna, A. & Wodniecka, Z. The relative balance between languages predicts the degree of engagement of global language control. *Cognition* **226**, 105169 (2022).

3.	Gollan, T. H., Montoya, R. I., Cera, C. & Sandoval, T. C. More use almost always means a smaller frequency effect: Aging, bilingualism, and the weaker links hypothesis☐. *J. Mem. Lang.* **58**, 787–814 (2008).

4.	Hanulová, J., Davidson, D. J. & Indefrey, P. Where does the delay in L2 picture naming come from? Psycholinguistic and neurocognitive evidence on second language word production. *Lang. Cogn. Process.* **26**, 902–934 (2011).

5.	Runnqvist, E., Strijkers, K., Sadat, J. & Costa, A. On the Temporal and Functional Origin of L2 Disadvantages in Speech Production: A Critical Review. *Front. Psychol.* **2**, (2011).

6.	Acheson, D. J., Ganushchak, L. Y., Christoffels, I. K. & Hagoort, P. Conflict monitoring in speech production: Physiological evidence from bilingual picture naming. *Brain Lang.* **123**, 131–136 (2012).

7.	Strijkers, K., Baus, C., Runnqvist, E., FitzPatrick, I. & Costa, A. The temporal dynamics of first versus second language production. *Brain Lang.* **127**, 6–11 (2013).

8.	Fu, Y. *et al.* Neural correlates for naming disadvantage of the dominant language in bilingual word production. *Brain Lang.* **175**, 123–129 (2017).

9.	Hernandez, A. E. Language switching in the bilingual brain: What's next? *Brain Lang.* **109**, 133–140 (2009).

10.	Liu, H., Hu, Z., Guo, T. & Peng, D. Speaking words in two languages with one brain: Neural overlap and dissociation. *Brain Res.* **1316**, 75–82 (2010).

11.	Rossi, E., Newman, S., Kroll, J. F. & Diaz, M. T. Neural signatures of inhibitory control in bilingual spoken production. *Cortex* **108**, 50–66 (2018).

12.	Videsott, G. *et al.* Speaking in multiple languages: Neural correlates of language proficiency in multilingual word production. *Brain Lang.* **113**, 103–112 (2010).

13.	Sulpizio, S., Del Maschio, N., Fedeli, D. & Abutalebi, J. Bilingual language processing: A meta-analysis of functional neuroimaging studies. *Neurosci. Biobehav. Rev.* **108**, 834–853 (2020).

14.	Green, D. W. Mental control of the bilingual lexico-semantic system. *Biling. Lang. Cogn.* **1**, 67 (1998).

15.	Abutalebi, J. & Green, D. W. Neuroimaging of language control in bilinguals: neural adaptation and reserve. *Biling. Lang. Cogn.* **19**, 689–698 (2016).

16.	Blanco-Elorrieta, E. & Caramazza, A. A common selection mechanism at each linguistic level in bilingual and monolingual language production. *Cognition* 104625 (2021) doi:10.1016/j.cognition.2021.104625.

17.	de Bruin, A., Roelofs, A., Dijkstra, T. & FitzPatrick, I. Domain-general inhibition areas of the brain are involved in language switching: FMRI evidence from trilingual speakers. *NeuroImage* **90**,

348–359 (2014).

18.   Green, D. W. & Abutalebi, J. Language control in bilinguals: The adaptive control hypothesis. *J. Cogn. Psychol.* **25**, 515–530 (2013).

19.   Tao, L., Wang, G., Zhu, M. & Cai, Q. Bilingualism and domain-general cognitive functions from a neural perspective: A systematic review. *Neurosci. Biobehav. Rev.* **125**, 264–295 (2021).

20.   Wu, J. *et al.* Brain network reconfiguration for language and domain-general cognitive control in bilinguals. *NeuroImage* **199**, 454–465 (2019).

21.   Duncan, J. The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. *Trends Cogn. Sci.* **14**, 172–179 (2010).

22.   Cole, M. W. *et al.* Multi-task connectivity reveals flexible hubs for adaptive task control. *Nat. Neurosci.* **16**, 1348–1355 (2013).

23.   Fedorenko, E., Duncan, J. & Kanwisher, N. Broad domain generality in focal regions of frontal and parietal cortex. *Proc. Natl. Acad. Sci.* **110**, 16616–16621 (2013).

24.   Woolgar, A., Duncan, J., Manes, F. & Fedorenko, E. Fluid intelligence is supported by the multiple-demand system not the language system. *Nat. Hum. Behav.* **2**, 200–204 (2018).

25.   Assem, M., Blank, I. A., Mineroff, Z., Ademoğlu, A. & Fedorenko, E. Activity in the fronto-parietal multiple-demand network is robustly associated with individual differences in working memory and fluid intelligence. *Cortex* **131**, 1–16 (2020).

26.   Shashidhara, S., Mitchell, D. J., Erez, Y. & Duncan, J. Progressive Recruitment of the Frontoparietal Multiple-demand System with Increased Task Complexity, Time Pressure, and Reward. *J. Cogn. Neurosci.* **31**, 1617–1630 (2019).

27.   Woolgar, A., Jackson, J. & Duncan, J. Coding of Visual, Auditory, Rule, and Response Information in the Brain: 10 Years of Multivoxel Pattern Analysis. *J. Cogn. Neurosci.* **28**, 1433–1454 (2016).

28.   Assem, M., Glasser, M. F., Van Essen, D. C. & Duncan, J. A Domain-General Cognitive Core Defined in Multimodally Parcellated Human Cortex. *Cereb. Cortex* **30**, 4361–4380 (2020).

29.   Assem, M., Shashidhara, S., Glasser, M. F. & Duncan, J. Precise Topology of Adjacent Domain-General and Sensory-Biased Regions in the Human Brain. *Cereb. Cortex* **32**, 2521–2537 (2022).

30.   Duncan, J., Assem, M. & Shashidhara, S. Integrated Intelligence from Distributed Brain Activity. *Trends Cogn. Sci.* **24**, 838–852 (2020).

31.   Fedorenko, E., Behr, M. K. & Kanwisher, N. Functional specificity for high-level linguistic processing in the human brain. *Proc. Natl. Acad. Sci.* **108**, 16428–16433 (2011).

32.   Fedorenko, E., Hsieh, P.-J., Nieto-Castañón, A., Whitfield-Gabrieli, S. & Kanwisher, N. New Method for fMRI Investigations of Language: Defining ROIs Functionally in Individual Subjects. *J. Neurophysiol.* **104**, 1177–1194 (2010).

33.   Scott, T. L., Gallée, J. & Fedorenko, E. A new fun and robust version of an fMRI localizer for the frontotemporal language system. *Cogn. Neurosci.* **8**, 167–176 (2017).

34.   Hu, J. *et al.* Precision fMRI reveals that the language-selective network supports both phrase-structure building and lexical access during language production. *Cereb. Cortex* bhac350 (2022) doi:10.1093/cercor/bhac350.

35.   Regev, T. I. *et al.* High-level language brain regions are sensitive to sub-lexical regularities. 2021.06.11.447786 Preprint at https://doi.org/10.1101/2021.06.11.447786 (2021).

36.   Blank, I., Balewski, Z., Mahowald, K. & Fedorenko, E. Syntactic processing is distributed across the language system. *NeuroImage* **127**, 307–323 (2016).

37.   Fedorenko, E., Blank, I. A., Siegelman, M. & Mineroff, Z. Lack of selectivity for syntax relateive to word meanings throughout the language network. *Cognition* **203**, 104348 (2020).

38.   Regev, T. I. *et al.* Intracranial recordings reveal three distinct neural response patterns in the language network. 2022.12.30.522216 Preprint at https://doi.org/10.1101/2022.12.30.522216 (2022).

39.   Shain, C., Blank, I. A., Fedorenko, E., Gibson, E. & Schuler, W. Robust Effects of Working Memory Demand during Naturalistic Language Comprehension in Language-Selective Cortex. *J. Neurosci.* **42**, 7412–7430 (2022).

40.   Wehbe, L. *et al.* Incremental Language Comprehension Difficulty Predicts Activity in the Language Network but Not the Multiple Demand Network. *Cereb. Cortex* **31**, 4006–4023 (2021).

41.   Malik-Moraleda, S. *et al.* Functional characterization of the language network of polyglots and hyperpolyglots with precision fMRI. 2023.01.19.524657 Preprint at https://doi.org/10.1101/2023.01.19.524657 (2023).

42.   Baus, C., Santesteban, M., Runnqvist, E., Strijkers, K. & Costa, A. Characterizing lexicalization and self-monitoring processes in bilingual speech production. *J. Neurolinguistics* **56**, 100934 (2020).

43.   De Baene, W., Duyck, W., Brass, M. & Carreiras, M. Brain Circuit for Cognitive Control Is Shared by Task and Language Switching. *J. Cogn. Neurosci.* **27**, 1752–1765 (2015).

44.   Guo, T., Liu, H., Misra, M. & Kroll, J. F. Local and global inhibition in bilingual word production: fMRI evidence from Chinese–English bilinguals. *NeuroImage* **56**, 2300–2309 (2011).

45.   Wang, Y., Xue, G., Chen, C., Xue, F. & Dong, Q. Neural bases of asymmetric language switching in second-language learners: An ER-fMRI study. *NeuroImage* **35**, 862–870 (2007).

46.   Liu, C., Jiao, L., Li, Z., Timmer, K. & Wang, R. Language control network adapts to second language learning: A longitudinal rs-fMRI study. *Neuropsychologia* **150**, 107688 (2021).

47.   Weissberger, G. H., Gollan, T. H., Bondi, M. W., Clark, L. R. & Wierenga, C. E. Language and task switching in the bilingual brain: Bilinguals are staying, not switching, experts. *Neuropsychologia* **66**, 193–203 (2015).

48.   Luk, G., Green, D. W., Abutalebi, J. & Grady, C. Cognitive control for language switching in bilinguals: A quantitative meta-analysis of functional neuroimaging studies. *Lang. Cogn. Process.* **27**, 1479–1488 (2012).

49.   Declerck, M., Kleinman, D. & Gollan, T. H. Which bilinguals reverse language dominance and why? *Cognition* **204**, 104384 (2020).

50.   De Bleser, R. *et al.* The organisation of the bilingual lexicon: a PET study. *J. Neurolinguistics* **16**, 439–456 (2003).

51.   Branzi, F. M., Della Rosa, P. A., Canini, M., Costa, A. & Abutalebi, J. Language Control in Bilinguals: Monitoring and Response Selection. *Cereb. Cortex* **26**, 2367–2380 (2016).

52.   Wang, R. *et al.* Functional and structural neuroplasticity associated with second language proficiency: An MRI study of Chinese-English bilinguals. *J. Neurolinguistics* **56**, 100940 (2020).

53.   Fedorenko, E. The early origins and the growing popularity of the individual-subject analytic approach in human neuroscience. *Curr. Opin. Behav. Sci.* **40**, 105–112 (2021).

54.   Michon, K. J., Khammash, D., Simmonite, M., Hamlin, A. M. & Polk, T. A. Person-specific and precision neuroimaging: Current methods and future directions. *NeuroImage* **263**, 119589 (2022).

55.   Hull, R. & Vaid, J. Laterality and language experience. *Laterality* **11**, 436–464 (2006).

56.   Hull, R. & Vaid, J. Bilingual language lateralization: A meta-analytic tale of two hemispheres. *Neuropsychologia* **45**, 1987–2008 (2007).

57.   Park, H. R. P., Badzakova-Trajkov, G. & Waldie, K. E. Language lateralisation in late proficient bilinguals: A lexical decision fMRI study. *Neuropsychologia* **50**, 688–695 (2012).

58.   Waldie, K. E. *et al.* The cognitive and neural correlates of written language: a selective review of bilingualism. *J. R. Soc. N. Z.* **51**, 81–96 (2021).

59.   Poldrack, R. A. Can cognitive processes be inferred from neuroimaging data? *Trends Cogn. Sci.* **10**, 59–63 (2006).

60.   Kanwisher, N., McDermott, J. & Chun, M. M. The Fusiform Face Area: A Module in Human Extrastriate Cortex Specialized for Face Perception. *J. Neurosci.* **17**, 4302–4311 (1997).

61.   Van de Putte, E., De Baene, W., Brass, M. & Duyck, W. Neural overlap of L1 and L2 semantic representations in speech: A decoding approach. *NeuroImage* **162**, 106–116 (2017).

62.   Van de Putte, E., De Baene, W., Price, C. J. & Duyck, W. "Neural overlap of L1 and L2 semantic representations across visual and auditory modalities: a decoding approach". *Neuropsychologia* **113**, 68–77 (2018).

63.   Xu, M., Baldauf, D., Chang, C. Q., Desimone, R. & Tan, L. H. Distinct distributed patterns of neural activity are associated with two languages in the bilingual brain. *Sci. Adv.* **3**, e1603309 (2017).

64.   Chee, M. W. L., Soon, C. S. & Lee, H. L. Common and Segregated Neuronal Networks for Different Languages Revealed Using Functional Magnetic Resonance Adaptation. *J. Cogn. Neurosci.* **15**, 85–97 (2003).

65.   Klein, D., Milner, B., Zatorre, R. J., Zhao, V. & Nikelski, J. Cerebral organization in bilinguals: A PET study of Chinese-English verb generation. *NeuroReport* **10**, 2841–2845 (1999).

66.   Indefrey, P. The Spatial and Temporal Signatures of Word Production Components: A Critical Update. *Front. Psychol.* **2**, (2011).

67.   Green, D. W. & Kroll, J. F. The Neurolinguistics of Bilingualism: Plasticity and Control. in *The Oxford Handbook of Neurolinguistics* (Oxford University Press, 2019).

68.   Papoutsi, M. *et al.* From Phonemes to Articulatory Codes: An fMRI Study of the Role of Broca's Area in Speech Production. *Cereb. Cortex* **19**, 2156–2165 (2009).

69.   Indefrey, P. A Meta-analysis of Hemodynamic Studies on First and Second Language Processing: Which Suggested Differences Can We Trust and What Do They Mean? *Lang. Learn.* **56**, 279–304 (2006).

70.   Indefrey, P. & Levelt, W. J. M. The spatial and temporal signatures of word production components. *Cognition* **92**, 101–144 (2004).

71.   Flinker, A. *et al.* Redefining the role of Broca's area in speech. *Proc. Natl. Acad. Sci.* **112**, 2871–2875 (2015).

72.   Thompson-Schill, S. L., D'Esposito, M., Aguirre, G. K. & Farah, M. J. Role of left inferior prefrontal cortex in retrieval of semantic knowledge: A reevaluation. *Proc. Natl. Acad. Sci.* **94**, 14792–14797 (1997).

73.   Schnur, T. T. *et al.* Localizing interference during naming: Convergent neuroimaging and neuropsychological evidence for the function of Broca's area. *Proc. Natl. Acad. Sci.* **106**, 322–327 (2009).

74.   Zhang, J. X., Feng, C.-M., Fox, P. T., Gao, J.-H. & Tan, L. H. Is left inferior frontal gyrus a general mechanism for selection? *NeuroImage* **23**, 596–603 (2004).

75.   Fedorenko, E. & Thompson-Schill, S. L. Reworking the language network. *Trends Cogn. Sci.* **18**, 120–126 (2014).

76.   Fedorenko, E., Duncan, J. & Kanwisher, N. Language-Selective and Domain-General Regions Lie Side by Side within Broca's Area. *Curr. Biol.* **22**, 2059–2062 (2012).

77.   Lindell, A. K. In Your Right Mind: Right Hemisphere Contributions to Language Processing and Production. *Neuropsychol. Rev.* **16**, 131–148 (2006).

78.   Birn, R. M. *et al.* Neural systems supporting lexical search guided by letter and semantic category cues: A self-paced overt response fMRI study of verbal fluency. *NeuroImage* **49**, 1099–1107 (2010).

79.   Wolna, A. *et al.* Tracking components of bilingual language control in speech production: an fMRI study using functional localizers. 2023.02.07.527469 Preprint at https://doi.org/10.1101/2023.02.07.527469 (2023).

80.   Lemhöfer, K. & Broersma, M. Introducing LexTALE: A quick and valid Lexical Test for Advanced Learners of English. *Behav. Res. Methods* **44**, 325–343 (2012).

81.   Wolna, A., Łuniewska, M., Haman, E. & Wodniecka, Z. Polish norms for a set of colored drawings of 168 objects and 146 actions with predictors of naming performance. *Behav. Res. Methods* (2022) doi:10.3758/s13428-022-01923-3.

82.   Dale, A. M. Optimal experimental design for event-related fMRI. *Hum. Brain Mapp.* **8**, 109–114 (1999).

83.   Xu, J. *et al.* Evaluation of slice accelerations using multiband echo planar imaging at 3T. *NeuroImage* **83**, 991–1001 (2013).

84.   Smith, S. M. Fast robust automated brain extraction. *Hum. Brain Mapp.* **17**, 143–155 (2002).

85.   Woolrich, M. W., Behrens, T. E. J., Beckmann, C. F., Jenkinson, M. & Smith, S. M. Multilevel linear modelling for FMRI group analysis using Bayesian inference. *NeuroImage* **21**, 1732–1747 (2004).

86.   Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage* **17**, 825–841 (2002).

87.   Jenkinson, M. & Smith, S. A global optimisation method for robust affine registration of brain images. *Med. Image Anal.* **5**, 143–156 (2001).

88.   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

89.   Bates, D., Kliegl, R., Vasishth, S. & Baayen, H. Parsimonious Mixed Models.

ArXiv150604967 Stat (2018).

Acknowledgements

We would like to thank all members of the Psychology of Language and Bilingualism Lab who contributed to this project by collecting and coding the data and to all participants who took part in the study. We are very grateful to Evelina Fedorenko for inspiration and insightful comments on the design and results. This research was possible thanks to an OPUS grant from the National Science Center Poland, awarded to ZW (2017/27/B/HS6/00959).

Author contributions

AW, JS, MD, MS, and ZW contributed to the conception, design, and methodology of the study. AW and AD collected the data and AW designed and conducted data analysis. AW, JS, MD and ZW interpreted the analyses. ZW was responsible for project supervision. AW, JS and ZW wrote the first draft of the manuscript and AD and MD provided critical review and revision the text. All authors approved the final version for publication.

Data availability statement

The raw neuroimaging that served as a basis for analyses in this paper along with the data on participants' language experience is available at https://openneuro.org/datasets/ds004456.

Data and code necessary to reproduce the presented ROI and behavioural analyses is available at the https://osf.io/s85hk/?view_only=f37a4a291bfd49f690f97548db66e80c.

Additional Information

Authors declare no competing interests.