Juvenile niches select between two distinct development
trajectories and symbiosis modes in vent shrimps

Author & Affiliations

Pierre Methou[1], Marion Guéganton[2], Jonathan T. Copley[3], Hiromi Kayama Watanabe[1], Florence
Pradillon[2], Marie-Anne Cambon-Bonavita[2], Chong Chen[1]

[1]X-STAR, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokosuka,
Kanagawa 237-0061, Japan

[2]Univ Brest, Ifremer, CNRS, Unité Biologie des Environnements Extrêmes marins Profonds, F-
29280, Plouzané, France

[3]Ocean and Earth Science, National Oceanography Centre Southampton, University of
Southampton, Southampton SO14 3ZH, United Kingdom

Abstract

Most animal species have a singular developmental pathway and adult ecology, but
developmental plasticity is well-known in some like honeybees where castes display profoundly
different morphology and ecology. An intriguing case is the Atlantic deep-sea hydrothermal
vent shrimp *Rimicaris hybisae/chacei* that share dominant COI haplotypes but develops into
either the symbiont-reliant *hybisae* with a hypertrophied head chamber (in the Mid-Cayman
Spreading Centre) or the mixotrophic *chacei* with a narrow head chamber (on the Mid-Atlantic
Ridge). Here, we use X-ray micro-computed tomography and fluorescent *in situ* hybridization to
show that key anatomical shifts in both occur between the juvenile-subadult transition, when
those developing into *hybisae* have fully established symbiosis but not *chacei*. On the Mid-
Atlantic Ridge the diet of *R. chacei* has been hypothetically linked to competition with the
obligatorily symbiotic congener *R. exoculata*, and we find anatomical evidences that *R.*
exoculata is indeed better adapted for symbiosis – suggesting the *chacei* morph could be an
adaptation to prevent competitive exclusion. Our results suggest that the two distinct
development trajectories are likely selected according to the diet available to the juveniles,
determined by whether energetically sufficient symbiont colonisation has occurred before
reaching subadult stage.

Introduction

Each animal species is generally constrained to a single mode of ontogenetic
development, leading to individuals (of the same sex) exhibiting similar ecology at each life
stage. Nevertheless, there are some well-documented cases where individuals of the same
species can develop into profoundly different forms with distinct 'ways of life.' Castes of

eusocial insects is a well-known example, where nutritional input typically has a key role in determining the development outcome. Whether a female larva of the honeybee *Apis mellifera* develops into worker or queen, for example, is almost entirely dependent on if it is fed a large quantity of royal jelly [1]. Another example is the cave fish *Astyanax mexicanus*, whose morphology differs greatly among individuals living in rivers and surrounding caves – cave morphotypes have lost their eyes and show reduced pigmentation but have larger body sizes and other sensory organs [2]. At hydrothermal vents, a significant case is the *Ridgeia piscesae* tubeworm, which develops into a short-fat or a long-skinny morphotype depending on the vent flow regime [3,4].

The formation of symbioses, intimate and long-term biological association between two or more organisms [5], is also a powerful evolutionary force that can have fundamental impacts on the developmental processes and ecology of the host animal [6–8]. For example, the bobtail squid *Euprymna scolopes* houses the bioluminescent symbiont *Vibrio fisheri* for counter-illumination in its light organ which morphogenesis requires signalling molecules from the symbiont [9]. Deep-sea hydrothermal vent ecosystems are unusual in that intimate symbiotic relationships between chemoautotrophic bacteria and endemic invertebrate animals form the basis for the entire system [10]. Vent holobionts usually acquire their symbionts from the environment after they settle in the vent field as juveniles – with the exception of vesicomyid clams which have maternal inheritance of chemosynthetic symbionts [11]. As significant transitions in ecological niche within the life cycle of a species often involve metamorphosis, where conspicuous anatomical and physiological shifts occur, symbiont acquisition is often linked to such metamorphosis involving the loss or the reduction of the digestive organs and an enlargement of the symbiont hosting structures [12–14]. These reconfigurations of internal organs post-settlement are not necessarily apparent from the external morphology, as seen with the 'cryptometamorphosis' of *Gigantopelta* vent snails where the symbiont-hosting 'trophosome' suddenly expands internally [15] or the straightening of the digestive tube in bathymodioline mussels [16].

Among vent animals, alvinocaridid shrimps exhibit a wide range of feeding strategies and reliance on symbiosis. A strict dependence on symbiosis is found in two closely related species, *Rimicaris exoculata* and *R. kairei*, hosting dense communities of filamentous bacterial symbionts inside their expanded 'head' (the cephalothorax) that provides most of their diet [17–20]. These symbiotic species exhibit striking morphological differences from other alvinocaridid shrimps, with the enlargement of their symbiont-hosting cephalothoracic cavities and mouthparts starting during the post-settlement metamorphosis between juveniles and adults [19,21]. Conversely, several other species of vent shrimps such as *Nautilocaris saintlaurentae* and *Rimicaris variabilis* harbour few to no symbiotic bacteria and instead feed on other food sources such as bacterial mats or detritus [22,23].

One species complex of alvinocaridid shrimps from Atlantic vents – the *Rimicaris hybisae/chacei* complex – is exceptional as *R. hybisae* and *R. chacei* share dominant *COI* haplotypes (electronic supplementary material, figure S1) but metamorphose into two distinct forms occupying disparate ecological niches depending on the environment [24,25]. The

apparent differences in morphology led to their descriptions as separate species [26], but
currently available genetic barcode data do not support the existence of two distinct species-
level clades [24,27]. *Rimicaris hybisae* occurs in the Mid-Cayman Spreading Centre (MCSC),
mostly rely on symbiosis and has an external morphology resembling *R. exoculata* with an
inflated cephalothorax [26,28]. It appears to still retain an ability to feed on alternative sources
outside of its symbiosis, with evidences of facultative carnivory/scavenging in individuals
distributed at the periphery of vent fluid emissions [25]. In contrast, *R. chacei* from the Mid-
Atlantic Ridge (MAR) co-occur with *R. exoculata* [29] and lacks an enlarged cephalothorax; it
has a mixed diet where it also feeds on bacterial mat and scavenges in addition to some
contribution from symbiosis [21,30].

Although *R. exoculata* and *R. chacei* co-occur at MAR vents, *R. exoculata* dominates the
vigorously venting 'black-smoker' chimneys whereas *R. chacei* mainly occurs in diffuse flow
environment with much weaker fluid input [29]. This is in stark contrast to the situation at
MCSC where *R. hybisae* is the only shrimp species dominating the surface of active smoker
chimneys [26]. This leads to a hypothesis that the *R. chacei* adult morphology is a result of
competitive exclusion during the juvenile stage from the vigorously venting environment by *R.*
exoculata [29], and that this environment is necessary for development into the *R. hybisae*
morphology. Anatomy and food source are closely linked in animals, and organ volumetric is a
useful estimator of feeding ecology for hydrothermal vent species where detailed *in situ*
studies are difficult [31]. Until now, no direct comparisons in the anatomy, ontogenetic changes,
and symbiont acquisition between *R. hybisae* vs *R. chacei* have been made, or how they
compare to the obligatorily symbiotic *R. exoculata*. In order to shed light on this hypothesis, we
combine quantitative 3D anatomy by X-ray micro-computed tomography (μ-CT) and
fluorescence in situ hybridisation (FISH) to answer three questions: 1) At which life stage does
the split in development trajectories occur in *R. hybisae/chacei*, 2) Whether this is correlated
with symbiont acquisition timing, and 3) Is there anatomical evidence that *R. hybisae/chacei*
would indeed be outcompeted by *R. exoculata*.

108   Methods

Specimens of the three main target shrimp species – *R. exoculata*, *R. hybisae*, and *R.*
chacei – were collected at hydrothermal vent fields during several sea-going expeditions: *R.*
exoculata and *R. chacei* were collected at the TAG and Snake Pit vent fields during the BICOSE 2
(R/V *Pourquoi pas?,* 2018, doi: https://doi.org/10.17600/18000004) expedition using suction
sampler of the human-occupied vehicle (HOV) *Nautile*; *R. hybisae* were collected at the Von
Damm and Beebe vent fields during the expeditions JC82 (RRS *James Cook*, 2013) and YK13-05
(R/V *Yokosuka*, 2013) expeditions, respectively, with suction samplers on ROV *Isis* or HOV
Shinkai 6500. In addition, we also studied the anatomy of *R. kairei* to confirm that it does not
differ significantly from that of its closest relative, *R. exoculata*; using specimens collected from
the Kairei and Edmond vent fields during expeditions YK09-13 (R/V *Yokosuka*, 2009) and YK16-
E02 (R/V *Yokosuka*, 2016) using suction samplers mounted on the HOV *Shinkai 6500*. Shrimps

were preserved in 10% formalin or 99% ethanol and measured with vernier calliper using
carapace length (CL) for standardized measurement, upon recovery on-board. Individuals were
identified to their life stages, according to Methou *et al.* [21]. Subadults of all species were
characterized as small individuals morphologically similar to adults with sizes below the onset
of sexual maturity whereas stage A and stage B juveniles of *R. exoculata* were distinguished by
the morphology of their eyes [21]. Given that *R. chacei* juveniles settle and begin their
anatomical transitions at smaller sizes than *R. exoculata*, their adult stages were separated into
two categories: 'small adults' of similar sizes to that of *R. exoculata* juvenile stages (CL = 8-10
mm) and 'adults' (CL= 14-16 mm). Life stages of *R. kairei* was defined following the established
nomenclature for *R. exoculata* and the same was done for *R. hybisae* based on *R. chacei*.
Additional subadult specimens of *R. exoculata*, *R. chacei*, and *R. hybisae* were fixed in 3%
paraformaldehyde (PFA) for 3 hours on board, rinsed with phosphate buffered saline (PBS) /
sterile seawater buffer and stored in 50:50 2x PBS/ethanol at -20°C for fluorescence *in situ*
hybridization (FISH) analyses (electronic supplementary material, table S1).

A total of 47 individuals, including a triplicate of each life stages for the four-shrimp
species (except where only two individuals were available for stage A juveniles of *R. hybisae*),
were used for X-ray micro-computed tomography (μ-CT) scanning. Shrimps preserved in 99%
ethanol were progressively rehydrated in Milli-Q water in four steps (25:75, 50:50 and 75:50
Mili-Q: Ethanol), and then stained for 24 hours in 0.05 mol/L iodine solution. Scanning was
performed using a ScanXmate-D160TSS105 (Comscantecno, Japan) commercial μ-CT at 60-90
kV/25-92 μA, with a resolution of 10.904-49.587 μm per pixel, at 992 x 992 pixels resolution
per slice (for details see electronic supplementary material, table S2). Images obtained were
imported into the specialist software Amira version 2022 (Thermo Fisher) for visualisation and
manual segmentation of the different organs of interest. This included the symbiont-hosting
branchiostegite (i.e., wall of the cephalothoracic gill chamber) and mouthparts
(scaphognathites and exopodites), as well as the digestive system (stomach, digestive tube, and
hepatopancreas). Only the anterior part of the cephalothoracic cavities were reconstructed,
since posterior components facing the gills are known to be not colonized by bacterial
symbionts [18,20,30]. Prior to smoothing, organ volumes and surface areas were calculated
also in Amira based on the segmentation. As pereopods and antennae were often fully or
partially broken, these parts were removed from all our 3D reconstructions and thus the
measurement of total body volumes and surface areas (see Table S3 for detailed body volumes
and surface areas measurements and supplementary material for 3D interactive models). The
final surface renderings used for the figures were generated post-smoothing and complexity-
reduction. Cephalothoracic cavities of five specimens at each life stage of *R. chacei* and *R.*
hybisae (except where only two individuals available for stage A juveniles of *R. hybisae*) were
also dissected to observe the distribution of bacteriophore setae on their mouthparts.
Mouthparts of one *R. exoculata* adult were also dissected and observed. As the anatomical
features and patterns seen in *R. kairei* generally matched *R. exoculata*, for simplicity and clarity
we only displayed the three main study species in the main figures and the *R. kairei* data are
figured in electronic supplementary material, figures S2-S4.

161   For FISH analyses of subadult symbiont colonization, the cephalothorax of *R. exoculata*
162 and *R. chacei* subadults were progressively dehydrated in a PBS-ethanol series (ambient
163 temperature) before embedding in polyethylene glycol distearate-1-hexadecanol (9: 1) resin
164 (Sigma, St. Louis, MO). Resin blocks were stored in -20°C until trimming into 10 μm sections
165 with a RM 2255 microtome (Leica Biosystems, Nussloch, Germany). Sections were placed on
166 adhesive glass slides (Menzel-Gläser Superfrost® Plus, USA), residues of resin were removed
167 prior to hybridization with three baths of 96% ethanol for 5 min each and tissues were
168 partially rehydrated with a bath of 70 ethanol. The cephalothorax of *R. hybisae* subadults
169 were cut from the abdomens and rehydrated in 50:50 PBS-ethanol bath followed by 75:25 PBS-
170 ethanol (30 min each) and then rinsed three times in 1x PBS buffer for 5 min each. Specimens
171 were then progressively transferred to an optimum temperature compound (OCT) (Tissue-Tek;
172 Sakura Finetek, Osaka, Japan): first in 15% sucrose-PBS, followed by 30% sucrose-PBS, and
173 finally in 30% sucrose-PBS/OCT, for 1h 30 min each, before embedding in OCT in a plastic
174 holder at 4°C. Sectioning of OCT-embedded specimens was done with a cryostat (CM1520;
175 Leica Biosystems) in a chamber at -20°C. Sections of 8 μm thickness were thaw-mounted on
176 adhesive glass slides. Before hybridization, mounted sections were washed in 1x PBS two times
177 for 5 min each, then postfixed 1x PBS-10% formalin for 10 min, and finally washed again in 1x
178 PBS for three times at 5 min each. These two different embedding and sectioning protocols
179 have already been used for FISH experiments with *Rimicaris* shrimps and are not expected to
180 impact the detection of symbionts [18,19].

181   For all species, sections were hybridized in a mix containing 0.5 mM (0.5 μM final
182 concentration) of the Eub338 (eubacteria) probe in a 30% (*R. chacei* & *R. exoculata*) or 35% (*R.*
183 *hybisae*) formamide hybridization buffer for 3 h at 46°C following a published protocol [19].
184 Sections were washed at 48°C for 30 min in a washing buffer [19], rinsed briefly with miliQ
185 water, air dried, and covered with SlowFade Gold antifade reagent mounting medium
186 containing DAPI (Invitrogen) and a cover slip. Observations of *R. exoculata* and *R. chacei*
187 subadults were made using a Zeiss Imager.Z2 microscope equipped with the Apotome.2®
188 sliding module and Colibri.7 light technology (Zeiss, Oberkochen, Germany) and processed with
189 the Zen software (Zeiss). Observations of *R. hybisae* subadults were made using a Leica Stellaris
190 STED confocal scanning microscope (Leica, Germany) and analysed with the LAS X software.

Results

193   Anatomy of *Rimicaris* species was reconstructed across four stages of development
194 from recently settled juvenile to adult stages (figure 1; electronic supplementary material,
195 figures S2 & S3). Early juvenile stages of all species had very small stomach volumes, ranging
196 from 0.12 ± 0.01% of the total body volume in *R. exoculata* to 0.32 ± 0.04% in *R. chacei* (Figures
197 1 and 2). In adults, *R. chacei* and *R. hybisae* both possessed large stomach volumes
198 representing 1.73 ± 0.84% and 2.18 ± 1.3% of their body volumes, respectively (Figures 2 and
199 3); whereas adult stages of *R. exoculata* had much smaller stomachs at 0.69 ± 0.32% of the
200 body volume (Figures 2 and 3). However, their stomachs expanded differently during their

transition from juvenile to subadult with a dramatic increase, although variable among
individuals, up to 2.03 ± 0.99% right from the subadult stages of *R. chacei* while it was more
gradual in *R. hybisae* with an increase up to 0.86 ± 0.09% in subadults and 0.77 ± 0.13% in
small adults (Figure 1 and 2). On the other hand, the hepatopancreas followed a similar
trajectory in the three species, occupying a very large volume in juvenile stages (*R. exoculata*:
18.6 ± 1.3%; *R. hybisae*: 17.2 ± 0.96%; *R. chacei*: 17.3 ± 4.0%) representing lipid reserves, and
then decreasing rapidly when metamorphosising into subadults (*R. exoculata*: 8.55 ± 1.93%; *R.*
hybisae: 6.61 ± 1.07%; *R. chacei*: 8.02 ± 2.85%). No clear variation could be seen in the volumes
of digestive tubes along the metamorphosis of the three species (figures 2 and S4). Overall, the
anatomical development of the *R. kairei* digestive system exhibited similar patterns as seen in *R.*
exoculata, although with an earlier stomach enlargement in stage B juveniles (*R. kairei*: 0.60 ±
0.26%; *R. exoculata*: 0.15 ± 0.04%) as well as an earlier decrease of hepatopancreas volume
(Juveniles B; *R. kairei*: 8.08 ± 3.23%; *R. exoculata*: 18.6 ± 1.3%) (electronic supplementary
material, figures S3-S4).

The symbiont-hosting branchiostegite (wall of the gill chamber) surface area was found
to exhibit a sudden expansion in *R. exoculata* between juvenile B (1.96 ± 0.52% of total body
surface) and subadult stages (3.72 ± 0.56% of the total body surface), and then up to 4.14 ±
0.17% in adult stages (figures 1 and 2). Surfaces of other symbiont-hosting mouthparts –
scaphognathites and exopodites – similarly expanded between juvenile B and subadult stages
of *R. exoculata* from 1.05 ± 0.28% to 1.82 ± 0.22% of the total body surfaces (Figure 2). The
pattern was also seen in *R. kairei* with rapid expansion before subadult, main differences being
in *R. kairei* the expansion was also evident between juveniles A and B and the expansion being
less and more variable in adults (see electronic supplementary material, figures S3-S4). The
increase of branchiostegite surfaces was more rapid in *R. hybisae* between juvenile and
subadult stages and more or less isometric thereafter, from 1.33 ± 0.06% of the total body
surface in stage A juveniles to 1.92 ± 0.33% in subadults and 2.74 ± 0.23% in adults. Conversely,
surfaces of *R. chacei* branchiostegites expanded only slightly along their ontogeny from 1.38 ±
0.05% to 1.60 ± 0.05%. For mouthparts, the increment was less marked than *R. exoculata* for
both *R. hybisae* and *R. chacei*, with a limited growth of both mouthparts from 0.69 ± 0.18% to
1.37 ± 0.15% between stage A juveniles and adults in *R. hybisae* and 0.79 ± 0.09% to 1.16 ±
0.30% in *R. chacei* (Figure S4). Dissection of the adult mouthparts between *R. hybisae* and *R.*
chacei however, showed a striking difference in the coverage of bacteria-hosting setae; being
much less dense on *R. chacei* mouthparts than on *R. hybisae* and *R. exoculata* mouthparts
(figure 3). The dense setae would mean a much greater colonisable surface area in *R. hybisae*
mouthparts than *R. chacei* (figure 3). Our FISH observations of subadult specimens indicate
that the three taxa differed in the timing of their complete bacterial colonization (figure 4).
Although subadult mouthparts were all colonized by filamentous bacteria in the three species,
the branchiostegites of *R. exoculata* and *R. hybisae* subadults were densely colonized across
the entire surface, but only partly in *R. chacei*.

Discussion

Our results confirm that *Rimicaris hybisae* and *R. chacei* indeed display striking
differences in anatomical development, despite the lack of genetic differentiation based on
mitochondrial *COI* gene suggesting they are likely two morphotypes of the same species-level
lineage [24,27]. The symbiont-hosting organs of the *R. hybisae* form increased in percentage
surface area (as a fraction of the overall body surface) throughout growth with the fastest
increase between juvenile and subadult stages, whereas there was almost no increase in *R.*
chacei. The symbiont-hosting mouthparts were not different in the overall surface area
between the two forms, but they were densely covered by setae only in *R. hybisae* (like *R.*
exoculata) meaning a much higher realised surface area for colonisation in *R. hybisae*. The
chacei morphotype also exhibits a dramatic increase in stomach size before reaching subadult
stage, whereas this was not seen in *R. hybisae*. As such, the most significant shifts in the
volume or surface area of key organs (figure 2) was clearly between juvenile and subadult
stages, indicating that the development trajectory into either form is determined in the 'critical
period' between these two stages. All species examined rapidly decreased in the volume of
hepatopancreas before reaching the subadult stage, reflecting a rapid depletion of the lipid
reserve accumulated before settlement [21]. This means the shrimps must become
energetically self-sustaining before reaching the subadult stage. Taken together, these results
strongly imply that the ecological niche occupied by juvenile *R. hybisae/chacei* before reaching
subadult size is key in deciding the development trajectory and therefore the niche occupied as
adults. The total anatomical shift between juvenile and subadult stages may be sufficient to be
considered a post-settlement metamorphosis [15].

Still, there are two key anatomical characters shared across *R. hybisae* and *R. chacei*
that are distinct from the anatomical characteristics of *R. exoculata* (and *R. kairei*). The final
stomach volumes of *R. hybisae* and *R. chacei* are equivalent at the adult stage, both being
much larger than that of *R. exoculata*. The overall mouthpart surface areas are also smaller in
both *R. hybisae* and *R. chacei* compared to *R. exoculata*. Furthermore, large variations in
stomach volumes of adults could be seen among individuals of both *R. hybisae* and *R. chacei*.
This inter-individual variability could be related to plasticity in feeding ecology with the
potential for facultative carnivory/scavenging reported for some *R. hybisae* individuals [25] or
mixed diets with variable contribution from food and symbiosis among *R. chacei* individuals
[21,30]. We recognise that potentially different fixation conditions of specimens collected
during different expeditions could be responsible for the observed variability to some extent,
but this should not influence the general trends observed. Thus, although *R. hybisae* is much
more similar to *R. exoculata* superficially at a glance from the external morphology, some
features of its internal anatomy are much more similar to *R. chacei* and are indicative of their
genetic relationship (electronic supplementary material, figure S1) [24,27].

Enlargement of symbiont-hosting organs and the correlated reduction of digestive
system are prime anatomical signals in hot vent species for increased reliance on
chemosymbiotic bacteria, resulting in increased energetic yield [31,32]. In the cephalothorax,
the surface areas of all symbiont-hosting organs in *R. exoculata* exhibited sudden, exponential
growth between stage B juveniles and subadults like between juvenile A and subadults for *R.*

hybisae, but to a much greater extent. The larger symbiont-hosting organs coupled with
reduced stomach volume in *R. exoculata* compared to *R. hybisae/chacei* are therefore
anatomical evidences that *R. exoculata* is indeed the more specialised species for
chemosynthetic symbiosis and would therefore outcompete the less specialised *R.*
hybisae/chacei, as previously hypothesised [17,18,25,28]. Since *R. exoculata* co-occurs with *R.*
chacei on MAR vent fields, juveniles of the two species have been suggested to directly
compete for space and thus access to the vent fluid source, with *R. exoculata* outcompeting *R.*
chacei from the hottest and most energetically superior habitable zone on active smoker
chimneys [29]. This is supported by their spatial segregation patterns in cooler areas for *R.*
chacei, and clear differences in their population demographics with a collapse of *R. chacei*
populations following settlement phase suggestive of a high juvenile mortality [29]. This
interspecific competition with *R. exoculata* may limit acquisition of symbionts for *R. chacei*
juveniles situated away from the influence of vent fluid and ultimately hamper the
development of an energetically sufficient chemosymbiosis; promoting a shift towards a
mixotrophic feeding strategy [29]. On the other hand, *R. hybisae* has no known competitor at
the MCSC vents [24,33] and can fuel its chemosymbiosis from its early acquisition in juvenile
stages. This niche differentiation between juveniles of *R. hybisae/chacei* in MCSC vs MAR could
therefore drive the separation of local individuals onto two ontogenetic development pathways.

Our observations show that *R. hybisae* and *R. chacei* differ in the timing of their
symbiont colonization on branchiostegites, being densely colonized in the subadult stages in *R.*
hybisae but not *R. chacei*, despite the organ being colonised by bacteria in the adults of both
forms [26,30]. This coincides well with the timing of key anatomical shifts before the subadult
stage, and suggests having well-colonized branchiostegites during this 'critical period' may be a
condition for taking the symbiosis-reliant developmental trajectory. The delayed symbiont
colonisation in *R. hybisae/chacei* at MAR vents during this competency window, due to being
outcompeted by *R. exoculata* and potentially unable to provide sufficient chemosynthetic
energy sources for its symbionts, may lead to individuals going down the mixotrophic *R. chacei*
pathway [21,29]. As such, symbionts may drive the morphogenesis of enlarged symbiont-
hosting organ in the shrimp. Though this is still speculative in our case, similar cases are known,
such as the morphogenesis of the light organ in juveniles of the bobtail squid *Euprymna*
scolopes which are mediated by the recognition of microorganism-associated molecular
patterns (MAMPs) from its *Vibrio fischeri* symbionts upon acquisition [9]. Moreover, *E. scolopes*
juveniles treated with antibiotics enter in a refractory state after five days and cannot be
recolonized by *V. fischeri* symbionts [34]. In juvenile *Rimicaris* shrimps, the transition from
juvenile to subadult is certainly aided by red lipid reserves stored in the hepatopancreas and
the depletion of this may mark the end of the 'critical period.' This would suggest that shifts in
the cephalothorax anatomy of *R. chacei* and *R. hybisae* are reversible either to a more or a less
developed symbiosis if exposed to different conditions during their juvenile development.

In species with differential adult phenotypes and therefore developmental pathways,
an individual's fate could be determined purely by environmental factors or genetic factors, or
anything in the spectrum in-between [1]. For example, phenotype variations in blind cavefishes

appear to be purely genetic [2,35] whereas caste determination in different species of eusocial
insects stem from a combination of various environmental stimuli and genetic basis mediated
by epigenetics [1]. The two distinct morphotypes of *Ridgeia piscesae* tubeworms is a similar
case in vents, influenced by vent flow dynamics [3]. This rapid divergence among populations
can be facilitated by phenotypic plasticity or by standing genetic variation with pre-existing
polymorphism of relevant alleles in ancestral populations [36–38]. Freshwater populations of
the stickleback fishes *Gasterosteus aculeatus* which recently and repeatedly diverged from
marine ancestors in several rivers and lake around the world after the last glaciation [39]
exhibit a high level of phenotypic variability with the freshwater ones exhibiting reduction in
bony plates linked to an allele that was already present at low frequency in the ancestral
population [37,39]. On the other hand, divergences among streams and lakes ecotypes of this
same species in terms of body depth, head shape, or morphology of the digestive system, were
found to be related to an interplay of phenotypic plasticity and genetic predisposition [40].
Though we hypothesise that the developmental fate of *R. hybisae/chacei* is largely
environmental and down to accessibility to the active smoker habitat and the timely
colonisation of symbionts on the branchiostegites, fine-scale genomic data contrasting the two
forms is still missing. Therefore, we cannot exclude the possibility of genetic factors having
important roles as well. Future work, including population genomics, will be required to
disentangle the respective influence of environment vs genetics on the phenotype and ecology
of these shrimps.

Understanding how chemosymbiotic relationships are established is a major challenge
because symbiosis is such a successful and effective strategy that most hosts energetically
reliant on symbionts have become obligate holobionts that cannot survive or develop into
adulthood to reproduce without their partners [13,41]. Our results highlight *R. hybisae/chacei*
as an exceptional case where the host is able to take a separate, mixotrophic development
trajectory if symbiont colonization is limited, as a 'back up plan' to survive and reproduce. We
show that individuals have a limited 'critical period' before reaching subadult stage that
determines the trajectory, likely by whether an energetically viable symbiosis has been
established by then. Importantly, with this new information, *R. hybisae/chacei* can be a
valuable model for understanding how chemosymbioses are established and how they function,
as it can provide comparable data for genetically equivalent individuals in different symbiotic
conditions and also during transitions – like how the blind cavefish model has proved useful for
studying eye loss [35]. Our work therefore lays the foundations for studying how symbiotic
relationships emerge and how dialogues can be established between animals and
microorganisms.

Acknowledgments

We thank the captains and crews of the different research cruises conducted on-board
R/V *Pourquoi pas?* (BICOSE 2), R/V *Yokosuka* (YK09-13, YK13-05 & YK16-E02), and R/V *James*
Cook (JC82). We also thank the pilots and the operation team of the HOV *Nautile, Shinkai 6500*

and of the ROV *ISIS* during these research cruises. We gratefully acknowledge the chief
scientists of the relevant expeditions: Marie-Anne Cambon-Bonavita (Ifremer; BICOSE 2), Ken
Takai (JAMSTEC; YK13-05, YK16-E02), Kentaro Nakamura and Satoshi Nakagawa (JAMSTEC;
YK09-13) and Jon Copley (NOC; JC82). We also thank Rika Horiuchi (JAMSTEC) for her
assistance with μ-CT scanning facilities at JAMSTEC. Faunal collections of *R. exoculata*, *R. chacei*
and *R. kairei* specimens were conducted within international waters. Permission for sampling *R.*
hybisae specimens in territorial waters of the Cayman Islands was issued by the appropriate
United Kingdom authorities to R/V Yokosuka during YK13-05.

Data Accessibility

373   3D Interactive models from each stage and shrimp species are available as
374 supplementary material data. Metadata associated to each shrimp individuals including
375 collection date, sampling location, storage conditions and body size are displayed in Table S1.
376 Surfaces and volumes measurement of shrimp organs obtained from 3D anatomical
377 reconstructions are given in Table S3.

Author's contribution

PM collected shrimps during the BICOSE 2 expedition, carried out the identification and
measurement of specimens, realized Fluorescent in situ hybridization experiments on *R.*
hybisae specimens, conducted data analysis of 3D anatomical reconstruction, lead the
conception and design of the study, and drafted the original manuscript; MG conducted FISH
analyses on *R. exoculata* and *R. chacei*, and critically revised the manuscript; JTC collected
specimens from the JC82 expedition and critically revised the manuscript. HKW collected
specimens from YK13-05 and critically revised the manuscript. FP participated in the design
and conception of the study and critically revised the manuscript. MACB participated in the
design and conception of the study and critically revised the manuscript. CC collected
specimens from YK16-06 expeditions, assisted in drafting the original manuscript, participated
in the design and conception of the study, and coordinated the study. All authors gave final
approval for submission and publication of the manuscript in its current form.

Conflict of interests

392   We declare we have no competing interests.

Funding

394   PM was supported by a JAMSTEC Young Research Fellow fellowship. MACB and FP, are
395 supported by Ifremer, REMIMA project, MG by Ifremer and Region Bretagne PhD grant. CC and
396 HKW were supported by a Grant-in-Aid for Scientific Research (KAKENHI) from the Japan
397 Society for the Promotion of Science (JSPS), grant code 18K06401.

Figure Captions

Figure 1. 3D anatomical reconstructions of symbiont hosting organs (branchiostegite,
scaphognathites, and exopodites) and digestive organs (stomach, hepatopancreas, digestive
tube) in three *Rimicaris* shrimps across post-settlement ontogeny of (A-C) *R. hybisae*, (D-F) *R.*
chacei, (G-I) *R. exoculata*.

Figure 2. Relationship between body size (carapace length) and the relative per cent body
volumes or body surfaces of symbiont-hosting organs (branchiostegite, mouthparts) and
digestive organs (stomach, hepatopancreas).

Figure 3. Symbiont hosting organs in adult stages of the three *Rimicaris* shrimps. (A-C). 3D reconstructions of hosting organs in frontal view. (D-F) Dissected mouthparts covered with bacteriophore setae.

419 **Figure 4.** FISH observations of cephalothoracic cavities of *Rimicaris* subadults with universal
420 bacterial probes (yellow) on semi-thin sections (7µm) stained with DAPI (blue). A. *R. hybisae*
421 subadult B. *R. chacei* subadult C. *R. exoculata* subadult

References

1. Schwander T, Lo N, Beekman M, Oldroyd BP, Keller L. 2010 Nature versus nurture in social insect caste differentiation. *Trends Ecol. Evol.* **25**, 275–282. (doi:10.1016/j.tree.2009.12.001)

2. McGaugh SE *et al.* 2014 The cavefish genome reveals candidate genes for eye loss. *Nat. Commun.* **5**. (doi:10.1038/ncomms6307)

3. Tunnicliffe V, St Germain C, Hilário A. 2014 Phenotypic variation and fitness in a metapopulation of tubeworms (Ridgeia piscesae jones) at hydrothermal vents. *PLoS One* **9**. (doi:10.1371/journal.pone.0110578)

4. Urcuyo IA, Bergquist DC, MacDonald IR, VanHorn M, Fisher CR. 2007 Growth and longevity of the tubeworm Ridgeia piscesae in the variable diffuse flow habitats of the Juan de Fuca Ridge. *Mar. Ecol. Prog. Ser.* **344**, 143–157. (doi:10.3354/meps06710)

5. McFall-Ngai M *et al.* 2013 Animals in a bacterial world, a new imperative for the life sciences. *Proc. Natl. Acad. Sci.* **110**, 3229–3236. (doi:10.1073/pnas.1218525110)

6. Bosch TCG, McFall-Ngai M. 2021 *Animal development in the microbial world: Re-thinking the conceptual framework.* 1st edn. Elsevier Inc. (doi:10.1016/bs.ctdb.2020.11.007)

7. Hammer TJ, Moran NA. 2019 Links between metamorphosis and symbiosis in holometabolous insects. *Philos. Trans. R. Soc. B Biol. Sci.* **374**, 20190068. (doi:10.1098/rstb.2019.0068)

8. Carrier TJ, Bosch TCG. 2022 Symbiosis: The other cells in development. *Dev.* **149**. (doi:10.1242/dev.200797)

9. Nyholm S V., McFall-Ngai MJ. 2021 A lasting symbiosis: how the Hawaiian bobtail squid finds and keeps its bioluminescent bacterial partner. *Nat. Rev. Microbiol.* **19**, 666–679. (doi:10.1038/s41579-021-00567-y)

10. Dubilier N, Bergin C, Lott C. 2008 Symbiotic diversity in marine animals: the art of harnessing chemosynthesis. *Nat. Rev. Microbiol.* **6**, 725–40. (doi:10.1038/nrmicro1992)

11. Ikuta T *et al.* 2016 Surfing the vegetal pole in a small population: extracellular vertical transmission of an 'intracellular' deep-sea clam symbiont. *R. Soc. Open Sci.* **3**, 160130. (doi:10.1098/rsos.160130)

12. Bright M, Bulgheresi S. 2010 A complex journey: transmission of microbial symbionts. *Nat. Rev. Microbiol.* **8**, 218–230. (doi:10.1038/nrmicro2262)

13. Nussbaumer AD, Fisher CR, Bright M. 2006 Horizontal endosymbiont transmission in hydrothermal vent tubeworms. *Nature* **441**, 345–348. (doi:10.1038/nature04793)

14. Wentrup C, Wendeberg A, Schimak M, Borowski C, Dubilier N. 2014 Forever competent: Deep-sea bivalves are colonized by their chemosynthetic symbionts throughout their lifetime. *Environ. Microbiol.* **16**, 3699–3713. (doi:10.1111/1462-2920.12597)

15. Chen C, Linse K, Uematsu K, Sigwart JD. 2018 Cryptic niche switching in a chemosymbiotic gastropod. *Proc. R. Soc. B Biol. Sci.* **285**, 1–6. (doi:10.1098/rspb.2018.1099)

16. Franke M, Geier B, Hammel JU, Dubilier N, Leisch N. 2021 Coming together—symbiont

acquisition and early development in deep-sea bathymodioline mussels. *Proc. R. Soc. B*
Biol. Sci. **288**, 20211044. (doi:10.1098/rspb.2021.1044)

17.   Ponsard J *et al.* 2013 Inorganic carbon fixation by chemosynthetic ectosymbionts and
nutritional transfers to the hydrothermal vent host-shrimp Rimicaris exoculata. *ISME J.*
7, 96–109. (doi:10.1038/ismej.2012.87)

18.   Methou P, Hikosaka M, Chen C, Watanabe HK, Miyamoto N, Makita H, Takahashi Y,
Jenkins RG. 2022 Symbiont Community Composition in Rimicaris kairei Shrimps from
Indian Ocean Vents with Notes on Mineralogy. *Appl. Environ. Microbiol.* **88**.
(doi:10.1128/aem.00185-22)

19.   Guri M, Durand L, Cueff-Gauchard V, Zbinden M, Crassous P, Shillito B, Cambon-
Bonavita M-A. 2012 Acquisition of epibiotic bacteria along the life cycle of the
hydrothermal shrimp Rimicaris exoculata. *ISME J.* **6**, 597–609.
(doi:10.1038/ismej.2011.133)

20.   Zbinden M, Le Bris N, Gaill F, Compère P. 2004 Distribution of bacteria and associated
minerals in the gill chamber of the vent shrimp Rimicaris exoculata and related
biogeochemical processes. *Mar. Ecol. Prog. Ser.* **284**, 237–251.
(doi:10.3354/meps284237)

21.   Methou P, Michel LN, Segonzac M, Cambon-Bonavita M-A, Pradillon F. 2020 Integrative
taxonomy revisits the ontogeny and trophic niches of Rimicaris vent shrimps. *R. Soc.*
Open Sci. **7**, 200837. (doi:10.1098/rsos.200837)

22.   Gebruk A V., Southward EC, Kennedy H, Southward AJ. 2000 Food sources, behaviour,
and distribution of hydrothermal vent shrimps at the Mid-Atlantic Ridge. *J. Mar. Biol.*
Assoc. United Kingdom **80**, 485–499. (doi:10.1017/S0025315400002186)

23.   Methou P, Cueff-Gauchard V, Michel LN, Gayet N, Pradillon F, Cambon-Bonavita M-A.
2023 Symbioses of alvinocaridid shrimps from the South West Pacific: No
chemosymbiotic diets but partially conserved gut microbiomes Authors and Affiliations.
bioRxiv (doi:10.1101/2023.02.08.527621)

24.   Plouviez S, Jacobson A, Wu M, Van Dover CL. 2015 Characterization of vent fauna at the
mid-cayman spreading center. *Deep. Res. Part I Oceanogr. Res. Pap.* **97**, 124–133.
(doi:10.1016/j.dsr.2014.11.011)

25.   Versteegh EAA, Dover CL Van, Audenhaege L Van, Coleman M. 2022 Multiple
nutritional strategies of hydrothermal vent shrimp (Rimicaris hybisae) assemblages at
the Mid-Cayman Rise. *Deep. Res. Part I* (doi:10.1016/j.dsr.2022.103915)

26.   Nye V, Copley J, Plouviez S. 2012 A new species of Rimicaris (Crustacea: Decapoda:
Caridea: Alvinocarididae) from hydrothermal vent fields on the Mid-Cayman Spreading
Centre, Caribbean. *J. Mar. Biol. Assoc. United Kingdom* **92**, 1057–1072.
(doi:10.1017/S0025315411002001)

27.   Teixeira S, Olu K, Decker C, Cunha RL, Fuchs S, Hourdez S, Serrão EA, Arnaud-Haond S.
2013 High connectivity across the fragmented chemosynthetic ecosystems of the deep
Atlantic Equatorial Belt: Efficient dispersal mechanisms or questionable endemism? *Mol.*
Ecol. **22**, 4663–4680. (doi:10.1111/mec.12419)

28.   Streit K, Bennett SA, Van Dover CL, Coleman M. 2015 Sources of organic carbon for
Rimicaris hybisae: Tracing individual fatty acids at two hydrothermal vent fields in the

Mid-Cayman rise. *Deep. Res. Part I Oceanogr. Res. Pap.* **100**, 13–20.
(doi:10.1016/j.dsr.2015.02.003)

29.   Methou P, Hernández-Ávila I, Cathalot C, Cambon-Bonavita M-A, Pradillon F. 2022
Population structure and environmental niches of Rimicaris shrimps from the Mid-
Atlantic Ridge. *Mar. Ecol. Prog. Ser.* **684**, 1–20. (doi:10.3354/meps13986)

30.   Apremont V, Cambon-Bonavita M-A, Cueff-Gauchard V, François D, Pradillon F, Corbari
L, Zbinden M. 2018 Gill chamber and gut microbial communities of the hydrothermal
shrimp Rimicaris chacei Williams and Rona 1986: A possible symbiosis. *PLoS One* **13**,
e0206084. (doi:10.1371/journal.pone.0206084)

31.   Chen C, Kayama Watanabe H, Gena K, Johnson SB. 2022 Anatomical shifts linked with
unusual diets in deep-sea snails. *Ecology* , 1–6. (doi:10.1002/ecy.3847)

32.   Sogin EM, Leisch N, Dubilier N. 2020 Chemosynthetic symbioses. *Curr. Biol.* **30**, R1137–
R1142. (doi:10.1016/j.cub.2020.07.050)

33.   Nye V, Copley JTP, Tyler PA. 2013 Spatial Variation in the Population Structure and
Reproductive Biology of Rimicaris hybisae (Caridea: Alvinocarididae) at Hydrothermal
Vents on the Mid-Cayman Spreading Centre. *PLoS One* **8**.
(doi:10.1371/journal.pone.0060319)

34.   Koch EJ, Miyashiro T, McFall-Ngai MJ, Ruby EG. 2014 Features governing symbiont
persistence in the squid-vibrio association. *Mol. Ecol.* **23**, 1624–1634.
(doi:10.1111/mec.12474)

35.   Krishnan J, Rohner N. 2017 Cavefish and the basis for eye loss. *Philos. Trans. R. Soc. B
Biol. Sci.* **372**. (doi:10.1098/rstb.2015.0487)

36.   Rohner N, Jarosz DF, Kowalko JE, Yoshizawa M, Jeffery WR, Borowsky RL, Lindquist S,
Tabin CJ. 2013 Cryptic variation in morphological evolution: HSP90 as a capacitor for
loss of eyes in cavefish. *Science (80-.).* **342**, 1372–1375. (doi:10.1126/science.1240276)

37.   Barrett RDH, Schluter D. 2008 Adaptation from standing genetic variation. *Trends Ecol.
Evol.* **23**, 38–44. (doi:10.1016/j.tree.2007.09.008)

38.   Pfennig DW, Wund MA, Snell-Rood EC, Cruickshank T, Schlichting CD, Moczek AP. 2010
Phenotypic plasticity's impacts on diversification and speciation. *Trends Ecol. Evol.* **25**,
459–467. (doi:10.1016/j.tree.2010.05.006)

39.   Colosimo PF *et al.* 2005 Widespread parallel evolution in sticklebacks by repeated
fixation of ectodysplasin alleles. *Science (80-.).* **307**, 1928–1933.
(doi:10.1126/science.1107239)

40.   Lucek K, Sivasundar A, Seehausen O. 2014 Disentangling the role of phenotypic
plasticity and genetic divergence in contemporary ecotype formation during a biological
invasion. *Evolution (N. Y).* **68**, 2619–2632. (doi:10.1111/evo.12443)

41.   Koga R, Tanahashi M, Nikoh N, Hosokawa T, Meng XY, Moriyama M, Fukatsu T. 2021
Host's guardian protein counters degenerative symbiont evolution. *Proc. Natl. Acad. Sci.
U. S. A.* **118**, 1–9. (doi:10.1073/pnas.2103957118)

558