GroEL-proteotyping of bacterial communities by tandem mass spectrometry

Simon Klaes[1,2#], Shobhit Madan[1,3], Darja Deobald[1], Myriel Cooper[4], Lorenz Adrian[1,2]

[1]Department of Environmental Biotechnology, Helmholtz-Centre for Environmental Research (UFZ), Leipzig, Germany

[2]Chair of Geobiotechnology, Technische Universität Berlin, Berlin, Germany

[3]Faculty of Engineering, Ansbach University of Applied Sciences, Ansbach, Germany

[4]Chair of Environmental Microbiology, Technische Universität Berlin, Berlin, Germany

#corresponding author: Simon Klaes (simon.klaes (AT) ufz.de)

Running title: GroEL-proteotyping

Abstract

Profiling bacterial populations in mixed communities is a common task in microbiology. 16S rRNA-based sequencing is a widely accepted and functional approach but relies on amplification primers and cannot quantify isotope incorporation. Tandem mass spectrometry proteotyping is an effective alternative for taxonomically profiling microorganisms. We suggest that targeted proteotyping approaches can complement traditional population analyses. Therefore, we describe an approach to assess bacterial community compositions at the family level using the taxonomic marker protein GroEL, which is ubiquitously found in bacteria, except few obligate intracellular species. We refer to our method as GroEL-proteotyping. GroEL-proteotyping is based on high-resolution tandem mass spectrometry of GroEL peptides and identification of GroEL-derived taxa via a Galaxy workflow and a subsequent Python-based analysis script. Its advantage is that it can be performed with a curated and extendable sample-independent database and that GroEL can be pre-separated by SDS-PAGE to reduce sample complexity, improving GroEL identification while simultaneously decreasing the instrument time. GroEL-proteotyping was validated by employing it on a comprehensive raw data set obtained through a metaproteome approach from synthetic microbial communities as well as real human gut samples. Our data show that GroEL-proteotyping enables fast and straightforward profiling of highly abundant taxa in bacterial communities at reasonable taxonomic resolution.

1   Introduction

Bacterial communities govern our planet, impacting essential ecosystem services such as soil fertility [1], carbon dioxide fixation [2], bioremediation [3], and wastewater treatment [4]. One aspect of understanding the functionality of microbial communities is the identification of their microbial composition. Commonly, microbial compositions are described by (high-throughput) sequencing of 16*S* small subunit ribosomal-RNA (16*S* rRNA) gene amplicons. Several 16*S* rRNA-gene-based fingerprinting techniques, such as denaturing gradient gel electrophoresis (DGGE) [5] and terminal restriction fragment length polymorphism (T-RFLP) analyses [6], have been developed to allow the monitoring of the microbial community dynamics at reasonable costs and speed. However, all these 16*S* rRNA gene-based methods are PCR-dependent and introduce a primer bias [7]. Additionally, the variability and multiplicity of the 16*S* rRNA gene as well as the low taxonomic resolution of its short-reads, can distort bacterial community analyses [8,9]. Analyzing several hypervariable regions of the 16*S* rRNA gene [10] or other taxonomic marker genes, such as *rpoB* [11] or *groEL* [12], can enhance taxonomic resolution.

Metaproteomics is the mass spectrometric analysis of proteins from microbial communities [13]. Given the central role of proteins in metabolic processes, metaproteomics serves as a tool to investigate the functional dynamics of microbiomes, shedding light on microbial networks and their interplay with the environment [14–16]. In combination with isotope labeling, metaproteomics can link microbial populations to physiological activities, *e.g.*, to the use of a specific carbon source [17], and monitor uptake, incorporation, and interspecies transfer of isotopically labeled substrates by protein-based stable isotope probing (protein-SIP) [18]. Metaproteomic datasets are typically extensive and can be used to derive species-specific taxonomic profiles (proteotyping) within microbiomes based on all identified proteins [19–21]. In the classical proteotyping approach, a sample-specific protein database derived from (meta)genome sequencing is used for the accurate identification and quantification of proteins from mass spectrometric raw data [22,23]. When metagenomics data is not available, alternatives like RNA sequencing data [24] or universal, broad-spectrum databases such as NCBInr or UniProtKB/Swiss-Prot can be employed [25–30]. However, the large size of broad-spectrum databases leads to extended computation times, potentially hampering protein identification via target-decoy strategies [31]. Iterative cascade searches can reduce the effect of hampered protein identifications but may at the cost of increased computation times [32,33].

As an alternative, initial taxonomic profiling can assist in constructing comprehensive sample-specific databases for subsequent functional analyses [34]. Lower-resolution taxonomic profiles can be obtained using universal reference databases of taxonomic marker proteins and focusing only on a small subset of a comprehensive metaproteomic dataset. Specifically, highly abundant proteins, such as GroEL, translation initiation factor 2, elongation factors, or ribosomal proteins allowed profiling microbial communities at order level in human gut samples [34], and at domain level in mock assemblies and soil samples [35]. Utilizing a universal reference database of taxonomic marker proteins offers the advantage of eliminating the need to generate sample-specific metagenomic databases. This labor-intensive process requires repetition for each new sample site, albeit at the cost of potentially limiting the depth of functional insights. Furthermore, taxonomic marker databases tend to be smaller compared to universal, broad-spectrum databases, resulting in reduced computation time requirements.

Our research aimed to develop and evaluate a workflow enabling a semiquantitative characterization

of microbial communities with high taxonomic resolution utilizing tandem mass spectrometric data

focusing on GroEL as a taxonomic marker protein for bacteria. To achieve this objective, we created a

sample-independent GroEL database for peptide identification and developed a Python script that

facilitates protein and taxonomy inference. Our approach was evaluated by applying it to synthetic

microbial communities and metaproteome data obtained from the human gut, ensuring its robustness

and applicability across different microbial ecosystems.

2 Materials and methods

2.1 Cultivation, harvesting, and protein extraction

All cultivations were done in 100 mL cotton plugged Erlenmeyer flasks with 50 mL of nitrate-free DSMZ 586 medium at 30 °C in the dark on a rotary shaker under aerobic conditions. *Thauera aromatica* K172 and *Pseudomonas putida* KT2440 were separately grown and harvested during the early stationary phase by centrifuging at 16 000 x g for 10 min. Cell pellets were washed with 50 mM ammonium bicarbonate (AMBIC) buffer, pH 7.9. For the co-cultivation of *P. putida* and *T. aromatica*, we first inoculated the medium with approximately 10^6 cells mL^{-1} of *T. aromatica*. After 15 h of incubation, we injected approximately 10^5 cells mL^{-1} of *P. putida*. Samples were taken directly before adding *P. putida* and after 0; 18; 25; 39; and 48 h incubation time. Cells were harvested from 1 mL samples by centrifugation at 16 000 x g for 10 min, and the pellet was washed with 50 mM AMBIC buffer. Cell pellets were stored at -80 °C until further analysis. For protein extraction, cell pellets were resuspended in 50 mM AMBIC buffer and lysed by three freeze and thaw cycles, utilizing liquid nitrogen and a thermal shaker operating at 40 °C. Additionally, a 30 s treatment in a sonication bath was applied to enhance lysis efficiency. Cell debris and insoluble proteins were removed by centrifuging at 16 000 x g for 10 min. Protein concentrations were determined with the enhanced protocol of the bicinchoninic acid (BCA) assay kit (Pierce, Thermo Scientific). For synthetic bicultures, crude extracts of *T. aromatica* and *P. putida* were mixed in varying protein ratios, resulting in total protein content of 5 µg in 30 µL AMBIC buffer.

2.2 Sample preparation for protein mass spectrometry

Crude extract samples containing 5 µg protein in 30 µL AMBIC buffer were first amended with 40 ng bovine serum albumin (BSA) as a quality control measure. Then, the samples were sequentially treated with 62.5 mM dithiothreitol and 128 mM 2-iodoacetamide to reduce and alkylate cysteine residues. Subsequently, proteins were digested overnight with 0.63 µg trypsin (Promega). Tryptic digestion was stopped by adding formic acid to a final concentration of 1.8% (v/v). Undigested and precipitated proteins were removed by centrifugation at 16 000 x g for 10 min, and samples were dried by vacuum centrifugation. Subsequently, the peptides were resuspended in 0.1% (v/v) formic acid and desalted using C$_{18}$ Zip tips (Pierce, Thermo Scientific). The peptides were again dried by vacuum centrifugation before being reconstituted in 50 µL of 0.1% (v/v) formic acid for nano-liquid chromatography coupled to tandem mass spectrometry (nLC-MS/MS) analysis.

In separate experiments, we decreased the heterogeneity of the crude extracts before nLC-MS/MS analysis by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). For SDS-PAGE, the crude extract containing 5 µg protein in 30 µL AMBIC buffer was mixed with 10 µL SDS reducing buffer and incubated at 95 °C for 10 min to fully denature proteins. After cooling to room temperature, samples were subjected to SDS-PAGE at 110 V for 60–90 min and Coomassie stained. Protein bands with a molecular weight of approximately 60 kDa, corresponding to the size of GroEL, were excised. The gel slices were destained with acetonitrile. Then, proteins captured within the gel slice were reduced and alkylated by sequential incubation in 50 µL of 10 mM dithiothreitol and 50 µL of 100 mM 2-iodoacetamide for 60 min each. Subsequently, 40 ng of reduced and alkylated BSA was added as a quality control measure. Proteins were then digested with 0.1 µg trypsin (Promega) at 37

121 °C overnight. After digestion, peptides were extracted from gel pieces with 50% (v/v) acetonitrile and
122 5% (v/v) formic acid and dried by vacuum centrifugation. Next, peptides were resuspended in 20 µL of
123 0.1% (v/v) formic acid, desalted using C_{18} Zip tips (Pierce, Thermo Scientific), and repeatedly dried by
124 vacuum centrifugation. Finally, the desalted peptides were resuspended in 50 µL of 0.1% (v/v) formic
125 acid for nLC-MS/MS analysis.

2.3 Mass spectrometry

Desalted peptides were separated on an UltiMate 3000 RSLCnano high-performance nano-UPLC
system (Thermo Scientific) coupled to an Orbitrap Fusion Tribrid mass spectrometer (Thermo
Scientific) via a TriVersa NanoMate nano-electrospray ionization (nano-ESI) ion source (Advion). For
in-solution digested samples, 3 µL of the peptide solution was injected, whereas 5 µL was injected for
in-gel digested samples. Peptides were first loaded for 3 min onto the Acclaim PepMap 100 C_{18} trap
column (75 µm x 2 cm, 3 µm material, Thermo Scientific) with 5 µL min^{-1} of 3.2% (v/v) acetonitrile in
water at 0.1% formic acid. Then, the trap column was switched into line with the Acclaim PepMap 100
C_{18} analytical column (75 µm x 25 cm, 3 µm material, Thermo Scientific) heated up to 35 °C to
separate peptides at a flow rate of 0.3 µL min^{-1} using a gradient of 145 min (for in-solution digested
samples) or 60 min (for in-gel digested samples) from 3.2% to 72% (v/v) acetonitrile in water at 0.1%
(v/v) formic acid. The ion source operated in positive mode at a spray voltage of 2.4 kV and a source
temperature of 275 °C. The mass spectrometer run in data-dependent mode with a cycle time of 3 s.
Internal mass calibration was performed using a lock mass of 445.12003 *m/z*. Precursor ions were
scanned in the Orbitrap mass analyzer over a range of 350–2000 *m/z* with a resolution of 120 000,
automatic gain control (AGC) target of 4 x 10^5 ions, and a maximum injection time of 50 ms. Precursor
ions with a minimum intensity of 5 x 10^4 and charge state of +2 and +3 (for in-solution digested
samples) or +2 to +7 (for in-gel digested samples) were selected and isolated by the quadrupole in a
window of 1.6 *m/z* accumulating to an AGC target of 5 x 10^4 ions with a maximum injection time of 54
ms (for in-solution digested samples) and 120 ms (for in-gel digested samples). The isolated
precursor ions were fragmented using higher energy collisional dissociation (HCD) at 30% relative
collision energy. Fragment ions were scanned with the Orbitrap mass analyzer at a resolution of
30 000 (for in-solution digested samples) or 60 000 (for in-gel digested samples), respectively.
Precursor ions with the same mass (±10 ppm) were excluded for further precursor selection for 30 s.

2.4 Databases for mass spectrometric analysis

We assembled a comprehensive dataset of bacterial GroEL protein sequences by downloading all
available amino acid sequences from the National Center for Biotechnology Information (NCBI) in
GenBank format on September 7, 2021 [36]. The dataset was used to generate a file in fasta format
as input for peptide identification search engines (see below) and to generate a second file with tab-
separated values (tsv) for protein and taxonomy inferences. The tsv-database included in separated
columns accession number, amino acid sequence, expected tryptic peptides, and taxonomic
classification at various levels (kingdom, phylum, class, order, family, and genus). Tryptic peptides
were calculated with pyOpenMS [37], accepting two missed cleavages and a peptide length of 6–144
amino acids. The taxonomic classification was updated by cross-referencing the "List of Prokaryotic
names with Standing in Nomenclature" (LPSN) on October 18, 2022 [38]. We also included GroEL
sequences from the separately published genomes of *Roseobacter* sp*.* AK199 and *Chromobacterium*

violaceum CV026 [20]. The common Repository of Adventitious Proteins (cRAP) was appended to the
database to identify common contaminants. Additionally, we downloaded the whole proteome
databases of *T. aromatica* K172 (CP028339.1) and *P. putida* KT2440 (NC_002947.4) from NCBI. The
detectability of GroEL peptides by tandem mass spectrometry was assessed using DeepMSPeptide
[39].

2.5   Analysis of mass-spectrometric raw data files

Thermo raw files were first converted into mzML format using msConvert (ProteoWizard) [40]. Next,
we analyzed the mzML files using a customized MetaProSIP workflow [41] on the Galaxy platform
[42]. Briefly, peptides were identified using MS-GF+ [43] and MetaProSIP [44]. We accepted two
missed trypsin cleavages, peptide lengths of 6–40 amino acids, and precursor *m/z* deviations of
5 ppm. Oxidation of methionine and carbamidomethylation of cysteine were set as dynamic and static
modifications, respectively. The false discovery rate of identified peptide sequences was kept below
0.01, based on the SpecEValue calculated by MS-GF+. The peptide-centric output of MetaProSIP
was used as input for a custom GroEL-proteotyping Python script. The isobaric amino acids leucine
(L) and isoleucine (I) were treated as indistinguishable. Peptides listed in the cRAP database were
excluded from the analysis with the customized GroEL-proteotyping Python script. The Python script
assigns detected peptides to each possible GroEL protein in the tsv-database based on matching
sequences. GroEL proteins with the same set of detected peptides are merged into protein groups
and sorted by the number of detected peptides in descending order. For each GroEL protein group,
only the detected peptides not present in another GroEL protein group containing more peptides are
counted, resulting in a Top Rank Count. GroEL protein groups are then filtered by the Top Rank
Count with a threshold (≥ 5), and the taxonomy of the remaining GroEL protein groups is read from
the tsv-file. For each taxonomic level, the most frequent taxonomic description and its frequency
within the GroEL protein group are calculated. The most frequent taxonomic description is used to
merge GroEL protein groups into taxonomic groups at the level of interest, which are then quantified
by the sum of the MS1 precursor ion intensities (based on peak area) or the number of non-redundant
peptides within the taxonomic group. In the current version of the Python script (version 1.0.0), non-
redundant peptides are only counted once for the same taxonomic group and multiple times for
different taxonomic groups. Furthermore, the precursor intensity of peptides assigned to multiple
taxonomic groups is distributed proportionately to the total number of peptides assigned to each
group. To evaluate our workflow, we used published metaproteome data from synthetic microbial
communities (PRIDE repository PXD006118). Specifically, we used the raw data acquired with an LC
gradient of 260 min (run 4 and 5) as this provided sufficient data for community analysis [20]. The
identified peptides of technical replicates were pooled before running our GroEL-proteotyping Python
script. Furthermore, we applied our analysis workflow to a metaproteome dataset of human gut
microbiota (PRIDE repository PXD020786) [45]. Identified peptides of biological replicates were
pooled before running the GroEL-proteotyping Python script to allow comparison with the original
study (Supplementary Table 3 of García-Durán et al. [45]).

2.6   Data availability

The mass spectrometric raw data and protein sequence databases have been deposited to the
ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the PRIDE partner

repository [46] with the data set identifier PXD043106 (Username: reviewer_pxd043106@ebi.ac.uk;

Password: KzeQ5W8r). The GroEL-proteotyping Python script is publicly available on GitLab

(https://git.ufz.de/klaes/groel-proteotyping/). Any other relevant data will be made available upon

request.

3 Results

3.1 Establishing a GroEL database and analyzing protein and peptide sequences

Our download of bacterial GroEL sequences from NCBI in September 2021 resulted in 284 351 GroEL homologous sequences, of which 72 759 were non-redundant. We restricted the download to sequences of a length of 6–1500 amino acids, resulting in a median length of 542 amino acids in the database. Prediction of trypsin digestion sites in the non-redundant protein sequences allowing up to two missed cleavages, a length of 6–144 amino acids, and no ambiguous amino acid codes (B, J, O, U, X, or Z) resulted in 9 091 897 peptides, of which 1 875 469 were non-redundant.

TABLE 1. Absolute and relative numbers of peptides with a standardized taxon name according to LPSN and taxon-specific peptides in the GroEL database at different taxonomic levels.

Number of non-redundant GroEL peptides	Taxonomic level				
	Phylum	Class	Order	Family	Genus
Standardized taxon name[a]	1 798 737 (95.9%)	1 564 290 (83.4%)	1 410 836 (75.2%)	1 343 631 (71.6%)	1 315 614 (70.1%)
Taxon-specific[b]	1 732 937 (96.3%)	1 485 993 (95.0%)	1 233 553 (87.4%)	1 205 161 (89.7%)	1 122 799 (85.3%)
Detectable by tandem mass spectrometry[c]	690 492 (39.8%)	587 287 (39.5%)	484 600 (39.3%)	470 112 (39.0%)	435 776 (38.8%)

[a]: Relative numbers based on the total number of non-redundant peptide sequences.
[b]: Relative numbers based on the number of peptides affiliated with an organism with a standardized taxon name.
[c]: Relative numbers based on the number of taxon-specific peptides. Detectability was predicted using DeepMSPeptide [39]

To evaluate the taxonomic relevance of GroEL-derived peptides, we calculated their taxon-specificity across different taxonomic levels. Additionally, we predicted their MS-detectability using DeepMSPeptide [39]. For this purpose, we initially standardized taxon names in our database according to the nomenclature provided by the "List of Prokaryotic names with Standing in Nomenclature" (LPSN). Taxon names matching the LPSN nomenclature were kept, while taxon names not matching the LPSN nomenclature were excluded. Subsequently, peptides exclusively attributed to a single taxon were counted as taxon-specific peptides, with leucine (L) and isoleucine (I) treated as indistinguishable. Our results show a decline in the count of peptides with standardized taxon names, the number of taxon-specific peptides, and their predicted detectability with increasing taxonomic resolution (Table 1).

3.2 Evaluating the sensitivity and specificity of GroEL-proteotyping and its protein filtering routine

To evaluate the performance of our database in detecting GroEL peptides, we analyzed pure cultures of *T. aromatica* K172 and *P. putida* KT2440. In this regard, each microorganism was separately cultivated, followed by protein extraction and tryptic digestion. The resulting peptides were analyzed by mass spectrometry and evaluated against two databases: *i)* a database representing the full proteome of *T. aromatica* K172 or *P. putida* KT2440, respectively, and *ii)* our GroEL database.

 TABLE 2. Peptides identified from protein extracts of *Thauera aromatica* K172 and *Pseudomonas putida*
KT2440. Peptides were identified using the whole proteome database of *T. aromatica* K172 ("K172") containing
3 335 entries or *P. putida* KT2440 ("KT2440") with 5 450 entries, respectively, or the GroEL database with 72
759 non-redundant entries. Values represent means of biological triplicates ± standard deviations.

Microorganism	Database: whole proteome		Database: GroEL	
	Identified peptides	Identified GroEL peptides	Identified GroEL peptides	Mean intensity of GroEL peptides (x 10^6)
T. aromatica **K172**	2441.0 ± 201.5	19.7 ± 1.2	14.7 ± 0.5	540 ± 120
P. putida **KT2440**	2961.7 ± 152.9	11.7 ± 1.7	8.3 ± 2.1	500 ± 20

When comparing our data against the whole proteome databases, the total number of identified
peptides was higher for *P. putida*, but the count of identified GroEL peptides was lower compared to
T. aromatica (Table 2). When using the GroEL database instead of the whole proteome database, we
observed a slight decrease in the number of detected GroEL peptides for both strains. However, the
average precursor peak intensity for GroEL peptides was similar for both organisms. In summary, this
initial experiment showed that in pure cultures, GroEL peptides of *T. aromatica* K172 exhibited
stronger signal intensities than GroEL peptides of *P. putida* KT2440. Furthermore, the use of our
GroEL database resulted in a slightly decreased detection of GroEL peptides.

To evaluate the applicability of GroEL peptide mass spectrometry for relative quantification of
subpopulations, we experimented with a synthetic mixture of *T. aromatica* K172 and *P. putida*
KT2440. Initially, each of the two strains was cultivated separately. Subsequently, proteins were
extracted from them and crude extracts of *T. aromatica* K172 and *P. putida* KT2440 were mixed in
predetermined protein ratios, ensuring a total protein mass of 5 µg. Before nLC-MS/MS analysis,
GroEL was separated from other proteins by SDS-PAGE, decreasing the heterogeneity of the
sample, which resulted in a 1.8-fold increase in the number of detected GroEL peptides. Peptides
often can be assigned to multiple proteins, known as the protein inference problem. Furthermore,
large databases tend to increase the number of false-positive detections. To remove false-positive
detections and to infer proteins and taxonomy unambiguously, we developed a custom Python script
that filters GroEL protein groups by the Top Rank Count, which only counts peptides once for the
largest GroEL protein group (for the description of this filtering routine, see Materials & Methods).

Various Top Rank Count thresholds were systematically evaluated to eliminate false positive
identifications while retaining the ability to detect low-abundant organisms (Figure 1). The
implementation of a Top Rank Count threshold ≥ 5 eliminated all false-positive identifications at the
genus level. Notably, both organisms were consistently detected at the genus level in all biological
replicates at different protein ratios, demonstrating a relative detection limit of 1% of total protein
content for low abundant organisms. Unfortunately, quantification based on the number of detected
GroEL peptides yielded relative abundance estimations for the two genera close to 50% across all
ratios, failing to reflect the actual mixing ratios. Quantification based on the sum of the precursor
intensities from detected GroEL peptides provided more accurate estimations. In mixtures containing
1%, 5%, and 10% *T. aromatica K172* proteins, we observed *Thauera* abundance values of 11.9% ±
3.1%, 8.9% ± 1.1%; and 22.4% ± 0.6% respectively.

FIGURE 1. Evaluating sensitivity **(A)** and specificity **(B)** of GroEL-proteotyping in detecting organisms at the genus level in mixed protein extracts. Mixed protein extracts were composed of *Thauera aromatica* K172 and *Pseudomonas putida* KT2440 proteins in different ratios, and different filtering criteria were applied. Bars represent means of three biological replicates.

Taken together, our findings indicate that GroEL-proteotyping allows for the differentiation of two organism mixtures at the genus level, achieving a relative detection limit of 1% of the total protein content, representing approximately 50 ng protein in the crude extract. Quantification based on the sum of the precursor intensities proved to be more reliable and closer to the actual mixing ratios than GroEL peptide count-based quantification. However, in all mixtures, we consistently detected a higher relative abundance of *Thauera* than originally added.

3.3 GroEL-proteotyping in action I: Reanalyzing raw data of synthetic microbial communities

In the previous experiments, communities of only two bacteria were investigated. To examine if the filtering and quantification techniques are applicable for characterizing more complex bacterial consortia, we reanalyzed proteomic raw data files of three synthetic microbial communities originally assembled by Kleiner *et al.* [20]. The synthetic communities comprised crude protein extracts of one archaeon, one eukaryote, five bacteriophages, three strains of Gram-positive bacteria, and 18 (Mock A and B) or 22 (Mock C) strains of Gram-negative bacteria mixed in different ratios (Supplementary Tables 1–3 of Kleiner et al. [20]). The bacterial population contained 12 (Mock A and B) to 15 (Mock C) bacterial families. Since our method is specialized in bacterial community characterization, we focused only on these bacterial families.

We again investigated different thresholds of the Top Rank Count to eliminate false-positive identifications while still detecting low-abundance microorganisms (Figure 2). Filtering by a Top Rank Count threshold ≥ 5 (as described above) resulted in some false identifications at the genus level. However, at the family level, this filter effectively eliminated all incorrect identifications while maintaining an acceptable sensitivity for detecting low-abundance organisms. Therefore, we used this threshold for subsequent community analysis at the family level.

FIGURE 2. Evaluating sensitivity **(A)** and specificity **(B)** of GroEL-proteotyping using different Top Rank Count filtering thresholds in detecting peptides assigned to bacterial families that were present in three synthetic microbial communities (Mock A–C) assembled by Kleiner *et al.* [20]. The detected peptides were assigned to specific GroEL proteins. GroEL proteins with the same set of detected peptides were merged into GroEL groups, which were then filtered by the Top Rank Count ≥ 5. Bars represent the means of four biological replicates.

In the raw data files of the three synthetic communities assembled by Kleiner *et al.* [20], we detected all bacterial families that were present at a relative bacterial protein content above 2.8%, while no absent family was falsely detected (Figure 3A). However, the detection of scarce families was inconsistent between the assessments of the different synthetic communities. Specifically, we detected all bacterial families in all four biological replicates of Mock B, which has the most balanced actual protein distribution of 4.8–19.1% of the total protein content for each family (Figure 3A, yellow).

In Mock A, all families except the lowest-abundant family *Staphylococcaceae*, accounting for 0.7% of the total bacterial protein content, were detected (Figure 3A, red). On the other hand, in Mock C, which had the highest number of low-abundant bacterial families (relative bacterial protein content ≤ 2.8%), and the most uneven actual protein distribution, with families representing 0.2–41.8% of the total bacterial protein content, we detected all highly abundant bacterial families (relative bacterial protein content > 2.8%) (Figure 3A, blue). However, the detection of low-abundant families with relative bacterial protein content ≤ 2.8% was inconsistent: we detected *Chromobacteriaceae* (1.3%), *Desulfovibrionaceae* (1.0%), *Rhodobacteraceae* (1.0%), *Roseobacteraceae* (1.7%), and *Thermaceae* (1.8%), but not *Staphylococcaceae* (2.8%), *Alteromondaceae (1.0%)*, *Bacillaceae (0.8%)*, *Nitrosomonadaceae (0.7%)*, and *Nitrospiraceae (0.2%)* (Figure 3A, blue).

FIGURE 3. Profiling the taxonomic composition of bacterial subpopulations in synthetic microbial communities using GroEL-proteotyping. The metaproteome dataset of synthetic communities was obtained from Kleiner *et al.* [20]. We used the same graphic layout to facilitate comparison. **(A)** Comparison of actual community shares (shaded values) with GroEL-based quantification: 'Peptide number' – quantification based on the number of non-redundant GroEL peptides assigned to that family; 'Intensities' – quantification based on the sum of the precursor intensities of the detected GroEL peptides. Quantities are represented as bubble areas. The figure includes information on families that were not added (NA) or not detected (ND) with our method. Results are means of four replicates. **(B)** Comparison of two quantification methods for GroEL peptides to the actual protein abundance of each family. $x\text{-}fold\ deviation\ from\ the\ expected\ value = \frac{measured\ value}{expected\ value} - 1$. Boxes show the 1st and 3rd quartile, the median, and the whiskers indicating the 10th and 90th percentile, with filled circles representing outliers. ND were removed before plotting. A value of 0 is depicted as a dotted line, indicating equal measurement and input. Negative values indicate underestimation, while positive values indicate overestimation.

In addition, we compared two quantification methods: *i)* analyzing the number of detected GroEL peptides and *ii)* analyzing the sum of the precursor intensities of detected GroEL peptides (Figure 3B). Both methods performed similarly for Mock A and Mock B, with the medians centering around zero, indicating high agreement of our analysis and input (Figure 3B, red and yellow). However, differences between the two quantification methods became apparent for Mock C. The method based on peptide counts overestimated most families (*e.g.*, *Thermaceae*) while underestimating others (*e.g.*, *Enterobacteriaceae*) (Figure 3B, blue). The deviation of the measured abundance from the actual input was smaller for the method based on precursor intensities than the method based on peptide counts. Overall, we show that the characterization of complex synthetic communities by GroEL-proteotyping is robust at the family level and more consistent when based on the sum precursor intensities of detected GroEL peptides than when based on peptide counts.

3.4 GroEL-proteotyping in action II: Reanalyzing raw data of human gut microbiomes

To test the applicability of GroEL-proteotyping for characterizing complex bacterial communities, we reanalyzed a gut metaproteomic dataset previously published by García-Durán *et al.* [45] pertained to six healthy individuals. We focused on the human gut microbiome because it harbors a diverse bacterial community of up to 1 150 species [47]. The original study used a protein database based on the human gut microbial gene catalog (9 878 647 sequences) [48], and human proteins from UniProt

(74 451 sequences) to detect an average of 11 712 peptides per sample, 56% of which were
assigned at the family level. García-Durán *et al.* identified 33 different bacterial families, of which 11
showed a relative family abundance of at least 1% (Supplementary Table 3 of García-Durán *et al.*
[45]).

In our reanalysis, we used our GroEL database to detect peptides and assign them to GroEL protein
groups. We filtered GroEL protein groups by the Top Rank Count with a threshold ≥ 5 and evaluated
the taxonomy at the family level to achieve high specificity. Our reanalysis closely mirrored the
identification of abundant families reported by García-Durán *et al.* (Figure 4) [45]. We detected all 11
families with a relative family abundance of at least 1% (in the original publication). Among the 22
families exhibiting a relative family abundance below 1%, our analysis successfully identified three
families: *Streptococcaceae*, *Enterobacteriaceae*, and *Coprobacillaceae* (identified as
Erysipelotrichaceae by [45]) (Table S1). We did not detect the remaining 19 families, identified with a
relative abundance below 1% in the original publication. We observed only minor differences in the
abundance of highly abundant families (at least 1% in the original publication), depending on the
quantification parameter used (peptide count, peptide spectrum matches (PSMs), or sum precursor
intensities). However, for a few families, we noted a substantial discrepancy between our
quantification based on GroEL and the quantification based on the human gut microbial gene catalog
applied in the original publication. Specifically, GroEL-proteotyping led to an overestimation of
Lachnospiraceae and *Clostridiaceae* and an underestimation of *Bacteriodaceae*, *Eubacteriaceae*, and
Prevotellaceae comparing to the original analysis (Figure 4).

FIGURE 4. The average composition of bacterial communities at the family level, as described from the gut
metaproteomic dataset of six healthy individuals by García-Durán *et al.* [45], compared with reanalysis of the
same data set using GroEL-proteotyping. Peptides that could not be assigned at the family level were excluded.
In the original analysis, the relative abundance of a family was determined by the number of peptide spectrum
matches (PSMs) or unique peptides assigned to that family as a proportion of the total number of PSMs or
unique peptides assigned at the family level, respectively. In our analysis, the relative abundance of a family
was calculated based on the sum of the precursor intensities or the number of non-redundant GroEL peptides
assigned to that family, relative to the total sum of the precursor intensities or the number of all filtered GroEL
peptides. Families with an abundance below 1% in the original analysis were merged and represented as 'other
families'.

In the second case study, we demonstrate that GroEL-proteotyping can be applied to real and
complex samples, as it yields results comparable to traditional metaproteomic approaches in
detecting highly abundant bacteria. However, we also identify certain limitations associated with our
method. Specifically, there are challenges in accurately quantifying and reliably detecting low-
abundance bacterial families.

4 Discussion

Our findings show that compositions of bacterial communities can be analyzed by using GroEL as a taxonomic marker protein and a sample-independent database. This aspect holds substantial advantages, particularly in situations where rapid or continuous monitoring of community compositions is required. In our work, highly abundant families were reliably identified in all tested samples. We propose a flexible workflow (Figure 5) that can be adapted to a variety of sample preparations, nLC-MS/MS setups, peptide search engines, and quantification strategies. This workflow facilitates the analysis and can be automatized, *e.g.*, as a Galaxy workflow. It provides quantitative identifications with both, sample-independent or sample-specific databases. While the sample-independent GroEL database gave satisfactory results on the family level, a sample-specific GroEL database can improve identifications, as shown with our defined bicultures. We propose introducing the general concept of 'targeted proteotyping' as a distinct subcategory of proteotyping, that can be applied to different taxonomic marker proteins. Adopting the term 'GroEL-protetyping would then differentiate this particular method from other (targeted) proteotyping approaches.

The analysis of our database shows that most tryptic GroEL peptides are highly taxon-specific similar to the nucleotide sequence of the *groEL* gene that has already been established as a barcode for bacteria [12]. Likewise, our workflow identified bacteria at the genus level in low-complexity samples, such as bicultures composed of *T. aromatica* K172 and *P. putida* KT2440. In more complex samples, the identification was reliable and robust at the family level. The classical metaproteomic approach allows for characterizing the same sample down to the species level [20]. However, this enhanced resolution is achieved at the expense of necessitating a sample-specific metagenomic database. In contrast, phylopeptidomics, a peptide-centric approach, achieves species-level characterization of the same sample but using a large, sample-independent NCBInr database, resulting in high computation demands [26]. Previous analyses based on GroEL or other taxonomic marker proteins without additional filtering procedures could differentiate the mock communities only at kingdom level [35]. This demonstrates the importance of effective peptide filtering and protein/taxonomy inference. Here, we achieved a higher taxonomic resolution by employing a filtering strategy for GroEL protein groups based on the number of peptides counted for top-scored proteins, which we refer to as 'Top Rank Count' (≥ 5). While this approach is not entirely novel, as similar filtering techniques have been implemented by Proteome Discoverer (Thermo Scientific) and MassSieve [49], its integration into our GroEL-proteotyping workflow enabled us to attain superior taxonomic resolution compared to previous GroEL-based approaches. Although GroEL-proteotyping currently has a lower taxonomic resolution and provides limited information beyond taxonomic composition, it stands out for its advantages in terms of speed and cost-efficiency compared to metaproteomics and phylopeptidomics. These benefits arise from a smaller sample-independent database and the reduced sample complexity allowing shorter instrument run times.

Taxa quantification was more robust based on the sum of the precursor intensities compared to the peptide count and a good semiquantitative estimate for the actual protein input in the complex synthetic communities. Surprisingly, none of the quantification approaches accurately reflected the relative protein amount of the strains in theIn our synthetic biculture, we consistently observed a higher signal for GroEL peptides of *Thauera* compared to those of *Pseudomonas*. One reason could be indicating that the relative GroEL expression differs between microbes, growth phases, and

environmental conditions and that the difference became more apparent for the biculture setup than in
the complex communities [50–52].

FIGURE 5. Proposed workflow for analyzing mixed bacterial communities using GroEL-proteotyping. The
workflow is adaptable to various sample preparations, nano-liquid chromatography-tandem mass spectrometry
(nLC-MS/MS) configurations, peptide search engines, and quantification strategies. While SDS-PAGE pre-
separation of GroEL proteins can enhance the detection limit, it is not compulsory. The lower part explains the
Top Rank Count approach used to filter hits. See the text for further explanations.

Our data indicate that the detection of low-abundant taxa strongly depends on the sample complexity
and protein distribution across taxa. In a complex synthetic community with an uneven protein
distribution, we only reliably detected bacterial families with a relative protein mass of more than
2.8%. However, pre-separation of GroEL by SDS-PAGE increased the identification of GroEL
peptides 1.8-fold, resulting in a relative detection limit of 1% in low-complex bicultures. Thus, we
hypothesize that using a separable marker protein allows for reducing sample complexity without
losing taxonomic information, consequently enhancing the detection of low-abundant taxa. Larger
protein input, pooling of gel bands, and longer LC gradients during mass spectrometry may further
improve the detection of low-abundant taxa. A fast and sensitive screening of taxa present in a
sample based on marker proteins could also aid in creating sample-specific databases for subsequent
functional analysis of the whole metaproteomic data as shown before [34].

In our approach, the detection of organisms depends on the presence of its GroEL sequence in the
database. For example, *Roseobacteraceae* was only detected in complex synthetic communities after
adding the GroEL sequence of *Roseobacter* sp. AK199 to the database, demonstrating that
incomplete databases bias the identification and quantification of taxa. Thus, applying GroEL-
proteotyping to environmental samples containing many non-sequenced organisms is still challenging
[53]. However, we successfully applied GroEL-proteotyping to human gut proteome data.
Furthermore, we are confident that the rapid growth of sequence databases will massively increase
database coverage. In addition, universal primers can amplify a 549–567 bp region of the *groEL*
gene, allowing a targeted, fast, and sample-specific extension of the database [54].

Our study introduces targeted proteotyping as a concept for proteotyping microbial communities using
taxonomic marker proteins. At present, our targeted proteotyping approach is limited to bacteria
because GroEL is highly abundant in bacteria [35,55–59] (except very few intracellular *Mycoplasma*
and *Ureaplasma* strains [60]), while it is only found in some archaea that most likely acquired it
through horizontal gene transfer [61]. Consequently, the current scope of our approach does not
encompass the detection of eukaryotes and archaea. To expand the applicability of our method to
also detect eukaryotes and archaea, the incorporation of additional putative marker proteins such as
ribosomal proteins, chaperonin TCP-1, or thermosome proteins should be considered.

In summary, we introduce GroEL-proteotyping as a rapid and cost-effective method for protein-based
profiling of bacterial communities at the family level. In comparison to classical protein-based
approaches, GroEL-proteotyping bypasses the need for sample-specific databases, saving time and
reducing costs associated with database generation while achieving higher taxonomic resolution than
previous targeted proteotyping approaches [34,35]. Although the implementation of our method
requires access to a protein mass spectrometer, the actual analysis process, with a 60 min LC

gradient, is fast, which enables the characterization of up to 20 complex samples per day, making our
approach highly efficient in particular for high-throughput analyses. Furthermore, our automatable
bioinformatics workflow enables the taxonomic profiling of bacterial communities within 48 hours. This
can particularly be advantageous for monitoring defined or highly enriched mixed communities over
time. Moreover, as the field progresses very fast, the applicability of GroEL-proteotyping will expand
with increasing GroEL protein sequence databases, the development of automated workflows, the
combination with stable-isotope probing, and the optimization of mass spectrometric techniques such
as ion mobility devices [62].

5 Acknowledgments

This project was funded by the German Research Foundation (DFG) as part of the Research Training Group "Urban Water Interfaces (UWI)" (GRK 2032/2, Project H8: "Microbial transformation of mobile halogenated aromatics in redox gradients of urban hyporheic zones"). Open access was supported by the UFZ library.

We acknowledge Benjamin Scheer for assistance with mass spectrometric measurements. Furthermore, we thank Matthias Bernt and Bryson Gibbons for rapid troubleshooting regarding the Galaxy server and MS-GF+.

6 Author contributions

Conceptualization, Simon Klaes, Darja Deobald, Myriel Cooper and Lorenz Adrian; Data curation, Simon Klaes; Formal analysis, Simon Klaes; Funding acquisition, Myriel Cooper and Lorenz Adrian; Investigation, Simon Klaes and Shobhit Madan; Methodology, Simon Klaes, Darja Deobald and Lorenz Adrian; Software, Simon Klaes and Lorenz Adrian; Validation, Simon Klaes and Shobhit Madan; Visualization, Simon Klaes; Writing – original draft, Simon Klaes; Writing – review & editing, Simon Klaes, Shobhit Madan, Darja Deobald, Myriel Cooper and Lorenz Adrian.

7 Competing interests

The authors declare that they have no competing financial interests.

8 References

1. Siciliano, S.D.; Palmer, A.S.; Winsley, T.; Lamb, E.; Bissett, A.; Brown, M. V.; van Dorst, J.; Ji, M.; Ferrari, B.C.; Grogan, P.; et al. Soil fertility is associated with fungal and bacterial richness, whereas pH is associated with community composition in polar soil microbial communities. *Soil Biol Biochem* **2014**, *78*, 10–20, doi:10.1016/J.SOILBIO.2014.07.005.

2. Berg, I.A. Ecological aspects of the distribution of different autotrophic CO_2 fixation pathways. *Appl Environ Microbiol* **2011**, *77*, 1925–1936, doi:10.1128/AEM.02473-10.

3. Lopes, P.R.M.; Cruz, V.H.; de Menezes, A.B.; Gadanhoto, B.P.; Moreira, B.R. de A.; Mendes, C.R.; Mazzeo, D.E.C.; Dilarri, G.; Montagnolli, R.N. Microbial bioremediation of pesticides in agricultural soils: an integrative review on natural attenuation, bioaugmentation and biostimulation. *Rev Environ Sci Biotechnol* **2022**, *21*, 851–876, doi:10.1007/s11157-022-09637-w.

4. Al Ali, A.A.; Naddeo, V.; Hasan, S.W.; Yousef, A.F. Correlation between bacterial community structure and performance efficiency of a full-scale wastewater treatment plant. *J Water Process Eng* **2020**, *37*, 101472, doi:10.1016/J.JWPE.2020.101472.

5. Muyzer, G.; De Waal, E.C.; Uitterlinden, A.G. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16*S* rRNA. *Appl Environ Microbiol* **1993**, *59*, 695, doi:10.1128/AEM.59.3.695-700.1993.

6. Liu, W.T.; Marsh, T.L.; Cheng, H.; Forney, L.J. Characterization of microbial diversity by determining terminal restriction fragment length polymorphisms of genes encoding 16*S* rRNA. *Appl Environ Microbiol* **1997**, *63*, 4516, doi:10.1128/AEM.63.11.4516-4522.1997.

7. Brooks, J.P.; Edwards, D.J.; Harwich, M.D.; Rivera, M.C.; Fettweis, J.M.; Serrano, M.G.; Reris, R.A.; Sheth, N.U.; Huang, B.; Girerd, P.; et al. The truth about metagenomics: quantifying and counteracting bias in 16*S* rRNA studies. *BMC Microbiol* **2015**, *15*, 66, doi:10.1186/s12866-015-0351-6.

8. Větrovský, T.; Baldrian, P. The variability of the 16*S* rRNA gene in bacterial genomes and its consequences for bacterial community analyses. *PLoS One* **2013**, *8*, e57923, doi:10.1371/JOURNAL.PONE.0057923.

9. Johnson, J.S.; Spakowicz, D.J.; Hong, B.Y.; Petersen, L.M.; Demkowicz, P.; Chen, L.; Leopold, S.R.; Hanson, B.M.; Agresta, H.O.; Gerstein, M.; et al. Evaluation of 16*S* rRNA gene sequencing for species and strain-level microbiome analysis. *Nat Commun* **2019**, *10*, 5029, doi:10.1038/s41467-019-13036-1.

10. Liu, Z.; Desantis, T.Z.; Andersen, G.L.; Knight, R. Accurate taxonomy assignments from 16*S* rRNA sequences produced by highly parallel pyrosequencers. *Nucleic Acids Res* **2008**, *36*, e120, doi:10.1093/NAR/GKN491.

11. Ogier, J.C.; Pagès, S.; Galan, M.; Barret, M.; Gaudriault, S. rpoB, a promising marker for analyzing the diversity of bacterial communities by amplicon sequencing. *BMC Microbiol* **2019**, *19*, 171, doi:10.1186/s12866-019-1546-z.

12. Links, M.G.; Dumonceaux, T.J.; Hemmingsen, S.M.; Hill, J.E. The chaperonin-60 universal target is a barcode for bacteria that enables *de novo* assembly of metagenomic sequence data. *PLoS One* **2012**, *7*, e49755, doi:10.1371/journal.pone.0049755.

13. Kleiner, M. Metaproteomics: much more than measuring gene expression in microbial communities. *mSystems* **2019**, *4*, e00115-19, doi:10.1128/msystems.00115-19.

14. Heyer, R.; Schallert, K.; Siewert, C.; Kohrs, F.; Greve, J.; Maus, I.; Klang, J.; Klocke, M.; Heiermann, M.; Hoffmann, M.; et al. Metaproteome analysis reveals that syntrophy, competition, and phage-host interaction shape microbial communities in biogas plants. *Microbiome* **2019**, *7*, 69, doi:10.1186/s40168-019-0673-y.

15. Li, L.; Ryan, J.; Ning, Z.; Zhang, X.; Mayne, J.; Lavallée-Adam, M.; Stintzi, A.; Figeys, D. A functional ecological network based on metaproteomics responses of individual gut microbiomes to resistant starches. *Comput Struct Biotechnol J* **2020**, *18*, 3833–3842, doi:10.1016/j.csbj.2020.10.042.

16. Shrestha, H.K.; Appidi, M.R.; Villalobos Solis, M.I.; Wang, J.; Carper, D.L.; Burdick, L.; Pelletier, D.A.; Doktycz, M.J.; Hettich, R.L.; Abraham, P.E. Metaproteomics reveals insights into microbial structure, interactions, and dynamic regulation in defined communities as they respond to environmental disturbance. *BMC Microbiol* **2021**, *21*, 308, doi:10.1186/s12866-021-02370-4.

17. Kleiner, M.; Dong, X.; Hinzke, T.; Wippler, J.; Thorson, E.; Mayer, B.; Strous, M. Metaproteomics method to determine carbon sources and assimilation pathways of species in microbial communities. *Proc Natl Acad Sci U S A* **2018**, *115*, E5576–E5584, doi:10.1073/pnas.1722325115.

18. Taubert, M.; Vogt, C.; Wubet, T.; Kleinsteuber, S.; Tarkka, M.T.; Harms, H.; Buscot, F.; Richnow, H.H.; Von Bergen, M.; Seifert, J. Protein-SIP enables time-resolved analysis of the carbon flux in a sulfate-reducing, benzene-degrading microbial consortium. *ISME J* **2012**, *6*, 2291–2301, doi:10.1038/ismej.2012.68.

19. Grassl, N.; Kulak, N.A.; Pichler, G.; Geyer, P.E.; Jung, J.; Schubert, S.; Sinitcyn, P.; Cox, J.; Mann, M. Ultra-deep and quantitative saliva proteome reveals dynamics of the oral microbiome. *Genome Med* **2016**, *8*, 44, doi:10.1186/s13073-016-0293-0.

20. Kleiner, M.; Thorson, E.; Sharp, C.E.; Dong, X.; Liu, D.; Li, C.; Strous, M. Assessing species biomass contributions in microbial communities via metaproteomics. *Nat Commun* **2017**, *8*, 1588, doi:10.1038/s41467-017-01544-x.

21. Ouyang, W.Y.; Su, J.Q.; Richnow, H.H.; Adrian, L. Identification of dominant sulfamethoxazole-degraders in pig farm-impacted soil by DNA and protein stable isotope probing. *Environ Int* **2019**, *126*, 118–126, doi:10.1016/j.envint.2019.02.001.

22. Blakeley-Ruiz, J.A.; Kleiner, M. Considerations for constructing a protein sequence database for metaproteomics. *Comput Struct Biotechnol J* 2022, *20*, 937–952, doi:10.1016/j.csbj.2022.01.018.

23. Karlsson, R.; Gonzales-Siles, L.; Gomila, M.; Busquets, A.; Salvà-Serra, F.; Jaén-Luchoro, D.; Jakobsson, H.E.; Karlsson, A.; Boulund, F.; Kristiansson, E.; et al. Proteotyping bacteria: Characterization, differentiation and identification of pneumococcus and other species within the Mitis Group of the genus *Streptococcus* by tandem mass spectrometry proteomics. *PLoS One* **2018**, *13*, e0208804, doi:10.1371/journal.pone.0208804.

24. Trapp, J.; Geffard, O.; Imbert, G.; Gaillard, J.C.; Davin, A.H.; Chaumot, A.; Armengaud, J. Proteogenomics of *Gammarus fossarum* to document the reproductive system of amphipods. *Mol Cell Proteomics* **2014**, *13*, 3612–3625, doi:10.1074/mcp.M114.038851.

25. Heyer, R.; Benndorf, D.; Kohrs, F.; De Vrieze, J.; Boon, N.; Hoffmann, M.; Rapp, E.; Schlüter, A.; Sczyrba, A.; Reichl, U. Proteotyping of biogas plant microbiomes separates biogas plants according to process temperature and reactor type. *Biotechnol Biofuels* **2016**, *9*, 155, doi:10.1186/s13068-016-0572-4.

26. Pible, O.; Allain, F.; Jouffret, V.; Culotta, K.; Miotello, G.; Armengaud, J. Estimating relative biomasses of organisms in microbiota using "phylopeptidomics." *Microbiome* **2020**, *8*, 30, doi:10.1186/s40168-020-00797-x.

27. Gouveia, D.; Pible, O.; Culotta, K.; Jouffret, V.; Geffard, O.; Chaumot, A.; Degli-Esposti, D.; Armengaud, J. Combining proteogenomics and metaproteomics for deep taxonomic and functional characterization of microbiomes from a non-sequenced host. *npj Biofilms Microbiomes* **2020**, *6*, 23, doi:10.1038/s41522-020-0133-2.

28. Pible, O.; Petit, P.; Steinmetz, G.; Rivasseau, C.; Armengaud, J. Taxonomical composition and functional analysis of biofilms sampled from a nuclear storage pool. *Front Microbiol* **2023**, *14*, 1148976, doi:10.3389/fmicb.2023.1148976.

29. Lozano, C.; Kielbasa, M.; Gaillard, J.C.; Miotello, G.; Pible, O.; Armengaud, J. Identification and characterization of marine microorganisms by tandem mass spectrometry proteotyping. *Microorganisms* **2022**, *10*, 719, doi:10.3390/microorganisms10040719.

30. Grenga, L.; Pible, O.; Miotello, G.; Culotta, K.; Ruat, S.; Roncato, M.A.; Gas, F.; Bellanger, L.; Claret, P.G.; Dunyach-Remy, C.; et al. Taxonomical and functional changes in COVID-19 faecal microbiome could be related to SARS-CoV-2 faecal load. *Environ Microbiol* **2022**, *24*,

4299–4316, doi:10.1111/1462-2920.16028.

31. Muth, T.; Kolmeder, C.A.; Salojärvi, J.; Keskitalo, S.; Varjosalo, M.; Verdam, F.J.; Rensen,
S.S.; Reichl, U.; de Vos, W.M.; Rapp, E.; et al. Navigating through metaproteomics data: a
logbook of database searching. *Proteomics* **2015**, *15*, 3439–3453,
doi:10.1002/pmic.201400560.

32. Bassignani, A.; Plancade, S.; Berland, M.; Blein-Nicolas, M.; Guillot, A.; Chevret, D.; Moritz, C.;
Huet, S.; Rizkalla, S.; Clément, K.; et al. Benefits of iterative searches of large databases to
interpret large human gut metaproteomic data sets. *J Proteome Res* **2021**, *20*, 1522–1534,
doi:10.1021/acs.jproteome.0c00669.

33. Jouffret, V.; Miotello, G.; Culotta, K.; Ayrault, S.; Pible, O.; Armengaud, J. Increasing the power
of interpretation for soil metaproteomics data. *Microbiome* **2021**, *9*, 195, doi:10.1186/s40168-
021-01139-1.

34. Stamboulian, M.; Li, S.; Ye, Y. Using high-abundance proteins as guides for fast and effective
peptide/protein identification from human gut metaproteomic data. *Microbiome* **2021**, *9*, 80,
doi:10.1186/s40168-021-01035-8.

35. Starke, R.; Fiore-Donno, A.M.; White, R.A.; Parente Fernandes, M.L.; Martinović, T.; Bastida,
F.; Delgado-Baquerizo, M.; Jehmlich, N. Biomarker metaproteomics for relative taxa
abundances across soil organisms. *Soil Biol Biochem* **2022**, *175*, 108861,
doi:10.1016/J.SOILBIO.2022.108861.

36. Sayers, E.W.; Beck, J.; Brister, J.R.; Bolton, E.E.; Canese, K.; Comeau, D.C.; Funk, K.; Ketter,
A.; Kim, S.; Kimchi, A.; et al. Database resources of the National Center for Biotechnology
Information. *Nucleic Acids Res* **2020**, *48*, D9–D16, doi:10.1093/nar/gkz899.

37. Röst, H.L.; Schmitt, U.; Aebersold, R.; Malmström, L. pyOpenMS: a Python-based interface to
the OpenMS mass-spectrometry algorithm library. *Proteomics* **2014**, *14*, 74–77,
doi:10.1002/pmic.201300246.

38. Parte, A.C.; Carbasse, J.S.; Meier-Kolthoff, J.P.; Reimer, L.C.; Göker, M. List of prokaryotic
names with standing in nomenclature (LPSN) moves to the DSMZ. *Int J Syst Evol Microbiol*
2020, *70*, 5607–5612, doi:10.1099/IJSEM.0.004332/CITE/REFWORKS.

39. Serrano, G.; Guruceaga, E.; Segura, V. DeepMSPeptide: Peptide detectability prediction using
deep learning. *Bioinformatics* **2020**, *36*, 1279–1280, doi:10.1093/bioinformatics/btz708.

40. Kessner, D.; Chambers, M.; Burke, R.; Agus, D.; Mallick, P. ProteoWizard: open source
software for rapid proteomics tools development. *Bioinformatics* **2008**, *24*, 2534–2536,
doi:10.1093/BIOINFORMATICS/BTN323.

41. Jehmlich, N.; von Bergen, M. Protein-based stable isotope probing (Protein-SIP): applications
for studying aromatic hydrocarbon degradation in microbial communities. In *Anaerobic*
utilization of hydrocarbons, oils, and lipids; Springer International Publishing: Cham, 2020; pp.
277–284.

42. Afgan, E.; Baker, D.; Batut, B.; Van Den Beek, M.; Bouvier, D.; Ech, M.; Chilton, J.; Clements,
D.; Coraor, N.; Grüning, B.A.; et al. The Galaxy platform for accessible, reproducible and
collaborative biomedical analyses: 2018 update. *Nucleic Acids Res* **2018**, *46*, W537–W544,
doi:10.1093/NAR/GKY379.

43. Kim, S.; Pevzner, P.A. MS-GF[+] makes progress towards a universal database search tool for
proteomics. *Nat Commun* **2014**, *5*, 5277, doi:10.1038/ncomms6277.

44. Sachsenberg, T.; Herbst, F.A.; Taubert, M.; Kermer, R.; Jehmlich, N.; Von Bergen, M.; Seifert,
J.; Kohlbacher, O. MetaProSIP: automated inference of stable isotope incorporation rates in
proteins for functional metaproteomics. *J Proteome Res* **2015**, *14*, 619–627,
doi:10.1021/pr500245w.

45. García-Durán, C.; Martínez-López, R.; Zapico, I.; Pérez, E.; Romeu, E.; Arroyo, J.; Hernáez,
M.L.; Pitarch, A.; Monteoliva, L.; Gil, C. Distinct human gut microbial taxonomic signatures
uncovered with different sample processing and microbial cell disruption methods for
metaproteomic analysis. *Front Microbiol* **2021**, *12*, 1849,
doi:10.3389/FMICB.2021.618566/BIBTEX.

46. Perez-Riverol, Y.; Bai, J.; Bandla, C.; García-Seisdedos, D.; Hewapathirana, S.;

Kamatchinathan, S.; Kundu, D.J.; Prakash, A.; Frericks-Zipper, A.; Eisenacher, M.; et al. The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic Acids Res* **2022**, *50*, D543–D552, doi:10.1093/nar/gkab1038.

47. Qin, J.; Li, R.; Raes, J.; Arumugam, M.; Burgdorf, K.S.; Manichanh, C.; Nielsen, T.; Pons, N.; Levenez, F.; Yamada, T.; et al. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* **2010**, *464*, 59–65, doi:10.1038/nature08821.

48. Li, J.; Jia, H.; Cai, X.; Zhong, H.; Feng, Q.; Sunagawa, S.; Arumugam, M.; Kultima, J.R.; Prifti, E.; Nielsen, T.; et al. An integrated catalog of reference genes in the human gut microbiome. *Nat Biotechnol* **2014**, *32*, 834–841, doi:10.1038/NBT.2942.

49. Slotta, D.J.; McFarland, M.A.; Markey, S.P. MassSieve: panning MS/MS peptide data for proteins. *Proteomics* **2010**, *10*, 3035–3039, doi:10.1002/PMIC.200900370.

50. Klančnik, A.; Botteldoorn, N.; Herman, L.; Možina, S.S. Survival and stress induced expression of *groEL* and *rpoD* of *Campylobacter jejuni* from different growth phases. *Int J Food Microbiol* **2006**, *112*, 200–207, doi:10.1016/j.ijfoodmicro.2006.03.015.

51. Kupper, M.; Gupta, S.K.; Feldhaar, H.; Gross, R. Versatile roles of the chaperonin GroEL in microorganism-insect interactions. *FEMS Microbiol Lett* **2014**, *353*, 1–10, doi:10.1111/1574-6968.12390.

52. Gifford, S.M.; Sharma, S.; Booth, M.; Moran, M.A. Expression patterns reveal niche diversification in a marine microbial assemblage. *ISME J* **2013**, *7*, 281–298, doi:10.1038/ismej.2012.96.

53. Rinke, C.; Schwientek, P.; Sczyrba, A.; Ivanova, N.N.; Anderson, I.J.; Cheng, J.F.; Darling, A.; Malfatti, S.; Swan, B.K.; Gies, E.A.; et al. Insights into the phylogeny and coding potential of microbial dark matter. *Nature* **2013**, *499*, 431–437, doi:10.1038/nature12352.

54. Hill, J.E.; Town, J.R.; Hemmingsen, S.M. Improved template representation in *cpn60* polymerase chain reaction (PCR) product libraries generated from complex templates by application of a specific mixture of PCR primers. *Environ Microbiol* **2006**, *8*, 741–746, doi:10.1111/J.1462-2920.2005.00944.X.

55. Tang, S.; Chan, W.W.M.; Fletcher, K.E.; Seifert, J.; Liang, X.; Löffler, F.E.; Edwards, E.A.; Adrian, L. Functional characterization of reductive dehalogenases by using blue native polyacrylamide gel electrophoresis. *Appl Environ Microbiol* **2013**, *79*, 974–981, doi:10.1128/AEM.01873-12.

56. Hemmingsen, S.M.; Woolford, C.; van der Vies, S.M.; Tilly, K.; Dennis, D.T.; Georgopoulos, C.P.; Hendrix, R.W.; Ellis, R.J. Homologous plant and bacterial proteins chaperone oligomeric protein assembly. *Nature* **1988**, *333*, 330–334, doi:10.1038/333330a0.

57. Schaffert, C.S.; Klatt, C.G.; Ward, D.M.; Pauley, M.; Steinke, L. Identification and distribution of high-abundance proteins in the octopus spring microbial mat community. *Appl Environ Microbiol* **2012**, *78*, 8481–8484, doi:10.1128/AEM.01695-12.

58. Hendrickson, E.L.; Beck, D.A.C.; Wang, T.; Lidstrom, M.E.; Hackett, M.; Chistoserdova, L. Expressed genome of *Methylobacillus flagellatus* as defined through comprehensive proteomics and new insights into methylotrophy. *J Bacteriol* **2010**, *192*, 4859–4867, doi:10.1128/JB.00512-10.

59. Gallois, N.; Alpha-Bazin, B.; Ortet, P.; Barakat, M.; Piette, L.; Long, J.; Berthomieu, C.; Armengaud, J.; Chapon, V. Proteogenomic insights into uranium tolerance of a Chernobyl's *Microbacterium* bacterial isolate. *J Proteomics* **2018**, *177*, 148–157, doi:10.1016/j.jprot.2017.11.021.

60. Musatovova, O.; Dhandayuthapani, S.; Baseman, J.B. Transcriptional heat shock response in the smallest known self-replicating cell, *Mycoplasma genitalium*. *J Bacteriol* **2006**, *188*, 2845–2855, doi:10.1128/JB.188.8.2845-2855.2006.

61. Deppenmeier, U.; Johann, A.; Hartsch, T.; Merkl, R.; Schmitz, R.; Martínez-Arias, R.; Henne, A.; Wiezer, A.; Bäumer, S.; Jacobi, C.; et al. The genome of *Methanosarcina mazei*: evidence for lateral gene transfer between bacteria and archaea. *J Mol Microbiol Biotechnol* **2002**, *4*, 453–461.

62. Armengaud, J. Metaproteomics to understand how microbiota function: The crystal ball predicts

714   a promising future. *Environ Microbiol* **2023**, *25*, 115–125, doi:10.1111/1462-2920.16238.

(A)

10% *T. aromatica* K172 | 5% *T. aromatica* K172 | 1% *T. aromatica* K172

Minimal Top Rank Count

Absolute number of assignments of detected GroEL peptides

■ *Pseudomonas*   ■ *Thauera*   □ other genera

(B)

Minimal Top Rank Count

Relative number of assignments of detected GroEL peptides [%]

(A)

Mock A | Mock B | Mock C

Minimal Top Rank Count

Absolute number of assignments of detected GroEL peptides

(B)

to a family:
- present in "Mock A"
- absent in "Mock A"
- present in "Mock B"
- absent in "Mock B"
- present in "Mock C"
- absent in "Mock C"

Minimal Top Rank Count

Relative number of assignments of detected GroEL peptides [%]

Sample collection → Cell lysis, *e.g.,* freeze-thaw → GroEL separation & digestion → GroEL peptides → nLC-MS/MS analysis → Spectral raw data

Interpretation of spectral raw data

Spectral raw data

>AccNo_I
Sequence_I
>AccNo_II
Sequence_II
>AccNo_III
Sequence_III
...

GroEL .fasta database (72,759 nr sequences)

GroEL peptide identification, *e.g.*, Galaxy workflow including MS-GF+ & MetaProSIP → GroEL-proteotyping Python script → Bacterial community composition

100%
50%
0%

1. Match   2. +1   Identified peptides   Generate   I/L equivalent

AccNo | Taxonomy | GroELSeq | TrypPeps | HitPepSeq | HitPepCount — **Merge** → HitPepSeq | HitPepCount | AccNos | Taxonomy — **Filter** → HitPepSeq | HitPepCount | **TopRankCount** | AccNos | Taxonomy — **Merge** → Taxonomy | HitPepSeq | HitPepCount | PrecursorInt

GroEL .tsv database (284,351 sequences)

GroEL protein groups with same set of HitPepSeq

GroEL protein groups with same taxonomy

Example Identified peptides: AA, AB, AC, AD, AE, BA, BB, BC, BD, BE, CA, CB, CC, CD, CE, DA

HitPepSeqs	HitPep Count	TopRank Count	AccNos	Taxonomy
AA, AB, AC, AD, AE	5	**5**	I	*Staphylococcaceae*
BA, BB, BC, BD, BE	5	**5**	II	*Desulfovibrionaceae*
CA, CB, CC, CD, CE	5	**5**	III	*Bacillaceae*
AA, AB, BA, CA	4	**0**	IV	*Enterobacteriaceae*
AA, AB, AC, DA	4	**1**	V	*Pseudomonadaceae*