Time-, tissue- and treatment-associated heterogeneity in tumour-residing migratory DCs

Colin YC Lee[1,2], Bethany C Kennedy[3], Nathan Richoz[1], Zewen K Tuong[1,2], Isaac Dean[3], Fabrina
Gaspal[3], Zhi Li[3], Claire Willis[3], Tetsuo Hasegawa[1], Sarah K Whiteside[4], David A Posner[1],
Gianluca Carlesso[5], Scott A Hammond[5], Simon J Dovedi[6], Rahul Roychoudhuri[4], David R
Withers[3#], Menna R Clatworthy[1,2#]

[1] Molecular Immunity Unit, Department of Medicine, Medical Research Council Laboratory of
Molecular Biology, University of Cambridge, Cambridge, UK

[2] Cellular Genetics, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, UK

[3] Institute of Immunology and Immunotherapy, College of Medical and Dental Sciences, University of
Birmingham, Birmingham, UK

[4] Department of Pathology, University of Cambridge, Cambridge, UK

[5] Early Oncology R&D, AstraZeneca, Gaithersburg, USA

[6] Early Oncology R&D, AstraZeneca, Cambridge, UK

[#] **Correspondence:**
Prof. David R Withers, D.Withers@bham.ac.uk;
Prof. Menna R Clatworthy, mrc38@medschl.cam.ac.uk

Abstract

CCR7-expressing mature DCs enriched in immunoregulatory molecules (mRegDCs), such as PD-L1,
have been identified in tumours and draining lymph nodes (dLN), but their spatiotemporal dynamics
and role in anti-tumour immune responses remain unclear. Using photoconvertible mice to precisely
track DC migration, we found that mRegDCs comprised the dominant DC population arriving in the
dLN, but a subset remained tumour-resident despite CCR7 expression. These tumour-retained
mRegDCs were phenotypically and transcriptionally distinct from their dLN counterparts and were
heterogeneous, with increasing features of an "exhausted" transcriptional profile with more prolonged
tumour residence. This was characterised by reduced expression of antigen presentation and pro-
inflammatory transcripts, and these "exhaustion" features were attenuated by anti-PD-L1 treatment.
Tumour mRegDCs spatially co-localised with effector CD8$^+$ T cells in human solid tumours, and anti-
PD-L1 treatment enhanced their expression of several T cell-stimulatory molecules. Altogether, these
data provide fundamental insights into mRegDC biology and their role in cancer immunotherapy.

Introduction

Dendritic cells (DCs) capture tumour antigens and upregulate the chemokine receptor CCR7, which directs their migration to secondary lymphoid organs where they present antigens to T cells[1–3]. Two major conventional DC (cDC) subsets have been identified in tumours; cDC1 that specialise in cross-presenting tumour antigens to CD8[+] T cells and are associated with improved survival[4], and cDC2 that present exogenous antigen to CD4[+] T cells and have variable associations with cancer prognosis and treatment responses[1]. The recent application of single-cell technologies to tissue samples has enabled the distinction of an intratumoral DC population that may arise from both cDC1 and cDC2, but demonstrates a conserved phenotype and transcriptional programme characterised by the expression of LAMP3, genes consistent with a mature and migratory state (CCR7, CD40, IL12), and immunoregulatory molecules including PD-L1 and PD-L2[5]. This DC subset or state has been variously labelled as "migratory DCs", "mature DCs enriched in immunoregulatory molecules" (mRegDCs, the term we adopted in the current study), as well as "LAMP3[+] DCs", or cDC3[6]. Regardless of nomenclature, single-cell atlasing efforts have identified mRegDCs in multiple human cancers[7–9], and the acquisition of this transcriptional programme appears dependent on the phagocytic receptor AXL and uptake of tumour antigens[5].

Intriguingly, despite the conserved expression of co-inhibitory molecules such as PD-L1 in mRegDCs, the expression of the mRegDC-associated marker LAMP3 is associated with improved prognosis in breast, lung cancer and metastatic melanoma[10–13]. However, the precise contribution of mRegDCs to anti-tumour responses, and whether they have a positive or negative effect on disease outcomes remains unclear. Certainly, mRegDCs are likely to be key targets of immune checkpoint blockade (ICB) by virtue of their high expression of co-inhibitory molecules. Consistent with this, murine studies demonstrate the importance of DC-expressed PD-L1 in determining anti-tumour cytotoxic responses to subcutaneously tumours[14–16], and the application of anti-IL4 to increase mRegDCs enhanced responses to anti-PD-L1 in a non-small cell lung cancer (NSCLC) model[5].

mRegDCs express high levels of CCR7, enabling trafficking from the tumour to draining lymph nodes (dLN)[5,17]. Indeed, LAMP3[+] DCs have been identified in tumour-dLN, where they may activate tumour antigen-specific T cells[18]. However, LAMP3-expressing DCs have also been described within tertiary lymphoid structures in tumours[13,19], suggesting that some mature DCs may be retained in tumours for prolonged periods. Thus, the precise temporal dynamic behaviour of mRegDCs, the extent to which they act within the tumour versus dLN, and how these dynamics and site-specific cellular interactions are influenced by ICB remain to be clarified. The concept that prolonged residence within tumours might influence mRegDC fate and function is worthy of consideration given the known effects of the tumour microenvironment (TME) on other immune cell populations. For example, CD8[+] T cells transition to a so-called 'exhausted' state with prolonged dwell time in the tumour[20], with three defining characteristics; reduced effector function, sustained expression of inhibitory molecules such as PD-1, and a transcriptional state distinct from that of functional effector cells[21]. These features have also been observed in other tumour-resident immune cells such as natural killer (NK) cells[22,23].

Here, we used a photo-tracking mouse model, combined with single-cell RNA sequencing (scRNA-seq), confocal imaging and spatial transcriptomics, in mouse and human tissues, to interrogate the spatio-temporal dynamics of mRegDCs, their roles within the tumour, and the effects of ICB. We found mRegDCs showed transcriptional heterogeneity, influenced by duration of tumour residence, location (tumour versus dLN), and anti-PDL1 treatment. Strikingly, tumour-retained mRegDCs showed similar expression of CCR7 to those that migrated the dLN but took on an increasingly "exhausted" transcriptional profile with more prolonged tumour dwell time. These intratumoural mRegDCs co-localised with cytotoxic CD8[+] T cells, and anti-PD-L1 treatment enhanced their expression of several

77 important T cell-stimulatory molecules. mRegDC-CD8$^+$ T cell engagement and their augmentation by
78 ICB was confirmed across a range of human cancers. Altogether, these data provide further insight to
79 tumour mRegDC dynamics and their role in ICB therapy.
80

Results

mRegDC signatures are associated with improved survival in human cancers

The presence of mature LAMP[+] DCs within lymphoid follicles has been associated with improved
prognosis in NSCLC[10], but the effect of tumour mRegDCs on prognosis in human cancers more
broadly has not been examined. To address this, we analysed the transcriptomes of 4,045 human solid
tumours from the cancer genome atlas (TCGA)[24]. Enrichment of an mRegDC signature was associated
with improved survival not only in lung cancer, but also in cutaneous melanoma, breast, and colorectal
cancer (CRC, **Supplementary Fig. 1A**), all of which harbour CCR7[+] mRegDCs (**Supplementary Fig.
1B-G**)[25–27]. Further analysis of 1,853 human breast tumours from METABRIC[28] revealed cancer
subtype-specific prognostic associations (**Supplementary Fig. 1H**). These data suggest that mRegDCs
contribute to anti-tumour responses across a range of human cancers.

mRegDCs are transcriptionally heterogenous, and some subpopulations remain within the
tumour despite CCR7 expression

To investigate the mechanisms by which mRegDCs promote anti-tumour immunity, we sought to
characterise their spatio-temporal dynamics. The mRegDC programme is thought to be driven by
acquisition of tumour antigen[5]. We asked whether establishment of this programme inevitably leads to
the trafficking of CCR7[+] mRegDCs to the dLN, or whether some cells remain in the tumour. To address
this, we established multiple syngeneic subcutaneous colorectal tumours (MC38, CT26) in a
photoconvertible Kaede transgenic model that enables site-specific temporal labelling of cells within
the tumour[29]. Tumours were transcutaneously photoconverted on day 13, converting infiltrating cells
from the default green fluorescence (Kaede-green) to a red fluorescent profile (Kaede-red)[20]. 24-72h
following photo-conversion, tumours were harvested, enabling the distinction of newly infiltrating
Kaede-green cells from tumour-retained Kaede-red cells (**Fig. 1A**). To address the effect of ICB on
mRegDC dynamics, we administered anti-PD-L1 antibodies in this model (**Supplementary Fig. 2A**),
selecting this target because mRegDCs in CRC showed the highest expression of PD-L1 among
immune cells (**Supplementary Fig. 1C**).

scRNA-seq of FACS-isolated Kaede-green[+] or Kaede-red[+] myeloid cells (CD45[+]Ter119[-]
CD11b[+]NK1.1[-]), as well as tumour-infiltrating lymphocytes (TIL, CD45[+]Ter119[-]CD11b[-]) isolated 48h
after photoconversion (**Supplementary Fig. 2B**) generated 80,556 single cell transcriptomes following
rigorous quality control, including 32,191 myeloid cells (**Fig. 1B**, **Supplementary Fig. 2C**). Unbiased
clustering of DCs revealed 8 distinct clusters, including cycling DCs, cDC1, cDC2s, and mRegDCs,
assigned based on expression of canonical marker genes and similarity to published transcriptomes
(**Fig. 1C-D**, **Supplementary Fig. 2D**), with mRegDC showing high expression of *Ccr7*, *Cd274* and
Pdcd1lg2. Indeed, mRegDCs expressed higher levels of surface PD-L1 than other immune cells
(**Supplementary Fig. 2E-F**).

Over 50% of tumour DCs were mRegDC, and surprisingly, they were predominantly Kaede-red (**Fig.
1E**, **Supplementary Fig. 2G**). In contrast, cDC1 and cDC2 were mostly Kaede-green, consistent with
the conclusion that these newly infiltrating cDCs differentiate into mRegDCs following uptake of
tumour antigen[5]. The majority of mRegDCs expressed IRF8, consistent with a cDC1 origin, with a
smaller number expressing the cDC2-associated IRF4 (**Fig. 1D**). The prevalence of Kaede-red
mRegDCs was validated by flow cytometry in both MC38 and CT26 subcutaneous tumours (**Fig. 1F**,
Supplementary Fig. 2H-J), where mRegDCs were the major Kaede-red DC cell-type at 24h, 48h and
72h post-photoconversion (**Fig. 1G**). Finally, we confirmed the presence of Kaede-red MHC-II[+]CCR7[+]
cells by immunofluorescence (IF) microscopy (**Fig. 1H**, **Supplementary Fig. 3A-B**).

To explore the relationship between tumour cDC subsets, we performed a pseudotime analysis rooted
in the Kaede-green dominant cluster (cDC2_1). This revealed a trajectory terminating in mRegDC_2/3
that transitioned through the mRegDC_1 state (**Fig. 1I**, **Supplementary Fig. 3C**), consistent with the
Kaede-green/red ratio of clusters that marks tumour dwell in real time. Kaede-red$^+$ cDC1 and cDC2
showed higher expression of migratory transcripts and *Ciita*, a master regulator of antigen
presentation[30], than Kaede-green$^+$ counterparts, suggesting that cDCs mature with duration in the TME
(**Supplementary Fig. 3D**). Principal component (PC) analysis showed that Kaede status (green versus
red) accounted for approximately 70% of the transcriptional variance in cDC2s (PC1, **Supplementary**
Fig. 3E). Genes driving PC1 were enriched in pathways relating to antigen presentation and myeloid
differentiation (**Supplementary Fig. 3F**). cDC2_3, the cDC cluster with most Kaede-red cells,
upregulated *Ciita*, class-II MHC (MHC-II) transcripts, and *Ccr7* compared with cDC2_2
(**Supplementary Fig. 3H**). The receptor tyrosine kinase *Axl*, which induces the mRegDC programme[5],
was also upregulated as cDC2_1 transitioned to cDC2_3 (**Figure 1J**). Altogether, these data suggest
time-associated maturation of cDCs in the TME towards an mRegDC state and that antigen-charged
mRegDCs may reside in the tumour for several days, despite the expression of genes involved in
tumour egress, including CCR7.

Migrated mRegDCs are transcriptionally distinct from tumour-residing mRegDCs

DCs expressing mRegDC transcripts have been identified in tumour-dLN[5,31], but given our
demonstration that mRegDCs in tumours represent the dominant Kaede-red long-dwelling DC
population, we sought to definitively track tumour DC egress to dLNs. Kaede-red DCs originating
from the tumour were readily detectable in the dLN, but not in the contralateral non-draining LN
(ndLN), and were CCR7$^+$PD-L2$^+$MHC-II^{high} (**Fig. 2A-B**). Kaede-red mRegDCs comprised a stable
population across time, making up 30% of mRegDCs in the dLN at 24h, 48h and 72h post-
photoconversion (**Fig. 2C**). scRNA-seq of sorted Kaede-red CD45$^+$ cells from tumour-dLNs, and
CD45$^+$ cells from control LNs included a prominent *Ccr7$^+$Cd274$^+$* cluster assigned as mRegDCs (**Fig.**
2D, **Supplementary Fig. 4A-B**). In this data, 94% of Kaede-red$^+$ myeloid cells in the dLN were
mRegDCs (**Fig. 2E**), placing them as the dominant myeloid cell tumour emigrants arriving in the dLN.

Compared to Kaede-red tumour emigrants in the dLN, mRegDCs in tumour were enriched in
'*interferon gamma response*' genes (**Fig. 2F**). There was no difference in mRegDC '*lymphocyte co-*
stimulation' geneset expression between sites, but the precise co-stimulatory molecules differed (**Fig.**
2F, **Supplementary Fig. 4B-C**). Specifically, CD80 and CD86 were higher in tumour mRegDCs,
confirmed at a protein level, while *Icosl* (ICOS ligand) was significantly higher in dLN mRegDCs
(**Fig. 2G-H**). Of note, CD80/86 and ICOSL have differing effects on T cell activation, where signalling
through CD28 but not ICOS induces IL-2 production to support the clonal expansion of T cells[32]. *Il12b*,
which drives anti-tumour cytotoxic T cell activity[33], was enriched in tumour mRegDCs, while *Il15ra*,
a marker of DC maturation[34], was higher in mRegDC in the dLN. Re-analysis of scRNA-seq from
B16-F10 tumours and their dLN[35] confirmed tissue-associated heterogeneity in mRegDCs
(**Supplementary Fig. 4D-G**).

We next asked whether the three transcriptionally distinct tumour mRegDC subsets contributed equally
to the dLN emigrants. To address this, we integrated the single-cell transcriptomes of tumour-
originating DCs in the dLN with the tumour DC landscape (**Fig. 2I**). Strikingly, 80% of Kaede-red
DCs from the dLN mapped to the tumour mRegDC_1 cluster (**Fig. 2J**). Only 9% of DCs in the dLN
were mRegDC_2/3, despite mRegDC_2/3 being the dominant Kaede-red$^+$ tumour mRegDC subsets.
Although *Ccr7* expression remained high throughout mRegDC_1, 2 and 3, we observed a decrease in
transcripts associated with DC chemotaxis and CCR7 signalling in mRegDC_2 and 3 (**Supplementary**
Fig. 4H-I).

These data suggest that mRegDC_1, which have most recently acquired the mRegDC programme, are
the main population to seed the dLN. Hence, we propose that mRegDC_2 and mRegDC_3 are terminal,
tumour-residing mRegDC states, and DCs that have transited through the intermediate mRegDC_1
state, become increasingly unlikely to egress.

Tumour-retained mRegDCs progress towards an "exhausted" state

Since tumour-residing mRegDC subsets are reduced in their migratory capacity, we sought to assess
how their transcriptional programmes changed across pseudotime with longer tumour dwell-time.
During the transition from mRegDC_1 to mRegDC_2, there was upregulation of *Il12b*, *Cd80*, *Cd40*
(**Supplementary Fig. 5A)** potentially augmenting their capacity to promote anti-tumour responses.
However, with further pseudotime progression, we observed a decrease in MHC-II expression in
mRegDC_2/3, including class-II invariant chain *Cd74*, (**Fig. 3A**, **Supplementary Fig. 5B**). Similarly,
DC antigen presentation genes decreased as cells transitioned towards the tumour-retained
mRegDC_2/3 states (**Fig. 3B-C**). This included downregulation of molecules involved in antigen
processing (*Ctsb*, *Ctsl*, *Ifi30*, *Lgmn*, *Psme2*), transport (*Tap1*, *Tap2*), chaperones (*Hsp1a1*, *Hspa2*,
Hspa8), MHC loading (*H2-DMa*, *H2-Oa*, *Tapbp*, *Calr*) and *Ciita* (**Supplementary Fig. 5C-D**).

Moreover, genes involved in several important immunological processes, including '*interferon gamma*
response', '*inflammatory response*' and '*cytokine-cytokine receptor signalling*' pathway genes,
significantly decreased as cells transitioned from the mRegDC_1 to mRegDC_3 states (**Fig. 3D**).
mRegDC_3 showed reduced expression of several innate immune genes (*Nlrp3*, *Tlr1*, *Cd14*), genes
involved in DC activation and migration (*Axl*, *Ccr7*, *Rgs1*, *Icam1*), lymphocyte co-stimulation (*Cd40*,
Tnfsf9, *Pvr*) and cytokines which may recruit and activate immune cells (*Cxcl9*, *Il1b*, *Tnf*) compared
with mRegDC1 (**Fig. 3E**, **Supplementary Fig. 5E**). Hence, mRegDCs retained in the tumour undergo
a transition, acquiring the transcriptional hallmarks of "exhaustion", as defined in T cells, namely
reduced expression of molecules enabling effector function (in this case, antigen presentation),
sustained expression of inhibitory molecules, and a transcriptional state distinct from that of functional
effector cells[21].

mRegDCs interact with CD8 T cells in human tumours

Given that a substantial proportion of mRegDCs in tumour failed to migrate to dLN, and that tumour
mRegDCs may associate with outcomes, we sought to identify which immune cells mRegDCs may
communicate within the tumour. We deconvolved 521 human CRC transcriptomes[24] and found the
highest correlation between mRegDC and CD8$^+$ T cell transcripts (**Fig. 4A**), an observation replicated
in melanoma, breast, and lung tumour biopsies (**Fig. 4B**). Furthermore, the
CCR7$^+$CD274$^+$PDCD1LG2$^+$ DC cluster (consistent with mRegDCs in human CRC[27])
(**Supplementary Fig. 1B-C**), had stronger predicted interactions with effector CD8$^+$ T cells than other
myeloid cells, including via CD28, CTLA-4 and PD-1 (*PDCD1*) engagement (**Fig. 4C**). Similar
observations were confirmed in scRNA-seq of human breast tumours and melanoma (**Supplementary
Fig. 1D-G**)[25,26].

Effective engagement of cell-surface ligand-receptor pairs requires co-localisation of mRegDCs and
effector CD8$^+$ T cells. Analysis of spatial transcriptomics data from human CRC tumours[36] revealed
hotspots with co-localised expression of mRegDC and effector CD8$^+$ T cell genes (**Fig. 4D-E**,

Supplementary Fig. 5F). Spatial correlation of mRegDC and effector CD8[+] T cell transcripts was also
evident in breast tumours and melanoma (**Supplementary Fig. 5G-H**).

Finally, we analysed data from RNA-seq of physically-interacting cells (PICs), consisting of sorted
myeloid-T cell doublets from NSCLC[37]. Of note, mRegDC-T cell PICs were more frequent in tumours
than normal tissue[37]. We found that mRegDC-CD8[+] T cell doublets were more frequent than other
mRegDC-T cell combinations (**Supplementary Fig. 5I-J**), and PICs containing mRegDC
(*CCR7[+]FSCN1[+]CD274[+]*) highly co-expressed *CD8B, PRF1, GZMB and PDCD1* (**Fig. 4F**), confirming
that in NSCLC, mRegDC and effector CD8[+] T cells physically interact.

Altogether, these data suggest that mRegDC-CD8[+] T cell interaction is conserved across multiple
human solid tumours. These interactions may regulate T cell activation and could be important targets
in cancer immunotherapy.

Anti-PD-L1 reduces the abundance of tumour mRegDCs with an 'exhaustion-like' state and
promotes CD8[+] T cell interactions

Loss of PD-L1 expression from host immune cells, not tumour cells, is an important driver of
inflammatory gene expression in the TME (**Supplementary Fig. 6A**), so we investigated the effects
of anti-PD-L1 treatment on tumour DCs. In cDC2s, anti-PD-L1 treatment led to the upregulation genes
involved in inflammation and adaptive immunity and an increased expression of *Axl* expression as
cDC2_1 transitioned to cDC2_3 (**Supplementary Fig. 3G-I**). Across all mRegDC subsets, similar
pathways were enriched following anti-PD-L1, including "*interferon gamma response*", necessary for
effective response to ICB[33], and "*antigen processing and presentation*" genes were upregulated in
mRegDC_1 (**Supplementary Fig. 6B-C**).

Assessment of differential abundance[38] showed a significant enrichment in cellular neighbourhoods of
mRegDC_2 following anti-PD-L1 treatment, but a relative depletion of mRegDC_3 (**Fig. 5A**).
Strikingly, mRegDC_2 was highly transcriptionally activated compared to mRegDC_3, including
marked upregulation of many transcripts involved in immune signalling, such as *Cd40*, *Fos*, *Nfkbia*,
Tnf, etc. (**Fig. 5B**). These collectively drove enrichment in multiple gene signatures relating to immune
activation, such as TNF and cytokine signalling (**Supplementary Fig. 6D**). Importantly, we identified
several ligands that were differentially expressed on mRegDC_2 compared to other tumour mRegDC
subsets, and also relative to mRegDCs in the dLN, including *Il12b*, *Tnfsf4*, *Tnfsf9*, *Cd70* and *Pvr*
(**Supplementary Fig. 6E-F**). These molecules have established roles in supporting T cell
activation[33,39,40] and were also preferentially expressed on tumour-residing mRegDCs in B16-F10
tumours (**Supplementary Fig. 4G**). These data suggest that an "activated" tumour-residing mRegDC
state is enriched following anti-PD-L1 treatment.

To investigate the effects of anti-PD-L1 on mRegDC-CD8 T cell interactions in tumour, we analysed
scRNA-seq of TILs (**Supplementary Fig. 2B**, **7A-B**), identifying a *Prf1[+]Gzm[+]Pdcd1[+]Havcr2[+]* CD8[+]
T cell cluster resembling "exhausted" T (T$_{EX}$) cells[41,42] (**Fig. 5C**, **Supplementary Fig. 7C-D**) that were
predominantly tumour-resident and expanded with anti-PD-L1 (**Fig. 5D**, **Supplementary Fig. 7E**).
T$_{EX}$ cells were evident among the cycling_CD8T cluster and increased in proliferation following anti-
PD-L1, potentially underpinning their increased number (**Supplementary Fig. 7F-I**). Among CD8 T
cells, the T$_{EX}$ cells also showed the largest transcriptional response to anti-PD-L1 (**Fig. 5E**), including
upregulation of genes involved in "*T cell receptor signalling*" and "*TNFα signalling via NFκB*", with
potential beneficial anti-tumour effects (**Supplementary Fig. 7J**). Indeed, Kaede-red[+]PD-1[+] CD8[+] T
cells showed a significant increase in Granzyme B and IFNγ protein expression (**Supplementary Fig.**
7H,K-L).

Since mRegDCs are the major source of PD-L1 in tumours, and CD8$^+$ T$_{EX}$ cells the primary responders
to anti-PD-L1 therapy, we interrogated the molecular interactions between these two cell types that
might promote CD8$^+$ T$_{EX}$ activation and proliferation post-anti-PD-L1 using ligand-receptor expression
analysis[43] (**Fig. 5F**). This revealed several previously described interactions, including *CXCL16* and
IL15 from CCR7$^+$ DCs which may recruit and sustain cytotoxic T cells (**Supplementary Fig. 8A**)[44].
mRegDC_3 had fewer activating interactions with CD8$^+$ T$_{EX}$ cells than mRegDC_1/2, supporting their
status as an "exhausted" mRegDC state (**Supplementary Fig. 8B**). Importantly, we identified TNF-
superfamily interactions, including *TNFSF4–TNFRSF4* (OX40L–OX40), *TNFSF9–TNFRSF9* (4-
1BBL–4-1BB), *CD70–CD27*, and *PVR*-mediated interactions between mRegDC_2s and CD8$^+$ T$_{EX}$
cells, that were enriched following anti-PD-L1 treatment. These TNF-superfamily molecules promote
T cell survival, proliferation, and activation[39]. Moreover, PVR engages activating receptor CD226 or
its competing inhibitory receptor TIGIT to control CD8$^+$ T cell effector function[40]. Using IF
microscopy, we observed frequent interactions between tumour-residing mRegDCs and Kaede-
red$^+$CD3$^+$CD8$^+$ T cells within a perivascular niche, including proliferating (Ki-67$^+$)4-1BB$^+$ cytotoxic
cells (**Fig. 5G**, **Supplementary Fig. 8C-F**), consistent with a role for mRegDCs in regulating anti-
tumour cytolytic activity.

Conserved mRegDC heterogeneity and CD8$^+$ T cell crosstalk in human cancers

We next asked if the time, tissue, and treatment-associated heterogeneity in mRegDCs observed in our
murine model was pertinent to human cancers. In mRegDCs from human CRC (**Supplementary Fig.
1B**), there was a gradient of MHC-II expression (**Fig. 6A**). Importantly, markers upregulated on
tumour-residing MHC-II^{low} mRegDCs in murine tumours were also preferentially expressed in the
human MHC-II^{low} mRegDCs, including *IL15*, *PVR*, *TNFSF4*, *TNFSF9*, and *CD70*.

To assess tissue-associated differences in mRegDCs, we analysed an independent human CRC dataset
with paired scRNA-seq of tumour, dLN and normal adjacent tissue (**Supplementary Fig. 9A-C**)[45].
Consistent with our findings, there was no difference in *CCR7* expression, but *CD80*, *CD86*, *TNFSF4*,
TNFSF9, *CD70* and *PVR* were higher in tumour-residing mRegDCs versus dLN and normal tissue,
while *ICOSLG* was enriched in the dLN (**Fig. 6B**, **Supplementary Fig. 9D-E**). Therefore, the
heterogeneity of mRegDCs in murine tumours are paralleled in human CRC.

Next, we sought to address whether intra-tumour mRegDCs, and their precise phenotype, influences
clinical response to ICB. Atezolizumab (anti-PD-L1 antibody) is widely used in the treatment of
metastatic urothelial carcinoma (mUC), but its efficacy is variable[46]. We analysed 208 bulk
transcriptomes of tumour biopsies from the IMvigor210 trial for mUC[47,48]. In both responders and non-
responders, mRegDC gene expression was positively correlated with effector CD8$^+$ T cell transcripts,
and was higher in responders (**Supplementary Fig. 9F**). To leverage the clinical response data from
this cohort, we integrated the bulk transcriptomes with scRNA-seq of myeloid cells in mUC[49] (**Fig.
6C**) using *Scissor*[50], which enables identification of clinically relevant cell subpopulations and gene
expression profiles. This analysis revealed that mRegDCs were enriched in responders (**Fig. 6D**).
Indeed, myeloid cells associated with a favourable clinical response expressed transcripts associated
with tumour-residing mRegDCs, including *CCR7*, *CD274*, *IL15*, *PVR* and *CD70* (**Fig. 6E**).

Moreover, in breast cancer (**Supplementary Fig. 1D**, **9G**)[25], mRegDCs from tumours that successfully
underwent T cell clonotype expansion following treatment with anti-PD-1 antibodies were enriched in
transcripts associated with our ICB-induced mRegDC_2 cluster, versus non-responders (**Fig. 6F**).
mRegDCs from T cell clonotype expanders also upregulated genes contributing to '*interferon gamma
response*', '*FcγR-mediated phagocytosis*', '*lymphocyte co-stimulation*', etc. (**Supplementary Fig.

9H). Altogether, data from these treatment cohorts support the importance of an ICB-activated
mRegDC state.

Next, we asked if the molecular crosstalk between mRegDCs and CD8[+] T cells in mice is conserved
in humans. In PIC-seq data from NSCLC[37], we found that doublets containing mRegDCs highly
expressed ligands we had identified in mice (**Fig. 5F**), and their corresponding T cell receptors were
highly expressed in the same mRegDC-T cell conjugates (**Fig. 6G**). Moreover, mRegDC-CD8[+] T cell
doublets were most enriched for T cell cytotoxicity genes and '*TNFα response via NFκB*' compared to
other myeloid-CD8[+] T cell combinations (**Fig. 6H**, **Supplementary Fig. 9I-J**).

Finally, we re-examined hotspots of mRegDC and effector CD8[+] T cell co-localisation in spatial
transcriptomics of CRC, breast cancer and melanoma (**Fig. 4E**, **Supplementary Fig. 5G-H**).
Molecules involved in mRegDC-CD8[+] T cell crosstalk were expressed in these voxels, with spatial
correlation of mRegDC-CD8[+] T cell ligand-receptor pairs in all 3 tumour types (**Fig. 6I-J**,
Supplementary Fig. 10A-C). Altogether, these data show that mRegDCs are critically positioned to
modulate anti-tumour cytolytic activity, including via TNF-superfamily ligands and PVR.

Discussion

In this study, we combined scRNA-seq with a photoconvertible murine tumour model to unravel the
spatio-temporal dynamics of tumour mRegDCs. Contrary to current assumptions, mRegDCs are
heterogeneous, including sub-populations that either primarily contribute to LN migration or are
retained in the tumour. We found that mRegDCs acquire transcriptional features consistent with
functional "exhaustion" with increasing tumour dwell-time, but following anti-PD-L1 treatment,
become skewed towards a state enriched in T cell stimulatory molecules. This heterogeneity, crosstalk
with cytotoxic T cells, and association with response to ICB was conserved across human cancers.

Our work suggests that mRegDCs serve as a critical immune hub within the TME, through the
provision of chemotactic, survival, activating, and regulatory factors. We identified an mRegDC-CD8$^+$
T$_{EX}$ cell axis, including specific interactions which may control anti-tumour cytolytic activity.
Consistent with this, a recent study showed that acquisition of the full effector CD8$^+$ T cell program
requires engagement of CD80^{high} DCs in the local TME[51], which we show are likely mRegDCs. The
success of ICB have led to combination immune checkpoint therapy[52], exemplified by the recent
success of tiragolumab (anti-TIGIT) + atezolizumab for NSCLC[53]. In CD8$^+$ T cells, mechanistic
convergence of PD-1 and TIGIT inhibitory pathways powerfully regulates cytotoxic function[54]. We
found that mRegDCs are the major suppliers of both PD-1 and TIGIT ligands to CD8$^+$ T cells, and
importantly, there was an increase in PVR expression following anti-PD-L1 treatment, potentially
regulating T cell activation via CD226/TIGIT engagement[40]. Independently, combining OX40 agonists
with anti-PD-L1 antibodies effectively treated breast cancer in mice[55]. Our data suggest that mRegDCs
may contribute to the synergistic mechanisms of novel combinatorial immunotherapies.

Several questions remain; While provision of PD-1 ligands by DCs are essential for effective ICB[14,16],
it remains unclear whether anti-PD-L1 antibodies directly alter DC function. Interestingly, 'reverse
signalling' through PD-L1 has been described[56,57], including effects on DC migration[58]. Moreover,
knowledge of mRegDC ontogeny remain incomplete[6,17], including the relative contribution of cDC
subsets or monocytes to heterogeneous mRegDC states. Finally, whether the retention of mRegDCs in
tumours is due to the acquisition of aberrant trafficking behaviour or due to intratumoral interactions,
such as chemoattraction to stromal cells or tumour cells expressing CCR7 ligands [59,60] remains to be
resolved. However, recent studies have reported that mRegDCs engage CXCL13$^+$ CD4$^+$ T cells and
regulatory T cells in tumours[37,61,62], and may reside in tertiary lymphoid structures[10,13].

Previous studies have drawn conflicting conclusions on the role of mRegDCs in tumours; Loss of TIM-
3 in DCs prevents the acquisition of the mRegDC programme, but facilitates maintenance of the
effector CD8$^+$ T cell pool[63]. Conversely, CXCL16 and IL-15 expressing CCR7$^+$ DCs may recruit and
sustain CXCR6$^+$ cytotoxic T cells crucial in cancer immunosurveillance[44]. Our data help to resolve
these contradictions; We propose that tumour mRegDCs are heterogeneous, with subset-specific
capacity to both support and inhibit the cytotoxic T cell niche, presenting new opportunities for
intervention.

Methods

Mice

Transgenic C57BL/6 Kaede and BALB/c Kaede mice are maintained and bred at the University of Birmingham Biomedical Services Unit. C57BL/6 mice were maintained and bred at the University of Birmingham Biomedical Services Unit or the University of Cambridge Biomedical Services Gurdon Institute animal facilities. Mice were culled between the ages of 8 and 14 weeks. All animal experiments were conducted in accordance with Home Office guidelines and were approved by the University of Birmingham Animal Welfare and Ethical Review Body or the University of Cambridge Animal Welfare and Ethics Review Board. Mice were housed at 21°C, 55% humidity, with 12 h light-dark cycles in 7-7 individually ventilated caging with environmental enrichment of plastic houses plus paper bedding.

Mouse subcutaneous tumour model

MC38 (kindly provided by Dr. Gregory Sonnenberg; Weill Cornell Medicine, New York, NY), CT26 (kindly provided by Professor Tim Elliot, University of Oxford, Oxford, UK) and MC38-Ova (obtained from AstraZeneca) murine colon adenocarcinoma cells were cultured in RPMI in supplemented with 2mM L-glutamine (Thermo Fisher Scientific), 10% FBS (F9665; Sigma-Aldrich), and penicillin-streptomycin (Sigma-Aldrich) at 37°C with 5% CO2. Cells grown in the log-phase were harvested and resuspended to 2.5×10^6 cells/ml in Dulbecco's PBS (Sigma-Aldrich) for tumour injection.

2.5×10^5 tumour cells in 100 μl were subcutaneously injected into mice in the pre-shaved left flank area under anaesthesia via 2% gaseous isoflurane. MC38-Ova tumours were used for scRNA-seq and flow cytometry experiments. MC38 tumours were used for microscopy. CT26 tumours were used for flow cytometry and microscopy. Tumour size was periodically measured with a digital Vernier calliper, and the volume was calculated using the formula $V = 0.5 \times a \times b^2$ in cubic millimetres, where a and b are the long and short diameters of the tumour respectively. Tumour weights were measured at the endpoint of the experiment. Mice were sacrificed on day 14, 15, or 16 (24h, 48h, 72h post-photoconversion respectively, where performed) and tumours were harvested for analysis. Where indicated for specific experiments, tumour-dLNs (i.e. left inguinal LN) and contralateral non-draining lymph nodes were also harvested.

Administration of anti-PD-L1 antibodies

Anti-PD-L1 mouse IgG1 (Clone 80, SP16-260; AstraZeneca) or NIP228 isotype control mouse IgG1 (SP16-017; Astra Zeneca) were administered on day 7, 10, and 13 after tumour injection by intraperitoneal injection. Each dose consisted of 200 μg antibodies diluted in 200 μl PBS (10 mg/kg body weight). Tumour volume was measured on day 7, 10, 13 and the experiment endpoint.

Labelling of tumour compartment by photoconversion

Photoconversion was performed as previously described[20,29]. Briefly, on day 13 after tumour injection, the subcutaneous tumour was exposed to a 405-nm wavelength focussed LED light (Dymax BlueWave QX4 outfitted with 8mm focussing length, DYM41572; Intertronics) for 3 minutes, with a 5-second break every 20 seconds, at a fixed distance of 1 cm. Black cardboard was used to shield the remainder of the mouse. We previously showed that this method resulted in complete conversion (99.9%) of host cells within the tumour from the default green fluorescence of the Kaede protein (Kaede-green) to the altered red fluorescence (Kaede-red), and that the cells in the dLN were fully protected from tumour photoconversion[20]. Of note, while converted cells express Kaede-red fluorescence, they also retain a

393 weak Kaede-green signal. Overall, this enabled the discrimination of newly infiltrating (Kaede-green)
394 and resident (Kaede-red) cell populations in the tumour, or cells that have egressed the tumour (Kaede-
395 red) by their fluorescence profile. Moreover, using i.v. administration of anti-CD45 antibodies prior to
396 culling, we previously showed that majority of cells (>95%) were in tumour tissue and were not
397 intravascular contaminants[20].

Confocal microscopy

Tumour or LN samples were fixed in 1% paraformaldehyde (Electron Microscopy Services) for 24h
at 4°C followed by 12h in 30% sucrose in PBS. 20μm sections were permeabilized and blocked in
0.1M TRIS, containing 0.1% Triton (Sigma), 1% normal mouse serum, 1% normal rat serum and 1%
BSA (R&D Systems). Samples were stained for 2h at RT in a humid chamber with the appropriate
antibodies, listed in Supplementary table 1, washed 3 times in PBS and mounted in Fluoromount-G®
(Southern Biotech). Images were acquired using a TCS SP8 (Leica microsystems, Milton Keynes, UK)
confocal microscope. Raw imaging data were processed using Imaris v9.7.2 (Bitplane).

Iterative staining of sections was performed as previously described[64]. Samples were prepared and
stained as detailed above. Following acquisition, the coverslips were removed, and slides were washed
3 times in PBS to remove residual mounting medium. Bleaching of the fluorochromes were achieved
by submerging the slide in a 1mg/mL solution of lithium borohydride in water (Acros Organics) for 15
minutes at room temperature. The slides were then washed 3 times in PBS prior to staining with a
different set of antibodies. The process was repeated twice. Raw imaging data were processed using
Imaris using CD31 as fiducial for the alignment of acquired images.

Tissue dissociation

Tumours were cut into small pieces using surgical scissors, and incubated with 1 mg/ml collagenase D
(Roche) and 0.1 mg/ml DNase I (Roche) in a volume of 1.2 ml RPMI media at 37°C on a thermomixer
(Eppendorf) for 20 min; or tumours were digested using Tumour Dissociation Kit (Miltenyi Biotec)
and gentleMACS Dissociator (Miltenyi Biotec) for 40 minutes at 37°C according to the manufacturer's
protocol. The gentleMACS protocol was used for scRNA-seq experiments. Subsequently, the sample
was filtered through a 70 μm strainer to remove undigested tissue debris. Next, dead cells were
removed using Dead Cell Removal Kit and LS Columns (Miltenyi Biotec), according to the
manufacturer's instructions. Lymph nodes were cleaned and dissected in RPMI 1640 medium (Thermo
Fisher Scientific) and crushed through a 70 μm strainer. Thereafter, cells were centrifuged at 400 *g* at
4°C for 5 min and resuspended in FACS staining buffer (2% FBS; 2mM EDTA in PBS) for flow
cytometry.

Flow cytometry

Cell suspensions were subjected to Fc block with anti-CD16/32 (BioLegend) diluted in FACS staining
buffer on ice for 15 min before staining with a Live/Dead stain and surface markers, listed in
Supplementary table 1, diluted in FACS staining buffer on ice for 30 min. Where applicable, cells were
then fixed with Cytofix fixation buffer (BD) for 45 min and stained for intracellular markers diluted in
eBioscience permeabilization buffer (Thermo Fisher) at room temperature overnight. 1×10^4 counting
beads (Spherotech) were added to stained samples at the final step, to calculate absolute cell numbers.
Data were acquired on the LSR Fortessa X-20 (BD) using FACSDiva v8.0.2 software (BD) or
CytoFLEX (Beckman Coulter) using CytExpert v2.5 (Beckman Coulter) and analysed with FlowJo
v10.8.1 (BD).

Single-cell isolation

MC38-Ova tumours were injected in age-matched female C57BL/6 mice, treated with anti-PD-L1
antibodies, and photoconverted on day 13 as described above. Mice with tumours of similar sizes were
collected 48h after tumour photoconversion. After tumour digestion, as described above, cell
suspensions were stained for CD45 BV786, TER119 PE-Cy7, CD11b BV421, NK1.1 BV650,
Live/dead APC-Cy7 on ice for 30 min (Supplementary table 1). Subsequently, cells were centrifuged
at 400 g at 4°C for 5 min and resuspended in FACS staining buffer for sorting. Tumour-infiltrating
lymphocytes (Live CD45$^+$TER119$^-$Kaede$^+$CD11b$^{-/lo}$NK1.1$^{lo/hi}$) and tumour-infiltrating myeloid cells
(Live CD45$^+$TER119$^-$Kaede$^+$CD11b$^+$NK1.1$^-$) from anti-PD-L1 or isotype-control treated tumours
were sorted with a FACS Aria II Cell Sorter (BD) into two groups per condition, based on the presence
or absence of Kaede-red signal.

Tumour-dLNs from mice treated with the same experimental protocol were harvested and digested as
described in the preceding paragraph. Cell suspensions were stained for CD45 BV785, TER119 PE-
Cy7, Live/dead APC-Cy7 on ice for 30 min. Live CD45$^+$Ter119$^-$Kaede$^+$Kaede-red$^+$ cells were sorted,
to identify immune cells that have migrated from photoconverted tumours (Kaede-red) to the tumour-
dLNs. Inguinal lymph nodes from control mice (no tumour cells injected) were also processed for
scRNA-seq to obtain a representation of homeostatic cell populations in the LN. This was done to
facilitate accurate annotation and integration of Kaede-red cells in the tumour-dLN, where specific
populations arriving from the tumour would be disproportionately over-represented, with the LN
cellular landscape. Control LN cell suspensions were stained for B220 BV421, CD11c BV786,
TER119 PE-Cy7, Live/dead APC-Cy7 on ice for 30 min. Thereafter, a 3-way sort was performed
consisting of Live Kaede$^+$TER119$^-$CD11c$^+$B220$^-$ (myeloid), Kaede$^+$TER119$^-$CD11c$^-$B220$^+$ (B cells),
and Kaede$^+$TER119$^-$CD11c$^-$B220$^-$ fractions. The three subsets were then mixed at a ratio of 1:1:1 to
generate a cellular suspension enriched for DCs and with reduced frequency of B cells. scRNA-seq of
control LNs and tumour-dLNs were sequenced and analysed together.

Single-cell library construction and sequencing

Gene expression libraries from were prepared from FACS-sorted populations of single cells using the
Chromium Controller and Chromium Single Cell 3' GEM Reagent Kits v3 (10x genomics, Inc.)
according to the manufacturer's protocol. The resulting sequencing libraries comprised of standard
Illumina paired-end constructs flanked with P5 and P7 sequences. The 16 bp 10x barcode and 10 bp
UMI were encoded in read 1, while read 2 was used to sequence the cDNA fragment. Sample index
sequences were incorporated as the i7 index read. Paired-end sequencing (2 x 150 bp) was performed
on the Illumina NovaSeq 6000 platform. The resulting .bcl sequence data were processed for QC
purposes using bcl2fastq software (v2.20.0.422) and the resulting fastq files were assessed using
FastQC (v0.11.3), FastqScreen (v0.9.2) and FastqStrand (v0.0.5) prior to alignment and processing
with the CellRanger (v6.1.2) pipeline.

Processing of scRNA-seq

Single-cell gene expression data from CellRanger count output (filtered features, barcodes, and
matrices) were analysed using the Scanpy[65] (v1.8.2) workflow. Raw count data from the myeloid and
TIL sorts were concatenated. Doublet detection was performed using Scrublet[66] (v0.2.1), with cells
from iterative sub-clustering flagged with outlier Scrublet scores labelled as potential doublets. Cells
with counts mapped to >6000 or <1000 genes were filtered. The percentage mitochondrial content cut-
off was set at <7.5%. Genes detected in fewer than 3 cells were filtered. Total gene counts for each cell
were normalised to a target sum of 10^4 and log1p transformed. There resulted in a working dataset of

80,556 cells. Next, highly variable features were selected based on a minimum and maximum mean
expression of ≥0.0125 and ≤3 respectively, with a minimum dispersion of 0.5. Total feature counts,
mitochondrial percentage, and cell cycle scores, where indicated, were regressed out. The number of
principal components used for neighbourhood graph construction was set to 50 initially, and
subsequently 30 for myeloid and TIL subgroup processing. Clustering was performed using the Leiden
algorithm with resolution set at 1.5 for initial annotations, but subsequently sub-clustering was
performed at lower resolutions (0.7-1.0) for analysis of myeloid cell, DC, T cell, and CD8[+] T cell
subsets. Uniform manifold approximation and projection (UMAP, v0.5.1) was used for dimensional
reduction and visualisation, with a minimum distance of 0.3, and all other parameters according to the
default settings in Scanpy.

Analysis of scRNA-seq from mouse tumour models

Broad cell types of interest were subset (eg. DCs from myeloid cells) and re-clustered as described
above. Resulting clusters were annotated using canonical marker gene expression and published
transcriptomic signatures. Unless otherwise indicated, log-transformed expression values were used
for plotting. Gene set scoring was performed using Scanpy's tl.score_genes tool. Gene sets were
obtained from the Molecular Signature Database (MSigDB) inventory, specifically Hallmark, KEGG,
or Gene Ontology (GO), using the R package msigdbr (v7.5.1) or published RNAseq data
(Supplementary table 2). The original published mRegDC signature gene list[5] was used for the initial
scoring and identification of mRegDCs. Differential gene testing was performed using the Wilcoxon
rank sum test implemented in Scanpy's tl.rank_genes_groups. Analysis of MC38-Ova and B16
tumours and their dLNs followed these same methods.

Trajectory analysis was performed using partition-based graph abstraction (PAGA)[67] and the Palantir
algorithm[68]. For Palantir pseudo-time analysis, the differentiation trajectory was rooted in the Kaede-
green dominant cluster, which represents the cellular subset most associated with newly-infiltrating
cells. Integration and label transfer of lymph node DC scRNA-seq data and tumour DCs, or cycling
CD8[+] T cells and non-cycling CD8[+] T cells was performed using Scanpy's tl.ingest tool. For analysis
of cycling CD8[+] T cells, cell cycle regression was first performed on the isolated cluster using Scanpy's
pp.regress_out function, before re-integration.

Gaussian kernel density estimation to compute density of cells in the UMAP embedding was performed
using Scanpy's tl.embedding_density. Differential abundance analysis was performed using Milo.
Spcifically, the k-nearest-neighbour (kNN) graph was constructed with a k parameter of 30 and initial
random sampling rate of 0.1. Cellular neighbourhoods with constituent cells comprising less than 70%
of a previously defined cluster were designated as mixed neighbourhoods. CellPhoneDB[43] (v2.1.7) was
used for cell-cell communication analysis, using the default parameters and normalised expression
values as the input. CellPhoneDB output was visualised using ktplots
(https://github.com/zktuong/ktplots).

For pathway analysis between Kaede-red versus Kaede-green or anti-PD-L1 versus isotype control
groups in the scRNA-seq dataset, a pseudo-bulk approach was first applied to single-cell gene
expression data (https://github.com/colin-leeyc/CLpseudobulk), to increase robustness for pathway
analysis and overcome limitations associated with differential expression testing on single cells[69].
Briefly, filtered, raw count data (prior to normalisation, transformation, or scaling) from each condition
were randomly sorted into artificial replicates, with iterative bootstrapping applied to random sampling
(n >10). Pseudo-bulked count matrices were normalised using the median-of-ratios method,
implemented in DESeq2[70] (v1.34), and differential gene expression testing was performed using the
Wald test. Pre-ranked gene set enrichment analysis[71] (GSEA) was implemented in fgsea (v1.24), using

the averaged Wald statistic over $n >10$ iterations of random sorting into pseudo-bulked replicates, as
the gene rank metric. Leading edge genes were identified for further analysis. GO term over-
representation analysis was implemented in topGO[72] (v2.50), using the top 100 genes with the highest
loadings for each principal component (PC) of interest, following PC analysis.

Analysis of scRNA-seq from human tumours

scRNA-seq data from human solid tumours were downloaded from public repositories, which are listed
under *Data availability* with references. Where necessary, data access permissions were sought and
approved prior to download. Data was analysed using the Scanpy (v1.8.2) workflow, as outlined in the
sections above. Filtering for quality control was performed according to the parameters outlined in the
original publications. scRNA-seq integration was performed using the batch-balanced KNN (with
ridge regression) approach or Harmony algorithm with sequencing batch as the batch term, where
available and sufficient, or patient ID if sequencing batch was not available. Cell annotations from
original publications were checked and refined, particularly where myeloid cell annotations were not
complete or the focus of the original publication.

Myeloid cells were subset and used for further analysis. In total, this included re-analysis of 41,624
and 43,193 myeloid cells from two independent CRC datasets, 16,688 myeloid cells from breast
tumours, 8,555 myeloid cells from cutaneous melanoma, 11,663 myeloid or T cells from NSCLC
including 901 myeloid-T cell doublets, and 1,988 myeloid cells from metastatic urothelial carcinoma.
mRegDCs were identified using expression of migratory transcripts and mRegDC signature genes[5].

The only exception to the Scanpy workflow was for analysis of NSCLC PICseq data, where the
Metacell workflow was utilised, as documented in the original publication, for analysis and annotation
of single-cells[37]. This was to ensure compatibility with the PICseq deconvolution algorithm[73], which
was used to analyse sorted doublets. Gene expression raw count data from PICs only were obtained
and input to the Scanpy workflow for visualisation and further analysis. Doublet deconvolution results
for contributing myeloid cell or T cell identities in each PIC were retained.

Subsequently, analysis of gene expression profiles, gene signature scoring, cell-cell communication,
were as described in earlier sections. For analysis of enrichment of mRegDC_2 genes in the breast
cancer dataset, where anti-PD-1 response data was available, mRegDC_2 genes were first identified
as DEGs between mRegDC_2 versus remaining mRegDCs (Wilcoxon rank-sum test) in scRNA-seq
from murine tumours. 85 differentially upregulated genes (p-adj < 0.05, $\log_2$Fold-change > 1) were
converted to equivalent human gene symbols using Ensembl and used to score mRegDCs from human
breast cancer (Fig. 6F).

Analysis of bulk transcriptomics

Bulk RNA-seq data were downloaded from public repositories, listed under *Data availability* with
references. Where necessary, data access permissions were sought and approved prior to download.
TCGA data was accessed via TCGAbiolinks, using STAR-aligned reads[24]. For bulk transcriptomics
data from both TCGA and the IMvigor 210 trial, transcripts-per-million (tpm) normalised values were
used for analysis. Cellular deconvolution was performed using xCell[74], implemented in the webtool
using default parameters. Gene signatures scoring was performed using single-sample GSEA[75]
(ssGSEA, v10.1.0), implemented in GenePattern, followed by normalisation (scaled between 0 and 1)
of gene set enrichment scores. Cell type signatures for ssGSEA were derived from cell-specific
differentially expressed genes in scRNA-seq of human cancers. Pearson correlation was used to assess
for correlation of cellular proportions or cell-specific signatures of interests in biopsy samples. Survival

analysis was performed on overall survival (months), using the median mRegDC signature score for
stratification to 'high' and 'low' groups, and log-rank test was applied to survival curves. For
integration of bulk transcriptomic data from the IMvigor210 trial with accompanying clinical response
data with scRNA-seq of myeloid cells in mUC, the Scissor pipeline was used[50]. Scissor enables
identification of single-cells and gene expression profiles within cell subpopulations that are
significantly associated with phenotypes obtained from bulk expression data. A logistic regression
model using the binary outcome of clinical responders (CR/PR) vs non-responders (SD/PD) was used,
and the alpha parameter was set at 0.05, per default settings.

Raw RNA-seq sequencing data from PD-L1 knockout (KO) mice injected with wild-type (WT) MC38
tumour cells, WT mice injected with PD-L1 KO tumours, PD-L1 KO mice injected with PD-L1 KO
tumours, and WT mice injected with WT tumours was aligned and quantified using Salmon and
tximport[15]. Analysis was performed using the standard DESeq2 workflow and GSEA was used for
pathway analysis.

Analysis of spatial transcriptomics

10x Visium spatial transcriptomics were analysed using the standard Scanpy (v1.8.2) workflow, using
the default SpaceRanger outputs including spot alignments to corresponding tissue haematoxylin and
eosin or immune-fluorescence images. Spots with fewer <500 counts, >80000 counts, and percentage
mitochondrial content >15% were filtered. Genes detected in fewer than 5 spots were filtered. Total
feature counts, mitochondrial percentage, and cell cycle scores were regressed out. Expression values
plotted are log1p-transformed 10^4-sum-normalised values. Gene set scoring was performed using
Scanpy's tl.score_genes tool. Calculation of spatial correlation was implemented using the
correlationSpot function in ktplots (https://github.com/zktuong/ktplots), as previously described[76].
Briefly, k = 6 nearest neighbourhoods were extracted from a kNN graph computed from the spatial
location of each Visium spot. Pearson correlation was performed on each neighbourhood using gene
expression values or gene signature scores and averaged across the neighbourhoods. Correlation values
were only returned if expression value or signature scores were detected in all spots. Spots with positive
correlation scores were plotted, which represent areas of co-localisation.

Statistical analysis

Mice were gender-matched (all female). Tumour growth curves for anti-PD-L1 and isotype control
treated groups were analysed using two-way ANOVA and Sidak's multiple comparisons test and are
presented as mean ± SEM. Analysis for RNA sequencing is as described above, including statistical
frameworks used. Statistical tests were implemented for two principal purposes; to compare expression
values (RNA or protein) between samples, or to compare proportions of defined cell subsets. All
experiments were performed with biological replicates, and the specific statistical tests applied are
indicated in the figure legends. All statistical tests applied were two-tailed. Paired statistical tests were
used where different populations from the same animal were compared. All animal experiments were
randomised prior to experimental intervention. Statistical significance are denoted as follows: ns, not
significant; $*P < 0.05$; $**P < 0.01$; $***P < 0.001$; $****P < 0.0001$. Statistical analyses were performed
in R or GraphPad Prism.

Online supplemental material

607 Supplementary table 1 list antibodies used. Supplementary table 2 list gene sets used.

Data and code availability

The scRNA-seq data have been deposited on the GEO public repository under accension numbers
GSExxxxxxx and GSExxxxxx (*available upon manuscript publication*). Additional data or
information will be made available upon reasonable request. Published data was accessed and
downloaded from public GEO, SRA, EGA and Synapse repositories using the following ascension
numbers: scRNA-seq of CRC (GSE178341 and syn26844071)[27,45]; scRNA-seq of breast cancer
(EGAS00001004809)[25]; scRNA-seq of melanoma (GSE123139)[26]; scRNA-seq of mUC
(HRA000212)[49]; scRNA-seq and PICseq of NSCLC (GSE160903)[37]; TCGA
(https://portal.gdc.cancer.gov, via TCGAbiolinks)[24]; IMvigor210 bulk RNA-seq
(EGAS00001004343)[47]; 10x Visium (https://www.10xgenomics.com/resources/datasets)[36]; PD-L1
knockout bulk RNA-seq (PRJNA356678)[15]. All analysis performed are described in detail under
Methods. Source code for key softwares or computational pipelines used have been cited. Additional
code used for analysis, as described above, have been made available on GitHub repositories.

References

1. Wculek, S. K. *et al.* Dendritic cells in cancer immunology and immunotherapy. *Nature Reviews Immunology 2019 20:1* **20**, 7–24 (2019).

2. Roberts, E. W. *et al.* Critical Role for CD103(+)/CD141(+) Dendritic Cells Bearing CCR7 for Tumor Antigen Trafficking and Priming of T Cell Immunity in Melanoma. *Cancer Cell* **30**, 324–336 (2016).

3. Sallusto, F. *et al.* Rapid and coordinated switch in chemokine receptor expression during dendritic cell maturation. *Eur J Immunol* **28**, 2760–2769 (1998).

4. Böttcher, J. P. & Reis e Sousa, C. The Role of Type 1 Conventional Dendritic Cells in Cancer Immunity. *Trends Cancer* **4**, 784–792 (2018).

5. Maier, B. *et al.* A conserved dendritic-cell regulatory program limits antitumour immunity. *Nature* **580**, 257–262 (2020).

6. Kvedaraite, E. & Ginhoux, F. Human dendritic cells in cancer. *Sci Immunol* **7**, (2022).

7. Gerhard, G. M., Bill, R., Messemaker, M., Klein, A. M. & Pittet, M. J. Tumor-infiltrating dendritic cell states are conserved across solid human cancers. *Journal of Experimental Medicine* **218**, (2021).

8. Cheng, S. *et al.* A pan-cancer single-cell transcriptional atlas of tumor infiltrating myeloid cells. *Cell* **184**, 792-809.e23 (2021).

9. Zilionis, R. *et al.* Single-Cell Transcriptomics of Human and Mouse Lung Cancers Reveals Conserved Myeloid Populations across Individuals and Species. *Immunity* **50**, 1317-1334.e10 (2019).

10. Goc, J. *et al.* Dendritic cells in tumor-associated tertiary lymphoid structures signal a th1 cytotoxic immune contexture and license the positive prognostic value of infiltrating CD8+ t cells. *Cancer Res* **74**, 705–715 (2014).

11. Movassagh, M. *et al.* Selective Accumulation of Mature DC-Lamp+ Dendritic Cells in Tumor Sites Is Associated with Efficient T-Cell-Mediated Antitumor Response and Control of Metastatic Dissemination in Melanoma. *Cancer Res* **64**, 2192–2198 (2004).

12. de Mingo Pulido, Á. *et al.* TIM-3 Regulates CD103+ Dendritic Cell Function and Response to Chemotherapy in Breast Cancer. *Cancer Cell* **33**, 60-74.e6 (2018).

13. Dieu-Nosjean, M. C. *et al.* Long-term survival for patients with non-small-cell lung cancer with intratumoral lymphoid structures. *J Clin Oncol* **26**, 4410–4417 (2008).

14. Peng, Q. *et al.* PD-L1 on dendritic cells attenuates T cell activation and regulates response to immune checkpoint blockade. *Nature Communications 2020 11:1* **11**, 1–8 (2020).

15. Lau, J. *et al.* Tumour and host cell PD-L1 is required to mediate suppression of anti-tumour immunity in mice. *Nature Communications 2017 8:1* **8**, 1–11 (2017).

16. Oh, S. A. *et al.* PD-L1 expression by dendritic cells is a key regulator of T-cell immunity in cancer. *Nature Cancer 2020 1:7* **1**, 681–691 (2020).

17. Ginhoux, F., Guilliams, M. & Merad, M. Expanding dendritic cell nomenclature in the single-cell era. *Nature Reviews Immunology 2022 22:2* **22**, 67–68 (2022).

18. Zhang, Q. *et al.* Landscape and Dynamics of Single Immune Cells in Hepatocellular Carcinoma. *Cell* **179**, 829-845.e20 (2019).

19. Germain, C. *et al.* Presence of B cells in tertiary lymphoid structures is associated with a protective immunity in patients with lung cancer. *Am J Respir Crit Care Med* **189**, 832–844 (2014).

20. Li, Z. *et al.* In vivo labeling reveals continuous trafficking of TCF-1+ T cells between tumor and lymphoid tissue. *J Exp Med* **219**, (2022).

21. Wherry, E. J. T cell exhaustion. *Nat Immunol* **12**, 492–499 (2011).

22. Gao, Y. *et al.* Tumor immunoevasion by the conversion of effector NK cells into type 1 innate lymphoid cells. *Nat Immunol* **18**, 1004–1015 (2017).

23. Platonova, S. *et al.* Profound coordinated alterations of intratumoral NK cell phenotype and function in lung carcinoma. *Cancer Res* **71**, 5412–5422 (2011).

669   24.   The Cancer Genoma Atlas. TCGA. *National Cancer Institute (NCI) and National Human Genome*
670   *Research Institute (NHGRI)* (2013).

671   25.   Bassez, A. *et al.* A single-cell map of intratumoral changes during anti-PD1 treatment of patients with
672   breast cancer. *Nat Med* **27**, 820–832 (2021).

673   26.   Li, H. *et al.* Dysfunctional CD8 T Cells Form a Proliferative, Dynamically Regulated Compartment
674   within Human Melanoma. *Cell* **176**, 775-789.e18 (2019).

675   27.   Pelka, K. *et al.* Spatially organized multicellular immune hubs in human colorectal cancer. *Cell* **184**,
676   4734-4752.e20 (2021).

677   28.   Curtis, C. *et al.* The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel
678   subgroups. *Nature* **486**, 346–352 (2012).

679   29.   Tomura, M. *et al.* Monitoring cellular movement in vivo with photoconvertible fluorescence protein
680   'Kaede' transgenic mice. *Proc Natl Acad Sci U S A* **105**, 10871–10876 (2008).

681   30.   Riley, J. L., Westerheide, S. D., Price, J. A., Brown, J. A. & Boss, J. M. Activation of class II MHC
682   genes requires both the X □ region and the class II transactivator (CIITA). *Immunity* **2**, 533–543 (1995).

683   31.   Miller, J. C. *et al.* Deciphering the transcriptional network of the DC lineage. *Nat Immunol* **13**, 888
684   (2012).

685   32.   Riley, J. L. *et al.* Modulation of TCR-induced transcriptional profiles by ligation of CD28, ICOS, and
686   CTLA-4 receptors. *Proc Natl Acad Sci U S A* **99**, 11790–11795 (2002).

687   33.   Garris, C. S. *et al.* Successful Anti-PD-1 Cancer Immunotherapy Requires T Cell-Dendritic Cell
688   Crosstalk Involving the Cytokines IFN-γ and IL-12. *Immunity* **49**, 1148-1161.e7 (2018).

689   34.   Dubois, S. P., Waldmann, T. A. & Müller, J. R. Survival adjustment of mature dendritic cells by IL-15.
690   *Proc Natl Acad Sci U S A* **102**, 8662 (2005).

691   35.   Davidson, S. *et al.* Single-Cell RNA Sequencing Reveals a Dynamic Stromal Niche That Supports
692   Tumor Growth. *Cell Rep* **31**, (2020).

693   36.   10x Genomics. 10x Genomics Spatial Gene Expression. *Visium demonstration dataset*
694   https://www.10xgenomics.com/resources/datasets (2022).

695   37.   Cohen, M. *et al.* The interaction of CD4+ helper T cells with dendritic cells shapes the tumor
696   microenvironment and immune checkpoint blockade response. *Nat Cancer* **3**, 303–317 (2022).

697   38.   Dann, E., Henderson, N. C., Teichmann, S. A., Morgan, M. D. & Marioni, J. C. Differential abundance
698   testing on single-cell data using k-nearest neighbor graphs. *Nat Biotechnol* **40**, 245–253 (2022).

699   39.   Croft, M. The role of TNF superfamily members in T-cell function and diseases. *Nature Reviews*
700   *Immunology 2009 9:4* **9**, 271–285 (2009).

701   40.   Johnston, R. J. *et al.* The Immunoreceptor TIGIT Regulates Antitumor and Antiviral CD8+ T Cell
702   Effector Function. *Cancer Cell* **26**, 923–937 (2014).

703   41.   Im, S. J. *et al.* Defining CD8+ T cells that provide the proliferative burst after PD-1 therapy. *Nature* **537**,
704   417–421 (2016).

705   42.   Singer, M. *et al.* A Distinct Gene Module for Dysfunction Uncoupled from Activation in Tumor-
706   Infiltrating T Cells. *Cell* **166**, 1500-1511.e9 (2016).

707   43.   Efremova, M., Vento-Tormo, M., Teichmann, S. A. & Vento-Tormo, R. CellPhoneDB: inferring cell–
708   cell communication from combined expression of multi-subunit ligand–receptor complexes. *Nature*
709   *Protocols 2020 15:4* **15**, 1484–1506 (2020).

710   44.   di Pilato, M. *et al.* CXCR6 positions cytotoxic T cells to receive critical survival signals in the tumor
711   microenvironment. *Cell* **184**, 4512-4530.e22 (2021).

712   45.   Joanito, I. *et al.* Single-cell and bulk transcriptome sequencing identifies two epithelial tumor cell states
713   and refines the consensus molecular classification of colorectal cancer. *Nature Genetics 2022 54:7* **54**,
714   963–975 (2022).

46.   Rosenberg, J. E. *et al.* Atezolizumab in patients with locally advanced and metastatic urothelial
carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm,
multicentre, phase 2 trial. *Lancet* **387**, 1909–1920 (2016).

47.   Banchereau, R. *et al.* Molecular determinants of response to PD-L1 blockade across tumor types. *Nature*
Communications 2021 12:1 **12**, 1–11 (2021).

48.   Balar, A. v. *et al.* Atezolizumab as first-line treatment in cisplatin-ineligible patients with locally
advanced and metastatic urothelial carcinoma: a single-arm, multicentre, phase 2 trial. *The Lancet* **389**,
67–76 (2017).

49.   Chen, Z. *et al.* Single-cell RNA sequencing highlights the role of inflammatory cancer-associated
fibroblasts in bladder urothelial carcinoma. *Nature Communications 2020 11:1* **11**, 1–12 (2020).

50.   Sun, D. *et al.* Identifying phenotype-associated subpopulations by integrating bulk and single-cell
sequencing data. *Nature Biotechnology 2021 40:4* **40**, 527–538 (2021).

51.   Prokhnevska, N. *et al.* CD8+ T cell activation in cancer comprises an initial activation phase in lymph
nodes followed by effector differentiation within the tumor. *Immunity* **56**, 107-124.e5 (2022).

52.   Yi, M. *et al.* Combination strategies with PD-1/PD-L1 blockade: current advances and future directions.
Molecular Cancer 2022 21:1 **21**, 1–27 (2022).

53.   Cho, B. C. *et al.* Tiragolumab plus atezolizumab versus placebo plus atezolizumab as a first-line
treatment for PD-L1-selected non-small-cell lung cancer (CITYSCAPE): primary and follow-up
analyses of a randomised, double-blind, phase 2 study. *Lancet Oncol* **23**, 781–792 (2022).

54.   Banta, K. L. *et al.* Mechanistic convergence of the TIGIT and PD-1 inhibitory pathways necessitates co-
blockade to optimize anti-tumor CD8+ T cell responses. *Immunity* **55**, 512-526.e9 (2022).

55.   Lee, J. v. *et al.* Combinatorial immunotherapies overcome MYC-driven immune evasion in triple
negative breast cancer. *Nature Communications 2022 13:1* **13**, 1–12 (2022).

56.   Gato-Cañas, M. *et al.* PDL1 Signals through Conserved Sequence Motifs to Overcome Interferon-
Mediated Cytotoxicity. *Cell Rep* **20**, 1818–1829 (2017).

57.   Clark, C. A. *et al.* Tumor-Intrinsic PD-L1 Signals Regulate Cell Growth, Pathogenesis, and Autophagy
in Ovarian Cancer and Melanoma. *Cancer Res* **76**, 6964–6974 (2016).

58.   Lucas, E. D. *et al.* PD-L1 Reverse Signaling in Dermal Dendritic Cells Promotes Dendritic Cell
Migration Required for Skin Immunity. *Cell Rep* **33**, (2020).

59.   Whyte, C. E. *et al.* ACKR4 restrains antitumor immunity by regulating CCL21. *Journal of Experimental*
Medicine **217**, (2020).

60.   Cheng, H. W. *et al.* CCL19-producing fibroblastic stromal cells restrain lung carcinoma growth by
promoting local antitumor T-cell responses. *Journal of Allergy and Clinical Immunology* **142**, 1257-
1271.e4 (2018).

61.   Thumkeo, D. *et al.* PGE2-EP2/EP4 signaling elicits immunosuppression by driving the mregDC-Treg
axis in inflammatory tumor microenvironment. *Cell Rep* **39**, (2022).

62.   Magen, A. *et al.* Intratumoral mregDC and CXCL13 T helper niches enable local differentiation of CD8
T cells following PD-1 blockade. *bioRxiv* 2022.06.22.497216 (2022) doi:10.1101/2022.06.22.497216.

63.   Dixon, K. O. *et al.* TIM-3 restrains anti-tumour immunity by regulating inflammasome activation. *Nature*
595, 101 (2021).

64.   Radtke, A. J. *et al.* IBEX: A versatile multiplex optical imaging approach for deep phenotyping and
spatial analysis of cells in complex tissues. *Proc Natl Acad Sci U S A* **117**, 33455–33465 (2020).

65.   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: Large-scale single-cell gene expression data analysis.
Genome Biol **19**, 1–5 (2018).

66.   Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of Cell Doublets in
Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291.e9 (2019).

67.   Wolf, F. A. *et al.* PAGA: graph abstraction reconciles clustering with trajectory inference through a
topology preserving map of single cells. *Genome Biol* **20**, 1–9 (2019).

68.   Setty, M. *et al.* Characterization of cell fate probabilities in single-cell data with Palantir. *Nat Biotechnol*
764   **37**, 451–460 (2019).

765   69.   Squair, J. W. *et al.* Confronting false discoveries in single-cell differential expression. *Nature*
766   *Communications 2021 12:1* **12**, 1–15 (2021).

767   70.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq
768   data with DESeq2. *Genome Biol* **15**, (2014).

769   71.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for interpreting
770   genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545–15550 (2005).

771   72.   Alexa, A. & Rahnenfuhrer, J. topGO: Enrichment Analysis for Gene Ontology. Preprint at (2022).

772   73.   Giladi, A. *et al.* Dissecting cellular crosstalk by sequencing physically interacting cells. *Nature*
773   *Biotechnology 2020 38:5* **38**, 629–637 (2020).

774   74.   Aran, D., Hu, Z. & Butte, A. J. xCell: Digitally portraying the tissue cellular heterogeneity landscape.
775   *Genome Biol* **18**, 220 (2017).

776   75.   Barbie, D. A. *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven cancers require
777   TBK1. *Nature* **462**, 108–112 (2009).

778   76.   Tuong, Z. K. *et al.* Resolving the immune landscape of human prostate at a single-cell level in health and
779   cancer. *Cell Rep* **37**, (2021).

Author Contributions

CYCL processed and analysed sequencing data, designed and performed experiments, analysed data, and wrote the manuscript. BCK, NR, ZL and ID performed experiments and analysed data. ZKT and TH provided project guidance. CW, SW and DP performed experiments. GC, SAH and SJD provided input and reviewed the manuscript. RR supported the project. DW conceived the project, designed experiments, provided guidance, and edited the manuscript. MRC conceived the project, designed experiments, provided guidance, and wrote the manuscript.

Funding

CYCL is funded by the Gates Cambridge scholarship trust and University of Cambridge School of Clinical Medicine Elmore fund. BCK was supported by Cancer Research UK (SEBSTF-2021/100002). ID is supported by an iCASE Studentship with AstraZeneca. DRW is funded by Cancer Research UK Immunology Project Award C54019/A27535, Cancer Research Institute CLIP grant CRI3128, Worldwide Cancer Research Grant 21-0073 and a Medical Research Council IMPACT iCASE Studentship with AstraZeneca. MRC is supported by the National Institute of Health Research (NIHR) Cambridge Biomedical Research Centre and the NIHR Blood and Transplant Research Unit and a Wellcome Investigator Award (220268/Z/20/Z).

Acknowledgments

We thank A. Ptasinska and other members of Genomics Birmingham at University of Birmingham for all their help with single-cell RNA-sequencing experiments, and the University of Birmingham Flow Cytometry Facility. We thank Dr. Y. Miwa (Tsukuba University, Tsukuba, Japan) and Dr. O. Kanagawa (RIKEN Research Center for Allergy and Immunology, Yokohama, Japan) and Dr. M. Tomura (Osaka Ohtani University, Tondabayashi, Japan) for the Kaede mice.

Disclosures

AstraZeneca provided therapeutic anti-PD-L1 antibodies, isotype control antibodies, and MC38-Ova cells. GC, SAH and SJD are full employees and share-holders in AstraZeneca. No other disclosures or conflicts of interest are reported.

Figure 1

Figure 1 | Landscape and temporal dynamics of DCs in murine subcutaneous tumours.

(A) Experiment design. Tumours from C57BL/6 Kaede transgenic mice were harvested 24h, 48h, or 72h after photoconversion.

(B) UMAP of 32,191 myeloid cells from scRNA-seq of FACS-sorted CD45$^+$CD11b$^+$ Kaede-green$^+$/Kaede-red$^+$ cells 48h after photoconversion of subcutaneous MC38-Ova tumours.

(C) UMAP of 10,336 DCs from (B), and canonical marker gene expression of respective clusters (D).

(E) Density embedding of DCs from each condition (Kaede-green/red, anti-PD-L1/isotype control antibodies).

(F) Representative flow cytometry plots of Kaede-green/red composition in DCs subsets from MC38-Ova tumours 48h post-photoconversion. Data is representative of 3 independent experiments (12 independent mice).

(G) Time-course of DC subsets from MC38-Ova tumours *x* hours after tumour photoconversion. Medians are shown; individual points represent independent mice. Paired student's t-test was used.

(H) Representative confocal microscopy images of CT26 tumours 72h after photoconversion. Scale bar, 40μm; arrows, tumour-residing mRegDCs (Kaede-red$^+$CCR7$^+$MHC-II$^+$, dendritic morphology).

(I) Tumour DCs ordered by pseudotime, rooted in the cluster with highest proportion of Kaede-green DCs.

(J) Expression of *Axl* in tumour DCs.

Figure 2

Figure 2 | mRegDCs that migrate to the dLN alter their co-stimulatory phenotype and are transcriptionally distinct from tumour-residing populations.

(A) Representative flow cytometry plots of tumour-dLNs and contralateral ndLNs 48h after tumour photoconversion. DCs that migrate to the LN from photoconverted tumours carry the Kaede-red fluorescent profile, which enables tracking to the dLN.

(B) Time-course of mRegDCs in the dLN versus ndLN x hours after tumour photoconversion; and (C) percentage of mRegDCs that are Kaede-red in dLNs. Medians are shown; individual points represent independent mice; paired student's t-test was used.

(D-E) UMAP of 7795 myeloid cells from scRNA-seq of FACS-sorted CD45$^+$ Kaede-red cells from tumour-dLNs and CD45$^+$ cells from control LNs.

(F) GSEA of Kaede-red mRegDCs in the dLN versus mRegDCs in the tumour (isotype control-treated). Signed P-adj indicate $\log_{10}$(adjusted P-value) with the direction of the enrichment score.

(G) Expression of selected genes between Kaede-red mRegDCs in the tumour-dLN and mRegDCs in the tumour (isotype control-treated). Wilcoxon rank-sum test was used.

(H) Flow cytometry of surface expression of CD80 and CD86 on mRegDCs 48h after tumour photoconversion. Representative histograms and MFI; means ± standard deviation shown; individual points represent independent mice; Mann-Whitney U test was used.

(I) Integration of Kaede-red DCs from the tumour-dLN with the tumour DC landscape and label transfer.

(J) Proportions of DCs by origin and Kaede profile.

Figure 3

Figure 3 | Tumour-residing mRegDCs undertake an "exhausted" state with duration in the tumour.
(A) Expression of MHC-II transcripts and *Cd74* over pseudotime in tumour DCs. Local polynomial regression (loess) was fit to range-standardised expression values.
(B) Gene signature scores for "*GO:BP dendritic cell antigen processing and presentation (GO:0002468)*" across mRegDC clusters. Wilcoxon rank-sum test was used.
(C) Gene signature scores of individual cells for "*GO:BP dendritic cell antigen processing and presentation (GO:0002468)*" and mRegDC signature genes, coloured by pseudotime. Range-standardised scores shown.
(D) GSEA of mRegDC_3 versus mRegDC_1. Only significantly enriched pathways (*P*-adj < 0.05) shown.
(E) Volcano plot of DEGs between mRegDC_1 and mRegDC_3. Significant DEGs from "*Hallmark inflammatory response*" are highlighted in red (*P*-adj < 0.01).

Figure 4

Figure 4 | mRegDCs interact with CD8⁺ T cells in human cancers.
(A) Pearson correlation between cell proportions, from deconvolution of 521 bulk transcriptomes of colorectal adenocarcinoma biopsies (TCGA), and mRegDC signature genes. Box highlights high correlation between mRegDC and CD8⁺ T cells. Only significant correlations ($p < 0.01$) shown.
(B) Pearson correlation between mRegDC signature genes and effector CD8⁺ T cell signature genes in colorectal cancer (COAD), breast cancer (BRCA), cutaneous melanoma (SKCM) and lung adenocarcinoma (LUAD) from TCGA, where enrichment of mRegDC genes shows survival benefit (SFig. 1A).
(C) CellPhoneDB[43] cell-cell communication analysis between myeloid cells and effector CD8⁺ T cells in scRNA-seq of human CRC. Width of edges represent standardised interaction scores. Only significant interactions ($p < 0.01$) shown.
(D) Gene signature scores for mRegDCs and effector CD8⁺ T cells in 10x Visium spatial transcriptomics of a human CRC tumour section.
(E) Spatial correlation of mRegDC and effector CD8⁺ T cell signature scores in spatial transcriptomics of independent human CRC tumour sections ($n = 2$).
(F) Expression of selected genes in myeloid-T cell PICs from NSCLC, grouped by the myeloid cell identity.

Figure 5

Figure 5 | anti-PD-L1 promotes activating mRegDC-cytotoxic CD8⁺ T cell interactions in the TME.

(A) Milo[38] differential abundance analysis of tumour DCs. Bee-swarm plot shows treatment-associated differential abundance in overlapping cellular neighbourhoods (individual points). Differentially abundant neighbourhoods at FDR < 0.05 are coloured. 'Mixed' refers to neighbourhoods where cells do not predominantly (>70%) belong to a single cluster.

(B) Volcano plot between mRegDC_2 and mRegDC_3. DEGs (P-adj < 0.01, $\log_2$Fold-change > 0.5) are coloured. Selected genes involved in immune and inflammatory signalling are labelled.

(C) UMAP of cells from scRNA-seq of 16,351 CD8⁺ T cells, from FACS-sorted CD45⁺CD11b⁻ Kaede-green/red TILs 48h after photoconversion of subcutaneous MC38-Ova tumours.

(D) Proportions of CD8⁺ T$_{EX}$ cells by Kaede profile and treatment.

(E) Number of significant DEGs in anti-PD-L1 versus isotype control-treated tumours. DEGs calculated using Wilcoxon rank-sum test.

(F) CellPhoneDB[43] ligand-receptor interaction analysis between tumour DCs and CD8⁺ T cells. Only significant interactions ($p < 0.05$) shown.

(G) Representative confocal microscopy images of MC38 tumours, showing co-localisation of CCR7⁺MHC-II⁺ mRegDCs and CD3⁺CD8⁺4-1BB⁺Ki-67⁺ T cells. Scale bar, 20 μm.

Figure 6

Human colorectal cancer

A — HLA-DRA, CD74, IL15, IL15RA, PVR, TNFSF4, TNFSF9, CD70, CXCL16 UMAP plots. n = 1572

B — Violin plots of CD80, CD86, TNFSF9, TNFSF4, PVR, CD70 expression across N, T, dLN tissues.

IMvigor 210 trial: anti-PD-L1 for metastatic urothelial carcinoma

C — Human mUC myeloid cells: Mac, cDC1, cDC2, mRegDC, pDC. n = 1988

D — Myeloid GEX profiles associated with response: Background cells, SDPD, CRPR

E — Non-responders (SD/PD) or Background cells ← → Responders (CR/PR). Volcano plot. p-adj < 0.05

Human breast cancer

F — mRegDC_2 genes; NE vs E; T cell clonal expansion; ***

Human NSCLC (*PICseq*)

G — mRegDC ligands / T cell receptors

H — GO:BP T cell cytotoxicity across Mono, Mac, MoDC, cDC1, cDC2, mRegDC; Myeloid-CD8T cell doublet

Human colorectal cancer (*10X Genomics Visium*)

I — mRegDC ~ CXCR6, mRegDC ~ TNFRSF4, mRegDC ~ TNFRSF9, mRegDC ~ CD27, mRegDC ~ IL12RB1, mRegDC ~ TIGIT

J — mRegDC ~ CXCR6, mRegDC ~ TNFRSF4, mRegDC ~ TNFRSF9, mRegDC ~ CD27, mRegDC ~ IL12RB1, mRegDC ~ TIGIT

Figure 6 | Conserved mRegDC heterogeneity and CD8$^+$ T cell crosstalk in human cancer.

(A) UMAP of 1572 mRegDCs from human CRC (n = 62 patients), from Fig. 1A.

(B) Expression of selected co-stimulatory molecules in mRegDCs between tumour (T), normal adjacent tissue (N) and tumour-dLN, (independent CRC scRNA-seq dataset from (A)[45]). Wilcoxon rank-sum test was used.

(C) UMAP of 1988 myeloid cells from scRNA-seq of human metastatic urothelial carcinoma (mUC, n = 11 patients).

(D) Cells associated with response (complete responder, CR; partial responder, PR) or non-responders (stable disease, SD; progressive disease, PD) in the IMvigor 210 trial (atezolizumab) for treatment of mUC. Scissor[50] was used to integrate bulk transcriptomes of patients (n = 208) from the IMvigor210 trial and scRNA-seq of myeloid cells from mUC (C).

(E) Volcano plot of gene expression in myeloid cells associated with responders (CR/PR) versus non-responders (SD/PD) from (D).

(F) Gene signature scores for mRegDC_2 transcripts in mRegDCs from human breast cancer treated with anti-PD-1 antibodies. Comparison between T cell clonotype expanders (responders, n = 9 patients) versus non-expanders (non-responders, n = 20 patients). Wilcoxon rank-sum test was used.

(G) PICseq (myeloid-T cell doublets) of human NSCLC: expression of mRegDC-T cell ligand-receptor pairs, grouped by myeloid cell identity in each myeloid-T cell doublet.

(H) Gene signature scores for "*GO:BP T cell mediated cytotoxicity (GO:0001913)*" (filtered for HLA genes), in myeloid-T cell PICs grouped by the myeloid identity. Wilcoxon rank-sum test was used.

(I-J) Spatial correlation of mRegDC signature scores and selected CD8$^+$ T$_{EX}$ cell receptors, in 10x Visium spatial transcriptomics of 2 independent human CRC tumour sections.

Supplementary Figure 1

Supplementary Figure 1 | mRegDCs in human cancer.
(A) Kaplan-Meier analysis of overall survival rate in skin cutaneous melanoma (SKCM, $n =$ 473), colorectal adenocarcinoma (COAD, $n = 521$), breast invasive carcinoma (BRCA, $n =$ 1226), and lung adenocarcinoma (LUAD, $n = 598$) from TCGA, stratified by enrichment of mRegDC signature genes.
(B) Uniform manifold approximation projection (UMAP) of 41,624 myeloid cells from scRNA-seq of human CRC ($n = 62$ patients) and expression of mRegDC genes (mRegDC signature score[5], *CCR7*, *LAMP3*, *FSCN1*).
(C) Expression of *CD274* (PD-L1), *PDCD1LG2* (PD-L2) and *PDCD1* (PD-1), *CCR7*, *CD80* and CD86 by cell-type in scRNA-seq of human CRC.
(D) UMAP of 16,688 myeloid cells from scRNA-seq of human breast cancer ($n = 29$ patients) and expression of mRegDC genes.
(E) Expression of selected genes by cell-type in scRNA-seq of human breast cancer.
(F) UMAP of 8,555 myeloid cells from scRNA-seq of human melanoma ($n = 25$ patients) and expression of mRegDC genes.
(G) Expression of selected genes by cell-type in scRNA-seq of human melanoma.
(H) Kaplan-Meier analysis of overall survival rate in hormone receptor-positive (oestrogen receptor, ER, or progesterone receptor, PR, $n = 1369$), human epidermal growth factor receptor 2 (HER2)-positive ($n = 185$), or triple-negative breast cancer (TNBC, $n = 299$) from METABRIC. *L*og-rank test was used.

Supplementary Figure 2

Supplementary Figure 2 | Single cell profiling of mouse subcutaneous tumours.
(A) Tumour growth curves comparing anti-PD-L1 antibody-treated versus isotype control antibody treatment, administered on day 7, 10 and 13, over 15 days. Data shown is related to the scRNA-seq experiment (from Fig. 1) and includes 5 mice per condition.
(B) Fluorescence-activated cell sorting (FACS) strategy for isolation of tumour myeloid cells and TILs, for scRNA-seq.
(C) Canonical marker gene expression in myeloid cell subsets, related to Fig. 2B.
(D) Gene signature scores of published DC transcriptomic signatures in DC clusters from Fig. 2C. Monocyte-derived DCs (moDC) signatures also scored highly in cDC2 clusters, consistent with known challenges in transcriptional distinction of cDC2 and moDC[7].
(E) *Cd274* (PD-L1) expression in myeloid cells.
(F) PD-L1 surface expression of immune cells in MC38-Ova tumours.
(G) Proportion of cells from Kaede-green/red fractions by DC cluster.
(H) Representative flow cytometry gating strategy for identification of tumour DC populations. mRegDCs were identified as Live, CD45$^+$ Lin$^-$ (NK1.1, B220, Ly6G, CD3, F4/80) Kaede$^+$ PD-L2$^+$ CCR7^{lo-hi} cells. PD-L2 expression was more restricted to mRegDCs than PD-L1 and used as the principal marker to identify mRegDCs.
(I) Representative flow cytometry plot; Live, CD45$^+$Lin$^-$F/80$^-$CD3$^-$PD-L2$^+$ cells are MHC-II$^+$ and CD11c^{lo-hi}, supporting the use of PD-L2$^+$ signal to identify mRegDCs (H). CD3$^+$ T cells shown in blue for comparison.
(J) Representative flow cytometry plots of Kaede-green/red composition in mRegDCs from CT26 tumours 48h post-photoconversion. Data is representative of tumours from 6 independent mice.

Supplementary Figure 3

Supplementary Figure 3 | Imaging of tumour mRegDCs and scRNA-seq analysis of pre-mRegDC populations.

(A) Representative confocal microscopy images of CT26 tumours 72h after tumour photoconversion, stained for CCR7 and class-II MHC (MHC-II). Scale bar, 500μm; insets highlight selected regions with CCR7$^+$MHC-II$^+$ DCs.

(B) Representative confocal microscopy images of MC38 tumours 48h after tumour photoconversion. Scale bar, 30μm; arrows, tumour-residing mRegDCs.

(C) Partition-based graph abstraction (PAGA) of scRNA-seq of tumour DC clusters, showing that mRegDC_1 are an intermediate state between cDCs and terminal mRegDC_2/3.

(D) Expression of selected migratory genes and *Ciita* in scRNA-seq data of cDC1 and cDC2, by Kaede profile. Wilcoxon rank-sum test was used.

(E) Principal component (PC) analysis of scRNA-seq of cDC2s, pseudo-bulked to 3 artificial replicates per condition. PC1; variance due to Kaede profile. PC2; variance due to treatment.

(F) Gene ontology (GO) term enrichment test for the top 100 loading genes of PC1 (F) and PC2 (G). O/E, observed/expected ratio; colour, direction of gene loading. Fisher-exact test was used.

(H) Volcano plot of DEGs between cDC2_2 and cDC2_3. Selected DEGs relating to DC maturation, antigen presentation, MHC-II and migration are highlighted in red (all significant, *P*-adj < 0.01).

(I) Expression of *Axl* in cDC_2 and cDC2_3.

Supplementary Figure 4

Supplementary Figure 4 | Comparison of mRegDCs in dLNs versus tumour.
(A) Expression of mRegDC or myeloid cell canonical marker genes in LN myeloid cells (Fig. 3C).
(B) Expression of selected genes in DCs from tumours (isotype control-treated) and their tumour-dLNs. DCs from the dLN were most similar to mRegDCs in tumours.
(C) GSEA of "*GO:biological process (BP) lymphocyte co-stimulation (GO:0031294)*", comparing Kaede-red mRegDCs from the dLN versus tumour mRegDCs.
(D) UMAP of 1303 myeloid cells from scRNA-seq of murine subcutaneous B16 tumours and tumour-dLNs, by myeloid cell-type and tissue.
(E) Expression of mRegDC genes.
(F) Expression of selected genes in scRNAseq data of mRegDCs from B16 tumours and tumour-dLNs. Wilcoxon rank-sum test was used.
(G) Expression of additional selected genes in mRegDCs from (D).
(H) Expression of *Ccr7* in DCs from MC38-Ova tumours.
(I) Gene signature enrichment of pathways relating to DC movement and CCR7 signalling.

Supplementary Figure 5

Supplementary Figure 5 | Transcriptional changes in mRegDCs with tumour dwell time and mRegDC CD8⁺ T cell interactions in human solid tumours.

(A) Dot plot of *Cd80, Cd40, Il12b* expression in tumour DCs.

(B) Dot plot of class-I and class-II MHC gene expression (log-transformed, unscaled) in mRegDCs.

(C) Volcano plot of DEGs between mRegDC_1 and mRegDC_3. Significant DEGs from "*KEGG antigen processing and presentation*" are highlighted in red (P-adj < 0.01).

(D) Expression of *Ciita* in tumour DCs.

(E) Volcano plot of DEGs between mRegDC_1 and mRegDC_3. Significant DEGs from "*GO: molecular function (MF) cytokine activity (GO:0005125)*" are highlighted in red (P-adj < 0.01).

(F) Gene signature scores for mRegDCs and effector CD8⁺ T cells in 10x Visium spatial transcriptomics of 1 human CRC tumour section associated with Fig. 4E.

(G) Spatial correlation of mRegDC and effector CD8⁺ T cell signature scores in spatial transcriptomics of independent human breast tumour sections ($n = 2$) and a human melanoma section (H, $n = 1$).

(I) Sequencing of physically interacting cells (PICseq, myeloid-T cell doublets) from human NSCLC ($n = 10$ patients). UMAP of PICs coloured by the deconvolved myeloid and T cell assignment for each doublet (PICseq algorithm). Each dot represents one myeloid-T cell doublet.

(J) Heatmap of frequency of myeloid-T cell doublet combinations in PICseq of NSCLC.

Supplementary Figure 6

Supplementary Figure 6 | mRegDC activation following anti-PD-L1 treatment.
(A) GSEA of pathways differentially enriched in bulk RNA-seq of PD-L1 knockout (KO) mice injected with wild-type (WT) MC38 tumour cells, WT mice injected with PD-L1 KO tumours, or PD-L1 KO mice injected with PD-L1 KO tumours; versus WT mice injected with WT tumours. Only significant pathways (FDR-q-val < 0.05) are shown; *n=5* biological replicates per condition.
(B) GSEA of mRegDCs by treatment arm in mRegDCs, using Hallmark (HM) and KEGG pathways. Only significant pathways (*P*-adj < 0.05) are shown.
(C) Dot plot of selected leading-edge genes from GSEA analysis of "*Hallmark interferon gamma response*", "*Hallmark inflammatory response*", and "*KEGG antigen processing and presentation*" pathways from (B).
(D) Gene signature scores of selected pathways in mRegDCs.
(E) DEGs in mRegDC_2 versus remaining DC clusters, subset for ligands with cytokine activity (from "*GO:MF cytokine activity (GO:0005125)*"), to identify differentially expressed ligands which may influence the local TME. DEGs calculated using Wilcoxon rank-sum test.
(F) Expression of selected ligands between tumour mRegDC clusters, and Kaede-red DCs that have migrated to the dLN (predominantly mRegDCs, Fig. 2). Wilcoxon rank-sum test was used.

Supplementary Figure 7

Supplementary Figure 7 | Single cell profiling of tumour CD8$^+$ T cells following anti-PD-L1 treatment.

(A) UMAP of 46,342 T cells from scRNA-seq of FACS-sorted CD45$^+$CD11b$^-$ Kaede-green/red cells 48h after photoconversion of subcutaneous MC38-Ova tumours, representing tumour-infiltrating lymphocytes (TILs), and canonical marker gene expression (B).

(C) Canonical marker gene expression of CD8$^+$ T cell clusters from Fig. 5A.

(D) Gene signature enrichment of published T cell exhaustion transcriptomic signatures in T cell clusters from Fig. 5A.

(E) Proportions of cells by Kaede-green/red and treatment groups in CD8$^+$ T cells.

(F) Regression of cell cycle genes from the cycling_CD8T cluster followed by re-integration, and label transfer. Majority of the cycling cluster embedded with the CD8$^+$ T$_{EX}$ cell cluster.

(G) Proportion of cycling CD8$^+$ T cells belonging to each CD8$^+$ T cell cluster, from (F). Chi-squared test for over-representation of T$_{EX}$ CD8$^+$ T cells was used.

(H) Representative flow cytometry plots of Ki-67$^+$, GzmB$^+$ and IFNγ$^+$ expression in CD3$^+$CD8$^+$Kaede-red$^+$PD-1$^+$ T$_{EX}$ cells, following anti-PD-L1 treatment.

(I) Flow cytometry of Kaede-red$^+$PD-1$^+$CD8$^+$ T cells, showing increased frequency of cycling cells (Ki-67$^+$) following anti-PD-L1.

(J) GSEA for T$_{EX}$ CD8$^+$ T cells from anti-PD-L1 versus isotype control-treated tumours.

(K) Flow cytometry of Kaede-red$^+$PD-1$^+$CD8$^+$ T cells, showing increased frequency of GzmB$^+$IFNγ$^+$ cells following anti-PD-L1.

(L) Flow cytometry of Kaede-red$^+$PD-1$^+$ CD8$^+$ T cells, showing increased mean fluorescence intensity of GzmB following anti-PD-L1. (I, K-L) Points represent independent mice, Mann-Whitney U test was used.

Supplementary Figure 8

Supplementary Figure 8 | mRegDCs interact with CD8$^+$ T cells in murine tumours.
(A) Expression of *Cxcl16* and *Il15* in tumour DCs.
(B) Number of significant ligand-receptor interactions (from CellPhoneDB), between mRegDC and T$_{EX}$ CD8$^+$ T cells. Interactions that are well-described in existing literature to influence CD8$^+$ T cells function were identified and classified based on whether engagement of the cognate T cell receptor was activating (increase in effector function, proliferation, or survival) or inhibitory in nature.
(C-D) Representative confocal microscopy images of MC38 tumours 48h after tumour photoconversion, showing interaction between Kaede-red CCR7$^+$MHC-II$^+$ mRegDCs and Kaede-red CD3$^+$ T cells (arrows). Scale bar, 20 μm (C), 15 μm (D).
(E) Representative confocal microscopy images of MC38 tumours, showing co-localisation of CCR7$^+$MHC-II$^+$ mRegDCs and CD3$^+$CD8$^+$ T cells (arrows). Scale bar, 80 μm.
(F) Representative confocal microscopy images of MC38 tumours, showing co-localisation of CCR7$^+$MHC-II$^+$ mRegDCs and CD3$^+$CD8$^+$4-1BB$^+$Ki-67$^+$ T cells. Scale bar, 15 μm.

Supplementary Figure 9

Human colorectal cancer (syn26844071)

A Myeloid cells

n = 43,193

- Mac
- Mast
- Mono
- cDC1
- cDC2
- mRegDC
- pDC

B mRegDC_sig   CCR7

LAMP3   FSCN1

C mRegDCs

- Normal
- Tumour
- LN

n = 946

D CCR7 expression / ICOSLG expression

ns ns

N T dLN
Tissue

ns *

N T dLN
Tissue

E CCR7   TNFSF4   TNFSF9   CD274

CD70   CD80   CD86   CD74

IMvigor 210 trial (mUC, *EGAS00001004343*)

F
mUC Atezolizumab (n = 208)
R = 0.8, p < 2.2e−16

mUC Atezolizumab CR/PR (n = 45)
R = 0.87, p = 1.3e−14

mUC Atezolizumab SD/PD (n = 163)
R = 0.78, p < 2.2e−16

Effector CD8+ T cell signature / mRegDC signature

Human breast cancer (*EGAS00001004809*)

G mRegDCs only

- E
- NE

n = 368

H
Pathway / Normalised enrichment score

- HM Interferon gamma response
- Kegg FcγR-mediated phagocytosis
- GO:BP Lymphocyte costimulation
- HM Inflammatory response
- Kegg Chemokine signaling pathway
- Kegg Endocytosis

P-adj
0.01
0.02

Human NSCLC (*GSE160903*)
PICseq of sorted myeloid cell-T cell doublets

I Hallmark TNFa response

*
ns
*
*

Mono Mac MoDC cDC1 cDC2 mRegDC
Myeloid-CD8T cell doublet

J Myeloid-CD8 T cell doublet

- KEGG_CELL_CYCLE
- KEGG_JAK_STAT_SIGNALING_PATHWAY
- HALLMARK_INTERFERON_GAMMA_RESPONSE
- HALLMARK_TNFA_SIGNALING_VIA_NFKB
- GOBP_T_CELL_MEDIATED_CYTOTOXICITY

Mono Mac MoDC cDC1 cDC2 mRegDC

Scaled enrichment
0 1

Supplementary Figure 9 | mRegDC heterogeneity and interaction with CD8$^+$ T cells in human cancer.

(A) UMAP of 43,193 myeloid cells from scRNA-seq of human CRC (n = 63 patients) with paired tumour (T), dLN and normal adjacent tissue (N), and expression of mRegDC signature genes (B).

(C) UMAP of 946 mRegDCs subsetted from (A), coloured by tissue origin.

(D) Expression of *CCR7* and *ICOSLG* in mRegDCs by tissue. Wilcoxon rank-sum test was used.

(E) Expression of selected genes in mRegDCs from (C). Molecules associated with tumour-residing mRegDCs in mice were selectively expressed in tumour mRegDCs in human CRC, but not the dLN.

(F) Pearson correlation between mRegDC signature genes and effector CD8$^+$ T cell signature genes in bulk RNA-seq of 208 mUC tumours treated with atezolizumab (IMvigor 210 trial). Left, middle and right panel show all patients, clinical responders, and non-responders respectively.

(G) UMAP of 368 mRegDCs from human breast cancer (from Supplementary Fig. 1E).

(H) GSEA of mRegDCs from T cell clonotype expanders (E, i.e. responders, n = 9 patients) versus non-expanders (NE, non-responders, n = 20 patients), in breast tumours treated with anti-PD-1.

(I) Gene signature score for "*Hallmark TNFa response via NFkB*" in PICseq of NSCLC, grouped by the myeloid cell identity in each myeloid-T cell doublet. Wilcoxon rank-sum test was used.

(J) Gene signature scores of selected pathways in PICs, grouped by the myeloid cell identity.

Supplementary Figure 10

Supplementary Figure 10 | Visium spatial transcriptomics of human cancer.
(A) Expression of selected T_{EX} CD8$^+$ T cell receptors, identified from significant interactions with tumour-residing mRegDCs in spatial transcriptomics of 2 independent human CRC tumour sections.
(B) Spatial correlation of mRegDC signature scores (SFig. 1) and T_{EX} CD8$^+$ T cell receptors of interest, in Visium spatial transcriptomics of 2 independent human breast tumour sections and 1 human melanoma section (C).