Identification and characterization of specific motifs in effector proteins of plant parasites using MOnSTER.

Giulia Calia[1,2 ¶], Paola Porracciolo[3,4¶], Yongpan Chen[4,7], Djampa Kozlowski[3,4], Hannes Schuler[1,5], Alessandro Cestaro[2,6], Michaël Quentin[4], Bruno Favery[4], Etienne G.J. Danchin[4&] and Silvia Bottini[3,8&*]

[1] Free University of Bolzano, Faculty of Agricultural Environmental and Food Science, Bolzano, Italy

[2] Fondazione Edmund Mach, Research and Innovation Centre, San Michele all'Adige, Italy

[3] Université Côte d'Azur, Center of Modeling, Simulation and Interactions, Nice, France

[4] INRAE, Université Côte d'Azur, CNRS, Institut Sophia Agrobiotech, Sophia-Antipolis, France

[5] Free University of Bolzano, Competence Centre for Plant Health, Bolzano, Italy

[6] Institute of Biomembranes, Bioenergetics and Molecular Biotechnologies (IBIOM), National Research Council (CNR), Bari, Italy

[7] Present address: Department of Plant Pathology, China Agricultural University, Beijing, China

[8] Present address: INRAE, Université Côte d'Azur, CNRS, Institut Sophia Agrobiotech, Sophia-Antipolis, France

[¶] contributed as co-first author

[#] contributed as co-last author

[*] Corresponding Author: silvia.bottini@univ-cotedazur.fr

Short title: MOnSTER allows identification of motifs in effectors of plant parasites

To whom correspondence should be addressed:

Silvia Bottini, PhD

Institut Sophia Agrobiotech

UMR 1355 INRAE/7254 CNRS/UniCA

400 route des Chappes – BP 167

06903 Sophia Antipolis Cedex

France

silvia.bottini@univ-cotedazur.fr

Abstract

Plant pathogens cause billions of dollars of crop loss every year and are a major threat to global food security. Identifying and characterizing pathogens effectors is crucial towards their improved control. Because of their poor sequence conservation, effector identification in protein sequences predicted from genomes is challenging and current methods generate too many candidates without indication for prioritizing further experimental studies. In most phyla, effectors contain specific sequence motifs which influence their localization and targets in the plant. Although bacterial, fungal and oomycetes effectors have been studied extensively and conserved characteristic motifs have been identified, research on plant-parasitic nematode effectors (PPN) identified some enriched degenerate motifs in only one species so far. The different lifestyles of PPNs might reflect effectors with different functions according to the nematode's specific needs, thus presenting a high variety of characteristic motifs.

To circumvent these limitations, we have developed MOnSTER a novel tool that identifies clusters of motifs of protein sequences (CLUMPs). MOnSTER can be fed with motifs identified by *de novo* tools or from databases such as Pfam and InterProScan. The advantage of MOnSTER is the reduction of motif redundancy by clustering them and associating a score. This score encompasses the physicochemical properties of AAs and the motif occurrences. We built up our method to identify discriminant CLUMPs in candidate parasitism proteins of plant-pathogenic oomycetes. We showed the reliability of MOnSTER by identifying five CLUMPs that correspond to the known motifs: RxLR, -dEER and LxLFLAK-HVLVxxP. Consequently, we applied MOnSTER on PPN candidate parasitism proteins and identified peculiar motifs in their sequences. We identified six CLUMPs in about 60% of the known nematode candidate parasitism proteins. Furthermore, we found that specific co-occurrences of at least two CLUMPs are present in PPN candidate parasitism protein sequences bearing protein domains important for invasion and pathogenicity.

The potentiality of this tool goes beyond the candidate parasitism proteins and can be used to easily cluster motifs and calculate the CLUMP-score on any set of protein sequences.

Keywords

Motif clustering – motif scoring – effectors – plant-parasite interaction – *oomycetes - nematodes*

Authors summary

Population growth, environmental degradation and climate change are already bringing harm to human communities and the natural world that needs to be addressed rapidly. Ensuring food security for a population that will exceed 9 billion people by 2050 while preserving the environment and biodiversity is a major challenge. Agricultural pathogens, to cause the infection, secrete effector proteins that promote colonization of the host plant. Identifying and characterizing pathogens' effectors is crucial towards understanding how they manipulate the plant and better combat them. Because of their poor sequence conservation, effector identification in protein sequences predicted from genomes is challenging and current methods generate too many candidates without indication for prioritizing further experimental studies. To address these challenges, we have developed a novel tool called MOnSTER, that identifies and score clusters of motifs of protein sequences (CLUMPs). MOnSTER is an easy to use tool that can be included in any pipeline needing motif calling and will be of great

72 use to accelerate both computational and experimental studies relating to protein motif
73 discovery. Altogether our findings provide improvements in the understanding of the
74 mechanisms set up by the pathogens to infect the plant and can elucidate important
75 signatures to block the development of plant-pathogen interactions and allow to engineer of
76 durable disease resistance.

Introduction

Plant pathogens are a major threat to global food security. To cause the infection, pathogenic
organisms secrete effector proteins that promote colonization of the host plant by overcoming the
physical barriers of plant cell walls, suppressing or evading immune perception, and deriving nutrients
from host tissues [1]. Therefore, identifying and characterizing pathogens effectors is crucial towards
understanding how they manipulate the plant and better combat them. Effector proteins are often
specific to pathogens and essential for causing plant pathology, constituting targets of choice for the
development of cleaner and more specific control methods [2]–[4]. Because of their poor sequence
conservation, effector identification among the set of predicted proteins from the genome (proteome) is
challenging and current methods generate too many candidates without further indication for
prioritizing experimental studies. Classically, effector proteins are indirectly identified among the
predicted secretome based on the presence of a signal peptide for secretion and a lack of
transmembrane region [5], [6]. However, these criteria alone suffer from two main limitations. On one
side, the secretome comprises many proteins that are not effectors, on the other side some known
effectors do not possess signal peptides for secretion. In most phyla, effectors contain specific
sequence motifs which target host proteins with distinct roles in the infection process and control
virulence [7]. The best-studied example is effectors secreted via the type III secretion system (T3SS)
class of Gram-negative bacterial pathogens which are characterized by a specific motif/domain
conferring a repertoire of molecular determinants with important roles during infection [8], [9].
However, these features are not conserved in other bacteria. Indeed, gram-positive pathogens and
certain phloem- and xylem-colonizers, such as *Candidatus liberibacter* and *Xylella spp.*, do not encode
the T3SS. In these bacteria, effector delivery is dependent on the presence of the N-terminal signal
peptide, which is required for protein secretion [10]. In fungi, often effectors are small in size and
present cysteine-rich sequences [11]. Another well-characterized example is the effectors of the
oomycetes pathogens. Oomycetes are eukaryotic filamentous and heterotrophic microorganisms
among which, more than 60% of them parasitize plants [12]. Well-known plant pathogens in
oomycetes include late blight of potato, sudden oak death, root rot agents (*Phytophthora* species), and
downy mildew *Peronospora* and *Bremia* species [13], [14]. These pathogens code for two notable
classes of effector proteins RxLR and Crinkler (CRN), that can be predicted by the occurrence of the
related motifs, RxLR, -dEER and LxLFLAK-HVLVxxP in the N-terminal region downstream the signal
peptide [15]–[17].

Although for some plant pathogens such as oomycetes, effectors have been studied extensively and
characteristics motifs have been identified [18], [19], research on Plant-Parasitic Nematode effectors
(PPN) did not identify any consensus motif, conserved across multiple species. The most
economically important PPNs are the sedentary Root-Knot Nematodes (RKNs) and cyst nematodes

[20]. These sedentary parasites induce the formation of a feeding structure that serves as a constant

food source for the nematode. Other PPNs are migratory and a whole spectrum of variations exists

between endo and ecto parasites, with semi-endoparasites an intermediate between the two extremes

[21]. The different lifestyles of PPNs are expected to be reflected in their secretions, which presumably

contain effectors with different functions according to the nematode's specific needs, thus presenting a

high variety of characteristic motifs complicating their identification.

A first step toward the identification of motif characteristics of RKN effectors was performed by Vens et

al. [22]. The authors developed a bioinformatic tool, called MERCI, to identify motifs with high

occurrences in a positive dataset (known effector sequences) and absent in the negative one (non-

effector sequences). MERCI uses a graph-based approach incorporating physicochemical features of

the amino acids composing protein sequences. By analyzing the known effector sequences of the

RKN species *Meloidogyne incognita*, one of the most important known crop pathogens among all [23],

they identified 4 motifs. However, at the time of their publication, very few genomes for RKN species

were available, and the study was therefore conducted on one single RKN species. Furthermore, the

genome used at that time was later shown to be partially incomplete [24]. These limitations prevent the

generalization of the previous findings. Da Rocha et al. identified a *cis*-regulatory promoter motif (Mel-

DOG box) characteristic of dorsal gland effectors [25]. Recently, Rocha et al. used this motif combined

with other criteria to select new putative effectors and validated 14 new dorsal gland-specific candidate

effectors expressed in adult females [26]. Although all these studies have contributed to enlarging the

list of known effectors, a global characterization of their properties is still missing. Therefore, there is

an urgent need for a novel study of the properties of PPN effector sequences and motif research.

By taking advantage of the multitude of proteomes available nowadays for several PPN, we developed

a comprehensive motif mining analysis to identify characteristic motifs of candidate parasitism protein

sequences of these species. Sequence motifs are usually of constant short size and are often

repeated and conserved. Typically, motifs conform to a particular sequence pattern, where certain

positions can be constrained to a specific amino acid, whereas others are not [27]. This confers a high

degeneration of the motifs yielding a huge list of non-redundant motif sequences and consequently,

some motifs that are not characteristics of effector sequences only [28]. Furthermore, different amino

acids (AAs) can have similar physicochemical properties, thus different motif sequences can share

similar properties. However, most available motif discovery tools do not consider these properties. To

circumvent these limitations, we have developed MOnSTER a novel tool that identifies clusters of

motifs of protein sequences (CLUMP) and associates a score to each CLUMP. This score

encompasses the physicochemical properties of AAs and the motif occurrences. Overall, one of the

key advantages of MOnSTER is that it reduces the redundancy of motifs found by *de novo* tools.

Furthermore, already known motifs available in publicly available databases such as Pfam [29] and/or

InterProScan [30] can also be used as input of MOnSTER to identify discriminant CLUMPs.

We built up our method to identify discriminant CLUMPs in 1743 candidate parasitism proteins of

plant-pathogenic oomycetes. We showed the reliability of MOnSTER by identifying 5 CLUMPs that

correspond to the known motifs: RxLR, -dEER and LxLFLAK-HVLVxxP. After this proof of concept, we

applied MOnSTER on PPN effector proteins and identified peculiar motifs in their sequences at an unprecedented level. We selected a set of 4395 protein sequences from 13 PPN species belonging to the genera *Meloidogyne, Globodera, Heterodera, Radopholus and Bursaphelenchus*. We identified 6 CLUMPs present in 60% of the known effectors (positive dataset). Of note these CLUMPs were found in only 5% of the sequences of the negative datasets, thus highlighting the enrichment of the identified motifs in effector sequences. Furthermore, we found a specific co-occurrence of at least two CLUMPs in PPN candidate parasitism protein sequences bearing protein domains important for invasion and pathogenicity.

The potentiality of this tool goes behind the candidate parasitism proteins and can be used to easily cluster motifs and calculate the CLUMPs score on any set of protein sequences. Furthermore, we also provide a new scoring system capable of measuring the physicochemical properties of motifs grouped in CLUMPs and a motif alignment algorithm to better explore chemical-physical properties within the CLUMPs. MOnSTER is freely available at https://github.com/paolaporracciolo/MOnSTER_PROMOCA.git.

Materials and methods

Datasets

Oomycetes
We used proteins from five oomycetes species to create the input datasets for MOnSTER, namely *Phytophthora infestans*, *Phytophthora sojae*, *Phytophthora ramorum*, *Hyaloperonospora arabidopsidis* and *Bremia lactucae*.

Positive dataset
The positive dataset consists of 1743 effector proteins belonging to the aforementioned oomycetes obtained from a concatenation of proteins selected from PHI-base database (v4.14) [31], Uniprot (release 2023_02)[32], and the work of Haas et al., (2009) [33], in which they have manually curated the annotations of the proteins. Since the proteins come from different sources, we used CD-HIT (v4.8.1) [34] with the parameters in **Supplementary information,** to filter out identical protein sequences. A total of 1283 proteins are annotated as RxLR effectors, 377 as Crinkler effectors and the last 83 sequences are proteins with no previously identified motif and known to be involved in the host-pathogen interaction.

Negative dataset
Proteins in the negative dataset derive all from Uniprot (release 2023_02) and from the oomycetes species cited before filtered from proteins included in the positive dataset and for evident effector-related annotations. Due to the large amount of non-effector proteins remaining from the filtering we firstly used 'cd-hit' to reduce protein sequence redundancy and then, to also reduce the unbalance of the final dataset we refined the selection taking only the representative sequences of the orthogroups

191 found with Orthofinder (v2.5.4) [35]. In total 3009 non effector proteins are included in the negative
192 dataset.
193
Motif Discovery
The last input file consists in a list of motifs identified as enriched in the sequences of the positive
dataset compared to the sequences of the negative one. We used MERCI and STREME (v5.5.1) [36],
with parameters detailed in **Supplementary information**. We imposed different lengths for motifs
prediction to be inclusive but more stringent on the motifs in which we are interested. STREME's
output is a list of motifs. Hence, we used the tool FIMO (v5.5.1) [37], with default parameters to
extract 246 degenerated motifs from the 4524 different motifs.
We obtained the following numbers of non-redundant motifs: 19 with MERCI and 246 with STREME.
Then, we removed the identical motifs and created a single non-redundant list containing all the motifs
in the same format, which resulted in 265 different motifs.
204
205 **Plant Parasitic Nematodes (PPNs)**
206
Positive dataset
The positive dataset contains candidate parasitism proteins selected to be likely secreted by PPNs in
their plant host and belonging to 13 species (*Meloidogyne incognita, Meloidogyne javanica,*
Meloidogyne arenaria, Meloidogyne hapla, Meloidogyne chitwoodi, Meloidogyne graminicola,
Globodera rostochiensis, Globodera pallida, Heterodera havenae, Heterodera glycines, Heterodera
schachtii, Radopholus similis, Bursaphelenchus xylophilus). We collected candidate parasitism protein
from literature mining. More precisely we considered as candidate parasitism protein those proteins for
which *in-situ* hybridization experiments showed that the corresponding transcript is present in
nematode secretory glands (dorsal or sub-ventral), implying that these proteins are likely secreted by
the nematodes into the host plant. The literature mining led to the extraction of 163 proteins from NCBI
GeneBank thanks to the NCBI 'entrez' API. We also manually extracted 41 sequences from the
publications' core text and Supplementary information. In addition, we downloaded 41 sequences from
WormBase ParaSite (www.parasite.wormbase.org, vWBPS17-WS282 [38], [39]), and eight
sequences from nematode.net [40]. In total we obtained 229 candidate parasitism protein. We
extended the positive dataset with proteins that are non-redundant homologs of the previous
candidate parasitism proteins in PPN proteomes. We first used cd-hit-2D with parameters in
Supplementary information, to cluster sequences from PPNs proteomes and candidate parasitism
proteins [41]. We then pooled all the candidate parasitism proteins from closely related *Meloidogyne*
species (e.g., *M. incognita*, *M. javanica* and *M. arenaria*) and scanned each corresponding proteome
with this multi-species set of sequences using cd-hit. Since the remaining species are genetically
distinct, we then scanned each proteome with the relative set of candidate parasitism proteins, except
for *H. havenae* and *M. chitwoodi* for which no proteomes were currently available. We merged the two
sets of selected candidate parasitism proteins and we performed CD-HIT intra- and inter-species to
reduce dataset redundancy (parameters in **Supplementary information**), retaining only sequences

having more than 1% divergence and aligning on more than 80% of their length (the longest sequence

from each cluster was kept). The final positive dataset includes 546 candidate parasitism proteins from

13 species.

Negative dataset

The negative dataset is composed of 3849 protein sequences that we obtained by selecting genes

widely conserved across the nematode tree of life and close outgroup species, including many species

that are non-parasites. Specifically, we filtered the results from a previous analysis [42] and only

retained genes from orthogroups i) conserved in more than 90% (62/64) of the analyzed species

including two tardigrade species (outgroups), and ii) presenting less than 10

genes/species/orthogroups to avoid multigenic families, which would lead to overrepresentation of

some proteins. To remove the redundancy, we used the same strategy as for the positive dataset

(cdhit2D first and then CD-HIT).

Motif Discovery

Using the aforementioned software in the same configuration we obtained the following numbers of

non-redundant motifs: 40 with MERCI and 229 with STREME applying FIMO. In total, we obtained 269

different motifs.

All datasets are available at https://github.com/paolaporracciolo/MOnSTER_PROMOCA.git and in

Supplementary tables 1.1-1.2 and 2.1-2.2.

MOnSTER pipeline

The MOnSTER (MOtifs of cluSTERs) pipeline is composed of three main steps as described in **Figure**

1 and in the following paragraphs.

Feature calculation

The first step of the pipeline concerns the calculation of parameters that describe protein sequences

(**Figure 1A**). To allow an easy calculation of the features on any dataset, we calculated the sequence

length and used *ProteinAnalysis* class from the *Bio.SeqUtils.ProtParam,* a python sub-package to

select 13 additional features based on individual AA properties, belonging to 4 categories:

- secondary structure propensity 'helix' (V, I, Y, F, W, L), 'turn' (N, P, G, S), and 'sheet' (E, M, A,

L)).

- amino-acids dimensions ('tiny' (A, C, G, S, T) and 'small' (A, C, F, G, I, L, M, P, V, W, Y)).

- pH ('basic' (H, K, R), 'acid' (B, D, E), and 'charged' (H, K, R, B, D, E)).

- physicochemical properties ('hydropathy-score', 'polar' (D, E, H, K, N, Q, R, S, T), 'non-polar'

(A, C, F, G, I, L, M, P, V, W, Y), 'aromatic' (F, H, W, Y), and 'aliphatic' (A, I, L, V)).

We performed feature calculations on the positive and negative datasets and the list of motifs. At the
end of this step, we obtained three tables of features, one for each of the input datasets (positive,
negative datasets and the list of motifs).
273

Clustering
This step allowed to cluster motifs based on their properties described by the 13 features. To make the
features comparable to each other, we performed data normalization by using the *StandardScaler*
method from *sklearn.preprocessing* [43]. This normalization consists of the removal of the mean and
the scaling to unit variance.
Then, we performed a hierarchical clustering of the motifs using the Euclidian distance. We then
divided the resulting tree into clusters of motifs of proteins (CLUMPs) selecting the threshold distance
that minimized the Davies-Bouldin score [44].
For each CLUMP, we removed the redundant motifs. Briefly, we identified motifs that shared a core
sequence (for example: 'HWT in HWTQ' and 'GHWTQ'), and we only retained the cores (for instance:
"HWT") in the CLUMPs.
285

Scoring
The final objective is to identify the CLUMP(s) with the highest discriminative power concerning the
positive dataset. Thus, we conceived a new score called the MOnSTER score, to rank the CLUMPs by
their discriminative power.
The MOnSTER score is composed of three parts: the CLUMP score and two modified versions of the
Jaccard index.
292

293 ***CLUMP score***
294 This score considers the AA composition of the motifs belonging to each CLUMP concerning the
295 preferences of the sequences of the positive dataset. The procedure that we implemented to calculate
296 this score is shown in **Figure 1B**.
297

298   a)   Feature selection
299 We used the Mann-Whitney test to identify the features whose values were significantly different
300 between the positive and negative datasets. We only retained the statistically significant features, with
301 a p-value < 0.05. Then, we assigned them a score, by calculating -Log(p-value) of each feature. We
302 will refer to it as the 'feature weight' hereafter.
303

304   b)   Average calculation
305 For each of the selected features (ranging from one to *f*), we calculated the average value for the
306 positive dataset, the negative dataset, and each CLUMP (ranging from zero to *c*). We will refer to
307 these values with the notation: μ_f^+, μ_f^- and $\mu_f^{CLUMP_c}$, respectively.
308

c)   CLUMPs sorting

We compared the averages of the positive and negative datasets for each feature and sorted CLUMPs
accordingly.

Thus, if the $\mu_f^+ \geq \mu_f^-$, the CLUMPs averages would be sorted in ascending order.

Otherwise ($\mu_f^+ < \mu_f^-$), CLUMPs averages would be sorted in descending order.

d)   CLUMPs voting

For each feature, and each CLUMP, we divided the CLUMP into two groups accordingly to the
following statements:

If $\mu_f^+ \geq \mu_f^-$: CLUMPs with $\mu_f^{CLUMP_c} \geq \mu_f^+$ have a vote from 1 to the number of CLUMPs with an
increment of 1, otherwise the score is set to 0.

If $\mu_f^+ < \mu_f^-$: CLUMPs with $\mu_f^{CLUMP_c} < \mu_f^+$ the vote attributed goes from 1 to the number of
CLUMPs, otherwise it is 0.

e)   CLUMPs scoring

For each CLUMP (ranging from zero to c), for each feature (ranging from one to f), we multiplied the
feature-vote by the 'feature weight' (W_f) and summed-up to obtain a CLUMP-vote. Then we scaled
each CLUMP-vote to a range from 0 to 1 using the following formula:

$$CLUMPscore_c = \frac{V_c - \min(V)}{(\max(V) - \min(V))}$$

where:

V is the list of CLUMPs votes and V_c is calculated as:

$V_c = \sum_{features[1,f]} \left(vote_f \subset CLUMP_c \right) W_f$

Occurrences indexes
The two indexes respectively consider: i) the occurrences of the motifs, for each CLUMP, in the
positive dataset compared to the negative, and ii) the number of positive sequences containing the
motifs in each CLUMP concerning the negatives (**Figure 1C**).

a)   CLUMPs occurrences

We calculated the occurrences of the motifs in each CLUMPs in the two datasets (positive and
negative).

b)   I's scores

We propose two ways to calculate the dissimilarity between two sets that will be called I$_1$ and I$_2$
hereafter.

To obtain I_1, we calculated the number of occurrences of the motifs for each CLUMP (ranging from

zero to c) in the negative dataset over the number of occurrences of the motifs of the same CLUMP in

the positive dataset, using the following equation:

$$I_{1 \, \forall \, CLUMP[0,c]} = \frac{1}{2} \left(1 - \frac{\sum \Delta_- \subset CLUMP_c}{\sum \Delta_+ \subset CLUMP_c} \right)$$

Where:

Δ_- and Δ_+ the number of occurrences of the motifs of the CLUMP in the negative or

in the positive dataset, respectively.

To obtain I_2, for each CLUMP (ranging from zero to c), we calculated the number of sequences of the

negative dataset that contain at least a motif of the CLUMP, over the number of sequences of the

positive dataset that contain at least a motif of the same CLUMP, accordingly to the following formula:

$$I_{2 \, \forall \, CLUMP[0,c]} = \frac{1}{2} \left(1 - \frac{\sum seq_- \subset CLUMP_c}{\sum seq_+ \subset CLUMP_c} \right)$$

Where:

seq_- is the number of sequences of the negative dataset containing at least a motif of the CLUMP.

seq_+ is the number of sequences of the positive dataset containing at least a motif of the CLUMP.

The ½ factor is applied to have values between 0 and 0.5 for each Index to have equal weight in the

final score, and (1 – Index) is to consider the degree of dissimilarity rather than similarity.

MOnSTER score
The MOnSTER score, for each CLUMP (from zero to c), is the sum of the corresponding CLUMP

score, and the two I indexes:

$$MOnSTERscore_c = CLUMPscore_c + I_{1c} + I_{2c}$$

368 **PRO-MOCA: a novel method to create motif logo of CLUMPs**

369
To create motif logos for each CLUMP, we developed a novel method. PRO-MOCA (PROtein-MOtifs

Characteristics Aligner) aligns protein motifs based on the characteristics of the amino acids as shown

in **Supplementary figure 1**. The first step is to define the alphabets associated with each

characteristic that can be used to represent the motifs (**Supplementary figure 1**a). We have defined

four alphabets, namely: "chemical", "hydrophobicity", "charge", "secondary structure propensity"

(details for each alphabet are included in the **Supplementary information**).

These alphabets are easily modifiable and other alphabets can be included. Different CLUMP logos

can be obtained depending on the alphabet chosen. Secondly, PRO-MOCA uses the selected

alphabet to translate the AA sequences of each motif in a CLUMP in the new alphabet

(**Supplementary figure 1**b). The third step is the alignment (**Supplementary figure 1**c). Briefly,

PRO-MOCA screens the translated motif sequences of a CLUMP looking for a "summit position" with

the highest frequency of the same "letter" of the novel alphabet (further details in supplementary
materials). Once this position is identified, all motifs are aligned accordingly (**Supplementary figure**
1d). Since the motifs of a CLUMP can have different lengths, after the alignment, PRO-MOCA
calculates the number of gaps to add at the extremities to make all motifs having the same length.
Importantly, gaps are not allowed inside the motif sequences. The last step of the method is to re-
translate the aligned motifs in the original AA sequences (**Supplementary figure 1**e). The alignment
is ready to feed a program to create logos. Here we used the tool Weblogo3 [45].

PPNs candidate parasitism protein domains mining analysis

389
390
To investigate the relationship between the selected CLUMPs and functional domains in candidate
parasitism proteins we first selected proteins from the positive datasets containing at least one
occurrence of a selected CLUMP (311 proteins in total). Then we predicted the protein domains with
InterProScan (v5.54-87.0) [30] with default parameters. From the results, we extracted the proteins
containing the most frequent predicted domains and considered only entries coming from: MobiDB-lite,
Coils, CDD, PANTHER, Pfam and ProSitePatterns. Afterwards, we also predicted the presence of
Signal Peptide (SP) (SignalP4.1 [46]) and TransMembrane (TM) domain regions (TMHMM2.0 [47]).
We obtained 258 proteins having at least a CLUMP and one of the aforementioned predicted
domains, SP or TM.

In situ hybridisation (ISH) and *N. benthamiana* agroinfiltration

401
M. incognita strain "Morelos" was multiplied on tomato (*Solanum lycopersicum* cv. "Saint Pierre")
growing in a growth chamber (25°C, 16 h photoperiod). Freshly hatched J2s were collected and ISH
performed as previously described [48], [49]. The *M. incognita Minc3s00056g02931/MiEFF72* coding
sequence (CDS) lacking the signal peptide for secretion was amplified by PCR with specific primers
(EFF72_F: 5'-AAAAAGCAGGCTTCACCATGAATACTGCTGACAAGACACAG-3' and EFF72_R: 5'-
AGAAAGCTGGGTGTTAGAACAAAGCTCGCACTGC-3') and inserted into the pDON207 entry vector.
Antisense probe was amplified using EFF72_R from entry vector. Sense probe was amplified using
EFF72_F and used as negative control. Images were obtained with a microscope (Axioplan2, Zeiss,
Germany).
The *M. incognita MiEFF72* CDS lacking the signal peptide was recombined in pK7FGW2
(P35S:eGFP-MiEFF72) with Gateway technology (Invitrogen). The construct was sequenced (GATC
Biotech) and transferred into *Agrobacterium tumefaciens* strain GV3101. Transient expression was
achieved by infiltrating *N. benthamiana* leaves with *A. tumefaciens* GV3101 strain harbouring the
GFP-fusion construct, as previously described [50]. Leaves were imaged 48 hours after
agroinfiltration, with an inverted confocal microscope (LSM880, Zeiss, Germany) equipped with an
Argon ion and HeNe laser as the excitation source. GFP emission was detected selectively with a 505-
530 nm band-pass emission filter.

Results & Discussion

MOnSTER identified five CLUMPs containing known motifs characteristics of oomycetes effector protein sequences

Characteristic motifs of oomycetes effector proteins are well-known in the literature, such as RxLR, -dEER and LxLFLAK-HVLVxxP [15]. Thus, we reasoned to apply our novel tool, MOnSTER, on oomycetes effectors to test its ability to recover well-characterized motifs. We compiled a set of 4752 oomycetes proteins, comprising 1743 effectors and 3009 non effectors, from five oomycetes species. We performed motif discovery on this set of proteins using MERCI and STREME and we identified 265 significantly enriched motifs (see methods for further details). Then we fed MOnSTER with these motifs and we obtained 11 CLUMPs (**Supplementary table 3**), employing the Davis-Bouldin score, as a criterion to cut the tree. By selecting CLUMPs having a MOnSTER score greater than the median of the overall scores we identified six CLUMPs (CLUMP7, 4, 10, 6, 2 and 9), the first five best-scoring CLUMPs, accordingly to the MOnSTER score, correspond to the known motifs (**Figure 2**). In **Supplementary figure 2** we can also observe that the motifs are respectively grouped in two clades, the two characteristics motifs of CRN-effectors (LxLFLAK and HVLVxxP), form a separate subclade on the right, while the RxLR and -dEER motifs fall into the left clade, resembling the family distinction of effectors to which they belong. More precisely RxLR motifs are divided into two different CLUMPs; CLUMP6 containing only RYLR and RFLR motifs, and CLUMP10, containing other RxLR motifs and included in the same sub-clade of the dEER motif (CLUMP2). The last best-scoring CLUMP contains no known motifs, perhaps suggesting a novel putative motif for oomycetes effectors to investigate. Since oomycetes effectors characterization is not in the scope of this article, we did not consider this last CLUMP for further analysis. In support of that, CLUMPs 7, 4, 10, 6 and 2 are present in 1205/1743 effectors (~70% of the sequences in the positive dataset) while in combination with the last significant CLUMP (CLUMP9) only two more sequences can be detected.

Thus, we investigated the occurrences and co-occurrences of the five selected CLUMPs in oomycetes effectors and non-effectors (**Supplementary figure 3**). For the effectors we deeply analyzed the two distinct families; in total we found that 68% of the RxLR-effectors in the positive dataset contain the motifs in CLUMPs associated with the RxLR motif (CLUMP10, 6 and 2). In particular, CLUMP10 and 6 are present alone in 41% of the RxLR-effectors (1238/1743 RxLR-effectors), while 19% of the RxLR-effectors contained the co-occurrence of these CLUMPs with the CLUMPs representing the dEER motif (CLUMP2). This reflects the importance of the RxLR motifs in the effector sequences and the role of the attached dEER [51]. On the other hand, the co-occurrence of CLUMPs specific for LxLFLAK and HVLVxxP (CLUMP7 and 4), in CRN-effector sequences accounts for 67% of the relative sequences in the positive dataset (377/1743). The high co-occurrences rate of CLUMP7 and 4 is strongly in agreement with the presence of LxLFLAK and HVLVxxP motif marking the beginning and the end of the DWL-domain in the Crinkler-effector family [33]. For the negative dataset, instead, only 15% of the sequences show the presence of CLUMP-motifs with a huge decrease in CLUMPs co-

occurrences. Overall co-occurrences, indeed, are present in around 30% of positive sequences and in
1% of negative ones.

Previous research showed that the motifs characteristics of oomycetes effectors have strong
sequence position preferences [52]–[54]. Thus, we plotted the CLUMPs occurrences in the positive
versus negative dataset (**Supplementary figure 4**). Indeed, we can observe that the CLUMPs are
concentrated at the beginning of the sequence in positive sequences and conversely spread around
the sequence of negative dataset proteins. More precisely the five most interesting CLUMPs are
condensed in the first 40% of the sequence with a higher preference at the very beginning and around
30% of the sequence probably corresponding to the N-terminal of the protein in which the target motifs
lie.

Altogether these results highlight the ability of MOnSTER to identify CLUMPs containing biologically
relevant motifs.

MOnSTER allowed to identify six CLUMPs characteristics of nematode candidate
parasitism proteins
474

The application of MOnSTER of the oomycetes effectors served as a proof of concept of our
methodology. Thus, we moved to the characterization of nematode candidate parasitism sequences
for which no characteristic motifs have been identified yet. We collected a set of 4395 proteins,
including 546 well-known candidate parasitism proteins and 3849 proteins in the negative dataset,
coming from 13 nematode species. By running motif discovery analysis as for the previous dataset, we
found 269 motifs enriched in the candidate parasitism protein sequences. By applying MOnSTER with
the previous configuration, the 269 input motifs were grouped into 11 CLUMPs. Six best-scoring
CLUMPs were selected using the median as the significant threshold (**Supplementary table 4**).
Similar to the oomycetes results, we observe two main clades (**Figure 3**): the second and the third
best scoring ones (CLUMP2 and 5 respectively) form a single clade while the other significant
CLUMPs (CLUMP1, 3, 7 and 10) are distributed in the bigger clade with the non-significant ones.
Overall, we found at least one occurrence of one of the six CLUMPs in almost 60% of sequences from
the positive dataset compared to 5% of sequences from the negative.

Then we investigated the presence of the six CLUMPs in each of the 13 PPN species present in the
dataset. **Figure 4** shows the abundance of the six best-scoring CLUMPs in the species according to
their phylogeny tree. The first three species are the most represented in the positive dataset.
Interestingly very distant species show similar CLUMPs frequencies thus suggesting that they might
share common characteristics at the sequence level for accomplishing similar functions. Furthermore,
we could identify characteristic CLUMPs also for species represented in the dataset with very few
sequences reinforcing the previous observation. Overall, this analysis suggests that CLUMPs might be
associated with the functional properties of PPN nematodes.

Finally, we focused on the positional sequence preferences of CLUMPs in candidate parasitism
protein sequences (**Supplementary figure 5**). In general, we observe a difference in the position
preferences of the best-scoring CLUMPs between positive and negative dataset sequences. The six
CLUMPs tend to occur more frequently in the middle of the sequences in candidate parasitism

proteins (positive dataset), with more abundance in central (around 50% of the sequence) and terminal (around 70%), positions. The same CLUMPs are rare in the central position of the negative dataset protein sequences (negative dataset). Contrary to the properties of oomycetes effectors, whose characteristics CLUMPs occur mainly at the beginning of the sequence, PPN candidate parasitism proteins showed a different pattern of occurrences, privileging a central – C terminal occurrence.

Co-occurrences of different CLUMPs are associated with functional protein domains.

We investigated the co-occurrence patterns of CLUMPs in the PPNs candidate parasitism protein sequences (all possible combinations of co-occurrences are reported in **Supplementary figure 6)**. Overall, we notice that CLUMPs tend to co-occur more frequently in the sequences of the positive dataset than in the negative one, despite the positive set being smaller than the negative one. 30% of candidate parasitism protein sequences show co-occurrences of the six selected CLUMPs, while in the sequences from the negative dataset, co-occurrences, are present in less than 1% of the sequences. As observed for oomycetes, some CLUMPs tend to be present alone, while others tend to co-occur with specific CLUMPs. This suggests that different classes of nematode candidate parasitism proteins might exist, similar to the oomycetes effectors. Interestingly, among the 311 candidate parasitism proteins bearing at least one occurrence of one of the six selected CLUMPs, 72 do not have a predicted signal peptide, consisting of 55% of the proteins in the positive dataset not having the signal peptide. Of note, this is a similar percentage to the percentage of proteins bearing both the CLUMPs and the signal peptide, suggesting that CLUMPs characterize sequence properties beyond the type of secretion. Furthermore, similar patterns of co-occurrences of CLUMPs in candidate parasitism proteins bearing or not the signal peptide are observed with slightly higher co-occurrence presences in the sequences not having the signal peptide (**Supplementary figure 7**). Importantly, there is no relationship between the sequence length and the number of co-occurrences possibly suggesting a functional role for CLUMPs co-occurrences (**Supplementary figure 8**).

To inspect further a putative functional role of CLUMPs in candidate parasitism protein sequences, we queried the sequences having at least one CLUMP or a co-occurrence of multiple CLUMPs against several protein domain databases (see Methods, results in **Figure 5** and **Supplementary table 5**). Among the 311 candidate parasitism protein sequences bearing at least one occurrence of at least one of the six CLUMPs, 84 also have at least an occurrence of a known protein domain. The most recurrent hits are the coil domain, intrinsically disordered domain and the presence of the signal peptide (SP) followed by the pectate lyase domain, glycosyl hydrolase family 5, Stichodactyla toxin (ShK) domain, 14-3-3 family and cysteine-rich domain. Importantly, none of these domains was also found in the sequences from the negative dataset bearing at least one occurrence of at least one of the six CLUMPs. Interestingly, we observe the almost exclusive association between CLUMPs and functional domains, mainly when multiple CLUMPs co-occur in candidate parasitism protein sequences.

The strongest association that we observe is between the co-occurrences of CLUMPs 7 and 10 and the glycosyl hydrolase family 5 domain on one hand and the co-occurrences of CLUMPs 3, 7, 10 and

the cysteine-rich domain, on the other hand. Specifically, all 23 candidate parasitism protein
sequences containing the co-occurrences of CLUMP 7 and 10 bear also the glycosyl hydrolase family
5 domain. By inspecting the position of CLUMPs occurrences within the sequences, we observed that
the two CLUMPs are flanking the domain: CLUMP7 is consistently present at the beginning of the
sequence and consequently of the domain, while CLUMP10 mostly concentrates at the end of the
domain, around 60-80% of the sequences (**Supplementary figure 9**). Examples of these genes in
nematodes is poorly characterized and likely resulting from horizontal transfer [55], [56]. Similarly, all
17 sequences presenting the co-occurrence of CLUMPs 3, 7,10 also contain the cysteine-rich domain.
Cysteine-rich domain and CAP protein are known to be involved in the virulence of nematodes [57].
They are expressed in both plants and pathogens; in the latter, they are important for their virulence by
suppressing the host's immune responses and promoting colonization. Interestingly, these sequences
do not contain disordered regions or coil domains, consistently with unique conserved sandwich fold
with a large central cavity of these kinds of proteins [58]. 16 out of 19 sequences presenting co-
occurrences of CLUMPs 2, 3 have also the 14-3-3 family domain, a eukaryotic-specific protein family
with a general role in the signal transduction [59]. We also observe only one motif from CLUMP 2 in
these sequences (KDKM) and 4 from CLUMP 3 (NKDKAC, KMKG, PTHPIR, PTHP). 13 out of 34
sequences bearing only CLUMP 1 also contain the pectate lyase domain. Of note, these sequences
do not contain coiled or disordinate regions, and only seven show the presence of the SP. Pectate
lyase enzymes in nematodes facilitate penetration in plant-cell walls made of pectin [60]. Numerous
recent reports showed that these enzymes are produced in specialized nematode gland cells and
secreted during the parasitism process. In the case of sedentary endo-parasitic nematodes, this
occurs mainly during juvenile migration through the root tissue, when these enzymes play a crucial
role in the maceration of the plant tissue facilitating the infection [61]. Finally, eight out of 22
sequences bear the co-occurrences of CLUMPs 2, 5 and the ShK domain. Although the exact
biological function of the ShK domain remains unclear, previous reports have shown that this domain
might be associated with immunosuppression [62], [63].
Overall, these findings highlight that specific CLUMPs co-occurrences are associated with specific
functional domains with roles in invasion and/or infection and might suggest different classes of
candidate parasitism proteins cross-species.
570

CLUMPs screening yielded the identification of a novel effector in *M. incognita* validated by *in*
***situ* hybridization.**
To inspect whether the novel-identified CLUMPs could also help to find new effectors, we focused on
the selection of a novel putative effector to validate experimentally. Thus, we selected all proteins of
Meloidogyne incognita proteome bearing the signal peptide for secreted proteins and no
transmembrane domain. Then we screened these sequences and retrieved the ones containing at
least one motif of the six significant CLUMPs. Among them, 23% contain at least one occurrence of
motifs in CLUMP5 (Supplementary Table 6). Since this is the most abundant CLUMP in this species,
we decided to focus on this one to identify a putative candidate to validate experimentally. By literature
mining, we refined our list, by sorting out seven sequences that were already experimentally validated

by previous studies (Supplementary Table 6). Then we filtered out any candidates having homologs in
species other than root-knot nematodes and more than two gene copies to avoid dealing with
multigene families according to [42]. Finally, among these eight new putative effector sequences, we
studied the pattern of expression of one candidate: *MiEFF72* (*Minc3s00056g02931*) by performing *in*
situ hybridisation (ISH). A specific signal was detected in the subventral oesophageal gland cells of
pre-parasitic J2s after hybridisation with digoxigenin-labelled MiEFF72 antisense probes (**Figure 6A**).
No signal was detected in pre-J2s with sense negative controls. MiEFF72 fused to the C-terminus of
GFP was transiently expressed in *N. benthamiana* leaf epidermis. GFP fluorescence was detected in
the cytoplasm and in cytoplasmic vesicles (**Figure 6B**). This finding suggests that MiEFF72 be
secreted and play a role *in planta* in nematode parasitism.
591

Conclusions

This work is structured around three main aims: (1) the development of a novel method to cluster and
score discriminant motifs of protein sequences called MOnSTER, (2) the validation of the MOnSTER
results by applying it to identify CLUMPs specific to candidate parasitism protein sequences of
oomycetes (3) the application of MOnSTER to protein sequences from plant-parasitic nematodes with
unprecedented discriminant motifs detection.
The application of MOnSTER on oomycetes yielded the identification of five CLUMPs corresponding
to the well-known effector-related motifs like RxLR-dEER and LxLFLAK-HVLVxxP motifs in
oomycetes. This demonstrated that the novel scoring method introduced by MOnSTER is a good
parameter with which calculate CLUMP specificity for effector protein sequences. When applied to the
nematodes candidate parasitism protein, MOnSTER found six novel CLUMPs, not previously
characterized. The main advantage of MOnSTER is that the definition of CLUMPs allowed us to
reduce the degeneration of 265 and 269 motifs (oomycetes and nematodes respectively), to 11
CLUMPs. Candidate parasitism protein sequences of both pathogens show some common
characteristics. Indeed, selected CLUMPs-motifs are present in about 70% of the input proteins for
oomycetes and 60% in PPN compared to 15% and 5% in the negative dataset proteins, respectively.
Furthermore, around 30% of candidate parasitism protein sequences have co-occurring CLUMPs, in
contrast with less than 1% of the negative dataset sequences, in both applications. The main
difference between candidate parasitism protein specific motifs of the two pathogens is the positional
preference: the beginning of the sequence for oomycetes and central C-terminal for PPNs. This
highlights MOnSTER ability to cluster motifs specifically relevant for candidate parasitism protein
sequences without privileging any portion of the sequence, like other motif discovery tools.
Concerning the novel identified motifs for PPNs candidate parasitism proteins, we observed that the
pattern of occurrences and co-occurrences of CLUMPs in candidate parasitism protein sequences is
associated with specific functional domains and might suggest the existence of different classes of
candidate parasitism proteins. Importantly we did not observe any species-related preferences thus
implying the generality of these results.

In conclusion, MOnSTER quantifies the motifs and sequence properties in each dataset provided, thus
allowing a wide application to other protein classes. Since the MOnSTER score considers the
physicochemical properties and occurrences of motifs in CLUMPs concerning the protein sequences
provided, it works without the need for a reference dataset. Furthermore, the MOnSTER scores are
normalized values, therefore, allowing direct comparison between different studies.

Our results highlighted that MOnSTER is a powerful new method to cluster and score discriminant
motifs in protein sequences according to their physicochemical properties and pattern of occurrences.
It is also a tool that can be easily used on any set of protein sequences and a list of motifs. Therefore,
by constructing a dataset of positive candidate parasitism protein sequences and a negative dataset,
MOnSTER can be also used to identify CLUMPs characteristics of fungal or bacterial candidate
parasitism proteins. As such, MOnSTER can be included in any pipeline needing motif calling and will
be of great use to accelerate both computational and experimental studies relating to protein motif
discovery.

Data availability

635 The source code and related data are available at:
636 https://github.com/paolaporracciolo/MOnSTER_PROMOCA.git

Funding

638 This work was supported by the French government, through the UCA JEDI Investments in the Future
639 project managed by the National Research Agency (ANR) under reference number ANR-15-IDEX-01.

Competing Interests

641 The authors declare that they have no competing interests.

Financial Disclosure statement

644 The funders had no role in study design, data collection and analysis, decision to publish, or prepara-
645 tion of the manuscript.

Acknowledgements

648 Microscopy work was performed at the SPIBOC imaging facility of Institut Sophia Agrobiotech, and we
649 thank Dr Olivier Pierre for is availability.

CRediT author contribution

GC: Methodology, Software, Validation, Formal analysis, Writing – original draft, Visualization. PP:
Methodology, Software, Writing – original draft. JC: Investigation, Resources. DK: Software,
Resources. HS: Writing – review & editing, Supervision. AC: Writing – review & editing, Supervision.
MQ: Investigation, Resources, Writing – review & editing, Supervision. BF: Investigation, Resources,
Writing – review & editing, Supervision. EGJD: Conceptualization, Methodology, Writing – review &

editing, Supervision. SB: Conceptualization, Methodology, Writing – original draft, Writing – review &
editing, Supervision, Project administration.

References

[1] T. Y. Toruño, I. Stergiopoulos, and G. Coaker, "Plant-Pathogen Effectors: Cellular Probes Interfering with Plant Defenses in Spatial and Temporal Manners," *Annu. Rev. Phytopathol.*, vol. 54, pp. 419–441, Aug. 2016, doi: 10.1146/annurev-phyto-080615-100204.

[2] A. Haegeman, S. Mantelin, J. T. Jones, and G. Gheysen, "Functional roles of effectors of plant-parasitic nematodes," *Gene*, vol. 492, no. 1, pp. 19–31, Jan. 2012, doi: 10.1016/j.gene.2011.10.040.

[3] C. Selin, T. R. de Kievit, M. F. Belmonte, and W. G. D. Fernando, "Elucidating the Role of Effectors in Plant-Fungal Interactions: Progress and Challenges," *Front. Microbiol.*, vol. 7, 2016, [Online]. Available: https://www.frontiersin.org/articles/10.3389/fmicb.2016.00600

[4] D. M. Bird, J. T. Jones, C. H. Opperman, T. Kikuchi, and E. G. J. Danchin, "Signatures of adaptation to plant parasitism in nematode genomes," *Parasitology*, vol. 142 Suppl 1, no. Suppl 1, pp. S71-84, Feb. 2015, doi: 10.1017/S0031182013002163.

[5] J. Sperschneider, A. H. Williams, J. K. Hane, K. B. Singh, and J. M. Taylor, "Evaluation of Secretion Prediction Highlights Differing Approaches Needed for Oomycete and Fungal Effectors," *Front. Plant Sci.*, vol. 6, p. 1168, 2015, doi: 10.3389/fpls.2015.01168.

[6] H. Sonah, R. K. Deshmukh, and R. R. Bélanger, "Computational Prediction of Effector Proteins in Fungi: Opportunities and Challenges," *Front. Plant Sci.*, vol. 7, p. 126, 2016, doi: 10.3389/fpls.2016.00126.

[7] L. Liu *et al.*, "Arms race: diverse effector proteins with conserved motifs," *Plant Signal. Behav.*, vol. 14, no. 2, p. 1557008, Jan. 2019, doi: 10.1080/15592324.2018.1557008.

[8] P. Dean, "Functional domains and motifs of bacterial type III effector proteins and their roles in infection," *FEMS Microbiol. Rev.*, vol. 35, no. 6, pp. 1100–1125, Nov. 2011, doi: 10.1111/j.1574-6976.2011.00271.x.

[9] E. R. Green and J. Mecsas, "Bacterial Secretion Systems: An Overview," *Microbiol. Spectr.*, vol. 4, no. 1, Feb. 2016, doi: 10.1128/microbiolspec.VMBF-0012-2015.

[10] P. Natale, T. Brüser, and A. J. M. Driessen, "Sec- and Tat-mediated protein secretion across the bacterial cytoplasmic membrane—Distinct translocases and mechanisms," *Biochim. Biophys. Acta BBA - Biomembr.*, vol. 1778, no. 9, pp. 1735–1756, Sep. 2008, doi: 10.1016/j.bbamem.2007.07.015.

[11] J. Sperschneider, P. N. Dodds, D. M. Gardiner, J. M. Manners, K. B. Singh, and J. M. Taylor, "Advances and Challenges in Computational Prediction of Effectors from Plant Pathogenic Fungi," *PLOS Pathog.*, vol. 11, no. 5, p. e1004806, May 2015, doi: 10.1371/journal.ppat.1004806.

[12] G. W. Beakes, S. L. Glockling, and S. Sekimoto, "The evolutionary phylogeny of the oomycete 'fungi,'" *Protoplasma*, vol. 249, no. 1, pp. 3–19, Jan. 2012, doi: 10.1007/s00709-011-0269-2.

[13] M. Thines and S. Kamoun, "Oomycete-plant coevolution: recent advances and future prospects," *Curr. Opin. Plant Biol.*, vol. 13, no. 4, pp. 427–433, Aug. 2010, doi: 10.1016/j.pbi.2010.04.001.

[14] K. J. Wood *et al.*, "Effector prediction and characterization in the oomycete pathogen Bremia lactucae reveal host-recognized WY domain proteins that lack the canonical RXLR motif," *PLOS Pathog.*, vol. 16, no. 10, p. e1009012, Oct. 2020, doi: 10.1371/journal.ppat.1009012.

[15] M. Franceschetti, A. Maqbool, M. J. Jiménez-Dalmaroni, H. G. Pennington, S. Kamoun,
and M. J. Banfield, "Effectors of Filamentous Plant Pathogens: Commonalities amid
Diversity," *Microbiol. Mol. Biol. Rev.*, vol. 81, no. 2, pp. e00066-16, Mar. 2017, doi:
10.1128/MMBR.00066-16.
[16] R. H. Y. Jiang, S. Tripathy, F. Govers, and B. M. Tyler, "RXLR effector reservoir in two
Phytophthora species is dominated by a single rapidly evolving superfamily with more
than 700 members," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 105, no. 12, pp. 4874–4879,
Mar. 2008, doi: 10.1073/pnas.0709303105.
[17] T. A. Torto *et al.*, "EST mining and functional expression assays identify extracellular
effector proteins from the plant pathogen Phytophthora," *Genome Res.*, vol. 13, no. 7,
pp. 1675–1685, Jul. 2003, doi: 10.1101/gr.910003.
[18] L. Pritchard and P. Birch, "A systems biology perspective on plant–microbe interactions:
Biochemical and structural targets of pathogen effectors," *Plant Sci.*, vol. 180, no. 4, pp.
584–603, Apr. 2011, doi: 10.1016/j.plantsci.2010.12.008.
[19] A. H. Lovelace, S. Dorhmi, M. T. Hulin, Y. Li, J. W. Mansfield, and W. Ma, "Effector
Identification in Plant Pathogens," *Phytopathology*, vol. 113, no. 4, pp. 637–650, Apr.
2023, doi: 10.1094/PHYTO-09-22-0337-KD.
[20] J. T. Jones *et al.*, "Top 10 plant-parasitic nematodes in molecular plant pathology," *Mol.
Plant Pathol.*, vol. 14, no. 9, pp. 946–961, Dec. 2013, doi: 10.1111/mpp.12057.
[21] M. Holterman *et al.*, "Disparate gain and loss of parasitic abilities among nematode
lineages," *PLoS One*, vol. 12, no. 9, p. e0185445, 2017, doi:
10.1371/journal.pone.0185445.
[22] C. Vens, M.-N. Rosso, and E. G. J. Danchin, "Identifying discriminative classification-
based motifs in biological sequences," *Bioinformatics*, vol. 27, no. 9, pp. 1231–1238,
May 2011, doi: 10.1093/bioinformatics/btr110.
[23] R. Blanc-Mathieu *et al.*, "Hybridization and polyploidy enable genomic plasticity
without sex in the most devastating plant-parasitic nematodes," *PLoS Genet.*, vol. 13, no.
6, p. e1006777, Jun. 2017, doi: 10.1371/journal.pgen.1006777.
[24] P. Abad *et al.*, "Genome sequence of the metazoan plant-parasitic nematode
Meloidogyne incognita," *Nat. Biotechnol.*, vol. 26, no. 8, Art. no. 8, Aug. 2008, doi:
10.1038/nbt.1482.
[25] M. Da Rocha *et al.*, "Genome Expression Dynamics Reveal the Parasitism Regulatory
Landscape of the Root-Knot Nematode Meloidogyne incognita and a Promoter Motif
Associated with Effector Genes," *Genes*, vol. 12, no. 5, p. 771, May 2021, doi:
10.3390/genes12050771.
[26] R. O. Rocha, R. S. Hussey, L. E. Pepi, P. Azadi, and M. G. Mitchum, "Discovery of
Novel Effector Protein Candidates Produced in the Dorsal Gland of Adult Female Root-
Knot Nematodes," *Mol. Plant-Microbe Interactions*, vol. 36, no. 6, pp. 372–380, Jun.
2023, doi: 10.1094/MPMI-11-22-0232-R.
[27] N. E. Davey, M. S. Cyert, and A. M. Moses, "Short linear motifs – ex nihilo evolution of
protein regulation," *Cell Commun. Signal.*, vol. 13, no. 1, p. 43, Nov. 2015, doi:
10.1186/s12964-015-0120-z.
[28] E. D. O. Roberson, "Motif scraper: a cross-platform, open-source tool for identifying
degenerate nucleotide motif matches in FASTA files," *Bioinforma. Oxf. Engl.*, vol. 34,
no. 22, pp. 3926–3928, Nov. 2018, doi: 10.1093/bioinformatics/bty437.
[29] J. Mistry *et al.*, "Pfam: The protein families database in 2021," *Nucleic Acids Res.*, vol.
49, no. D1, pp. D412–D419, Jan. 2021, doi: 10.1093/nar/gkaa913.
[30] P. Jones *et al.*, "InterProScan 5: genome-scale protein function classification,"
Bioinformatics, vol. 30, no. 9, pp. 1236–1240, May 2014, doi:
10.1093/bioinformatics/btu031.

[31] M. Urban, R. Pant, A. Raghunath, A. G. Irvine, H. Pedro, and K. E. Hammond-Kosack,
"The Pathogen-Host Interactions database (PHI-base): additions and future
developments," *Nucleic Acids Res.*, vol. 43, no. Database issue, pp. D645-655, Jan.
2015, doi: 10.1093/nar/gku1165.
[32] The UniProt Consortium, "UniProt: the Universal Protein Knowledgebase in 2023,"
Nucleic Acids Res., vol. 51, no. D1, pp. D523–D531, Jan. 2023, doi:
10.1093/nar/gkac1052.
[33] B. J. Haas *et al.*, "Genome sequence and analysis of the Irish potato famine pathogen
Phytophthora infestans," *Nature*, vol. 461, no. 7262, Art. no. 7262, Sep. 2009, doi:
10.1038/nature08358.
[34] L. Fu, B. Niu, Z. Zhu, S. Wu, and W. Li, "CD-HIT: accelerated for clustering the next-
generation sequencing data," *Bioinformatics*, vol. 28, no. 23, pp. 3150–3152, Dec. 2012,
doi: 10.1093/bioinformatics/bts565.
[35] D. M. Emms and S. Kelly, "OrthoFinder: phylogenetic orthology inference for
comparative genomics," *Genome Biol.*, vol. 20, no. 1, p. 238, Nov. 2019, doi:
10.1186/s13059-019-1832-y.
[36] T. L. Bailey, "STREME: accurate and versatile sequence motif discovery,"
Bioinformatics, vol. 37, no. 18, pp. 2834–2840, Sep. 2021, doi:
10.1093/bioinformatics/btab203.
[37] C. E. Grant, T. L. Bailey, and W. S. Noble, "FIMO: scanning for occurrences of a given
motif," *Bioinformatics*, vol. 27, no. 7, pp. 1017–1018, Apr. 2011, doi:
10.1093/bioinformatics/btr064.
[38] K. L. Howe *et al.*, "WormBase 2016: expanding to enable helminth genomic research,"
Nucleic Acids Res., vol. 44, no. D1, pp. D774–D780, Jan. 2016, doi:
10.1093/nar/gkv1217.
[39] K. L. Howe, B. J. Bolt, M. Shafie, P. Kersey, and M. Berriman, "WormBase ParaSite –
a comprehensive resource for helminth genomics," *Mol. Biochem. Parasitol.*, vol. 215,
pp. 2–10, Jul. 2017, doi: 10.1016/j.molbiopara.2016.11.005.
[40] J. Martin *et al.*, "Helminth.net: expansions to Nematode.net and an introduction to
Trematode.net," *Nucleic Acids Res.*, vol. 43, no. D1, pp. D698–D706, Jan. 2015, doi:
10.1093/nar/gku1128.
[41] W. Li and A. Godzik, "Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences," *Bioinformatics*, vol. 22, no. 13, pp. 1658–1659, Jul.
2006, doi: 10.1093/bioinformatics/btl158.
[42] P. Grynberg *et al.*, "Comparative Genomics Reveals Novel Target Genes towards
Specific Control of Plant-Parasitic Nematodes," *Genes*, vol. 11, no. 11, p. 1347, Nov.
2020, doi: 10.3390/genes11111347.
[43] F. Pedregosa *et al.*, "Scikit-learn: Machine Learning in Python," *Mach. Learn. PYTHON*.
[44] D. L. Davies and D. W. Bouldin, "A Cluster Separation Measure," *IEEE Trans. Pattern
Anal. Mach. Intell.*, vol. PAMI-1, no. 2, pp. 224–227, Apr. 1979, doi:
10.1109/TPAMI.1979.4766909.
[45] G. E. Crooks, G. Hon, J.-M. Chandonia, and S. E. Brenner, "WebLogo: a sequence logo
generator," *Genome Res.*, vol. 14, no. 6, pp. 1188–1190, Jun. 2004, doi:
10.1101/gr.849004.
[46] H. Nielsen, "Predicting Secretory Proteins with SignalP," in *Protein Function
Prediction: Methods and Protocols*, D. Kihara, Ed., in Methods in Molecular Biology.
New York, NY: Springer, 2017, pp. 59–73. doi: 10.1007/978-1-4939-7015-5_6.
[47] A. Krogh, B. Larsson, G. von Heijne, and E. L. Sonnhammer, "Predicting
transmembrane protein topology with a hidden Markov model: application to complete

genomes," *J. Mol. Biol.*, vol. 305, no. 3, pp. 567–580, Jan. 2001, doi: 10.1006/jmbi.2000.4315.

[48] M.-C. Caillaud and B. Favery, "In Vivo Imaging of Microtubule Organization in Dividing Giant Cell," in *Plant Cell Division: Methods and Protocols*, M.-C. Caillaud, Ed., in Methods in Molecular Biology. New York, NY: Springer, 2016, pp. 137–144. doi: 10.1007/978-1-4939-3142-2_11.

[49] M. Jaouannet, C.-N. Nguyen, M. Quentin, S. Jaubert-Possamai, M.-N. Rosso, and B. Favery, "In situ Hybridization (ISH) in Preparasitic and Parasitic Stages of the Plant-parasitic Nematode Meloidogyne spp," *Bio-Protoc.*, vol. 8, no. 6, p. e2766, Mar. 2018, doi: 10.21769/BioProtoc.2766.

[50] J. Mejias *et al.*, "The root-knot nematode effector MiEFF18 interacts with the plant core spliceosomal protein SmD1 required for giant cell formation," *New Phytol.*, vol. 229, no. 6, pp. 3408–3423, 2021, doi: 10.1111/nph.17089.

[51] L. S. Boutemy *et al.*, "Structures of Phytophthora RXLR effector proteins: a conserved but adaptable fold underpins functional diversity," *J. Biol. Chem.*, vol. 286, no. 41, pp. 35834–35842, Oct. 2011, doi: 10.1074/jbc.M111.262303.

[52] S. Schornack *et al.*, "Ancient class of translocated oomycete effectors targets the host nucleus," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 107, no. 40, pp. 17421–17426, Oct. 2010, doi: 10.1073/pnas.1008491107.

[53] A. P. Rehmany *et al.*, "Differential recognition of highly divergent downy mildew avirulence gene alleles by RPP1 resistance genes from two Arabidopsis lines," *Plant Cell*, vol. 17, no. 6, pp. 1839–1850, Jun. 2005, doi: 10.1105/tpc.105.031807.

[54] T. M. M. M. Amaro, G. J. A. Thilliez, G. B. Motion, and E. Huitema, "A Perspective on CRN Proteins in the Genomics Age: Evolution, Classification, Delivery and Function Revisited," *Front. Plant Sci.*, vol. 8, 2017, Accessed: Jun. 01, 2023. [Online]. Available: https://www.frontiersin.org/articles/10.3389/fpls.2017.00099

[55] E. G. J. Danchin *et al.*, "Multiple lateral gene transfers and duplications have promoted plant parasitism ability in nematodes," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 107, no. 41, pp. 17651–17656, Oct. 2010, doi: 10.1073/pnas.1008486107.

[56] H. Aspeborg, P. M. Coutinho, Y. Wang, H. Brumer, and B. Henrissat, "Evolution, substrate specificity and subfamily classification of glycoside hydrolase family 5 (GH5)," *BMC Evol. Biol.*, vol. 12, no. 1, p. 186, Sep. 2012, doi: 10.1186/1471-2148-12-186.

[57] Z. Han, D. Xiong, R. Schneiter, and C. Tian, "The function of plant PR1 and other members of the CAP protein superfamily in plant–pathogen interactions," *Mol. Plant Pathol.*, vol. 24, no. 6, pp. 651–668, 2023, doi: 10.1111/mpp.13320.

[58] G. M. Gibbs, K. Roelants, and M. K. O'Bryan, "The CAP superfamily: cysteine-rich secretory proteins, antigen 5, and pathogenesis-related 1 proteins--roles in reproduction, cancer, and immune defense," *Endocr. Rev.*, vol. 29, no. 7, pp. 865–897, Dec. 2008, doi: 10.1210/er.2008-0032.

[59] R. Lozano-Durán and S. Robatzek, "14-3-3 Proteins in Plant-Pathogen Interactions," *Mol. Plant-Microbe Interactions*, vol. 28, no. 5, pp. 511–518, May 2015, doi: 10.1094/MPMI-10-14-0322-CR.

[60] K. Wieczorek *et al.*, "A Distinct Role of Pectate Lyases in the Formation of Feeding Structures Induced by Cyst and Root-Knot Nematodes," *Mol. Plant-Microbe Interactions*, vol. 27, no. 9, pp. 901–912, Sep. 2014, doi: 10.1094/MPMI-01-14-0005-R.

[61] T. Hewezi and T. J. Baum, "Manipulation of plant cells by cyst and root-knot nematode effectors," *Mol. Plant-Microbe Interact. MPMI*, vol. 26, no. 1, pp. 9–16, Jan. 2013, doi: 10.1094/MPMI-05-12-0106-FI.

[62] H. Song *et al.*, "The Meloidogyne javanica effector Mj2G02 interferes with jasmonic
acid signalling to suppress cell death and promote parasitism in Arabidopsis," *Mol. Plant*
Pathol., vol. 22, no. 10, pp. 1288–1301, Oct. 2021, doi: 10.1111/mpp.13111.
[63] J. Niu *et al.*, "Msp40 effector of root-knot nematode manipulates plant immunity to
facilitate parasitism," *Sci. Rep.*, vol. 6, no. 1, Art. no. 1, Jan. 2016, doi:
10.1038/srep19443.
860
861

862
863

864 **Figure 1: MOnSTER pipeline scheme.**

865 (A) MOnSTER pipeline is composed of three steps. It takes two FASTA protein sequences datasets
866 (positive and negative) and a list of predicted motifs (enriched in the positive dataset) as input. The
867 output is a list of CLUMPs and an associated MOnSTER score. The MOnSTER score is constituted
868 by: (B) CLUMP$_{score}$ calculation. (C) Two occurrences Indexes.

| RxLR | dEER | LxLFLAK | HVLVxxP |

Figure 2: Motif logos of CLUMPs compared to the target motifs.
Upper-panel: alignments of motifs in the respective CLUMP are produced by PROMOCA, and then the aligned motif sequences are used to produce the logos with WebLogo3. The x-axis represents the AA position in the motif, while the y-axis represents log-transformed frequencies translated into bits of information. Lower-panel: characteristic motifs of oomycetes effectors families from literature.

885
886
Figure 3: Dendrogram of CLUMPs in Plant Parasitic Nematodes (PPNs)
11 CLUMPs produced by MOnSTER (indicated with "/" sign). The coloured ones are those selected as
best-scoring CLUMPs after MOnSTER-score calculation. Each best-scoring CLUMP is associated with
the corresponding motif logo; alignment of motifs in each CLUMP is produced by PROMOCA and then
WebLogo 3 is used to produce the image (the x-axis shows the AA position of the motif and the y-axis
represents the log-transformed frequency of each AA in terms of bits of information).
893

CLUMP/s	dc1	dc2	dc3	dc4	dc5
2	.	.	.	.	.
1	13	.	.	.	.
7	.	2	.	.	.
7,10	.	23	.	.	.
2,5	.	.	8	.	.
3	.	.	1	.	.
2,3	.	.	3	16	.
5	.	.	.	.	.
3,7,10	.	.	.	.	17
10	.	1	.	.	.

901
Figure 5: Candidate parasitism proteins showing the presence of CLUMP/s associated with pathogenicity-related protein domain/s.
The table on the right shows the co-occurrence of CLUMP or CLUMPs with specific domain classes (dc); dc1, pectate lyase domain class, dc2, glycosyl hydrolase family 5 domain class, dc3 Stichodactyla toxin (ShK) domain class, dc4 14-3-3 family domain class and dc5, cysteine-rich domain class. The upset plot on the left represents the occurrences and co-occurrences of respective CLUMPs in the positive dataset, highlighting the sequences that also have an interesting protein domain following the table counts.
910

Figure 6: MiEFF72 is specifically expressed in the subventral glands.

(A) *In situ* hybridisation showing EFF72 transcripts in the subventral glands (SvG) of J2s of *M. incognita* (two left pictures). Sense probe for the MiEFF72 transcripts was used as a negative control (right picture). (B) MiEFF72 localised to the cytoplasm of plant cells and in cytoplasmic vesicles (red arrows). The MiEFF72 sequence was fused to C-terminal end of the GFP and expressed in *N. benthamiana* leaves by agroinfiltration. Bars = 20 µm.

918
919

Supporting information files legend

Supplementary figure 1: PRO-MOCA algorithm.
(a) DEFINITION OF THE ALPHABET. The user defines the alphabet to use to translate the protein sequences into patterns. Alphabet are constructed as pairs of keys, a letter defined by the user, and values, a list of amino acids to include in the group. Amino acids from the group will be replaced by the key in the sequence. PRO-MOCA already include the following alphabets. Chemical alphabet: 'P' = 'polar', 'H' = hydrophobic, 'N'= 'neutral', 'A' = 'acid', 'B' = 'basic'. Hydrophobicity alphabet: 'P' = 'hydrophilic', 'N' = 'neutral', 'B' = 'hydrophobic'. Charge alphabet: 'P' = 'positive', 'N' = 'negative', 'U' = 'neutral'. Secondary structure propensity alphabet: 'H' = 'helix (α)', 'S' = 'sheet (β)', 'T' = 'turn'.
(b) TRANSLATION. Using the chosen alphabet (step (a)), the motifs in the CLUMP are translated from the protein alphabet (amino acids) into patterns of the given alphabet. For instance, using the 'chemical' alphabet, the 'GHWT' motif is translated into 'PBHP'.
(c) ALIGNMENT. (c1) PRO-MOCA: (1) takes the list of patterns, (2) selects the first position of each pattern, (3) calculates how many times each letter is repeated at that position and finds the most repeated letter (PRO-MOCA stores this result at each iteration) (4) After that, patterns with the most repeated letter at the first position, lose that letter, and the rest of the pattern is shifted of -1 position. (5) The result is a list of patterns where part of the patterns has a new first position. From this point, the process (3)-(5) is repeated until (6) the result is a list where at least one pattern does not have any more letters. Until there is a pattern with no more letters (golden rectangle). (c2) Once the first part of the alignment is done and PRO-MOCA gets to (6), it finds the central position of the alignment. To do that, PRO-MOCA stored in (3) the number of repetitions of the most repeated letter. In (c2) PRO-MOCA calculates which is the highest value, and gets the corresponding list of patterns with those first positions.
(d) **GAP INSERTION**. The gap insertion is only allowed at the extremities of the patterns and allows to have an alignment where all the patterns have the same length. To do that, PRO-MOCA calculates, for each pattern, how many letters there are before (x_i) and after (y_i) the central position and calculates the maximum (m and n) of these values. After that, it subtracts $m - x_i$ and $n - y_i$. The result of the subtraction is the number of gaps each pattern has before and after the central position.
(e) RE-TRANSLATION. In this step, PRO-MOCA re-translates the aligned patterns into aligned motifs

Supplementary figure 2: Dendrogram of CLUMPs in Oomycetes.
11 CLUMPs produced by MOnSTER (indicated with "/" sign). The coloured ones are those selected as best-scoring CLUMPs after MOnSTER-score calculation. Each best-scoring CLUMP is associated the motif logo; the alignment of the motif in each CLUMP is produced by PROMOCA and then WebLogo 3 is used to produce the image (the x-axis shows the AA position of the motif and the y-axis represents the log-transformed frequency of each AA in terms of bits of information).

Supplementary figure 3: Occurrences and co-occurrences of CLUMPs in positive and negative dataset – Oomycetes.
Each upset-plot represents the number of sequences in the respective dataset according to the presence of one or more CLUMPs ordered by MOnSTER-score. **(a)** positive dataset, **(b)** negative dataset, (c-d) specifically show the occurrence and co-occurrence of CLUMPs corresponding to RxLR and dEER (CLUMP10, 6, 2), and LxLFLAK-HVLVxxP (CLUMP7,4), in RxLR and Crinkler-effector sequences, respectively.

Supplementary figure 4: Sequence position preference of motifs in CLUMPs – Oomycetes.
(a) Shows the overall preference position of CLUMPs in the positive (blu trace) compared to the negative (orange trace) dataset. In the rug-plot each line corresponds to a CLUMP-motif position occurrence. **(b)** represents the same distribution in the positive dataset but divided by each CLUMP abundance in the sequence bin. **(c)** shows the same concept but in the negative dataset; **(c.1)** is a

zoom-in of the original plot to better visualize the occurrences of CLUMPs. The distributions are all represented in bins of 10% each as absolute positions on the sequence.

Supplementary figure 5: Sequence position preference of motifs in CLUMPs – PPNs.
(a) Shows the overall preference of CLUMPs in the positive (blu trace) compared to the negative (orange trace) dataset. In the rug-plot each line corresponds to a CLUMP-motif position occurrence. **(b)** represents the same distribution in the positive dataset but divided by each CLUMP abundance in the sequence bin. **(c)** shows the same concept but in the negative dataset; **(c.1)** is a zoom-in of the original plot to better visualize the occurrences of CLUMPs. The distributions are all represented in bins of 10% each as absolute positions on the sequence.

Supplementary figure 6: Occurrence and co-occurrence of CLUMPs in PPNs.
Representation of occurrence and co-occurrence of each best-scoring CLUMP in protein sequences, according to their MOnSTER-score. (a) shows the number of sequences and the occurrence of CLUMPs in the positive dataset, while (b) represents the same measures in the negative dataset.

Supplementary figure 7. Occurrence and co-occurrence of CLUMPs in proteins with or without the signal peptide.
Representation of occurrence and co-occurrence of each best-scoring CLUMP in protein sequences, accordingly to their MOnSTER-score. (a) shows the number of sequences and the occurrence of CLUMPs in the sequences bearing the signal peptide, while (b) represents the same measures in the sequences without the signal peptide.

Supplementary figure 8: Relation between CLUMPs abundance and protein sequence length – PPNs.
(a) Shows a non-linear relationship between the occurrences of CLUMPs and the length of sequences in the positive dataset, while **(b)** represents the same but in the negative dataset.

Supplementary figure 9: Protein sequence position of motifs in CLUMP7 and 10 associated with glycosyl hydrolase family 5 domain (dc2).
(a) CLUMP7 and CLUMP10 motifs-position and respective occurrence along selected protein sequences divided into bins representing 10% sequence portion. (**b**) rug-plot representing the start and end position of the domains (PS00659-ProSitePatterns and PF00150-Pfam) belonging to the class 2 of domains, namely glycosyl hydrolase family 5 domains.

Supplementary table 1.1: Oomycetes' positive dataset.
Detailed description of the protein sequences in the oomycetes' positive dataset including species, retrieving sources, and corresponding amino acid sequence.

Supplementary table 1.2: Oomycetes' negative dataset.
Detailed description of the protein sequences in the oomycetes' negative dataset including species, retrieving sources, and corresponding amino acid sequence.

Supplementary table 2.1: PPNs' positive dataset.
Detailed description of the protein sequences in the PPNs' positive dataset including species, retrieving sources, and corresponding amino acid sequence.

Supplementary table 2.2: PPNs' negative dataset.
Detailed description of the protein sequences in the PPNs' negative dataset including species, retrieving sources, and corresponding amino acid sequence.

Supplementary table 3: Oomycetes' MOnSTER-score.

1026 Summary table of MOnSTER-score for each CLUMP found in oomycetes dataset, ranked by higher
1027 score.
1028
1029 **Supplementary table 4: PPNs' MOnSTER-score.**
1030 Summary table of MOnSTER-score for each CLUMP found in PPNs dataset, ranked by higher score.
1031
1032 **Supplementary table 5: InterProScan domains and CLUMPs co-occurrences.**
1033 Summary of co-occurrences of InterProScan domains in the six characteristic CLUMPs for PPN
1034 species.
1035
1036 **Supplementary table 6: *Meloidogyne incognita* proteins for validation of putative effector.**
1037 List of proteins in *M. incognita* proteome presenting at least one motif belonging to CLUMP5 (the most
1038 abundant in the proteome) and predicted to possess signal peptide and no transmembrane domain.
1039 Further details on the selection criteria can be found in the "Results & Discussion" section of the main
1040 text.
1041
1042
1043
1044

31