RNA-directed DNA Methylation is required for seed development in outbreeding species

Kelly J. Dew-Budd[1], Hiu Tung Chow[1], Timmy Kendall[1], Brandon C. David[1], James A. Rozelle[1], Rebecca A. Mosher[1], and Mark A. Beilstein[1]

[1] School of Plant Sciences, University of Arizona, Tucson, Arizona.

Corresponding authors: RAM (rmosher@arizona.edu) and MAB (mbeilstein@arizona.edu)

Abstract

In plants, *de novo* DNA methylation is guided by 24-nt short interfering (si)RNAs in a process called RNA-directed DNA methylation (RdDM). Primarily targeted at transposons, RdDM causes transcriptional silencing and can indirectly influence expression of neighboring genes. During reproduction, a small number of siRNA loci are dramatically upregulation in the maternally-derived seed coat, suggesting that RdDM might have a special function during reproduction. However, the developmental consequence of RdDM has been difficult to dissect because disruption of RdDM does not result in overt phenotypes in *Arabidopsis thaliana*, where the pathway has been most thoroughly studied. In contrast, *Brassica rapa* mutants lacking RdDM have a severe seed production defect, which is determined by the maternal sporophytic genotype. To explore the factors that underlie the different phenotypes of these closely-related species, we produced RdDM mutations in three additional members of the Brassicaceae family: *Camelina sativa*, *Capsella rubella*, and *Capsella grandiflora*. Among these three species, only mutations in *Capsella grandiflora* displayed a seed production defect similar to *Brassica rapa* mutants, suggesting that mating system is a key determinant for reproductive phenotypes in RdDM mutants.

Introduction

RNA-directed DNA methylation (RdDM) triggers de novo DNA methylation and maintains non-symmetric methylation at euchromatic transposons (Matzke and Mosher, 2014; Erdmann and Picard, 2020). This methylation causes transcriptional silencing to prevent the mobilization of transposons and can indirectly influence expression of neighboring genes (Hollister and Gaut, 2009; Hollister et al., 2011; Forestan et al., 2017). RdDM employs 24-nt small interfering (si)RNAs, which are produced though the sequential activities of RNA Polymerase IV (Pol IV), RNA-DEPENDENT RNA POLYMERASE2 (RDR2), and DICER-LIKE3 (Singh et al., 2019; Fukudome et al., 2021; Huang et al., 2021; Wang et al., 2021; Loffer et al., 2022). These siRNAs are loaded into ARGONAUTE4 and direct the protein to nascent transcripts produced by RNA Polymerase V (Wierzbicki et al., 2009; Liu et al., 2018; Singh et al., 2019; Wang et al., 2023). Once assembled at the chromatin, ARGONAUTE4 recruits DOMAINS REARRANGED METHYLTRANSFERASE to methylate DNA proximal to the Pol V transcript (Zemach et al., 2013; Stroud et al., 2014; Zhong et al., 2014). Because non-symmetric DNA methylation is not

maintained on both daughter strands during DNA replication, repeated *de novo* methylation is necessary to maintain DNA methylation (Law and Jacobsen, 2010; Law et al., 2013).

Twenty-four nt siRNAs are particularly abundant in reproductive tissues, suggesting that RdDM has a special role during reproduction (Chow and Mosher, 2023). Increased accumulation of 24-nt siRNAs is due to expression of reproductive-specific loci, rather than upregulation of all RdDM sites (Mosher et al., 2009; Rodrigues et al., 2013; Grover et al., 2020; Li et al., 2020; Long et al., 2021; Zhou et al., 2022). In particular, ovule- and endosperm-specific siRNA loci have been described in rice (Rodrigues et al., 2013), *Brassica rapa* (Grover et al., 2020), and Arabidopsis (Zhou et al., 2022). Known as siren loci, these siRNAs comprise approximately 90% of the siRNA population in unfertilized ovules and trigger methylation of protein-coding genes during seed development (Grover et al., 2020; Burgess et al., 2022). In some cases, siren-induced *trans*-methylation impacts gene expression (Burgess et al., 2022).

The developmental consequence of RdDM action during reproduction has been difficult to dissect because in *Arabidopsis thaliana*, where the pathway has been most thoroughly studied, disruption of RdDM does not result in overt phenotypes. There is a slight reduction in seed weight for homozygous mutant seed (Grover et al., 2018) and an increase in the success of paternal-excess crosses when RdDM is eliminated from the maternal parent (Erdmann et al., 2017; Martinez et al., 2018; Satyaki and Gehring, 2019). In contrast, *Brassica rapa* RdDM mutants are almost completely sterile and produce seeds that are smaller than wild type (Grover et al., 2018). The different phenotypes for these relatively closely-related species raises the question: what underlies the phenotypic differences upon loss of RdDM? It has been proposed that RdDM might play a larger developmental role in species with a higher proportion of transposons, or in genomes where transposons are in closer proximity to protein-coding genes, as these factors increase the possibility that transcriptional silencing of a transposon will have an indirect impact on a protein-coding gene (Wei et al., 2014; Gouil and Baulcombe, 2016; Wang et al., 2020). We propose two additional hypotheses to explain the different phenotypes of *A. thaliana* and *B. rapa* RdDM mutants.

Firstly, RdDM might play a role in mediating conflicts among the subgenomes of polyploids since *A. thaliana* is diploid while *B. rapa* is a recent allohexaploid (Wang et al., 2011). A common result of allopolyploidization is genome dominance, where the non-dominant subgenome experiences reduced expression and increased fractionation (Grover et al., 2012; Garsmeur et al., 2014). In *B. rapa*, the homeolog with more transposons within 500 bp of the transcription start or stop site was more likely to have lower expression and be present in the non-dominant subgenome (Woodhouse et al., 2014), suggesting that RdDM is associated with reduced expression from the non-dominant genome. This idea is supported by the higher proportion of transposon-derived 24-nt siRNAs mapping to regions surrounding genes of the non-dominant genome in *B. rapa*, cotton, and maize (Woodhouse et al., 2014; Renny-Byfield et al., 2015; Cheng et al., 2016; Forestan et al., 2017). Consequently, RdDM might be required to maintain the balance among subgenomes during reproduction, and increased expression of genes on the non-dominant subgenome in RdDM mutants might cause dosage imbalances that result in detrimental phenotypes (Alger and Edger, 2020). Species without a recent history of genome duplication have fewer homeologous gene pairs and less need for subgenome balance and thus might have fewer phenotypes when RdDM is eliminated.

Alternatively, RdDM's role during reproduction might relate to balancing parental genomes in the endosperm, a process linked to RdDM (reviewed in (Chow and Mosher, 2023)). The triploid endosperm develops following the fertilization of the diploid central cell by a haploid sperm cell and therefore has a 2:1 maternal:paternal genome ratio. Distortion of this ratio results in failure during endosperm development, likely due to an imbalance in parental conflict over resource allocation to seeds (Scott et al., 1998; Haig, 2013). Similar imbalances are seen in interspecific crosses of the same ploidy, leading to the conclusion that it is not the *absolute* ploidy from each parent that determines balance, but rather the *effective* ploidy – a combination of ploidy and the strength of parental conflict (Johnston et al., 1980). The WISO hypothesis (Weak Inbreeder, Strong Outbreeder) proposes that parental conflict is proportional to the extent of outbreeding in a species (Brandvain and Haig, 2005; Brandvain and Haig, 2018). Until recent domestication, *B. rapa* is an outbreeding species, implying it will have strong conflict, while *A. thaliana* evolved inbreeding ~ 500 thousand years ago (KYA) and should have weak conflict (de la Chaux et al., 2012). RdDM may be a necessary referee to mediate conflict between strong maternal and paternal interests in outbreeders, but loss of that balance is less important in inbreeders who have only weak conflict.

To test these hypotheses, we used a CRISPR/Cas9 DNA editing system to introduce mutations into two essential genes in the RdDM pathway in three Brassicaceae species that differed from each other in ploidy or breeding system. Our target species were: 1) *Camelina sativa,* a self-compatible recent allopolyploid; 2) the diploid obligate outcrosser *Capsella grandiflora*; and 3) *Capsella rubella*, a diploid inbreeder (**Supplementary Figure 1**). *C. sativa* underwent a polyploidization event ~ 5.5 million years ago (MYA), retains three distinct subgenomes, and has a clear subgenomic expression bias (Kagale et al., 2014; Brock et al., 2018; Mandáková et al., 2019). We reasoned that if RdDM is critical in mediating conflict among subgenomes, *C. sativa* RdDM mutants would have severely reduced seed set similar to *B. rapa*. *C. rubella* split from its congener, *C. grandiflora*, < 50-200 KYA, soon after its founders lost self-incompatibility (Foxe et al., 2009; Guo et al., 2009; Slotte et al., 2013) yielding a fully self-compatible *C. rubella*. The sister relationship between *C. grandiflora* and *C. rubella* allowed us to test whether outcrossers exhibit a stronger requirement for RdDM during reproduction than do inbreeders. Here, we show that RdDM is not critical for seed production in *C. sativa*, suggesting that a recent history of polyploidy is not an important factor for RdDM mutant phenotypes. Instead, we observed a significant and striking reduction in seed set in *C. grandiflora*, suggesting that breeding system is a major factor determining reproductive impacts upon loss of RdDM.

Results

Creation of RdDM mutants in Brassicaceae species

To explore the role of RdDM during reproduction, we used CRISPR/Cas9-nickase (Fauser et al., 2014) to generate loss of function mutations in *RDR2* and *NRPE1* in three Brassicaceae species that vary in breeding system and history of genome duplication. *NRPE1* encodes the largest subunit of RNA Pol V (Ream et al., 2009). *C. rubella* and *C. grandiflora* encode single copies of *RDR2* and *NRPE1*, while *C. sativa* retains three copies of these genes due to its

Figure 1: Creation and characterization of RdDM mutants in Brassicaceae species
(**A**) Schematic diagrams of *Capsella NRPE1* and *RDR2*. *Camelina sativa* has the same number and size of exons, but intron length varies. Cas9 guide RNAs are depicted as orange arrows. (**B**) *Camelina* and *Capsella* RdDM mutants have no or mild vegetative growth phenotypes. (**C**) Size profiles of genome-mapping small RNAs from wild-type and RdDM mutant leaves of each species. Each biological replicate is shown as a separate bar. (**D**) The largest class of leaf siRNA loci in each species are 24-nt dominant loci. (**E**) In each species, *rdr2* mutations dramatically reduce expression from 80-90% of siRNA loci, while *nrpe1* mutations have a less pronounced effect on siRNA expression.

recent whole genome triplication. We identified guide RNAs that flank functionally important regions of the proteins and target all five versions of each gene (**Figure 1A**). Following floral-dip transformation of the genome editing constructs, T1 individuals were genotyped to identify strong mutations in target genes and these mutations were bred to homozygosity (**Supplementary Table 1, Figure S2**). For *C. sativa*, mutations in all three gene copies were combined and the resulting triple mutants are hereafter designated *Cs rdr2-1*, *Cs rdr2-2*, *Cs nrpe1-1*, and *Cs nrpe1-2*. Similar to *rdr2* and *nrpe1* mutations in *A. thaliana* and *B. rapa*, the *C. sativa* and *C. grandiflora* mutants did not exhibit consistent vegetative growth defects, however all RdDM mutants in *C. rubella* were moderately smaller than wild type (**Figure 1B**) (Grover et al., 2018).

To confirm whether these mutations impact RdDM, we sequenced small RNAs from leaf tissue of each allele and mapped them to their respective genomes (**Supplementary Table 2**). To increase the proportion of mapped *C. grandiflora* sRNA reads, we used Oxford Nanopore Technology to sequence the genome of *C. grandiflora* accession 83.17, and then *de novo* assembled the reads, polished the assembly with publicly-available Illumina reads (Williamson et al., 2014), and annotated it. Our resequencing efforts yielded a *C. grandiflora* genome that is less fragmented compared to the previously published genome, with an increase in linearity to L/N50 of 17/2.3 Mb from 256/29.1 kb (**Supplementary Table 3**). The size profile of mapped sRNA reads demonstrates that there is strong loss of 23-24-nt siRNA in the *rdr2* lines of all three species, and a smaller reduction in 23-24-nt accumulation in *nrpe1* plants (**Figure 1C**). A partial loss of siRNA accumulation in *nrpe1* is expected, as Pol V is not required for siRNA biosynthesis, but the failure to target methylation in *nrpe1* mutants results in reduced production of siRNAs via Pol IV (Kanno et al., 2005; Pontier et al., 2005; Mosher et al., 2008; Gouil and Baulcombe, 2016; Grover et al., 2018; Zheng et al., 2021). There is also an observable increase in all other size classes in the *rdr2* mutants, likely due to over-sampling these species following loss of the major size class.

Next, we used ShortStack to define small (s)RNA-producing loci in each genome. The *Capsella* species have 30-40,000 sRNA loci, while *C. sativa* has approximately 3 times that number, perhaps due to its recent whole genome triplication (**Figure 1D**). Categorizing these loci by the dominant size of sRNA demonstrates that 24-nt sRNA loci form the largest group (61-68%). We then assessed the accumulation of sRNA from each locus in the mutants. In each species, *rdr2* mutant exhibit a strong reduction in sRNA from 80-90% of sRNA loci and increased sRNA accumulation from a small proportion of loci (**Figure 1E**). In contrast, fewer sRNA loci were strongly reduced in *nrpe1* mutants, and many loci displayed only a moderate reduction in sRNA accumulation. This pattern is consistent with sRNA size profiles and sRNA locus analysis of similar mutations in *A. thaliana*, tomato, *B. rapa*, and rice (Mosher et al., 2008; Gouil and Baulcombe, 2016; Grover et al., 2018; Zheng et al., 2021), indicating that the mutations generated here disrupt the RdDM pathway.

Mating system determines seed production phenotype upon loss of RdDM

To investigate reproductive phenotypes upon loss of RdDM, we first assessed seed set, defined here as the number of healthy seeds per fruit. Because *C. sativa* fruits do not lose their valves (shatter) at maturity, we counted the number of healthy seeds in mature fruits. Seeds

that were shriveled or darkly pigmented were considered inviable (**Supplementary Figure 3A**). Fruits in both *Capsella* species shatter, particularly when they contain many mature seeds. For these species we therefore counted developing seeds at 7 days after pollination (7 DAP), a timepoint that allows us to measure seed development before shattering. Seeds in these fruits were considered healthy if they were plump and green; white, brown, and/or unexpanded seeds were counted as aborted or unfertilized (**Supplementary Figure 3B, C**).

RdDM mutant lines in all three species exhibited reduced seed set (**Figure 2A**), however the scale of reduction varied. *Cs rdr2* plants had 25-32% reduction in seed set compared with wild type, while *Cs nrpe1-1* showed no change in seed set. RdDM mutations in *C. rubella* reduced

Figure 2. Seed production in Brassicaceae RdDM mutants
(**A**) Histograms of the number of mature seeds per fruit in *Camelina* or healthy seeds at 7 DAP in *Capsella*. The average seed count is shown with a black dot; black bars represent the standard deviation around the mean. (**B**) Seed weight of mature seeds. Pools of 20 seeds were weighed three times and the average of each pool was divided by the number of seeds to obtain an average seed weight. The average and standard deviation of these seed weights for n pools are shown. * = p<0.01 after Bonferroni FDR correction; n.s. = not significant.

healthy seeds at 7 DAP by approximately half. In comparison, *C. grandiflora rdr2* alleles had 97-99% fewer healthy seeds in the RdDM mutant fruits compared to wild type, while *Cg nrpe1* mutants showed approximately 70% reduction. We also measured the weight of mature seeds collected in bulk, which demonstrated that even seeds scored as healthy are slightly smaller for RdDM mutants (**Figure 2B**). Taken together with observations in *A. thaliana* and *B. rapa*, these results indicate that the RdDM pathway plays a role during seed development in each of the species tested, with the strongest effect in outbreeding *C. grandiflora*.

<u>Maternal factors are responsible for seed set reductions in *C. rubella* and *C. grandiflora*</u>

To better understand the role of RdDM in species with different mating systems, we further compared the reproductive phenotypes of RdDM mutants in the two *Capsella* species. Wang et al. (2020) reported defective pollen in *C. rubella nrpd1* mutants, with approximately 80% of the pollen arresting prior to maturity (Wang et al., 2020). We therefore imaged mature pollen grains from each *C. rubella* and *C. grandiflora* RdDM mutant line to determine if reduction in seed set resulted from defects in pollen maturation. We observed no difference in the proportion of mature pollen compared to wild type (**Figure 3A, B**). Consistent with this observation, pollination with wild-type pollen was unable to rescue the seed production defect in any of the *Capsella* RdDM mutant lines (**Figure 4A**). These observations indicate that decreased seed production in *Capsella* RdDM mutants is primarily due to maternal factors, a feature that is shared with seed production phenotypes of *B. rapa* RdDM mutants (Grover et al., 2018).

Figure 3: Normal pollen development in *Capsella* RdDM mutants
(**A**) Representative images of pollen from *Capsella* RdDM mutants demonstrates that most pollen is mature and trinucleate, with a single diffuse vegetative nucleus and two condensed sperm cells (top, DAPI; bottom, brightfield). (**B**) Quantification of mature (trinucleate) pollen and the number of assessed pollen grains.

Reduction in seed set might result from fewer ovules, decreased fertilization, or increased seed abortion. To distinguish among these possibilities, we analyzed the number of ovules in unfertilized pistils from each mutant (**Figure 4B**). In *C. grandiflora*, there is no statistically significant difference in ovule number between wild type and RdDM mutants, however *C. rubella* RdDM mutants exhibit approximately 25% fewer ovules than wild type ($p < 0.01$, Bonferonni corrected Student's T test), a reduction that accounts for most of the reduced seed set in these mutants. The higher number of ovules in *C. rubella* wild type relative to *C. grandiflora* wild type is consistent with the decrease in pollen:ovule ratio commonly observed following the transition to self-fertilization (Sicard and Lenhard, 2011). To further explore the seed set reduction in *Capsella* RdDM mutants, we counted the number of inviable seeds in 7 DAP fruits and expressed this as a fraction of all objects in a fruit to account for differences in ovule number (**Figure 4C**). Inviable seeds included those that had clearly begun development before aborting as well as small white seeds that might be unfertilized ovules or result from abortion shortly after fertilization (**Supplemental Figure 3**). While *C. rubella* plants displayed relatively high levels of seed failure, this was independent of genotype. In contrast, *C. grandiflora rdr2* and *nrpe1* fruits exhibited substantially higher levels of seed failure than wild type (**Figure 4C**), including 14-56% clearly resulting from seed abortion. From this we conclude that reduction in seed set in *C. rubella* RdDM mutants is primarily due to the production of fewer ovules before fertilization, while abortion after fertilization is the major contributor to reduced seed set observed in *C. grandiflora* mutants.

B. rapa RdDM mutants also exhibit extensive seed abortion after fertilization, a trait that is controlled by the maternal sporophytic genotype (Grover et al., 2018). To test whether seed abortion is sporophytically determined in *C. grandiflora*, we crossed heterozygous plants with wild-type or homozygous mutant pollen and scored seed development at 7 DAP. Seed production in such crosses is indistinguishable from wild type (**Figure 4D**), indicating that mutant embryo sacs develop normally even when fertilized by mutant pollen, as long as they are supported by a maternal sporophyte capable of RdDM.

Discussion

We sought to test two hypotheses to explain the different seed production phenotypes of RdDM mutants in Brassicaceae. First, that failure of seed development in RdDM mutants was associated with recent polyploidy and ongoing balance between subgenomes, and secondly, that breeding system and the increased parental conflict expected in outbreeding species results in a requirement for RdDM. Here, we show that RdDM is critical for seed development in outbreeding species while its role is less important in inbreeding species, including inbreeding polyploids.

Seed production was disrupted when RdDM was eliminated in the three Brassicaceae species, but to a different extent in each (**Figure 2**). RdDM mutants in polyploid *C. sativa* had only a very slight decrease in seed set. RdDM might be important for maintaining balance among its three subgenomes, however loss of this balance does not confer a reproductive phenotype. Similarly, although loss of RdDM causes a reduced size of rosette and fewer ovules in the diploid inbreeder *C. rubella* (**Figure 1B, 4B**), upon fertilization, those ovules produce healthy seeds at the same frequency as wild type plants (**Figure 4C**). In contrast, *C. grandiflora*

RdDM mutants had a strong post-fertilization defect that was dependent on the maternal sporophytic genotype; this defect showed similar characteristics and genetic control as RdDM mutants in the recently outbreeding *B. rapa* (Grover et al., 2018).

Figure 4: Seed set reduction in *Capsella* RdDM mutants is controlled maternally
(**A**) Number of "healthy" seeds per fruit at 7 DAP in *Capsella* RdDM mutants crossed with wild-type pollen. WT x WT are the same data presented in Figure 2A. (**B**) Number of ovules in unfertilized pistils (n > 20). Colored bars represent the average; black dots are individual datapoints. All *C. rubella* RdDM mutants are significantly different from wild type (WT) at p < 0.01 after Bonferroni correction (Student's T test). *C. grandiflora* mutants are not significantly different at this threshold. (C) Fraction of things inside a fruit at 7 DAP that are not scored as "healthy", representing either failure to fertilize or seed abortion. (**D**) Number of "healthy" seeds per fruit at 7 DAP in *C. grandiflora rdr2* heterozygotes crossed with homozygous mutant or wild-type pollen. *Cg rdr2-1* x *Cg rdr2-1* are the same data presented in Figure 2A.

A maternal sporophytic defect impacting seed number and size in *C. rubella nrpd1* mutants has also been reported (Wang et al., 2020), however it is not clear whether this results from a decreased number of ovules (as we detect for *Cr rdr2* and *Cr nrpe1*) or another mechanism. Instead, the authors focused on a reported defect in pollen development. We do not detect a pollen defect in either *C. rubella* or *C. grandiflora* RdDM mutants (**Figure 3**), consistent with the fact that wild-type pollen is unable to rescue seed production from *Cg rdr2* or *Cg nrpe1* mutants (**Figure 4**). This discrepancy in pollen phenotype might be due to the mutation of different siRNA biosynthesis genes (*NRPD1* versus *RDR2*), however these proteins function together to generate dsRNA and would be expected to yield highly similar, if not identical, phenotypes (Singh et al., 2019; Fukudome et al., 2021; Huang et al., 2021). Variation in pollen phenotypes might also result from the use of different *C. rubella* accessions or different growth conditions.

The requirement for maternal sporophytic RdDM for successful seed production in *C. grandiflora* implicates maternally-derived siRNAs. The most abundant of these are siren siRNAs, which are expressed in the developing seed coat and might be trafficked into the developing endosperm (Grover et al., 2020). Siren siRNAs trigger non-CG methylation of protein-coding genes, which can result in altered gene expression (Burgess et al., 2022). Further research is needed to determine whether ovule siRNA populations and levels of methylation differ between *C. rubella* and *C. grandiflora*, or whether specific sRNA loci are required for seed development in outbreeding species.

Loss of RdDM has been linked to reduction of effective ploidy in diploid by tetraploid crosses in *A. thaliana* (Erdmann et al., 2017; Martinez et al., 2018; Satyaki and Gehring, 2019). Interspecific crosses indicate that *C. grandiflora* has a higher effective ploidy than *C. rubella*, which is consistent with the WISO hypothesis and indicative of stronger parental conflict in *C. grandiflora* (Brandvain and Haig, 2005; Rebernig et al., 2015; Brandvain and Haig, 2018; Lafon-Placette et al., 2018). We therefore hypothesize that loss of RdDM reduces effective ploidy and disrupts the balance between parental genomes after fertilization. For outbreeding species with high effective ploidy, this disruption results in a greater developmental defect because parental conflict is high. Inbreeding species have limited parental conflict and therefore disruption of the maternal:paternal ratio has less impact.

Transposable element content, and the proximity of transposons to protein-coding genes, has also been proposed to explain variation in phenotypes upon loss of RdDM in different species (Wei et al., 2014; Gouil and Baulcombe, 2016; Wang et al., 2020). However, *C. rubella* and *C. grandiflora* genomes have similar transposon number, expression, and proximity to genes (Slotte et al., 2013; Agren et al., 2014), and both *Capsella* genomes have a much lower transposon content than *B. rapa* (Wang et al., 2011). Therefore, transposon density cannot account for the dramatic difference in reproductive phenotype when RdDM is eliminated, although we cannot eliminate the possibility that it is linked to the likelihood of vegetative phenotypes.

There are a suite of floral development changes that commonly occur following the transition to self-compatibility, together referred to as the selfing syndrome (Sicard and Lenhard, 2011). Reduction in parental conflict, effective ploidy, and the requirement for RdDM could therefore be considered an epigenomic selfing syndrome – a suite of epigenomic changes that commonly develop over evolutionary time following the transition to self-fertilization. Our work provides the foundation for further study of this phenomenon.

Methods

<u>Plant material and growth conditions</u>

C. sativa cultivar Ames (a gift of John McKay, Colorado State University), *C. rubella* Monte Gargano, and *C. grandiflora* accession 83.17 (both donated by Stephen Wright, University of Toronto) were grown at 20°C under 16h:8h light:dark conditions. *C. sativa* seeds were stratified for one day at 4˚C prior to germination; *C. rubella* and *C. grandiflora* were both stratified for two weeks at 4˚C. *C. rubella* also required three weeks of vernalization at 4˚C to induce flowering. For the collection of unfertilized ovules, *C. rubella* flowers were emasculated approximately 24 hours before anthesis; pistils were manually dissected 2 days later. *C. grandiflora* ovules were collected identically to *C. rubella* except emasculation was not performed.

<u>CRISPR design and plasmid construction</u>

The T-DNA cassette used for transformation contained Cas9 nickase under a ribosomal protein promoter (*AtRPS5A::Cas9-D10A*), two guide RNAs, EGFP-tagged oleosin under its native promoter (*AtOLE::OLE-GFP*), and a BASTA resistance gene. The vector was constructed using traditional cloning techniques in *E. coli* to combine desired promoters, reporters, and proteins into a single vector with a variety of source materials. Assembled T-DNA plasmids were purified and transformed into *Agrobacterium tumefaciens* GV3101.

Two protospacers for both *RDR2* and *NRPE1* were selected manually from regions conserved in all three species with target specificity verified using CRISPOR (Haeussler et al., 2016) and CRISPRdirect (Naito et al., 2015). The protospacers for *RDR2* were 5'-ccrTACCTCTCAATGATGCTTCG - 3' and 5' - GATCGGTTGCATGGATGAAAtgg - 3' (PAM sequences in lowercase). The protospacers for *NRPE1* were 5' - ccaGTGCCAATATACCATCCTGC - 3' and 5' - GAAATGTCTGAAGATAAAGAagg - 3'.

<u>Plant transformation, genotyping, and phenotyping</u>

C. grandiflora plants were transformed using *Agrobacterium*-mediated floral dip as described in (Dew-Budd et al., 2019). Briefly, developing fruits and opened flowers were removed prior to the first *Agrobacterium* dipping. Plants were dipped 3-4 times at an interval of 4-7 days, and were hand pollinated daily following the dipping. *C. rubella* and *C. sativa* were transformed via standard floral dip (Clough and Bent, 1998). Positive transformants were selected through EGFP visualization in mature seeds using a Zeiss Axiozoom 16 fluorescent stereo microscope with a 488 nm filter.

RdDM mutants were genotyped with PCR primers flanking the protospacers (**Supplementary Table 4**). Size differences between wild-type and mutant amplicons were visualized on an agarose gel and bands were sequenced to identify the exact mutation.

For *C. sativa*, dried fruits were collected from five individuals of each genotype to measure seed set. To prevent bias due to preferential dehiscence of the largest fruits, *C. rubella* and *C. grandiflora* seed set was measured 7 days after manual pollination (DAP), which is

approximately torpedo stage in wild type. For *C. grandiflora* all plants were genotyped and only crossed if they shared no more than one S allele to avoid confounding effects from S-locus incompatibility. Upon maturation, seeds were dried for at least two weeks under vacuum before measuring in batches of twenty on an ultra-microbalance (Sartorius Cubis MSE3.6P-000-DM).

Pollen grains were harvested by vortexing and centrifuging flowers in pollen isolation buffer (100 mM NaPO$_4$, pH 7.5; 1 mM EDTA; 0.1% (v/v) Triton X-100. The flowers and the fixing solution were removed and replaced with 1 µg/mL DAPI (4′,6-diamidino-2-phenylindole, BD Biosciences) diluted in pollen isolation buffer before visualizing under a fluorescence microscope using a 420 nm filter (Zeiss LSM880 Inverted Confocal Microscope). Pollen was considered normal if two sperm cell nuclei were distinguishable from the vegetative nucleus. The majority of abnormal pollen were small and did not fluoresce.

Capsella grandiflora genome sequencing, assembly, and annotation

Genomic DNA was isolated using a modified protocol of the Circulomics Nanobind Plant Nuclei Big DNA Kit to account for *C. grandiflora*'s small genome. The modified extraction replaced the nuclei isolation with Circulomics' supplementary direct tissue lysis protocol. Approximately 1 µg of genomic DNA was prepared using the Oxford Nanopore LSK-109 kit without shearing and sequenced on a MinION system using a R9.4 flow cell. Reads were called during sequencing using Guppy (v3.2.6) with a MinIT device (ont-minit-release 19.10.3) used for processing.

Adaptors were trimmed from the reads with Porechop (v0.2.3, https://github.com/rrwick/Porechop) using default parameters, and then filtered for reads > 500 bp in length with an average quality > 9 using Nanofilt (v2.6.0) (De Coster et al., 2018). Reads matching either the *C. grandiflora* chloroplast genome or the *C. rubella* mitochondrial genome were removed using minimap2 (v2.17) (Li, 2018) and SAMtools (v1.4) (Li et al., 2009). The remaining reads were *de novo* assembled using Canu (v1.9) (Koren et al., 2017) with default parameters and an estimated genome size of 120 Mb. The resulting genome was polished using the filtered long read data with Racon (v.1.4.3, non-default parameters: -m 8 -x 6 -g 8) (Vaser et al., 2017) and Medaka (v0.12.1, https://github.com/nanoporetech/medaka). Accession-specific publicly available short-read data from Williamson (2014, SRR1508428) was used to reiteratively polish three times with Pilon (v1.23) (Walker et al., 2014; Williamson et al., 2014). Due to the heterozygosity of *C. grandiflora*, the polished genome was further processed with Purge Haplotigs to separate the main genome and the haplotigs (Roach et al., 2018). Thirty-seven contigs less than 1 kb were removed from the main assembly and seven contigs less than 1 kb were removed from the haplotig assembly. The assembly is available at NCBI (BioProject PRJNA988139).

Small RNA-seq generation and analysis

Small RNA sequencing libraries were prepared using a standard protocol (Grover et al., 2018). Briefly, after RNA precipitation, small RNA was enriched with a mirVana miRNA isolation kit (Thermo Fisher Scientific) followed by NEBNext small RNA library preparation (New England Biolabs). Three independent biological replicates were prepared for each genotype then

sequenced at the University of Arizona Genomics Core on a NextSeq500. Raw reads are available at NCBI SRA (BioProject number PRJNA986946).

Small RNA reads were processed and aligned to the appropriate genome using the sRNA snakemake workflow (v1.0) (Bose and Grover, 2019). Briefly, raw reads were trimmed for size and quality using Trim Galore! (v0.6.2) (Krueger, 2012), contaminating non-coding RNAs were removed using bowtie (v1.2.2) (Langmead et al., 2009) with the Rfam database entries for *A. thaliana* and *C. rubella* (Kalvari et al., 2018), then reads aligning to the species-specific chloroplast and *C. rubella* mitochondrial genomes were removed with bowtie. The remaining reads were aligned to the reference genome using ShortStack (v3.8.5) (Axtell, 2013) with the snakemake workflow's default parameters except one mismatch was allowed during ShortStack alignment. Biological replicates were pooled for locus-level analyses.

Acknowledgements and Funding

The authors wish to thank the following colleagues for their assistance and expertise: University of Arizona Genomics Core for Illumina sequencing, Professor David Baltrus (University of Arizona) for Nanopore expertise, Professor Steven Wright (University of Toronto) for seeds of *Capsella* species, Professor John McKay (Colorado State University) for *Camelina sativa* seed, Arizona Laboratory for Emerging Contaminants for access to the microbalance and Danielle Barrientes for training on its use, and undergraduate assistants Jeff Clark, Jack Stearns, and Cecelia Martinez. The authors are grateful for support from the National Science Foundation (IOS-1546825 to R.A.M. and M.A.B.), the USDA National Institute of Food and Agriculture (AFRI 2021-67013-33797 to R.A.M.), the USDA Hatch funding (ARZT 1361510-H25-249 to R.A.M and M.A.B), and The University of Arizona College of Agriculture and Life Sciences iVIP award (to M.A.B.).

Author Contributions

RAM and MAB designed the research; KDB, HTC, TK, BCD, and JAR performed research; KDB, HTC, RAM, and MAB analyzed data and wrote the manuscript.

References

Agren JÅ, Wang W, Koenig D, Neuffer B, Weigel D, Wright SI (2014) Mating system shifts and transposable element evolution in the plant genus Capsella. BMC Genomics **15**: 602–609

Alger EI, Edger PP (2020) One subgenome to rule them all: underlying mechanisms of subgenome dominance. Curr Opin Plant Biol **54**: 108–113

Axtell MJ (2013) ShortStack: Comprehensive annotation and quantification of small RNA genes. RNA **19**: 740–751

Bose M, Grover JW (2019) sRNA Snakemake Workflow.

Brandvain Y, Haig D (2005) Divergent mating systems and parental conflict as a barrier to hybridization in flowering plants. Am Nat **166**: 330–338

Brandvain Y, Haig D (2018) Outbreeders pull harder in a parental tug-of-war. Proc Natl Acad Sci U S A **115**: 11354–11356

Brock JR, Dönmez AA, Beilstein MA, Olsen KM (2018) Phylogenetics of Camelina Crantz. (Brassicaceae) and insights on the origin of gold-of-pleasure (Camelina sativa). Mol Phylogenet Evol **127**: 834–842

Burgess D, Chow HT, Grover JW, Freeling M, Mosher RA (2022) Ovule siRNAs methylate protein-coding genes in trans. Plant Cell. doi: 10.1093/plcell/koac197

de la Chaux N, Tsuchimatsu T, Shimizu KK, Wagner A (2012) The predominantly selfing plant Arabidopsis thaliana experienced a recent reduction in transposable element abundance compared to its outcrossing relative Arabidopsis lyrata. Mob DNA **3**: 2

Cheng F, Sun C, Wu J, Schnable J, Woodhouse MR, Liang J, Cai C, Freeling M, Wang X (2016) Epigenetic regulation of subgenome dominance following whole genome triplication in Brassica rapa. New Phytol **211**: 288–299

Chow HT, Mosher RA (2023) Small RNA-mediated DNA methylation during plant reproduction. Plant Cell. doi: 10.1093/plcell/koad010

Clough SJ, Bent AF (1998) Floral dip: a simplified method forAgrobacterium-mediated transformation ofArabidopsis thaliana. The Plant Journal **16**: 735–743

De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C (2018) NanoPack: visualizing and processing long-read sequencing data. Bioinformatics **34**: 2666–2669

Dew-Budd K, David B, Beilstein MA (2019) Agrobacterium-mediated floral-dip transformation of the obligate outcrosser Capsella grandiflora. bioRxiv. doi: 10.1101/757328

Erdmann RM, Picard CL (2020) RNA-directed DNA Methylation. PLoS Genet **16**: e1009034

Erdmann RM, Satyaki PRV, Klosinska M, Gehring M (2017) A Small RNA Pathway Mediates Allelic Dosage in Endosperm. Cell Rep **21**: 3364–3372

Fauser F, Schiml S, Puchta H (2014) Both CRISPR/Cas-based nucleases and nickases can be used efficiently for genome engineering in Arabidopsis thaliana. Plant Journal **79**: 348–359

Forestan C, Farinati S, Aiese Cigliano R, Lunardon A, Sanseverino W, Varotto S (2017) Maize RNA PolIV affects the expression of genes with nearby TE insertions and has a genome-wide repressive impact on transcription. BMC Plant Biol **17**: 161

Foxe JP, Slotte T, Stahl EA, Neuffer B, Hurka H, Wright SI (2009) Recent speciation associated with the evolution of selfing in Capsella. Proceedings of the National Academy of Sciences **106**: 5241–5245

Fukudome A, Singh J, Mishra V, Reddem E, Martinez-Marquez F, Wenzel S, Yan R, Shiozaki M, Yu Z, Wang JC-Y, et al (2021) Structure and RNA template requirements of *Arabidopsis* RNA-DEPENDENT RNA POLYMERASE 2. Proceedings of the National Academy of Sciences. doi: 10.1073/pnas.2115899118

Garsmeur O, Schnable JC, Almeida A, Jourda C, D'Hont A, Freeling M (2014) Two evolutionarily distinct classes of paleopolyploidy. Molecular Biology and Evolution **31**: 448–454

Gouil Q, Baulcombe DC (2016) DNA methylation signatures of the plant chromomethyltransferases. PLoS Genet **12**: e1006526

Grover CE, Gallagher JP, Szadkowski E, Yoo MJ, Flagel L, Wendel JF (2012) Homoeolog expression bias and expression level dominance in allopolyploids. New Phytologist **196**: 966–971

Grover JW, Burgess D, Kendall T, Baten A, Pokhrel S, King GJ, Meyers BC, Freeling M, Mosher RA (2020) Abundant expression of maternal siRNAs is a conserved feature of seed development. Proc Natl Acad Sci U S A **117**: 15305–15315

Grover JW, Kendall T, Baten A, Burgess D, Freeling M, King GJ, Mosher RA (2018) Maternal components of RNA-directed DNA methylation are required for seed development in Brassica rapa. Plant J **94**: 575–582

Guo Y-L, Bechsgaard JS, Slotte T, Neuffer B, Lascoux M, Weigel DD, Schierup MH (2009) Recent speciation of Capsella rubella from Capsella grandiflora, associated with loss of self-incompatibility and an extreme bottleneck. Proc Natl Acad Sci U S A **106**: 5246–5251

Haeussler M, Schönig K, Eckert H, Eschstruth A, Mianné J, Renaud J-B, Schneider-Maunoury S, Shkumatava A, Teboul L, Kent J, et al (2016) Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. Genome Biology **17**: 148

Haig D (2013) Kin conflict in seed development: an interdependent but fractious collective. Annu Rev Cell Dev Biol **29**: 189–211

Hollister JD, Gaut BS (2009) Epigenetic silencing of transposable elements: A trade-off between reduced transposition and deleterious effects on neighboring gene expression. Genome Res **19**: 1419–1428

Hollister JD, Smith LM, Guo Y-L, Ott F, Weigel D, Gaut BS (2011) Transposable elements and small RNAs contribute to gene expression divergence between Arabidopsis thaliana and Arabidopsis lyrata. Proc Natl Acad Sci U S A **108**: 2322–2327

Huang K, Wu X-X, Fang C-L, Xu Z-G, Zhang H-W, Gao J, Zhou C-M, You L-L, Gu Z-X, Mu W-H, et al (2021) Pol IV and RDR2: A two-RNA-polymerase machine that produces double-stranded RNA. Science **374**: 1579–1586

Johnston SA, den Nijs TP, Peloquin SJ, Hanneman RE Jr (1980) The significance of genic balance to endosperm development in interspecific crosses. Theor Appl Genet **57**: 5–9

Kagale S, Koh C, Nixon J, Bollina V, Clarke WE, Tuteja R, Spillane C, Robinson SJ, Links MG, Clarke C, et al (2014) The emerging biofuel crop Camelina sativa retains a highly undifferentiated hexaploid genome structure. Nature Communications **5**: 3706

Kalvari I, Argasinska J, Quinones-Olvera N, Nawrocki EP, Rivas E, Eddy SR, Bateman A, Finn RD, Petrov AI (2018) Rfam 13.0: shifting to a genome-centric resource for non-coding RNA families. Nucleic Acids Research **46**: D335–D342

Kanno T, Huettel B, Mette MF, Aufsatz W, Jaligot E, Daxinger L, Kreil DP, Matzke M, Matzke AJM (2005) Atypical RNA polymerase subunits required for RNA-directed DNA methylation. Nat Genet **37**: 761–765

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM (2017) Canu: scalable and accurate long-read assembly via adaptive k -mer weighting and repeat separation. Genome Research **27**: 722–736

Krueger F (2012) Trim Galore!: A wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files, with some extra functionality for MspI-digested RRBS-type (Reduced Representation Bisufite-Seq) libraries. http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/

Lafon-Placette C, Hatorangan MR, Steige KA, Cornille A, Lascoux M, Slotte T, Köhler C (2018) Paternally expressed imprinted genes associate with hybridization barriers in Capsella. Nat Plants **4**: 352–357

Langmead B, Trapnell C, Pop M, Salzberg SL (2009) Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biol **10**: R25

Law JA, Du J, Hale CJ, Feng S, Krajewski K, Palanca AMS, Strahl BD, Patel DJ, Jacobsen SE (2013) Polymerase IV occupancy at RNA-directed DNA methylation sites requires SHH1. Nature **498**: 385–389

Law JA, Jacobsen SE (2010) Establishing, maintaining and modifying DNA methylation patterns in plants and animals. Nat Rev Genet **11**: 204–220

Li C, Xu H, Fu F-F, Russell SD, Sundaresan V, Gent JI (2020) Genome-wide redistribution of 24-nt siRNAs in rice gametes. Genome Res **30**: 173–184

Li H (2018) Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics **34**: 3094–3100

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup (2009) The Sequence Alignment/Map format and SAMtools. Bioinformatics **25**: 2078–2079

Liu W, Duttke SH, Hetzel J, Groth M, Feng S, Gallego-Bartolome J, Zhong Z, Kuo HY, Wang Z, Zhai J, et al (2018) RNA-directed DNA methylation involves co-transcriptional small-RNA-guided slicing of polymerase V transcripts in Arabidopsis. Nat Plants **4**: 181–188

Loffer A, Singh J, Fukudome A, Mishra V, Wang F, Pikaard CS (2022) A DCL3 dicing code within Pol IV-RDR2 transcripts diversifies the siRNA pool guiding RNA-directed DNA methylation. Elife. doi: 10.7554/eLife.73260

Long J, Walker J, She W, Aldridge B, Gao H, Deans S, Vickers M, Feng X (2021) Nurse cell--derived small RNAs define paternal epigenetic inheritance in Arabidopsis. Science **373**: eabh0556

Mandáková T, Pouch M, Brock JR, Al-Shehbaz IA, Lysak MA (2019) Origin and evolution of diploid and allopolyploid Camelina genomes were accompanied by chromosome shattering. Plant Cell **31**: 2596–2612

Martinez G, Wolff P, Wang Z, Moreno-Romero J, Santos-González J, Conze LL, DeFraia C, Slotkin RK, Köhler C (2018) Paternal easiRNAs regulate parental genome dosage in Arabidopsis. Nat Genet **50**: 193–198

Matzke MA, Mosher RA (2014) RNA-directed DNA methylation: an epigenetic pathway of increasing complexity. Nat Rev Genet **15**: 394–408

Mosher RA, Melnyk CW, Kelly KA, Dunn RM, Studholme DJ, Baulcombe DC (2009) Uniparental expression of PolIV-dependent siRNAs in developing endosperm of Arabidopsis. Nature **460**: 283–286

Mosher RA, Schwach F, Studholme D, Baulcombe DC (2008) PolIVb influences RNA-directed DNA methylation independently of its role in siRNA biogenesis. Proc Natl Acad Sci U S A **105**: 3145–3150

Naito Y, Hino K, Bono H, Ui-Tei K (2015) CRISPRdirect: software for designing CRISPR/Cas guide RNA with reduced off-target sites. Bioinformatics **31**: 1120–1123

Pontier D, Yahubyan G, Vega D, Bulski A, Saez-Vasquez J, Hakimi M-A, Lerbs-Mache S, Colot V, Lagrange T (2005) Reinforcement of silencing at transposons and highly repeated sequences requires the concerted action of two distinct RNA polymerases IV in Arabidopsis. Genes Dev **19**: 2030–2040

Ream TS, Haag JR, Wierzbicki AT, Nicora CD, Norbeck AD, Zhu J-K, Hagen G, Guilfoyle TJ, Pasa-Tolić L, Pikaard CS (2009) Subunit compositions of the RNA-silencing enzymes Pol IV and Pol V reveal their origins as specialized forms of RNA polymerase II. Mol Cell **33**: 192–203

Rebernig CA, Lafon-Placette C, Hatorangan MR, Slotte T, Köhler C (2015) Non-reciprocal Interspecies Hybridization Barriers in the Capsella Genus Are Established in the Endosperm. PLoS Genet **11**: e1005295-19

Renny-Byfield S, Gong L, Gallagher JP, Wendel JF (2015) Persistence of Subgenomes in Paleopolyploid Cotton after 60 My of Evolution. Molecular Biology and Evolution **32**: 1063–1071

Roach MJ, Schmidt SA, Borneman AR (2018) Purge Haplotigs: allelic contig reassignment for third-gen diploid genome assemblies. BMC Bioinformatics **19**: 460

Rodrigues JA, Ruan R, Nishimura T, Sharma MK, Sharma R, Ronald PC, Fischer RL, Zilberman D (2013) Imprinted expression of genes and small RNA is associated with localized hypomethylation of the maternal genome in rice endosperm. Proc Natl Acad Sci U S A **110**: 7934–7939

Satyaki PRV, Gehring M (2019) Paternally Acting Canonical RNA-Directed DNA Methylation Pathway Genes Sensitize Arabidopsis Endosperm to Paternal Genome Dosage. Plant Cell **31**: 1563–1578

Scott RJ, Spielman M, Bailey J, Dickinson HG (1998) Parent-of-origin effects on seed development in Arabidopsis thaliana. Development **125**: 3329–3341

Sicard A, Lenhard M (2011) The selfing syndrome: a model for studying the genetic and evolutionary basis of morphological adaptation in plants. Annals of Botany **107**: 1433–1443

Singh J, Mishra V, Wang F, Huang H-Y, Pikaard CS (2019) Reaction Mechanisms of Pol IV, RDR2, and DCL3 Drive RNA Channeling in the siRNA-Directed DNA Methylation Pathway. Mol Cell **75**: 576-589.e5

Slotte T, Hazzouri KM, Ågren JA, Koenig D, Maumus F, Guo Y-L, Steige K, Platts AE, Escobar JS, Newman LK, et al (2013) The Capsella rubella genome and the genomic consequences of rapid mating system evolution. Nat Genet **45**: 831–835

Stroud H, Do T, Du J, Zhong X, Feng S, Johnson L, Patel DJ, Jacobsen SE (2014) Non-CG methylation patterns shape the epigenetic landscape in Arabidopsis. Nat Struct Mol Biol **21**: 64–72

Vaser R, Sović I, Nagarajan N, Šikić M (2017) Fast and accurate de novo genome assembly from long uncorrected reads. Genome Research **27**: 737–746

Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, et al (2014) Pilon: An Integrated Tool for Comprehensive Microbial Variant Detection and Genome Assembly Improvement. PLoS ONE **9**: e112963

Wang F, Huang H-Y, Huang J, Singh J, Pikaard CS (2023) Enzymatic reactions of AGO4 in RNA-directed DNA methylation: siRNA duplex loading, passenger strand elimination, target RNA slicing, and sliced target retention. Genes Dev. doi: 10.1101/gad.350240.122

Wang Q, Xue Y, Zhang L, Zhong Z, Feng S, Wang C, Xiao L, Yang Z, Harris CJ, Wu Z, et al (2021) Mechanism of siRNA production by a plant Dicer-RNA complex in dicing-competent conformation. Science **374**: 1152–1157

Wang X, Wang H, Wang J, Sun R, Wu J, Liu S, Bai Y, Mun J-H, Bancroft I, Cheng F, et al (2011) The genome of the mesopolyploid crop species Brassica rapa. Nat Genet **43**: 1035–1039

Wang Z, Butel N, Santos-González J, Borges F, Yi J, Martienssen RA, Martinez G, Köhler C (2020) Polymerase IV plays a crucial role in pollen development in Capsella. Plant Cell **32**: 950–966

Wei L, Gu L, Song X, Cui X, Lu Z, Zhou M, Wang L, Hu F, Zhai J, Meyers BC, et al (2014) Dicer-like 3 produces transposable element-associated 24-nt siRNAs that control agricultural traits in rice. Proc Natl Acad Sci U S A **111**: 3877–3882

Wierzbicki AT, Ream TS, Haag JR, Pikaard CS (2009) RNA polymerase V transcription guides ARGONAUTE4 to chromatin. Nat Genet **41**: 630–634

Williamson RJ, Josephs EB, Platts AE, Hazzouri KM, Haudry A, Blanchette M, Wright SI (2014) Evidence for Widespread Positive and Negative Selection in Coding and Conserved Noncoding Regions of Capsella grandiflora. PLoS Genetics **10**: e1004622

Woodhouse MR, Cheng F, Pires JC, Lisch D (2014) Origin, inheritance, and gene regulatory consequences of genome dominance in polyploids. Proceedings of the National Academy of Sciences **111**: 6527–6527

Zemach A, Kim MY, Hsieh P-H, Coleman-Derr D, Eshed-Williams L, Thao K, Harmer SL, Zilberman D (2013) The Arabidopsis nucleosome remodeler DDM1 allows DNA methyltransferases to access H1-containing heterochromatin. Cell **153**: 193–205

Zheng K, Wang L, Zeng L, Xu D, Guo Z, Gao X, Yang D-L (2021) The effect of RNA polymerase V on 24-nt siRNA accumulation depends on DNA methylation contexts and histone modifications in rice. Proc Natl Acad Sci U S A. doi: 10.1073/pnas.2100709118

Zhong X, Du J, Hale CJ, Gallego-Bartolome J, Feng S, Vashisht AA, Chory J, Wohlschlegel JA, Patel DJ, Jacobsen SE (2014) Molecular mechanism of action of plant DRM de novo DNA methyltransferases. Cell **157**: 1050–1060

Zhou M, Coruh C, Xu G, Martins LM, Bourbousse C, Lambolez A, Law JA (2022) The CLASSY family controls tissue-specific DNA methylation patterns in Arabidopsis. Nat Commun **13**: 244