Allelic bias when performing in-solution enrichment of ancient human DNA

Authors: Roberta Davidson[1#], Matthew P. Williams[1,2], Xavier Roca-Rada[1], Kalina

Kassadjikova[3], Raymond Tobler[1,4,5], Lars Fehren-Schmitz[3,6#^], Bastien Llamas[1,5,7,8#^]

Corresponding authors: roberta.davidson@adelaide.edu.au,

bastien.llamas@adelaide.edu.au, lfehrens@ucsc.edu

^ Equal contributor

Affiliations

    1. The Australian Centre for Ancient DNA and the Environment Institute, School of

    Biological Sciences, The University of Adelaide, Adelaide, SA 5005, Australia

    2. Biology Department, The Pennsylvania State University, University Park,

    Pennsylvania, USA

    3. UCSC Paleogenomics, Department of Anthropology, University of California,

    Santa Cruz, California, USA

    4. Evolution of Cultural Diversity Initiative, Australian National University,

    Canberra, ACT 2601, Australia

    5. Centre of Excellence for Australian Biodiversity and Heritage, The University of

    Adelaide, Adelaide, SA 5005, Australia

    6. UCSC Genomics Institute, University of California, Santa Cruz, California, USA

    7. National Centre for Indigenous Genomics, Australian National University,

    Canberra, ACT 0200, Australia

    8. Indigenous Genomics, Telethon Kids Institute, Adelaide, SA 5000, Australia

Abstract

In-solution hybridisation enrichment of genetic variation is a valuable methodology in human paleogenomics. It allows enrichment of endogenous DNA by targeting genetic markers that are comparable between sequencing libraries. Many studies have used the 1240k reagent—which enriches 1,237,207 genome-wide SNPs—since 2015, though access was restricted. In 2021, Twist Biosciences and Daicel Arbor Biosciences independently released commercial kits that enabled all researchers to perform enrichments for the same 1240k SNPs. We used the Daicel Arbor Biosciences Prime Plus kit to enrich 268 ancient samples from four continents. We identified a systematic assay bias that increases genetic similarity between enriched samples and that cannot be explained by batch effects. We present the impact of the bias on population genetics inferences (e.g., Principal Components Analysis, f-statistics) and genetic relatedness (READ). We compare the Prime Plus bias to that previously reported of the legacy 1240k enrichment assay. In f-statistics, we find that all Prime-Plus-generated data exhibit artefactual excess shared drift, such that within-continent relationships cannot be correctly determined. In PCA, the first PCs may capture the assay bias rather than the underlying genetic structure. The bias is more subtle in READ, though interpretation of the results can still be misleading in specific contexts. We expect the bias may affect analyses we have not yet tested. Our observations support previously reported concerns for the integration of different data types in paleogenomics. We also caution that technological solutions to generate 1240k data necessitate a thorough validation process before their adoption in the paleogenomic community.

46 **Keywords**

47 Bias, enrichment, 1240k, ancient DNA

INTRODUCTION

One of the major challenges faced when working with ancient DNA is the high

proportion of exogenous DNA contamination present in the DNA extract. This

contamination is mostly from microbes that invade the organism post-mortem or are

present in the soil where the specimen was buried. It can also be introduced during

laboratory experiments. To address this challenge, one method that has become popular in

paleogenomic studies is the in-solution enrichment of target genomic regions using pre-

designed oligonucleotides as molecular "probes" or "baits". This technique increases the

proportion of endogenous DNA of interest in a sequencing library, reducing sequencing

cost and leading to larger quantities of comparable data across individual samples

compared to using shotgun sequencing. Ultimately, the increase in analytical power

resulting from target enrichment has enabled large-scale ancient human population

genetic (i.e., paleogenomic) studies (Fumagalli, 2013).

In human paleogenomic studies, in-solution enrichment assays can be designed to target

complete mitochondrial genomes (Briggs et al., 2009; Brotherton et al., 2013; Llamas et al.,

2016; Maricic et al., 2010), the non-recombining portion of chromosome Y (Cruz-Dávalos

et al., 2018; Rohrlach et al., 2021), genome-wide SNPs (Fu et al., 2015; Haak et al., 2015;

Mathieson et al., 2015), exomes (Castellano et al., 2014), autosomes (Fu et al., 2013), or

complete genomes (Ávila-Arcos et al., 2015; Carpenter et al., 2013). In 2012, Patterson

and colleagues proposed a molecular bait design that used a specific ascertainment strategy

to enable unbiased population genetics studies of global human populations through time

(Patterson et al., 2012). The first baits using this design were applied in a handful of

landmark paleogenomic studies of European populations (Fu et al., 2015; Haak et al., 2015;

Mathieson et al., 2015). These bait sets were subsequently restricted to 1240k genome-

wide SNPs, which paved the way for the generation of thousands of individual datasets

and launched the paleogenomic era (Marciniak & Perry, 2017; Olalde & Posth, 2020;

Skoglund & Mathieson, 2018). Hereafter, we will refer to this set of baits as the 'legacy

1240k reagent'.

Since the initial publication of the molecular baits sequences in 2015, the legacy 1240k

reagent was only available to a handful of laboratories through a commercial agreement.

Subsequently, in 2021 two biotechnology companies—Daicel Arbor Biosciences and Twist

Bioscience—released commercial in-solution enrichment kits that target the same set of

1240k SNPs (Rohland et al., 2022). In particular, the release of the suite of myBaits Expert

Human Affinities kits by Daicel Arbor Biosciences aimed to make the 1240k enrichment

assay accessible to all research groups worldwide, including those with limited human

resources and funding. Importantly, the molecular design of both the new bait sets differs

from the legacy 1240k design, with the differences specific to the Twist reagents described

in a recently published study (Rohland et al., 2022). While the bait design used in the

myBaits Expert Human Affinities kits is also known to differ from both the legacy 1240k

and Twist assays (Daicel Arbor Biosciences, personal communication), it is a proprietary

product and the specific details are unknown to the present authors. The bait design and

associated laboratory protocol steps of any enrichment reagent are key to its efficacy to

enrich endogenous DNA and to achieve an unbiased enrichment that respects the true

proportions of reference and alternate alleles at heterozygous sites.

Here, we report genomic data from 268 ancient human individuals from western Eurasia,

South and Central America (from periods preceding the colonial contact period between

these populations), which were generated using myBaits Expert Human Affinities Kit

"Prime Plus" (Daicel Arbor Biosciences) for various independent studies. In addition to

the standard set of 1240k SNPs, the Prime Plus (PP) probes target 46,218 Y Chromosome

SNPs, as well as human, Neanderthal and Denisovan whole mitochondrial genomes.

Notably, when performing routine population genetic analyses on the PP-generated

dataset using f-statistics (Patterson et al., 2012), we observed a significant excess of

shared genetic drift between any two populations enriched with the PP bait set, compared

with published data generated with the legacy 1240k reagent or shotgun sequencing.

These unexpected affinities could not be explained by batch effects or previously reported

demographic events.

In the following sections, we outline our investigation and characterisation of the bias

observed in PP-enriched ancient DNA data via principal components analyses,

genetic relatedness analyses and f-statistics. In total, we attempted 21 strategies to

reduce or negate the observed bias, which included two wet-lab protocol adjustments

and numerous bioinformatic techniques, but we were unable to correct the f-

statistics scores to a satisfactory standard. We conclude by recommending a series of

principles for researchers to apply when co-analysing PP-enriched data with 1240k

legacy and shotgun (SG) data, which mitigate the assay bias by adjusting the

construction of f-statistics. These guidelines may also be useful for reviewers and general

readers when assessing the robustness of studies that have used data generated with the

myBaits Expert Human Affinities Kit "Prime Plus" kit.

MATERIALS AND METHODS

The data used in this study were initially generated for various studies intended by the

authors. However, we collated all independent datasets into one overarching dataset after

identifying some problems with Prime Plus data, resulting in the present study.

Samples

We generated PP-enriched data for a total of 268 ancient samples, sourced from Spain,

Portugal, South-Western Asia, Mexico, Guatemala, Peru, Bolivia, Argentina and Chile. All

samples from the Americas pre-date 16th-century European contact (see Table S2 for

sample metadata). In our analyses, we use a comparative dataset of 106 shotgun data

samples and 291 legacy 1240k samples (Bergström et al., 2020; Bongers et al., 2020; de la

Fuente et al., 2018; González-Fortes et al., 2019; Günther et al., 2015; Lindo et al., 2018;

Lipson et al., 2017; Mallick et al., 2016; Martiniano et al., 2017; Mathieson et al., 2015;

Moreno-Mayar et al., 2018; Nakatsuka, Lazaridis, et al., 2020; Nakatsuka, Luisi, et al.,

2020; Olalde et al., 2018, 2019; Popović et al., 2021; Posth et al., 2018; Prüfer et al., 2014;

Raghavan et al., 2015; Valdiosera et al., 2018; Villalba-Mouco et al., 2019). A complete list

of the comparative samples used in this study and relevant citations are provided in Table

S3.

Data generation

DNA extraction

For the samples from Spain, DNA was extracted from tooth root powder using a method

optimised to retrieve highly degraded ancient DNA fragments (Dabney et al., 2013) at the

Autonomous University of Barcelona (UAB). For Southwest Asian, Argentinian,

Portuguese and some Mexican samples, the same methods were undertaken at the

Australian Centre for Ancient DNA (ACAD) at the University of Adelaide. For the rest of

the samples from the Americas, DNA was extracted similarly, but with a modification to

the protocol that includes a bleach pre-wash step (Boessenkool et al., 2017; Dabney et al.,

2013), at the University of California, Santa Cruz (UCSC) Human Paleogenomics Lab.

Table S2 provides a full summary of which lab processed which samples at each step.

Library preparation

For the Southwest Asian, Spanish, Portuguese, Argentinian and some Mexican samples,

partially UDG-treated double-stranded DNA libraries (Rohland et al., 2015) were

generated at ACAD. For the Bolivian, Peruvian, Chilean, Guatemalan and the rest of the

Mexican samples, partially UDG-treated single-stranded DNA libraries (Kapp et al., 2021)

were prepared at UCSC. Note that the difference in library structures should not impact

the enrichment bias as libraries are first denatured and then hybridised to the baits.

Library enrichment

Before enrichment, libraries were over-amplified in order to reach the minimum required

DNA input of 500 ng per sample. Library enrichment was then performed according to

the suggested myBaits protocol of two enrichment rounds with hybridisation temperature

at 70°C and 20 amplification cycles in the first enrichment followed by 10 cycles in the

second round. Samples were grouped in pools of 2–4 libraries according to the endogenous

DNA proportions obtained from screening of the initial shotgun sequences. Samples were

processed at UCSC or at ACAD (see Table S2 for a breakdown of which lab processed

which samples at each step).

Additionally, two experiments that examined the utility of adjustments to standard

enrichment protocols were performed on a subset of libraries at UCSC (Table 1). In

experiment A, a single round of hybridisation at 70°C was performed with 12 cycles of

amplification. A single round of hybridisation was also used in Experiment B but at 62.5°C

and with 13 cycles of amplification.

Sequencing

Libraries enriched at UCSC were paired-end sequenced on a HiSeq 4000 at Fulgent

Genomics (Los Angeles, USA). Paired-end sequencing using an Illumina NovaSeq 6000

(200 cycles) was performed by the Kinghorn Centre for Clinical Genomics (Sydney,

Australia) for libraries enriched at ACAD (Table S2).

Data processing and genotype calling

Data were processed either using nf-core/eager (Fellows Yates et al., 2021), or UCSC's in-

house processing pipeline, Batpipe (https://github.com/mjobin/batpipe). The terminal 2bp

at either end of each read were trimmed to remove ancient DNA damage and then

mapped to the GRCh37d5 reference genome using bwa-aln with parameters -l 1024 -n

0.01 -o 2 (Oliva et al., 2021). Other mapping parameters vary slightly between different

datasets used in this study, such as mapping quality threshold 20-30 and minimum read

length 20-30 (see Table S2 for details of processing parameters for each population).

Pseudo-haploid genotypes were called at the enrichment target sites using the random

haploid mode within pileupCaller (https://github.com/stschiff/sequenceTools).

Population genomics assessment of capture bias

PCA

We computed principal components analyses using both EMU (Meisner et al., 2021) and smartPCA (Patterson et al., 2006; Price et al., 2006) with and without projection to test how the data bias affected each type of PCA.

f-statistics

Outgroup-f_3 and f_4 statistics were computed using the ADMIXTOOLS package (Patterson et al., 2012).

READ

We tested the impact of the enrichment bias on genetic relatedness using READ, i.e. Relatedness Estimation for Ancient DNA (Monroy Kuhn et al., 2018), which is a widely used PMR-based estimator for genetic relatedness for ancient DNA data.

Filters and strategies to resolve bias

In order to estimate the effect of various bioinformatic approaches to mitigate the bias observed in f_4 statistics, we calculated f-statistics in the following four forms:

1) f_4(Mbuti, Spain.PP.2rnd; Target, Peru.Shotgun)

2) f_4(Mbuti, Spain.PP.2rnd; Target, Peru.1240k)

3) f_4(Mbuti, Spain.1240k; Target, Peru.Shotgun)

4) f_4(Mbuti, Spain.1240k; Target, Peru.1240k)

In each, the target population was the Peru.PP population after application of one of the

21 strategies to reduce bias detailed below.

SNP Missingness

Amongst the set of 1240k SNPs assayed by the PP probes, each SNP was missing at least

once in our dataset. Hence, we used the full data matrix with missing genotypes and

reported the number of overlapping SNPs included in each f-statistic comparison where

relevant.

To the target population (Peru.PP), we applied two data missingness filters using the --

geno flag in PLINK 1.9 (Chang et al., 2015)—namely, 0.0 and 0.5, which correspond to

removing SNPs with >0% or >50% missingness, respectively.

Minor allele frequency

We tested some simple filters for minor allele frequency (i.e., 0.3, 0.1 and 0.01) using the -

-*maf* flag in PLINK 1.9 (Chang et al., 2015).

Linkage disequilibrium

We filtered for linkage disequilibrium (LD), with the intention to remove potential

artefactual linkage introduced by one bait targeting multiple SNP sites. We used the

PLINK 1.9 LD pruning parameter --*indep-pairwise* with a window size of 3 variants,

sliding in 1 variant steps and tested correlation threshold values of $r^2 = 0.1, 0.3, 0.5, 0.7$

and 0.9 (Chang et al., 2015). These filters resulted in datasets of 300,838, 346,428, 379,345,

388,038, and 388,678 SNPs, respectively.

Given that each r^2 threshold produced approximately 300,000 SNPs, to assess if the

improvement in f_4-value was driven by the reduction in bias attributable to artefactual

linkage, rather than changes to other statistical properties of the data (such as decreased

power due to a reduced number of SNPs), we repeated our analyses with random sub-

samples of 300,000 SNPs, replicated five times, using the PLINK 1.9 flag *--thin-count*

(Chang et al., 2015).

Transversions

Sequencing libraries were partially treated with UDG (Kapp et al., 2021) and the terminal

two nucleotides at either end of all mapped reads were trimmed when processing the data

with nf-core/eager (Fellows Yates et al., 2021). Therefore, we did not expect to observe a

strong impact of ancient DNA damage in the analyses. Nevertheless, we filtered out

transitions to minimise the potential impact of ancient DNA damage, which is typically

characterised by C-to-T substitutions at the 5' end of sequencing reads and C-to-T or G-

to-A substitutions at the 3' end of sequencing reads when using single-stranded or double-

stranded libraries, respectively (Krause et al., 2010; Meyer et al., 2012).

Application of Rohland et al. 2022 filter

We applied the filter suggested by Rohland and colleagues (Rohland et al., 2022) to extract

SNPs that will allow co-analysis between 1240K data generated with shotgun sequencing,

or the legacy 1240k, Prime Plus or Twist enrichments.

PCA derived filter

PCAs computed for a combination of PP-, legacy-1240k- and shotgun-generated ancient

samples with smartPCA (Patterson et al., 2006; Price et al., 2006) displayed patterns

consistent with assay bias along PC2. Hence, we used this principal component to

construct an assay-based filter by extracting SNPs weighted $> |1|$ in PC2 of Fig 2B (n =

129,982 SNPs). The SNP weight threshold was validated by the observed removal of the

assay bias along PC2 when computing PCA after applying this filter (Fig 2C).

Read length and mapping quality threshold

We filtered out reads that were shorter than 50 bp and/or with mapping quality less than

20, as suggested by Günther & Nettelblad to reduce reference bias when mapping

paleogenomic data (Günther & Nettelblad, 2019).

Off-target genotypes

Given the hypothesis that the bias is caused by bait-target interactions and that the

enriched library efficacy is not 100%, we called off-target (non-specific) genotypes on the

assumption that they would not be impacted by any bait-mediated mechanism driving the

bias. The genotyping of off-target sites was performed using a standard pseudo-haploid

pileupCaller call with the dbSNP set 138 (Sherry et al., 2001) after filtering out all PP

target sites. Additionally, we called genotypes that were at least 100 bp away from target

sites to eliminate SNPs that may have been covered by a bait but were not the target

locus.

Majority call pseudo-haploid and random diploid genotypes

In addition to standard pseudo-haploid calls based on the random selection of a single

allele at prespecified SNPs, majority allele pseudo-haploid genotypes were also called at

sites with greater than 5X depth-of-coverage using the majority call mode in pileupCaller.

Finally, diploid genotypes were called using the random diploid call mode in pileupCaller,

where one read is chosen at random, followed by a second chosen at random without

replacement, and sites covered by only one read are considered missing.

Investigation of allele call bias

We investigated the rate of each allele call at heterozygous SNP sites for the most

common substitutions: (A/C, T/G, A/G, T/C), where the inverse substitution was also

included such that i.e., A/C and C/A SNPs are collated together. The relative allele call

frequencies were calculated per SNP (ignoring missing SNPs) in aggregate across all

population datasets for shotgun-, legacy-1240k- and PP-enriched genotypes. SNP

missingness was quantified across 100 bins with equal numbers of SNPs, and the mean

allele frequency per bin was plotted against SNP missingness.

RESULTS

Global bias assessment

To quantify the extent of the PP-associated bias in outgroup-f_3 statistics, we

calculated statistics of the form f_3(Mbuti; Target, X), where Target refers to samples that

were enriched using PP and originate either from ancient Spain (Spain.PP.2rnd;

unpublished) or Peru (Peru.PP.2rnd; unpublished), and X denotes a set of South American

and European ancient reference populations either generated from shotgun sequencing or

enriched with the PP or legacy 1240k kits.

For f_3 statistics that use Spain.PP.2rnd as the Target population (Fig 1A), all non-PP

European populations (shotgun or legacy 1240k data) exhibit more shared drift with the

Spain.PP.2rnd population than do the South American populations of either enrichment

type (Fig 1A), and the relative relationships between the European populations are

consistent with previous reports (Martiniano et al., 2017; Olalde et al., 2019; Valdiosera et

al., 2018). However, the four South American PP populations share more genetic drift

with the Spain.PP.2rnd target population relative to other non-PP South American

populations. Similarly, the inter-continental relationship is preserved when Peru.PP.2rnd

is used as the target population, as all South American populations have substantially more

shared drift with the Peru.PP.2rnd target population than with European populations, yet

the intra-continental relationships appear to be violated by the biased shared drift

between PP populations (Fig 1B). Further, Spain.PP.2rnd and Portugal.PP.2rnd exhibit a

strong attraction toward Peru.PP.2rnd samples relative to the other European non-PP

populations.

We also observe variance between 1240k and shotgun datasets, which is likely due to a

combination of different ancestries and 1240k bias (Rohland et al., 2022), however in most

cases the shotgun and 1240k error bars overlap, whereas the Prime Plus bias is more

pronounced. Clearly, the bias observed in PP-generated data is not unique to South

American populations but is ubiquitous across the highly diverged populations of Europe

and South America, strongly suggesting that the bias affects all global populations.

Principal Components Analysis

We tested the impact of PP bias on multiple PCA approaches, including the widely used

smartPCA (Patterson et al., 2006; Price et al., 2006) and EMU, which is reported to handle

data missingness well (Meisner et al., 2021). For each approach, we computed PCAs at

continental and global scales to investigate how missingness and the PP bias manifested at

different geographical scales.

Continental PCA

Initially, we performed a PCA using EMU on a dataset of 386 ancient genomes from South

America including shotgun (n = 38), legacy 1240k (n = 223) and PP-enriched (n = 125)

data (see Table S1 for a detailed list of samples). Sample clustering in the first three

principal components of the PCA is driven by population structure (Fig S1, right) and

there is no apparent bias exhibited by the data source type (Fig S1, left). Note that no

individual samples were assayed with each of the three methods investigated, but many

samples of different data types derive from the same localised ancestry groups. This result

concurs with a recent study (Rohland et al., 2022), which also reported that PCA is robust

to combinations of these three data types. However, when we used smartPCA on the same

386 ancient genomes (without projection onto modern diversity), the resulting PCA

produced sample clustering along PC2 and PC3 that was driven by the assay type rather

than ancestry (Fig S2). Additionally, samples moved toward the origin due to SNP

missingness (Fig S4), an issue known to affect smartPCA that is largely circumvented by

EMU in the present study.

Global PCA

Subsequently, we performed another PCA with EMU that included the 386 ancient South

American genomes described above, and an additional 107 ancient Iberian individuals

from Spain, Portugal and Gibraltar of the three investigated data types (Table S1). While

PC1 is driven by the ancestral difference between South American and Iberian

populations, as expected, PC2 is driven by the assay type and PC3 again represents

ancestry according to expected relationships (Fig 2A).

We then repeated this PCA but using smartPCA with outlier removal and no projection

(Fig 2B). While this PCA looks significantly different there is still a clear separation by

continent along PC1, assay along PC2 and by South American regional ancestry along

PC3. This strongly suggests that the allele frequency variation driving PC2 comes from

SNPs that are strongly affected by the assay bias, at least for this dataset. Hence, we

leveraged the SNP loadings outputted by smartPCA to construct a PCA-based filter that

removed all SNPs weighted > |1| from the dataset (n = 129,982 SNPs). After applying this

filter to the data and recomputing the PCA (Fig 2C), the 'assay component' is no longer

evident along PC2—which instead captures South American regional variation, as did

PC3 before the filtering (Fig 2B) — whereas PC3 is driven by variation among just a few

individuals. Lastly, we made use of the smartPCA projection functionality and projected

the PP samples onto the PCA generated from the non-PP samples (Fig 2D). This PCA

looks strikingly similar to the PCA constructed from the filtered data in Fig 2C.

Global PCA with smartPCA projection

To assess the effect of the enrichment bias on a global scale, we applied both EMU and

smartPCA projection methods to 612 ancient genomes from South and Central America,

Europe and Southwest Asia, either generated by shotgun sequencing or enriched using the

PP or legacy 1240k kits, onto a PCA of modern diversity that used 1,071 individuals from

the HGDP dataset (Bergström et al., 2020) (Fig 3).

When using EMU, the ancient individuals are projected with similar modern ancestries

(Fig 3, left). However when using smartPCA (Fig 3, right), the ancient samples are

projected into the correct quadrant of the PCA, i.e., clustering within the correct

continent, but data missingness seems to drive other discernible patterns, as both the

ancient American and Eurasian samples form clines from the plot origin to their

respective modern sample clusters located at either end of PC2. While PP samples are

situated closer to the PCA origin than the legacy 1240k samples, this can be explained by

the higher rate of data missingness in the PP dataset (see Fig S4).

Genetic relatedness

We ran READ for pairwise combinations of four individuals from Peru for which we had

generated shotgun data and single round PP-enriched data as a part of Experiment B (see

Methods). These individuals represent the only samples for which both shotgun data and

PP-enriched data were available. Given that only a single round of enrichment was used

in these PP assays, technical bias is expected to be reduced relative to double-round

enrichments.

The four artefactual "true twins" pairs (i.e. the same individual assayed with different

methods; e.g., IndA.PP.1rnd.B and IndA.Shotgun) were the most related of all

combinations with all average P0 values <0.4 (Fig S3). The remaining "non-twin" pairs all

had a P0 value of close to 0.5, with small increments between them. Notably, PP-enriched

pairs (e.g., Ind1.PP.1rnd.B and Ind2.PP.1rnd.B) were always more closely related (lower

P0) than the respective shotgun pairing (e.g., Ind1.Shotgun and Ind2.Shotgun) (Fig S3).

This suggests that the PP bias effects READ inferences, although it is subtle in this

instance.

Comparison of strategies to reduce bias

We conducted two wet-lab experiments with altered enrichment protocols and applied a

total of 19 bioinformatic strategies, aiming to find a strategy that best reduced the Prime

Plus bias. Bioinformatic strategies are categorised according to whether they 1) reduce

mapping reference bias, 2) reduce the impact of ancient DNA damage, 3) remove specific

"biased" SNPs, 4) employ statistical filters based on allele frequencies or 5) call genotypes

with alternate methods. Our test population comprised the ancient Peruvian samples, the

only population for which we generated both Prime Plus and shotgun data (Bongers et al.,

2020). This allowed us to compare the effect of bias-reduction strategies in an f_4 test

using statistics of the form f_4(Mbuti, Spain.PP.2rnd; Target, Peru.Shotgun), where the

target rotates between each of the different bias reduction strategies applied to the

Peruvian population. We chose Spain.PP.2rnd for this assessment for the following

reasons: i) the Spanish dataset was generated in a different lab, by different individuals,

and was sequenced on a different sequencing machine, than the Peruvian dataset (Table

S2), thereby ruling out batch effects and isolating the Prime Plus enrichment kit as the

underlying cause of any bias; and ii) the Spanish and Peruvian populations are sufficiently

divergent and lack evidence of recent admixture (the Peruvian data date to pre-colonial

times) such that the impact of Prime Plus bias is readily observable, but is not stronger

than the population affinities, as would be the case if we were comparing two more

closely related groups (such as populations from the same continent). To quantify the

impact of Prime Plus bias relative to population affinity between the Spanish and

Peruvian samples, we included Spain.1240k and Spain.Shotgun from publicly available

data as target populations.

In total, we calculated f_4 statistics in the following four configurations to robustly

estimate the Prime Plus bias and determine the best strategies to reduce it (Fig 4):

1) f_4(Mbuti, Spain.PP.2rnd; Target, Peru.Shotgun)

2) f_4(Mbuti, Spain.PP.2rnd; Target, Peru.1240k)

3) f_4(Mbuti, Spain.1240k; Target, Peru.Shotgun)

4) f_4(Mbuti, Spain.1240k; Target, Peru.1240k)

The reason for using the first two configurations is that by comparing the target

populations with Peruvian shotgun data, we are measuring our attempts to resolve the

Prime Plus bias against the least biased data currently available. However, this may be an

unreasonable standard to meet, given that it is known that legacy 1240k data also has

some biases (Rohland et al., 2022), and the bulk of published ancient DNA data is

legacy 1240k data. Therefore, while this comparison may not be a perfect estimation

of total bias reduction, it will provide more useful insights for researchers given that

most projects will leverage large proportions of 1240k reference data. Following the

same logic, in the third and fourth configurations, we also computed f_4 statistics with

Spanish legacy 1240k data (Fig 4), to gain a deeper understanding of the merits of

alternate strategies for reducing bias in different combinations of PP, legacy 1240k and

shotgun data.

As a positive control, we included a different set of ancient shotgun genomes from

Peru (Peru.other.Shotgun) as a Target population, which resulted in f_4 results that

were not significantly different from zero (Z < 3), as expected. The main aim of this

process was to identify a strategy that would behave like the positive control, i.e., that

would reduce the enrichment bias (Peru.PP.2rnd; dashed vertical line, Fig 4) so the f_4

values are not significantly different from 0. In the first configuration f_4(Mbuti,

Spain.PP.2rnd; Target, Peru.Shotgun), a result not statistically different from 0 would

indicate the bias was alleviated, and in the second configuration f_4(Mbuti,

Spain.PP.2rnd; Target, Peru.1240k), the same result would suggest that the PP data is at

least co-analysable with 1240k data under specific configurations. Importantly, the third

configuration f_4(Mbuti, Spain.1240k; Peru.PP.2rnd, Peru.Shotgun) (Fig 4) resulted in an

f_4 result that was not significantly different from zero, indicating that there was no

significant bias between these PP and legacy 1240k datasets. Finally, for the fourth

configuration f_4(Mbuti, Spain.1240k; Peru.PP.2rnd, Peru.1240k) (Fig 4) the f_4 result is

significantly positive, providing further evidence for the previously reported bias in the

legacy 1240k enrichment (Rohland et al., 2022). Therefore, the purpose of including

Spain.1240k among our tested configurations is to find a bias-reduction strategy that

does not significantly impact these f_4 statistics, and that allows the confident

application of bias reduction strategies to different combinations of the Prime Plus

and other data types and under alternate f_4 statistic configurations.

Ancient DNA damage or reference bias

The filter for minimum read length (Peru.PP.2rnd.>50bp) suggested by (Günther &

Nettelblad, 2019) to reduce mapping reference bias had minimal effect on the f_4

statistics, suggesting that the observed assay bias is not strongly impacted by very short

reads.

Filtering out transition SNPs (A/G and C/T), thus removing any possible ancient DNA

damage (Peru.PP.2rnd.transversion) had a negligible effect on the f_4 statistics scores.

Indeed, because all libraries were half-UDG treated and the ends of sequencing reads

trimmed, we did not anticipate that ancient DNA damage would have a large impact.

SNP-set based filters

Strategies to reduce bias by removing certain SNPs included filtering to include only the

Human Origins SNP set (Peru.PP.2rnd.HO; 593,120 SNPs retained), filtering according to

the SNPs driving PC2 (the "assay component" in Fig 2B, Peru.PP.PCAfilter; 486,445 SNPs

retained), and a filter previously reported (Rohland et al., 2022) to enable the co-analysis

of legacy 1240k and Prime Plus data (Peru.PP.2rnd.Rohland; 803,662 SNPs retained).

Retaining only the Human Origins SNPs or applying the PCA- based filter had minimal

effect on the f_4 results. Notably, the application of the Rohland filter did not alleviate

bias in any of the configurations of the f_4 statistics that we tested. In some cases

application of the Rohland filter resulted in the f_4 values shifting slightly toward zero,

but in other cases slightly away, which is likely due to an increase in uncertainty from the

approximately 2-fold reduction in the number of SNPs. Overall, this indicates the

Rohland filter does not work as reported on the present dataset, suggesting that it is not a

universally applicable solution for capture bias.

Allele frequency-based filters.

We further examined the bias reduction efficacy of a series of filters based on allele

frequencies, including filters for SNP missingness, minor allele frequency and linkage

disequilibrium. Of the SNP missingness filters, a strategy that excluded sites missing

in >50% of individuals (Peru.PP.2rnd.miss0.5) did not notably shift the f_4 value

towards zero. However, excluding all SNPs exhibiting any missingness

(Peru.PP.2rnd.miss0) did move the value toward zero, but it also reduced the number of

SNPs so drastically that this strategy would probably not be suitable for most

paleogenomic studies. For SNP filters based on minor allele frequency

(Peru.PP.2rnd.MAF0.3, Peru.PP.2rnd.MAF0.1 and Peru.PP.2rnd.MAF0.01), f_4 results

were shifted toward zero as the stringency of the filter increased, but this also led to

increasingly large reductions in the number of analysable SNPs. Interestingly, in the

f_4 configurations with Spain.1240k, the minor allele frequency filters produced

positive f_4 values indicating a bias towards the Peruvian data regardless of whether

shotgun or legacy 1240k are used.

When applying LD-based filters, we identified that the shared drift with Spain.PP.2rnd

reduces as the stringency of the filter increases, though this once again came at the cost

of increasingly fewer retained SNPs. In the configuration f_4(Mbuti, Spain.PP.2rnd;

Target, Peru.Shotgun), the f_4 result for $r^2 = 0.1$ was very close to 0 (Peru.PP.2rnd.LDr0.1).

Filters for less stringent r^2 values filtered out fewer SNPs but did not improve f_4 results to

the same degree (Fig 4). In the configuration f_4(Mbuti, Spain.PP; Target, Peru.1240k), the

less stringent $r^2 = 0.3$ produces an f_4 value not significantly different from 0, with the

retention of more SNPs than the $r^2 = 0.1$ filter. The use of this filter to alleviate PP bias

could be considered by users in specific cases with high sequencing depth to accommodate

the expected SNP loss. Randomly downsampling the Peru.PP data to an equivalent

number of SNPs (300,000) yielded f_4 values with additional variance around the value of

the unfiltered Peru.PP data (dashed lines, Fig 4) and higher absolute Z-scores (Fig 4). This

suggests that the bias may in fact be creating artificial "linkage" among neighbouring

SNPs, but that sufficiently stringent LD pruning to "correct" the value to zero requires

discarding the majority of targeted SNPs. Accordingly, the value of linkage disequilibrium

pruning may be considered as a potential solution to the PP bias on a case-by-case basis,

with the caveat that improved bias correction will be offset to some degree by the loss of

precision from reduced statistical power, and consideration will need to be given to the

effect of LD pruning on the results of various analyses.

Alternative genotyping

We considered several genotyping approaches alternative to the standard random pseudo-

haploidisation, including calling random diploid genotypes, using majority call pseudo-

haploid genotypes at sites with > 5X coverage, genotyping off-target SNPs, and genotyping

'out-of-range' SNPs that were > 100 bp from the target sites. Calling off-target genotypes

(Peru.PP.2rnd.OffTarget) and out-of-range genotypes

(Peru.PP.2rnd.>100bpOffTarget) both resulted in a small shift of the f_4 values toward

zero. Majority call genotypes (Peru.PP.2rnd.>5X.maj) behaved very differently between

configurations. In configurations with Spain.1240k, the filter had a minimal effect, but

with Spain.PP.2rnd, the majority call genotyping drastically shifted the f_4 results

towards negative values, indicating that this approach increases the artefactual

"shared drift" between PP data from Spain and Peru. Random diploid genotyping

produced an f_4 value that was even more biased (Peru.PP.2rnd.diploid), mildly so when

compared with Spain.1240k, but much more strongly for Spain.PP.2rnd. In fact, the

latter value is close to the f_4 result where the target dataset is Spain.Shotgun. This

means that using random diploid genotype calls exacerbates the artefactual shared

drift driven by the PP enrichment bias to approximately the same magnitude as true

shared drift between Spanish populations (Fig 4).

Laboratory experiments

In our final set of bias reduction strategies, two non-standard lab experiment protocols that used a single round of enrichment (Peru.PP.1rnd.A and Peru.PP.1rnd.B) both led to minor bias reductions. Protocol B, which lowered the hybridisation temperature performed slightly better than Protocol A. While the overall improvement exhibited by both methods was minor, the bias remained considerably greater than the bias observed in the legacy 1240k data (Peru.1240k).

Allele calls

We observe strong GC bias amongst the data generated with the legacy 1240k baits, whereas the PP data display comparatively lower levels of bias that are broadly comparable with shotgun datasets (Fig 5). It is important to note that the bias apparent in the legacy 1240k data was not a concern prior to the release of alternate enrichment kits, simply because all the data generated using the legacy 1240k baits are comparable. We observe that across all three methods, the GC bias is stronger for transitions—i.e., allele pairs A/G and C/T where both alleles are either purines or pyrimidines, respectively— which represent the majority of target sites (80%). Though our understanding of the biochemical mechanisms underlying the PP bias is limited by a lack of knowledge of the bait design, our data clearly demonstrate a systematic difference in the pattern of allelic biases between the legacy 1240k and PP assays.

DISCUSSION

Results presented in this study support the previous report of a bias when enriching SNPs at ~1240k sites using the Daicel Arbor Biosciences Expert Human Affinities Prime Plus enrichment kit, and extend the previous findings to ancient Indigenous American (i.e., non-Eurasian) populations. In accordance with (Rohland et al., 2022) we find that the biased data strongly affect f-statistic calculations, where the shared genetic drift (i.e., the evolutionary history) between groups is explicitly estimated. Note that we consider the effect strong, not because of the magnitude of any quantitative measurements, but due to the corruption of expected demographic relationships between populations that is driven by this bias. In contrast with results reported by (Rohland et al., 2022), we find that the Prime Plus (PP) bias does affect PCA when PP-enriched samples are included in the PCA calculation (i.e. not projected onto PCs computed from modern genomes), as might be expected given the close relationship between PCA and f-statistics (Peter, 2022). Additionally, for the first time, we observe a (in this case mild) bias effect on the predicted relatedness between individuals using the ancient DNA relatedness test READ.

Batch effects

The initial observation that PP data had an unexpected excess of shared drift with other PP-generated data in f-statistics prompted us to investigate if this artefact was due to a batch effect arising from certain samples, specific labs having done the work, DNA sequencing machines, or resulted from the distinct demographic history of populations in

question. We can confidently rule out each of these hypotheses, as the lab work was

performed on numerous sample sets, in different labs, by multiple individuals, sequenced

in different facilities using different sequencers (Table S2), with the only common factor

to all biased samples being the PP enrichment. Our outgroup-f_3 statistics calculated

between European and South American populations (Fig 1) confirmed that the bias

affected these two highly diverged populations suggesting that it is most likely a

systematic bias that affects all human ancestries. However, we did not have access to the

appropriate shotgun data to determine if the enrichment bias has differential effects by

ancestry source, and this hypothesis requires additional testing.

Describing the Prime Plus bias

PCA

We performed principal components analysis in several ways with the aim of addressing

multiple analytical approaches that researchers might take. To accommodate differing

geographical scales, we computed separate PCAs at continental and global scales. While

the approach of using smartPCA with projection has long been considered the gold

standard, it should be noted that this method works best with high-quality ancient

samples that have low missingness and a well-developed reference database of modern

diversity that captures the genetic variation present in the ancient populations, therefore

enabling the most accurate projection. Hence, the PCA published by Rohland et al. 2022

could be expected to show good results as they projected 15 high-quality ancient genomes

onto the modern diversity of western Eurasia, the most studied region of the world.

However, this is generally less achievable in understudied regions where modern

reference data is lacking and/or when working with low-quality samples. This is what

motivated the simultaneous use of EMU (designed to handle data missingness) (Meisner et

al., 2021) and smartPCA both with and without projection onto modern diversity.

At the continental scale, EMU produced a PCA that reflects South American population

structure in accordance with previous results (Nakatsuka, Lazaridis, et al., 2020;

Nakatsuka, Luisi, et al., 2020; Posth et al., 2018) and patterns suggesting assay bias are not

evident within the first three principal components (Fig S1). In contrast, PC2 in the

smartPCA (without projection) clearly distinguished data generated from the legacy

1240k enrichment from PP data (Fig S2), and exhibited patterns reflective of data

missingness (i.e., samples with more missingness being drawn toward the origin), which

was not apparent in any EMU PCA (Fig S2).

However, at the global scale, the PP bias was apparent in PCAs produced by both

smartPCA and EMU (Fig 2A and 2B, top row), although the resulting patterns look very

different. In both cases, the PP and legacy 1240k samples occupy largely distinct regions

of PCA space, although this divergence is less pronounced in South America than in

Iberia, suggesting that the bias may have a variable effect on different ancestries. Plots of

PC1 against PC3 (Fig 2A and 2B, bottom row) then reflect the clustering that we would

expect, separating South America from Iberia along PC1 and then the Andes from the

Southern Cone of South America along PC3. It is notable that the smartPCA (without

projection) output is still highly affected by missing data whereas this did not obviously

affect the EMU results.

The most striking discrepancy between the two PCA methods came from smartPCA's

projection of ancient genomes onto the modern diversity of the HGDP data (Bergström et

al., 2020). In the smartPCA output (Fig 3, right), while the shotgun genomes are in

accordance with previous results, both the legacy 1240k and PP samples are drawn toward

the origin, most likely due to missingness in the data (Fig S4). Similarly, the apparent

increased clustering of the PP samples relative to the legacy 1240k data may also be driven

by the higher rate of missingness amongst the PP data in this study (Fig S4). Ultimately,

these underlying issues mean that this PCA is not suitable for making demographic

inferences. In contrast, the EMU PCA places all ancient individuals within the proximity

of their associated modern population. Note that EMU does not have projection

functionality and instead includes all individuals in the calculation of eigenvectors while

imputing sites with missing data (Meisner et al., 2021). There does not seem to be any

obvious assay bias in the EMU PCA output, although this might still be present in a

lower-level PC.

Overall, we suggest that PCAs built with both smartPCA and EMU methods should be

produced and compared, especially in cases where PP data, or data with high rates of

missingness are included. PCAs are likely to be more robust at a smaller geographical

scale, and if computed from ancient samples, will benefit from including a majority of

non-PP samples to drive the assay bias component to a lower-level PC, thereby enabling

broad interpretations of the first two to three PCs.

Genetic relatedness

We assessed the effect of PP bias on results from the genetic relatedness estimation

software READ (Monroy Kuhn et al., 2018). We found that pairwise estimations between

PP libraries were always more closely related than shotgun library pairs of the same four

samples (Fig S3). Given that these PP libraries were only enriched for a single round, we

would expect this bias to be even stronger in PP libraries that follow the standard two-

round enrichment protocol (reflecting patterns observed in the f_4 statistics; Fig 4). In

the present case, the bias is subtle and unlikely to lead to, for example, a false-positive

first-degree relationship from an unrelated pair. While our ability to make more general

conclusions is clearly limited by the sample set, our results tentatively suggest that READ

inferences should be reasonably robust if run on a dataset of only PP data, but users

should exhibit caution when combining different data types.

f-statistics

Our analyses of the PP bias in both outgroup-f_3 and f_4 statistics showed that the bias is

strong enough to corrupt expected relationships between populations from the same

continent (Fig 1). We reproduced the legacy 1240k bias that has been previously reported

(Rohland et al., 2022) and showed that it is relatively subtle compared to the PP bias,

indicating that it is less impactful on population genetic inferences.

Importantly, our analyses suggest that it should be possible to use Prime Plus data in

a comparative outgroup-f_3 if the target population is the only one enriched with Prime

Plus, that is statistics of the form f_3(Outgroup.notPP; target.PP, X.notPP) where both

the outgroup and population X are not enriched with Prime Plus. This is suggested by

our f_3 statistics (Fig 1A) where Spain.PP.2rnd is the Target population, as the three

European populations reproduce the expected relative relationships previously reported

(Martiniano et al., 2017; Olalde et al., 2019; Valdiosera et al., 2018) despite none having

been enriched with the Prime Plus baits. However, the limitation is that a population

enriched with Prime Plus could not be split into subpopulations, such as Spain_A.PP.2rnd

and Spain_B.PP.2rnd, and rotated in the X population position, or in scenarios where

one is X and the other Test, but then compared to f_3 statistics of a different form.

Presumably, an f-statistic would also be viable if all X groups were PP-enriched because

the enrichment bias would be shared by all data and should not corrupt demographic

interpretations of the results. See Table S4 for a comprehensive list of acceptable

outgroup-f_3 and f_4 statistic forms and, though not directly investigated here, using the

same logic we also list qpWave and qpAdm configurations and their relative reliability

under PP bias.

Strategies to reduce bias

The bias observed in f-statistics is both persistent and systematic, which allowed us to

test a total of 21 experimental and analytical protocols to characterise the bias and

alleviate its disruptive impact on inferences from common population genetic analyses.

Many of these analytical protocols produced marginal improvements, but no protocol was

sufficient to remove all biases (Z-score remained such that $|Z| > 3$, and/or resulted in low

SNP retention).

711   *Ancient DNA damage or reference bias*

712   Neither ancient DNA damage nor reference bias appeared to contribute to the bias, as

713   removing transition sites (all SNPs that could potentially be ancient DNA damage)

714   and removing reads shorter than 50 bp had little impact on bias in the f_4 statistics (Fig

715   4).

717   *SNP-set based filters*

Next, we investigated if a subset of the 1240k SNPs was driving the bias and could

therefore be removed to allow safe co-analysis of PP enriched samples with other data

types. This idea was initially suggested by Rohland and colleagues (Rohland et al., 2022),

who published a list of SNPs that they suggested should be suitable for co-analyses of

shotgun, legacy 1240k, Twist, and PP data. However, our tests of the Rohland et al. filter

produced no notable improvement on the f_4 statistics when co-analysed with 1240k or PP

data (Peru.PP.2rnd.Rohland, Fig 4). Note that to build this filter Rohland and colleagues

looked for improved Z-scores, i.e., closer to zero, but did not report the f_4 statistics that

are informative to estimate the effect on population relatedness. When using any SNP

filter, the removal of SNPs will likely increase the standard error of the f_4 statistic,

leading to decreased statistical power and precision. However, it is not possible to

predict the deviation from the true f_4 statistic. Therefore Z-scores alone do not serve as a

measure of the bias-reduction of a filtering strategy. This emphasises the importance of

testing hypotheses in the context of the study and understanding the expected

relationships of the populations in question.

Subsetting our data to include only the ˜600k SNPs in the Human Origins set had a

similar effect to the Rohland filter. This may be related to the overlap between the two

sets, 61% of the Human Origins SNPs are also present in the Rohland filter set. (367,332

intersecting SNPs).

Additionally, we generated our own list of SNPs to filter by extracting the SNPs that were

driving variation evident in PC2, the "assay component" in the smartPCA output (Fig 2B).

Although this filtering approach did seem to remove the biased component from the

PCA (Fig 2C), it had a negligible impact on any of the f_4 statistics. In summary, none of

these filtering strategies corrected the PP bias satisfactorily. Furthermore, while Rohland

et al. acknowledge there is room for improvement in their filtering approach, it is possible

that their filter is well-calibrated for the Eurasian ancestries present in their dataset but

does not represent a universal filter to mitigate Prime Plus capture bias in all ancestries. In

any case, it is less than ideal to have enrichment kits on the market that would require

discarding a subset of target SNPs when combining multiple data types. This will

negatively affect not only users of the Daicel Arbor Biosciences Human Affinities kits —

and potentially other commercial kits — but also impact subsequent analyses by other

researchers that incorporate published PP datasets.

Alternative genotyping

Genotype calling in paleogenomic studies typically involves randomly sampling one read

covering a targeted SNP site, known as 'pseudo-haploidisation', making it possible to

generate allele calls at sites with low coverage (Morris et al., 2011). Given that our

experiments with SNP filtering did not produce a universal solution and following the

assumption that every targeted SNP site may be affected by the bias, we investigated the

efficacy of alternate genotype calling for bias reduction. In our first assessment, we called

pseudo-haploid genotypes at off-target SNPs, which we hypothesised would be equivalent

to shotgun data and therefore unbiased if the bias was driven by the bait design. Similarly,

we also called random pseudo-haploid genotypes at sites more than 100 bp away from a

target locus to mitigate the possibility of an "off-target" site being covered by a probe that

had correctly hybridised to a nearby target locus. The f_4 statistic for both off-target and

out-of-range genotypes did improve compared to the on-target results, producing

values that were not significantly different from zero (Fig 4). Interestingly, the f_4

statistic for the out-of-range genotypes was not notably different from the off-target

estimation. Given that all non-bound DNA fragments are theoretically washed away

during the enrichment protocol, the off-target bait bias is likely due to an as-yet-

unknown biochemical bias that systematically affects bait binding to DNA at non-target

locations, but it is unlikely this is caused by on-target hybridised baits covering nearby

SNPs within a 100 bp range.

Additionally, we used an alternate pseudo-haploid calling approach which requires

target SNPs to have a higher minimum coverage (5X) and takes the allele

represented by the majority of reads as the final call. This experiment

(Peru.PP.2rnd.>5X.maj, Fig 4) yielded a dramatic shift of the f_4 results towards large

negative values, further exacerbating the artefactual shared drift and bias. This is strong

evidence that the bias is conferred by the bait design, and results in the persistent

overrepresentation of one allele over the other at heterozygous sites. Concordantly, our

random diploid genotype test produced even more biased f_4 statistics, such that the

estimated levels of shared drift observed between the Spanish PP data and Peruvian PP

data that used random diploid genotype calls is of a similar magnitude to the true shared

drift between Spanish PP generated and Spanish shotgun data. This is evidence of a

remarkably strong systematic bias that has the potential to corrupt all paleogenomic

studies combining PP with other data types if not handled carefully. These two results are

strongly suggestive of a systematic allelic bias that impacts all heterozygous SNPs and is

potentially caused by the molecular design of the PP baits.

Laboratory protocol experiments

Our two experiments with single-round PP enrichment yielded f_4 values closer to zero

than those produced by the standard two-round enrichment. This improvement is likely

because whatever bias occurs during the first round of bait–DNA binding becomes

amplified during the second hybridisation step because it is acting on an already biased set

of DNA fragments. This interpretation is supported by the sequencing statistics (Fig S5)

showing that repeated capture rounds result in an increased endogenous DNA percentage

but decreased library complexity (observed as fewer unique reads when sequenced).

Additionally, any biased sampling that has occurred during the initial binding will be

exponentially amplified during the PCR step, so we would expect a reduction in PCR

cycles to improve this. The reduced hybridisation temperature should also decrease the

specificity of hybridisation and thus reduce allelic bias, but simultaneously increase off-

target binding, thus reducing the percentage of endogenous DNA enriched.

Characterising the bias

Given the evidence that the bias affects all targeted SNPs, we investigated whether this

effect was independent from the SNP transition, hoping that we might be able to apply

some relatively simple statistical scaling of the genotype calls that would provide co-

analysable genotypes. However, our investigations of allelic bias (Fig 5) clearly show that

the capture bias produces complex effects on allele calls. We can only confidently observe

that the pattern of GC bias in PP data is different to the 1240k and shotgun data. Further,

the patterns of missingness evident in the PCA analyses consistently differ between data

types, indicating that there may be several layers of biochemical bias at play including

that each assay preferentially captures certain SNPs, and/or favours the capture of

different alleles at heterozygous sites, with allelic preference being more exacerbated in

PP data.

Study limitations

The main limitation of this study is that we did not have samples for which data was

generated in all of the three methods addressed. This is due to the opportunistic nature of

the study in that we collated this dataset when the PP bias was initially observed, after the

data generation process. However, we have made every effort to combat this limitation by

maximising the size of the dataset and explicitly stating assumptions where we've made

them.

In our READ analyses we were limited to a dataset of only four individuals for which we

had shotgun and PP data, and a further limitation is that this PP data was from one of our

single-round enrichment experiments and thus does not fully represent the PP bias under

investigation. To further understand the impact on READ, both shotgun and two-round

PP enriched data would need to be generated for a larger number of samples, from a more

diverse set of relationships to test the impact of bias on truly related individuals.

The main outstanding hypothesis to investigate is whether this bias has differential effects

on different ancestries. This would require a large dataset of PP and shotgun data from the

same individuals across more ancestries than in the present study. While an interesting

question, we believe this study presents sufficient evidence that the PP assay is biased.

Implications

For the past decade in-solution hybridisation enrichment has been a go-to method for

obtaining affordable, high-quality, and informative data for paleogenomic studies.

However, irrespective of the bait design, the biochemistry of DNA hybridisation means

that one bait will always have a higher binding affinity with its targeted allele than any

other. Greater differences in these affinities result in increased allelic biases in the

generated datasets. This is evidenced by our assessment of alternative enrichment

protocols, where the single-round enrichments have less endogenous DNA (Fig S3), but

also appear to be less biased (Fig 4). Therefore, we hypothesise that the enrichment

efficiency of the PP reagent is in direct trade-off with the biochemical allelic bias.

It is worth reiterating here that all capture methods used in paleogenomic studies will lead

to ascertainment bias, meaning that one must carefully consider both the selection of

methods and the use of comparative published data. Previously, the monopoly on data

generation by labs with access to the legacy 1240k enrichment protocol somewhat

insulated the community from this compatibility issue. However, now that data can be

generated using different enrichment kits, researchers face a new set of challenges

whenever they integrate data generated with new capture assays with previously

published data generated using the legacy 1240k—the most comprehensive dataset to date

that underpins many of the findings in paleogenomics. Additionally, researchers utilising

the Twist reagent should benchmark the data similarly to the present study to establish its

co-analysability with other data types.

Here, we have tested some of the most used analyses in paleogenomic studies,

though not an exhaustive list. Principal Components Analysis is a standard way to

understand population structure, and outgroup-f_3 and f_4 statistics, along with their

more complex derivatives, are integral to reconstructing population relationships and

evolutionary history, while READ is a standard for genetic relatedness analysis. We have

found that PP-enriched data bias the results of all these analyses when used in

combination with other non-PP datasets, and therefore we caution potential PP users that

analyses of such multi-assayed datasets may lead to erroneous conclusions if not

interpreted carefully.

Having comparative shotgun data for a subset of samples in a study, or from samples

expected to have a very similar ancestry should allow researchers to estimate and control

for the PP bias in the specific methods they wish to utilise. Further, while we explicitly

test PP data, we advise researchers to exhibit similar caution when using the Daicel Arbor

Biosciences Expert Human Affinities "Complete" and "Ancestral Plus" enrichment kits, as

the bait chemistry and enrichment protocol is the same as the PP kit, and only the

number of target loci differ.

Guidelines

- When calculating f-statistics only compare Prime-Plus-generated data with other

Prime-Plus-generated data; OR

- Only compare one Prime-Plus-generated target population/individual with other

non-Prime-Plus-generated populations/individuals (See Table S4 for a full

description).

- Beware that Prime-Plus-generated data attract each other and that this may

artificially confirm/reject your hypothesis depending on the study design.

- Explicitly assess the impact of Prime-Plus bias in your data when running analyses

that have not yet been investigated.

- Recommend to shotgun sequence a subset of enriched libraries as a control set if possible, ideally select libraries to represent a range of ancestries.
- Always annotate your data as Prime-Plus-generated when publishing, sharing with other researchers, or incorporating into any public database.
- Always annotate the data type of both newly generated and published data used in your study so this can be factored into interpretations.
- For best results compute PCA using EMU and smartPCA and compare results; if clustering appears driven by capture assay, pursue alternative analyses.
- In general, pursue multiple lines of evidence to support your findings.

Acknowledgements

We thank Adam Rohrlach for insightful manuscript revisions and assistance with statistical interpretation. We thank members of the Australian Centre for Ancient DNA's Thesis Writing Group for their contributions to manuscript revisions, especially Olivia Johnson for assistance with R coding. We thank members of the Unitat Antropologia Biològica at the Universitat Autònoma de Barcelona (Spain) for their contribution to data generation. LFS was supported by funding from the National Science Foundation (NSF - 1515138), RT by the Australian Research Council (DE190101069), BL by an Australian Research Council Future Fellowship (FT170100448).

References

Ávila-Arcos, M. C., Sandoval-Velasco, M., Schroeder, H., Carpenter, M. L., Malaspinas, A.-S.,
Wales, N., Peñaloza, F., Bustamante, C. D., & Gilbert, M. T. P. (2015). Comparative
performance of two whole-genome capture methodologies on ancient DNA Illumina
libraries. *Methods in Ecology and Evolution / British Ecological Society*, *6*(6), 725–734.
Bergström, A., McCarthy, S. A., Hui, R., Almarri, M. A., Ayub, Q., Danecek, P., Chen, Y., Felkel,
S., Hallast, P., Kamm, J., Blanché, H., Deleuze, J.-F., Cann, H., Mallick, S., Reich, D., Sandhu,
M. S., Skoglund, P., Scally, A., Xue, Y., … Tyler-Smith, C. (2020). Insights into human
genetic variation and population history from 929 diverse genomes. *Science* , *367*(6484),
eaay5012.
Boessenkool, S., Hanghøj, K., Nistelberger, H. M., Der Sarkissian, C., Gondek, A. T., Orlando, L.,
Barrett, J. H., & Star, B. (2017). Combining bleach and mild predigestion improves ancient
DNA recovery from bones. *Molecular Ecology Resources*, *17*(4), 742–751.
Bongers, J. L., Nakatsuka, N., O'Shea, C., Harper, T. K., Tantaleán, H., Stanish, C., & Fehren-
Schmitz, L. (2020). Integration of ancient DNA with transdisciplinary dataset finds strong
support for Inca resettlement in the south Peruvian coast. *Proceedings of the National*
Academy of Sciences, *117*(31), 18359–18368;
Briggs, A. W., Good, J. M., Green, R. E., Krause, J., Maricic, T., Stenzel, U., Lalueza-Fox, C.,
Rudan, P., Brajkovic, D., Kucan, Z., Gusic, I., Schmitz, R., Doronichev, V. B., Golovanova, L.
V., de la Rasilla, M., Fortea, J., Rosas, A., & Pääbo, S. (2009). Targeted retrieval and analysis of
five Neandertal mtDNA genomes. *Science*, *325*(5938), 318–321.
Brotherton, P., Haak, W., Templeton, J., Brandt, G., Soubrier, J., Jane Adler, C., Richards, S. M.,
Sarkissian, C. D., Ganslmeier, R., Friederich, S., Dresely, V., van Oven, M., Kenyon, R., Van
der Hoek, M. B., Korlach, J., Luong, K., Ho, S. Y. W., Quintana-Murci, L., Behar, D. M., …
Cooper, A. (2013). Neolithic mitochondrial haplogroup H genomes and the genetic origins of
Europeans. *Nature Communications*, *4*, 1764.
Carpenter, M. L., Buenrostro, J. D., Valdiosera, C., Schroeder, H., Allentoft, M. E., Sikora, M.,
Rasmussen, M., Gravel, S., Guillén, S., Nekhrizov, G., Leshtakov, K., Dimitrova, D.,
Theodossiev, N., Pettener, D., Luiselli, D., Sandoval, K., Moreno-Estrada, A., Li, Y., Wang, J.,
… Bustamante, C. D. (2013). Pulling out the 1%: Whole-genome capture for the targeted
enrichment of ancient DNA sequencing libraries. *American Journal of Human Genetics*,
93(5), 852–864.
Castellano, S., Parra, G., Sánchez-Quinto, F. A., Racimo, F., Kuhlwilm, M., Kircher, M., Sawyer,
S., Fu, Q., Heinze, A., Nickel, B., Dabney, J., Siebauer, M., White, L., Burbano, H. A., Renaud,
G., Stenzel, U., Lalueza-Fox, C., Rasilla, M. de la, Rosas, A., … Pääbo, S. (2014). Patterns of
coding variation in the complete exomes of three Neandertals. *Proceedings of the National*
Academy of Sciences of the United States of America, *111*(18), 6666–6671.
Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-
generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*, *4*, 7.
Cruz-Dávalos, D. I., Nieves-Colón, M. A., Sockell, A., Poznik, G. D., Schroeder, H., Stone, A. C.,
Bustamante, C. D., Malaspinas, A.-S., & Ávila-Arcos, M. C. (2018). In-solution Y-
chromosome capture-enrichment on ancient DNA libraries. *BMC Genomics*, *19*.
https://doi.org/10.1186/s12864-018-4945-x
Dabney, J., Knapp, M., Glocke, I., Gansauge, M.-T., Weihmann, A., Nickel, B., Valdiosera, C.,
García, N., Pääbo, S., Arsuaga, J.-L., & Meyer, M. (2013). Complete mitochondrial genome
sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments.
Proceedings of the National Academy of Sciences of the United States of America, *110*(39),

15758–15763.

de la Fuente, C., Ávila-Arcos, M. C., Galimany, J., Carpenter, M. L., Homburger, J. R., Blanco, A.,
Contreras, P., Cruz Dávalos, D., Reyes, O., San Roman, M., Moreno-Estrada, A., Campos, P.
F., Eng, C., Huntsman, S., Burchard, E. G., Malaspinas, A.-S., Bustamante, C. D., Willerslev,
E., Llop, E., … Moraga, M. (2018). Genomic insights into the origin and diversification of late
maritime hunter-gatherers from the Chilean Patagonia. *Proceedings of the National Academy*
of Sciences of the United States of America, *115*(17), E4006–E4012.

Fellows Yates, J. A., Lamnidis, T. C., Borry, M., Andrades Valtueña, A., Fagernäs, Z., Clayton, S.,
Garcia, M. U., Neukamm, J., & Peltzer, A. (2021). Reproducible, portable, and efficient
ancient genome reconstruction with nf-core/eager. *PeerJ*, *9*(e10947), e10947.

Fumagalli, M. (2013). Assessing the effect of sequencing depth and sample size in population
genetics inferences. *PloS One*, *8*(11), e79667.

Fu, Q., Hajdinjak, M., Moldovan, O. T., Constantin, S., Mallick, S., Skoglund, P., Patterson, N.,
Rohland, N., Lazaridis, I., Nickel, B., Viola, B., Prüfer, K., Meyer, M., Kelso, J., Reich, D., &
Pääbo, S. (2015). An early modern human from Romania with a recent Neanderthal ancestor.
Nature, *524*(7564), 216–219.

Fu, Q., Meyer, M., Gao, X., Stenzel, U., Burbano, H. A., Kelso, J., & Pääbo, S. (2013). DNA analysis
of an early modern human from Tianyuan Cave, China. *Proceedings of the National Academy*
of Sciences, *110*(6), 2223–2227.

González-Fortes, G., Tassi, F., Trucchi, E., Henneberger, K., Paijmans, J. L. A., Díez-Del-Molino,
D., Schroeder, H., Susca, R. R., Barroso-Ruíz, C., Bermudez, F. J., Barroso-Medina, C.,
Bettencourt, A. M. S., Sampaio, H. A., Grandal-d'Anglade, A., Salas, A., de Lombera-Hermida,
A., Fabregas Valcarce, R., Vaquero, M., Alonso, S., … Barbujani, G. (2019). A western route of
prehistoric human migration from Africa into the Iberian Peninsula. *Proceedings. Biological*
Sciences / The Royal Society, *286*(1895), 20182288.

Günther, T., & Nettelblad, C. (2019). The presence and impact of reference bias on population
genomic studies of prehistoric human populations. *PLoS Genetics*, *15*(7), e1008302.

Günther, T., Valdiosera, C., Malmström, H., Ureña, I., Rodriguez-Varela, R., Sverrisdóttir, Ó. O.,
Daskalaki, E. A., Skoglund, P., Naidoo, T., Svensson, E. M., Bermúdez de Castro, J. M.,
Carbonell, E., Dunn, M., Storå, J., Iriarte, E., Arsuaga, J. L., Carretero, J.-M., Götherström, A.,
& Jakobsson, M. (2015). Ancient genomes link early farmers from Atapuerca in Spain to
modern-day Basques. *Proceedings of the National Academy of Sciences of the United States*
of America, *112*(38), 11917–11922.

Haak, W., Lazaridis, I., Patterson, N., Rohland, N., Mallick, S., Llamas, B., Brandt, G., Nordenfelt,
S., Harney, E., & Stewardson, K. (2015). Massive migration from the steppe was a source for
Indo-European languages in Europe. *Nature*, *522*(7555), 207.

Kapp, J. D., Green, R. E., & Shapiro, B. (2021). A fast and efficient single-stranded genomic library
preparation method optimized for ancient DNA. *The Journal of Heredity*, *112*(3), 241–249.

Krause, J., Briggs, A. W., Kircher, M., Maricic, T., Zwyns, N., Derevianko, A., & Pääbo, S. (2010).
A complete mtDNA genome of an early modern human from Kostenki, Russia. *Current*
Biology: CB, *20*(3), 231–236.

Lindo, J., Haas, R., Hofman, C., Apata, M., Moraga, M., Verdugo, R. A., Watson, J. T., Viviano
Llave, C., Witonsky, D., Beall, C., Warinner, C., Novembre, J., Aldenderfer, M., & Di Rienzo,
A. (2018). The genetic prehistory of the Andean highlands 7000 years BP though European
contact. *Science Advances*, *4*(11), eaau4921.

Lipson, M., Szécsényi-Nagy, A., Mallick, S., Pósa, A., Stégmár, B., Keerl, V., Rohland, N.,

Stewardson, K., Ferry, M., Michel, M., Oppenheimer, J., Broomandkhoshbacht, N., Harney, E., Nordenfelt, S., Llamas, B., Gusztáv Mende, B., Köhler, K., Oross, K., Bondár, M., … Reich, D. (2017). Parallel palaeogenomic transects reveal complex genetic history of early European farmers. *Nature*, *551*(7680), 368–372.

Llamas, B., Fehren-Schmitz, L., Valverde, G., Soubrier, J., Mallick, S., Rohland, N., Nordenfelt, S., Valdiosera, C., Richards, S. M., Rohrlach, A., Romero, M. I. B., Espinoza, I. F., Cagigao, E. T., Jiménez, L. W., Makowski, K., Reyna, I. S. L., Lory, J. M., Torrez, J. A. B., Rivera, M. A., … Haak, W. (2016). Ancient mitochondrial DNA provides high-resolution time scale of the peopling of the Americas. *Science Advances*, *2*(4), e1501385.

Mallick, S., Li, H., Lipson, M., Mathieson, I., Gymrek, M., Racimo, F., Zhao, M., Chennagiri, N., Nordenfelt, S., Tandon, A., Skoglund, P., Lazaridis, I., Sankararaman, S., Fu, Q., Rohland, N., Renaud, G., Erlich, Y., Willems, T., Gallo, C., … Reich, D. (2016). The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature*, *538*(7624), 201–206.

Marciniak, S., & Perry, G. H. (2017). Harnessing ancient genomes to study the history of human adaptation. *Nature Reviews. Genetics*, *18*(11), 659–674.

Maricic, T., Whitten, M., & Pääbo, S. (2010). Multiplexed DNA sequence capture of mitochondrial genomes using PCR products. *PloS One*, *5*(11), e14004.

Martiniano, R., Cassidy, L. M., Ó'Maoldúin, R., McLaughlin, R., Silva, N. M., Manco, L., Fidalgo, D., Pereira, T., Coelho, M. J., Serra, M., Burger, J., Parreira, R., Moran, E., Valera, A. C., Porfirio, E., Boaventura, R., Silva, A. M., & Bradley, D. G. (2017). The population genomics of archaeological transition in west Iberia: Investigation of ancient substructure using imputation and haplotype-based methods. *PLoS Genetics*, *13*(7), e1006852.

Mathieson, I., Lazaridis, I., Rohland, N., Mallick, S., Patterson, N., Roodenberg, S. A., Harney, E., Stewardson, K., Fernandes, D., & Novak, M. (2015). Genome-wide patterns of selection in 230 ancient Eurasians. *Nature*, *528*(7583), 499–503.

Meisner, J., Liu, S., Huang, M., & Albrechtsen, A. (2021). Large-scale Inference of Population Structure in Presence of Missingness using PCA. *Bioinformatics* . https://doi.org/10.1093/bioinformatics/btab027

Meyer, M., Kircher, M., Gansauge, M.-T., Li, H., Racimo, F., Mallick, S., Schraiber, J. G., Jay, F., Pruefer, K., de Filippo, C., Sudmant, P. H., Alkan, C., Fu, Q., Do, R., Rohland, N., Tandon, A., Siebauer, M., Green, R. E., Bryc, K., … Pääbo, S. (2012). A high-coverage genome sequence from an archaic Denisovan individual. *Science*, *338*(6104), 222–226.

Monroy Kuhn, J. M., Jakobsson, M., & Günther, T. (2018). Estimating genetic kin relationships in prehistoric populations. *PloS One*, *13*(4), e0195491.

Moreno-Mayar, J. V., Vinner, L., de Barros Damgaard, P., de la Fuente, C., Chan, J., Spence, J. P., Allentoft, M. E., Vimala, T., Racimo, F., Pinotti, T., Rasmussen, S., Margaryan, A., Iraeta Orbegozo, M., Mylopotamitaki, D., Wooller, M., Bataille, C., Becerra-Valdivia, L., Chivall, D., Comeskey, D., … Willerslev, E. (2018). Early human dispersals within the Americas. *Science*, *362*(6419), eaav2621.

Morris, G. P., Grabowski, P. P., & Borevitz, J. O. (2011). Genomic diversity in switchgrass (Panicum virgatum): from the continental scale to a dune landscape. *Molecular Ecology*, *20*(23), 4938–4952.

Nakatsuka, N., Lazaridis, I., Barbieri, C., Skoglund, P., Rohland, N., Mallick, S., Posth, C., Harkins-Kinkaid, K., Ferry, M., Harney, É., Michel, M., Stewardson, K., Novak-Forst, J., Capriles, J. M., Alfonso Durruty, M., Aranda Álvarez, K., Beresford-Jones, D., Burger, R. L., Cadwallader, L., … Fehren-Schmitz, L. (2020). A paleogenomic reconstruction of the deep population

history of the Andes. *Cell*, *181*(5), 1131–1145.e21.

Nakatsuka, N., Luisi, P., Motti, J. M. B., Salemme, M., Santiago, F., D'Angelo Del Campo, M. D.,
Vecchi, R. J., Espinosa-Parrilla, Y., Prieto, A., Adamski, N., Lawson, A. M., Harper, T. K.,
Culleton, B. J., Kennett, D. J., Lalueza-Fox, C., Mallick, S., Rohland, N., Guichón, R. A.,
Cabana, G. S., … Reich, D. (2020). Ancient genomes in South Patagonia reveal population
movements associated with technological shifts and geography. *Nature Communications*,
11(1), 3868.

Olalde, I., Brace, S., Allentoft, M. E., Armit, I., Kristiansen, K., Booth, T., Rohland, N., Mallick, S.,
Szécsényi-Nagy, A., Mittnik, A., Altena, E., Lipson, M., Lazaridis, I., Harper, T. K., Patterson,
N., Broomandkhoshbacht, N., Diekmann, Y., Faltyskova, Z., Fernandes, D., … Reich, D.
(2018). The Beaker phenomenon and the genomic transformation of northwest Europe.
Nature, *555*(7695), 190–196.

Olalde, I., Mallick, S., Patterson, N., Rohland, N., Villalba-Mouco, V., Silva, M., Dulias, K.,
Edwards, C. J., Gandini, F., Pala, M., Soares, P., Ferrando-Bernal, M., Adamski, N.,
Broomandkhoshbacht, N., Cheronet, O., Culleton, B. J., Fernandes, D., Lawson, A. M., Mah,
M., … Reich, D. (2019). The genomic history of the Iberian Peninsula over the past 8000
years. *Science*, *363*(6432), 1230–1234.

Olalde, I., & Posth, C. (2020). Latest trends in archaeogenetic research of west Eurasians. *Current*
Opinion in Genetics & Development, *62*, 36–43.

Oliva, A., Tobler, R., Cooper, A., Llamas, B., & Souilmi, Y. (2021). Systematic benchmark of
ancient DNA read mapping. *Briefings in Bioinformatics*, *22*(5), bbab076.

Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T., Webster,
T., & Reich, D. (2012). Ancient admixture in human history. *Genetics*, *192*(3), 1065–1093.

Patterson, N., Price, A. L., & Reich, D. (2006). Population structure and eigenanalysis. *PLoS*
Genetics, *2*(12), e190.

Peter, B. M. (2022). A geometric relationship of F2, F3 and F4-statistics with principal component
analysis. *Philosophical Transactions of the Royal Society of London. Series B, Biological*
Sciences, *377*(1852), 20200413.

Popović, D., Molak, M., Ziółkowski, M., Vranich, A., Sobczyk, M., Vidaurre, D. U., Agresti, G.,
Skrzypczak, M., Ginalski, K., & Lamnidis, T. C. (2021). Ancient genomes reveal long-range
influence of the pre-Columbian culture and site of Tiwanaku. *Science Advances*, *7*(39),
eabg7261.

Posth, C., Nakatsuka, N., Lazaridis, I., Skoglund, P., Mallick, S., Lamnidis, T. C., Rohland, N.,
Nägele, K., Adamski, N., & Bertolini, E. (2018). Reconstructing the deep population history of
Central and South America. *Cell*, *175*(5), 1185–1197. e22.

Price, A. L., Patterson, N. J., Plenge, R. M., Weinblatt, M. E., Shadick, N. A., & Reich, D. (2006).
Principal components analysis corrects for stratification in genome-wide association studies.
Nature Genetics, *38*(8), 904–909.

Prüfer, K., Racimo, F., Patterson, N., Jay, F., Sankararaman, S., Sawyer, S., Heinze, A., Renaud, G.,
Sudmant, P. H., de Filippo, C., Li, H., Mallick, S., Dannemann, M., Fu, Q., Kircher, M.,
Kuhlwilm, M., Lachmann, M., Meyer, M., Ongyerth, M., … Pääbo, S. (2014). The complete
genome sequence of a Neanderthal from the Altai Mountains. *Nature*, *505*(7481), 43–49.

Raghavan, M., Steinrücken, M., Harris, K., Schiffels, S., Rasmussen, S., DeGiorgio, M.,
Albrechtsen, A., Valdiosera, C., Ávila-Arcos, M. C., Malaspinas, A.-S., Eriksson, A., Moltke,
I., Metspalu, M., Homburger, J. R., Wall, J., Cornejo, O. E., Víctor Moreno-Mayar, J.,
Korneliussen, T. S., Pierre, T., … Willerslev, E. (2015). Genomic evidence for the Pleistocene

and recent population history of Native Americans. *Science*.

https://doi.org/10.1126/science.aab3884

Rohland, N., Harney, E., Mallick, S., Nordenfelt, S., & Reich, D. (2015). Partial uracil-DNA-

glycosylase treatment for screening of ancient DNA. *Philosophical Transactions of the Royal*

Society of London. Series B, Biological Sciences, *370*(1660), 20130624.

Rohland, N., Mallick, S., Mah, M., Maier, R., Patterson, N., & Reich, D. (2022). Three assays for

in-solution enrichment of ancient human DNA at more than a million SNPs. *Genome*

Research. https://doi.org/10.1101/gr.276728.122

Rohrlach, A. B., Papac, L., Childebayeva, A., Rivollat, M., Villalba-Mouco, V., Neumann, G. U.,

Penske, S., Skourtanioti, E., van de Loosdrecht, M., Akar, M., Boyadzhiev, K., Boyadzhiev, Y.,

Deguilloux, M.-F., Dobeš, M., Erdal, Y. S., Ernée, M., Frangipane, M., Furmanek, M.,

Friederich, S., … Haak, W. (2021). Using Y-chromosome capture enrichment to resolve

haplogroup H2 shows new evidence for a two-path Neolithic expansion to Western Europe.

Scientific Reports, *11*(1), 15005.

Sherry, S. T., Ward, M. H., Kholodov, M., Baker, J., Phan, L., Smigielski, E. M., & Sirotkin, K.

(2001). dbSNP: the NCBI database of genetic variation. *Nucleic Acids Research*, *29*(1), 308–

311.

Skoglund, P., & Mathieson, I. (2018). Ancient Genomics of Modern Humans: The First Decade.

Annual Review of Genomics and Human Genetics, *19*(1), 381–404.

Valdiosera, C., Günther, T., Vera-Rodríguez, J. C., Ureña, I., Iriarte, E., Rodríguez-Varela, R.,

Simões, L. G., Martínez-Sánchez, R. M., Svensson, E. M., Malmström, H., Rodríguez, L.,

Bermúdez de Castro, J.-M., Carbonell, E., Alday, A., Hernández Vera, J. A., Götherström, A.,

Carretero, J.-M., Arsuaga, J. L., Smith, C. I., & Jakobsson, M. (2018). Four millennia of Iberian

biomolecular prehistory illustrate the impact of prehistoric migrations at the far end of

Eurasia. *Proceedings of the National Academy of Sciences of the United States of America*,

115(13), 3428–3433.

Villalba-Mouco, V., van de Loosdrecht, M. S., Posth, C., Mora, R., Martínez-Moreno, J., Rojo-

Guerra, M., Salazar-García, D. C., Royo-Guillén, J. I., Kunst, M., Rougier, H., Crevecoeur, I.,

Arcusa-Magallón, H., Tejedor-Rodríguez, C., García-Martínez de Lagrán, I., Garrido-Pena, R.,

Alt, K. W., Jeong, C., Schiffels, S., Utrilla, P., … Haak, W. (2019). Survival of Late Pleistocene

Hunter-Gatherer Ancestry in the Iberian Peninsula. *Current Biology: CB*, *29*(7), 1169–

1177.e7.

1126

Author Contributions

R.D., B.L., L.F-S., M.P.W. and X.R-R. conceptualised the study. R.D., L.F-S., K.K., M.P.W.

and X.R-R. performed laboratory work. R.D. and L.F-S. performed computational

analyses. R.T., B.L., and L.F-S. funded the study. All authors contributed to manuscript

writing and revisions.

Data Accessibility and Benefit-Sharing Section

<u>Genomic Data:</u>

The newly generated genomic data used in this study will be made available upon request,

only to replicate the results presented in this study.

1136

1137 <u>Code Availability:</u>

1138 Code used for analyses and plot generation is available at https://github.com/roberta-

1139 davidson/Davidson_etal_2023-Capture-bias-paper.

1140

1141 **Tables and Figures**

1142 **Table 1:** Relevant modified variables for each of the laboratory experiments.

Variable	Standard Protocol	Experiment A	Experiment B
Enrichment rounds	2	1	1
Hybridisation temperature	70°C	70°C	62.5°C
Amplification cycles	20 (1st round) + 10 (2nd round)	12	13

1143

1144

Figure 1: Outgroup f_3 statistics comparing ancient South American and European populations by data type. Panels (A) and (B) show results for Spain.PP.2rnd and Peru.PP.2rnd as the target population, respectively. Note that not all regions have data of every type and different data types from the same region are not the same individuals but are expected to share broadly similar ancestries.

1151

Figure 2: Comparing PCA methods using a collated dataset of 493 ancient genomes from
South America and Iberia. Samples are coloured by region (left column of each panel) or
data type (right column of each panel). (A) PCA computed with EMU (Meisner et al.,
2021). (B) PCA computed with smartPCA, no projection (Patterson et al., 2006; Price et
al., 2006). (C) PCA computed on data passed through a PCA-based filter—which removed
SNPs that drove PC2 in panel B—using smartPCA without projection (Patterson et al.,
2006; Price et al., 2006). (D) PCA computed with smartPCA where Prime Plus data (PP;
green squares) were projected onto non-PP data.

1160

1161 **Figure 3:** PCAs computed with EMU (left) and smartPCA (right) of ancient genomes from
1162 South America, Central America, Southwest Asia and Iberia coloured by region (top row)
1163 or data type (bottom row). Ancient samples are projected by smartPCA onto the modern
1164 diversity of the HGDP shown in grey (Bergström et al., 2020).

 Figure 4: f_4 statistics showing the effectiveness of 29 bias-reduction strategies. Test
configurations are, from left to right: f_4(Mbuti, Spain.PP.2rnd; Target, Peru.Shotgun),
f_4(Mbuti, Spain.PP.2rnd; Target, Peru.1240k), f_4(Mbuti, Spain.1240k; Target,
Peru.Shotgun), f_4(Mbuti, Spain.1240k; Target, Peru.1240k). "Target" groups are
shown along the y-axis, being either one of the 29 strategies used to resolve the Prime
Plus bias, or shotgun or 1240k data from Peru or Spain. Numbers above the points
indicate the number of SNPs used in each f_4 statistic and the associated Z-score. Non-
significant Z-scores ($|Z| < 3$) are shown in blue, and significant Z-scores ($|Z| > 3$) are
indicated with a yellow-red gradient. Error bars represent ± 2 se. The vertical dashed
line represents the observed f_4 statistic in each configuration when Peru.PP.2rnd was
the Target group. Note that bias-reduction strategies are only applied to the target
population and the f_4 statistic is calculated from SNPs overlapping between the target and
the other populations that remain unchanged between tests. Note that in the f_4 statistics
where the target population is off-target or out-of-range genotypes
(Peru.PP.2rnd.OffTarget, Peru.PP.2rnd.>100bpOffTarget), all four populations were
genotyped in the respective alternative way to ensure overlapping SNPs, and therefore
these are not exactly comparable.

1183

1184

Figure 5: Mean frequency of A1 allele calls in different data types as a function of SNP
missingness. Results for the four different A1 allele pairs are provided in separate panels,
with inset data points showing the mean allele frequencies calculated across all individuals
in successive SNP missingness bins. SNPs that are fixed within each dataset were removed

to avoid skewing the mean A1 calls. The header of each panel shows the number of target

SNPs of the corresponding A1/A2 pair observed in the Prime Plus assay.

Allelic bias when performing in-solution enrichment of ancient human DNA

Roberta Davidson, Matthew P. Williams, Xavier Roca-Rada, Kalina Kassadjikova, Raymond Tobler, Lars Fehren-Schmitz, Bastien Llamas

Figure S1: PCA of ancient genomes from South America coloured by region (left) and data type (right). Shapes are consistent between rows. Computed with EMU (Meisner et al., 2021).

Figure S2: PCA of ancient genomes from South America coloured by region (left) and data type (right). Shapes are consistent between rows. Computed with smartPCA without projection (Patterson et al., 2006; Price et al., 2006).

Figure S3: READ results for pairwise combinations of four individuals from Peru for which we had generated shotgun and single round Prime Plus enriched data as a part of Experiment B. Each point is labelled with the number of SNPs used in the pairwise comparison.

Fig S4: PCA of SNP missingness between data types (bottom row) computed with smartPCA. The top row is coloured by missingness frequency where yellow is high missingness and blue is low missingness.

Figure S5: Key summary statistics from sequencing libraries at different levels of enrichment (x-axis), contrasting increasing endogenous DNA retrieved (top) with decreasing complexity, taken as % unique reads (bottom). Values are standardised by the total reads.

Tables S1-S3 and S6 are in the Supplementary Excel spreadsheet.

Table S1: Number of ancient samples per region for each data type represented in this study.

Table S2: Processing details of all Prime Plus enriched samples in this study.

Table S3: Metadata for all published samples used as comparative data in this study

Table S4: List of possible ways to structure outgroup-f_3 and f_4 statistics as well as qpAdm and qpWave configurations using PP data and their respective reliability.

f_3 statistic structure	Reliability and Expectation
Where Outgroup is kept constant between comparisons and X and/or Y rotate between populations or individuals	
f_3 (Outgroup.PP; X.PP, Y.PP)	Reliable, bias affects all equally
f_3 (Outgroup.notPP; X.PP, Y.PP)	Reliable, under the assumption that bias does not significantly differentially affect different ancestries.
f_3 (Outgroup.notPP; X.sometimesPP, Y.sometimesPP)	Unreliable, X.PP will always have more shared drift with Y.PP than X.notPP.
f_3 (Outgroup.notPP; X.notPP, Y.PP)	Reliable, the bias in Target.PP is not shared with X or Outgroup so the test should be agnostic to it.
f_3 (Outgroup.notPP; X.notPP, Y.notPP)	Reliable when using shotgun or legacy 1240k, standard practice.

f_4 statistic structure	Reliability and Expectation
Where A is usually an outgroup population and B, C and D may rotate between populations or individuals	
f_4 (A.PP, B.PP; C.PP, D.PP)	Reliable, bias affects all equally
f_4 (A.notPP, B.PP; C.PP, D.PP)	Reliable, under the assumption that bias does not significantly differentially affect different ancestries.
f_4 (A.notPP, B.PP; C.notPP, D.PP)	Unreliable, B.PP and D.PP will be artificially attracted to each other
f_4 (A.notPP, B.notPP; C.PP, D.PP)	Reliable, as C and D share the PP bias, the test will reveal whether either received gene flow from B.
f_4 (A.notPP, B.PP; C.PP, D.notPP)	Unreliable, C.PP will be more attracted to the other PP population than it truly is.

f_4(A.notPP, B.notPP; C.sometimesPP, D.PP) f_4(A.notPP, B.notPP; C.PP, D.sometimesPP)	Unreliable
f_4(A.PP, B.notPP; C.notPP, D.notPP) f_4(A.notPP, B.PP; C.notPP, D.notPP) f_4(A.notPP, B.notPP; C.PP, D.notPP) f_4(A.notPP, B.notPP; C.notPP, D.PP)	Reliable, the bias in the one PP population is not shared with the others so the test should be agnostic to it.
f_4(A.notPP, B.notPP; C.notPP, D.notPP)	Reliable when using shotgun or legacy 1240k, standard practice
qpAdm configuration	**Reliability and Expectation**
Where T = target, S = source, R = reference	
(T) = T.PP, (S) = S.notPP, (R) = R.notPP	Reliable, under the assumption that the PP bias does not significantly differentially affect different ancestries or either of the shotgun and 1240k technologies.
(T) = T.PP, (S) = S.sometimesPP, (R) = R.notPP	Unreliable, bias in the matrix of f_4 statistics underlying qpAdm could lead to artificial relationships and weight estimations from the source to the target population.
(T) = T.PP, (S) = S.sometimesPP, (R) = R.sometimesPP	Unreliable, bias in the matrix of f_4 statistics could lead to incorrect estimation of the matrix rank in addition to the problems outlined above.
(T) = T.PP, (S) = S.notPP, (R) = R.sometimesPP	Unreliable, bias in the rank of the matrix of f_4 statistics can be driven by the artificial affinity between the T.PP target and R.PP right-group leading to a false inference in an increase in the matrix rank.
qpWave configuration	**Reliability and Expectation**
Where L = left, R = right	
(L) = L.PP, (R) = R.PP (L) = L.PP, (R) = R.notPP (L) = L.notPP, (R) = R.PP	Reliable.
(L) = L.PP, (R) = R.sometimesPP (L) = L.sometimesPP, (R) = R.PP (L) = L .sometimesPP, (R) = R	Unreliable, bias in the matrix of f_4 statistics between L.PP and R.PP underlying qpWave could lead to a false estimation of the matrix rank.

.sometimesPP	

Table S5: List of all strategies tested to reduce Prime Plus assay bias and their predicted impact on bias. Filters tried with different thresholds have variable values listed inside braces {}.

Strategy	Hypothesised impact or reason for testing.
Remove transition SNPs	Remove possible impact of ancient DNA damage.
Remove reads <50 bp	Reduce impact of reference bias.
Allele frequency-based filters	
SNP missingness {>0%, >50%}	Reduce impact of bias caused by preferential capture of certain SNPs by each enrichment, causing differential patterns of data missingness.
Minor allele frequency (MAF) {> 1%, >10%, >30%}	Explore the effect of MAF filters on bias. Informative about the relationship of bias to segregating sites.
Linkage disequilibrium filters Window size = 3 variants, slide = 1 variant, r^2 = {0.1,0.3,0.5,0.7,0.9}	Remove artefactual 'linkage' between neighbouring variants that could potentially be captured by one bait. Exploratory tested with a range of r^2 values due to unknown expected correlation.
SNP-set based filters	
Rohland et al. filter	Reported in Rohland et al. 2022 to allow co-analysis of Legacy 1240k and shotgun data.
Filter to Human Origins SNPs	Explorative, to understand the bias impact on these most commonly used SNPs
PCA-derived filter	Remove the bias conferred by SNPs driving the "assay bias".
Alternative genotyping	
Random diploid	Explore the capture of alleles at heterozygous sites.
Pseudohaploid by majority (>5X sites)	
Off-target sites	Remove bias that could be due to the biochemical binding of bait to target DNA, focussing on off-target by-catch.
>100 bp out of range sites	Same as above, plus removing biased by-catch where one bait may cover two target sites.
Laboratory protocol adjustments	
Protocol A Single capture round, less amplification cycles	Overall reduction of PCR cycles reduces amplification of any preferential binding bias.
Protocol B Single capture round, less amplification cycles, lower	Same as above, and lower hybridisation temperature decreases specificity of bait binding, thus reducing bias in the allele captured.

hybridisation temperature	

Table S6: Count of number of SNPs used in each f_4 test in Figure 4 and corresponding Z scores.