Title: Dynamic ventral disc contraction is necessary for *Giardia* attachment and host pathology

Authors: [1]Nosala,C, [1]Hagen, KD, [1]Guest, SL, [1]Hilton, NA, [2]Müller, A, [3]Laue, M, [2]Klotz, C., [2]Aebischer, A

and [1]Dawson,SC

[1]Department of Microbiology and Molecular Genetics

One Shields Avenue

UC Davis

Davis, CA 95616

[2]Robert Koch-Institut (RKI)

FG16 Erreger von Pilz- und Parasiteninfektionen und Mykobakteriosen

Seestraße 10

13353 Berlin

[3]Robert Koch-Institut (RKI)

Advanced Light and Electron Microscopy (ZBS 4),

Centre for Biological Threats and Special Pathogens,

Seestraße 10

13353 Berlin

Corresponding author: scdawson@ucdavis.edu

Keywords: *Giardia*, parasite, ventral disc, microtubule organelle.

<u>Abstract:</u>

Giardia lamblia is a common parasitic protist that infects the small intestine and causes giardiasis,

resulting in diarrhea, vomiting, weight loss, and malabsorption. Giardiasis leads to cellular damage,

including loss of microvilli, disruption of tight junctions, impaired barrier function, enzyme inhibition,

malabsorption, and apoptosis. In the host, motile *Giardia* trophozoites attach to the duodenal

microvilli using a unique microtubule organelle called the ventral disc. Despite early observations of

disc-shaped depressions in microvilli after parasite detachment, little is known about disc-mediated

attachment mechanisms and there little direct evidence showing that parasite attachment causes

cellular damage. However, advancements in *in vitro* organoid models of infection and genetic tools

have opened new possibilities for studying molecular mechanisms of attachment and the impact of

attachment on the host. Through high-resolution live imaging and a novel disc mutant, we provide

direct evidence for disc contraction during attachment, resolving the long-standing controversy of its

existence. Specifically, we identify three types of disc movements that characterize contraction, which

in combination result in a decrease in disc diameter and volume. Additionally, we investigate the

consequences of attachment and disc contractility using an attachment mutant that has abnormal disc

architecture. In a human organoid model, we demonstrate that this mutant has a limited ability to

break down the epithelial barrier as compared to wild type. Based on this direct evidence, we propose

a model of attachment that incorporates disc contraction to generates the forces required for the

observed "grasping" of trophozoites on the host epithelium. Overall, this work highlights the

importance of disc contractility in establishing and maintaining parasite attachment, leading to

intestinal barrier breakdown.

Introduction

Giardia lamblia is a common zoonotic protistan parasite that colonizes the small intestine, causing the

diarrheal disease giardiasis . Giardiasis occurs worldwide and is especially prevalent where water

quality is poor [1]. Early and recurrent childhood *Giardia* infection can exacerbate malnutrition and

cause development delays [2]. Hosts ingest *Giardia* cysts, which then excyst to release flagellated

trophozoites that colonize and attach to the duodenal microvilli with the "ventral disc", a conspicuous

cup-like microtubule (MT) organelle unique to *Giardia* species [3]. Common symptoms of *Giardia*

infection include diarrhea, vomiting, weight loss, and malabsorption [2]. At the cellular level, the

consequences of *Giardia* infection are reported to include damage to and loss of enterocyte microvilli,

disrupted tight junctions and barrier function, inhibition of brush border enzymes, malabsorption, and

apoptosis [2].

The pathophysiological consequences of *Giardia* infection have traditionally been attributed to

indirect, rather than direct interactions of the trophozoite with the host epithelium during parasite

attachment [2]. These indirect interactions include factors such as the unique substrates and products

of parasite metabolism, secretion of cysteine proteases, or disruptions to the host microbiome [2,4].

The intricate interplay between parasite attachment, colonization, and direct damage to the

epithelium was a key component of early observations of *Giardia*-host interactions, and researchers

specifically noted the presence of ventral disc-shaped depressions that remained in the microvilli after

parasites detached from the epithelium [5,6] [6].

Initially, actin-based mechanisms of the disc-associated lateral crest, rather than movements of

the disc itself, were proposed to generate the underlying biophysical forces required for the grasping

behavior of trophozoites on the epithelium [7], and lateral crest mediated attachment of the disc was

assumed to either directly cause tissue damage or indirectly limit absorption of nutrient in the small

intestine. Thus actin-based disc contractility has since been generally discounted. In contrast, the oft-

cited hydrodynamic suction model of Holberton [8] introduced an innovative hypothesis for

attachment that discounted contractile forces of the disc [6] and instead proposed that beating of the

ventral flagella produced a hydrodynamic current drawn underneath a rigid, inflexible disc. Using

flagellar motility mutants, we challenged this hypothesis [9], and demonstrated that attachment to

surfaces was unaffected when flagellar motility was impaired. Thus, ventral flagellar beating alone is

not necessary for maintaining hydrodynamic suction.

In the absence of either actin-based contraction of the lateral crest [7] or hydrodynamic suction

mediated by flagellar motility, the disc itself becomes the obvious candidate for force generation for

attachment. Consequently, a thorough comprehension of disc architecture is essential to unravel

attachment mechanisms and understand their impacts on the host. Attachment mechanisms likely

derive from the novel structural components or protein complexes comprising the disc. As an organelle

of attachment, the ventral disc is an elaborate domed structure defined by approximately 100 parallel

MTs that spiral until they overlap in the "overlap zone", a region where the upper and lower disc layers

are connected (Figure 1A and [3]). The spiral MTs scaffold additional unique protein complexes

including the ribbon-like microribbon-crossbridge (MR-CB) complex. The asymmetric trilaminar

microribbons track the entire MT spiral and extend 400 nm into the cytoplasm, while the crossbridges

connect parallel microribbons. Additional protein complexes comprise the disc perimeter, or "disc

margin," and the lateral crest, which is an additional repetitive structure that lies outside the MT spiral

on the disc circumference and been predicted to grasp the host microvilli [6]. We previously identified

close to 90 disc-associated proteins (DAPs) that associate with specific disc substructures, such as the

lateral crest or overlap zone [10], that could be involved in force generation for attachment (Figure 1).

With respect to the role of DAPs in disc structure, we have recently used both CRISPR-based

knockdowns and CRISPR-based quadruple KOs to show that some DAPs are critical for the disc domed

architecture[11,12], and others are essential in providing for disc MT stability and overall disc

hyperstability [10].

Combined with new *in vitro* models such as human gastrointestinal epithelium iPSC organoids

that recapitulate epithelial damage and barrier function, the advent of new genetic tools to create

stable *Giardia* attachment mutants offers promise for discovering the impact of trophozoite

attachment and colonization on the host. We recently developed robust CRISPR-based genetic tools to

create highly penetrant disc attachment null mutants, and have used these nulls with the organoid

infection model to test the impact of attachment [11,13]. Our results indicated that disc-mediated

attachment is necessary for the disruption of epithelial barrier integrity and functioning[11].

Nonetheless, with respect to cellular mechanisms of attachment, we still lack direct evidence for the

disc contractility initially reported [6] and we also lack direct evidence that disc-mediated attachment

and/or contractile functions cause cellular damage to the host epithelium [2].

Yet to paraphrase the classic maxim: the ***absence of evidence*** for disc contraction ***is not***

evidence of the absence of disc contraction. Here, using high resolution live imaging of disc dynamics

with DAPs that mark the lateral crest, we provide direct evidence for disc contractility, resolving the

long-standing controversy regarding disc contraction during parasite attachment. Live imaging of disc

"marker" strains allowed us to track and quantify movements of disc components relative to each

other in seconds in live attaching trophozoites. We define three types of disc and lateral crest

movements that characterize overall disc contraction that decreases the disc diameter by about 0.5

µm and disc volume by roughly 10-15%. Further, we test the consequences of attachment and disc

contractility for the first time, using a DAP quadruple null mutant (DAP7268) that lacks contractility and

has an aberrant disc "Baumkuchen" architecture with multiple disc overlap zones and lateral crests.

Using the infections with human organoid derived monolayers [13] we show that Baumkuchen

mutants have a limited ability to cause epithelial barrier breakdown . Based on this new, direct

empirical molecular genetic evidence, we propose a model of attachment with a mechanism for disc

contraction that could generate forces for the grasping behavior initially observed over 50 years ago.

Overall, this work demonstrates that disc contractility is necessary for establishing and maintaining

parasite attachment and that such contractility-based attachment is necessary for intestinal barrier

breakdown.

Materials and Methods

Giardia cultivation, electroporation, and isolate cloning

All strains used in this study, including tagged, knockout, and other variants, were derived from *Giardia lamblia* WBC6 strain ATCC 50803. Strains were thawed from frozen stocks and cultivated at least 24 to 48 hours prior to phenotypic analysis unless otherwise specified. Routine cultivation was carried out in 16-ml screw-cap plastic tubes (BD Falcon) without agitation. TYI-S-33 growth medium was supplemented with bovine bile, 5% adult bovine serum, and 5% fetal bovine serum, and the cultures were maintained at 37°C. Upon reaching confluency, culture tubes were incubated on ice for 30 minutes, and 0.5 to 1 ml of detached trophozoites was added to 12 ml of pre-warmed medium. Antibiotics were added into the culture medium when selecting for episomal plasmids (GFP-tagged strains) or gene knockouts ([11] and further details provided below).

Electroporation of DNA constructs (plasmids or linear homology-directed repair (HDR) templates) into

Giardia were performed under conditions described previously [12], using 20 to 40 µg DNA and 1 x 10^7

trophozoites per electroporation. Electroporated cells were then incubated in standard culture tubes

at 37C, with medium replacement every 48 hours. Lower antibiotic concentrations were used for initial

selection and antibiotics were increased to maintenance levels once culture confluency reached >50%.

Concentrations used were: puromycin (10 µg/ml and 50 µg/ml), blasticidin (75 µg/ml and 150 µg/ml),

and hygromycin B Gold (600 µg/ml and 1200 µg/ml). All antibiotics were purchased from Invivogen

except G418 (Fisher Scientific).

Clonal cell lines were obtained by limiting dilution. Iced, detached trophozoites were counted

with a Luna-II cell counter (Logos Biosystems). Trophozoites were diluted to 2 cells per ml in fresh TYI-

S-33 containing antibiotics used for selection, and 250 µl aliquots (0.5 cells/well) were added to 96 well

tissue culture plates (BD Falcon). Plates were incubated at 37C in Mitsubishi AnaeroPack 2.5L

rectangular jars with a Mitsubishi AnaeroPack-Anaero Gas Generator sachet (ThermoScientific) for 7 to

10 days. Typically, 20 to 30% of plate wells contained cells, and most wells with cells had reached 80 to

100% confluency. Plates were iced 30 minutes and detached trophozoites were transferred to 8 ml

screw-capped tubes (BD Falcon). Once these tubes reached >50% confluency, cells were propagated in

standard 12 ml tubes.

Design and creation of DAP7268 and DAP7268_12139mNG quadruple knockout (null) strains

Our CRISPR quadruple knockout strategy has been described elsewhere [11]. Briefly, a guide RNA

targeting the coding region of DAP7268 at position 908 (gRNA908R, 5'-AGTTGATCCGGAACGCTGTG-3') was

selected using Benchling's CRISPR 'Design and Analyze Guides' tool (https://benchling.com/crispr) with

an NGG PAM sequence and the *Giardia lamblia* ATCC 50803 genome (GenBank Assembly

GCA_000002435.1). A slightly modified vector, Cas9U6g1pac, was used to express both the gRNA and

Cas9 fused to the *Giardia* GL50803_2340 NLS—the gRNA expression cassette in this new vector differs

from our previously published CRISPR and CRISPRi vectors [11,14] in that 12 bases encoding U6 snRNA

that were immediately downstream from the U6 promoter have been removed. Complementary gRNA

oligos (5'-cggcAGTTGATCCGGAACGCTGTG-3' and 5'-aaacCACAGCGTTCCGGATCAACT-3') with 4 base

overhangs were annealed and ligated to BbsI-digested Cas9U6g1pac as previously described [11]. This

vector was introduced into wild-type *Giardia* WBC6 as described above, using puromycin selection.

To make the DAP7268 quadruple knockout strain DAP7268quadKO, two *Giardia* antibiotic

cassettes, BsdRgo and HygRgo were introduced sequentially into the Cas9DAP7268_gRNA908R

expression strain above by electroporation of 20 to 40 ug of linear HDR template as previously

described [11]. Each HDR template consisted of an antibiotic cassette flanked by the 750 bp of

DAP7268 sequence immediately up- and downstream from the DSB site at position 908 (Supplemental

Figure 1). The templates were synthesized by Twist Biosciences and cloned into their pTwist Amp High

Copy vector. Linear HDR templates were amplified from this vector using Q5 DNA polymerase (NEB)

and M13Forward and M13Reverse primers and purified using Zymo Research Clean and Concentrator-

25 columns. Once knockout strains reached >50% confluence on full antibiotic selection, total DNA was

extracted using the Zymo Research Quick DNA Miniprep Plus kit. Knock in of markers and loss of wild-

type DAP7268 alleles was confirmed by PCR with primers 7268LeftF (5'-CTTATGCTGGCAGTGCGATCC-

3') and 7268RightR (5'-CTCTATAAGCTCCTCAACACTCTGC-3'), which bind roughly 800 bp from the DSB

site, outside the homology arms. In contrast to our knockout of DAP16343[11], which required three

markers, wild-type DAP7268 alleles were undetectable after knock-in of the second (HygRgo) cassette.

To make the DAP7268quadKO_DAP12139mNG strain, which stably expresses DAP12139 as a

marker of the disc margin, we designed a fragment containing a DAP12139-mNeonGreen C terminal

fusion, with DAP12139 expression driven by its native promoter (Supplemental Figure 2). The 2.9 kb

fragment was synthesized (Twist Biosciences) and cloned into a unique MluI site that lies just upstream

of the rpS28 promoter in the HygRgo cassette. The entire 6.3 kb DAP12139mNG_HygRgo HDR template

was amplified, purified, and introduced via CRISPR knockout after the introduction of BsdRgo. The

DAP7268quadKO_DAP12139mNG strain was confirmed by PCR as described above. For both strains,

clonal cell lines were obtained by limiting dilution (see above).

Live imaging of fluorescently tagged strains

The DAP12139 disc margin GFP fusion strain and DAP7268KO-DAP12139mNG strains were thawed

from frozen stocks and cultured at least 24 hours prior to live imaging. Cells were iced, harvested, and

resuspended in 1 ml warmed medium, and 300 µl aliquots were incubated in 96-well black glass

bottom imaging plates (Cellvis, Mountain View, CA) or MatTek dishes for up to one hour at 37°C to

promote attachment. Prior to imaging, the medium was replaced with warmed 1X HBS, and

trophozoites were incubated under the same conditions for 30 minutes. Additional warmed 1X HBS

washes were performed as needed during imaging to remove detached cells. DIC and epifluorescence

imaging was performed with a Leica DMI6000B microscope as described previously [11]. Images were

processed and disc contraction was quantified using kymographs using FIJI [8].

Immunostaining and quantification of ventral disc dimensions in mutants

Prior to immunostaining, strains were grown to confluency and fixed in 12 ml tubes with a final

concentration of 1% paraformaldehyde, pH 7.4 (37°C, 15 min). Detached cells were pelleted at 900 x g

for five minutes. The pellet was washed three times with 2 ml PEM, pH 6.9 [12], incubated in 0.125M

glycine (15 min, 25°C), and washed three more times with PEM. Fixed cell suspensions were then

settled on to coverslips treated with 1mg/ml poly-L lysine for one hour. Coverslips were then washed

3X in PEM, and permeabilized with 0.1% Triton X-100/PEM for 10 minutes. Coverslips were blocked in

PEMBALG for 30 minutes and incubated overnight at 4°C with primary antibodies for microtubules

using anti-TAT1 (1:250, gift of K. Gull, University of Oxford) or ventral disc microribbons using anti-

delta-giardin (1:100) or anti-beta-giardin (1:1000, gift of M. Jenkins, USDA) antibodies. Following the

primary antibody incubation, coverslips were washed three times in PEMBALG and incubated with Alex

Fluor 488 goat anti-rabbit and/or Alex Fluor 594 goat anti-mouse antibodies (1:1000; Life Technologies)

for four hours at room temperature. Following incubation with the secondary antibodies, coverslips

were washed three times each with PEMBALG and PEM and mounted in Prolong Diamond antifade

reagent (Life Technologies). All imaging experiments were performed with three biologically

independent samples.

Overall disc curvature, disc and bare area dimensions including perimeter and area, and surface

contacts dimensions were quantified from > 300 immunostained images per condition using Fiji [15].

Disc surface contacts were quantified from CellMask-stained images.

Super-resolution SIM 3D imaging of fixed cells

For super-resolution imaging of strains, three-dimensional stacks were captured with 0.125 μm

intervals between slices. SIM microscopy used the Nikon N-SIM Structured Illumination Super-

resolution Microscope with the 100X/NA 1.49 objective, 100 EX V-R diffraction grating, and an Andor

iXon3 DU-897E EMCCD camera. The collected images were then processed using the "3D-SIM" mode,

which involved reconstructing the images using three diffraction grating angles and three translations

each. To evaluate the quality of the images, a reconstruction score was assigned to each image using

the "Reconstruct Slice" mode. Only images with a score of 8 or higher were deemed acceptable for

analysis. Additionally, both the raw and reconstructed images underwent further assessment using

SIMcheck, a quality assessment tool developed by Ball et al. (. The final images are displayed either as

maximum intensity projections or as 3D stacks, depending on the desired representation.

Quantification of disc doming (angle of curvature)

To quantify the range of disc curvature, cells were attached to imaging dishes (MatTek) by incubating

at 37°C for 30 min in 1X HBS, washed three times with 1X HBS to remove unattached cells, and

resuspended in 100 µl of warm 1X HBS. 100 µl of warmed (37°C) 3% low melt agarose in 1X HBS was

added to wells (final concentration of 1.5% agarose) to immobilize trophozoites for live imaging. To

capture the entire disc, a Leica wide-field DMI6000B microscope was used to collect optical slices at 0.2

µm intervals for up to 8 µm. Disc doming or curvature images were generated by reslicing the stack

laterally across the posterior portion of the bare area and angles were measured in ImageJ/Fiji [15].

For each experimental condition, at least 30 cells were measured on three separate days totaling 90

cells imaged and quantified.

Quantification of shear forces of using a flow cell assay of attachment

To quantify their ability to resist shear forces in microscopic flow assays, the DAP7268KO strain, Cas9

vector control, DAP12139GFP, and wild-type trophozoites were initially incubated in Ibidi mSlide VI 0.4

flow chambers (Ibidi). Single focal plane images of attaching 1,000,000 cells were acquired using

the µManager acquisition software [16] with the 10X objective using a Leica DMI 6000 wide-field

inverted fluorescence microscope. Cells were allowed to attach for ten minutes prior to initiation of

flow. Attached cells were counted by overlaying the pre-flow image over the post-flow image. The

overall percent attached was calculated by quantifying the percentage of shear force resisting cells

relative to the total number of cells.

Organoid growth and TEER measurements

Human organoids were generated and maintained from duodenal biopsy specimens from healthy

volunteers undergoing routine examinations at Charité-Universitätmedizin Berlin (ethics approval

#EA4-015-13 by local authorities) were used to generate human organoids are previously as described

[13]. Matrigel-coated (Corning, Tewksbury, NY) transwell cell culture inserts (0.6 cm2, 0.4-mm pores;

Merck-Millipore, Burlington, MA) were then used to culture organoid derived monolayers (ODMs)

using previously described methods [13]. For infections, wild-type and DAP7268KO trophozoites were

passaged the day before infection experiments to guarantee logarithmic growth phase. Prior to

infection, parasites were detached from culture tubes by incubation at 4°C for 20 minutes, pelleted

and quantified using a Neubauer counting chamber. Parasites were then harvested by pelleting at 1000

x g for five minutes at °C and resuspended at the final desired concentration. The apical compartment

medium of 8- to 10-day-old ODMs was replaced with complete TYI-S-33 the evening before infection to

adapt cells to the TYI-S-33 medium. Medium was renewed before infection (i.e., TYI-S-33 in the apical

and organoid differentiation medium in the basal compartment). *Giardia* trophozoites were added

apically and incubated for up to 72 hours.

Transepithelial electric resistance (TEER) measurements of epithelial barrier integrity

To assess the barrier integrity of disc mutants, transepithelial electric resistance (TEER) measurements

were conducted. A Millicell ERS-2 Voltohmmeter (Merck-Millipore) equipped with an Ag/AgCl

electrode (STX01; Merck-Millipore) was used for these measurements. The instrument was placed on a

37°C heating block to maintain a consistent temperature. To quantify the TEER values, the electric

resistance of a blank transwell insert without cells was measured and subtracted from the raw

resistance values obtained from the samples. Furthermore, the resulting values were standardized to

account for a surface area of 1 cm2.

Results

Disc attachment models are defined by essential disc structural elements and flagellar movements

The ventral disc is a composite structure composed of an extensive MT spiral array and associated

protein complexes including the MR-CB complex (body, Figure 1A). The disc also has a region of

overlap, the overlap zone (oz, Figure 1A), that is key to its domed architecture. The disc margin marks

the outside edge of the spiral MT array, including a gap at the point of the disc overlap zone (Figure 1A

and [9]). The lateral crest is an additional repetitive structure is bound to the disc perimeter outside of

the MT array, and has previously been argued to be contractile (Figure 1A, [3]). The ventral groove is a

deformed region of the disc overlapping the ventral flagella, which were proposed in the

hydrodynamic model to create a hydrodynamic current required for attachment by drawing fluid

underneath an inflexible, non-contractile disc (Figure 1B).

Is the disc an inflexible, passive structure that merely maintains attachment forces generated

by flagellar motility as assumed by Holberton, or is the disc an actively dynamic and flexible structure

that independently mediates attachment forces? Overall disc contraction has been proposed to occur

through shortening or compression of the crossbridges, reducing the spacing between the MTs of the

disc spiral (e.g., disc compression, Figure 1C). This proposal implicates the MR-CB complex in an overall

and evenly spaced compression of the disc during attachment, resulting in a lower volume underneath

the disc and a smaller disc surface area and perimeter. Alternatively, the lateral crest has been

proposed to constrict, resulting in an overall contraction of the disc. In this model, the spacing between

the disc MT spiral may or may not be reduced. Lastly, the upper and lower layers of the disc could

rotate or slide over each other, again resulting in an overall disc contraction. Spacing between disc MTs

again may or may not be altered. Of course, the three models are not mutually exclusive.

Direct evidence for disc contraction during trophozoite attachment to surfaces.

To test the prediction that disc contraction occurs during parasite attachment to surfaces and to

quantify disc contractile dynamics, we used quantitative time-lapse imaging to track disc movements in

DAP12139GFP, which expresses a fluorescent tag marking the lateral crest. DAP12139GFP does not

colocalize with primary components of the disc including the microribbon-crossbridge complex (Figure

2A and [3]). Three-dimensional line scans were drawn at three regions of the disc margin including the

overlap zone (OZ), ventral groove (VG), and the disc body, to define disc margin movements with

kymograph analyses (Figure 2B,C,D and Supplemental Figures S1-S3) corresponding to disc contractile

movements associated with overlap zone rotation (Figure 2B), disc compression (Figure 2C) and disc

constriction (Figure 2D). First, we observed that the overlap zone region rotates clockwise toward the

disc center (Figure 2B, asterisk and Supplemental Figure S1). Second, by measuring changes in the

horizontal distance between the disc edges (Figure 2C), we observed disc compression (Figure 2D and

Supplemental Figure S2) that averaged 0.3-0.5 μm during attachment but ranged up to about 0.8 μm

(Figure 3E). Lastly, lateral crest constriction also occurred during attachment (Figure 2F,G,H and

Supplemental Figure S3), mirroring disc overlap zone rotation with clockwise constriction of the disc

margin toward the ventral groove and causing the opening of the disc margin to increase on average

about 0.4 μm (Figure 2G,H). In total, we quantify disc contraction of about 0.5 μm on average during

attachment that is attributable to the three types of quantifiable movements (Figure 2I) that occur

within 15 seconds (overlap zone rotation) to 30 seconds (disc compression and constriction).

Interrogating the role of the disc overlap zone ankyrin repeat protein DAP7268 in disc contractility.

Using disc-based proteomics with subsequent protein tagging, we have previously shown that

DAP7268 is a disc-associated protein that localizes to the overlap zone and disc body (Figure 3A-B).

DAP7268 also has one N-terminal and one C-terminal ankyrin repeat domain (Figure 3C), like 30 other

DAPs [3]. A predicted alpha helical domain lies between the terminal ankyrin repeat domains. To

determine the localization role of the ankyrin repeat domains, we expressed truncated forms of

DAP7268 fused to a C-terminal GFP. This ankyrin domain analysis (Figure 3D,E,F,G) indicates that the N

terminal ankyrin domain is necessary for localization of DAP7268 to the disc overlap zone (Figure 3E,

no oz localization). In contrast, the C-terminal ankyrin domain is necessary for localization to the disc

body (Figure 3F). Deletion of both ankyrin repeats has a similar localization as just the C-terminal

ankyrin.

DAP7268 quad knockouts have a "Baumkuchen" disc phenotype defined by multiple overlapping disc

layers

In initial DAP morpholino-based transient knockdown screens, we found that the DAP7268 KD strain

had aberrant disc structure (Figure 4C,D). Using our lab's new methods to create quadruple allele

deletion mutants ("quad knockouts", Methods), we confirmed a quad knockout using blasticidin (B)

and hygromycin (H) repair templates. Confirmation through genomic sequencing indicated that two

alleles were disrupted by the B marker, and two by the H marker (Supplemental Figures 4 and 5).

Immunostaining of the DAP7268quadKO strain using anti-delta Giardia antibodies that mark

disc MT-CB complexes indicated that the disc structure was severely disrupted (SIM, Figure 4C;

negative staining, 4D) as compared to the wild-type control strain. To visualize the lateral crest region

in DAP7268 quad knockouts, we also created an additional strain (DAP7268quadKO_DAP12139mNG)

that included a mNeonGreen (mNG) C-terminally tagged lateral crest marker (DAP12139, Figure 4)

adjacent to the hygromycin cassette (Supplemental Figure 6). By immunostaining for both the

microtubules (anti-TAT1) and the MT-CB complex of the disc (anti-delta-giardin and anti-beta-giardin),

we discovered that DAP7268 quad KOs had an aberrant disc with disc multiple layers as well as a large

central region (bare area). The extra layers are readily observable as extra disc margins (marked by

DAP12139mNG). DAP7268KO mutant disc phenotypes are 100% penetrant (Figure 3D). We termed the

layered disc phenotype "Baumkuchen" after the traditional multilayered German cake of the same

name (Figure 3D). Baumkuchen phenotypes are the most penetrant of DAP7268KO disc defects, as

compared to "small" discs or discs with extra partial spirals ("looped"). Overall, the area of

DAP7268KO discs was smaller (Figure 4O) and the discs were more flattened (Figure 4N). DAP7268KD

discs are also much less domed that wild type (Figure 4N). Furthermore, DAP7268KO mutant discs are

on average about 80% smaller with respect to disc perimeter, disc area and the height of the ventral

groove (Figure 4O-Q)

Knocking in DAP12139mNG in the context of the DAP7268 quad KO allowed us to quantify

aberrant lateral crests in this DAP mutant. We observed multiple lateral crest regions (Figure 4) that in

some cases bulged out into additional aberrant lateral crests. Using TEM, we confirmed that the

aberrant disc structure of the Baumkuchen phenotype included additional layers (Figure 5). We

observed upwards of four or five overlapping layers in some cases, each with MTs and MR-CB

complexes. In addition, each extra layer had aberrant microribbons that were of different heights and

irregular spacing (Figure 5). We confirmed the duplication of the lateral crest using extracted

cytoskeletons of the DAP7268 morpholino knockdown visualized with negative staining. In extracted

cytoskeletons, the extra discs layers were again visible, with additional overlap zones and duplicated

lateral crest structures causing a more circular, doubled disc (Supplemental Figure S7, arrows).

DAP7268KO Baumkuchen mutants lack disc contractility

The presence of multiple or aberrant disc edges/lateral crests in Baumkuchen mutants implied that

mutants could also have defect in disc edge contractility as demonstrated in Figure 2. Imaging

attaching DAP7268KO_DAP12139mNG strain mutants allowed us to quantify disc contractile dynamics

in DA7268KO mutants using their integrated DAP12139mNG the lateral crest marker (Figure 6). As

seen in live imaging (Figure 6A and Supplemental Figures S8-S11) and associated kymographs (Figure

6B), Baumkuchen mutants have significantly decreased overall disc contraction. With respect to

overlap zone rotation, mutants had slight rotation in the opposite direction as we observed in

DAP12139GFP strains (Figure 2). With respect to both disc compression and constriction

measurements, we determined DAP7268KO mutants have on average only 0.2 μm contraction as

compared to wild-type strains (Figure 6C,D).

Baumkuchen mutants have decreased attachment surface contacts and resistance to shear forces

Membrane contacts are diagnostic of the level of cellular attachment to surfaces during early

attachment "stages" [12]. Initial cellular contacts in Stages I and II include the ventrolateral flange and

disc edge, and later Stage III and IV contacts include the "tail" regions and the lateral shields and the

bare area in the center of the disc . After staining with the membrane dye CellMask Orange, we

determined that membrane contacts with the surface were altered in the DAP7268KO mutant. We

observed larger membrane regions of the bare area contacting the surface (Stage IV+ba), and in some

cases the bare area membrane lacked any contacts with the surface (Figure 7A). In comparison to wild-

type surface contacts which are primarily at stage IV (including lateral crest, lateral shield, and bare

area contacts), the majority of DAP7268KO mutants reached a modified stage IV stage which lacked

surface contacts associated with the bare area (Stage IV-LC). We also noted in light and electron

microscopy that the central disc region (bare area) was larger in mutants, and this is reflected in the

size of the bare area surface contacts observed (Figure 7A, Stage IV+B).

To evaluate shear forces of attachment, we used a novel quantitative shear stress assay that uses time

lapse imaging of trophozoites in commercial flow chambers (Methods). Wild-type, the Baumkuchen

quad KO, DAP12139GFP and the DAP7268KO_DAP12139mNG strains were flowed into chambers,

allowed to attach, and then challenged with a range of shear forces. Mutants and tagged strains were

normalized to wild-type. Less than 40 percent of the attached parasites resisted 3ml/min of flow

(~4dyn/cm2) Baumkuchen DAP7268KO and DAP7268KO/DAP12139mNG strains trophozoites had

significant decreases in their ability to resist 3ml/min of flow (Figure 7B,C).

Significantly reduced epithelial barrier breakdown in Baumkuchen mutants using human intestinal

organoid derived monolayers

2D human intestinal organoid-derived monolayers (ODMs) undergo maturation over a period of

approximately 9 days in culture. During this time, the ODMs reach a height of around 20 μm and

exhibit the characteristic polarized shape of enterocytes, including the presence of microvilli and well-

developed cellular junctions [36]. As the ODMs mature, there is a gradual increase in trans-epithelial

electrical resistance (TEER), which stabilizes at approximately 250 Ω · cm2 after 9 days of culture. TEER

serves as a measure of the integrity of the barrier function, and its decrease is observed in *Giardia*

infections of ODMs, indicating disruption of the barrier function [36].

To investigate the impact of wild-type and Baumkuchen mutant *Giardia* trophozoites on the

degradation of barrier function, ODMs were infected with both wild-type trophozoites and two

DAP7268KO clones (7268-B2 and 7268-C7) or treated with vehicle in mock infections (no *Giardia*) at

different multiplicities of infection (MOI) ranging from 1 to 10. In wild-type infections, the normalized

TEER exhibited a dose- and time-dependent decrease after 24 hours of infection, reaching its peak at

48-54 hours (Figure 8A, B). In contrast, both DAP7268KO clones showed significantly higher normalized

TEER compared to the wild-type control infections.

Discussion

Giardia's attachment via the ventral disc to either the host epithelium or to inert surfaces is defined by

its rapid, reversible, and dynamic nature, enabling the parasite to resist peristaltic flow in the

gastrointestinal tract and colonize the host. Numerous hypotheses have been proposed over the past

50 years to explain attachment mechanisms involving the ventral disc and/or lateral crest. These

hypotheses encompass hydrodynamic suction forces generated by flagellar motility [5,8,17],

cytoskeletal-based contraction of the disc, lateral crest, or ventrolateral flange [7,18–20], microtubule

sliding dynamics [21], and lectin-based [22–25] or osmolarity-based mechanisms [26]. Among these,

the "hydrodynamic suction model" has gained prominence based on early observations of *Giardia*

muris attached to glass slides and modelling of fluid dynamics modeling [8].

An early assertion of the hydrodynamic suction model is that the ventral disc serves as a rigid,

non-contractile structure, yet other early observations suggested a more contractile disc and lateral

crest that engages and directly grips or "clutches" the host epithelium [7]. This disc contractility has

been proposed to account for the formation of disc-shaped tissue damage or "craters" when

trophozoites attach to microvilli. Such conflicting observations and insufficient empirical evidence

supporting attachment mechanisms such as flagellar-based hydrodynamic suction have led to

incongruent attachment models, however   [5,7,8,17–25]. Moreover, the lack of advanced genetic

tools for creating stable attachment mutants and a lack of robust *in vitro* and *in vivo* infection models

has hindered investigations into whether contractility-based attachment directly contributes to the

observed host cellular damage.

To gain insights into disc attachment mechanisms, we examined the contractility of the disc and

lateral crest. Through live imaging (Figure 2) and analysis of a DAP contractility mutant with what we

term the "Baumkuchen" phenotype (Figures 3-7), we provide direct evidence of disc and lateral crest

contractility. While the specific molecular mechanisms underlying these contractile movements are yet

to be determined, our results are consistent with earlier models put forth by Feely et al. [7] and Narcisi

et al. [18], yet in the absence of actin-based contractility. This discovery of disc movements associated

with contractility is pivotal for advancing our understanding of *Giardia* attachment mechanisms and

their impacts on the host epithelium.

Three types of dynamic movements of the ventral disc during attachment

We define and quantify three types of disc dynamic movements that occur in seconds in attaching cells

and result in overall disc contraction: "**disc compression**", "**lateral crest constriction**", and "**overlap**

zone rotation" (Figure 1 and Figure 2). Each of these movements are not mutually exclusive, and all

may contribute to overall disc contraction we report here. Overall disc contraction in attaching cells

reduces the diameter and the volume of discs by roughly 10-15 % (Figure 2). Changes in the relative

positions of various disc components, such as MR-CB complexes, the overlap zone region, and the

lateral crest have the potential to either expand or contract the diameter of the disc resulting in the

characteristic "gripping" of the host microvilli by the ventral disc [7]. Although indirect mechanisms

involving anterior flagellar movements may also contribute to disc contractility, there is currently no

supporting evidence for this notion.

The first mode of disc contractile movement, which we term "**disc compression**," is expected to

act uniformly throughout the disc. A potential mechanism could involve the compression of

crossbridges connecting adjacent microtubules (MTs) while maintaining a 50-nanometer spacing in the

disc MT spiral. Dynamic movements of microribbons, composed of SF-assemblins (beta-giardin, delta-

giardin, SALP1 [3], could also contribute to overall disc compression. Although Holberton [5] initially

proposed this hypothesis regarding disc MR-CB complex compressions, it remains unconfirmed.

The type of second contractile movement we observe involves the direct "**constriction**" of the

lateral crest or disc margin area of the disc (Figure 1C). Early observations of attaching trophozoites

proposed that contractile activity in the lateral crest leads to the compression of the brush border

microvilli [27](Erlandsen and Chase 1974, Takano and Yardley 1967, Balazs and Szatloczky 1978). These

contractile forces were initially attributed actin and actin-associated proteins including myosin-based

immunostaining of actin-binding proteins using heterologous antibodies (Feely 1982). However, later

genome sequencing revealed that these localizations were artifacts, as only actin itself and no

canonical actin binding proteins are present in *Giardia*. Furthermore, *Giardia* actin does not directly

localize to the disc itself, but may interact with the disc through actin-associated disc proteins or

DAAPs [28]. Alternatively, the constriction of the lateral crest could reduce the disc perimeter, and the

mechanisms of constriction could be governed by specific lateral crest protein complexes or DAPs. Our

laboratory has identified 30 disc-associated proteins (DAPs) that localize to the lateral crest and disc

margin including ankyrin repeat proteins that may contribute to the lateral crest constriction that

ultimately results in disc contraction. The ankyrin repeat protein DAP12139 used to track contractile

movements (Figures 2 and 7) is likely associated with the lateral crest as the GFP staining does not

colocalize with disc components including microribbons (Figure 2).

The third dynamic disc movement involves the rotation, or possible sliding, of the upper and

lower layers of the overlap zone, a movement we term "overlap zone rotation" (Figure 1C). The

physical connection between the upper and lower layers is not well understood, and the mechanism by

which overlap zone DAPs or other DAPs may mediate these movements remains unclear. Additionally,

these disc movements could be associated with the constriction of the lateral crest.

Phenotypic analyses of DAP7268KO Baumkuchen mutants support an evidence-based model of

attachment that requires disc and lateral crest contractility

The game-changing new genetic methods used here to create the stable, isogenic DAP7268 quadruple

deletion mutant alleviate issues of mutant phenotypic transience [29] or incomplete penetrance [14]

that can confound and limit both *in vitro* and *in vivo* assays of attachment. Because plasmids, such as

CRISPRi plasmids, are readily lost off positive selection [14], DAP knockouts are also critical for *in vivo*

analyses of *Giardia*-host interactions and pathology.

Initially identified in DAP morpholino screens, DAP7268 knockdown and knock out mutants of

result in gross structural defects of the disc spiral including duplication or additional "layering" of the

disc and lateral crest (Figure 4). We have coined the term "Baumkuchen" to describe this mutant disc

phenotype, as it resembles a traditional German cake with multiple thin layers that resemble tree

rings. DAP7268 consists of one N-terminal and one C-terminal ankyrin repeat domain linked by a

central alpha-helical yet non-homologous region (Figure 3) and is among more than 30 ankyrin repeat

containing disc-associated proteins (DAPs). DAP7268 is also one of the 31 DAPs that specifically localize

to the "overlap zone," and is specifically positioned at the outer edge of the overlap zone (Figure 1).

Ankyrin repeats are generally recognized for their involvement in protein-protein interactions (Li et al.,

2006), and some ankyrin proteins have been shown to directly interact with or bind to microtubules

(Bennett and Davis, 1981; Davis and Bennett, 1984). Furthermore, in the apicomplexan protist

Toxoplasma gondii, ankyrin repeat proteins link structural and motor proteins with the tubulin-rich

conoid fibers (Long et al., 2017). In *Giardia*, our previous genetic analysis of another disc associated

ankyrin-repeat protein DAP5188, which localizes to the entire disc and overlap zone, revealed its role

in limiting interphase disc edge MT dynamics and promoting disc hyperstability [10].

DAP7268 knockout mutants also have irregular microribbons with uneven heights and spacing

(Figures 4 and 5), disrupting the microribbon-crossbridge (MR-CB) complexes crucial for the overall

domed architecture of the disc [30]. The disc spiral MT array's edge is defined by the "disc margin" and

is surrounded by the "lateral crest," which makes contact with the ventral surface and has a gap at the

overlap zone ([3] and Figure 1). Depletion of DAP7268 leads additional disc layers and additional lateral

crests (Figures 4 and 5). The Baumkuchen phenotype supports a negative regulatory role for DAP7268

has in disc assembly, preventing additional layers of the disc MT spiral and lateral crest from forming

during cell division. The presence of duplicated lateral crest regions, marked by DAP12139mNG in both

wild-type and DAP7268 knockout mutants, further supports the idea that the lateral crest assembles

after the disc MT array (Figures 4 and 5).

DAP7268KO Baumkuchen mutants also have significantly reduced (about 60% less) disc contraction

and lack overlap zone rotation or other contractile movements (Figure 6). The lack of disc contractility

in mutants is likely due to aberrant and extra disc edges/lateral crests. Disruption of the MR-CB

complex in the DAP7268KO is another mechanism that may contribute to the observed lack of disc

contractility and attachment (Figures 6 and 7). The lack of disc contractility in Baumkuchen mutants importantly results in decreased ability of parasites to resist shear forces in flow chambers (Figure 7) or create proper surface membrane contacts that require e a lateral crest seal [9]. During the early stages of attachment, trophozoites make initial contact with the surface using a specialized anterior structure called the ventrolateral flange (VLF), as first described by Holberton [5] and later confirmed by Erlandsen [19]. Recent studies have further supported the existence of focal contacts initiated by the VLF in the adhesion of parasites on laboratory surfaces [20]. These focal contacts contribute to the formation of a seal, which plays a crucial role in maintaining negative pressure underneath the disc, independent of flagellar beating. The presence of this seal marks the transition from "attaching" trophozoites to "attached" trophozoites (Figure 7 and [9]). The lateral crest seal serves two main purposes: it firmly secures the trophozoite to the attachment surface and restricts the flow of fluid entering or exiting from beneath the disc on smooth, non-elastic surfaces such as slides. Additionally, this seal positions the dorsal ventral disc membrane and peripheral vacuoles in direct contact with the microvilli of the host's brush border.

Thus, we contend that disc contraction dynamics are necessary and likely directly responsible for the force generation required for attachment. We propose a "**disc contraction**" model of attachment that incorporates overall disc contraction with subsequent constriction of the lateral crest to create a closed seal around the disc perimeter. Such dynamic movements are necessary for attachment and resistance to hydrodynamic flow on inert surfaces, and likely result in "gripping" or "grasping" of parasites on host enterocytes.

Disc contraction and attachment are necessary for intestinal barrier function breakdown associated with giardiasis

Direct cellular damage caused by extracellular attachment via the ventral disc has been assumed for decades, but not directly proven. Early reports also suggested that upward curvature of the brush border membrane of epithelial cells seen underneath the ventral disc in attached trophozoites was caused by attachment suction-based forces. *Giardia* infections are known to either directly or indirectly cause various types of damage to the host epithelium. Attached trophozoites could directly cause observed epithelial damage including reduced barrier function and integrity, to microvilli damage and shortening, degradation of the mucus layer, decreased nutrient absorption, and apoptosis. The direct or indirect impact of *Giardia's* extracellular attachment on the observed cellular pathology in *Giardia* infections has been largely overlooked in previous investigations of the disease's mechanisms.

The recent development of primary organoids, derived from small intestinal stem cells [6, 36] has opened up new opportunities for understanding *Giardia* pathogenesis. 3D cultures, derived from human duodenal biopsy specimens, are used to create 2-dimensional transwell systems known as organoid-derived monolayers (ODMs). Infections of ODMs with *Giardia* cause a gradual disruption of the epithelial barrier, directly correlated with the duration of infection and the parasite load. Infections trigger changes in gene expression related to ion transport and the structure of tight junctions in ODMs. Recent studies using ODM models have also revealed alternative mechanisms by which *Giardia* induces cellular pathologies. Inhibiting the activity of cysteine proteases and caspases, which are enzymes suggested to be virulence factors in giardiasis, does not prevent the breakdown of the epithelial barrier as measured using trans-epithelial electrical resistance (TEER) or prevent the occurrence of apoptosis. Further, we recently showed that attachment is required for complete barrier breakdown. Infections with the disc MBPquadKO mutant with significant structural and attachment

1   defects, have markedly reduced impacts on barrier integrity as compared to wild-type *Giardia*

2   infections of ODMs.

3   Here we provide additional support for the direct link between attachment and intestinal

barrier breakdown. Using infections with the 100% penetrant Baumkuchen mutants that lack disc

contract contractility, we again see a similar reduction the degree of barrier breakdown in mutants as

compared to wild type infections (Figure 8). Both DAP7268KO clones show significantly limited ability

to induce barrier breakdowns, which highlights the crucial role of parasite attachment in causing

complete barrier breakdown in this ODM model of giardiasis. Other factors are likely still involved since

we do not observe a complete absence of barrier breakdown, but the close and intimate association

provided by extracellular contractility-based attachment is necessary for host cellular pathology and

likely contributes to the systemic symptoms observed in giardiasis.

Conclusions

The modern molecular genetic and live imaging approaches used in this work bridge the gap between

early observations of disc contraction and the presumed effects of attachment on the host epithelium.

While live imaging of movements of the disc resulting in disc contraction on inert surfaces would likely

result in a gripping or grasping behavior of parasites on host enterocytes, the precise mechanism

underlying these movements remains unknown. DAPs comprising specific structural elements of the

disc may generate the forces required for disc contraction, yet these also remain to be discovered.

Future investigations should aim to identify the specific DAPs involved in the contractile mechanisms of

the disc, such as those present in the overlap zone, the MR-CB complex, or the lateral crest.

Furthermore, it still remains a possibility that interactions between the disc and the actin cytoskeleton

may also play a role in these movements. Overall, this work emphasizes the significance of disc

contractility in establishing and maintaining parasite attachment, ultimately leading to key aspects of

host pathology.

<u>Acknowledgements</u>

This work was supported by a EuPATH/GiardiaDB award and a R01AI077571 award to SCD. We

graciously thank the UC Davis MCB Microscopy Core for helpful advice on the SDC and SIM

microscopes.

Figure Legends

Figure 1. Three types of disc movements may be associated with disc contraction

As illustrated in the negative stained disc preparation (A) the ventral disc microtubules are nucleated

from the basal bodies (bb) located in the cytoplasm. Distinct structural regions of the disc and lateral

crest (lc) are noted, including the disc body (body), the ventral groove (vg), the disc margin (dm), the

overlap zone (oz), and the dense bands (db). In B, hydrodynamic currents generated by the ventral

flagellar beating and fluid flowing through a gap in the disc margin (dm) underneath the ventral groove

(vg) were proposed to generate suction-based attachment. In C, three different types of movements

associated with specific disc structural elements have also been predicted to generate contractile

forces that mediate attachment. Microribbon-crossbridge complex compression in "disc compression"

movements; the constriction of the lateral crest; or the rotation and sliding of the upper and lower

layers of the disc in the overlap zone region.

Figure 2: Three types of disc movements contribute to overall disc contraction. Using a lateral crest

marker to quantify disc movements (panel A, DAP12139GFP), disc movements were quantified using

live imaging of over 20 cells with kymographic analyses. In B, the rotational movements of the overlap

zone are defined and quantified. In C, overall disc compression movements (green line) that occur

within 25 seconds in attaching trophozoites are quantified using kymography analysis (D) with

summaries of movements of over 15 cells (E). In F, movements leading to constriction of the lateral

crest (magenta line) resulting in disc contraction are again quantified in kymographs (G) and with over

15 cells summarized (H). In panel I, disc contraction measures indicate contraction of up 5-10% of disc

diameter.

Figure 3: *Both N- and C-terminal ankyrin repeat domains are necessary for localization of DAP7268 to*

the disc body and overlap zone. Structured illumination (SIM) imaging (A) and wide field imaging

mNeonGreen-tagged DAP7268mNG (green) highlights the localization of this DAP to the disc overlap

zone and disc body relative to membrane -stained cells (B, CellMask, red). In C, an AlphaFold structural

predictions shows high confidence with the N- and C-terminal ankyrin repeats, but less confidence with

other alpha helical regions of DAP7268 (C). In panels D-G, localization of various DAP7268mNG

constructs with N-, C-, or both ankyrin repeat regions are indicated in inverted false colored images

(mNG = black).

Figure 4. Quadruple allele DAP7268KO mutants have aberrant multi-layered, yet smaller discs with

multiple lateral crests. SIM imaging of DAP7268KO and wild-type discs immunostained with delta-

giardin or beta-giardin highlight aberrant discs. The characteristic spiral array of wild type discs is

shown in A and B. (A) SIM with delta giardin immunostaining (B) negative staining of extracted disc. In

C and D, the aberrant structure of the DAP7268KO shown (C) DAP7268KO SIM, delta giardin

immunostaining. (D) DAP7268KD, negative staining of extracted disc. The 100% penetrance of the

DAP7268KO phenotype is seen in immunostained beta- and delta-giardin images (E-G) and with anti-

delta giardin (magenta) and anti-alpha tubulin (green) immunostaining (H). To visualize the lateral crest

in the DAP7268KO mutant, the DAP12139mNG lateral crest marker (green) was integrated into

DAP7268KO strains enabling the visualization of multiple lateral crests in the DAP7268KO mutant

background. Discs were immunostained with delta-giardin (magenta, J, K) and microtubules were also

immunostained with anti-alpha tubulin (TAT1) antibodies (yellow, L). In M, DAP7268KO mutant

phenotypes were scored based on the three most common mutant phenotypes (small, Baumkuchen,

or looped). Quantitation of overall disc doming angle of curvature (N), disc body area (O), disc

perimeter (P), and ventral groove height (Q) are summarized for delta-giardin GFP strains (dGGFP),

DAP7268 morpholino knockdown strains (7268KD), 7268KO, and wild type strains.

Figure 5: DAP7268KO mutants are defined by a multilayered "Baumkuchen" phenotype with

multiple aberrant MR-CB complexes and multiple lateral crests

Transmission electron microscopy with sectioning DAP7268KO whole cells highlight the aberrant and

extra spiral MT arrays (MTs) (A). In mutants, the microribbon-crossbridge complexes have aberrant

spacing (B). DAP7268KO strains have multiple overlap zone layers as seen in the comparison of TEM

cross sections of wild-type MR-CB complexes (C) with extra layered and shortened MR-CB complexes

(D). Panel E shows the additional overlap zone layers, each with MR-CB complexes stacked above one

another in serial block face SEM sections.

Figure 6: DAP7268KO "Baumkuchen" mutants lack proper disc contractility

To test disc contractility or other disc movements the Baumkuchen mutant, we integrated the

DAP12139mNG marker into the DAP7268KO strain and tracked disc contraction over several seconds

(A). The schematic in B indicated the doubled and aberrant lateral crests that showed reduced and

backwards overlap zone rotation movements as demonstrated using live imaging and kymograph

analysis (B). Baumkuchen mutants also have markedly reduced disc constriction or compression (C, D).

Figure 7: DAP7268KO strains have significantly reduced surface contacts and reduced ability to resist

flow stress associated with proper attachment.

In panel A, surface contacts associated with various early stages of trophozoite attachment (stages I-IV,

[12]) are scored in both DAP7268KO and wild-type strains. DAP7268KO mutants lack canonical surface

contacts, and instead have three additional attachment stage categories: stage III-LC and stage IV-LC

which both lack lateral crest contacts, and stage IV+B which is defined by aberrant bare area contacts

connecting to the lateral crest. Shear stress assays using 3 ml/minute flow (B) highlight the severe

defects in the ability of DAP7268KO mutants to resist shear stress, specifically, the reduced proportion

of attached cells following 20 seconds of shear stress as compared to the 7268Cas9 plasmid alone

strain, wild-type, DAP12139GFP tagged, and DAP7268KO/DAP12139mNG strains(C). N = 7 to 8

Asterisks indicate p values < 0.01 in t-tests.

Figure 8: DAP7268 attachment mutants have a significantly reduced impact on enterocyte barrier

degradation.

TEER measurements of ODM-based infections of wild-type (WBC6) trophozoites and two DAP7268KO

disc mutant clones (DAP7268-B2 and DAP7268-C7) are compared with mock (no *Giardia*) infections in

an 80-hour time period (A). In B, relative TEER measurements are compared at representative time

points (0, 48, 56, 72 hours post infection (hpi)), with significant differences indicated by T-tests (****).

Supplemental Figure S1. Movie of ventral disc contraction movements

AVI file associated with image stack used in Figure 2B.

Supplemental Figure S2. Movie of ventral disc compression movements

AVI file associated with image stack used in Figure 2D.

Supplemental Figure S3. Movie of ventral disc constriction movements

AVI file associated with image stack used in Figure 2F.

Supplemental Figure S4. Sequential knockout of all four DAP7268 alleles with antibiotic cassettes. Genomic
DNA from partial or complete DAP7268 knockout strains and from WBC6 was PCR-amplified with primers
7268LeftF and 7268RightR, which bind outside the homology arms of the hygromycin (Hyg), neomycin (Neo) and
blasticidin (Bsd) repair templates. Electroporation (EP) of the 7268Bsd repair template into the
Cas9/7268gRNA908R expression strain resulted in partial knockout of DAP7268, with a 1.6 kb wild-type
DAP7268 band and a 2.5 kb band indicating insertion of the 0.9 kb Bsd cassette into the gene. Quadruple
knockout was achieved after a second electroporation of either the 7268Neo or the 7268Hyg repair templates
into the 7268Bsd partial KO strain. The resulting quadruple KO strains lack the 1.6 kb wild-type DAP7268 band
and have either a 2.9 kb or 3.1 kb band indicating insertion of the 1.3 kb Neo cassette or the 1.5 kb Hyg
antibiotic cassette, respectively.

Supplemental Figure S5. Screening of quad KO in 7268BH clones with expression strain and wild-type controls.
Genomic DNA from 7268BH quadruple knockout clones (B2 through E11), the Cas9 and 7268gRNA908R
expression strain (Cas) and WBC6 (Wt) was PCR-amplified with primers 7268LeftF and 7268RightR, which bind
outside the homology arms of the hygromycin (Hyg) and blasticidin (Bsd) repair templates. The 7268BH
quadruple KO clones lack the 1.6 kb wild-type DAP7268 band and instead contain 3.1 kb and 2.5 kb bands
indicating insertion of the 1.5 kb hygromycin and 0.9 kb blasticidin antibiotic cassettes into all four copies of the
DAP7268 gene.

Supplemental Figure S6. Quadruple KO clones 7268BH B2 and quadKO plus knock-in of disc edge marker in
clones 7268BH_12139mNG 3B11 and 3E9. Genomic DNA from DAP7268 quadruple knockout strains and WBC6
was PCR-amplified with primers 7268LeftF and 7268RightR, which bind outside the homology arms of repair
templates used for knockout. All DAP7268 quadruple knockout strains lack the 1.6 kb wild-type DAP7268 band
present in WBC6 and have a 2.5 kb band indicating insertion of the 0.9 kb Bsd cassette into the gene. Insertion of
either the hygromycin resistance cassette (Hyg) or the hygromycin cassette plus DAP12139-mNeonGreen, a
marker of the disc edge, into the gene encoding DAP7268 results in 3.1 kb or 6.2 kb bands, respectively.

Supplemental Figure S7. DAP7268 morpholino-based knockdowns have severe disc and lateral crest structural defects resulting in flattened, closed disc conformation. Negative staining of isolated discs from both wild type (A, D) and DAP7268KD (B,C,E) demonstrate disc body (body), and multiple overlap zone (oz) and multiple and extraneous regions of the lateral crest (lc) defects as compared to the wild type disc conformation. Other components of the microtubule cytoskeleton are noted, including the dense bands (db), the basal bodies (bb), the funis (fn) and the four flagellar pairs (AF = anterior, CF = caudal, VF = ventral, and PF = posteriolateral). Scale bars = 1 µm (A-C) or 200 nm (D).

Supplemental Figure S8. Baumkuchen disc edge movie

AVI file associated with image stack used in Figure 6A.

Supplemental Figure S9. Wt compression of disc edge movie

AVI file associated with image stack used in Figure 6C.

Supplemental Figure S10. Wt constriction of disc edge movie

AVI file associated with image stack used in Figure 6C.

Supplemental Figure S11. 7268KO+ 12139 aberrant disc edge dynamics

AVI file associated with image stack used in Figure 6C.

<u>References</u>

1.   Einarsson E, Ma'ayeh S, Svard SG. An up-date on Giardia and giardiasis. Current opinion in

microbiology. 2016;34:47-52. doi: 10.1016/j.mib.2016.07.019. PubMed PMID: 27501461.

2.   Nosala C, Dawson SC. The Critical Role of the Cytoskeleton in the Pathogenesis of Giardia. Curr

Clin Microbiol Rep. 2015;2(4):155-62. doi: 10.1007/s40588-015-0026-y. PubMed PMID: 27347476;

PubMed Central PMCID: PMCPMC4918636.

3.   Savioli L, Smith H, Thompson A. *Giardia* and *Cryptosporidium* join the 'Neglected Diseases

Initiative'. Trends Parasitol. 2006;22(5):203-8. PubMed PMID: 16545611.

4.   Upcroft P, Upcroft JA. Drug targets and mechanisms of resistance in the anaerobic protozoa.

Clinical microbiology reviews. 2001;14(1):150-64.

5.   Land KM, Johnson PJ. Molecular basis of metronidazole resistance in pathogenic bacteria and

protozoa. Drug Resist Updat. 1999;2(5):289-94. PubMed PMID: 11504503.

6.   Barat LM, Bloland PB. Drug resistance among malaria and other parasites. Infect Dis Clin North

Am. 1997;11(4):969-87. PubMed PMID: 9421710.

7.   Upcroft J, Samarawickrema N, Brown D, Upcroft P. Mechanisms of metronidazole resistance in

Giardia and *Entamoeba*. Abstracts of the Interscience Conference on Antimicrobial Agents and

Chemotherapy. 1996;36(0):47.

8.   Bartelt LA, Sartor RB. Advances in understanding Giardia: determinants and mechanisms of

chronic sequelae. F1000Prime Rep. 2015;7:62. doi: 10.12703/P7-62. PubMed PMID: 26097735;

PubMed Central PMCID: PMCPMC4447054.

9.   Nosala C, Hagen KD, Dawson SC. 'Disc-o-Fever': Getting Down with Giardia's Groovy

Microtubule Organelle. Trends Cell Biol. 2018;28(2):99-112. Epub 2017/11/21. doi:

10.1016/j.tcb.2017.10.007. PubMed PMID: 29153830.

10.   Feely DE, Schollmeyer JV, Erlandsen SL. *Giardia* spp.: distribution of contractile proteins in the

attachment organelle. Exp Parasitol. 1982;53(1):145-54. PubMed PMID: 7056342.

11.   Crossley R, Holberton DV. Assembly of 2.5 nm filaments from giardin, a protein associated with

cytoskeletal microtubules in *Giardia*. Journal of cell science. 1985;78:205-31. PubMed PMID: 4093472.

12.   Crossley R, Holberton DV. Characterization of proteins from the cytoskeleton of *Giardia lamblia*.

Journal of cell science. 1983;59:81-103. PubMed PMID: 6863412.

13.   Holberton DV. Fine structure of the ventral disk apparatus and the mechanism of attachment in

the flagellate *Giardia muris*. Journal of cell science. 1973;13(1):11-41. PubMed PMID: 4729933.

14.   Holberton DV. Arrangement of subunits in microribbons from *Giardia*. Journal of cell science.

1981;47:167-85. PubMed PMID: 7263776.

15.   Friend DS. The fine structure of *Giardia muris*. J Cell Biol. 1966;29(2):317-32.

16.   Schwartz CL, Heumann JM, Dawson SC, Hoenger A. A detailed, hierarchical study of Giardia

lamblia's ventral disc reveals novel microtubule-associated protein complexes. PLoS One.

2012;7(9):e43783. doi: 10.1371/journal.pone.0043783. PubMed PMID: 22984443; PubMed Central

PMCID: PMC3439489.

17.   Brown JR, Schwartz CL, Heumann JM, Dawson SC, Hoenger A. A detailed look at the cytoskeletal

architecture of the Giardia lamblia ventral disc. J Struct Biol. 2016;194(1):38-48. doi:

10.1016/j.jsb.2016.01.011. PubMed PMID: 26821343; PubMed Central PMCID: PMCPMC4764415.

18.   Feely DE, Hoberton, D.V., Erlandsen, S.L. The Biology of *Giardia*. In: Meyer EA, editor. Giardiasis.

Human Parasitic Diseases. 3. Amsterdam: Elsevier; 1990. p. 11-50.

19.   House SA, Richter DJ, Pham JK, Dawson SC. Giardia flagellar motility is not directly required to

maintain attachment to surfaces. PLoS Pathog. 2011;7(8):e1002167. Epub 2011/08/11. doi:

10.1371/journal.ppat.1002167. PubMed PMID: 21829364; PubMed Central PMCID: PMC3150270.

20.   Holberton DV. Attachment of *Giardia*-a hydrodynamic model based on flagellar activity. J Exp

Biol. 1974;60(1):207-21. PubMed PMID: 4594683.

21.   Jones RD, Lemanski CL, Jones TJ. Theory of attachment in Giardia. Biophys J. 1983;44(2):185-90.

Epub 1983/11/01. doi: 10.1016/S0006-3495(83)84290-8. PubMed PMID: 6652213; PubMed Central

PMCID: PMCPMC1434824.

22.   Cheissin EM. Ultrastructure of Lamblia Duodenalis. I. Body Surface, Sucking Disc and Median

Bodies. J Protozool. 1964;11:91-8. PubMed PMID: 14119567.

23.   Holberton DV. Mechanism of attachment of *Giardia* to the wall of the small intestine.

Transactions of the Royal Society of Tropical Medicine and Hygiene. 1973;67(1):29-30. PubMed PMID:

4777416.

24.   Hansen WR, Fletcher DA. Tonic shock induces detachment of *Giardia lamblia*. PLoS Negl Trop

Dis. 2008;2(2):e169. PubMed PMID: 18270543.

25.   Hansen WR, Tulyathan O, Dawson SC, Cande WZ, Fletcher DA. Giardia lamblia attachment force

is insensitive to surface treatments. Eukaryot Cell. 2006;5(4):781-3. Epub 2006/04/12. doi: 5/4/781 [pii]

10.1128/EC.5.4.781-783.2006. PubMed PMID: 16607025; PubMed Central PMCID: PMC1459675.

26.   Woessner DJ, Dawson SC. The Giardia median body protein is a ventral disc protein that is

critical for maintaining a domed disc conformation during attachment. Eukaryot Cell. 2012;11(3):292-

301. doi: 10.1128/EC.05262-11. PubMed PMID: 22247266; PubMed Central PMCID: PMC3294447.

27.   Holberton DV, Ward AP. Isolation of the cytoskeleton from *Giardia*. Tubulin and a low-

molecular-weight protein associated with microribbon structures. Journal of cell science. 1981;47:139-

66. PubMed PMID: 7263775.

28.   Keister DB. Axenic culture of *Giardia lamblia* in TYI-S-33 medium supplemented with bile. Trans

R Soc Trop Med Hyg. 1983;77:487-8.

29.   Hagen KD, Hirakawa MP, House SA, Schwartz CL, Pham JK, Cipriano MJ, et al. Novel structural

components of the ventral disc and lateral crest in Giardia intestinalis. PLoS Negl Trop Dis.

2011;5(12):e1442. doi: 10.1371/journal.pntd.0001442. PubMed PMID: 22206034; PubMed Central

PMCID: PMC3243723.

30.   McInally SG, Hagen KD, Nosala C, Williams J, Nguyen K, Booker J, et al. Robust and stable

transcriptional repression in *Giardia* using CRISPRi. Molecular Biology of the Cell. 2018;in press.

31.   Schindelin J, Rueden CT, Hiner MC, Eliceiri KW. The ImageJ ecosystem: An open platform for

biomedical image analysis. Mol Reprod Dev. 2015;82(7-8):518-29. Epub 2015/07/15. doi:

10.1002/mrd.22489. PubMed PMID: 26153368; PubMed Central PMCID: PMCPMC5428984.

32.   Gillin FD, Reiner DS. Attachment of the flagellate Giardia lamblia: role of reducing agents,

serum, temperature, and ionic composition. Mol Cell Biol. 1982;2(4):369-77. PubMed PMID: 7110136.

33.   Nosala C, Hagen KD, Jones K, Loudermilk R, Nguyen K, Dawson SC. Giardia's ventral disc is

hyperstable and composed of over 80 disc-associated proteins. BioRxiv. 2018;361105 doi:

https://doi.org/10.1101/361105.

34.   Elmendorf HG, Dawson SC, McCaffery JM. The cytoskeleton of *Giardia lamblia*. Int J Parasitol.

2003;33(1):3-28. Epub 2003/01/28. doi: S002075190200228X [pii]. PubMed PMID: 12547343.

35.   Feely DE, Erlandsen SL. Isolation and purification of *Giardia* trophozoites from rat intestine. J

Parasitol. 1981;67(1):59-64. PubMed PMID: 7229820.

36.   Feely DE, Erlandsen SL. Effect of cytochalasin-B, low Ca++ concentration, iodoacetic acid, and

quinacrine-HCl on the attachment of *Giardia* trophozoites *in vitro*. J Parasitol. 1982;68(5):869-73.

PubMed PMID: 7131192.

37.   Inge PM, Edson CM, Farthing MJ. Attachment of *Giardia lamblia* to rat intestinal epithelial cells.

Gut. 1988;29(6):795-801. PubMed PMID: 3384364.

38.   Mariante RM, Vancini RG, Melo AL, Benchimol M. *Giardia lamblia*: evaluation of the *in vitro*

effects of nocodazole and colchicine on trophozoites. Exp Parasitol. 2005;110(1):62-72. PubMed PMID:

15804380.

39.   Sousa M, Concalves, CA, Bairos, VA, Poiares-Da-Silva, J. Adherence of *Giardia lamblia*

trophozoites to Int-407 human intestinal cells. Clin Diagn Lab Immunol. 2001;8:258-65.

40.   Davis-Hayman SR, Nash TE. Genetic manipulation of Giardia lamblia. Mol Biochem Parasitol.

2002;122(1):1-7. PubMed PMID: 12076765.

41.   Nosala C, Dawson SC. The ventral disc is a flexible microtubule structure that depends on a

domed conformation for functional attachment of *Giardia lamblia*. BioRxiv. 2017;213421. doi:

https://doi.org/10.1101/213421.

42.   Li J, Mahajan A, Tsai MD. Ankyrin repeat: a unique motif mediating protein-protein interactions.

Biochemistry. 2006;45(51):15168-78. Epub 2006/12/21. doi: 10.1021/bi062188q. PubMed PMID:

17176038.