Mechanosensing and Sphingolipid-Docking Mediate Lipopeptide-Induced Immunity in *Arabidopsis*

Jelena Pršić[1†], Guillaume Gilliard[2†], Heba Ibrahim[3], Anthony Argüelles-Arias[1], Valeria Rondelli[4], Jean-Marc Crowet[2], Manon Genva[2], W. Patricio Luzuriaga-Loaiza[1], Estelle Deboever[2], M. Nail Nasir[2], Laurence Lins[2], Marion Mathelie-Guinlet[5], Farah Boubsi[1], Sabine Eschrig[6], Stefanie Ranf[7], Stephan Dorey[8], Barbara De Coninck[3], Thorsten Nürnberger[9], Sébastien Mongrand[10], Monica Höfte[11], Cyril Zipfel[12], Yves F. Dufrêne[5], Alexandros Koutsioubas[13], Paola Brocca[4], Magali Deleu[2T]*, and Marc Ongena[1T]*

[1]Microbial Processes and Interactions laboratory and [2]Laboratory of Molecular Biophysics at Interfaces, TERRA teaching and research centre, Gembloux Agro-Bio Tech, University of Liège, Gembloux, 5030, Belgium

[3]Division of Plant Biotechnics, Department of Biosystems, KU Leuven and KU Leuven Plant Institute, Heverlee, 3001, Belgium

[4]Department of Medical Biotechnologies and Translational Medicine, Università degli Studi di Milano, Segrate, 20090, Italy

[5]Institute of Biomolecular Science and Technology, Louvain University, Louvain-la-Neuve, 1348, Belgium

[6] Chair of Phytopathology, School of Life Sciences Weihenstephan, Technical University of Munich, Freising, 85354, Germany

[7]Department of Biology, Faculty of Science and Medicine, University of Fribourg, Fribourg, 1700, Switzerland

[8]Université de Reims Champagne Ardenne, RIBP-USC INRAE 1488, 51100 Reims , France

[9]Centre of Plant Molecular Biology, Eberhard-Karls-University Tübingen, Tübingen, 72076,

Germany

[10]Laboratoire de Biogenèse Membranaire, University of Bordeaux, Villenave d'Ornon, 33882,

France

[11]Laboratory of Phytopathology, Faculty of Bioscience engineering, Ghent University, Ghent,

9000, Belgium

[12]Department of Plant and Microbial Biology, Zurich-Basel Plant Science Center, University of

Zurich, Zurich, 8006, Switzerland

[13]Jülich Centre for Neutron Science, Heinz Maier-Leibnitz Zentrum, Forschungszentrum Jülich

GmbH, Garching, 85748, Germany

[†] JP and GG contributed equally to this article and share first authorship.

[ᵀ] MD and MO contributed equally to the design and global supervision of the study and share

last authorship.

*Corresponding authors: M.D. (magali.deleu@uliege.be) and M.O. (marc.ongena@uliege.be)

Abstract

Bacteria-derived lipopeptides are immunogenic triggers of host defenses in metazoans and plants. Root-associated rhizobacteria produce cyclic lipopeptides that activate systemically induced resistance (IR) against microbial infection in various plants. How these molecules are perceived by plant cells remains elusive. Here, we reveal that immunity activation in *Arabidopsis thaliana* by the lipopeptide elicitor surfactin is mediated by docking into specific sphingolipid-enriched domains and relies on host membrane deformation and subsequent activation of mechanosensitive ion channels. This mechanism leads to host defense potentiation and resistance to the necrotroph *B. cinerea* but is distinct from host pattern recognition receptor-mediated immune activation and reminiscent of damage-induced plant immunity.

Main Text

Lipopeptides (LPs) represent a prominent and structurally heterogeneous class of molecules among the broad spectrum of small specialized metabolites synthesized by bacteria. Besides serving key functions for the ecological fitness of the producer (motility, biofilm formation, colonization, nutrient acquisition, or antagonism towards competing neighbors), some LPs also act as triggers of immune responses that restrict pathogen infection of metazoans and plants[1,2]. The vast majority of LPs formed by plant-associated bacteria are comprised of a partly or fully cyclized oligopeptide linked to a single fatty acid chain. Some of these cyclic lipopeptides (CLP) formed by beneficial species belonging to the *Pseudomonas* and *Bacillus* genera are potent elicitors of immune responses in the host plant leading to a systemically induced resistance (IR) against infection by microbial pathogens[2,3].

This CLP-induced plant resistance is a key process for biocontrol of crop diseases [4], but, in

contrast to pattern-triggered immunity (PTI), its molecular basis remains poorly understood.

Like in animals, PTI in plants relies on the detection of specific molecular motifs (Microbe-

Associated Molecular Patterns (MAMPs) via cell-surface plasma membrane (PM)-localized

Pattern-Recognition Receptors (PRRs)[5]. Upon assembly of higher order receptor complexes

involving conserved co-receptors, PRRs activate receptor-like cytoplasmic kinases (RLCKs)

such as BIK1 and its closest homolog PBL1 described as key convergent signaling hubs. This

leads to phosphorylation of numerous substrate proteins and subsequent induction of a

well-characterized immune response[6]. Early hallmarks of PTI signaling in plants include

apoplastic burst of reactive oxygen species ($[ROS]_{apo}$), calcium influx, medium alkalinization

indicating H^+/K^+ exchange and membrane depolarization, MAPK phosphorylation cascade

and initiation of transcriptional reprogramming [7,8,9,10].

The CLP surfactin (Srf, **Fig 1A**) is well conserved in plant beneficial bacilli[11] and is

among the bacterial compounds best described as immunity elicitor in several plant

species[2]. In *Arabidopsis thaliana* ecotype Col-0 (hereafter, *Arabidopsis*), root treatment with

purified Srf (at 10 µM as minimal active concentration previously determined[12] and used as a

mix of naturally produced homologues slightly differing in the length of the fatty acid tail,

see **Suppl Fig 1**) triggers IR and significantly reduces leaf infection by the grey mold pathogen

Botrytis cinerea (**Fig 1B**). Therefore, we used Srf as a model to further investigate the

molecular mechanisms determining CLP perception and immunity stimulation in *Arabidopsis*

root cells*.

We first performed quantitative and time-resolved measurements of early responses

commonly associated with MAMP perception in *Arabidopsis* and other plants. $[ROS]_{apo}$ burst

is almost invariably associated with PTI[7] but, by contrast to treatment with the MAMP

flagellin-derived peptide flg22 or with chitin, we did not observe a $[ROS]_{apo}$ burst in *Arabidopsis* root cells treated with Srf based on a horseradish peroxidase-luminol assay (**Suppl Fig 2**). Srf-mediated IR against *B. cinerea* is fully conserved in the *rbohD* mutant lacking functional plasma membrane NADPH oxidase RBOHD responsible for MAMP-induced $[ROS]_{apo}$ burst[13] (**Suppl Fig 3**). Hence, Srf-mediated activation of IR in the root does not require RBOHD[14]. However, Srf triggered a fast and consistent increase in intracellular ROS ($[ROS]_{intra}$) in root loaded with the fluorescent probe DCFH-DA (**Fig 1C**). This Srf-triggered $[ROS]_{intra}$ burst is also observed in the *rbohD* mutant (**Suppl Fig 4**), suggesting it is not caused by the uptake of apoplastic ROS via aquaporins (see **Suppl Fig 5** for response to flg22) but may originate from different organelles as reported for abiotic stresses or other small microbial compounds[15,16,7,17]. Calcium influx typically associated with PTI in plants[8] was tested upon elicitation by Srf using an aequorin-based bioluminescence assay. It did not reveal any significant Ca^{2+} increase ($[Ca^{2+}]_{cyt}$) in root of the Col-0^{AEQ} reporter line in contrast to the increase observed upon flg22 treatment (**Fig 1D**) or in response to chitin (**Suppl Fig 6**). On the other hand, medium alkalinization occurs within minutes after Srf treatment (**Fig 1E**), which indicates H^+/K^+ exchange possibly leading to membrane depolarization[9]. However, no significant increase in conductivity was measured in the medium following Srf treatment (**Suppl Fig 7**) indicating that the lipopeptide does not affect plasma membrane (PM) integrity and does not cause massive electrolyte leakage. Cell viability tests confirmed that Srf is not toxic for *Arabidopsis* root cells at concentrations up to 50 µM (**Suppl Fig 8**).

Next, we explored early changes in the root transcriptome profile induced by Srf via time course RNAseq analysis (30 min, 1h, 3h and 6h post treatment) using the same setup previously reported for flg22 and the fungal MAMP chitin[18]. Data revealed a relatively low transcriptional response to Srf elicitation over all sampling times with a total of 564

differentially expressed genes (DEGs, Log$_2$ Fold Change > 2, p<0.05; **Fig 1F**) compared to approximately 5000 DEGs and 2000 DEGs reported upon flg22 and chitin treatment respectively[18]. While MAMPs mainly up-regulate early responsive genes (30 min – 1 h)[18,19], an almost equal number of up- and down-regulated DEGs were observed upon Srf treatment at all time points (**Fig 1F**), with about half of the transcriptional changes specific to Srf elicitation (47,9% and 58% compared with flg22 and chitin respectively)[18]. Strikingly, many of the Srf down-regulated genes are upregulated by flg22 and chitin[18] (**Fig 1F, Suppl Table 1**). Differential expression was confirmed by quantitative RT-PCR performed on some selected genes in plantlets elicited with the lipopeptide and with chitin (**Suppl Fig 9**). More specifically, the expression of genes typically associated with early immune signaling (receptor-like kinases, [ROS]$_{apo}$ burst, calcium signaling or MAPK phosphorylation cascade[10]) or defense mechanisms (pathogenesis-related (PR) proteins, callose deposition, lignification) is not modulated or down-regulated by Srf by contrast with MAMP treatment (**Fig 1G, Suppl Table 1**).. However, *CYP71A12*, encoding a key enzyme of the camalexin biosynthesis pathway[20], is among the late-responsive genes (6h) strongly stimulated by Srf. In accordance, we measured significantly higher amounts of this phytoalexin, which is toxic to *B. cinerea*[21,22], in infected leaves of Srf-treated plants compared with mock treatment (**Fig 1H**). The key role of camalexin in disease control was confirmed by the loss of Srf-triggered resistance in the *pad3* mutant[23] unable to form camalexin[22] (**Fig 1I**). Thus, by contrast to PTI which is associated with substantial transcriptional reprogramming[19], immunity stimulation by Srf does not lead to major changes in the expression of genes involved in signaling and defense.

Since the molecular basis of Srf-induced immune activation is signal-specific, we hypothesized that plant cells perceive lipopeptides by a mechanism that differs from pattern

sensing. Srf possesses both a peptidic moiety and a fatty acid tail, but its IR-eliciting potential

is fully conserved in *Arabidopsis* mutants lacking functional PRRs that recognize either

bacterial proteinaceous immunogenic patterns or acyl chain epitopes such as medium chain

3-hydroxy fatty acids and HAAs[24,25] (**Suppl Fig 10**). Srf elicitation is not significantly affected

either in mutants lacking co-receptors required for proper functioning of a wider range of

PRRs detecting immunogenic peptides such as Pep1[26], nlp20[27] and IF1[28] nor in the *bik1 pbl1*

double mutant lacking RLCKs that act downstream of the PRR-co-receptor complexes (**Suppl**

Fig 10). Although we only tested a small subset of the multitude of PRRs potentially

expressed in *Arabidopsis*[6] and although early cellular signaling may be BIK1/PBL1-

independent[29], our data strongly suggest that *Arabidopsis* does not sense Srf via PRR-type

cell surface sentinels. This is in accordance with previous data from tobacco, which showed

that Srf is still active on protease-treated cells and that there is no refractory state upon

repeated Srf treatment unlike typically observed for PTI[30].

Due to their amphipathicity, CLPs readily interact with biological membranes, causing

pore formation and membrane disruption responsible for their antimicrobial activities[31].

Such an adverse effect is not expected on plant membranes, but we hypothesized that Srf

perception by root cells might primarily rely on its interaction with the lipid phase of the PM.

Complex sphingolipids glucosylceramides (GluCer) and glycosyl inositol phosphorylceramides

(GIPC) constitute more than 30% of *Arabidopsis* PM lipids and are key components required

for membrane integrity and functionality, notably by forming ordered nano-domains with

sterols[32,33,34]. *In silico* docking simulation first revealed a more favorable interaction of Srf

with GluCer or GIPCs than with the other typical plant PM lipids PLPC (1-Palmitoyl-2-

linoleoyl-sn-glycero-3-phosphocholine as phospholipid) and ß-sitosterol (as main sterol) (**Fig**

2A). To test this experimentally, we generated biomimetic liposomes using commercially

available GluCer, PLPC and ß-sitosterol. Isothermal titration calorimetry performed on liposomes with increasing composition complexity in such lipids showed the highest binding affinity of Srf to model membranes containing GluCer (**Fig 2B**). In support of a preferential interaction with sphingolipids, molecular dynamic (MD) simulation on the same ternary lipid system showed the specific insertion of Srf in the vicinity of GluCer molecules or in GluCer-enriched areas in the membrane (**Fig 2C**). In light of these results, we tested Srf elicitor activity on the *Arabidopsis* ceramide synthase mutant *loh1* (LONGEVITY ASSURANCE 1 HOMOLOG1) which is depleted in these complex sphingolipids[35,36]. We observed strongly reduced $[ROS]_{intra}$ responses (**Fig 2D**) as well as loss of IR to *B. cinerea* infection in *loh1* compared to wild-type plants (**Fig 2E**). Such lipid-dependent $[ROS]_{intra}$ elicitation was also observed for other IR-eliciting CLPs such as orfamide and WLIP[2] isolated from beneficial pseudomonads that resemble Srf in size and amphiphilic character (**Suppl Fig 11**). The CLP immunogenic activity thus relies on an intricate interaction with PM sphingolipids as reported for other microbial compounds[36,37,38].

By inserting into lipid bilayers, Srf may transiently affect the local structure of membranes. Indeed, neutron reflectivity (NR) experiments (see **Suppl Fig 12** for deuterated Srf synthesis and characterization) demonstrate that Srf exclusively inserts into the outer leaflet of PLPC-ß-sitosterol-GluCer model membranes (**Fig 3A**). This is supported by MD simulation showing that the Srf peptide backbone preferentially positions at the level of the polar lipid heads of the membrane (**Suppl Fig 13**). Srf insertion does not affect the lipid chain-chain interaction as shown by WAXS and FTIR (**Suppl Fig 13**). In addition, NR data indicated that Srf insertion results in a decrease in membrane thickness (from 40 to 36Å), which is more pronounced in ternary membranes than in membranes lacking GluCer (from 43 to 41Å) (**Suppl Table 4**). Analysis of the nanoscale morphology of supported PLPC-ß-

sitosterol-GluCer bilayers by Atomic Force Microscopy confirmed this membrane thinning

caused by Srf insertion (**Suppl Fig 14**). An additional impact of the lipopeptide on PM

physical properties was derived from coarse-grained MD simulation which revealed a strong

curvature-inducing effect mediated by Srf docking on ternary membranes (**Fig 3B**).

In light of these biophysical data, a clear impact of Srf on PM structure can be

predicted but in integral root cells, the PM is physically connected to the thick and

mechanically strong cell wall polymer matrix, which provides structural support and might

stabilize the membrane into a flat conformation under low tension[39]. We thus next tested

early Srf-induced immune responses in cell wall-free protoplasts. Use of PPs renders the PM

more susceptible to deformation, which was used to study responses to cell swelling or

shrinkage/expansion during osmotic stresses[40,41]. As in root cells, Srf triggered a consistent

$[ROS]_{intra}$ burst in freshly isolated protoplasts (**Suppl Fig 15**). However, in contrast to roots, a

significant calcium influx, was observed in Srf-treated protoplasts by using the Col-0^{AEQ}

reporter line and also by loading Col-0 with the Fluo4-AM probe (**Fig3C and Suppl Fig 16**).

This Srf-induced Ca^{2+} influx is comparable in amplitude to the one induced by MAMPs (**Suppl**

Fig 17). It involves some PM channels since it is abolished in protoplasts pre-treated with the

general channel blocker $LaCl_3$ (**Suppl Fig 18**) and considering that the lipopeptide does not

cause any detrimental effect on protoplast viability at the concentration used (**Suppl Fig 19**).

Additional assays on protoplasts revealed that activation of early responses by Srf requires

threshold concentrations of 5-10 μM both for calcium influx (**Fig 3D**) and $[ROS]_{intra}$ burst

(**Suppl Fig 20**), which is much higher than MAMPs detected at nanomolar concentrations.

This further indicates that Srf perception is not mediated by a high-affinity receptor-based

detection system and is in accordance with the mechanism predicted from biophysics in

which threshold amounts of Srf molecules must dock into sphingolipid domains in order to

modulate PM structure. We tested the impact of the lipopeptide on protoplast membrane fluidity via measurements of laurdan generalized polarization (laurdan GP) related to the lipid bilayer order. Our results show that Srf treatment led to a significant increase in ΔGP values indicating a clear membrane rigidification effect as also observed upon interaction of the lipopeptide with PM mimicking liposomes (**Fig 3E**).

Altogether, these data obtained with protoplasts support the relevance of PM deformation in the response to Srf. We therefore hypothesized that insertion of the CLP could induce physical constrains resulting in increased lateral tension sufficient for activating mechano-sensitive (MS) ion channels, in a process similar to the one observed for some anionic amphipathic chemicals[42,43]. This was supported by the reduced calcium influx observed upon pre-treatment of Col-0^{AEQ} protoplasts with the specific MS channel blocker GsMTX-4 (**Suppl Fig 21**). Among stretch sensitive mechanosensors identified so far in plant cells, MSL9, MSL10 and MCA1/2 localize in the PM [44,45,46] but do not require RLCK-mediated phosphorylation of the cytoplasmic domains for gating unlike other MS ion channels such as OSCA1.3 which needs BIK1 phosphorylation to be activated [47]. Using Fluo-4, we thus tested protoplasts prepared from the quintuple *msl4/5/6/9/10*[48] and the double *mca1/2*[49] mutants for their response to Srf and observed a significantly decreased calcium influx, to the same extent as chemical inactivation with GsMTX-4 in Col-0 (**Fig 3F**).

We next evaluated the effect of inactivation or knock-out of *msl* and *mca* channels on intracellular ROS burst as early response of root tissues elicited by Srf. Pre-treatment with GsMTX-4, LaCl$_3$ or with the Ca^{2+} chelator EGTA eliminated the ROS burst triggered by Srf in root tissues (**Fig 3G**), supporting the importance of MS channels in the response and indicating that ion fluxes acts upstream of or are interdependent of [ROS]$_{intra}$[8]. An almost complete loss of [ROS]$_{intra}$ burst was also observed upon Srf treatment in the *msl4/5/6/9/10*

and *mca1/2* mutants as compared to Col-0 (**Fig 3H**). In addition, *mca1/2* and *msl4/5/6/9/10*

plants were strongly impaired in mounting systemic resistance against *B. cinerea* upon Srf

treatment (**Fig 3I**), further indicating that functional MS channels are necessary for full

response of *Arabidopsis* to Srf elicitation on roots. Data on protoplasts show that Srf may

trigger some calcium transients as early immune-related event but a detectable Ca^{2+} influx is

not required for defense activation in plantlets, which correlates with the fact that no

downstream components of calcium signaling are up-regulated upon perception of the

lipopeptide.

Collectively, although contributions of other channels cannot be ruled out[47], our data

provides evidence for a key role of PM-located mechanosensors in lipopeptide-induced plant

defenses. The relative contribution of each channel remains to be determined as they

display specific properties in terms of sensitivity to membrane tension and ion selectivity.

MCA1/2 are described as genuine transporters of Ca^{2+}[46,50] while MSL10 is regarded as a non-

selective ion transporter that is indirectly involved in calcium signaling upon wounding[51] and

response to hypo-osmotic shock in cell swelling[41]. Both channels may thus act in a

coordinated fashion to tailor ion fluxes leading to cellular responses and PM depolarization.

As previously reported for other plant species[2], treatment with Srf prepares *Arabidopsis* to

mount defense responses culminating in the systemically expressed IR phenotype. We

provide new insights into the molecular basis of the well-known long-standing process of

CLP-triggered plant immunity activation by unveiling a new lipid-mediated mechanism for

the detection of these molecules at the cell surface. We infer from our data that CLP

insertion into sphingolipid-enriched PM domains causes deformation and increases lateral

tension in the membrane leading to rearrangement of the MS protein complexes and gating

of the channels. This allows ion influx and initiates chemical signaling that can be integrated

by root cells to activate early immune responses in a process that remains to be deciphered.

Such a lipid-dependent perception at the cell surface may apply also to other bacterial

amphiphilic IR elicitors such as acyl-homoserine lactones and rhamnolipids which also

readily interact with membrane lipids and may thus be perceived via similar

mechanisms[2,52,53,54,55]. The nature of PM lipids widely varies across plant species[34]. This could

explain, at least in the case of Srf, why this molecule triggers immunity in dicots but is not

very active on monocots[2]. We assume that the effect of a CLP on a particular target

membrane is also fine-tuned by precise structural traits in the molecule. It may explain why

some CLPs produced by *Pseudomonas* leaf pathogens act as virulence factors in a wide range

of plants by causing necrosis via pore formation in cellular membranes[56]. However, further

investigation is required to capture the physico-chemical rules governing lipid selectivity and

CLP insertion dynamics.

As the two components of the plant immune system, PTI works in concert with

effector-triggered immunity (ETI) for mounting robust defense responses to biotrophic

invaders but ETI is not efficient against necrotrophic pathogens[5,57]. Here, we describe a

novel molecular mechanism of defense activation in plants, which provides resistance to the

necrotroph *B. cinerea* via a unique process not related to the receptor-based surveillance

system involved in the recognition of MAMPs by plant cells or in the perception of the

Pam_3CSK_4 analog of triacylated lipopeptides produced by *Staphylococcus aureus* and acting

as agonists of Toll Like-type PRRs in metazoans[58]. It therefore provides new insights in plant-

microbe interactions mediated by small chemicals from beneficial bacteria. Collectively, our

data show that Srf perception leads to specific immune activation signature regarding the

type, timing, and amplitude of early defense-related events and the weak transcriptional

reprogramming as compared to PTI. This may explain why elicitation by Srf is cost-effective

for the host plant as it does not result in growth-defense trade-off[59,60] nor does it cause a

strong response associated with the alertness state or a hypersensitive reaction leading to

cell death. Using CLPs as elicitors would enable bacteria to bypass a strong immune response

and avoid their rejection as undesirable associate. Further investigations are needed for a

comprehensive understanding of the whole process from perception to systemic signaling

but the mechanistic basis of CLP-induced plant resistance reported here should contribute to

rationally implement the use of these compounds or their producers as bio-sourced

alternatives to chemicals in sustainable agriculture.

Fig. 1. Surfactin triggers systemic resistance in *Arabidopsis* associated with atypical immune responses. (A) Structural model of the heptapeptide Srf (C14 acyl chain homologue) in water (Gromacs v.4.5.4). Red: oxygen, white: hydrogen, dark blue: nitrogen, light blue: carbon. The polar amino acids are circled in yellow, and other amino acids and the acyl chain constitute the non-polar part of the molecule. **(B)** Disease incidence caused by *Botrytis cinerea* in *Arabidopsis* Col-0 plants pre-treated with Srf (10 μM) or not (mock treatment, 0.1 % ethanol) (n=28 replicates from three independent experiments). The box plots encompass the 1st and 3rd quartiles, the horizontal line indicates the median, and bars extend from the lower to the higher values. Disease reduction (D.R.) is calculated from the mean values of both treatments. Significant difference ***$P<0.001$, two-tailed *t*-test. **(C)** Burst in intracellular ROS species in *Arabidopsis* Col-0 roots upon treatment with Srf (10 μM). Left, time course of $[ROS]_{intra}$ accumulation (Relative Fluorescence Units, RFU) with data at each time point representing mean ± SD, n=3 (independent root samples). Right, fold increase in fluorescence values ± SD, at 30 min after the addition of Srf compared to mock-treated roots. Data are pooled from three independent experiments (total n=9) and asterisks indicate significant difference (***$P<0.001$, two-tailed *t*-test). **(D)** $[Ca^{2+}]_{cyt}$ kinetics in Srf-treated (10 μM) or flg22-treated root tissues (1 μM) (n=6) compared to mock treatment in the *Arabidopsis* Col-0^{AEQ} reporter line. Results are represented as luminescence counts per second relative to total luminescence counts remaining (L/Lmax; mean ± SD). Experiments were repeated three times with similar results. **(E)** pH variation in Col-0 root medium following mock treatment or addition of 10 μM Srf. Values on the graph are normalized to pH of the first time point ± SD and are from one representative experiment (n=4) out of 2 independent experiments showing similar results. **(F)** Number of DEGs (Log$_2$ Fold Change > 2, $P<0.05$) in *Arabidopsis* root cells determined via RNAseq for each time point in response to

Srf treatment (10 μM). Our data were compared with those reported for DEGs in response

to flg22 (1 μM) and chitin (Chi, 1 mg/ml)[18] and bars are subdivided by the number of genes

specifically responding to Srf and by the number of genes differentially (oppositely)

regulated by Srf and the two MAMPs. **(G)** Heatmap of the expression of genes putatively

associated with plant immune responses (listed in Supp table 1) that were modulated upon

Srf treatment (S, left) (10 μM) and compared with their expression in response to flg22 (1

μM) and chitin (1 mg/ml) (F and C respectively, right) based on published data[18]. Colour

scale represents Log_2 FC (> 2, P<0.05). **(H)** Camalexin response associated with IR triggered

by Srf. Camalexin accumulation 96 hours post *B. cinerea* inoculation (hpi) in *Arabidopsis* Col-

0 leaves of mock- or 10 μM Srf-treated plants at the root level. Graph shows values obtained

in one experiment with each value representing a sample of five plants pooled together.

Asterisks indicate significant difference with ns, not significant; *$P<0.05$; ***$P<0.001$; two-

tailed *t*-test. **(I)** Disease incidence of *B. cinerea* in *pad3* mutant pretreated with 10 μM Srf or

mock-treated at the root level (n=30, values obtained from three independent experiments,

presented as differently shaded grey values). Data are represented as in fig 1B. Asterisks

indicate significant difference with ns, not significant; *$P<0.05$; ***$P<0.001$; two-way ANOVA

and Sidak's multiple-comparison post-test.

Fig. 2. Affinity for sphingolipids determines CLP-triggered immunity. **(A)** *In silico* docking

simulation of the interaction between Srf and plant PM lipids with their associated energy of

interaction (E_{int}). A lower E_{int} value indicates a more favorable interaction. Hydrogen, oxygen

and phosphate atoms are respectively represented in grey, red and blue. Carbon atoms of

Srf are in yellow and carbon atoms of GluCer, Sito and PLPC are in pink. **(B)** Binding

coefficient (K) of Srf to liposomes with different lipid compositions. Graph presents values

from two independent experiments, mean ± SD. **(C)** Molecular dynamics simulation of Srf

insertion in GluCer-enriched domains of a PLPC-Sito-GluCer bilayer. Left: Top views of

bilayers before and after Srf insertion (right). **(D)** [ROS]$_{intra}$ accumulation in roots of Col-0 and

loh1 mutant. Data represents fold increase in fluorescence values ± SD (n=6 from two

independent experiments) at 30 min after Srf addition (10 μM) or not . Significant difference

***$P<0.001$, two-tailed *t*-test. **(E)** Disease incidence of *B. cinerea* in *Arabidopsis* Col-0 and

loh1 mutant plants, pre-treated with Srf (10 μM) compared with mock treatment (n=30 from

two independent experiments). Data are represented as in fig 1B. ns = not significant,

***$P<0.001$, two-way ANOVA and Sidak's multiple-comparison post-test.

Fig. 3. Srf causes membrane deformation and activates mechanosensitive channel-

dependent immune responses. (A) Membrane thickness determined via neutron scattering

length density (SLD) profiles of supported PLPC-Sito-GluCer membrane before (black) and

after (green) Srf addition to the final 95:5 membrane: Srf molar proportion (0.24 μM)

(below). Illustration (above) presents the correspondence between regions in the SLD profile

and specific zones in the membrane. **(B)** Molecular dynamics simulation of Srf-induced

membrane curvature. **(C)** Left, [Ca^{2+}]$_{cyt}$ kinetics in Srf-treated (10 μM) root cell protoplasts

(n=6) compared to mock treatment in the *Arabidopsis* Col-0^{AEQ} reporter line. Results are

represented as luminescence counts per second relative to total luminescence counts

remaining (L/Lmax; mean ± SD). Experiments were repeated three times with similar results

(see Suppl Fig 16 for additional experiments). Right, increase in [Ca^{2+}]$_{cyt}$ detected upon

loading root protoplasts of Col-0 with Fluo-4 in mock- or Srf-treated (10 μM). Experiments

were repeated three times with similar results. **(D)** Dose-dependent [Ca^{2+}]$_{cyt}$ increase

induced by Srf in root protoplasts of *Arabidopsis* Col-0^{AEQ}. Values are the average of L/Lmax

values from 1.5 to 4 min after treatment corresponding to the top of the peak. Mean ± SD of at least 10 technical replicates from at least five independent experiments. Asterisks indicate statistically significant differences to the mock treatment (ns= no significant difference; *P<0.05; ***P < 0.001; (a) two-tailed *t*-test; (b) Welch and Brown-Forsythe ANOVA). **(E)** Change of laurdan generalized polarization (ΔGP) in Srf-treated (10 µM) Col-0 root protoplasts and in liposomes reflecting a change of membrane rigidity. ΔGP is defined as the subtraction of GP measured at 10 min following treatment and GP measured before treatment. Mean ± SD of 12 (for protoplasts) and 15 (for liposomes) replicates from 8 (for protoplasts) and 5 (for liposomes) independent experiments. ***P<0.001, two-way ANOVA and Sidak's multiple comparison test. **(F)** $[Ca^{2+}]_{cyt}$ response measured with Fluo-4 upon Srf elicitation (10 µM) in *Arabidopsis* Col-0 root protoplasts with and without pre-treatment with the mechanosensitive channel blocker GsMTX-4 (10 min incubation, 7.5 µM)(n=10) and in root protoplasts of the *mca1/2*, and *msl4/5/6/9/10* mutants (n=14). Mean ± SD from four independent experiments. Letters represent statistically different groups at α = 0.05 (two-way ANOVA and Tukey's multiple-comparison post-test). **(G)** $[ROS]_{intra}$ accumulation upon addition of 10 µM Srf to *Arabidopsis* Col-0 roots upon pre-treatment or not (Col-0, n=16) with the mechanosensitive channel blocker GsMTX-4 (10 min incubation, 7.5 µM) (n=12), with the non-selective Ca^{2+} channel blocker $LaCl_3$ (10 mM) (n=7) and the Ca^{2+} chelator EGTA (1 mM) (n=7). Data represent fold increase in fluorescence values 30 min after Srf addition compared to mock-treated roots. Mean ± SD calculated from data from two independent experiments. Letters represent statistically different groups at α = 0.05 (two-way ANOVA and Tukey's multiple-comparison post-test). **(H)** $[ROS]_{intra}$ accumulation in *Arabidopsis* Col-0 (n=6), *mca1/2* (n=7), and *msl4/5/6/9/10* (n=8) roots following Srf treatment (10 µM). Data represent fold increase in fluorescence values 30 min after Srf addition compared to mock-

treated roots. Mean ± SD from two independent experiments. **P<0.01, two-tailed *t*-test. **(I)**

Disease incidence of *B. cinerea* in *Arabidopsis* Col-0, *mca1/2*, and *msl4/5/6/9/10* mutant

plants, mock- or Srf pre-treated (10 μM)(each n=30 from two independent experiments

represented as differently shaded grey values). Data are represented as in fig 1B.

Acknowledgments

We are grateful to Prof. Ivo Feussner at University of Goettingen for providing us with

Arabidopsis lipid mutants. We thank Aurélien Legras, Sébastien Steels, Catherine Helmus, and

Adiilah Mamode-Cassim for their excellent technical support.

Funding

This work was supported by the EU Interreg V France-Wallonie-Vlaanderen portfolio

SmartBiocontrol (Bioscreen and Bioprotect projects, avec le soutien du Fonds européen de

développement régional - Met steun van het Europees Fonds voor Regionale Ontwikkeling),

by the European Union Horizon 2020 research and innovation program under grant

agreement No. 731077, by the PDR surfasymm (T.0063.19) from F.R.S.-FNRS (National Funds

for Scientific Research in Belgium) and by the EOS project ID 30650620 from the FWO/F.R.S.-

FNRS. G.G. is recipient of a F.R.I.A. fellowship (F.R.S.-FNRS), M.D. and M.O. are respectively

senior research associate and research director at the F.R.S.-FNRS.

Author contributions

J.P., G.G., M.D. and M.O. conceived and designed experiments; J.P., G.G., H.I., A.A., V.R.,

W.P.L-L, E.D., M.N.N., M.M-G., S.E., A.K., P.B. and Y.F.D. performed experiments and analyzed

data; J-M.C., M.G. and L.L. performed modeling; S.D., B.D.C, S.R., T.N., S.R., M.H. and C.Z.

substantially revised the manuscript and were involved in the discussion of the work; M.D.

and M.O. supervised the study and provided funding.

Competing interests: The authors declare no competing interests.

References

1. Mohammad, M. *et al.* Staphylococcus aureus lipoproteins in infectious diseases.

Front. Microbiol. **13**, 1–17 (2022).

2. Pršić, J. & Ongena, M. Elicitors of Plant Immunity Triggered by Beneficial Bacteria.

Front. Plant Sci. **11**, 1–12 (2020).

3. Cesa-Luna, C. *et al.* Charting the Lipopeptidome of Nonpathogenic Pseudomonas.

mSystems **8**, (2023).

4. Blake, C., Christensen, M. N. & Kovacs, A. T. Molecular aspects of plant growth

promotion and protection by bacillus subtilis. *Mol. Plant-Microbe Interact.* **34**, 15–25

(2021).

5. Zhang, L., Hua, C., Janocha, D., Fliegmann, J. & Nürnberger, T. Plant cell surface

immune receptors-Novels insights into function and evolution. *Curr. Opin. Plant Biol.*

74, 102384 (2023).

6. DeFalco, T. A. & Zipfel, C. Molecular mechanisms of early plant pattern-triggered

immune signaling. *Mol. Cell* **81**, 3449–3467 (2021).

7.   Waszczak, C., Carmody, M. & Kangasjärvi, J. Reactive Oxygen Species in Plant

Signaling. *Annu. Rev. Plant Biol.* **69**, 1–28 (2018).

8.   Köster, P., DeFalco, T. A. & Zipfel, C. Ca 2+ signals in plant immunity . *EMBO J.* **41**,

(2022).

9.   Falhof, J., Pedersen, J. T., Fuglsang, A. T. & Palmgren, M. Plasma Membrane H+-

ATPase Regulation in the Center of Plant Physiology. *Mol. Plant* **9**, 323–337 (2016).

10.   Meng, X. & Zhang, S. MAPK cascades in plant disease resistance signaling. *Annu. Rev.*

Phytopathol. **51**, 245–266 (2013).

11.   Harwood, C. R., Mouillon, J.-M. M., Pohl, S. & Arnau, J. Secondary metabolite

production and the safety of industrially important members of the Bacillus subtilis

group. *FEMS Microbiol. Rev.* **42**, 721–738 (2018).

12.   Cawoy, H. *et al.* Plant Defense Stimulation by Natural Isolates of *Bacillus* Depends on

Efficient Surfactin Production. *Mol. Plant-Microbe Interact.* **27**, 87–100 (2014).

13.   Torres, M. A., Dangl, J. L. & Jones, J. D. G. Arabidopsis gp91phox homologues Atrbohd

and Atrbohf are required for accumulation of reactive oxygen intermediates in the

plant defense response. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 517–522 (2002).

14.   Morales, J., Kadota, Y., Zipfel, C., Molina, A. & Torres, M. A. The Arabidopsis NADPH

oxidases RbohD and RbohF display differential expression patterns and contributions

during plant immunity. *J. Exp. Bot.* **67**, 1663–1676 (2016).

15.   Shang-Guan, K. *et al.* Lipopolysaccharides trigger two successive bursts of reactive

oxygen species at distinct cellular locations. *Plant Physiol.* **176**, 2543–2556 (2018).

16.   Arnaud, D., Deeks, M. J. & Smirnoff, N. Organelle-targeted biosensors reveal distinct

oxidative events during pattern-triggered immune responses. *Plant Physiol.* **191**,

2551–2569 (2022).

17.   Ashtamker, C., Kiss, V., Sagi, M., Davydov, O. & Fluhr, R. Diverse subcellular locations

of cryptogein-induced reactive oxygen species production in tobacco bright yellow-2

cells. *Plant Physiol.* **143**, 1817–1826 (2007).

18.   Stringlis, I. A. *et al.* Root transcriptional dynamics induced by beneficial rhizobacteria

and microbial immune elicitors reveal signatures of adaptation to mutualists. *Plant J.*

93, 166–180 (2018).

19.   Bjornson, M., Pimprikar, P., Nürnberger, T. & Zipfel, C. The transcriptional landscape

of Arabidopsis thaliana pattern-triggered immunity. *Nat. Plants* **7**, 579–586 (2021).

20.   Müller, T. M. *et al.* Transcription activator-like effector nuclease-mediated generation

and metabolic analysis of camalexin-deficient cyp71a12 cyp71a13 double knockout

lines. *Plant Physiol.* **168**, 849–858 (2015).

21.   Ferrari, S., Plotnikova, J. M., De Lorenzo, G. & Ausubel, F. M. Arabidopsis local

resistance to Botrytis cinerea involves salicylic acid and camalexin and requires EDS4

and PAD2, but not SID2, EDS5 or PAD4. *Plant J.* **35**, 193–205 (2003).

22.   Denoux, C. *et al.* Activation of defense response pathways by OGs and Flg22 elicitors

in Arabidopsis seedlings. *Mol. Plant* **1**, 423–445 (2008).

23.   Glazebrook, J. & Ausubel, F. M. Isolation of phytoalexin-deficient mutants of

Arabidopsis thaliana and characterization of their interactions with bacterial

pathogens. *Proc. Natl. Acad. Sci. U. S. A.* **91**, 8955–8959 (1994).

24.   Kutschera, A. *et al.* Bacterial medium-chain 3-hydroxy fatty acid metabolites trigger

immunity in Arabidopsis plants. *Science (80-.).* **364**, 178–181 (2019).

25.   Schellenberger, R. *et al.* Bacterial rhamnolipids and their 3-hydroxyalkanoate

precursors activate Arabidopsis innate immunity through two independent

mechanisms. *Proc. Natl. Acad. Sci. U. S. A.* **118**, 1–10 (2021).

26.   Liang, X. & Zhou, J. M. Receptor-Like Cytoplasmic Kinases: Central Players in Plant

482   Receptor Kinase-Mediated Signaling. *Annu. Rev. Plant Biol.* **69**, 267–299 (2018).

483   27.   Albert, I. *et al.* An RLP23–SOBIR1–BAK1 complex mediates NLP-triggered immunity.

484   *Nat. Plants* **1**, 15140 (2015).

485   28.   Fan, L. *et al.* Genotyping-by-sequencing-based identification of Arabidopsis pattern

486   recognition receptor RLP32 recognizing proteobacterial translation initiation factor

487   IF1. *Nat. Commun.* **13**, (2022).

488   29.   Wan, W. L. *et al.* Comparing Arabidopsis receptor kinase and receptor protein-

489   mediated immune signaling reveals BIK1-dependent differences. *New Phytol.* **221**,

490   2080–2095 (2019).

491   30.   Henry, G., Deleu, M., Jourdan, E., Thonart, P. & Ongena, M. The bacterial lipopeptide

492   surfactin targets the lipid fraction of the plant plasma membrane to trigger immune-

493   related defence responses. *Cell. Microbiol.* **13**, 1824–1837 (2011).

494   31.   Balleza, D., Alessandrini, A. & Beltrán García, M. J. Role of Lipid Composition,

495   Physicochemical Interactions, and Membrane Mechanics in the Molecular Actions of

496   Microbial Cyclic Lipopeptides. *J. Membr. Biol.* **252**, 131–157 (2019).

497   32.   Rondelli, V. *et al.* Sitosterol and glucosylceramide cooperative transversal and lateral

498   uneven distribution in plant membranes. *Sci. Rep.* **11**, 1–11 (2021).

499   33.   Gronnier, J., Gerbeau-Pissot, P., Germain, V., Mongrand, S. & Simon-Plas, F. Divide

500   and Rule: Plant Plasma Membrane Organization. *Trends Plant Sci.* **23**, 899–917 (2018).

501   34.   Haslam, T. M. & Feussner, I. Diversity in sphingolipid metabolism across land plants. *J.*

502   *Exp. Bot.* **73**, 2785–2798 (2022).

503   35.   Ternes, P. *et al.* Disruption of the ceramide synthase LOH1 causes spontaneous cell

504   death in Arabidopsis thaliana. *New Phytol.* **192**, 841–854 (2011).

36.   Lenarčič, T. *et al.* Eudicot plant-specific sphingolipids determine host selectivity of

microbial NLP cytolysins. *Science (80-.).* **358**, 1431–1434 (2017).

37.   Gerbeau-Pissot, P. *et al.* Modification of plasma membrane organization in tobacco

cells elicited by cryptogein. *Plant Physiol.* **164**, 273–286 (2014).

38.   Sandor, R. *et al.* Plasma membrane order and fluidity are diversely triggered by

elicitors of plant defence. *J. Exp. Bot.* **67**, 5173–5185 (2016).

39.   McKenna, J. F. *et al.* The cell wall regulates dynamics and size of plasma-membrane

nanodomains in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 12857–12862 (2019).

40.   Vaahtera, L., Schulz, J. & Hamann, T. Cell wall integrity maintenance during plant

development and interaction with the environment. *Nat. Plants* **5**, 924–932 (2019).

41.   Basu, D. & Haswell, E. S. The Mechanosensitive Ion Channel MSL10 Potentiates

Responses to Cell Swelling in Arabidopsis Seedlings. *Curr. Biol.* **30**, 2716-2728.e6

(2020).

42.   Martinac, B., Adler, J. & Kung, C. Mechanosensitive ion channels of E. coli activated by

amphipaths. *Nature* **348**, 261–263 (1990).

43.   Hamilton, E. S., Schlegel, A. M. & Haswell, E. S. United in Diversity: Mechanosensitive

Ion Channels in Plants. *Annu. Rev. Plant Biol.* **66**, 113–137 (2014).

44.   Hamant, O. & Haswell, E. S. Life behind the wall: Sensing mechanical cues in plants.

BMC Biol. **15**, 1–9 (2017).

45.   Demidchik, V., Shabala, S., Isayenkov, S., Cuin, T. A. & Pottosin, I. Calcium transport

across plant membranes: mechanisms and functions. *New Phytol.* **220**, 49–69 (2018).

46.   Yoshimura, K., Iida, K. & Iida, H. MCAs in Arabidopsis are Ca^{2+}-permeable

mechanosensitive channels inherently sensitive to membrane tension. *Nat. Commun.*

12, 6074 (2021).

47.   Thor, K. *et al.* The calcium-permeable channel OSCA1.3 regulates plant stomatal

immunity. *Nature* **585**, 569–573 (2020).

48.   Haswell, E. S., Peyronnet, R., Barbier-Brygoo, H., Meyerowitz, E. M. & Frachisse, J. M.

Two MscS Homologs Provide Mechanosensitive Channel Activities in the Arabidopsis

Root. *Curr. Biol.* **18**, 730–734 (2008).

49.   Yamanaka, T. *et al.* MCA1 and MCA2 That Mediate Ca2+ Uptake Have Distinct and

Overlapping Roles in Arabidopsis. *Plant Physiol.* **152**, 1284–1296 (2010).

50.   Mori, K. *et al.* Ca2+-permeable mechanosensitive channels MCA1 and MCA2 mediate

cold-induced cytosolic Ca2+ increase and cold tolerance in Arabidopsis. *Sci. Rep.* **8**,

550 (2018).

51.   Moe-Lange, J. *et al.* Interdependence of a mechanosensitive anion channel and

glutamate receptors in distal wound signaling. *Sci. Adv.* **7**, (2021).

52.   Schellenberger, R. *et al.* Bacterial rhamnolipids and their 3-hydroxyalkanoate

precursors activate Arabidopsis innate immunity through two independent

mechanisms. *Proc. Natl. Acad. Sci. U. S. A.* **118**, 2101366118 (2021).

53.   Schellenberger, R. *et al.* Apoplastic invasion patterns triggering plant immunity:

plasma membrane sensing at the frontline. *Mol. Plant Pathol.* **20**, 1602–1616 (2019).

54.   Schikora, A., Schenk, S. T. & Hartmann, A. Beneficial effects of bacteria-plant

communication based on quorum sensing molecules of the N-acyl homoserine lactone

group. *Plant Mol. Biol.* **90**, 605–612 (2016).

55.   Davis, B. M., Jensen, R., Williams, P. & O'Shea, P. The Interaction of N-Acylhomoserine

Lactone Quorum Sensing Signaling Molecules with Biological Membranes:

Implications for Inter-Kingdom Signaling. *PLoS One* **5**, e13522 (2010).

56.   Girard, L., Höfte, M. & Mot, R. De. Lipopeptide families at the interface between

pathogenic and beneficial Pseudomonas-plant interactions. *Critical Reviews in*

Microbiology **46**, 397–419 (2020).

57.   Ngou, B. P. M., Ding, P. & Jones, J. D. G. Thirty years of resistance: Zig-zag through the

plant immune system. *Plant Cell* **34**, 1447–1478 (2022).

58.   Jin, M. S. *et al.* Crystal Structure of the TLR1-TLR2 Heterodimer Induced by Binding of

a Tri-Acylated Lipopeptide. *Cell* **130**, 1071–1082 (2007).

59.   Debois, D. *et al.* Plant polysaccharides initiate underground crosstalk with bacilli by

inducing synthesis of the immunogenic lipopeptide surfactin. *Environ. Microbiol. Rep.*

7, 570–582 (2015).

60.   He, Z., Webster, S. & He, S. Y. Growth–defense trade-offs in plants. *Current Biology* **32**,

R634–R639 (2022).

Figure 1

A — Cyclized peptide ELLVDLL / Acyl chain

B — D.R. 57,4% / *** / Spreading lesions/plant / Mock / Srf

C — Fluorescence increase (RFU) / Mock / Srf / Time (min) / Fluorescence fold increase / *** / *** / Mock / Srf / Mock / Srf

D — $[Ca^{2+}]_{cyt}$ (L/Lmax) / Srf / Mock / flg22 / Time (sec)

E — Δ pH / Mock / Srf / Time (min)

F — Total DEGs / Srf specific / Srf + / Flg22 - / Srf - / Flg22 + / Srf + / Chi - / Srf - / Chi + / DEGs / Time (hours)

96h PI / * / Camalexin (ng/g LDW) / Mock / Srf / * / Mock / Srf / Spreading lesions/plant / Mock / Srf / **** / ns / Col-0 / *pad3*

G — Calcium binding / ROS / MAPK / PR / Receptor Receptor-like kinases / Cell wall / Srf / C F / Hours (post treatment)

Figure 2

Srf-PLPC
E_{int} = -20 kcal/mol

Srf-sitosterol
E_{int} = -2 kcal/mol

Srf-GIPC
E_{int} = -52 kcal/mol

Srf-GluCer
E_{int} = -37 kcal/mol

Figure 3