1

Highly regenerative species-specific genes improve age-associated

features in the adult *Drosophila* midgut

Hiroki Nagai[1], Yuya Adachi[1], Tenki Nakasugi[1], Ema Takigawa[1], Junichiro Ui[1],

Takashi Makino[2], Masayuki Miura[1], and Yu-ichiro Nakajima[1] *

1. Graduate School of Pharmaceutical Sciences, The University of Tokyo

2. Graduate School of Life Science, Tohoku University

*Author for correspondence e-mail: nakaji97@g.ecc.u-tokyo.ac.jp

Tel: +81-3-5841-4863

Text: 35580 characters

Figures: 7

Supporting information: Supplemental figures S1-S4, Supplemental tables S1-S2,

Supplemental texts S1

Abstract

Background

The remarkable regenerative abilities observed in planarians and cnidarians are closely linked to the active proliferation of adult stem cells and the precise differentiation of their progeny, both of which typically deteriorate during aging in low regenerative animals. While regeneration-specific genes conserved in highly regenerative organisms may confer regenerative abilities and long-term maintenance of tissue homeostasis, it remains unclear whether introducing these regenerative genes into low regenerative animals can improve their regeneration and aging processes.

Results

Here we ectopically express high regenerative species-specific JmjC domain-encoding genes (HRJDs) in *Drosophila*, a widely used low regenerative model organism. Surprisingly, HRJD expression impedes tissue regeneration in the developing wing disc but extends organismal lifespan when expressed in the intestinal stem cell lineages of the adult midgut under non-regenerative conditions. Notably, HRJDs enhance the proliferative activity of intestinal stem cells while maintaining their differentiation fidelity, ameliorating age-related decline in gut barrier functions.

Conclusions

These findings together suggest that the introduction of highly regenerative species-specific genes can improve stem cell functions and promote a healthy lifespan when expressed in aging animals.

Keywords

Synthetic biology, aging, healthy lifespan, regeneration, *Drosophila*, intestinal stem cells

INTRODUCTION

Regeneration, an intricate process that rebuilds lost body parts, is a widespread

phenomenon among metazoans, but the capacity for regeneration displays significant

variation across different groups and species [1–5]. While certain animals like planarians

and hydras possess the remarkable ability to regenerate their entire body from a small

fragment, other groups with more complex body structures, such as mammals and insects,

exhibit a diminished regenerative potential and can only regenerate specific tissues and/or

organs to a limited extent. Furthermore, regenerative capacity often declines with aging

in most species with limited regeneration abilities [2], resulting in increased susceptibility

to organismal death upon injury. In contrast, animals that can achieve whole body

regeneration, along with developmental reversion observed in the jellyfish *Turritopsis*,

exhibit potential immortality [2, 5, 6]. Understanding the mechanisms underlying high

regenerative ability and their relationship with aging represents a fundamental challenge

in the field of developmental biology and gerontology with implications for regenerative

medicine.

Several cellular and molecular factors have been identified as determinants of

regeneration capacity. Highly regenerative animals such as planarians and cnidarian

polyps rely on pluripotent adult stem cells, called neoblasts and interstitial cells (i-cells),

respectively [2–5, 7]. These stem cells migrate to the injury sites and contribute to the

formation of a blastema, an undifferentiated cellular mass, enabling the restoration of

amputated body structures. Some vertebrates like salamanders and fish, which do not

possess adult pluripotent stem cells, can regenerate organs after injury by recruiting

blastema cells through dedifferentiation and/or the activation of quiescent lineage-

restricted stem cells [1, 2, 4, 5, 8]. At the molecular level, the evolutionary conserved

WNT signaling pathway promotes a wide range of regenerative events across species,

including blastema formation in newts and *Hydra* [1–5, 8].

In contrast to the conserved regulators of regeneration, several genes are

specific to highly regenerative animal groups and species: for instance, the newt gene

Prod1 regulates re-patterning during limb regeneration [9, 10], and viropana family

(*viropana 1-5*) is up-regulated during lens regeneration [11, 12]. These species/group-

specific genes might explain differences in regeneration capacity between species.

Remarkably, ectopic expression of *viropana 1-5* can enhance regeneration of the

primordium of *Drosophila* eyes that maintain regenerative capacity during development

[12]. This finding raises the possibility that heterologous induction of regenerative genes

may accelerate tissue regeneration, at least in developing animals, and potentially provide

a cue for developing novel regenerative therapies. However, it remains unknown whether

heterologously-induced regenerative genes can improve regenerative and/or aging

processes even when induced in post-developmental mature adults.

Notably, given that basal metazoans such as Porifera, Ctenophore, Placozoa,

and Cnidaria all exhibit robust regenerative abilities, it is conceivable that a common

ancestor of all metazoans once possessed a high regenerative potential and independently

lost genes related to high regenerative capacity in multiple phyla. Building upon this

hypothesis, bioinformatics analysis has identified genes that are common among species

with high regenerative abilities and absent in species with limited regenerative capacities

(Fig. 1A) [13]. The highly regenerative species-specific JmjC domain-encoding genes

(HRJDs) are a group of such genes (with typically two or three orthologs per species)

characterized by their JmjC domain (Fig. 1A), yet their molecular functions remain

unknown. Given their potential influence on the regenerative process, HRJDs may

contribute to the high regeneration potential of highly regenerative animals. With this in

mind, a question arises: what would happen if a low regenerative species, which has lost

HRJDs, were to acquire them again? By ectopically expressing HRJDs in low

regenerative animal models, we can investigate their impacts on regeneration as well as

on aging processes, providing insight into the role of HRJDs.

Here we express HRJDs in the fruit fly *Drosophila melanogaster* and evaluate

their impact *in vivo*, especially by focusing on two epithelial tissues: developing wing

discs and post-developmental adult midguts, both of which exhibit regeneration potential

and can replenish damaged epithelial cells. In contrast to the predicted contribution of

HRJDs in regeneration as observed in planarian, ectopic HRJD induction impedes

regenerative responses and decreases organismal survival upon injury in *Drosophila*.

Surprisingly, however, HRJD expression in the stem/progenitor population of the adult

midguts extends organismal lifespan under the non-regenerative condition. Further

investigations reveal that HRJDs enhance the proliferative activity of intestinal stem cells

while keeping their differentiation fidelity in aged guts, ameliorating age-related decline

in gut barrier functions. These findings provide evidence that genes specific to highly-

regenerative animals can improve stem cell function as well as increase healthy lifespan

upon heterologous expression in aging animals.

RESULTS

Planarian HRJDs expression do not affect gross morphology of fly adults

Planarians are one of the most highly regenerative animals; they are capable of

regenerating most body parts upon amputation and can even reconstruct their whole

body from fragments [1–4, 8, 14]. Previous work has identified two HRJD orthologs

HRJDa and *HRJDb* from two planarian species, *Dugesia japonica* and *Schmidtea*

mediterranea, where both HRJDs contain only the JmjC domain and the EF hand motif

(Fig. 1B) [13]. In functional assays using RNAi-mediated knockdown in *D. japonica*,

these two HRJDs affect viability of organisms after amputation [13], suggesting that

planarian HRJDs are associated with regeneration processes. We thus utilized these

functional HRJDs in *D. japonica* as representatives for the following studies and

hereafter simply named them as HRJDa and HRJDb.

To examine potential benefits and/or disadvantages of acquiring HRJDs, we

then ectopically expressed HRJDs in *Drosophila melanogaster*, which has lost HRJD

genes during evolution (Fig. 1A), using the Gal4/UAS system (Fig. S1) [15]. We first

introduced HRJDs in the whole body throughout development with the ubiquitous driver

Act5C-Gal4 (Fig. 1C, *Act5C-Gal4>UAS-HRJDa/b*). The whole body expression of

HRJDs neither caused developmental lethality nor changed the body weight of mature

adults compared with the *Act5C-Gal4>UAS-GFP* control (Fig. 1D and 1E). We further

assessed the gross morphology of adult wings and eyes under the ubiquitous expression

of HRJDs. The wing size was not altered by HRJD expression (Fig. 1F and 1G). Similarly,

Act5C-Gal4>UAS-HRJDa/b flies did not change the size of adult compound eyes (Fig.

1H and 1I). These results indicate that HRJD expression does not disturb gross

morphology of adult flies under homeostatic conditions, likely due to minimal impacts

on developmental processes.

HRJD expression hampers tissue regeneration in the developing wing disc

Given that HRJDs are conserved only among highly regenerative animals, their primary

functions may be related to regeneration processes. Indeed, both planarian HRJDa and

HRJDb function in whole-body regeneration while their relative contribution is likely

context-dependent: HRJDa is indispensable for the regeneration of amputated heads

while HRJDb promotes organismal survival after two consecutive amputations [13]. To

test whether heterologously-induced HRJDs can enhance regenerative responses in

Drosophila, we examined their impacts on regeneration after ablation of the developing

wing imaginal disc. The *Drosophila* larval imaginal discs, including wing discs, which

are composed of columnar epithelial cells, exhibit regenerative capacity and restore

morphology even after massive cell death [16, 17] (Fig. 2A and 2B). We utilized the

genetic ablation system in which transient overexpression of *eiger*, a *Drosophil*a TNF

ligand, induces apoptosis in the wing pouch region [17, 18]. In this tissue ablation system,

the temperature-sensitive form of the Gal4 repressor Gal80 (*tub-Gal80^ts*) allows transient

tissue ablation under the control of a wing pouch driver *rn-Gal4*. To circumvent the

temporal expression associated with the Gal4/Gal80^ts system, we further introduced an

additional binary expression system: the QF/QUAS system for HRJD induction (Fig. 2A)

[19, 20]. When expressing HRJDs in the wing pouch with the *WP-QF* driver during the

entire process of recovery, we found that *WP-QF>QUAS-HRJDs* flies exhibit severe

defects compared to controls (Fig. 2C). Immunostaining of a mitotic marker, phospho-

histone H3 (PH3), revealed that HRJD expression resulted in a slight but not statistically

significant decrease in regenerative proliferation in the wing pouch, implying the

possibility of insufficient damaged cell replenishment (Fig. 2D and 2E). Importantly,

HRJD induction in the wing disc affected neither the development of the adult wing nor

proliferation in wing discs under homeostatic condition (Fig. 2F-2I). These results

suggest that HRJD expression does not facilitate regeneration in the developing wing disc

epithelium.

HRJD expression compromises regeneration in the adult midgut

Most highly regenerative animals exhibit regeneration potential in mature adult stages

[2–4, 8], and thus one possibility is that HRJDs stimulate regenerative responses only in

adults, where tissue-resident stem cells play an important role in homeostasis and

regeneration. We thus tested the impact of HRJD expression on the regenerative capacity

of the *Drosophila* adult midgut. In the adult midgut, intestinal stem cells (ISCs) self-

renew ISCs themselves and also generate differentiating progenitor cells called

enteroblasts (EBs) and enteroendocrine progenitors (EEPs), which eventually

differentiate into absorptive enterocytes (ECs) and secretory enteroendocrine cells (EEs),

respectively (Fig. 3A) [21–24]. The adult midgut activates proliferation of ISCs in

response to orally-treated harmful chemicals such as paraquat and dextran sulfate sodium

(DSS), a regenerative response that is essential for organismal survival [21, 25–28]. We

thus expressed HRJDs in ISCs and EBs using the *esg-Gal4* (*esg-Gal4>UAS-HRJDa/b*)

driver and tested survival against paraquat/DSS damage. Similar to the wing regeneration

assay, continuous expression of HRJDs via *esg-Gal4* significantly impaired organismal

survival upon the chemical challenges, with a stronger effect for HRJDa (Fig. S2A and

S2B). Survival under control feeding (5% sucrose) was also slightly decreased by HRJD

expression, albeit not to a statistically significant degree (Fig. S2C). We further examined

ISC proliferation after paraquat feeding by counting the number of cells marked with a

mitotic marker, anti-PH3 staining, and found that induction of HRJDa significantly

suppressed ISC division (Fig. S2D), which is consistent with the stronger decline in

survival rate. Concordant with the wing ablation experiments, these results suggest that

continuous HRJD expression causes detrimental effects to regenerative capacity.

Because the *esg-Gal4* driver is active not only in adult ISCs/EBs but also in the

proliferative adult midgut progenitors in the larval midguts and larval imaginal discs [29,

30], continuous HRJD expression may cause internal organ defects associated with

development. To temporally restrict HRJD expression to the adult stage, we next set up

an experiment where HRJD expression starts only in adults after eclosion from pupa by

combining *esg-Gal4* with *tub-Gal80^{ts}* (*esg-Gal4^{ts}>UAS-HRJDa/b*) (Fig. 3B); however,

neither HRJDa nor HRJDb improved organismal survival upon paraquat/DSS feeding

(Fig. 3C and 3D). Instead, HRJDa expression caused a slight but significant decrease in

survival rate compared to controls (*esg-Gal4^{ts}>UAS-GFP*). HRJDb expression similarly

resulted in decreased survival at late phase (Day 4 and Day 5 in paraquat feeding, Day 5

and Day 6 in DSS feeding) , though not to a statistically significant degree (Fig. 3C and

3D). Survival under control feeding (5% sucrose) was comparable between *esg-*

Gal4^{ts}>UAS-GFP and *esg-Gal4^{ts}>UAS-HRJDa/b* (Fig. 3E), suggesting that the impaired

survival of *esg-Gal4^{ts}>UAS-HRJDs* is specific to regenerative contexts. We also found

that, after paraquat feeding, mitotic cell numbers in HRJD expressing flies decreased

slightly (Fig. 3F and 3G), suggesting the possibility of impaired regeneration. These

results indicate that post-developmental expression of HRJDs does not enhance intestinal

regeneration in adult flies.

Post-developmental HRJD expression in ISC lineages extends organismal lifespan

Highly-regenerative animals are often resistant to organismal aging and have long

lifespan; some species like the jellyfish *Turritopsis* even revert their early developmental

stages under harsh environments and are considered to be potentially immortal [2, 6].

Although heterologous expression of HRJDs does not augment tissue regeneration in

either developing or adult *Drosophila* tissues, we further investigated whether HRJDs

influence organismal lifespan under homeostatic conditions in which no experimental

injuries are applied to fly adults. We observed that post-developmental HRJD expression

in adult ISCs/EBs (*esg-Gal4^{ts}>UAS-HRJDa/b*) significantly extended organismal

lifespan both in females (Fig. 4A) and males (Fig. 4B). In contrast, continuous HRJD

expression in the whole body (*Act5C-Gal4>UAS-HRJDa/b*) shortened males' lifespan

(Fig. 4C and 4D). Moreover, HRJD expression in differentiated ECs (*NP1-Gal4>UAS-*

HRJDa/b) also negatively impacted female lifespan, likely due to developmental

abnormalities since EC-specific HRJD expression in the adult stage did not decrease

survival (Fig. S3A-S3D). These results suggest that spatio-temporally regulated induction

of HRJDs in adult ISCs/EBs can be beneficial for adult flies under homeostatic conditions.

Further investigation revealed the distinct impacts of HRJDa and HRJDb on

organismal lifespan: while continuous expression of HRJDa via *esg-Gal4 (esg-*

Gal4>UAS-HRJDa) shortened lifespans for both males and females, *esg-Gal4>UAS-*

HRJDb prolonged female lifespans but shortened male lifespans (Fig. S3E and S3F). The

ortholog-dependent phenotypic differences were also reported in planarian regeneration

[13]. To address the mechanisms of lifespan extension associated with post-

developmental HRJD expression in ISCs/EBs, we focused on *esg-Gal4^{ts}>UAS-HRJDs*

midguts in the following experiments.

HRJD expression improves barrier function in the aged intestine

Age-related mortality accompanies intestinal barrier dysfunction and disruption of cell-

cell junction in the gut epithelium across species [31–34]. Given that aging phenotypes

are more severe in female flies than male flies [35], we first tested barrier function of the

esg-Gal4^{ts}>UAS-HRJDa/b female midgut using the Smurf assay, in which orally-

administered blue dye spreads throughout the entire body when the intestinal barrier is

disrupted (Smurf+, Fig. 5A) [31, 35]. Consistent with lifespan extension, post-

developmental HRJD expression in ISCs/EBs significantly decreased the ratio of Smurf+

adults at 30-days old, a time point where age-related organismal death starts (Fig. 4A and

5A), suggesting that barrier function is maintained in aged flies that express HRJDs in

ISCs/EBs. We further assessed the localization of septate junction markers (Dlg, Tsp2A,

Mesh, Ssk), which diffuse in cytoplasm in the aged midgut [33, 36–38]. The diffusion of

the Dlg protein in aged midguts was significantly suppressed in *esg-Gal4^ts^>UAS-*

HRJDa/b midguts (Fig. 5B and 5C). Similarly, peripheral localization of Tsp2A, Mesh,

and Ssk proteins was also maintained in *esg-Gal4^ts^>UAS-HRJDa/b* midguts compared

with control (*esg-Gal4^ts^>UAS-GFP*) midguts (Fig. 5D-5I). Given that HRJDs in

ISCs/EBs affect localization of junctional components in ECs, we next tested if HRJDs

non-autonomously affect junctional integrity. To this end, we performed a mosaic

experiment where HRJDs were clonally induced using the esgFLPout system [26]. Clonal

induction of HRJDs improved junctional localization of Ssk both inside and outside the

clones (Fig. S4A and S4B), supporting the conclusion that HRJDs' effect is non-cell

autonomous. We then examined JNK activation in ECs, which is associated with barrier

dysfunction in the aged intestine [27, 36]. HRJD expression in ISCs/EBs significantly

repressed transcription of *puckered* (*puc*), a downstream target of JNK signaling in the

whole midgut (Fig. 5J), and indeed the intensity of *puc-lacZ* reporter decreased in ECs

upon HRJD induction (Fig. 5K and 5L). HRJD expression in ISCs/EBs thus suppresses

JNK signaling in ECs in the aged midgut, which is mediated by either a non-autonomous

function of HRJDs or residual HRJD proteins in newly differentiated ECs. These results

indicate that HRJD expression in adult ISCs/EBs contributes to the prolonged

maintenance of the junctional integrity as well as gut barrier function in the aged intestine.

HRJD expression attenuates mis-differentiation of stem cell lineage in the aged

intestine

In addition to the intestinal barrier dysfunction, aberrant activation of ISC mitosis and

accumulation of mis-differentiated ISC progenies are also common age-related

phenotypes in the *Drosophila* adult midgut [27, 31, 33, 39]. In 30-days old control

midguts, *esg-Gal4^{ts}>UAS-GFP$^+$* cells exhibited hallmarks of age-related mis-

differentiation into ECs such as increased ploidy and enlarged nuclear size (Fig. 6A) [27].

By contrast, we found that HRJD expression in adult ISCs/EBs suppressed these mis-

differentiation phenotypes (Fig. 6A and 6B), which are consistent with the amelioration

of age-related barrier dysfunction in *esg-Gal4^{ts}>UAS-HRJDa/b* midguts. Surprisingly,

however, *esg-Gal4^{ts}>UAS-HRJDa/b* did not suppress age-related increase in ISC

proliferation, but rather enhanced it in 30-day old midguts (Fig. 6C and 6D). The

enhancement of ISC proliferation was specific to the aged intestine since *esg-*

Gal4^{ts}>UAS-HRJDs did not affect PH3$^+$ cell number in 10-day-old young midguts (Fig.

S4E). These results raised the possibility that HRJD expression augments the mitotic

activity of stem cells while maintaining their differentiation fidelity during aging.

To test this hypothesis, we first measured the expression of cell type markers

(*Delta* for ISCs, *nub* for ECs, and *pros* for EEs) by RT-qPCR of whole midguts. Both *nub*

and *pros* are negatively regulated by Esg [40, 41], and accumulation of *esg$^+$* mis-

differentiated cells accompanies the down-regulation of EC-related genes in aged midguts

[27]. Consistent with enhanced ISC proliferation, *esg-Gal4^ts^>UAS-HRJDa* significantly

up-regulated *Delta* (*Dl*) expression in 30-day old midguts (Fig. 6E). More importantly,

esg-Gal4^ts^>UAS-HRJDa also up-regulated *nub* and *pros*, which supports our hypothesis

that HRJDs improve proper differentiation of ISC progenies. In *esg-Gal4^ts^>UAS-HRJDb*,

only *Dl*, but not *nub* and *pros*, was up-regulated when whole midguts were used as sample

(Fig. 6E). Next, we focused on the posterior midgut where both HRJDa and HRJDb

prevent mis-differentiation of EBs (Fig. 6A), and examined cell type composition using

the combination of markers *esg^ts^>GFP* (ISC/EB), *Dl-lacZ* (ISC/EEP), and anti-Pros

(EEP/EE). Consistent with the prevention of EB mis-differentiation, *esg^ts^*-mediated

HRJD expression decreased EBs ($esg^+Dl^-Pros^-$) and increased polyploid ECs

($esg^-Dl^-Pros^-$) compared to the control midgut (Fig. 6F and 6G). These results indicate

that HRJDs improve differentiation fidelity in ISC lineage.

To address the potential mechanism of HRJD-dependent maintenance of ISC

functions, we focused on the JAK-STAT pathway since its activation promotes both ISC

proliferation and differentiation into ECs [26, 42, 43]. In young mature midguts, STAT

activity is largely restricted to esg^+ ISCs/EBs [26, 42, 44]. In aged midguts, however, a

subset of polyploid cells exhibited weak signal of the STAT reporter *10×STAT-GFP*,

likely due to the accumulation of mis-differentiated EBs (Fig. 6H). Notably, HRJD

induction canceled such ectopic STAT activation in polyploid cells and rather enhanced

STAT activity in diploid ISCs/EBs (Fig. 6H and 6I), which is consistent with our

observations that HRJDs boost ISC functions during aging. These results suggest that

HRJD expression accelerates the generation of differentiated ECs via JAK-STAT activation. In the healthy homeostatic intestine, ISC division and subsequent differentiation into ECs is coupled with the loss of old ECs [45, 46]. Interestingly, we found that *esg*-lineage clones that express HRJDs rapidly replaced pre-existing cells by generating polyploid ECs (Fig. S4A, S4C, and S4D). In addition, more cells exhibited cleaved Dcp1, an apoptotic marker, in 30-day old *esg^ts* >*HRJDs* midguts (Fig. S4F and S4G), implying that HRJD induction promotes turnover of midgut epithelial cells. Collectively, our data show that post-developmental expression of HRJDs improves stem cell functions in the aged intestine.

DISCUSSION

In this study, we demonstrate that post-developmental expression of planarian HRJD genes in the *Drosophila* adult ISCs/EBs can suppress age-related intestinal dysfunctions and extend organismal lifespan, while their continuous expression throughout the entire developmental process hampers regenerative responses and principally shortens lifespan. Notably, HRJDs in adult ISCs/EBs boost the age-related increase of ISC proliferation but do not cause age-related mis-differentiation of ISC progenies (Fig. 7). These HRJD-mediated outcomes are distinct from those mediated by typical anti-aging manipulations such as antibiotic treatment and metabolic intervention, which ameliorate both ISC over-proliferation and mis-differentiation [24, 31, 47]. Therefore, heterologously induced HRJDs create the neomorphic state in the aged intestine (Fig. 7). Given the age-related increase of chronic cellular stresses [24, 27, 36, 47, 48], we speculate that the HRJD-

dependent ISC activation and their proper differentiation improve organismal fitness by

enabling active turnover of damaged intestinal cells.

Our investigation implies that the activation of the JAK-STAT pathway

underlies the HRJD-mediated enhancement of ISC functions. Surprisingly, however, the

expression of most mitogens including upstream ligands of JAK-STAT signaling (*upd1,*

upd2, upd3) were not up-regulated by HRJD induction (Fig. S4H), raising the possibility

that HRJDs activate STAT by modulating intracellular signaling factors. Such ligand-

independent STAT activation can be achieved via non-receptor type tyrosine kinases like

Src and Abl [49]. Given that the typical JmjC family proteins function as histone

demethylase or protein hydroxylase [50], HRJDs may epigenetically target these tyrosine

kinases. Notably, *Drosophila* Jarid2 (Jumonji, AT rich interactive domain 2) activates

EGFR signaling upon overexpression in ISCs/EBs without changing ligand expression

[51]. Our findings, together with these reports, suggest a cell-autonomous role of JmjC

family proteins in regulating proliferative signaling in stem cell lineages. Intriguingly,

Jarid2 expression in ISCs/EBs leads to ISC over-proliferation as well as barrier

dysfunction, resulting in reduced lifespan [51], which is the opposite adult phenotype of

HRJD expression in ISC lineages. Moreover, in contrast to the nuclear localization of

histone demethylase KDM8 [52, 53], the closest paralog of HRJDs, ectopically-induced

HRJDs localize in the cytoplasm but not in the nucleus both *in vitro* (S2 cells) and *in vivo*

(wing discs and adult midguts) (Fig. S1B-S1D). This is a common localization pattern of

JmjC-domain only proteins [53, 54] and implies that an unknown mechanism may operate

when HRJDs work as a potential histone demethylase. In the future, it will thus be critical

to examine the detailed molecular function of HRJDs to understand their impact on stem

cell lineages.

Although HRJDs have been identified as genes conserved between highly

regenerative animals [13], it is unclear whether HRJDs alone are sufficient to enhance

regenerative responses. Our investigations revealed that ectopic expression of HRJDs

failed to improve regeneration of developing wing discs and adult midguts in *Drosophila*

(Fig. 2 and 3), likely through the attenuation of regenerative growth. However, it should

be noted that we induced HRJDs before tissue injury and maintained their expression

during regeneration period. Given that *HRJDb* is up-regulated in the late phase of

planarian regeneration (3 days after amputation) [13], strict regulation of HRJD induction

might be important for their proper function as true regeneration regulators. On the other

hand, the expression level of *HRJDa* remains constant during planarian regeneration [13],

which is recapitulated in our experiments. Another possibility is that a high regeneration

ability can be achieved by cooperation between HRJDs and other unidentified genes that

have been lost during evolution, which should be addressed in future studies.

During the normal aging process in the *Drosophila* adult midgut, ISC over-

proliferation is closely linked with other aging phenotypes, and suppression of ISC

division can prevent the accumulation of mis-differentiated ISC progenies and extend

organismal lifespan [24, 36, 47, 48]. In contrast, HRJDs enhance age-related ISC

activation but can alleviate mis-differentiation and extend lifespan, suggesting that it is

not ISC activation itself but rather mis-differentiation of ISC progenies that principally

drives age-related mortality. Consistently, manipulation of mitotic spindle orientation,

which affects the cell fate of daughter cells [55–57], can extend organismal lifespan

without changing mitotic activity of ISCs [57]. Of note, in contrast to *Drosophila* ISCs

which over-proliferate in aged midguts, many adult stem cells decrease their activity and

abundance during aging both in *Drosophila* and mammals [24, 58, 59]. Future

investigations for mechanisms of HRJD-mediated stem cell rejuvenation will provide

clues to develop new anti-aging strategies.

CONCLUSIONS

In this study, we established a *Drosophila* model in which HRJDs are heterologously

expressed in specific tissue or cell type using binary expression systems, and

demonstrated that HRJDs can improve proliferative/differentiation capacity of ISCs in

the aged midgut. Although continuous HRJD expression in the wing disc and in the adult

midgut impairs tissue regeneration upon injury, restricted HRJD expression in post-

developmental adult ISCs/EBs enhances mitotic activity of ISCs as well as maintains

their differentiation fidelity in the aged flies, leading to the prevention of age-related

intestinal barrier dysfunction and the extension of organismal lifespan. Our HRJD-

expressing model will serve as a valuable resource to understand unprecedented

mechanisms of stem cell rejuvenation in the future.

METHODS

***Drosophila* stocks**

All stocks were maintained on a standard diet containing 4% cornmeal, 6% baker's yeast
(Saf Yeast), 6% glucose (Wako, 049-31177), and 0.8% agar (Kishida chemical, 260-
01705) with 0.3% propionic acid (Tokyo Chemical Industry, P0500) and 0.05% nipagin
(Wako, 132-02635). Canton S was utilized as the wildtype strain. Transgenic fly lines
were obtained from the Bloomington Drosophila Stock Center (BDSC) and the Kyoto
Stock Center: *Act5C-Gal4* (BDSC 3954), *rn-Gal4, tub-Gal80ts, UAS-egr* (BDSC 51280),
WP-QF2 [60], *tub-Gal80ts* (BDSC 7017, 7019), *esg-Gal4* (Kyoto 109126), *NP1-Gal4*
(Kyoto 112001), *UAS-GFP* (BDSC 1522), *M{3xP3-RFP.attP}ZH-86Fb* (Kyoto 130437),
UAS-HRJDa (this study), *UAS-HRJDb* (this study), *QUAS-HRJDa* (this study), *QUAS-*
HRJDb (this study), *UAS-FLAG-HRJDa* (this study), *UAS-FLAG-HRJDb* (this study),
puc-lacZ (BDSC 98329), *10×STAT-GFP* (BDSC 26198), *Dl-lacZ* (BDSC 11651). *esg-*
Gal4, UAS-GFP, tub-Gal80ts; UAS-FLP, Act5C-FRT.CD2-Gal4 (esgFLPout) is a gift
from Irene Miguel-Aliaga. See **Table S1** for genotypes in each Figure. We used female
flies unless otherwise noted in the Figures.
409
***Drosophila* genetics**
Experimental crosses that did not involve Gal80ts-mediated inhibition of Gal4 were
performed at 25°C. When using Gal80ts, experimental crosses were maintained at 18°C.
For genetic wing ablation experiments (Fig. 2), F1 larvae were raised at 18ºC until day 7,
incubated at 29ºC for the next 40 hr, and then maintained at 18ºC until the adult hatched
or dissection. For lifespan assays, midgut staining, and Smurf assays (Fig. 3-6); F1 adults
were transferred to 29°C three days after eclosion until the experiments. Midgut staining
and Smurf assays were performed after 7 or 27 days of 29°C incubation (young: Day 10
adults, old: Day 30 adults). For esgFLPout experiments, flies were maintained at 18°C
until 50 days, and 50-day old adults were transferred to 29°C and analyzed after 10 days
(final 60-day old).
421
Generation of HRJD expressing lines
To express HRJDs using the Gal4/UAS or the QF/QUAS system, we constructed a vector
containing *HRJDa* or *HRJDb* under the control of either the UAS sequence or QUAS
sequence. Namely we amplified HRJDa and HRJDb sequence from codon-optimized
synthesized DNAs (pUCIDT-HRJDa and pUCIDT-HRJDb, IDT, Supplementary texts
S1) for *DjHRJDa* (Genbank # LC408963) and *DjHRJDb* (Genbank # LC408964),

respectively. UAS-HRJDa and UAS-HRJDb were constructed by ligating HRJDa and
HRJDb into pUAST-attB (DGRC 1419), respectively (digested by EcoRI/NotI).
Similarly, QUAS-HRJDa and QUAS-HRJDb were constructed by ligating HRJDa and
HRJDb into pQUAS-WALIUM20 (DGRC 1474), respectively (digested by NheI/EcoRI).
To construct UAS-3xFLAG-HRJDs, we PCR-amplified the coding sequences of HRJDs
from pUCIDT-HRJDs by gene-specific primers with 3xFLAG tag at 5'-end of the
forward primers, and the amplicons were then ligated into a pUAST-attB vector digested
by EcoRI/NotI using In-Fusion HD Cloning Kit (Takara, 639649). The landing site for
each construct was {3xP3-RFP.attP}ZH-86Fb. Injection and selection were performed
by WellGenetics (Taiwan, R.O.C.). Please see also Table S2 and Supplementary texts S1
for primer sequences and HRJDs sequences.
439

440   **S2 cell culture**
441   *Drosophila* S2 cells were grown at 25°C in Schneider's *Drosophila* medium (GIBCO,
442   21720001) supplemented with 10% (v/v) heat-inactivated fetal bovine serum (FBS), 100
443   U/mL penicillin, and 100 μg/mL streptomycin (FUJIFILM Wako, 168–23191).
444

S2 cell immunostaining
To examine the expression pattern of HRJDs through immunostaining, 1×10^6 cells were
seeded in 6-well plates containing Schneider's *Drosophila* medium supplemented with
10% (v/v) heat-inactivated fetal bovine serum, 100 U/mL penicillin, and 100 μg/mL
streptomycin and were transfected with 800 ng of pAc5-3xFLAG-HRJDs using Effectene
Transfection Reagent (QIAGEN, 301427) following the manufacturer's protocol. To
prepare pAc5-3xFLAG-HRJDs, we amplified the coding sequences of HRJDs from
pUCIDT-HRJDs (Supplementary texts S1) by gene-specific primers with 3xFLAG tag at
5'-end of the forward primers, and the amplicons were then ligated into a pAc5-
STABLE2-neo (Addgene, 32426) [61] digested with EcoRI/XhoI by In-Fusion HD
Cloning Kit (Takara, 639649). Twenty-four hours after transfection, the cells were
washed with PBS and the medium was replaced with fresh Schneider's *Drosophila*
medium supplemented with 10% (v/v) heat-inactivated FBS, 100 U/mL penicillin, and
100 μg/mL streptomycin. Forty-eight hours after transfection, the cells were fixed with
4% paraformaldehyde (PFA) in PBS, washed with PBS containing 0.1% Triton X-100
(PBST), blocked in PBST with 5% normal donkey serum (PBSTn), and incubated with

primary antibodies in PBSTn overnight at 4 °C. The samples were then washed with
PBST, incubated for 1 h at room temperature with secondary antibodies and Hoechst
33342 suspended in PBSTn and washed again with PBST. Images were captured using
an LSM880 (Zeiss). The primary antibody used was a mouse anti-FLAG M2 monoclonal
antibody (1 : 2000, Sigma, F1804). The secondary antibodies used were Goat anti-Mouse
IgG2b Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 555 (1:2000, Thermo Fisher
Scientific, A-21147). Hoechst 33342 (0.4 µM; Invitrogen, H3570) was used for nuclear
staining. Images were analyzed and edited using Fiji/ImageJ software (NIH).
469
S2 cell western blotting
To examine the expression of HRJDs, 2.5×10^5 cells were seeded in 24-well plates
containing Schneider's *Drosophila* medium supplemented with 10% (v/v) heat-
inactivated FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin and were transfected
with 200 ng of pAc5-3xFLAG-HRJDs using Effectene Transfection Reagent (QIAGEN,
301427) following the manufacturer's protocol. Twenty-four hours after transfection, the
cells were washed with PBS and the medium was replaced with fresh Schneider's
Drosophila medium supplemented with 10% (v/v) heat-inactivated FBS, 100 U/mL
penicillin, and 100 µg/mL streptomycin. Forty-eight hours after transfection, the cells
were washed with PBS and lysed with 50 µL RIPA buffer supplemented with cOmplete
ULTRA EDTA-free protease inhibitor cocktail (Roche, 05892953001). The lysate was
sonicated and centrifuged at 20,000 g for 5 minutes. Protein concentration was
determined by BCA assay. The supernatant was mixed with SDS, boiled at 95 °C for 5
minutes, and subjected to western blotting.
Proteins were separated using SDS-PAGE and transferred onto Immobilon-P
PVDF membranes (Millipore, IPVH00010) for immunoblotting. Membranes were
blocked with 4% skimmed milk diluted in 1×Tris buffered saline containing 0.1% Tween-
20. Immunoblotting was performed using the below-mentioned antibodies, which were
diluted with 4% skim milk. The signals were visualized using Immobilon Western
Chemiluminescent HRP Substrate (Millipore, WBKLS0500) and FUSION SOLO. 7S.
EDGE (Vilber-Lourmat). Contrast and brightness adjustments were applied using the
Fiji/ImageJ software (NIH).
The primary antibody used was mouse anti-FLAG M2 monoclonal antibody
(1 : 5000, Sigma, F1804). Mouse anti-alpha tubulin (DM1A) monoclonal antibody

(1 : 5000, Sigma, T9026) was used as loading control. The secondary antibody used was

HRP-conjugated goat/rabbit/donkey anti-mouse IgG (1 : 10000, Promega, W402B).

Drug treatment

5 mM paraquat (Sigma, 856177) and 5% (w/v) DSS (MP Biomedicals, 160110) were

dissolved in 5% (w/v) sucrose solution. Filter paper (Whatman 3MM) was soaked with

400 µl of these reagents and placed into empty vials. For histological analyses, flies were

fed with the reagent solution for 1 day. For the survival assay, flies were transferred to

new vials and dead flies (determined by immobility and showing no response to tapping)

were counted every day. 5% sucrose was used for control feeding.

Immunofluorescence

Samples were dissected in 1×PBS and fixed in 4% PFA for 20 minutes (wing discs) and

30-45 minutes (adult midgut) at room temperature (RT), respectively. The following

primary antibodies were used with indicated dilution into 1×PBS containing 0.5% BSA

and 0.1% Triton X-100: rabbit anti-PH3 (Millipore 06-570, 1:1000), mouse anti-FLAG

(Sigma F1804, 1:1000), mouse anti-Dlg (DSHB 4F3, 1:100), rabbit anti-GFP (MBL 598,

1:500), rat anti-GFP (Nacalai tesque 04404-26, 1:400), rabbit anti-Tsp2A (Izumi et al.,

2016, 1:1000) [62], rabbit anti-Mesh (Izumi et al., 2012, 1:1000) [63], rabbit anti-Ssk

(Izumi et al., 2016, 1:1000) [62], chicken anti-β-galactosidase (Abcam ab9361, 1:500),

anti-cDcp1 (Cell Signaling Technology 9578, 1:200). After overnight incubation with

primary antibodies at 4°C, samples were incubated with fluorescent secondary antibodies

(Jackson ImmunoResearch and Invitrogen, 1:500) for 1 hour at RT. Hoechst 33342

(Invitrogen, final concentration: 10 µg/ml) was used to visualize DNA. Wing discs were

mounted as described previously [64, 65]. Samples were mounted in Slowfade Diamond

(ThermoFisher, S36963) and imaged with confocal microscopy Zeiss LSM880 or Zeiss

LSM980.

Smurf assay

We referred to Rera et al. (2012) for the Smurf assay conditions [35]. To prepare the

feeding medium, 100 µl of 50% (w/v) brilliant blue FCF (Wako, 027-12842, final 2.5%)

and 100 µl of 5% (w/v) sucrose were added to a vial containing 2 ml of cornmeal-agar

food. After mixing with a spatula, an Whatman 1 filter (1001-020) was put on the feeding

medium. Flies were fed with this medium at 25°C for one day, after which they were
transferred to a new vial containing cornmeal-agar food without blue dye to clean the
epidermis. Two hours after the transfer, Smurf phenotype was checked. We classified the
Smurf phenotype as blue dye leakage outside abdomen (thorax, head, legs).

RT-qPCR

Total RNA was purified from 10-15 midguts using the ReliaPrep RNA Tissue Miniprep
System (Promega). cDNA was made from 100 or 200 ng of RNA using PrimeScript RT
Reagent Kit (TaKaRa). Quantitative PCR was performed using TB Green Premix Ex
Taq II (TaKaRa) and the QuantStudio 6 Flex Real-Time PCR System (ThermoFisher).
RpL32 was used as an internal control. Primer sequences are listed in Table S2.

Phylogenetic tree

We created the phylogenetic tree of representative organisms using the NCBI taxonomy
browser   (https://www.ncbi.nlm.nih.gov/Taxonomy/CommonTree/wwwcmt.cgi).   The
species examined are *Amphimedon queenslandica* (Porifera), *Mnemiopsis leidyi*
(Ctenophora), *Trichoplax adhaerens* (Placozoa), *Hydra vulgaris* (Cnidaria), *Ptychodera
flava* (Hemichordata), *Acanthaster planci* (Echinodermata), *Branchiostoma floridae*
(Cephalochordata), *Ciona intestinalis* (Tunicata), *Danio rerio* (Teleostei), *Homo sapiens*,
Mus musculus (Mammalia), *Schmidtea mediterranea* (Platyhelminthes), *Caenorhabditis
elegans* (Nematoda), and *Drosophila melanogaster* (Arthropoda). These species, except
for *D. rerio, H. sapiens, M. musculus, C. elegans,* and *D. melanogaster,* are known to
have HRJDs and exhibit high regeneration ability [13]. The pictures were downloaded
from PhyloPic, and the color of some were changed from black to blue. Credits: Bennet
McComish (*B. floridae*), Malio Kodis (*M. leidyi*), and Markus A. Grohme (*S.
mediterranea*,   https://www.phylopic.org/images/93f9611c-0cbd-4a14-92da-
004a4521e21f/schmidtea-mediterranea).

Quantification of PH3 positive cells

For wing discs, we counted PH3 positive cells in the pouch region, which develops into
the adult wing. After binarization, the number of PH3 positive cells was counted using
the Analyze Particles function in Fiji/ImageJ, and subsequently confirmed or corrected

559 by visual review of the images. For adult midguts, the number of PH3 positive cells in
560 the posterior midguts was manually counted.
561

Quantification of junctional proteins
To quantify junctional and cytoplasmic intensity, a line that was orthogonal to one side
of a cell was drawn, and then the plot profile was calculated for the line. We defined the
junctional intensity as the highest value in that prolife, and defined cytoplasmic intensity
as the value 20 pixels (~2.6 μm) away from the highest value. Three lines were drawn for
one cell and the average of the three was used to represent the junctional/cytoplasmic
ratio for the cell. Six polyploid cells were randomly selected from each image for
quantification.
570

571 **Quantification of reporter intensity**
572 The signal intensity of *puc-lacZ* and *10×STAT-GFP* reporters was quantified using
573 Fiji/ImageJ. The nuclei of large polyploid cells (for *puc-lacZ*) and Pros$^-$ diploid cells (for
574 *10×STAT-GFP*) were selected as ROIs using the polygon selection tool. The reporter
575 intensity in each ROI was quantified using the Measure command.
576

577 **Quantification of nuclear size**
578 Nuclei of *esg-Gal4>UAS-GFP* positive cells were traced by the polygon selection tool
579 and added as ROI in Fiji/ImageJ. The size of each ROI was quantified using the Measure
580 command.
581

582 **Quantification of cell number**
583 Cell number was quantified using Fiji/ImageJ. To count the total cell number, Hoechst
584 signal was processed as following: (1) despeckle, (2) binarization, (3) fill hole, (4)
585 watershed, (5) analyze particle. The following cell number was manually counted using
586 the cell counter function: *esg>GFP*$^+$ cells, *Dl-lacZ*$^+$ cells, Pros$^+$ cells, and cDcp1$^+$ cells.
587

Statistics
Statistical analyses were performed using Excel and RStudio. Two tailed *t* tests were used
for comparisons between two groups. One-way ANOVAs with post hoc Tukey tests were
performed when comparing three or more groups. Log-rank tests were used for

comparison of survival curve. Significance is indicated in the Fig.s as follows: *P≤0.05,

**P≤0.01, ***P≤0.001, Not Significant (N.S.): P>0.05. Bar graphs show mean ± standard

error. Boxplots show median (thick line in the box), first and third quartiles (bottom and

top of the box), minimum value (lower whisker), and maximum value (upper whisker).

Dots in bar graphs and boxplots indicate individual values. Violin plots portray the

distribution of individual values.

DECLARATIONS

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets and *Drosophila* stocks used and/or analyzed during the current study are

available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by JSPS/MEXT KAKENHI (grant numbers JP22J01430 to

H.N., JP21H04774, JP23H04766, JP24H00567 to M.M., and JP17H06332, JP22H02762,

JP23K18134, JP23H04696 to Y.N.), AMED-Aging (JP21gm5010001 to M.M.), AMED-

PRIME (JP22gm6110025 to Y.N.), and Sadako O. Hirai Ban Award for Young

Researchers (H.N.)

Author contributions

Conceptualization: H.N., Y.N.

Investigation: H.N., Y.A., T.N., E.T., J.U., Y.N.

Methodology: H.N., Y.N.

Validation: H.N., Y.A., T.N., E.T., J.U., T.M., M.M., Y.N.

Data curation: H.N., T.M., Y.N.

Writing – original draft: H.N., J.U., T.M., Y.N.

Writing – review & editing: H.N., Y.A., T.N., J.U., T.M., M.M., Y.N.

Supervision: M.M., Y.N.

Funding acquisition: H.N., T.M., M.M., Y.N.

Acknowledgements

We thank I. Miguel-Aliaga, N. Shinoda, M. Furuse, Y. Izumi, BDSC, Kyoto Stock Center,

Transgenic RNAi Project (TRiP), and DSHB for fly stocks and reagents.

FIGURE LEGENDS

Figure 1. HRJD expression in whole body does not affect gross morphology.

(A) Phylogenetic tree of HRJD conservation. Green indicates species that possess HRJD

gene(s) and gray indicates species that have lost HRJD gene(s). High regenerative ability

indicates that the species can regenerate their whole body or anterior/posterior body parts,

and low regenerative ability indicates that the species can only regenerate their appendage

(limbs, tails, fins) or much smaller scale of tissues/organs. We referred to Cao et al., 2019

for the definition of regenerative ability [13].

(B) Protein sequence of HRJDs used in this study, which derived from *Dugesia japonica*

(DjHRJDa/b, hereafter simply described as HRJDa/b). Conserved residues characteristic

of the JmjC domain are shown.

(C) Schematics of genetic experiments for whole body induction of HRJDs. The Gal4-

UAS system enables gene expression downstream of the UAS sequence, which is

regulated by the transcription factor Gal4 [15]. In this case, the ubiquitously active *Act5C*

promoter is used for Gal4 expression.

(D) Representative images of mature adults for whole body expression of HRJDs.

(E) Whole body induction of HRJDs did not change body weight of mature adults.

(F) Representative images of adult wings.

(G) Whole body induction of HRJDs did not change wing size.

(H) Representative images for adult heads.

(I) Whole body induction of HRJDs did not change eye size of adult flies.

N.S., not significant: P>0.05. One-way ANOVAs with post hoc Tukey test. *n* indicates

the number of flies examined. See also Fig. S1.

Figure 2. HRJD expression impedes tissue regeneration in the developing wind disc

(A) Schematics of genetic ablation of the wing disc. TNF ligand *eiger* was overexpressed

in the wing pouch for 40 hrs during the larval stage.

(B) Wing regeneration was assessed in the adult stage by checking the size of wing (0%:

no wing, 100%: intact wing).

(C) Line graph for wing regeneration. Expression of HRJDa/b increased the rate of low

regeneration (0% and 25%) at the expense of the rate of full regeneration (100%).

(D-E) Representative images for PH3 staining of wing discs during regeneration at 48 hrs

after ablation (D). The number of PH3 positive cells is quantified in (E).

(F-G) Representative images for adult wings of *WP-QF>QUAS-HRJDa/b* flies under

homeostatic (non-regenerative) condition (F). The wing size is quantified in (G).

 (H-I) Representative images for PH3 staining of wing discs (3rd instar larvae) during

normal development (H). The number of PH3 positive cells is quantified in (I).

N.S., not significant: P>0.05, ***P ≤0.001. Chi-square test (C) and one-way ANOVAs

with post hoc Tukey test (E, G, I). *n* indicates the number of wings (C, G) and wing discs

(E, I) examined. Scale bars: 20 μm.

Figure 3. HRJDs expression compromises regeneration in the adult midgut

(A) Schematics of the *Drosophila* adult midgut. The adult midgut is a pseudostratified

epithelium, in which intestinal stem cells (ISCs) generate both absorptive enterocytes

(ECs) and secretory enteroendocrine cells (EEs) through progenitor cell enteroblasts

(EBs) and EE progenitor cells (EEPs).

(B) Experimental scheme for ISC/EB specific induction of HRJDs under the *esg-Gal4*

driver in the adult stage. Adult flies were transferred to 29°C at Day 3 to induce HRJD

expression.

(C-E) Survival curve during paraquat (C), DSS (D), and sucrose (E) feeding.

(F) Representative images for PH3 staining upon paraquat feeding.

(G) Quantification of PH3 positive cells in posterior midguts.

N.S., not significant: P>0.05, *P<0.05, **P≤0.01. Log-rank test. *n* indicates the number

of flies (C-E) and midguts (G). Scale bars: 50 μm. See also Fig. S2.

Figure 4. Post-developmental HRJD expression in ISC lineages extends organismal

lifespan

(A-D) Survival curve for organismal lifespan by sex. HRJDs are induced by *esg^{ts}-Gal4*

(A and B) and *Act5C-Gal4* (C and D). For temporal regulation of HRJD induction using

tub-Gal80^{ts} (A and B), adult flies were transferred to 29°C at Day 3.

N.S., not significant: P>0.05, *P<0.05, **P≤0.01, ***P≤0.001, Log-rank tests. *n*

indicates the number of flies. See also Fig. S3.

Figure 5. HRJD expression in adult ISCs/EBs suppress age-related gut barrier

dysfunction

(A) *esg^{ts}*-mediated HRJD induction decreased Smurf(+) ratio at Day 30. The

representative images of the Smurf phenotype (leakage of blue dye throughout whole

body) is shown on the left.

(B-C) HRJD expression in adult ISCs/EBs maintained junctional localization of Dlg

protein in ECs, which is quantified in (C).

(D-I) HRJD expression in adult ISCs/EBs maintained junctional localization of Tsp2A

(D), Mesh (E), and Ssk (F) in ECs, which is quantified in (G-I).

(J) RT-qPCR of *puc* in the whole midgut. *esg^{ts}*-mediated induction of HRJDa significantly

repressed *puc*.

(K-L) *puc-lacZ* intensity in polyploid cells was decreased by HRJD expression (K), which

is quantified in (L).

*P<0.05, **P<0.01, ***P≤0.001, One-way ANOVAs with post hoc Tukey test. *n*

indicates the number of flies (A), cells (C, G, H, I, L), and biological replicates (J). Scale

bars: 50 μm.

Figure 6. HRJD expression in adult ISCs/EBs improves stem cell functions in aged

midguts

(A) Representative images for *esg-Gal4* positive cells in Day 30 midguts. HRJDs

suppressed mis-differentiation of *esg*⁺ cells (cellular enlargement with polyploid large

nuclei).

(B) Quantification of nuclear size of *esg-Gal4* positive cells.

(C-D) Representative images (C) and quantification (D) of PH3 positive cells in Day 30

posterior midguts.

(E) RT-qPCR of *Dl*, *nub*, and *pros* in Day 30 midguts. Expression levels were normalized

to those of GFP control.

(F-G) Cell type composition in aged midguts were assessed with *esg^{ts}>GFP* (ISC, EB,

and early EC), *Dl-lacZ* (ISC and EEP), and anti-Pros staining (EEP and EE). HRJD

expression in adult ISCs/EBs reduced the ratio of EB/early EC and increased the ratio of

EC (G).

(H-I) *esg^{ts}*-mediated HRJD induction enhanced STAT-GFP intensity in Pros⁻ diploid cells

(ISCs/EBs), which is quantified in (I). Additionally, HRJDs suppressed STAT activation

in polyploid ECs (arrowheads).

N.S., not significant: P>0.05, *P≤0.05, **P≤0.01, ***P≤0.001, one-way ANOVAs with

post hoc Tukey test. *n* indicates the number of cells (B, I), guts (D, G), and biological

replicates (E). Scale bars: 50 μm.

Figure 7. Graphical summary.

Schematic model for the impact of HRJDs on intestinal homeostasis. In aged wildtype

midguts, ISCs over-proliferate, and their daughters fail to differentiate into mature ECs,

resulting in an accumulation of mis-differentiated cells (esg^+ large polyploid). HRJD

expression in adult ISCs/EBs further enhances ISC proliferation but suppresses mis-

differentiation likely through upregulation of JAK-STAT signaling, which results in

successful maintenance of the gut barrier and an extension of organismal lifespan.

Supplemental figure legends

Figure S1. Expression and localization of HRJD proteins

(A) Western blots of S2 cell samples upon transfection of 3×FLAG-HRJDa/b. Anti-FLAG

was used to detect 3×FLAG-HRJDa/b proteins and anti-tubulin was used as the loading

control. Expected molecular weight: 52.0 kDa for 3×FLAG-HRJDa, 52.9 kDa for

3×FLAG-HRJDb.

(B) Representative images of anti-FLAG staining using S2 cells transfected with

3×FLAG-HRJDa/b. Both HRJDa and HRJDb localized cytoplasm but not in nuclei. Scale

bar: 30 μm in wide-view images (left), 5 μm in magnified view images (yellow frames).

(C-D) Representative images of 3×FLAG-HRJDa/b localization in wing discs and adult

midguts. FLAG-tagged HRJDs were induced using *WP-Gal4* (C) or *esg-Gal4* (D). Scale

bars: 10 μm.

Figure S2. Continuous expression of HRJDs throughout development hampers intestinal regeneration

(A-C) Organismal survival upon paraquat (A), DSS (B), and sucrose (C) feeding. Drug feeding started at Day 7 after eclosion. HRJD induction using *esg-Gal4* throughout development significantly decreased survival upon paraquat/DSS feeding.

(D) Expression of HRJDa using *esg-Gal4* decreased the number of PH3 positive cells in the posterior midguts, while HRJDb did not significantly change PH3 positive cell number.

N.S., not significant: $P>0.05$, **$P\leq0.01$, ***$P\leq0.001$, one-way ANOVAs with post hoc Tukey test. *n* indicates the number of flies (A-C) and guts (D).

Figure S3. Organismal lifespan of flies expressing HRJDs in ECs and ISC/EBs

(A-F) Survival curve for organismal lifespan by sex. HRJDs are induced by *NP1-Gal4* (A and B), *NP1ts-Gal4* (C and D), and *esg-Gal4* (E and F).

N.S., not significant: $P>0.05$, ***$P\leq0.001$, Log-rank tests. *n* indicates the number of flies.

Figure S4. HRJD induction in adult ISCs/EBs promotes turnover of midgut epithelial cells.

(A) Representative images of esgFLPout aged midguts in which *esg*-lineage clones express HRJDs together with GFP. Clones were induced 50 day after eclosion and traced for 10 days (final 60-day old adults). Ssk was immunostained to compare the septate

801 **Table S1. Detailed genotypes in each experiment**

Fig.	Panel	Genotype
Fig. 1	**D-I**	*w;; Act5C-Gal4 / +*
		w; UAS-GFP / +; Act5C-Gal4 / UAS-HRJDb

		w;; Act5C-Gal4 / UAS-HRJDa
		w;; Act5C-Gal4 / UAS-HRJDb
Fig. 2	**A-E**	*w; rn-Gal4, tub-Gal80ts, UAS-egr / +; WP-QF2 / +*
		w; rn-Gal4, tub-Gal80ts, UAS-egr / +; WP-QF2 / QUAS-HRJDa
		w; rn-Gal4, tub-Gal80ts, UAS-egr / +; WP-QF2 / QUAS-HRJDb
	F-I	*w;; WP-QF2 / +*
		w;; WP-QF2 / QUAS-HRJDa
		w;; WP-QF2 / QUAS-HRJDb
Fig. 3	**C-G**	*w; esg-Gal4, tub-Gal80ts / UAS-GFP*
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDa / +
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDb / +
Fig. 4	**A-B**	*w; esg-Gal4, tub-Gal80ts / UAS-GFP*
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDa / +
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDb / +
	C-D	*w; UAS-GFP / +; Act5C-Gal4 / +*
		w;; Act5C-Gal4 / UAS-HRJDa
		w;; Act5C-Gal4 / UAS-HRJDb
Fig. 5	**A-I**	*w; esg-Gal4, tub-Gal80ts / UAS-GFP*
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDa / +
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDb / +
	J	*w, Su(H)GBE-lacZ / w; esg-Gal4, tub-Gal80ts / UAS-GFP*
		w, Su(H)GBE-lacZ / w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDa / +
		w, Su(H)GBE-lacZ / w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDb / +
	K-L	*w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; puc-lacZ / +*
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; puc-lacZ / UAS-HRJDa
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; puc-lacZ / UAS-HRJDb
Fig. 6	**A-E**	*w; esg-Gal4, tub-Gal80ts, UAS-GFP / +*
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; UAS-HRJDa / +
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; UAS-HRJDb / +
	F-G	*w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; Dl-lacZ / +*
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; Dl-lacZ / UAS-HRJDa
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; Dl-lacZ / UAS-HRJDb

	H-I	*w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; 10×STAT-GFP / +*
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; 10×STAT-GFP / UAS-HRJDa
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; 10×STAT-GFP / UAS-HRJDb
Fig. S1	**C**	*w;; WP-Gal4 / +*
		w;; WP-Gal4 / UAS-FLAG-HRJDa
		w;; WP-Gal4 / UAS-FLAG-HRJDb
	D	*w; esg-Gal4, tub-Gal80ts, UAS-GFP / +*
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; UAS-FLAG-HRJDa / +
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; UAS-FLAG-HRJDb / +
Fig. S2	**All**	*w; esg-Gal4 / UAS-GFP*
		w; esg-Gal4 / +; UAS-HRJDa / +
		w; esg-Gal4 / +; UAS-HRJDb / +
Fig. S3	**A-B**	*w; NP1-Gal4 / UAS-GFP*
		w; NP1-Gal4 / +; UAS-HRJDa / +
		w; NP1-Gal4 / +; UAS-HRJDb / +
	C-D	*w; NP1-Gal4, tub-Gal80ts / UAS-GFP*
		w; NP1-Gal4, tub-Gal80ts / +; UAS-HRJDa / +
		w; NP1-Gal4, tub-Gal80ts / +; UAS-HRJDb / +
	E-F	*w; esg-Gal4 / UAS-GFP*
		w; esg-Gal4 / +; UAS-HRJDa / +
		w; esg-Gal4 / +; UAS-HRJDb / +
Fig. S4	**A-D**	*w; esg-Gal4, UAS-GFP, tub-Gal80ts / +; UAS-FLP, Act5C-FRT-CD2-FRT-Gal4 / +*
		w; esg-Gal4, UAS-GFP, tub-Gal80ts / +; UAS-FLP, Act5C-FRT-CD2-FRT-Gal4 / UAS-HRJDa
		w; esg-Gal4, UAS-GFP, tub-Gal80ts / +; UAS-FLP, Act5C-FRT-CD2-FRT-Gal4 / UAS-HRJDb
	E-G	*w; esg-Gal4, tub-Gal80ts, UAS-GFP / +*
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; UAS-HRJDa / +
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; UAS-HRJDb / +
	H	*w, Su(H)GBE-lacZ / w; esg-Gal4, tub-Gal80ts / UAS-GFP*
		w, Su(H)GBE-lacZ / w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDa / +

		w, Su(H)GBE-lacZ / w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDb / +

802 **Table S2. Primer sequences**

804 **Supplementary texts S1. Sequences for pUCIDT-HRJDa and pUCIDT-HRJDb**

Abbreviations

i-cell: interstitial cell

HRJD: highly regenerative species-specific JmjC domain-encoding gene

PH3: phospho-histone H3

ISC: intestinal stem cell

EB: enteroblast

EEP: enteroendocrine progenitor

EC: enterocyte

EE: enteroendocrine cell

DSS: dextran sulfate sodium

puc: puckered

Dl: Delta

Jarid2: Jumonji, AT rich interactive domain 2

References

1. Bely AE, Nyberg KG. Evolution of animal regeneration: re-emergence of a field.
Trends Ecol Evol. 2010;25:161–70.

2. Poss KD. Advances in understanding tissue regenerative capacity and mechanisms in
animals. Nat Rev Genet. 2010;11:710–22.

3. Tiozzo S, Copley RR. Reconsidering regeneration in metazoans: an evo-devo approach.
Front Ecol Evol. 2015;3.

4. Bideau L, Kerner P, Hui J, Vervoort M, Gazave E. Animal regeneration in the era of
transcriptomics. Cell Mol Life Sci. 2021;78:3941–56.

5. Fujita S, Kuranaga E, Nakajima Y. Regeneration Potential of Jellyfish: Cellular
Mechanisms and Molecular Insights. Genes. 2021;12:758.

6. Piraino S, Boero F, Aeschbach B, Schmid V. Reversing the Life Cycle: Medusae

Transforming into Polyps and Cell Transdifferentiation in *Turritopsis nutricula* (Cnidaria,

Hydrozoa). Biol Bull. 1996;190:302–12.

7. Varley Á, Horkan HR, McMahon ET, Krasovec G, Frank U. Pluripotent, germ cell

competent adult stem cells underlie cnidarian regenerative ability and clonal growth. Curr

Biol. 2023;33:1883-1892.e3.

8. Gazave E, Röttinger E. 7th Euro Evo Devo meeting: Report on the "Evolution of

regeneration in Metazoa" symposium. J Exp Zoolog B Mol Dev Evol. 2021;336:89–93.

9. Kumar A, Godwin JW, Gates PB, Garza-Garcia AA, Brockes JP. Molecular Basis for

the Nerve Dependence of Limb Regeneration in an Adult Vertebrate. Science.

2007;318:772–7.

10. Da Silva SM, Gates PB, Brockes JP. The Newt Ortholog of CD59 Is Implicated in

Proximodistal Identity during Amphibian Limb Regeneration. Dev Cell. 2002;3:547–55.

11. Looso M, Preussner J, Sousounis K, Bruckskotten M, Michel CS, Lignelli E, et al. A

de novo assembly of the newt transcriptome combined with proteomic validation

identifies new protein families expressed during tissue regeneration. Genome Biol.

2013;14:R16.

12. Mehta AS, Deshpande P, Chimata AV, Tsonis PA, Singh A. Newt regeneration genes

regulate Wingless signaling to restore patterning in Drosophila eye. iScience.

2021;24:103166.

13. Cao P-L, Kumagai N, Inoue T, Agata K, Makino T. JmjC Domain-Encoding Genes

Are Conserved in Highly Regenerative Metazoans and Are Associated with Planarian

Whole-Body Regeneration. Genome Biol Evol. 2019;11:552–64.

14. Umesono Y, Tasaki J, Nishimura Y, Hrouda M, Kawaguchi E, Yazawa S, et al. The

molecular logic for planarian regeneration along the anterior–posterior axis. Nature.

2013;500:73–6.

15. Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates

and generating dominant phenotypes. Development. 1993;118:401–15.

16. Hariharan IK, Serras F. Imaginal disc regeneration takes flight. Curr Opin Cell Biol.

2017;48:10–6.

17. Smith-Bolton RK, Worley MI, Kanda H, Hariharan IK. Regenerative Growth in

Drosophila Imaginal Discs Is Regulated by Wingless and Myc. Dev Cell. 2009;16:797–

809.

18. Worley MI, Alexander LA, Hariharan IK. CtBP impedes JNK- and Upd/STAT-driven

cell fate misspecifications in regenerating Drosophila imaginal discs. eLife.
2018;7:e30391.

19. Potter CJ, Tasic B, Russler EV, Liang L, Luo L. The Q System: A Repressible Binary
System for Transgene Expression, Lineage Tracing, and Mosaic Analysis. Cell.
2010;141:536–48.

20. Riabinina O, Luginbuhl D, Marr E, Liu S, Wu MN, Luo L, et al. Improved and
expanded Q-system reagents for genetic manipulations. Nat Methods. 2015;12:219–22.

21. Nagai H, Miura M, Nakajima Y. Cellular mechanisms underlying adult tissue
plasticity in Drosophila. Fly (Austin). 2022;16:190–206.

22. Miguel-Aliaga I, Jasper H, Lemaitre B. Anatomy and Physiology of the Digestive
Tract of *Drosophila melanogaster*. Genetics. 2018;210:357–96.

23. Zhang P, Edgar BA. Insect Gut Regeneration. Cold Spring Harb Perspect Biol.
2021;:a040915.

24. Jasper H. Intestinal Stem Cell Aging: Origins and Interventions. Annu Rev Physiol.
2020;82:203–26.

25. Buchon N, Broderick NA, Chakrabarti S, Lemaitre B. Invasive and indigenous
microbiota impact intestinal stem cell activity through multiple pathways in *Drosophila*.
Genes Dev. 2009;23:2333–44.

26. Jiang H, Patel PH, Kohlmaier A, Grenley MO, McEwen DG, Edgar BA.
Cytokine/Jak/Stat Signaling Mediates Regeneration and Homeostasis in the Drosophila
Midgut. Cell. 2009;137:1343–55.

27. Biteau B, Hochmuth CE, Jasper H. JNK Activity in Somatic Stem Cells Causes Loss
of Tissue Homeostasis in the Aging Drosophila Gut. Cell Stem Cell. 2008;3:442–55.

28. Amcheslavsky A, Jiang J, Ip YT. Tissue Damage-Induced Intestinal Stem Cell
Division in Drosophila. Cell Stem Cell. 2009;4:49–61.

29. Jiang H, Edgar BA. EGFR signaling regulates the proliferation of *Drosophila* adult
midgut progenitors. Development. 2009;136:483–93.

30. Houtz P, Bonfini A, Bing X, Buchon N. Recruitment of Adult Precursor Cells
Underlies Limited Repair of the Infected Larval Midgut in Drosophila. Cell Host Microbe.
2019;26:412-425.e5.

31. Rera M, Bahadorani S, Cho J, Koehler CL, Ulgherait M, Hur JH, et al. Modulation of
Longevity and Tissue Homeostasis by the Drosophila PGC-1 Homolog. Cell Metab.
2011;14:623–34.

32. Thevaranjan N, Puchta A, Schulz C, Naidoo A, Szamosi JC, Verschoor CP, et al.

Age-Associated Microbial Dysbiosis Promotes Intestinal Permeability, Systemic

Inflammation, and Macrophage Dysfunction. Cell Host Microbe. 2017;21:455-466.e4.

33. Nagai H, Tatara H, Tanaka-Furuhashi K, Kurata S, Yano T. Homeostatic Regulation

of ROS-Triggered Hippo-Yki Pathway via Autophagic Clearance of Ref(2)P/p62 in the

Drosophila Intestine. Dev Cell. 2021;56:81-94.e10.

34. Egge N, Arneaud SLB, Wales P, Mihelakis M, McClendon J, Fonseca RS, et al. Age-

Onset Phosphorylation of a Minor Actin Variant Promotes Intestinal Barrier Dysfunction.

Dev Cell. 2019;51:587-601.e7.

35. Rera M, Clark RI, Walker DW. Intestinal barrier dysfunction links metabolic and

inflammatory markers of aging to death in Drosophila. Proc Natl Acad Sci.

2012;109:21528–33.

36. Resnik-Docampo M, Koehler CL, Clark RI, Schinaman JM, Sauer V, Wong DM, et

al. Tricellular junctions regulate intestinal stem cell behaviour to maintain homeostasis.

Nat Cell Biol. 2017;19:52–9.

37. Salazar AM, Resnik-Docampo M, Ulgherait M, Clark RI, Shirasu-Hiza M, Jones DL,

et al. Intestinal Snakeskin Limits Microbial Dysbiosis during Aging and Promotes

Longevity. iScience. 2018;9:229–43.

38. Izumi Y, Furuse K, Furuse M. Septate junctions regulate gut homeostasis through

regulation of stem cell proliferation and enterocyte behavior in *Drosophila*. J Cell Sci.

2019;:jcs.232108.

39. Chen H, Zheng X, Zheng Y. Age-Associated Loss of Lamin-B Leads to Systemic

Inflammation and Gut Hyperplasia. Cell. 2014;159:829–43.

40. Korzelius J, Naumann SK, Loza-Coll MA, Chan JS, Dutta D, Oberheim J, et al.

Escargot maintains stemness and suppresses differentiation in *Drosophila* intestinal stem

cells. EMBO J. 2014;33:2967–82.

41. Li Y, Pang Z, Huang H, Wang C, Cai T, Xi R. Transcription Factor Antagonism

Controls Enteroendocrine Cell Specification from Intestinal Stem Cells. Sci Rep.

2017;7:988.

42. Zhou F, Rasmussen A, Lee S, Agaisse H. The UPD3 cytokine couples environmental

challenge and intestinal stem cell division through modulation of JAK/STAT signaling

in the stem cell microenvironment. Dev Biol. 2013;373:383–93.

43. Zhai Z, Boquete J-P, Lemaitre B. A genetic framework controlling the differentiation

of intestinal stem cells during regeneration in Drosophila. PLOS Genet.
2017;13:e1006854.

44. Hudry B, De Goeij E, Mineo A, Gaspar P, Hadjieconomou D, Studd C, et al. Sex
Differences in Intestinal Carbohydrate Metabolism Promote Food Intake and Sperm
Maturation. Cell. 2019;178:901-918.e16.

45. Liang J, Balachandra S, Ngo S, O'Brien LE. Feedback regulation of steady-state
epithelial turnover and organ size. Nature. 2017;548:588–91.

46. Antonello ZA, Reiff T, Ballesta-Illan E, Dominguez M. Robust intestinal homeostasis
relies on cellular plasticity in enteroblasts mediated by miR-8–Escargot switch. EMBO J.
2015;34:2025–41.

47. Clark RI, Salazar A, Yamada R, Fitz-Gibbon S, Morselli M, Alcaraz J, et al. Distinct
Shifts in Microbiota Composition during Drosophila Aging Impair Intestinal Function
and Drive Mortality. Cell Rep. 2015;12:1656–67.

48. Wang L, Ryoo HD, Qi Y, Jasper H. PERK Limits Drosophila Lifespan by Promoting
Intestinal Stem Cell Proliferation in Response to ER Stress. PLOS Genet.
2015;11:e1005220.

49. Hu X, Li J, Fu M, Zhao X, Wang W. The JAK/STAT signaling pathway: from bench
to clinic. Signal Transduct Target Ther. 2021;6:402.

50. Markolovic S, Leissing TM, Chowdhury R, Wilkins SE, Lu X, Schofield CJ.
Structure–function relationships of human JmjC oxygenases—demethylases versus
hydroxylases. Curr Opin Struct Biol. 2016;41:62–72.

51. Suong DNA, Shimaji K, Pyo J-H, Park J-S, Yoshida H, Yoo M-A, et al.
Overexpression of dJmj differentially affects intestinal stem cells and differentiated
enterocytes. Cell Signal. 2018;42:194–210.

52. Hsia DA, Tepper CG, Pochampalli MR, Hsia EYC, Izumiya C, Huerta SB, et al.
KDM8, a H3K36me2 histone demethylase that acts in the cyclin A1 coding region to
regulate cancer cell proliferation. Proc Natl Acad Sci. 2010;107:9671–6.

53. Shalaby NA, Sayed R, Zhang Q, Scoggin S, Eliazer S, Rothenfluh A, et al. Systematic
discovery of genetic modulation by Jumonji histone demethylases in Drosophila. Sci Rep.
2017;7:5240.

54. Klose RJ, Kallin EM, Zhang Y. JmjC-domain-containing proteins and histone
demethylation. Nat Rev Genet. 2006;7:715–27.

55. Nakajima Y. Mitotic spindle orientation in epithelial homeostasis and plasticity. J

Biochem (Tokyo). 2018;164:277–84.

56. Goulas S, Conder R, Knoblich JA. The Par Complex and Integrins Direct Asymmetric
Cell Division in Adult Intestinal Stem Cells. Cell Stem Cell. 2012;11:529–40.

57. Hu DJ-K, Jasper H. Control of Intestinal Cell Fate by Dynamic Mitotic Spindle
Repositioning Influences Epithelial Homeostasis and Longevity. Cell Rep.
2019;28:2807-2823.e5.

58. Brunet A, Goodell MA, Rando TA. Ageing and rejuvenation of tissue stem cells and
their niches. Nat Rev Mol Cell Biol. 2023;24:45–62.

59. Campisi J, Kapahi P, Lithgow GJ, Melov S, Newman JC, Verdin E. From discoveries
in ageing research to therapeutics for healthy ageing. Nature. 2019;571:183–92.

60. Kashio S, Obata F, Zhang L, Katsuyama T, Chihara T, Miura M. Tissue
nonautonomous effects of fat body methionine metabolism on imaginal disc repair in
Drosophila. Proc Natl Acad Sci. 2016;113:1835–40.

61. González M, Martín-Ruíz I, Jiménez S, Pirone L, Barrio R, Sutherland JD. Generation
of stable Drosophila cell lines using multicistronic vectors. Sci Rep. 2011;1:75.

62. Izumi Y, Motoishi M, Furuse K, Furuse M. A tetraspanin regulates septate junction
formation in *Drosophila* midgut. J Cell Sci. 2016;:jcs.180448.

63. Izumi Y, Yanagihashi Y, Furuse M. A novel protein complex, mesh-ssk, is required
for septate junction formation in *drosophila* midgut. J Cell Sci. 2012;:jcs.112243.

64. Nakajima Y. Analysis of Epithelial Architecture and Planar Spindle Orientation in
the Drosophila Wing Disc. In: Turksen K, editor. Stem Cell Renewal and Cell-Cell
Communication. Methods Mol Biol. 2021;2346:51–62.

65. Nakajima Y, Lee ZT, McKinney SA, Swanson SK, Florens L, Gibson MC. Junctional
tumor suppressors interact with 14-3-3 proteins to control planar spindle alignment. J Cell
Biol. 2019;218:1824–38.

A

HRJD (+) ▮   HRJD (−) ▮   Regeneration ability

Ctenophora *Mnemiopsis leidyi*	High
Porifera *Amphimedon queenslandica*	High
Placozoa *Trichoplax adhaerens*	High
Cnidaria *Hydra vulgaris*	High
Hemichordata *Ptychodera flava*	High
Echinodermata *Acanthaster planci*	High
Cephalochordata *Branchiostoma floridae*	High
Tunicata *Ciona intestinalis*	High
Teleostei *Danio rerio*	Low
Mammalia *Homo sapiens, Mus musculus*	Low
Platyhelminthes *Schmidtea mediterranea*	High
Nematoda *Caenorhabditis elegans*	Low
Arthropoda *Drosophila melanogaster*	Low

B

DjHRJDa
JmjC domain · EF hand
(aa) 115 · 164 · 66 · 36 · 36

DjHRJDb
JmjC domain · EF hand
(aa) 141 · 174 · 63 · 36 · 16

S ··· H ··· D ··· K ··· H
conserved residues

C

Whole body overexpression of HRJDs

♀ *Act5C-Gal4* × ♂ *UAS-HRJDa/b*
→ *Act5C-Gal4>UAS-HRJDa/b*

Act5C – Gal4 → GAL4 → UAS – HRJDa or b

D

Act-Gal4>

+   GFP   HRJDa   HRJDb

E

N.S.

Body weight (mg)

+ (20)   GFP (13)   HRJDa (34)   HRJDb (18)

F

+   GFP   HRJDa   HRJDb

G

N.S.

Relative wing size

+ (22)   GFP (13)   HRJDa (28)   HRJDb (17)

H

+   GFP   HRJDa   HRJDb

I

N.S.

Relative eye size

+ (20)   GFP (13)   HRJDa (30)   HRJDb (18)

A

Lumen
Epithelia
Visceral muscle

ISC (esg⁺) → EB (esg⁺) → EC (NP1⁺)
ISC (esg⁺) → EEp → EE (Pros⁺)

B ISC/EB specific overexpression of HRJDs in adult stage

esg-Gal4, tub-Gal80ᵗˢ>UAS-HRHDa/b

29°C
PH3
Paraquat or DSS
18°C
Eclosion Day 3 D10 D11
HRJD overexpression

C Paraquat feeding, *esgᵗˢ>*

□ GFP (n=144)
◇ HRJDa (n=158)
▲ HRJDb (n=153)
* N.S.

D DSS feeding, *esgᵗˢ>*

□ GFP (n=86)
◇ HRJDa (n=105)
▲ HRJDb (n=95)
** N.S.

E Control sucrose feeding, *esgᵗˢ>*

□ GFP (n=115)
◇ HRJDa (n=118)
▲ HRJDb (n=107)
N.S.

F Paraquat feeding, *esgᵗˢ>*

PH3 / Hoechst
PH3
GFP HRJDa HRJDb

G Paraquat feeding, *esgᵗˢ>*

PH3 positive cells in posterior
N.S.
(21) (20) (16)
GFP HRJDa HRJDb

A Female lifespan, *esg^ts-Gal4>*

B Male lifespan, *esg^ts-Gal4>*

C Female lifespan, *Act-Gal4>*

D Male lifespan, *Act-Gal4>*