1

Highly regenerative species-specific genes improve age-associated

features in the adult *Drosophila* midgut

Hiroki Nagai[1], Junichiro Ui[1], Tenki Nakasugi[1], Takashi Makino[2], Masayuki

Miura[1], and Yu-ichiro Nakajima[1] *

1. Graduate School of Pharmaceutical Sciences, The University of Tokyo

2. Graduate School of Life Science, Tohoku University

*Author for correspondence e-mail: nakaji97@g.ecc.u-tokyo.ac.jp

Tel: +81-3-5841-4863

Text: 29632 characters

Figures: 6

Supporting information: supplemental figures S1-S3, supplemental tables S1-S2,

supplemental texts S1

Abstract

The remarkable regenerative abilities observed in planarians and cnidarians are closely linked to the active proliferation of adult stem cells and the precise differentiation of their progeny, both of which typically deteriorate during aging in low regenerative animals. While regeneration-specific genes conserved in highly regenerative organisms may confer regenerative abilities and long-term maintenance of tissue homeostasis, it remains unclear whether introducing these regenerative genes into low regenerative animals can improve their regeneration and aging processes. Here we ectopically express high regenerative species-specific JmjC domain-encoding genes (HRJDs) in *Drosophila*, a widely used low regenerative model organism. Surprisingly, HRJD expression impedes tissue regeneration in the developing wing disc but extends organismal lifespan when expressed in the intestinal stem cell lineages of the adult midgut under non-regenerative conditions. Notably, HRJDs enhance the proliferative activity of intestinal stem cells while maintaining their differentiation fidelity, ameliorating age-related decline in gut barrier functions. These findings together suggest that the introduction of highly regenerative species-specific genes can improve stem cell functions and promote a healthy lifespan when expressed in aging animals.

Keywords

Synthetic biology, aging, healthy lifespan, regeneration, *Drosophila*, intestinal stem cells

INTRODUCTION

Regeneration, an intricate process that rebuilds lost body parts, is a widespread phenomenon among metazoans, but the capacity for regeneration displays significant variation across different groups and species [1–5]. While certain animals like planarians and hydras possess the remarkable ability to regenerate their entire body from a small fragment, other groups with more complex body structures, such as mammals and insects, exhibit a diminished regenerative potential and can only regenerate specific tissues and/or organs to a limited extent. Furthermore, regenerative capacity often declines with aging in most species with limited regeneration abilities [2], resulting in increased susceptibility to organismal death upon injury. In contrast, animals that can achieve whole body regeneration, along with developmental reversion observed in the jellyfish *Turritopsis*, exhibit potential immortality [2, 5, 6]. Understanding the mechanisms underlying high regenerative ability and their relationship with aging represents a fundamental challenge in the field of developmental biology and gerontology with implications for regenerative medicine.

Several cellular and molecular factors have been identified as determinants of regeneration capacity. Highly regenerative animals such as planarians and cnidarian polyps rely on pluripotent adult stem cells, called neoblasts and interstitial cells (i-cells), respectively [2–5, 7]. These stem cells migrate to the injury sites and contribute to the formation of a blastema, an undifferentiated cellular mass, enabling the restoration of amputated body structures. Some vertebrates like salamanders and fish, which do not possess adult pluripotent stem cells, can regenerate organs after injury by recruiting

blastema cells through dedifferentiation and/or the activation of quiescent lineage-

restricted stem cells [1, 2, 4, 5, 8]. At the molecular level, the evolutionary conserved

WNT signaling pathway promotes a wide range of regenerative events across species,

including blastema formation in newts and *Hydra* [1–5, 8].

In contrast to the conserved regulators of regeneration, several genes are

specific to highly regenerative animal groups and species: for instance, the newt gene

Prod1 regulates re-patterning during limb regeneration [9, 10], and viropana family

(*viropana 1-5*) is up-regulated during lens regeneration [11, 12]. These species/group-

specific genes might explain differences in regeneration capacity between species.

Remarkably, ectopic expression of *viropana 1-5* can enhance regeneration of the

primordium of *Drosophila* eyes that maintain regenerative capacity during development

[12]. This finding raises the possibility that heterologous induction of regenerative genes

may accelerate tissue regeneration, at least in developing animals, and potentially provide

a cue for developing novel regenerative therapies. However, it remains unknown whether

heterologously-induced regenerative genes can improve regenerative and/or aging

processes even when induced in post-developmental mature adults.

Notably, given that basal metazoans such as Porifera, Ctenophore, Placozoa,

and Cnidaria all exhibit robust regenerative abilities, it is conceivable that a common

ancestor of all metazoans once possessed a high regenerative potential and independently

lost genes related to high regenerative capacity in multiple phyla. Building upon this

hypothesis, bioinformatics analysis has identified genes that are common among species

with high regenerative abilities and absent in species with limited regenerative capacities

(Fig. 1A) [13]. The highly regenerative species-specific JmjC domain-encoding genes

(HRJDs) are a group of such genes characterized by their JmjC domain (Fig. 1A), yet

their molecular functions remain unknown. Given their potential influence on the

regenerative process, HRJDs may contribute to the high regeneration potential of highly

regenerative animals. With this in mind, a question arises: what would happen if a low

regenerative species, which has lost HRJDs, were to acquire them again? By ectopically

expressing HRJDs in low regenerative animal models, we can investigate their impacts

on regeneration as well as on aging processes, providing insight into the role of HRJDs.

Here we express HRJDs in the fruit fly *Drosophila melanogaster* and evaluate

their impact *in vivo*, especially by focusing on two epithelial tissues: developing wing

discs and post-developmental adult midguts, both of which exhibit regeneration potential

and can replenish damaged epithelial cells. In contrast to the predicted contribution of

HRJDs in regeneration as observed in planarian, ectopic HRJD induction impedes

regenerative responses and decreases organismal survival upon injury in *Drosophila*.

Surprisingly, however, HRJD expression in the stem/progenitor population of the adult

midguts extends organismal lifespan under the non-regenerative condition. Further

investigations reveal that HRJDs enhance the proliferative activity of intestinal stem cells

while keeping their differentiation fidelity in aged guts, ameliorating age-related decline

in gut barrier functions. These findings provide evidence that genes specific to highly-

regenerative animals can improve stem cell function as well as increase healthy lifespan

upon heterologous expression in aging animals.

RESULTS

Planarian HRJDs expression do not affect gross morphology of fly adults

Planarians are one of the most highly regenerative animals; they are capable of

regenerating most body parts upon amputation and can even reconstruct their whole

body from fragments [1–4, 8, 14]. Previous work has identified two HRJD orthologs

HRJDa and *HRJDb* from two planarian species, *Dugesia japonica* and *Schmidtea*

mediterranea (Fig. 1B) [13]. In functional assays using RNAi-mediated knockdown in

D. japonica, these two HRJDs affect viability of organisms after amputation [13],

suggesting that planarian HRJDs are associated with regeneration processes. We thus

utilized these functional HRJDs in *D. japonica* as representatives for the following

studies and hereafter simply named them as HRJDa and HRJDb.

To examine potential benefits and/or disadvantages of acquiring HRJDs, we

then ectopically expressed HRJDs in *Drosophila melanogaster*, which has lost HRJD

genes during evolution (Fig. 1A), using the Gal4/UAS system (Fig. S1) [15]. We first

introduced HRJDs in the whole body throughout development with the ubiquitous driver

Act5C-Gal4 (Fig. 1C, *Act5C-Gal4>UAS-HRJDa/b*). The whole body expression of

HRJDs neither caused developmental lethality nor changed the body weight of mature

adults compared with the *Act5C-Gal4>UAS-GFP* control (Fig. 1D and 1E). We further

assessed the gross morphology of adult wings and eyes under the ubiquitous expression

of HRJDs. The wing size was not altered by HRJD expression (Fig. 1F and 1G). Similarly,

Act5C-Gal4>UAS-HRJDa/b flies did not change the size of adult compound eyes (Fig.

1H and 1I). These results indicate that HRJD expression does not disturb gross

morphology of adult flies under homeostatic conditions, likely due to minimal impacts

on developmental processes.

HRJD expression hampers tissue regeneration in the developing wing disc

Given that HRJDs are conserved only among highly regenerative animals, their primary

functions may be related to regeneration processes. Indeed, both planarian HRJDa and

HRJDb function in whole-body regeneration while their relative contribution is likely

context-dependent: HRJDa is indispensable for the regeneration of amputated heads

while HRJDb promotes organismal survival after two consecutive amputations [13]. To

test whether heterologously-induced HRJDs can enhance regenerative responses in

Drosophila, we examined their impacts on regeneration after ablation of the developing

wing imaginal disc. The *Drosophila* larval imaginal discs, including wing discs, which

are composed of columnar epithelial cells, exhibit regenerative capacity and restore

morphology even after massive cell death [16, 17] (Fig. 2A and 2B). We utilized the

genetic ablation system in which transient overexpression of *eiger*, a *Drosophil*a TNF

ligand, induces apoptosis in the wing pouch region [17, 18]. In this tissue ablation system,

the temperature-sensitive form of the Gal4 repressor Gal80 (*tub-Gal80^{ts}*) allows transient

tissue ablation under the control of a wing pouch driver *rn-Gal4*. To circumvent the

temporal expression associated with the Gal4/Gal80^{ts} system, we further introduced an

additional binary expression system: the QF/QUAS system for HRJD induction (Fig. 2A)

[19, 20]. When expressing HRJDs in the wing pouch with the *WP-QF* driver during the

entire process of recovery, we found that *WP-QF>QUAS-HRJDs* flies exhibit severe

defects compared to controls (Fig. 2C). Importantly, HRJD induction in the wing disc did

not affect the development of adult wing under homeostatic condition (Fig. 2D and 2E).

These results suggest that HRJD expression does not facilitate regeneration in the

developing wing disc epithelium.

HRJD expression compromises regeneration in the adult midgut

Most highly regenerative animals exhibit regeneration potential in mature adult stages

[2–4, 8], and thus one possibility is that HRJDs stimulate regenerative responses only in

adults, where tissue-resident stem cells play an important role in homeostasis and

regeneration. We thus tested the impact of HRJD expression on the regenerative capacity

of the *Drosophila* adult midgut. In the adult midgut, intestinal stem cells (ISCs) self-

renew ISCs themselves and also generate differentiating progenitor cells called

enteroblasts (EBs) and enteroendocrine progenitors (EEPs), which eventually

differentiate into absorptive enterocytes (ECs) and secretory enteroendocrine cells (EEs),

respectively (Fig. 3A) [21–24]. The adult midgut activates proliferation of ISCs in

response to orally-treated harmful chemicals such as paraquat and dextran sulfate sodium

(DSS), a regenerative response that is essential for organismal survival [21, 25–28]. We

thus expressed HRJDs in ISCs and EBs using the *esg-Gal4* (*esg-Gal4>UAS-HRJDa/b*)

driver and tested survival against paraquat/DSS damage. Similar to the wing regeneration

assay, continuous expression of HRJDs via *esg-Gal4* significantly impaired organismal

survival upon the chemical challenges, with a stronger effect for HRJDa (Fig. S2A and

S2B). Survival under control feeding (5% sucrose) was also slightly decreased by HRJD

expression, albeit not to a statistically significant degree (Fig. S2C). We further examined

ISC proliferation after paraquat feeding by counting the number of cells marked with a

mitotic marker, anti-phospho histone H3 (PH3) staining, and found that induction of

HRJDa significantly suppressed ISC division (Fig. S2D), which is consistent with the

stronger decline in survival rate. Concordant with the wing ablation experiments, these

results suggest that continuous HRJD expression causes detrimental effects to

regenerative capacity.

Because the *esg-Gal4* driver is active not only in adult ISCs/EBs but also in the

proliferative adult midgut progenitors in the larval midguts and larval imaginal discs [29,

30], continuous HRJD expression may cause internal organ defects associated with

development. To temporally restrict HRJD expression to the adult stage, we next set up

an experiment where HRJD expression starts only in adults after eclosion from pupa by

combining *esg-Gal4* with *tub-Gal80^{ts}* (*esg-Gal4^{ts}>UAS-HRJDa/b*) (Fig. 3B); however,

neither HRJDa nor HRJDb improved organismal survival upon paraquat/DSS feeding

(Fig. 3C and 3D). Instead, HRJDa expression caused a slight but significant decrease in

survival rate compared with control (*esg-Gal4^{ts}>UAS-GFP*) (Fig. 3C and 3D). Survival

under control feeding (5% sucrose) was comparable between *esg-Gal4^{ts}>UAS-GFP* and

esg-Gal4^{ts}>UAS-HRJDa/b (Fig. 3E), suggesting that the impaired survival of *esg-*

Gal4^{ts}>UAS-HRJDa is specific to regenerative contexts. We also confirmed that there

were no significant changes in mitotic cell numbers in HRJD expressing flies compared

to controls (Fig. 3F and 3G). These results indicate that post-developmental expression

of HRJDs does not enhance intestinal regeneration in adult flies.

Post-developmental HRJD expression in ISC lineages extends organismal lifespan

Highly-regenerative animals are often resistant to organismal aging and have long

lifespan; some species like the jellyfish *Turritopsis* even revert their early developmental

stages under harsh environments and are considered to be potentially immortal [2, 6].

Although heterologous expression of HRJDs does not augment tissue regeneration in

either developing or adult *Drosophila* tissues, we further investigated whether HRJDs

influence organismal lifespan under homeostatic conditions in which no experimental

injuries are applied to fly adults. We observed that post-developmental HRJD expression

in adult ISCs/EBs (*esg-Gal4^{ts}>UAS-HRJDa/b*) significantly extended organismal

lifespan both in females (Fig. 4A) and males (Fig. 4B). In contrast, continuous HRJD

expression in the whole body (*Act5C-Gal4>UAS-HRJDa/b*) shortened males' lifespan

(Fig. 4C and 4D). Moreover, HRJD expression in differentiated ECs (*NP1-Gal4>UAS-*

HRJDa/b) also negatively impacted female lifespan, likely due to developmental

abnormalities since EC-specific HRJD expression in the adult stage did not affect survival

(Fig. S3A-S3D). These results suggest that spatio-temporally regulated induction of

HRJDs in adult ISCs/EBs can be beneficial for adult flies under homeostatic conditions.

Further investigation revealed the distinct impacts of HRJDa and HRJDb on

organismal lifespan: while continuous expression of HRJDa via *esg-Gal4 (esg-*

Gal4>UAS-HRJDa) shortened lifespans for both males and females, *esg-Gal4>UAS-*

HRJDb prolonged female lifespans but shortened male lifespans (Fig. S3E and S3F). The

ortholog-dependent phenotypic differences were also reported in planarian regeneration

[13]. To address the mechanisms of lifespan extension associated with post-

developmental HRJD expression in ISCs/EBs, we focused on *esg-Gal4^ts>UAS-HRJDs*

midguts in the following experiments.

HRJD expression attenuates mis-differentiation of stem cell lineage in the aged

intestine

Age-related mortality accompanies intestinal barrier dysfunction and disruption of cell-

cell junction in the gut epithelium across species [31–34]. Given that aging phenotypes

are more severe in female flies than male flies [35], we first tested barrier function of the

esg-Gal4^ts>UAS-HRJDa/b female midgut using the Smurf assay, in which orally-

administered blue dye spreads throughout the entire body when the intestinal barrier is

disrupted (Smurf+, Fig. 5A) [31, 35]. Consistent with lifespan extension, post-

developmental HRJD expression in ISCs/EBs significantly decreased the ratio of Smurf+

adults at 30-days old, a time point where age-related organismal death starts (Fig. 4A and

5A), suggesting that barrier function is maintained in aged flies that express HRJDs in

ISCs/EBs. We further assessed the localization of septate junction markers (Dlg, Tsp2A,

Mesh, Ssk), which diffuse in cytoplasm in the aged midgut [33, 36–38]. The diffusion of

the Dlg protein in aged midguts was significantly suppressed in *esg-Gal4^ts>UAS-*

HRJDa/b midguts (Fig. 5B and 5C). Similarly, peripheral localization of Tsp2A, Mesh,

and Ssk proteins was also maintained in *esg-Gal4^ts>UAS-HRJDa/b* midguts compared

with control (*esg-Gal4^ts>UAS-GFP*) midguts (Fig. 5D-5I). These results indicate that

HRJD expression in adult ISCs/EBs contributes to the prolonged maintenance of the

junctional integrity as well as gut barrier function in the aged intestine.

In addition to the intestinal barrier dysfunction, aberrant activation of ISC

mitosis and accumulation of mis-differentiated ISC progenies are also common age-

related phenotypes in the *Drosophila* adult midgut [27, 31, 33, 39]. In 30-days old control

midguts, *esg-Gal4^{ts}>UAS-GFP$^+$* cells exhibited hallmarks of age-related mis-

differentiation into enterocytes such as increased ploidy and enlarged nuclear size (Fig.

6A) [27]. By contrast, we found that HRJD expression in adult ISCs/EBs suppressed these

mis-differentiation phenotypes (Fig. 6A and 6B), which are consistent with the

amelioration of age-related barrier dysfunction in *esg-Gal4^{ts}>UAS-HRJDa/b* midguts.

Surprisingly, however, *esg-Gal4^{ts}>UAS-HRJDa/b* did not suppress age-related increase

in ISC proliferation, but rather enhanced it in 30-day old midguts (Fig. 6C and 6D). The

enhancement of ISC proliferation was specific to the aged intestine since *esg-*

Gal4^{ts}>UAS-HRJDs did not affect PH3$^+$ cell number in 10-day-old young midguts (Fig.

6D). These results raised the possibility that HRJD expression augments the mitotic

activity of stem cells while maintaining their differentiation fidelity during aging. To test

this hypothesis, we measured the expression of cell type markers (*Delta* for ISCs, *nub* for

enterocytes, and *pros* for enteroendocrine cells) by RT-qPCR of whole midguts. Both *nub*

and *pros* are negatively regulated by Esg [40, 41], and accumulation of *esg$^+$* mis-

differentiated cells accompanies the down-regulation of EC-related genes in aged midguts

[27]. Consistent with enhanced ISC proliferation, *esg-Gal4^{ts}>UAS-HRJDa* significantly

up-regulated *Delta* expression in 30-day old midguts (Fig. 6E). More importantly, *esg-*

Gal4ts>UAS-HRJDa also up-regulated *nub* and *pros*, which supports our hypothesis that

HRJDs improve proper differentiation of ISC progenies. On the other hand, *esg-*

Gal4ts>UAS-HRJDb mildly increased *Delta* expression, although *nub* and *pros* were not

significantly up-regulated (Fig. 6E). Taken together, our results show that post-

developmental expression of HRJDs improves stem cell functions in aged intestine, while

the underlying mechanisms between HRJDa and HRJDb likely differ.

DISCUSSION

In this study, we demonstrate that post-developmental expression of planarian HRJD

genes in the *Drosophila* adult ISCs/EBs can suppress age-related intestinal dysfunctions

and extend organismal lifespan, while their continuous expression throughout the entire

developmental process hampers regenerative responses and principally shortens lifespan.

Notably, HRJDs in adult ISCs/EBs boost the age-related increase of ISC proliferation but

do not cause age-related mis-differentiation of ISC progenies (Fig. 6F). These HRJD-

mediated outcomes are distinct from those mediated by typical anti-aging manipulations

such as antibiotic treatment and metabolic intervention, which ameliorate both ISC over-

proliferation and mis-differentiation [24, 31, 42]. Therefore, heterologously induced

HRJDs create the neomorphic state in the aged intestine (Fig. 6F). Given the age-related

increase of chronic cellular stresses [24, 27, 36, 42, 43], we speculate that the HRJD-

dependent ISC activation and their proper differentiation improve organismal fitness by

enabling active turnover of damaged intestinal cells.

The typical JmjC family proteins function as histone demethylase or nucleic

acid/protein hydroxylase. Domains other than the JmjC domain determine which

enzymatic functions are exerted; for instance, the tudor domain and PHD domain confer

demethylase activity [44]. However, HRJDs contain only the JmjC domain and the EF

hand motif (Fig. 1B), which makes it difficult to address the molecular functions of

HRJDs. Among the JmjC-domain containing genes, HRJDs have been predicted to be

closest to KDM8 [13], and human KDM8 activates cancer cell proliferation via

demethylation of H3K36me2 on the cyclin A1 locus [45]. These reports imply the

possibility that HRJDs function as histone demethylase like KDM8. If this is the case,

HRJDs must access chromatins; however, we found that ectopically-induced HRJDs

localize in the cytoplasm but not in the nucleus both *in vitro* (S2 cells) and *in vivo* (wing

discs and adult midguts) (Fig. S1B-S1D), which is a common localization pattern of

JmjC-domain only proteins [46, 47]. Interestingly, *Drosophila* JmjC-domain only genes

such as *JMJD4/5/7* genetically interact with loss of function of *snr1*, a component of the

SWI/SNF complex [48], and the mammalian ortholog of *snr1*, SMARCB1, binds to

mitotic chromatin to ensure proliferating activity of daughter cells [49]. It is thus tempting

to speculate that HRJDs may access chromatins during mitosis, when nuclear membrane

disappears, to regulate stem cell functions in the *Drosophila* adult midgut.

Although HRJDs have been identified as genes conserved between highly

regenerative animals [13], it is unclear whether HRJDs alone are sufficient to enhance

regenerative responses. Our investigations revealed that ectopic expression of HRJDs

failed to improve regeneration of developing wing discs and adult midguts in *Drosophila*

(Fig. 2 and 3). However, it should be noted that we induced HRJDs before tissue injury

and maintained their expression during regeneration period. Given that *HRJDb* is up-

regulated in the late phase of planarian regeneration (3 days after amputation) [13], strict

regulation of HRJD induction might be important for their proper function as true

regeneration regulators. On the other hand, the expression level of *HRJDa* remains

constant during planarian regeneration [13], which is recapitulated in our experiments.

Another possibility is that a high regeneration ability can be achieved by cooperation

between HRJDs and other unidentified genes that have been lost during evolution, which

should be addressed in future studies.

During the normal aging process in the *Drosophila* adult midgut, ISC over-

proliferation is closely linked with other aging phenotypes, and suppression of ISC

division can prevent the accumulation of mis-differentiated ISC progenies and extend

organismal lifespan [24, 36, 42, 43]. In contrast, HRJDs enhance age-related ISC

activation but can alleviate mis-differentiation and extend lifespan, suggesting that it is

not ISC activation itself but rather mis-differentiation of ISC progenies that principally

drives age-related mortality. Consistently, manipulation of mitotic spindle orientation,

which affects the cell fate of daughter cells [50–52], can extend organismal lifespan

without changing mitotic activity of ISCs [52]. Of note, in contrast to *Drosophila* ISCs

which over-proliferate in aged midguts, many adult stem cells decrease their activity and

abundance during aging both in *Drosophila* and mammals [24, 53, 54]. Future

investigations for mechanisms of HRJD-mediated stem cell rejuvenation will provide

clues to develop new anti-aging strategies.

Methods

***Drosophila* stocks**

All stocks were maintained on a standard diet containing 4% cornmeal, 6% baker's

yeast (Saf Yeast), 6% glucose (Wako, 049-31177), and 0.8% agar (Kishida chemical,

260-01705) with 0.3% propionic acid (Tokyo Chemical Industry, P0500) and 0.05%

nipagin (Wako, 132-02635). Canton S was utilized as the wildtype strain. Transgenic fly

lines were obtained from the Bloomington Drosophila Stock Center (BDSC) and the

Kyoto Stock Center: *Act5C-Gal4* (BDSC 3954), *rn-Gal4, tub-Gal80ts, UAS-egr* (BDSC

51280), *WP-QF2* (Kashio 2016) [55], *tub-Gal80ts* (BDSC 7017, 7019), *esg-Gal4* (Kyoto

109126), *NP1-Gal4* (Kyoto 112001), *UAS-GFP* (BDSC 1522), M{3xP3-RFP.attP}ZH-

86Fb (Kyoto 130437), UAS-HRJDa (this study), UAS-HRJDb (this study), QUAS-

HRJDa (this study), QUAS-HRJDb (this study), UAS-FLAG-HRJDa (this study), UAS-

FLAG-HRJDb (this study). See **Table S1** for genotypes in each Figure. We used female

flies unless otherwise noted in the Figures.

***Drosophila* genetics**

Experimental crosses that did not involve Gal80^{ts}-mediated inhibition of Gal4 were

performed at 25°C. When using Gal80ts, experimental crosses were maintained at 18°C,

and F1 adults were transferred to 29°C three days after eclosion. Midgut staining and

Smurf assays were performed after 7 or 27 days of 29°C incubation (young: Day 10 adults,

old: Day 30 adults).

Generation of HRJD expressing lines

To express HRJDs using the Gal4/UAS or the QF/QUAS system, we constructed a vector

containing *HRJDa* or *HRJDb* under the control of either the UAS sequence or QUAS

sequence. Namely we amplified HRJDa and HRJDb sequence from codon-optimized

synthesized DNAs (pUCIDT-HRJDa and pUCIDT-HRJDb, IDT, Supplementary texts

S1) for *DjHRJDa* and *DjHRJDb*, respectively. UAS-HRJDa and UAS-HRJDb were

constructed by ligating HRJDa and HRJDb into pUAST-attB (DGRC 1419), respectively

(digested by EcoRI/NotI). Similarly, QUAS-HRJDa and QUAS-HRJDb were constructed

by ligating HRJDa and HRJDb into pQUAS-WALIUM20 (DGRC 1474), respectively

(digested by NheI/EcoRI). To construct UAS-3xFLAG-HRJDs, we PCR-amplified the
coding sequences of HRJDs from pUCIDT-HRJDs by gene-specific primers with
3xFLAG tag at 5'-end of the forward primers, and the amplicons were then ligated into a
pUAST-attB vector digested by EcoRI/NotI using In-Fusion HD Cloning Kit (Takara,
639649). The landing site for each construct was {3xP3-RFP.attP}ZH-86Fb. Injection
and selection were performed by WellGenetics (Taiwan, R.O.C.). Please see also Table
S2 and Supplementary texts S1 for primer sequences and HRJDs sequences.
366
367 **S2 cell culture**
368 *Drosophila* S2 cells were grown at 25°C in Schneider's *Drosophila* medium (GIBCO,
369 21720001) supplemented with 10% (v/v) heat-inactivated fetal bovine serum (FBS), 100
370 U/mL penicillin, and 100 µg/mL streptomycin (FUJIFILM Wako, 168–23191).
371
S2 cell immunostaining
To examine the expression pattern of HRJDs through immunostaining, 1×10^6 cells were
seeded in 6-well plates containing Schneider's *Drosophila* medium supplemented with
10% (v/v) heat-inactivated fetal bovine serum, 100 U/mL penicillin, and 100 µg/mL
streptomycin and were transfected with 800 ng of pAc5-3xFLAG-HRJDs using Effectene
Transfection Reagent (QIAGEN, 301427) following the manufacturer's protocol. To
prepare pAc5-3xFLAG-HRJDs, we amplified the coding sequences of HRJDs from
pUCIDT-HRJDs (Supplementary texts S1) by gene-specific primers with 3xFLAG tag at
5'-end of the forward primers, and the amplicons were then ligated into a pAc5-
STABLE2-neo (Addgene, 32426) [56] digested with EcoRI/XhoI by In-Fusion HD
Cloning Kit (Takara, 639649). Twenty-four hours after transfection, the cells were
washed with PBS and the medium was replaced with fresh Schneider's *Drosophila*
medium supplemented with 10% (v/v) heat-inactivated FBS, 100 U/mL penicillin, and
100 µg/mL streptomycin. Forty-eight hours after transfection, the cells were fixed with
4% paraformaldehyde (PFA) in PBS, washed with PBS containing 0.1% Triton X-100
(PBST), blocked in PBST with 5% normal donkey serum (PBSTn), and incubated with
primary antibodies in PBSTn overnight at 4 °C. The samples were then washed with
PBST, incubated for 1 h at room temperature with secondary antibodies and Hoechst
33342 suspended in PBSTn and washed again with PBST. Images were captured using
an LSM880 with Airyscan (Zeiss). The primary antibody used was a mouse anti-FLAG

M2 monoclonal antibody (1 : 2000, Sigma, F1804). The secondary antibodies used were

Goat anti-Mouse IgG2b Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 555

(1:2000, Thermo Fisher Scientific, A-21147). Hoechst 33342 (0.4 µM; Invitrogen,

H3570) was used for nuclear staining. Images were analyzed and edited using Fiji/ImageJ

software (NIH).

S2 cell western blotting

To examine the expression of HRJDs, 2.5×10^5 cells were seeded in 24-well plates

containing Schneider's *Drosophila* medium supplemented with 10% (v/v) heat-

inactivated FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin and were transfected

with 200 ng of pAc5-3xFLAG-HRJDs using Effectene Transfection Reagent (QIAGEN,

301427) following the manufacturer's protocol. Twenty-four hours after transfection, the

cells were washed with PBS and the medium was replaced with fresh Schneider's

Drosophila medium supplemented with 10% (v/v) heat-inactivated FBS, 100 U/mL

penicillin, and 100 µg/mL streptomycin. Forty-eight hours after transfection, the cells

were washed with PBS and lysed with 50 µL RIPA buffer supplemented with cOmplete

ULTRA EDTA-free protease inhibitor cocktail (Roche, 05892953001). The lysate was

sonicated and centrifuged at 20,000 g for 5 minutes. Protein concentration was

determined by BCA assay. The supernatant was mixed with SDS, boiled at 95 °C for 5

minutes, and subjected to western blotting.

Proteins were separated using SDS-PAGE and transferred onto Immobilon-P

PVDF membranes (Millipore, IPVH00010) for immunoblotting. Membranes were

blocked with 4% skimmed milk diluted in 1×Tris buffered saline containing 0.1% Tween-

20. Immunoblotting was performed using the below-mentioned antibodies, which were

diluted with 4% skim milk. The signals were visualized using Immobilon Western

Chemiluminescent HRP Substrate (Millipore, WBKLS0500) and FUSION SOLO. 7S.

EDGE (Vilber-Lourmat). Contrast and brightness adjustments were applied using the

Fiji/ImageJ software (NIH).

The primary antibody used was mouse anti-FLAG M2 monoclonal antibody

(1 : 5000, Sigma, F1804). Mouse anti-alpha tubulin (DM1A) monoclonal antibody

(1 : 5000, Sigma, T9026) was used as loading control. The secondary antibody used was

HRP-conjugated goat/rabbit/donkey anti-mouse IgG (1 : 10000, Promega, W402B).

Drug treatment

5 mM paraquat (Sigma, 856177) and 5% (w/v) DSS (MP Biomedicals, 160110) were dissolved in 5% (w/v) sucrose solution. Filter paper (Whatman 3MM) was soaked with 400 µl of these reagents and placed into empty vials. For histological analyses, flies were fed with the reagent solution for 1 day. For the survival assay, flies were transferred to new vials and dead flies (determined by immobility and showing no response to tapping) were counted every day. 5% sucrose was used for control feeding.

Immunofluorescence

Samples were dissected in 1X PBS and fixed in 4% PFA for 20 minutes (wing discs) and 30-45 minutes (adult midgut) at room temperature (RT), respectively. The following primary antibodies were used with indicated dilution into 1X PBS containing 0.5% BSA and 0.1% Triton X-100: rabbit anti-PH3 (Millipore 06-570, 1:1000), mouse anti-FLAG (Sigma F1804, 1:1000), mouse anti-Dlg (DSHB 4F3, 1:100), rabbit anti-GFP (MBL 598, 1:500), rat anti-GFP (Nacalai tesque 04404-26, 1:400), rabbit anti-Tsp2A (Izumi et al., 2016, 1:1000) [57], rabbit anti-Mesh (Izumi et al., 2012, 1:1000) [58], rabbit anti-Ssk (Izumi et al., 2016, 1:1000) [57]. After overnight incubation with primary antibodies at 4°C, samples were incubated with fluorescent secondary antibodies (Jackson ImmunoResearch and Invitrogen, 1:500) for 1 hour at RT. Hoechst 33342 (Invitrogen, final concentration: 10 µg/ml) was used to visualize DNA. Wing discs were mounted as described previously [59, 60]. Samples were mounted in Slowfade Diamond (ThermoFisher, S36963) and imaged with confocal microscopy (Zeiss LSM880).

Smurf assay

We referred to Rera et al. (2012) for the Smurf assay conditions [35]. To prepare the feeding medium, 100 µl of 50% (w/v) brilliant blue FCF (Wako, 027-12842, final 2.5%) and 100 µl of 5% (w/v) sucrose were added to a vial containing 2 ml of cornmeal-agar food. After mixing with a spatula, an Whatman 1 filter (1001-020) was put on the feeding medium. Flies were fed with this medium at 25°C for one day, after which they were transferred to a new vial containing cornmeal-agar food without blue dye to clean the epidermis. Two hours after the transfer, Smurf phenotype was checked. We classified the Smurf phenotype as blue dye leakage outside abdomen (thorax, head, legs).

Phylogenetic tree

Phylogenetic tree

We created the phylogenetic tree of representative organisms using the NCBI taxonomy

browser (https://www.ncbi.nlm.nih.gov/Taxonomy/CommonTree/wwwcmt.cgi). The

species examined are *Amphimedon queenslandica* (Porifera), *Mnemiopsis leidyi*

(Ctenophora), *Trichoplax adhaerens* (Placozoa), *Hydra vulgaris* (Cnidaria), *Ptychodera*

flava (Hemichordata), *Acanthaster planci* (Echinodermata), *Branchiostoma floridae*

(Cephalochordata), *Ciona intestinalis* (Tunicata), *Danio rerio* (Teleostei), *Homo sapiens*,

Mus musculus (Mammalia), *Schmidtea mediterranea* (Platyhelminthes), *Caenorhabditis*

elegans (Nematoda), and *Drosophila melanogaster* (Arthropoda). These species, except

for *D. rerio, H. sapiens, M. musculus, C. elegans,* and *D. melanogaster,* are known to

have HRJDs and exhibit high regeneration ability [13]. The pictures were downloaded

from PhyloPic, and the color of some were changed from black to blue. Credits: Bennet

McComish (*B. floridae*), Malio Kodis (*M. leidyi*), and Markus A. Grohme (*S.*

mediterranea, https://www.phylopic.org/images/93f9611c-0cbd-4a14-92da-

004a4521e21f/schmidtea-mediterranea).

Quantification of junctional proteins

To quantify junctional and cytoplasmic intensity, a line that was orthogonal to one side

of a cell was drawn, and then the plot profile was calculated for the line. We defined the

junctional intensity as the highest value in that prolife, and defined cytoplasmic intensity

as the value 20 pixels (~2.6 μm) away from the highest value. Three lines were drawn for

one cell and the average of the three was used to represent the junctional/cytoplasmic

ratio for the cell. Six polyploid cells were randomly selected from each image for

quantification.

Quantification of nuclear size

Nuclei of *esg-Gal4>UAS-GFP* positive cells were traced by polygon selection tool and

added as ROI in Fiji/ImageJ. The size of each ROI was quantified using the Measure

command in Fiji/ImageJ.

Statistics

Statistical analyses were performed using Excel and RStudio. Two tailed *t* tests were used

for comparisons between two groups. One-way ANOVAs with post hoc Tukey tests were

performed when comparing three or more groups. Log-rank tests were used for
comparison of survival curve. Significance is indicated in the Fig.s as follows: *P≤0.05,
**P≤0.01, ***P≤0.001, Not Significant (N.S.): P>0.05. Bar graphs show mean ± standard
error. Boxplots show median (thick line in the box), first and third quartiles (bottom and
top of the box), minimum value (lower whisker), and maximum value (upper whisker).
Dots in bar graphs and boxplots indicate individual values. Violin plots portray the
distribution of individual values.

DECLARATIONS

Ethics approval and consent to participate
Not applicable.

Consent for publication
Not applicable.

Availability of data and materials
The datasets and *Drosophila* stocks used and/or analysed during the current study are
available from the corresponding author on reasonable request.

Competing interests
The authors declare that they have no competing interests.

Funding
This work was supported by JSPS/MEXT KAKENHI (grant numbers JP22J01430 to
H.N., JP21H04774, JP21K19206 to M.M., and JP17H06332, JP19K22550, JP22H02762
to Y.N.), AMED-Aging (JP21gm5010001 to M.M.), AMED-PRIME (JP22gm6110025
to Y.N.), and Sadako O. Hirai Ban Award for Young Researchers (H.N.)

Author contributions
Conceptualization: H.N., Y.N.

Investigation: H.N., J.U., T.N., Y.N.

Methodology: H.N., Y.N.

Validation: H.N., J.U., T.N., T.M., M.M., Y.N.

Data curation: H.N., T.M., Y.N.

Writing – original draft: H.N., J.U., T.M., Y.N.

Writing – review & editing: H.N., J.U., T.N., T.M., M.M., Y.N.

Supervision: M.M., Y.N.

Funding acquisition: H.N., T.M., M.M., Y.N.

Acknowledgements

We thank N. Shinoda, M. Furuse, Y. Izumi, BDSC, Kyoto Stock Center, Transgenic RNAi

Project (TRiP), and DSHB for fly stocks and reagents.

 FIGURE LEGENDS

Figure 1. HRJD expression in whole body does not affect gross morphology.

(A) Phylogenetic tree of HRJD conservation. Green indicates species that possess HRJD

gene(s) and gray indicates species that have lost HRJD gene(s). High regenerative ability

indicates that the species can regenerate their whole body or anterior/posterior body parts,

and low regenerative ability indicates that the species can only regenerate their appendage

(limbs, tails, fins) or much smaller scale of tissues/organs. We referred to Cao et al., 2019

for the definition of regenerative ability [13].

(B) Protein sequence of HRJDs used in this study, which derived from *Dugesia japonica*

(DjHRJDa/b, hereafter simply described as HRJDa/b). Conserved residues characteristic

of the JmjC domain are shown.

(C) Schematics of genetic experiments for whole body induction of HRJDs. The Gal4-

UAS system enables gene expression downstream of the UAS sequence, which is

regulated by the transcription factor Gal4 [15]. In this case, the ubiquitously active *Act5C*

promoter is used for Gal4 expression.

(D) Representative images of mature adults for whole body expression of HRJDs.

(E) Whole body induction of HRJDs did not change body weight of mature adults.

(F) Representative images of adult wings.

(G) Whole body induction of HRJDs did not change wing size.

(H) Representative images for adult heads.

(I) Whole body induction of HRJDs did not change eye size of adult flies.

N.S., not significant: $P > 0.05$. One-way ANOVAs with post hoc Tukey test. *n* indicates

the number of flies examined. See also Fig. S1.

Figure 2. HRJD expression impedes tissue regeneration in the developing wind disc

(A) Schematics of genetic ablation of the wing disc. TNF ligand *eiger* was overexpressed

in the wing pouch for 40 hrs during the larval stage.

(B) Wing regeneration was assessed in the adult stage by checking the size of wing (0%:

no wing, 100%: intact wing).

(C) Line graph for wing regeneration. Expression of HRJDa/b increased the rate of low

regeneration (0% and 25%) at the expense of the rate of full regeneration (100%).

(D-E) Representative images for adult wings of *WP-QF>QUAS-HRJDa/b* flies under

homeostatic (non-regenerative) condition (D). The wing size is quantified in (E).

N.S., not significant: $P>0.05$, ***$P \leq 0.001$. Chi-square test (C) and one-way ANOVAs

with post hoc Tukey test (E). *n* indicates the number of wings examined.

Figure 3. HRJDs expression compromises regeneration in the adult midgut

(A) Schematics of the *Drosophila* adult midgut. The adult midgut is a pseudostratified

epithelium, in which intestinal stem cells (ISCs) generate both absorptive enterocytes

(ECs) and secretory enteroendocrine cells (EEs) through progenitor cell enteroblasts

(EBs) and EE progenitor cells (EEPs).

(B) Experimental scheme for ISC/EB specific induction of HRJDs under the *esg-Gal4*

driver in the adult stage. Adult flies were transferred to 29°C at Day 3 to induce HRJD

expression.

(C-E) Survival curve during paraquat (C), DSS (D), and sucrose (E) feeding.

(F) Representative images for PH3 staining upon paraquat feeding.

(G) Quantification of PH3 positive cells in posterior midguts.

N.S., not significant: P>0.05, *P<0.05, **P≤0.01. Log-rank test. *n* indicates the number

of flies (C-E) and midguts (G). Scale bars: 50 μm. See also Fig. S2.

Figure 4. Post-developmental HRJD expression in ISC lineages extends organismal

lifespan

(A-D) Survival curve for organismal lifespan by sex. HRJDs are induced by *esg^ts-Gal4*

(A and B) and *Act5C-Gal4* (C and D). For temporal regulation of HRJD induction using

tub-Gal80^ts (A and B), adult flies were transferred to 29°C at Day 3.

N.S., not significant: P>0.05, *P<0.05, **P≤0.01, ***P≤0.001, Log-rank tests. *n*

indicates the number of flies. See also Fig. S3.

Figure 5. HRJD expression in adult ISCs/EBs suppress age-related gut barrier

dysfunction

(A) *esg^ts*-mediated HRJD induction decreased Smurf(+) ratio at Day 30. The

representative images of the Smurf phenotype (leakage of blue dye throughout whole

body) is shown on the left.

(B-C) HRJD expression in adult ISCs/EBs maintained junctional localization of Dlg

protein in ECs, which is quantified in (C).

(D-I) HRJD expression in adult ISCs/EBs maintained junctional localization of Tsp2A

(D), Mesh (E), and Ssk (F) in ECs, which is quantified in (G-I).

*P<0.05, ***P≤0.001, One-way ANOVAs with post hoc Tukey test. *n* indicates the

number of flies (A) and cells (C, G, H, I) examined. Scale bars: 50 μm.

Figure 6. HRJD expression in adult ISCs/EBs improves stem cell functions in aged

midguts

(A) Representative images for *esg-Gal4* positive cells in Day 30 midguts. HRJDs

suppressed mis-differentiation of esg^+ cells (cellular enlargement with polyploid large

nuclei).

(B) Quantification of nuclear size of *esg-Gal4* positive cells.

(C-D) Representative images (C) and quantification (D) of PH3 positive cells in Day 30

posterior midguts. Day 10 young guts are also quantified in (D).

(E) RT-qPCR of *Dl*, *nub*, and *pros* in Day 30 midguts. Expression levels were normalized

to those of GFP control.

(F) Schematic model for the impact of HRJDs on intestinal homeostasis. In old wildtype

midguts, ISCs over-proliferate, and their daughters fail to differentiate into ECs, resulting

in accumulation of mis-differentiated cells (*esg*+ large polyploid). HRJD expression in

adult ISCs/EBs further enhances ISC proliferation but suppresses mis-differentiation.

N.S., not significant: P>0.05, *P≤0.05, **P≤0.01, ***P≤0.001, one-way ANOVAs with

post hoc Tukey test. *n* indicates the number of cells (B), guts (D), and biological replicates

(E). Scale bars: 50 μm.

 Supplemental figure legends

Figure S1. Expression and localization of HRJD proteins

(A) Western blots of S2 cell samples upon transfection of 3×FLAG-HRJDa/b. Anti-

FLAG was used to detect 3×FLAG-HRJDa/b proteins and anti-tubulin was used as the

loading control. Expected molecular weight: 52.0 kDa for 3×FLAG-HRJDa, 52.9 kDa

for 3×FLAG-HRJDb.

(B) Representative images of anti-FLAG staining using S2 cells transfected with

3×FLAG-HRJDa/b. Both HRJDa and HRJDb localized cytoplasm but not in nuclei. Scale

bar: 30 μm in wide-view images (left), 5 μm in magnified view images (yellow frames).

(C-D) Representative images of 3×FLAG-HRJDa/b localization in wing discs and adult

midguts. FLAG-tagged HRJDs were induced using *WP-Gal4* (C) or *esg-Gal4* (D). Scale

bars: 10 μm.

Figure S2. Continuous expression of HRJDs throughout development hampers

intestinal regeneration

(A-C) Organismal survival upon paraquat (A), DSS (B), and sucrose (C) feeding. Drug

feeding started at Day 7 after eclosion. HRJD induction using *esg-Gal4* throughout

development significantly decreased survival upon paraquat/DSS feeding.

(D) Expression of HRJDa using *esg-Gal4* decreased the number of PH3 positive cells in

the posterior midguts, while HRJDb did not change PH3 positive cell number.

N.S., not significant: P>0.05, **P≤0.01, ***P≤0.001, one-way ANOVAs with post hoc

Tukey test. *n* indicates the number of flies (A-C) and guts (D).

Figure S3. Organismal lifespan of flies expressing HRJDs in ECs and ISC/EBs

(A-F) Survival curve for organismal lifespan by sex. HRJDs are induced by *NP1-Gal4*

(A and B), *NP1^{ts}-Gal4* (C and D), and *esg-Gal4* (E and F).

N.S., not significant: P>0.05, ***P≤0.001, Log-rank tests. *n* indicates the number of flies.

Table S1. Detailed genotypes in each experiment

Fig.	Panel	Genotype
Fig. 1	**D-I**	*w;; Act5C-Gal4 / +*
		w; UAS-GFP / +; Act5C-Gal4 / UAS-HRJDb
		w;; Act5C-Gal4 / UAS-HRJDa
		w;; Act5C-Gal4 / UAS-HRJDb
Fig. 2	**A-C**	*w; rn-Gal4, tub-Gal80ts, UAS-egr / +; WP-QF2 / +*
		w; rn-Gal4, tub-Gal80ts, UAS-egr / +; WP-QF2 / QUAS-HRJDa
		w; rn-Gal4, tub-Gal80ts, UAS-egr / +; WP-QF2 / QUAS-HRJDb
	D-E	*w;; WP-QF2 / +*
		w;; WP-QF2 / QUAS-HRJDa
		w;; WP-QF2 / QUAS-HRJDb
Fig. 3	**C-G**	*w; esg-Gal4, tub-Gal80ts / UAS-GFP*
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDa / +
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDb / +
Fig. 4	**A-B**	*w; esg-Gal4, tub-Gal80ts / UAS-GFP*
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDa / +
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDb / +
	C-D	*w; UAS-GFP / +; Act5C-Gal4 / +*
		w;; Act5C-Gal4 / UAS-HRJDa
		w;; Act5C-Gal4 / UAS-HRJDb
Fig. 5, 6	**All**	*w; esg-Gal4, tub-Gal80ts / UAS-GFP*
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDa / +
		w; esg-Gal4, tub-Gal80ts / +; UAS-HRJDb / +
Fig. S1	**C**	*w;; WP-Gal4 / +*
		w;; WP-Gal4 / UAS-FLAG-HRJDa
		w;; WP-Gal4 / UAS-FLAG-HRJDb
	D	*w; esg-Gal4, tub-Gal80ts, UAS-GFP / +*
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; UAS-FLAG-HRJDa / +
		w; esg-Gal4, tub-Gal80ts, UAS-GFP / +; UAS-FLAG-HRJDb / +
Fig. S2	**All**	*w; esg-Gal4 / UAS-GFP*

		w; esg-Gal4 / +; UAS-HRJDa / +	
		w; esg-Gal4 / +; UAS-HRJDb / +	
Fig. S3	**A-B**	*w; NP1-Gal4 / UAS-GFP*	
		w; NP1-Gal4 / +; UAS-HRJDa / +	
		w; NP1-Gal4 / +; UAS-HRJDb / +	
	C-D	*w; NP1-Gal4, tub-Gal80ts / UAS-GFP*	
		w; NP1-Gal4, tub-Gal80ts / +; UAS-HRJDa / +	
		w; NP1-Gal4, tub-Gal80ts / +; UAS-HRJDb / +	
	E-F	*w; esg-Gal4 / UAS-GFP*	
		w; esg-Gal4 / +; UAS-HRJDa / +	
		w; esg-Gal4 / +; UAS-HRJDb / +	

References

1. Bely AE, Nyberg KG. Evolution of animal regeneration: re-emergence of a field. Trends Ecol Evol. 2010;25:161–70.

2. Poss KD. Advances in understanding tissue regenerative capacity and mechanisms in animals. Nat Rev Genet. 2010;11:710–22.

3. Tiozzo S, Copley RR. Reconsidering regeneration in metazoans: an evo-devo approach. Front Ecol Evol. 2015;3.

4. Bideau L, Kerner P, Hui J, Vervoort M, Gazave E. Animal regeneration in the era of transcriptomics. Cell Mol Life Sci. 2021;78:3941–56.

5. Fujita S, Kuranaga E, Nakajima Y. Regeneration Potential of Jellyfish: Cellular Mechanisms and Molecular Insights. Genes. 2021;12:758.

6. Piraino S, Boero F, Aeschbach B, Schmid V. Reversing the Life Cycle: Medusae Transforming into Polyps and Cell Transdifferentiation in *Turritopsis nutricula* (Cnidaria, Hydrozoa). Biol Bull. 1996;190:302–12.

7. Varley Á, Horkan HR, McMahon ET, Krasovec G, Frank U. Pluripotent, germ cell competent adult stem cells underlie cnidarian regenerative ability and clonal growth. Curr Biol. 2023;33:1883-1892.e3.

8. Gazave E, Röttinger E. 7th Euro Evo Devo meeting: Report on the "Evolution of regeneration in Metazoa" symposium. J Exp Zoolog B Mol Dev Evol. 2021;336:89–93.

9. Kumar A, Godwin JW, Gates PB, Garza-Garcia AA, Brockes JP. Molecular Basis for

the Nerve Dependence of Limb Regeneration in an Adult Vertebrate. Science.

2007;318:772–7.

10. Da Silva SM, Gates PB, Brockes JP. The Newt Ortholog of CD59 Is Implicated in

Proximodistal Identity during Amphibian Limb Regeneration. Dev Cell. 2002;3:547–

55.

11. Looso M, Preussner J, Sousounis K, Bruckskotten M, Michel CS, Lignelli E, et al.

A de novo assembly of the newt transcriptome combined with proteomic validation

identifies new protein families expressed during tissue regeneration. Genome Biol.

2013;14:R16.

12. Mehta AS, Deshpande P, Chimata AV, Tsonis PA, Singh A. Newt regeneration

genes regulate Wingless signaling to restore patterning in Drosophila eye. iScience.

2021;24:103166.

13. Cao P-L, Kumagai N, Inoue T, Agata K, Makino T. JmjC Domain-Encoding Genes

Are Conserved in Highly Regenerative Metazoans and Are Associated with Planarian

Whole-Body Regeneration. Genome Biol Evol. 2019;11:552–64.

14. Umesono Y, Tasaki J, Nishimura Y, Hrouda M, Kawaguchi E, Yazawa S, et al. The

molecular logic for planarian regeneration along the anterior–posterior axis. Nature.

2013;500:73–6.

15. Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates

and generating dominant phenotypes. Development. 1993;118:401–15.

16. Hariharan IK, Serras F. Imaginal disc regeneration takes flight. Curr Opin Cell Biol.

2017;48:10–6.

17. Smith-Bolton RK, Worley MI, Kanda H, Hariharan IK. Regenerative Growth in

Drosophila Imaginal Discs Is Regulated by Wingless and Myc. Dev Cell. 2009;16:797–

809.

18. Worley MI, Alexander LA, Hariharan IK. CtBP impedes JNK- and Upd/STAT-

driven cell fate misspecifications in regenerating Drosophila imaginal discs. eLife.

2018;7:e30391.

19. Potter CJ, Tasic B, Russler EV, Liang L, Luo L. The Q System: A Repressible

Binary System for Transgene Expression, Lineage Tracing, and Mosaic Analysis. Cell.

2010;141:536–48.

20. Riabinina O, Luginbuhl D, Marr E, Liu S, Wu MN, Luo L, et al. Improved and

expanded Q-system reagents for genetic manipulations. Nat Methods. 2015;12:219–22.

21. Nagai H, Miura M, Nakajima Y. Cellular mechanisms underlying adult tissue

plasticity in Drosophila. Fly (Austin). 2022;16:190–206.

22. Miguel-Aliaga I, Jasper H, Lemaitre B. Anatomy and Physiology of the Digestive

Tract of *Drosophila melanogaster*. Genetics. 2018;210:357–96.

23. Zhang P, Edgar BA. Insect Gut Regeneration. Cold Spring Harb Perspect Biol.

2021;:a040915.

24. Jasper H. Intestinal Stem Cell Aging: Origins and Interventions. Annu Rev Physiol.

2020;82:203–26.

25. Buchon N, Broderick NA, Chakrabarti S, Lemaitre B. Invasive and indigenous

microbiota impact intestinal stem cell activity through multiple pathways in *Drosophila*.

Genes Dev. 2009;23:2333–44.

26. Jiang H, Patel PH, Kohlmaier A, Grenley MO, McEwen DG, Edgar BA.

Cytokine/Jak/Stat Signaling Mediates Regeneration and Homeostasis in the Drosophila

Midgut. Cell. 2009;137:1343–55.

27. Biteau B, Hochmuth CE, Jasper H. JNK Activity in Somatic Stem Cells Causes

Loss of Tissue Homeostasis in the Aging Drosophila Gut. Cell Stem Cell. 2008;3:442–

55.

28. Amcheslavsky A, Jiang J, Ip YT. Tissue Damage-Induced Intestinal Stem Cell

Division in Drosophila. Cell Stem Cell. 2009;4:49–61.

29. Jiang H, Edgar BA. EGFR signaling regulates the proliferation of *Drosophila* adult

midgut progenitors. Development. 2009;136:483–93.

30. Houtz P, Bonfini A, Bing X, Buchon N. Recruitment of Adult Precursor Cells

Underlies Limited Repair of the Infected Larval Midgut in Drosophila. Cell Host

Microbe. 2019;26:412-425.e5.

31. Rera M, Bahadorani S, Cho J, Koehler CL, Ulgherait M, Hur JH, et al. Modulation

of Longevity and Tissue Homeostasis by the Drosophila PGC-1 Homolog. Cell Metab.

2011;14:623–34.

32. Thevaranjan N, Puchta A, Schulz C, Naidoo A, Szamosi JC, Verschoor CP, et al.

Age-Associated Microbial Dysbiosis Promotes Intestinal Permeability, Systemic

Inflammation, and Macrophage Dysfunction. Cell Host Microbe. 2017;21:455-466.e4.

33. Nagai H, Tatara H, Tanaka-Furuhashi K, Kurata S, Yano T. Homeostatic Regulation

of ROS-Triggered Hippo-Yki Pathway via Autophagic Clearance of Ref(2)P/p62 in the

Drosophila Intestine. Dev Cell. 2021;56:81-94.e10.

34. Egge N, Arneaud SLB, Wales P, Mihelakis M, McClendon J, Fonseca RS, et al.

Age-Onset Phosphorylation of a Minor Actin Variant Promotes Intestinal Barrier

Dysfunction. Dev Cell. 2019;51:587-601.e7.

35. Rera M, Clark RI, Walker DW. Intestinal barrier dysfunction links metabolic and

inflammatory markers of aging to death in Drosophila. Proc Natl Acad Sci.

2012;109:21528–33.

36. Resnik-Docampo M, Koehler CL, Clark RI, Schinaman JM, Sauer V, Wong DM, et

al. Tricellular junctions regulate intestinal stem cell behaviour to maintain homeostasis.

Nat Cell Biol. 2017;19:52–9.

37. Salazar AM, Resnik-Docampo M, Ulgherait M, Clark RI, Shirasu-Hiza M, Jones

DL, et al. Intestinal Snakeskin Limits Microbial Dysbiosis during Aging and Promotes

Longevity. iScience. 2018;9:229–43.

38. Izumi Y, Furuse K, Furuse M. Septate junctions regulate gut homeostasis through

regulation of stem cell proliferation and enterocyte behavior in *Drosophila*. J Cell Sci.

2019;:jcs.232108.

39. Chen H, Zheng X, Zheng Y. Age-Associated Loss of Lamin-B Leads to Systemic

Inflammation and Gut Hyperplasia. Cell. 2014;159:829–43.

40. Korzelius J, Naumann SK, Loza-Coll MA, Chan JS, Dutta D, Oberheim J, et al.

Escargot maintains stemness and suppresses differentiation in *Drosophila* intestinal

stem cells. EMBO J. 2014;33:2967–82.

41. Li Y, Pang Z, Huang H, Wang C, Cai T, Xi R. Transcription Factor Antagonism

Controls Enteroendocrine Cell Specification from Intestinal Stem Cells. Sci Rep.

2017;7:988.

42. Clark RI, Salazar A, Yamada R, Fitz-Gibbon S, Morselli M, Alcaraz J, et al.

Distinct Shifts in Microbiota Composition during Drosophila Aging Impair Intestinal

Function and Drive Mortality. Cell Rep. 2015;12:1656–67.

43. Wang L, Ryoo HD, Qi Y, Jasper H. PERK Limits Drosophila Lifespan by

Promoting Intestinal Stem Cell Proliferation in Response to ER Stress. PLOS Genet.

2015;11:e1005220.

44. Markolovic S, Leissing TM, Chowdhury R, Wilkins SE, Lu X, Schofield CJ.

Structure–function relationships of human JmjC oxygenases—demethylases versus

hydroxylases. Curr Opin Struct Biol. 2016;41:62–72.

45. Hsia DA, Tepper CG, Pochampalli MR, Hsia EYC, Izumiya C, Huerta SB, et al.

KDM8, a H3K36me2 histone demethylase that acts in the cyclin A1 coding region to

regulate cancer cell proliferation. Proc Natl Acad Sci. 2010;107:9671–6.

46. Shalaby NA, Sayed R, Zhang Q, Scoggin S, Eliazer S, Rothenfluh A, et al.

Systematic discovery of genetic modulation by Jumonji histone demethylases in

Drosophila. Sci Rep. 2017;7:5240.

47. Klose RJ, Kallin EM, Zhang Y. JmjC-domain-containing proteins and histone

demethylation. Nat Rev Genet. 2006;7:715–27.

48. Curtis BJ, Zraly CB, Marenda DR, Dingwall AK. Histone lysine demethylases

function as co-repressors of SWI/SNF remodeling activities during Drosophila wing

development. Dev Biol. 2011;350:534–47.

49. Zhu Z, Chen X, Guo A, Manzano T, Walsh PJ, Wills KM, et al. Mitotic

bookmarking by SWI/SNF subunits. Nature. 2023;618:180–7.

50. Nakajima Y. Mitotic spindle orientation in epithelial homeostasis and plasticity. J

Biochem (Tokyo). 2018;164:277–84.

51. Goulas S, Conder R, Knoblich JA. The Par Complex and Integrins Direct

Asymmetric Cell Division in Adult Intestinal Stem Cells. Cell Stem Cell. 2012;11:529–

40.

52. Hu DJ-K, Jasper H. Control of Intestinal Cell Fate by Dynamic Mitotic Spindle

Repositioning Influences Epithelial Homeostasis and Longevity. Cell Rep.

2019;28:2807-2823.e5.

53. Brunet A, Goodell MA, Rando TA. Ageing and rejuvenation of tissue stem cells and

their niches. Nat Rev Mol Cell Biol. 2023;24:45–62.

54. Campisi J, Kapahi P, Lithgow GJ, Melov S, Newman JC, Verdin E. From

discoveries in ageing research to therapeutics for healthy ageing. Nature.

2019;571:183–92.

55. Kashio S, Obata F, Zhang L, Katsuyama T, Chihara T, Miura M. Tissue

nonautonomous effects of fat body methionine metabolism on imaginal disc repair in

Drosophila. Proc Natl Acad Sci. 2016;113:1835–40.

56. González M, Martín-Ruíz I, Jiménez S, Pirone L, Barrio R, Sutherland JD.

Generation of stable Drosophila cell lines using multicistronic vectors. Sci Rep.

2011;1:75.

57. Izumi Y, Motoishi M, Furuse K, Furuse M. A tetraspanin regulates septate junction

formation in *Drosophila* midgut. J Cell Sci. 2016;:jcs.180448.

58. Izumi Y, Yanagihashi Y, Furuse M. A novel protein complex, mesh-ssk, is required

for septate junction formation in *drosophila* midgut. J Cell Sci. 2012;:jcs.112243.

59. Nakajima Y. Analysis of Epithelial Architecture and Planar Spindle Orientation in

the Drosophila Wing Disc. In: Turksen K, editor. Stem Cell Renewal and Cell-Cell

Communication. New York, NY: Springer US; 2020. p. 51–62.

60. Nakajima Y, Lee ZT, McKinney SA, Swanson SK, Florens L, Gibson MC.

Junctional tumor suppressors interact with 14-3-3 proteins to control planar spindle

alignment. J Cell Biol. 2019;218:1824–38.

A

rn-Gal4, tub-Gal80ᵗˢ>UAS-eiger

B

No wing

0%   25%   50%   75%   100%

C

□ WP-QF (n=606)
◇ WP-QF, HRJDa (n=603)
▲ WP-QF, HRJDb (n=602)

***

Percent of wings

Regeneration (%)

D

Normal development

WP-QF2

WP-QF2 HRJDa

WP-QF2 HRJDb

E

N.S.

Wing size (mm²)

29   30   26

WP-QF2   WP-QF2, HRJDa   WP-QF2, HRJDb

A

Lumen
Epithelia
Visceral muscle

ISC (esg⁺) → EB (esg⁺) → EC (NP1⁺)
ISC (esg⁺) → EEp → EE (Pros⁺)

B ISC/EB specific overexpression of HRJDs in adult stage

esg-Gal4, tub-Gal80ᵗˢ>UAS-HRHDa/b

29°C — PH3 — Paraquat or DSS
18°C
Eclosion   Day 3   D10 D11
HRJD overexpression

C Paraquat feeding, *esgᵗˢ>*

□ GFP (n=144)
◇ HRJDa (n=158)
▲ HRJDb (n=153)

D DSS feeding, *esgᵗˢ>*

□ GFP (n=86)
◇ HRJDa (n=105)
▲ HRJDb (n=95)

E Control sucrose feeding, *esgᵗˢ>*

□ GFP (n=115)
◇ HRJDa (n=118)
▲ HRJDb (n=107)

F Paraquat feeding, *esgᵗˢ>*

PH3 / Hoechst
PH3

GFP   HRJDa   HRJDb

G Paraquat feeding, *esgᵗˢ>*

N.S.
(21) (20) (16)
GFP HRJDa HRJDb

A Female lifespan, *esg^ts-Gal4>*

GFP (n=177)
HRJDa (n=175) ***
HRJDb (n=153) **

B Male lifespan, *esg^ts-Gal4>*

GFP (n=179)
HRJDa (n=178) **
HRJDb (n=153) *

C Female lifespan, *Act-Gal4>*

GFP (n=218)
HRJDa (n=208) N.S.
HRJDb (n=216)

D Male lifespan, *Act-Gal4>*

GFP (n=235)
HRJDa (n=198) ***
HRJDb (n=230)

A

esg^ts>GFP (Day 30)

GFP Hoechst

GFP

\+   HRJDa   HRJDb

B

esg^ts>
(Day 30)   GFP   HRJDa   HRJDb

C

esg^ts>GFP (Day 30)

PH3 Hoechst

PH3

\+   HRJDa   HRJDb

D

esg^ts>   GFP   HRJDa   HRJDb   GFP   HRJDa   HRJDb
              Day 10   Day 30

E

n=6

GFP
HRJDa
HRJDb

Dl   nub   pros

F

WT
(young)

WT
(old)

Barrier function ↓

HRJDs in ISCs/EBs
(old)

Barrier function ↑
Lifespan ↑