TITLE: Functional variations in gamma-secretase activity are critical determinants of the

clinical, biomarker, and cognitive progression of autosomal dominant Alzheimer's

disease

AUTHOR LIST Stephanie A. Schultz, PhD[1], Lei Liu, MD, PhD[2,3], Aaron P. Schultz,

PhD[1], Colleen D. Fitzpatrick, MA[1], Raina Levin, BS[1], Jean-Pierre Bellier, PhD[2,3], Zahra

Shirzadi, PhD[2,3], Nelly Joseph-Mathurin, PhD[4], Charles D. Chen, PhD[4], Tammie L.S.

Benzinger, MD, PhD[4], Gregory S. Day, MD[5], Martin R. Farlow, MD[6], Brian A. Gordon,

PhD[4], Jason J. Hassenstab, PhD[4], Clifford R. Jack Jr., MD[7], Mathias Jucker, PhD[8],

Celeste M. Karch, PhD[4], Jae-Hong Lee, MD, PhD[9], Johannes Levin, MD[10-12], Richard J.

Perrin, MD, PhD[4], Peter R Schofield, PhD, DSc[13,14], Chengjie Xiong, PhD[4], Keith A.

Johnson, MD[1,2], Eric McDade, DO[4], Randall J. Bateman, MD[4], Reisa A. Sperling, MD[1,2],

Dennis J. Selkoe, MD[2,3], Jasmeer P. Chhatwal, MD, PhD, MMSc*[1,2,3], and the

Dominantly Inherited Alzheimer Network Investigators

[1]Massachusetts General Hospital, Harvard Medical School; Boston, MA, USA

[2]Brigham and Women's Hospital; Boston, MA, USA

[3]Ann Romney Center for Neurologic Diseases; Boston, MA, USA

[4]Washington University in St. Louis School of Medicine; St. Louis, MO, USA

[5]Mayo Clinic; Jacksonville, FL, USA

[6]Indiana Alzheimer's Disease Research Center; Indianapolis, IN, USA

[7]Mayo Clinic, Rochester; MN, USA

[8]German Center for Neurodegenerative Diseases (DZNE); Tuebingen, Germany

[9]Asan Medical Center, University of Ulsan College of Medicine; Seoul, Korea

[10]German Center for Neurodegenerative Diseases (DZNE); Munich, Germany

[11]Department of Neurology, Ludwig-Maximilians-Universität München; Munich,
Germany

[12]Munich Cluster for Systems Neurology (SyNergy); Munich, Germany

[13]Neuroscience Research Australia; Randwick, NSW, Australia

[14]School of Medical Sciences, University of New South Wales; Sydney, NSW, Australia
30

31 *Corresponding author: Jasmeer P. Chhatwal MD, PhD, MMSc; Email:
32 Chhatwal.Jasmeer@mgh.harvard.edu

36
ABSTRACT

Background: The balance between production, clearance, and toxicity of Aβ peptides

is central to Alzheimer's disease (AD) pathobiology. Though highly variable in terms of

age at symptom onset (AAO), hundreds of variants in *PSEN1* cause autosomal

dominant forms of AD (ADAD) with nearly complete penetrance. PSEN1 forms the

catalytic core of the γ-secretase complex and thereby directly mediates the production

of longer, aggregation-prone Aβ peptides relative to shorter, non-aggregating peptides.

We hypothesized that the broad AAO and biomarker heterogeneity seen across ADAD

would be predictable based on mutation-specific differences in the production of Aβ

species.

Methods: Aβ-37, 38, 40, 42, and 43 production was quantified from 162 unique *PSEN1*

variants expressed in HEK293 cells. Prediction of AAO was carried out in 107 variants

with available AAO and then replicated in 55 variants represented across 190 PSEN1

mutation carriers who have detailed cognitive and biomarker data from the Dominantly

Inherited Alzheimer's Network (DIAN).

Results: Variations in Aβ production across the 162 mutations examined in cell-based

models were highly predictive of AAO. In those with corresponding *in vivo* data from the

DIAN study, our cell-based γ-secretase composite was strongly associated with

biomarker and cognitive trajectories.

Conclusions: These findings elucidate the critical link between γ-secretase function,

Aβ production, and AD progression and offer mechanistic support for the amyloid

hypothesis. The approach used here represents a powerful tool to account for

heterogeneity in disease progression in ADAD clinical trials and to assess the

pathogenicity of variants of unknown significance or with limited family history.

INTRODUCTION

An increase in longer, aggregation-prone Aβ fragments (e.g., Aβ 42 and 43) relative to shorter, non-aggregating fragments (e.g., Aβ 37 and 38) is a critical initiating pathogenic event in both late-onset, sporadic AD[1] (LOAD) and Autosomal Dominant AD[2–4] (ADAD). The balance between production of aggregating and non-aggregating forms of Aβ is a direct result of the efficiency and kinetics with which the γ-secretase proteolytic complex sequentially cleaves β-amyloid precursor protein[5–9] (APP). Subtle alterations in γ-secretase function can lead to profound neurodegenerative and cognitive consequences, while modulation of γ-secretase has therapeutic potential in both LOAD and ADAD[8–12]. Over 200 pathogenic variants in *Presenilin-1* (*PSEN1*), the key catalytic subunit in the γ-secretase complex, have been identified and are the most common causes of ADAD[13]. Mirroring the heterogeneity in clinical and AD biomarker progression seen in LOAD, there is striking heterogeneity in age of symptom onset (AAO)[14–16] and rates of cognitive and biomarker change[17–21] across carriers of *PSEN1* pathogenic variants. We hypothesized that this heterogeneity may be strongly tied to underlying mutation-level variation in γ-secretase function.

Intriguingly, prior work suggests some pathogenic *PSEN1* variants may not significantly alter the absolute amounts of Aβ42 production[22], indicating that more thorough investigation is needed to understand how these pathogenic variants lead to ADAD and the mechanisms that cause earlier vs. later disease onset and progression. To this end, findings from our group[23] and others[24] suggest that probing the ratios of short-to-long Aβ peptide production across ADAD pathogenic variants may help us understand the observed heterogeneity in these populations.

Here, we leverage a live-cell-based model system and newly developed

immunoassays to functionally characterize APP processing by γ-secretase across a

large set of *PSEN1* pathogenic variants with AAOs ranging from the 20s to the 70s. We

then assess how this mutation-level characterization of γ-secretase function may

explain heterogeneity in neuroimaging and biofluid biomarkers, as well as clinical

measures among *PSEN1* carriers with available data from the Dominantly Inherited

Alzheimer's Network Observational Study (DIAN-Obs).

METHODS

Functional assays of γ-secretase function

162 unique *PSEN1* variants, including 55 unique variants represented in DIAN-Obs and

data from 107 unique variants previously characterized in Liu et al., 2022,[23] and wild-

type (normal) *PSEN1* was characterized for Aβ peptide production and γ-secretase

function. Using methods detailed in supplemental methods and a prior report using this

model system, each *PSEN1* variant or normal *PSEN1* was sub-cloned and placed into

the PcDNA 3.1 expression vector[23]. Subsequently, each variant was co-transfected with

a plasmid encoding wild-type (normal) human APP-C99 into HEK293T cells genetically

depleted of *PSEN1* and *PSEN2*[23]. Creation of this cell line and transient transfection

procedures are described in supplemental methods. Conditioned media from

transfected HEK cells was harvested and diluted with 1% BSA in wash buffer (TBS

supplemented with 0.05% Tween). Aβ37, 38, 40, 42, and 43 immunoassays were

performed in triplicate and averaged as previously described[23] (see Figure S1 and

supplemental methods). Absolute peptide production levels and relevant ratios were

examined as percent relative to the production seen in the same model system

following transfection with normal (wild-type) *PSEN1.*

Participants

Participants provided written informed consent prior to the completion of any study

procedures, in accordance with Institution Review Board policy at each participating

site. Data were included from 190 *PSEN1* pathogenic variant carriers enrolled in DIAN-

Obs, each carrying one of 55 unique *PSEN1* pathogenic variants characterized via cell-

based methods (see Figure 1A, Table S1, and Table S2). Detailed protocols for DIAN

have previously been published and are described in supplemental methods. For DIAN-

Obs participants, the familial AAO was determined through structured interviews to

determine the age of onset of progressive cognitive decline for the participant's first

degree relative(s)[15].

Clinical evaluators were blind to the mutation status of participants. Participants

underwent comprehensive clinical and cognitive evaluations. Clinical Dementia Rating

(CDR®) Sum of Boxes (SB) was assessed for each participant using structured

interviews, as previously described[25]. Mini-Mental State Examination (MMSE) and

Wechsler Memory Scale-Revised Logical Memory Delayed Recall[26] were selected a

priori as our cognitive measures of interest. All DIAN-Obs participants with available Aβ

positron emission tomography (PET), magnetic resonance imaging (MRI), and at least

one cognitive assessment were included (Figure 1A; DIAN-Obs data freeze version 15;

last data from June 30, 2020; see supplemental methods). A subset of these individuals

also had [^{18}F]Fluorodeoxyglucose (FDG)-PET (n=162) and cerebrospinal fluid (CSF)

(n=157) available for analyses. Longitudinal data were available from 154 DIAN

participants (Table S3). For the 107 *PSEN1* variants studies previously in Liu et al.

(non-overlapping with the 55 variants from DIAN-Obs), AAO was derived from prior

literature, as previously described[23].

Imaging and Fluid Biomarker Analyses

Protocols for Aβ and brain metabolism PET, MRI, and CSF studies are described in

detail in supplemental methods and in previously published work[21,25,27–31]. Cerebral Aβ

load was measured using [^{11}C]Pittsburgh Compound B (PiB) PET, and brain glucose

metabolism was measured with [^{18}F]FDG-PET. Partial volume corrected values were

used in all analyses. FreeSurfer v 5.3 defined regional measures were derived from MRI

data. CSF Aβ42, Aβ40, and phosphorylated-tau181 levels were measured using an

automated immunoassay system (LUMIPULSE G1200, Fujirebio, Malvern, PA).

Phosphorylated-tau 217 (pT217/T217) values were derived from immunoprecipitation-

mass spectrometry (IP-MS) as previously described[27]. See supplemental methods for

additional details.

Statistical Analyses

Analyses were conducted using R (version 4.0.3, R Foundation for Statistical

Computing). Pearson correlations were used to assess associations of Aβ production

and γ-secretase function with AAO. A summary measure of variant-level Aβ production

and γ-secretase function (γ-secretase composite) was developed to reduce the number

of statistical comparisons and facilitate comparisons between mutant and wild-type

PSEN1, as described in the Results.

Linear mixed effect models including fixed effects for age, sex, and γ-secretase

composite score were employed to assess associations between the cell-based γ-

secretase function and multi-modal biomarker and clinical data from DIAN. Similar to

prior work from DIAN, a random effect for family membership was included in each

linear mixed effects model. Separate models were used to assess associations with

each clinical, cognitive, or biomarker outcome measure, focusing on CSF Aβ 42/40,

CSF phosphorylated-tau 181, CSF phosphorylated-tau 217, hippocampal volume, CDR-

SB, MMSE, and logical memory delayed (see supplemental methods). Years of

education was included as an additional fixed effect term in models examining cognitive

and clinical outcomes. Longitudinal models were similarly constructed but additionally

included interactions of each fixed effect with time and terms allowing both a random

intercept and slope. Additional exploratory analyses made use of a tertile split of γ-

secretase composite scores to visualize how clinical, cognitive, and biomarker

trajectories can be understood in the context of γ-secretase function.

RESULTS
184

Cell-based measures of γ-secretase function are associated with age of symptom

onset.

Aβ 37, 38, 40, 42, and 43 production was assessed in 162 unique *PSEN1* variants,

including 55 *PSEN1* variants represented in DIAN (Table S1) and 107

non-overlapping variants previously characterized in Liu et al., 2022[23]. Grouping

together Aβ production for all 162 variants, we observed reductions in Aβ37, 38, and 40

compared to normal levels (median % reduction relative to normal = -56.8, -48.6, and -

22.8%, respectively), suggesting that, as a group, *PSEN1* pathogenic variants decrease

the processive γ-cleavage of Aβ by γ-secretase (γ-processivity), leading to a decrease in

these fully processed (shorter) Aβ peptides. Additionally, relative levels of Aβ42 and 43

across variants were higher in *PSEN1* pathogenic variants compared to normal levels

(median % increase relative to normal= 66.5% and 47.1%; respectively).

We first assessed relationships between the relative concentrations of Aβ

species produced by each variant and AAO in a set of 107 variants not represented in

the DIAN study. To limit comparisons with multiple ratios when assessing relationships

between γ-secretase function and AAO, we leveraged prior literature[1,23,24,32,33]

suggesting the Aβ 42/40 ratio and a measure of the efficiency of successive cleavage

(γ-processivity) of Aβ fragments can be used to summarize γ-secretase function. These

two measures were operationalized as $\log_{10}(\frac{A\beta42}{A\beta40})$ (a measure of the relative production

of aggregation-prone Aβ) and a ratio of short (fully γ-secretase processed) to long

(incompletely γ-secretase processed) Aβ fragments ($\frac{\text{short } A\beta37+38}{\text{long } A\beta42+43}$). As expected, $\frac{A\beta42}{A\beta40}$,

and $\frac{\text{short } A\beta37+38}{\text{long } A\beta42+43}$ ratio levels across *PSEN1* pathogenic variants were both abnormal

(decreased processivity) compared to normal levels (median % change relative to the

normal = 108.9 and -71.3%). In the initial sample of 107 *PSEN1* pathogenic variants

from Liu et al., both higher $\log_{10}(\frac{A\beta42}{A\beta40})$ (r[105] = -0.64, p = 2.20e-13) and lower

$\frac{short\ A\beta 37+38}{long\ A\beta 42+43}$ (r[105] = 0.66, p =2.28e-14) levels were associated with earlier AAO (Figure

1B-C). Both ratios remained significantly associated when entered together into a single

model predicting AAO (Table S4), suggesting that they capture unique variance with

respect to the pathogenicity of each variant. Accordingly, these two ratios were

combined into a single summary measure of variant-level γ-secretase function ($\gamma -$

secretase $Composite = \frac{short\ A\beta 37+38}{long\ A\beta 42+43} - log_{10}\frac{A\beta 42}{A\beta 40}$). To facilitate visualization and

comparison, the γ-secretase composite is expressed as a percentage of wild-type γ-

secretase function.

219   In both the subset of variants represented in the DIAN-Obs and remaining

variants characterized previously by Liu et al[23], γ-secretase composite scores were

lower in all pathogenic variants compared to normal (average of 60.4% reduction

relative to normal). The γ-secretase composite had a the highest degree of correlation

with AAO (Liu et al cohort: r[105] = 0.71, p <2.2e-16; DIAN-Obs cohort: r[53] = 0.61, p =

6.10e-07; Figure 1E), as compared to using individual Aβ peptide levels, $\frac{A\beta 42}{A\beta 40}$ levels, or

$\frac{short\ A\beta 37+38}{long\ A\beta 42+43}$ ratio levels in the initial set of 107 *PSEN1* variants and in the 55 variants

represented in DIAN (see above). To demonstrate the potential utility of using the γ-

secretase composite to predict an approximate AAO for novel pathogenic variants

and/or when a clinical history is not available, we list the γ-secretase composite derived

AAO (AAO$_{gsc}$) values for the 55 DIAN-Obs variants examined here (Table S1). Notably,

γ-secretase composite values did not map in a straightforward manner onto the location

of the corresponding pathogenic variant within *PSEN1*, and substantial variation within a

single codon in terms of gamma-secretase function was observed for several sites in

which more than one variant was characterized (Table S1; Figure 1D).

Lower cell-based γ-secretase composite scores are associated with more

abnormal AD imaging and fluid biomarkers.

Lower γ-secretase composite scores (decreased γ-secretase function relative to

normal) were associated with higher levels of mean cortical Aβ burden (B[SE] = -

0.03[0.01], p =4.06e-07; Figure 2A) as assessed by Aβ PET, after controlling for

demographic factors. Exploratory regional analyses revealed widespread associations

between *in vivo* Aβ burden and the cell-based γ-secretase composite, with the strongest

associations seen in the precuneus, and anterior and posterior cingulate (Figure 2D).

We next examined associations between the variant-level γ-secretase composite

and commonly used imaging biomarkers of neurodegeneration, including structural MRI

and FDG-PET, in DIAN-Obs participants carrying one of the 55 of the *PSEN1* variants

characterized *in vitro*. Both the MRI-based measures hippocampal volume (B[SE] =

44.22[6.4], p = 5.15e-10; Figure 2B) and precuneus FDG-PET signal (B[SE] =

0.006[0.001], p = 5.38e-05; Figure 2C) were highly associated with the γ-secretase

composite, after controlling for demographic factors. Exploratory regional analyses

revealed that the cell-based γ-secretase composite significantly accounted for

widespread grey matter volume differences across *PSEN1* variant carriers, most

notably in the hippocampus, amygdala, temporal regions, precuneus, and posterior

cingulate (Figure 2E), as well as differences in glucose metabolism including the

precuneus, parietal, and frontal regions (Figure 2F).

Using established CSF measures of AD pathology, we observed that lower

variant-specific γ-secretase composite scores were associated with lower CSF Aβ 42/40

(B[SE] = 6.3e-04[1.5e-04], p = 4.46e-05; Figure 3A), higher CSF ELISA

$\log_{10}$(phosphorylated-tauT181) levels (B[SE] = -0.009[0.002], p =4.47e-06; Figure 3B),

and higher CSF IP-MS $\log_{10}$(pT217/T217) levels (B[SE] = -0.01[0.002], p =1.98e-05;

Figure 3C).

γ-secretase composite scores are associated with clinical and cognitive measures

in *PSEN1* pathogenic variant carriers in DIAN-Obs.

The cell-derived γ-secretase composite levels predicted changes of MMSE

(B[SE] = 0.11[.03], p = 2.17e-04; Figure 3D), CDR-SB (B[SE] = -0.07[0.02], p = 4.86e-

05; Figure 3E), and logical memory-delayed scores (B[SE] = 0.11[.03], p = 6.51e-05;

Figure 3F).

γ-secretase composite scores predict rate of change in biomarker, clinical and

cognitive features in *PSEN1* ADAD carriers.

We next assessed how longitudinal rates of change in core biomarker, clinical,

and cognitive measures of interest (Figure 4A, C, E, and G) may be related to cell-

based measures of γ-secretase function. Lower (more pathogenic) γ-secretase

composite measures were associated with faster increase in β-amyloid PET signal

(B[SE] = -7.2e-04[2e-04], p = 0.003; Figure 4B), as well as more rapid decreases in

hippocampal volume (B[SE] = 4.56[0.7], p = 2.96e-10; Figure 4D), MMSE (B[SE] =

0.02[0.007], p = 3.02e-04; Figure 4F), and logical memory-delayed (B[SE] =

0.006[0.001], p = 2.63e-06; Figure 4H).

Multimodal models suggest variant level variations in γ-secretase function may

impact the clinical, cognitive, and biomarker progression of ADAD.

To assess how differences in γ-secretase function may alter the observed

clinical, cognitive, and biomarker course of ADAD, we performed a tertile split of

variants represented in DIAN and plotted the standardized change in core clinical,

cognitive, and biomarker measures across the lifespan for individuals carrying *PSEN1*

pathogenic mutations in the lowest, middle, or highest tertile of γ-secretase function

(Figure 5A-C). To complement this group-level analysis, we used all available data to

visualize putative disease trajectories for carriers of 2 hypothetical variants with 25%

and 75% of normal γ-secretase function, respectively (Figure 5D-I). Consistent with the

foregoing analyses, these visualizations demonstrate an earlier shift in AAO, clinical and

biomarker abnormalities in carriers of variants with lower (more pathogenic) γ-secretase

composite scores. A similar pattern was seen when using estimated years to symptom

onset (EYO; calculated by subtracting history-derived AAO from the participant's age)

rather than age (Figure S2A-B). This suggests that cell-based assessments of γ-

secretase add to the prediction of disease trajectories even after history-derived

information about AAO is included in statistical models. However, while a shift in the

timing of symptom and biomarker changes was seen across a range of γ-secretase

function, it is notable that elevations in pathological Aβ burden (higher Aβ PET or lower

CSF Aβ42) and tau (higher CSF phosphorylated-tau181) appear to be early events

across all tertiles and precede changes in neurodegenerative markers (hippocampal

volume, FDG PET) and clinical measures of impairment (CDR-SB). Together, these

results suggest that, while mutation-level variations in γ-secretase function may impact

the age of symptom onset and rate of ADAD progression, the ordering of clinical,

cognitive, and biomarker changes is not itself fundamentally altered by differences in γ-

secretase function.

DISCUSSION:

The study of ADAD forms a cornerstone for understanding AD pathobiology, informing

the development of new AD biomarkers, broadly influencing the design of animal

models of AD, and advancing therapeutic approaches for both LOAD and ADAD. While

ADAD-causing *PSEN1* pathogenic variants are highly penetrant, remarkable

heterogeneity in the AAO, rates of disease progression, and biomarker changes is seen

across variants. Here we examined whether variant level differences in γ-secretase

processing of APP may be critical determinants of the age at which *PSEN1* pathogenic

variant carriers manifest cognitive changes and develop abnormalities in core

biomarkers for AD. Using a cell-based model system to assess the production of a

broad set of Aβ peptides, we observed that mutation level differences in γ-secretase

processing of APP strongly predicted the age at which progressive cognitive symptoms

are manifest (AAO) for a given variant. In addition to predicting AAO, we observed that

a summary measure of γ-secretase function was associated with the rates of functional

(CDR-SB) and cognitive decline. This composite measure of γ-secretase function was

also associated with all fluid and imaging biomarker measures of AD pathobiology and

neurodegeneration we examined, including CSF measures of AD pathology, Aβ PET,

FDG-PET, and MRI gray matter volumes. These results suggest that accounting for

mutation-level γ-secretase dysfunction may clarify treatment effects in ADAD clinical

trials and may also be useful in predicting approximate AAO and disease trajectories for

PSEN1 pathogenic variants that are novel or lack sufficient clinical history.

Mechanistically, these results highlight the direct importance of γ-secretase function to

AD pathobiology and provide support for γ-secretase modulation as a potentially

powerful AD therapeutic and preventative strategy[34].

Using a human cell-based model system, we found substantial variability in the

production of shorter (Aβ-37 and 38) to longer (Aβ-42 and 43) Aβ peptides across a

large number of *PSEN1* pathogenic variants. In this same system, we observed that the

Aβ42/40 ratio was increased in a majority but not all variants examined. We first

examined 107 *PSEN1* variants with available AAO but no available clinical and

biomarker data and observed that both the ratio of short-to-long Aβ peptide production

$(\frac{A\beta37+38}{A\beta42+43})$ and the more commonly used Aβ 42/40 ratio were predictive of AAO.

Intriguingly, each of these ratios was independently associated with AAO, and a

composite measure derived from these two ratios provided stronger associations with

AAO than either alone. This composite measure was carried forward to examine AD

progression in a non-overlapping set of *PSEN1* carriers with available clinical and

biomarker measures of AD progression.

We found that the γ-secretase composite measure explained substantial

biomarker heterogeneity across *PSEN1* pathogenic mutation carriers using available

data from DIAN. Focusing on associations between the γ-secretase composite and Aβ

PET, we observed that individuals who carried pathogenic variants in the lowest (most

pathogenic) tertile had, on average, 1.4 SUVR units greater Aβ PET signal compared to

those with variants in the highest (least pathogenic) tertile, after accounting for age and

sex. Putting these differences in the context of the conventionally used threshold for Aβ

PET positivity in this cohort and with this Aβ PET tracer (cortical PiB PET SUVR positive

at >= 1.42 SUVR[28]), carriers of pathogenic variants in the lowest (most pathogenic)

tertile of the γ-secretase composite would, on average, become amyloid positive ~15

years sooner than those carrying *PSEN1* variants in the highest (least pathogenic)

tertile (estimated mean age of 22.6 and 37.2 years for low and high tertiles,

respectively). Consistent with cross-sectional results, analysis of longitudinal data

demonstrated that Aβ PET signal increased more quickly in individuals carrying *PSEN1*

variants with lower (more pathogenic) γ-secretase composite scores compared to those

with higher composite scores. Though further work is needed to elucidate regional

differences, the results here also suggest that variants with low γ-secretase composite

values may have greater striatal Aβ PET signal compared to carriers of *PSEN1* variants

with high composite values.

Cerebrospinal fluid measures of AD pathology also had significant associations

with the variant-level γ-secretase composite, including CSF measures of Aβ42/40 and

phosphorylated-tau181, and phosphorylated-tau217. In addition to associations with

core measures of AD pathobiology, γ-secretase composite scores correlated with FDG

PET and structural MRI, two commonly used neuroimaging measures of

neurodegeneration. Together with the described associations with AAO and functional

impairment measures, this broad pattern of association between γ-secretase function

with core AD and neurodegenerative markers further support the hypothesis that

heterogeneity in ADAD can broadly be understood in the context of mutation-level

differences in γ-secretase dysfunction. Importantly, despite variations in AAO and

biomarker trajectories across variants, the results here also support prior observations

that pathologic changes in Aβ and tau consistently precede neurodegenerative changes

and clinical impairment across *PSEN1* variants[23,24,35]. Additionally, as the γ-secretase

composite used here was derived to predict AAO, it remains quite possible that different

formulations of this summary measure of γ-secretase function will be better optimized to

predict particular biomarker trajectories in clinical trial settings.

The results here need to be considered in the context of certain limitations.

Though AAO information was available on all *PSEN1* variants examined, only a subset

had available clinical, cognitive, and biomarker data from the DIAN-Obs study.

Accordingly, data from new pathogenic variants in DIAN and in the broader scientific

literature will need to be considered. Similarly, tau PET data will need to be considered

in future analyses as they become available. Further, it will be important to determine if

processing of γ-secretase substrates besides APP is additionally informative for

understanding the differential pathogenicity across *PSEN1* variants. Lastly, it will be

critical to extend the cell-based model system to allow for the characterization of

pathogenic variants in *PSEN2* and *APP*, as ~20% of ADAD-causing variants are found

within these genes.

Despite these limitations, these findings indicate that variant level differences in

γ-secretase function are highly associated with core measures of clinical, biomarker,

and cognitive, progression of ADAD, and suggest that these variant-level differences

explain a large portion of the heterogeneity seen across *PSEN1* pathogenic mutation

carriers. As the biochemical measures of γ-secretase function used here are derived

outside the context of family polygenic factors, age, and estimated years to symptom

onset, they offer a potentially unique channel of information with respect to disease

progression compared to these other factors and may thus represent useful

complements to these measures in ADAD clinical trials. Similarly, as these cell-based

measures are divorced from family polygenic factors and age, they may prove to be

valuable in gauging the impact of genetic and lifestyle factors that may confer resilience

or risk for ADAD onset and progression. Cell-based measures of γ-secretase function

may be particularly useful to assess pathogenicity, potential AAO, and predicted AD

trajectories for newly discovered *PSEN1* pathogenic variants or for variants lacking

sufficient clinical history. Importantly, these results also strongly support the continued

investigation of γ-secretase modulators as an AD therapeutic strategy. Lastly, by

demonstrating a tight linkage between the processing of APP by γ-secretase in a cell-

based model system and ADAD disease trajectories *in vivo*, the results here provide

strong support for the hypothesis that dysregulation of APP processing is central to AD

pathogenesis, offing support for some formulations of the "Aβ hypothesis" of AD

pathogenesis.

AUTHORS' CONFLICT OF INTERESTS

Partial funding from the Alzheimer's Association (SAS), NIH grants (CRJ, GSD, JMN,

JCM, RJB, TLSB, JPC), NIA grants (GSD), GHR Foundation (CRJ), Alexander Family

Alzheimer's Disease Research Professorship of Mayo Clinic (CRJ), grants from Lilly

(NGR), Biogen (NGR), AbbVie (NGR, JL), Chan-Zuckerberg Initiative (GSD), DZNE (JL), Eisai (CR), Instituto de Salud Carlos III (RSV), Anonymous Foundation (PRS).

Consultations have been declared for Humana Healthcare (JPC), Roche (NCF, JL, CR, RSV, RJB), Biogen (NCF, JL, CR), Parabon Nanolabs Inc. (GSD), DynaMed (EBSCO Health) (GSD), medical testimony on legal cases pertaining to management of Wernicke Encephalopathy (GSD), MODAG (JL), GmbH (JL), Bayer Vital (JL), Axon Neuroscience (JL), Thieme Medical Publishers (JL), Prothena (CR), Wave Pharmaceuticals (RSV), Janssen (RSV), Neuraxpharm (RSV), Eisai (RJB), and Amgen (RJB). Other financial interests include owning stocks greater than $10000 are held in ANI Pharmaceuticals (GSD), being a member of the DIAN-TU Pharma Consortium which includes funding and non-financial support for the DIAN-TU-001 trial from Avid Radiopharmaceuticals, Janssen Hoffman Roche/Genentech, Eli Lilly & Co, Eisai, Biogen, AbbVie, and Bristol Meyer Squibbs (RJB, TLSB), and royalties from C2N Diagnostics with equity ownership interest (RJB).

Non-financial interests include serving on an independent data monitoring board for Roche (CRJ), support from Lilly (NCF), support from Ionis (NCF), serving as director of Anti- NMDA Receptor Encephalitis Foundation (GSD), Eisai (TLSB), Siemens (TLSB), and a member of speaker's bureau for Biogen (TLSB).

All other authors have nothing to disclose.

ACKNOWLEDGMENTS:

Data collection and sharing for this project was supported by the Dominantly Inherited

Alzheimer Network (DIAN, U19AG032438) funded by the National Institute on Aging

(NIA), the Alzheimer's Association (SG-20-690363-DIAN), the German Center for

Neurodegenerative Diseases (DZNE), Raul Carrea Institute for Neurological Research

(FLENI). Partial support by the Research and Development Grants for Dementia from

Japan Agency for Medical Research and Development, AMED, and the Korea Health

Technology R&D Project through Korea Health Industry Development Institute (KHIDI),

Spanish Institute of Health Carlos III (ISCIII), Canadian Institutes of Health Research

(CIHR), Canadian Consortium of Neurodegeneration and Aging, Brain Canada

Foundation and Fonds de Recherche du Québec – Santé. This manuscript has been

reviewed by DIAN Study investigators for scientific content and consistency of data

interpretation with previous DIAN Study publications. We acknowledge the altruism of

the participants and their families, and contributions of the DIAN research and support

staff at each of the participating sites for their contributions to this study.

REFERENCES:

1. Lagomarsino VN, Pearse RV, Liu L, et al. Stem cell-derived neurons reflect features of protein networks, neuropathology, and cognitive outcome of their aged human donors. *Neuron*. 2021;109(21):3402-3420.e9. doi:10.1016/j.neuron.2021.08.003

2. Hsu S, Gordon BA, Hornbeck R, et al. Discovery and validation of autosomal dominant Alzheimer's disease mutations. *Alz Res Therapy*. 2018;10(1):67. doi:10.1186/s13195-018-0392-9

3. Karch CM, Hernández D, Wang JC, et al. Human fibroblast and stem cell resource from the Dominantly Inherited Alzheimer Network. *Alzheimers Res Ther*. 2018;10(1):69. doi:10.1186/s13195-018-0400-0

4. Lanoiselée HM, Nicolas G, Wallon D, et al. APP, PSEN1, and PSEN2 mutations in early-onset Alzheimer disease: A genetic screening study of familial and sporadic cases. *PLoS Med*. 2017;14(3):e1002270. doi:10.1371/journal.pmed.1002270

5. De Strooper B, Iwatsubo T, Wolfe MS. Presenilins and γ-secretase: structure, function, and role in Alzheimer Disease. *Cold Spring Harb Perspect Med*. 2012;2(1):a006304. doi:10.1101/cshperspect.a006304

6. Fernandez MA, Klutkowski JA, Freret T, Wolfe MS. Alzheimer presenilin-1 mutations dramatically reduce trimming of long amyloid β-peptides (Aβ) by γ-secretase to increase 42-to-40-residue Aβ. *J Biol Chem*. 2014;289(45):31043-31052. doi:10.1074/jbc.M114.581165

7. Liu L, Lauro BM, Wolfe MS, Selkoe DJ. Hydrophilic loop 1 of Presenilin-1 and the APP GxxxG transmembrane motif regulate γ-secretase function in generating Alzheimer-causing Aβ peptides. *Journal of Biological Chemistry*. 2021;296:100393. doi:10.1016/j.jbc.2021.100393

8. Wolfe MS. Dysfunctional γ-Secretase in Familial Alzheimer's Disease. *Neurochem Res*. 2019;44(1):5-11. doi:10.1007/s11064-018-2511-1

9. Wolfe MS, Selkoe DJ. Giving Alzheimer's the old one-two. *Cell*. 2010;142(2):194-196. doi:10.1016/j.cell.2010.07.006

10. Wolfe MS. Probing Mechanisms and Therapeutic Potential of γ-Secretase in Alzheimer's Disease. *Molecules*. 2021;26(2):E388. doi:10.3390/molecules26020388

11. Nie P, Vartak A, Li YM. γ-Secretase inhibitors and modulators: Mechanistic insights into the function and regulation of γ-Secretase. *Semin Cell Dev Biol*. 2020;105:43-53. doi:10.1016/j.semcdb.2020.03.002

12. Raven F, Ward JF, Zoltowska KM, et al. Soluble Gamma-secretase Modulators Attenuate Alzheimer's β-amyloid Pathology and Induce Conformational Changes in Presenilin 1. *EBioMedicine*. 2017;24:93-101. doi:10.1016/j.ebiom.2017.08.028

13. *Psen-1| Alzforum. (n.d.). Retrieved August 09, 2021, from*
Https://Www.Alzforum.Org/Mutations/Psen-1.

14. Ryan NS, Nicholas JM, Weston PSJ, et al. Clinical phenotype and genetic
associations in autosomal dominant familial Alzheimer's disease: a case series. *The*
Lancet Neurology. 2016;15(13):1326-1335. doi:10.1016/S1474-4422(16)30193-4

15. Ryman DC, Acosta-Baena N, Aisen PS, et al. Symptom onset in autosomal
dominant Alzheimer disease: a systematic review and meta-analysis. *Neurology*.
2014;83(3):253-260.

16. Lippa CF, Swearer JM, Kane KJ, et al. Familial Alzheimer's disease: site of mutation
influences clinical phenotype. *Ann Neurol*. 2000;48(3):376-379.

17. Wegiel J, Wisniewski HM, Kuchna I, et al. Cell-Type-Specific Enhancement of
Amyloid-β Deposition in a Novel Presenilin-1 Mutation (P117L): *Journal of*
Neuropathology and Experimental Neurology. 1998;57(9):831-838.
doi:10.1097/00005072-199809000-00004

18. Klunk WE, Price JC, Mathis CA, et al. Amyloid Deposition Begins in the Striatum of
Presenilin-1 Mutation Carriers from Two Unrelated Pedigrees. *Journal of*
Neuroscience. 2007;27(23):6174-6184. doi:10.1523/JNEUROSCI.0730-07.2007

19. Tang M, Ryman DC, McDade E, et al. Neurological manifestations of autosomal
dominant familial Alzheimer's disease: a comparison of the published literature with
the Dominantly Inherited Alzheimer Network observational study (DIAN-OBS). *The*
Lancet Neurology. 2016;15(13):1317-1325.

20. Ryan NS, Rossor MN. Correlating familial Alzheimer's disease gene mutations with
clinical phenotype. *Biomarkers Med*. 2010;4(1):99-112. doi:10.2217/bmm.09.92

21. Chhatwal JP, Schultz SA, McDade E, et al. Variant-dependent heterogeneity in
amyloid β burden in autosomal dominant Alzheimer's disease: cross-sectional and
longitudinal analyses of an observational study. *Lancet Neurol*. 2022;21(2):140-152.
doi:10.1016/S1474-4422(21)00375-6

22. Kumar-Singh S, Theuns J, Van Broeck B, et al. Mean age-of-onset of familial
alzheimer disease caused by presenilin mutations correlates with both increased
Aβ42 and decreased Aβ40. *Hum Mutat*. 2006;27(7):686-695.
doi:10.1002/humu.20336

23. Liu L, Lauro BM, He A, et al. Identification of the Aβ37/42 peptide ratio in CSF as an
improved Aβ biomarker for Alzheimer's disease. *Alzheimers Dement*. Published
online March 12, 2022. doi:10.1002/alz.12646

24. Petit D, Fernández SG, Zoltowska KM, et al. Aβ profiles generated by Alzheimer's
disease causing PSEN1 variants determine the pathogenicity of the mutation and

predict age at disease onset. *Mol Psychiatry*. 2022;27(6):2821-2832.
doi:10.1038/s41380-022-01518-6

25. Bateman RJ, Xiong C, Benzinger TLS, et al. Clinical and Biomarker Changes in
Dominantly Inherited Alzheimer's Disease. *New England Journal of Medicine*.
2012;367(9):795-804. doi:10.1056/NEJMoa1202753

26. Wechsler D. Wechsler Memory Scale (3rd ed.): Administration and scoring manual.
Published online 1997.

27. Barthélemy NR, Li Y, Joseph-Mathurin N, et al. A soluble phosphorylated tau
signature links tau, amyloid and the evolution of stages of dominantly inherited
Alzheimer's disease. *Nat Med*. 2020;26(3):398-407. doi:10.1038/s41591-020-0781-z

28. Su Y, D'Angelo GM, Vlassenko AG, et al. Quantitative Analysis of PiB-PET with
FreeSurfer ROIs. *PLOS ONE*. 2013;8(11):e73377.
doi:10.1371/journal.pone.0073377

29. Su Y, Blazey TM, Owen CJ, et al. Quantitative Amyloid Imaging in Autosomal
Dominant Alzheimer's Disease: Results from the DIAN Study Group. *PLoS One*.
2016;11(3):e0152082. doi:10.1371/journal.pone.0152082

30. Gordon BA, Blazey TM, Su Y, et al. Spatial patterns of neuroimaging biomarker
change in individuals from families with autosomal dominant Alzheimer's disease: a
longitudinal study. *The Lancet Neurology*. 2018;17(3):241-250.

31. McKay NS, Gordon BA, Hornbeck RC, et al. *Neuroimaging within the Dominantly
Inherited Alzheimer's Network (DIAN): PET and MRI*. Neuroscience; 2022.
doi:10.1101/2022.03.25.485799

32. Gu L, Guo Z. Alzheimer's Aβ42 and Aβ40 peptides form interlaced amyloid fibrils. *J
Neurochem*. 2013;126(3):305-311. doi:10.1111/jnc.12202

33. Cullen N, Janelidze S, Palmqvist S, et al. Association of CSF Aβ38 Levels With Risk
of Alzheimer Disease-Related Decline. *Neurology*. 2022;98(9):e958-e967.
doi:10.1212/WNL.0000000000013228

34. Toyn JH, Boy KM, Raybon J, et al. Robust Translation of -Secretase Modulator
Pharmacology across Preclinical Species and Human Subjects. *Journal of
Pharmacology and Experimental Therapeutics*. 2016;358(1):125-137.
doi:10.1124/jpet.116.232249

35. Sun L, Zhou R, Yang G, Shi Y. Analysis of 138 pathogenic mutations in presenilin-1
on the in vitro production of Aβ42 and Aβ40 peptides by γ-secretase. *Proc Natl Acad
Sci U S A*. 2017;114(4):E476-E485. doi:10.1073/pnas.1618657114

Figure 1. Cell-based measures of γ-secretase function are tightly associated with age of symptom onset across two cohorts representing 162 unique *PSEN1* variants. (A) Schematic summarizing the characterization of variant-level γ-secretase function and correlations with age of symptom onset, cognitive, and biomarker data for each variant (created with BioRender.com). γ-secretase function was summarized using two ratios of Aβ peptide production derived from cell models **(B)** Cell-based $\log_{10}(\frac{A\beta42}{A\beta40})$ levels and **(C)** a ratio of short (fully γ-secretase processed) to long (incompletely γ-secretase processed) Aβ fragments $(\frac{\text{short } A\beta37+38}{\text{long } A\beta42+43})$. Both ratios were independently associated with age of symptom onset in an initial set of 107 *PSEN1* variants (B-C). To reduce comparisons going forward, both ratios were combined into a single composite

571 measure (γ-secretase composite, expressed as a percentage of normal γ-secretase
572 function; D). This composite measure correlated with age of symptom onset in both the
573 initial set of 107 *PSEN1* variants and in a second set of 55 *PSEN1* variants represented
574 in DIAN-Obs (E). Individual data points are jittered to maintain blinding.
575

Figure 2. Lower γ-secretase function across variants is associated with more

abnormal AD imaging biomarkers. Association between variant-level γ-secretase

function and in *vivo* imaging biomarker data from carriers of corresponding *PSEN1*

pathogenic variants is shown for cross-sectional Aβ PET (age-adjusted PiB-PET SUVR

cortical composite, **A**), age- and intracranial volume-adjusted bilateral hippocampal grey

matter volume derived from structural MRI (**B**), and age-adjusted bilateral precuneus

FDG-PET SUVR (**C**) from 190 *PSEN1* pathogenic variant carriers (55 unique variants

represented). Individual data points are jittered to maintain blinding of the data.

Exploratory, regional analyses comparing PiB-PET SUVRs (**D**), grey matter volumes

(**E**), and FDG-PET SUVRs (**F**) across a range of γ-secretase function, with colors

indicating t- and p-values derived from the comparison of neuroimaging data from

PSEN1 carriers with variants in the lowest (most pathogenic) vs. highest (least

pathogenic) tertiles of γ-secretase composite function. Models additionally adjusted for

age and sex. Individual data points are jittered to maintain blinding.

591
592

Figure 3. Lower γ-secretase function across variants is associated with more abstract biofluid, clinical, and cognitive measures in ADAD. Associations between γ-secretase function for each *PSEN1* variant and cross-sectional CSF Aβ 42/40 (**A**), age-adjusted CSF $\log_{10}$ pT181 (**B**), CSF $\log_{10}$ pT217/T217 (**C**), age-adjusted MMSE (**D**), age-adjusted CDR-SB (**E**), and age-adjusted logical memory-delayed (**F**) values from 190 *PSEN1* pathogenic variant carriers (55 unique variants represented) are shown. Individual data points are jittered to maintain blinding.

CSF = Cerebrospinal Fluid; pT = phosphorylated tau; MMSE = Mini Mental State Examination scores; CDR-SB = Clinical Dementia Rating Scale- Sum of Boxes.

Figure 4. Lower γ-secretase function across *PSEN1* variants is associated with

more rapid worsening in biomarker, clinical and cognitive measures in ADAD.

Individual longitudinal measures are plotted for PiB-PET SUVR (Aβ burden; **A**),

hippocampal volume (HV; mm^3; **C**), Mini Mental Status Exam (MMSE; **E**), and logical

memory delayed (Log. Mem.; **G**). Colors indicate whether an individual's *PSEN1* variant

is in the low (red; most pathogenic tertile), middle (black; intermediate pathogenicity), or

high (blue; least pathogenic) tertile for γ-secretase function. To facilitate statistical

comparison, age-adjusted, individual-level slopes for each biomarker and cognitive

outcomes of interest were extracted from linear mixed effects models and plotted

against γ-secretase function (Aβ PET – **B**; Hippocampal Volume – **D**; MMSE – **F**;

Logical Memory – **H**). Significant associations were observed between rate-of-change

in each measure and variant-level γ-secretase function. Individual data points are

jittered to maintain blinding.

Figure 5. Variant-level variations in γ-secretase function are broadly associated

with the clinical and biomarker course of ADAD and can be used to predict ADAD

progression for novel *PSEN1* pathogenic variants. To visualize how variant-level

differences in γ-secretase function may broadly alter the clinical, cognitive, and

biomarker course of ADAD, we plotted the standardized differences in core clinical,

cognitive, and biomarker measures across the lifespan for DIAN-Obs participants

carrying *PSEN1* pathogenic variants in the lowest (**A**), middle (**B**), or highest (**C**) tertile

of γ-secretase function compared to non-carriers. This group-level analysis was followed

627   by an individual-level visualization of disease trajectories for pathogenic variant carriers

628   of 2 possible variants with 25% and 75% of normal γ-secretase function, respectively

629   (D-I).