**A cerebellar population coding model for sensorimotor learning**

Tianhe Wang[*], Richard B. Ivry

Department of Psychology and Helen Wills Neuroscience Institute, University of California, Berkeley, California

**Corresponding authors (*):**

Tianhe Wang (tianhewang@berkeley.edu)

**1 Abstract**

The cerebellum is crucial for sensorimotor adaptation, using error information to keep the sensorimotor

system well-calibrated. Here we introduce a population-coding model to explain how cerebellar-

dependent learning is modulated by contextual variation. The model consists of a two-layer network,

designed to capture activity in both the cerebellar cortex and deep cerebellar nuclei. A core feature of the

model is that within each layer, the processing units are tuned to both movement direction and the

direction of movement error. The model captures a large range of contextual effects including

interference from prior learning and the influence of error uncertainty and volatility. While these effects

have traditionally been taken to indicate meta learning or context-dependent memory within the

adaptation system, our results show that they are emergent properties that arise from the population

dynamics within the cerebellum. Our results provide a novel framework to understand how the nervous

system responds to variable environments.

**Introduction**

Humans are incredibly flexible in how we adapt our motor behavior across variable environments. We readily compensate for the added weight of a heavy winter coat when reaching for an object or adjust the force required as we sip on our morning coffee. The cerebellum is recognized as playing a key role in this adaptation process[1,2]. Impaired adaptation is one of the hallmarks of cerebellar pathology, observed across a range of tasks from experimentally-induced lesions in animal models[3–5] or neurological disorders in humans[6,7]. Moreover, anatomical and physiological studies have led to computational models in which the cerebellum uses error information to improve subsequent, similar movements[8,9]. This form of learning operates implicitly, automatically recalibrating the sensorimotor system without the need for awareness or drawing on cognitive resources[7,10,11]. The current paper aims to understand the neural computations that support flexible adaptation and how this recalibration process is modified by context and environmental uncertainty.

Previous research has suggested that cerebellum-dependent learning is cognitively impenetrable, responding to error in a "rigid" manner even when the correction fails to improve task performance[10,12–15]. Moreover, unlike many learning processes, adaptation is not sensitive to the statistical properties of the perturbations[16,17]. However, this view of a rigid, inflexible system has been challenged by recent evidence showing that implicit adaptation is modulated by experience[18]. For instance, when participants are exposed to a previously experienced perturbation, the rate of relearning is slower than had been originally observed[18]. Not only does this result suggest a degree of flexibility in adaptation, but, interestingly, this context effect is opposite what is typically observed in studies of relearning: Across a broad range of task domains (e.g., reward-based learning, language acquisition), relearning is typically faster, a phenomenon known as savings[19–21].

The rigidity and atypical effect of experience point to the need for considering the unique properties of

the cerebellum to understand how adaptation is modulated by context and environmental variability. To

this end, we develop a novel computational model of the cerebellum. This model incorporates two core

observations from cerebellar physiology. First, recent studies of oculomotor control have revealed a

fundamental tuning property of Purkinje cells, the primary integrative unit in the cerebellar cortex: These

cells are not only tuned to movement direction but also to the direction of error relative to that

movement[22–24]. Second, the model includes a two layer network, with the second layer designed to

capture activity in the deep cerebellar nuclei[25–28]. We posit that units in the DCN exhibit similar tuning

properties as Purkinje cells and that units with similar tuning profiles are linked across these two layers.

We used the model to generate predictions regarding a range of contextual manipulations and evaluated

the predictions with a series of behavioral experiments. Specifically, we systematically examined the

effect of past experience, error uncertainty, and variation in temporal dynamics in evaluating our model.

Where relevant, we consider several alternative models that have been proposed to elucidate how

context and environmental uncertainty modulate sensorimotor learning[29–31]. Our population-coding

model provides an excellent fit of the behavioral results, even without positing the direct representation

of context or latent state variables or having the capability to modulate learning parameters through meta

learning. As such, our model provides a comprehensive explanation of core computations that account

for how the cerebellum can keep the sensorimotor system well-calibrated across variable environments.

**Results**

<u>Cerebellar Population Coding (CPC) Model</u>

The basic principles of cerebellar-dependent error-based learning are articulated in the classic Marr-Albus

model[2,32]. Purkinje cells (PC) in the cerebellar cortex receive two types of input (Fig 1a). One source

originates in the pontine nuclei. This pathway is hypothesized to provides contextual information,

including an efference copy of the motor command. PCs operate as an internal model, utilizing the input

to predict the consequences of the motor command[33,34]. The second source originates in the inferior olive

with activation of the climbing fibers indicating a mismatch between the predicted and expected sensory

feedback, a teaching signal that is used to update the internal model.

Here we extend the model to provide a general explanation of how learning is modulated by contextual

variability. A foundational idea for our model is inspired by a recent work showing how PCs in the

oculomotor cerebellar cortex are simultaneously tuned to both movement direction and the direction of

a visual error that arises during that movement[22–24]. Tuning in terms of movement direction is reflected

in the simple spike activity of the Purkinje cells and tuning in terms of movement error is reflected in the

complex spike activity of these cells (Fig 1b). The latter one induces long-term depression (LTD) of parallel

fiber-PC (PF-PC) synapses, reducing the future efficacy of similar input on PC activity (Fig 1c-d). Importantly,

because the two tuning profiles are in opposite directions[22–24], error-related activation reduces the simple

spike activity, resulting in a change that can reduce the error in the future movement (Fig 1e-f).

A second prominent feature of our model is that plasticity occurs within the cerebellar cortex and deep

cerebellar nuclei (DCN)[25,35]. Lesion studies of eyeblink conditioning provide one line of evidence indicating

that some aspect of consolidated learning is centered in the DCN. Ablation of the cerebellar cortex can

completely block *de novo* cerebellar-dependent learning[26,36]. However, once the learned behavior is

established, it can persist after lesions to the cerebellar cortex even though the kinematics are

disrupted[37,38]. This dissociation can be explained by the dual-effect of pontine projections to the

cerebellum[39]: a polysynaptic projection through the granule layer to PC, and a direct excitatory projection

of the mossy fibers to the DCN. We assume that PC and DCN neurons are connected such that they share

the same tuning direction for movement[40], and that learning in the DCN is gated by learning at the

cerebellar cortex (Fig 1a). Specifically, LTD at parallel fiber-PC (PF-PC) synapses will reduce inhibitory PC

input to the DCN, facilitating the emergence of long term potentiation (LTP) at mossy fiber-DCN synapses

(Fig 1e)[41].

In summary, an error signal will decrease the efficacy of parallel fiber input to PCs and increase the efficacy

of mossy fiber input to the DCN (Fig 1d, e). Correspondingly, the net output of the cerebellum will provide

a signal of the required change in movement direction to correct for the error (Fig 1f).

**Fig. 1 Illustration of the CPC model. a)** Structure of the cerebellar circuit incorporated in the CPC model.

**b)** Each Gaussian-shaped curve represents the tuning function of a single Purkinje cell (PC) based on that

cell's preferred error direction. For the simulations, we used 1000 units with preferred directions that

covered 0-π in a uniform manner. **c)** Illustration of visual errors, with the direction of the error specified

in polar coordinates. **d-e)** Model-generated adaptation in the cerebellar cortex (d) and deep cerebellar nuclei

(DCN) (e). After experiencing an error in 0 direction, PCs with a preferred direction close to 0 will have high

probability of generating a complex spike (CS) (d, left) which will result in long-term depression (LTD) for active

synapses from granule cell inputs to that PC (d, right). During the preparation of the next movement, the

strength of the input from the parallel fibers (PF) will decrease due LTD, attenuating the SS activity of the PC.

We assume that attenuation of the inhibitory PC output to the DCN will facilitate long-term potentiation (LTP)

resulting from excitatory mossy fiber (MF) input to the DCN (e, left). DCN activation is determined by the

excitatory input from the MF and the inhibitory signal from the PC (e, right). The frame colors indicate the

corresponding pathway in Panel a. **f)** DCN activation plotted in a polar coordinate. Activation across the

population of cells results in a vector (purple arrow) indicating the change in hand angle (Δ hand). Note that

the vector points in the same direction as the error (c, left), and thus serves to compensate for the error.

<u>Behavioral Task and Model Parameterization</u>

We first aim to examine the behavior that emerges from the population dynamics of a network in which

the individual units are tuned to both movement direction and the direction of movement error. For the

work discussed in this section, a single-layer network with this form of representation is sufficient. In the

second half of the Results, we will turn to phenomena that motivate a two layered network that maps on

to the properties of the Purkinje cells within the cerebellar cortex and the deep cerebellar nuclei.

To measure implicit adaptation, we used a variant of a visuomotor rotation task. Participants reached to

a visual target and feedback, when present, was limited to a cursor. To restrict learning to implicit

adaptation, we used task-irrelevant clamped feedback in which the radial position of the cursor was

locked to the hand, but the angular position was predetermined for each trial. In most experiments, the

cursor was shifted by a constant angle relative to the target (Fig 2a-b)[10]. Despite being fully informed of

the non-contingent nature of the feedback and explicitly instructed to ignore the feedback, the reach

angle gradually shifts in the opposite direction of the clamp[10,18,42,43]. Clamp-induced adaptation has all of

the hallmarks of implicit adaptation and, as with other forms of this type of learning, is dependent on the

integrity of the cerebellum[10,44].

To determine the learning and forgetting rates for the units of the CPC model, we conducted an

experiment (Exp 1) in which participants were exposed to 100 trials with clamped feedback (30°) followed

by 60 trials without any feedback. We measured the retention rate from the washout block and then fit

the learning rate from the training phase. The data were also used to determine the scaling factor, a

parameter that transforms neural activation into hand angle. The CPC model provided an excellent fit for

the observed change of hand position (Fig 2c). To rigorously test the model, we fixed the parameters

described above when running simulations to generate predictions for the other experiments.

**Fig. 2 Cerebellar population coding captures learning, forgetting, and anterograde interference during**

**implicit adaptation. a)** For online testing, stimuli are presented on the participant's laptop computer and

movements are made on the trackpad. **b)** For clamped feedback, the angular position of the cursor is rotated

by 30° with respect to the target, regardless of the heading direction of the hand. **c)** Perturbation schedule (top)

and results (bottom) for Exp 1. Time course of the mean hand angle is shown in light violet. The CPC model

provides a good fit in both the training and no-feedback washout phases. **d)** To examine anterograde

interference, the direction of the clamp was reversed during the training section of Exp 2. The behavioral results match the prediction of the CPC, using the parameters estimated from Exp 1. **e)** We mark the direction of a clockwise error (30°) as 0 in the PC/DCN tuning space, and the direction of an opposite (counterclockwise) error (-30°) error as π. Memory of the original perturbation (top row) persists in the anti-clamp training phase, indicated by the activation of neurons tuned to 0 in the bottom row. This residual memory causes anterograde interference. Shaded area in c, d, e indicates standard error.

Anterograde Interference

In the following sections, we will examine several key predictions derived from the CPC model concerning how the experimental context should modulate implicit adaptation. One basic feature of the CPC model is that, for each unit, learning is fast due to the potent impact of complex spikes whereas forgetting in the absence of feedback occurs relatively slowly due to a passive decay process (Fig 2c). Given the slow forgetting, the population activation will be influenced by the persistent activation of units that were tuned to a recent error. As such, when the perturbation direction is abruptly reversed, the model predicts that the observed rate of change in performance will be attenuated due to the persistent activation, even if the learning and forgetting rates are invariant in the model (see Fig 2e).

To examine this model prediction, we used a task in which the sign of the clamp was reversed after an initial training block (e.g., 30° followed by -30°, Exp 2, see Fig 2d). Consistent with the model prediction, the results showed that the rate of adaptation was slower in response to the reversed clamp compared to the original clamp[45–49]. Strikingly, the degree of attenuation closely matched the CPC model's prediction based on the parameter values estimated from Exp 1. These results indicate the population dynamics within the CPC model can be useful to explain the contextual effects in adaptation.

Absence of Spontaneous Recovery

Here we aim to compare the CPC model with several alternative models that have been proposed to

account for contextual effects in implicit adaptation. Anterograde interference has typically been

explained by models positing context-dependent learning mechanisms[30,50,51]. For example, the contextual

inference (COIN) model assumes that the motor system forms separate memories for different contexts

and chooses which memory to use based on the inferred context[29]. To account for the results of Exp 2,

COIN would first build a memory for the 30° perturbation and then a second, distinct memory for the -30°

perturbation. Anterograde interference would arise because the introduction of the -30° perturbation

would lead to some degree of recall of the response to the initial perturbation. Over time, this would shift

to a bias to recall the response to the second memory. In contrast to the COIN model, the CPC model does

not posit distinct memories for different perturbations; rather anterograde interference emerges from

the population dynamics.

The dual state-space (Dual SS) model[21,39] offers another account of anterograde interference. This model

posits that learning involves two processes that operate at different rates (Fig S1a). When the

perturbation is reversed, a fast process will quickly respond to the new perturbation and drive adaption

in the opposite direction. However, a slow process will continue to be dominated by its response to the

original perturbation, thus producing anterograde interference. In contrast, the CPC model can account

for anterograde interference without positing different learning/retention rates across units.

While the COIN, Dual SS, and CPC models make similar predictions about anterograde interference, they

make differential predictions on another memory phenomenon, spontaneous recovery. A paradigmatic

design to elicit spontaneous recovery in sensorimotor learning studies would be to train participants with

a perturbation in one direction, extinguish the adapted behavior by shifting the perturbation in the

opposite direction, and then examine behavior in the absence of feedback (Fig 3a)[21]. Spontaneous

recovery refers to the fact that the initial movements during the no-feedback phase are in the opposite

direction of the initial perturbation (Fig 3b top left). By the COIN model, spontaneous recovery occurs

since there is some degree of recall of the original context during the no-feedback phase. By the Dual SS

model, the state of the fast process will decay back to zero (i.e., baseline). However, the state of the slow

process is still shifted in the direction induced by the initial perturbation, resulting in the manifestation of

spontaneous recovery in the no-feedback phase (Fig 3b, S1b). In contrast, the CPC model predicts that

spontaneous recovery will not occur when learning is restricted to the implicit system since the model

does not have a mechanism for context-dependent memory (Fig 3b bottom right).

To compare the three models, participants in Exp 3 were trained with a 30° clamp in one direction for 100

trials and then presented with the opposite clamp for 15 trials (Fig 3a). Pilot testing had shown that a

reversal of this duration is sufficient to extinguish the shift in hand angle observed to the initial

perturbation. The critical test was the subsequent 30-trials no-feedback block. At odds with the prediction

of COIN or the Dual SS model, we failed to observe spontaneous recovery (Fig 3c; Fig S2b; $t(33)=1.2$,

$p=0.23$). While being cautious in drawing inferences from a null result, the absence of spontaneous

recovery is consistent with the CPC model.

<u>Attenuation in Relearning</u>

As probably an even stronger comparison, all three models make unique predictions in the scenario in

which the initial perturbation is reintroduced after the no-feedback trials (Fig 3a-b, relearning). The COIN

model predicts that relearning should be faster (i.e., exhibit savings) because the system has stored a

memory of the initial perturbation. The Dual SS model predicts that the relearning will be identical to the

original learning, with no saving or attenuation since the system stores no memory of error. The CPC

model predicts that exposure to the opposite error during the washout phase will induce anterograde

interference; as such, relearning will now be attenuated. As with spontaneous recovery, the results are at

odds with the COIN and Dual SS models, and consistent with the CPC model: Adaptation during the re-

exposure block was slower compared to initial learning (Fig 3c; Fig S2a; t(33)=3.1, p=0.004). These results

suggest that population activation within the CPC model accounts for contextual effects that cannot be

captured by alternative models of sensorimotor adaptation.

**Fig. 3 Context as an emergent property of the CPC model. a)** Exp 3 perturbation schedule. To examine

spontaneous recovery, a perturbation in one direction is presented for an extended phase and then reversed

for a short phase. The critical test is in the subsequent no-feedback phase. To test for savings, the original

perturbation is then reintroduced. **b)** The COIN model predicts spontaneous recovery and savings; the Dual SS

model predicts spontaneous recovery but no saving; the MoE model predicts no spontaneous recovery but

savings; the CPC model predicts no spontaneous recovery and attenuation upon relearning. Note that, for

visualization, the data from the relearning phase are plotted on top of the original learning phase. **c)** Empirical

results match both predictions of the CPC model. Shaded area in c indicates standard error. **d)** Population

activation in the CPC model at three time points. While the hand angle is close to zero at the end of the no-

feedback washout, residual activation resulting from both the original and reversed clamps are evident in the

population, indicated by the peak at 0 and $\pi$, respectively (middle). The memory of the opposite clamp slowed

down the relearning (bottom). **e)** In Exp 4, the learning and relearning phases are separated by a variable phase

in which the probability of a perturbation switch is manipulated between participants. **f)** The COIN and MoE

models predict that during relearning, the learning rate will be modulated by the prior switching rate (i.e.,

perturbation variability). The Dual SS model predicts that learning rate will be identical in learning and

relearning regardless of the prior switching rate. The CPC model predicts that, while relearning will be slower

than original learning, the learning rate will not be modulated by switching rate. **g)** Empirical results are in

accord with the CPC model, showing attenuation during relearning and insensitivity to switching rate.

<u>Insensitivity to Error Consistency</u>

The preceding sections demonstrate how the CPC model can account for the dynamics of implicit

adaptation in various contexts without positing flexible context-dependent memory like the COIN model.

A different form of flexibility concerns the operation of meta-learning processes that modulate model

parameters across contexts. This idea is central to the Memory of Error (MoE) model[52], which posits an

optimization the learning parameters based on experienced errors during the training. For example, the

learning rate should increase when recently experienced errors are consistent (stable context), and the

rate should decrease when recently experienced errors are inconsistent. In contrast, the CPC model does

not include a meta-learning process of this sort. Experience-dependent changes in the response to an

error will arise because recently experienced errors have transiently altered the population dynamics of

the system (e.g., Fig 3d).

To compare the MoE and CPC models, we examined how the consistency of error modulates adaptation.

In Exp 4, we tested the response to a clamp with a fixed sign (e.g., 30°) before and after a phase in which

the sign of the clamp varied. To manipulate consistency, we varied the switching probability in the variable

phase, setting it to 12.5%, 50%, or 90% in a between-subject manipulation (Fig 3e). A key prediction of

MoE is that the rate of relearning will be modulated by the switching frequency (Fig 3f bottom). In contrast,

the CPC model predicts that the rate of relearning will be independent of switching frequency.

The results were consistent with the CPC model: The learning rate during the relearning phase was not

modulated by the switching rate during the preceding variable phase (Fig 3g; Fig S2c; $F_{(2,101)}=0.18$,

p=0.84). Interestingly, relearning was markedly slower during the relearning phase compared to original

learning ($F_{(1,101)}=37.7$, p<0.001). This attenuation is another manifestation of anterograde interference

resulting from the opposite errors experienced in the variable-clamp block. We note that these results are

not only at odds with the MoE model, but also with the COIN and Dual SS models. As with the MoE model,

COIN predicts that the learning rate will be inversely related to perturbation variability (Fig 3f); The Dual

SS model predicts that relearning will be identical to that observed during the initial learning phase for all

conditions.

To summarize the results from Experiments 1-4, we have examined a variety of tasks used to examine the

effect of context on sensorimotor adaptation, comparing the CPC model to three prominent alternative

models (Table 1). The results suggest that the population dynamics within the CPC model provide a

parsimonious explanation of how perturbation history influences adaption without positing context-

dependent memory or meta-learning processes.

| | CPC | Dual State Space[21,39] | MoE[52] | COIN[29] |
|---|---|---|---|---|
| Anterograde interference (EXP 2) | ✓ | ✓ | ✗ | ✓ |
| Attenuation in relearning (Exp 3) | ✓ | ✗ | ✗ | ✗ |
| No spontaneous recovery (Exp 3) | ✓ | ✗ | ✓ | ✗ |
| Invariant to error consistency (Exp 4) | ✓ | ✓ | ✗ | ✗ |
| Slower adaptation after variable perturbation (Exp 4) | ✓ | ✗ | ✗ | ✗ |
| Fast single trial learning for variable perturbation (Exp 4/5/8) | ✓ | ✓ | ✗[a] | ✗[a] |
| Different retention rates for variable and fixed perturbations | ✓ | ✓ | ✗ | ✓ |
| Minimal attenuation of the asymptote in half washout (Exp 6) | ✓ | ✗ | ✗ | ✓ |
| Fast single trial learning around the asymptote (Exp 6) | ✓ | ✓ | ✗ | ✗ |
| Stable process is gated by the volatile Process (Exp 7). | ✓ | ✗ | ✗ | ✗ |

**Table 1. Comparison of the CPC model and Other Models of Sensorimotor Adaptation.**

Summary of model performance on a set of core phenomena. In evaluating each of the alternative models, we

used an implementation based on that presented in the associated paper (recognizing that a reasonable variant

might be possible to capture more of the phenomena). The credit assignment model assumes that the agent

performs Bayesian inference to decompose the observed error into perturbation sources that vary across

different time scales and estimate the optimal policy to compensate for them.

[a] Both COIN and the MoE models have difficulty explaining why a large change in behavior would be observed

in response to random perturbations given their optimality assumptions. One would expect an optimal

system to show a minimal or attenuated response to a random perturbation.

## Tuning properties result in variable changing rates across the population units

A central proposition of the Dual SS model is that adaptation includes two processes with different

learning rates (Fig 4a)[21,39,53]. However, within a single layer of the CPC model, an epiphenomenon of

population coding is that units will appear to operate in different time scales even if the learning rate

parameters are identical across all units (Fig 4b). When an error is observed, cells with a preferred

direction centered on this error will display relatively fast learning and quickly saturate (Fig 4c-d). In

contrast, cells with a preferred direction slightly misaligned with the error direction will not only learn

slower due to the weaker climbing fiber input but will also take longer to saturate. As such, by the CPC

model, the change in movement direction is determined by the collective activation of all units, and thus,

can be regarded as a composite process of units with different learning trajectories arising from their error

tuning.

**Fig. 4 Emergent variation in learning rate across the population units. a)** Learning curve from Tsay et al (2022)

in which participants were exposed to a 30° clamp. This function can be described as sum of a fast process that

contributes to the rapid change in hand angle early in learning and a slow process that continues to accumulate

over time. The best-fitted Dual SS model and the relative contribution of the fast and slow processes are shown.

**b)** CPC model can account for learning function without positing different learning rates. **c)** By the CPC model,

the contribution from different units will vary over time due to their tuning for error direction. We mark the

direction of the error as 0 in the unit tuning space. Cells tuned to error direction (i.e., 0) respond strongly,

driving rapid early learning while saturating quickly. Cells with tuning slightly misaligned with the error direction

(e.g., 0.2π) have a small error response but saturate slower; as such they make a relatively large contribution

late in training. **d)** Panel c plotted in polar space. Each vector corresponds to a cell with the orientation indicating

the cell's preferred error direction and the length indicating its activation strength.

<u>Different retention rates in adaptation</u>

We have shown the tuning features of cerebellar units and their population dynamics within the CPC

model explain a wide range of contextual effects in implicit adaptation, outperforming alternative models.

While we have used a two-layered model in these simulations, the predictions would also hold in a single-

layered CPC model. However, the anatomy and physiology of the cerebellum suggest that plasticity effects

in the cerebellar cortex and deep cerebellar nuclei can be quite distinct, perhaps constrained by different

computational principles[21,39,40]. We explore this issue in the following section, first examining the evidence

to suggest computational differences between the layers and then testing predictions to ask how the

layers interact with each other.

As shown above, experimental tasks that focus on learning rates are not ideal for assessing the existence

and/or dynamics of different layers: A single-layered system can appear to be operating at multiple speeds.

As an alternative, we employed a variable perturbation task in which the sign and size of the perturbation

were varied across trials (Exp 5, Fig 5a). With this design, the forgetting rate of the system can be

empirically measured as the ratio of the change of hand angle in response to the perturbation just

experienced (1-back) relative to the previous perturbation (2-back, see Methods). Consistent with

previous studies[10,54,55], we found prominent trial-by-trial adaptation in response to the random

perturbations. Surprisingly, the observed retention rate of 0.5 indicates that about half of the learning

from the previous trial was forgotten over the 3 s inter-trial interval (Fig 5c). This low value stands in

marked contrast with the empirically estimated retention rate from designs in which the perturbation is

fixed (e.g., Exp 1, 0.98, Fig 4b,c). When we ran the same analysis on data from published studies using

variable or fixed perturbations, we observed a similar marked difference in the retention rate[18,55–57].

This result could suggest that there are two adaptation processes with different retention rates.

Alternatively, it might be taken to indicate the operation of a meta-learning process, where the forgetting

rate is ramped up in response to a variable environment. To differentiate those two hypotheses, we ran

an experiment (Exp 6) in which we first employed a fixed perturbation for an extended block and then

followed this with a half-washout phase in which 50% of the trials had no feedback and 50% had clamped

feedback (Fig 5d). Interestingly, there was a considerable drop in hand angle after each no-feedback trial,

much larger than what would be predicted by a single-layered model parameterized with a retention rate

estimated from a typical fixed design (Fig 5e). Indeed, the magnitude of those single-trial changes is

comparable to what is observed in a variable design. A meta-learning model would attribute the drop in

retention rate to the variable context (namely, the mixture of no-feedback and clamp trials). However, at

odds with this hypothesis, the asymptote in the half-washout block largely persisted (Fig 5d). Thus, the

retention rate of the system appears to have remained fixed between the learning and half washout

blocks. As such, the large trial-by-trial changes in hand angle and a persistent asymptote are consistent

with the dual rate hypothesis.

**Fig. 5 Operation of volatile and stable processes in cerebellum-dependent adaptation. a)** Trial-by-trial change

of hand angle (Δhand) as a function of the perturbation size on trial n-1 (1-back), n-2 (2-back), and n-3 (3-back)

in Exp 5. **b)** Left: Similar analysis applied to the data from the variable phase of Exp 4 for the 50% switching

condition. Right: The two-layer CPC model can account for the large, yet transient change in hand angle

observed in response to a random perturbation. A single-layered CPC model predicts negligible change due to

its high retention rate measured from Exp 1. **c)** Estimate of retention rate in experiments using variable or fixed

perturbations. Re-analysis of data from Wei: [55]; Tsay-1: Exp 2 [56]; Tsay-2: EXP 2 [58], Avraham: Exp1 [43]. All of the

depicted experiments used a design with a single target location. **d)** In Exp 6, half washout phase entails a 50/50

mix of clamp and no-feedback trials. Consistent with the CPC model, only a small reduction in hand angle was

observed during the half washout phase whereas the meta learning model with a changeable retention rate

predicts the hand angle will be reduced by 40%. Purple functions indicate behavioral results. **e)** Large trial-by-

trial changes of the hand angle in the half washout phase can only be predicted by the two-layered CPC model

rather than a single-layered model. **f)** The volatile process is hypothesized to produce LTD at the parallel fiber-

PC synapse; the stable process is hypothesized to produce LTP at the mossy fiber-DCN synapse**. g-h)** The dual

rate version of the CPC model (h) provides a better fit of the learning function in Exp 1 compared to the classic

dual SS model (g).

<u>Differential Plasticity in the Cerebellar Cortex and DCN</u>

Physiological studies have shown that learning in the cerebellar cortex appears to be more stable

compared to the DCN. For example, learning-induced changes in the firing rates of simple spikes in

Purkinje cells may decrease after a few trials[39,59,60] whereas changes in the DCN typically last for days[25,35].

Given the evidence reviewed in the previous section pointing to the existence of a system composed of

units with distinct retention rates, we instantiated this difference in the two-layer CPC model such that

plasticity within the cerebellar cortex/PC can produce changes that are weakly retained whereas plasticity

within the DCN is more persistent (Fig 5f and S3).

This conjecture bears similarity to that proposed in previous dual rate models (e.g., Dual SS model)[21,39,61].

However, there are significant differences between these previous models and the two-layer CPC model

characterizes the dynamics of the two processes. First, a key feature of state-space models is that

adaptation reaches an asymptote when the trial-by-trial effects of learning and forgetting cancel each

other out. However, it is difficult to simultaneously fit both the acquisition and washout phases in

response to a fixed perturbation with a state-space model, even with the degrees of freedom conferred

by a dual rate variant (Fig 5g). In contrast, we posit that learning saturates because of a limit to

neuroplasticity within the units[54]. Given this assumption, the CPC model can readily fit the full learning

and forgetting function in adaptation (Fig 5h).

A stronger comparison of the two models is provided by the half-washout phase of Exp 6. The SS model

predicts that during this phase, the asymptote will eventually drop to 50% of the original asymptote

because learning occurs in just 50% of the trials, those with feedback (Fig S6). This prediction holds even

in state-space models that posit learning at multiple time scales[21,62]. However, as noted above, the

asymptote only showed only a slight decrease when clamped feedback was presented in 50% of the trials,

consistent with the predictions of the CPC model (Fig S6b).

A second difference is that, unlike the state space model, the population dynamics within the CPC model

enabled it to account for various context effects discussed in Exp 2-4 (See Table 1). Moreover, by

extending the dual coding feature in a two-layered system, the CPC model predicts a novel context effect

in a random design experiment, one in which the sign of the perturbation is randomized: The trial-by-trial

change in hand angle will decrease over the course of an experimental session even if the

learning/retention rates of both layers are fixed. Early in the session, learning within both a volatile and

stable layer will contribute to the change in performance. However, gradual changes will accumulate in

units tuned to both directions within the stable layer, eventually reaching their asymptotic level of

plasticity. Thus, late in the session, learning is essentially dependent solely on the rapid changes occurring

within the volatile layer. The net effect is that the overall learning rate will decrease over the course of

training (Fig S7a-b), another example in which an apparent change in learning rate is emergent from the

dynamics of the system. In contrast, the dual state-space model predicts that the overall learning rate

should be constant since only the fast process makes a significant response to random perturbations. The

data are again consistent with the CPC model: The overall learning rate decreases over the course of a

block of trials in which the size and direction of the perturbation is randomized (Fig S7c).

<u>Learning within the DCN is gated by the cerebellar cortex.</u>

Given that the output of the PCs is the primary input to the DCN, we can ask how learning within the DCN

is modulated by activity in the cerebellar cortex. By the CPC model, learning in the DCN is scaled by the

change in simple spike activity of the PCs; in effect, the stable process is gated by the labile process. While

this hierarchical organization has been proposed previously in the discussion of dual-rate models of

adaptation[39], it has not been tested empirically.

One way to test the gating hypothesis is to manipulate the duration of the inter-trial interval. The change

in hand angle arising from a volatile process decreases with the passage of time. If the volatile process

gates the stable process, increasing the ITI should result in a much slower change in hand angle in a fixed

design (Fig 6b). In contrast, if the two processes operate in parallel (PARALLEL model), the operation of

the stable process will not be influenced by variation in ITI. Higher asymptotic values in a short ITI

condition would be solely due to the greater contribution of the volatile process.

To evaluate the gating hypothesis, we employed a variable design in Exp 8, using a 7 s ITI. By comparing

this long ITI condition to a short one (Exp 4) which involved the same design but in which the next trial

started as soon as the hand was repositioned at the start location (0 s ITI), we estimated the decay rate

of the volatile process across time. Based on the estimated parameter, we predicted the learning function

in response to a fixed perturbation (Exp 9), again comparing a 7 s and 0 s ITI conditions (Fig 6b). To measure

learning within the stable process (DCN), we ignored the first five trials since these would have a significant

contribution from the volatile process. Over the next five trials, we found that the change in hand angle

was much higher in the short ITI condition (Fig 6d), suggesting that the learning in the stable layer (DCN)

is modulated by the volatile layer (PC).

Furthermore, consistent with the prediction of the CPC model, the long ITI condition showed a slower

decrease in hand angle compared to the short ITI condition in the initial washout trials (Fig 6e), suggesting

a smaller contribution of the volatile process in the long ITI condition. This difference diminished in late

washout trials since the residual memory here comes from the state of the stable process.

However, we note that this version of the CPC model fails to capture one prominent feature in these data,

the convergence of the two functions at asymptote (Fig 6c & f). The CPC model predicts that the advantage

for the short ITI condition should persist, resulting in a lower asymptote in the long ITI condition. We

modified the CPC model (Fig 6g), adding an inhibitory connection from the DCN to the inferior olive, a

pathway that has been identified in physiological studies[63,64]. The inhibitory signal to IO will reduce its

input strength to the cerebellar cortex, thus decreasing the number of PCs that generate complex spikes[65].

Assuming the strength of this inhibition decays across time (ITIs), more learning units can be recruited in

the long ITI condition though plasticity within each unit is reduced due to the decay of memory during the

long ITI (Fig S5). The revised CPC generates learning functions that provide good fits in both ITI conditions

(Fig 6h & i). Importantly, after re-estimating all of the parameters using this variant of the CPC, we

observed negligible effects on the predictions reported for the other experiments (Fig S8). In sum, these

results point to a hierarchical arrangement in which the volatile process gates the operation of the stable

process.

The revised two-layer CPC model can account for another classic learning effect, contextual interference,

referring to the phenomenon in which performance gains are slower when training involves multiple

contexts (e.g., reaching towards multiple directions) compared to training in a single context, but

retention is better in the former[69,70]. This phenomenon, at least in the context of implicit adaptation, is an

emergent property of the parallel operation of volatile and stable learning processes (see Supplementary

result 1, Fig S9).

**Fig. 6 The stable process is gated by the volatile process. a)** Trial-by-trial change in response to a variable perturbation with a short (data taken from Exp 4, p(switch)=0.5 condition) or long ITI (Exp 8). **b)** Predictions of learning functions under the gating assumption of the CPC model and alternative model in which the two processes operate independently (PARALLEL). **c)** Learning functions in Exp 9 using either a short or long ITI. Consistent with the CPC model, the difference between the two functions is reduced over time. **d)** Model predictions and results from Exp 9 for the change of hand angle across trials 5-10. The change of hand angle is higher in the short ITI condition. **e)** The retention rate is larger in the initial no-feedback trials in the long ITI condition since the volatile process is weakened by the passage of time. However, the retention rate is similar across the two ITI conditions late in washout, consistent with the hypothesis that only the stable process remains operative. **f)** Hand angle ratio between short ITI and long ITI conditions deviates from the predictions of both models. The ratio falls between the two model predictions early in training and is smaller than predicted by both models late in training. **g)** Revised CPC model includes inhibitory projection from DCN to the IO. This suppresses the error signal conveyed by the climbing fibers. This suppression is assumed to decay with time, becoming negligible in the long ITI condition in the revised CPC model. **h-i)** Predictions of the revised CPC model

provide a good fit to the learning curve (h) as well as the change in the ratio between the long and short ITI

conditions (i) in Exp 9. Shaded area and error bar indicate standard error.

**Discussion**

To support flexible behavior, an organism needs to choose an action appropriate for a given context and

execute a movement to achieve the desired outcome. A large body of work has sought to delineate the

principles of these learning processes, with one prominent question centering on how the processes

incorporate context and respond to uncertainty. Here we address this question with respect to the

cerebellum, a subcortical structure long recognized as essential for keeping the sensorimotor system

precisely calibrated in the face of fluctuations in the environment or state of the agent. We developed a

population-coding model incorporating two key features: 1) Units are tuned to both movement direction

and error direction, and 2) learning occurs at different rates in the cerebellar cortex and deep cerebellar

nuclei, with the former characterized by a fast, volatile process and the latter characterized by a slower,

stable process. Our CPC model provides a parsimonious account of a diverse range of learning phenomena

and offers new insight into the temporal dynamics of learning (Table S1).

<u>Context Dependency as an Emergent Property of Population Coding</u>

Importantly, there is no explicit role of context in the CPC model in the sense that a context does not

trigger the retrieval of its associated response. Rather the signatures of context-dependent learning and

environmental uncertainty emerge naturally from a population of tuned elements that operate in an

inflexible manner. In this way, the CPC model diverges from classic models in the behavior that emerges

when a previously encountered context is re-experienced. Under such circumstances, classic models

predict savings in relearning given that the context facilitates the retrieval of the appropriate response[29].

In contrast, the CPC model accounts for the fact that when a previously experienced perturbation is

encountered, implicit adaptation not only fails to show savings, but actually can show attenuation[18]. This

attenuation can be seen as another manifestation of anterograde interference: Due to the tuning

properties of neurons in the cerebellar cortex and DCN, persistent activation in response to error in one

direction will interfere with the response to an error in a different direction.

Given the impressive flexibility in human motor learning, it might be surprising that implicit adaptation

does not explicitly track the context or uncertainty of the environment[29,50,66]. We propose that this rigidity

reflects a degree of modularity between processes associated with action selection and those related to

movement implementation. The cerebellum is part of a system designed to use error information to

ensure the accurate execution of a planned movement. The emphasis here is on "planned movement"

rather than "desired outcome" to underscore the point that this system appears to operate independent

of the task goal; indeed, participants will adapt to sensory prediction errors even when the change in

behavior is detrimental to task success[10,12]. This modularity provides a means to keep the system properly

calibrated across changes in the internal state of the organism (e.g., perceptual biases, fatigue), factors

that need not require a change in action planning. In contrast, other learning systems are designed to use

error information related to task success to determine if the selected action was optimal given the current

context. These systems would be optimized to track contextual shifts in determining the appropriate

policy. Consistent with this hypothesis, contextual effects such as savings and sensitivity to uncertainty

are observed in adaptation tasks that benefit from changes in action selection[52,67,68].

<u>Hierarchical Organization Within the Cerebellum for Implicit Adaptation</u>

The behavior observed in adaptation studies is assumed to reflect the function of learning processes that

operate at different time scales[21,61]. It has been suggested that fast and slow processes correspond to

explicit and implicit learning processes[62]. By using variable and fixed designs, we provide evidence that

learning limited to just the implicit system operates at different timescales, a notion similar to the original

framing of the Dual SS model.[21,39] However, rather than view these as processes that operate in parallel,

our empirical and modeling results highlight a hierarchical organization in which accumulated learning

from a volatile process will constrain the learning rate of a stable process. This organization readily maps

onto a two-layered network formed by the cerebellar cortex and DCN, with the output from the former

gating learning within the latter. The neurophysiological evidence is consistent with this assumption.

While the change of SS activation in the PCs can happen within a few trials[22,59], changes within the DCN

can maintain learning across days[25,35]. Reflective of the hierarchical organization, we showed that there is

an asymmetric dependency such that the synaptic strength in the cerebellar cortex determines the PC

output that modulates learning within the DCN.

We recognize that a two-layered model is clearly a simplification. Indeed, to explain the asymptotic

convergence in the long and short ITI conditions, we had to incorporate a third layer into the model,

creating a closed loop by adding a projection from the DCN to the IO. While the anatomy supports the

existence of this pathway, to achieve convergence, we added two specific features to the dynamics of this

pathway. First, the intensity of the inhibition from the DCN to the IO exhibits intensity decrease over

time[71,72]. Second, the projection is generic, inhibiting IO units independent of the directional tuning of the

DCN neuron. These two assumptions need to be tested in future physiological studies.

<u>Generalization of the CPC model</u>

Though our model focuses on the cerebellum and sensorimotor learning, the core computational

principles may offer insight into how the nervous system responds to environmental uncertainty. The

population-coding aspect of the CPC model is similar to models of perceptual learning that include a basis

set of tuned elements[73]. For example, in models of time perception, contextual effects on perceived

duration have been proposed to reflect the interaction of units tuned to different durations[74,75]. Applied

to this domain, the CPC model could be used to derive specific predictions of how temporal perception is

modulated by uncertainty and establish boundary conditions for interference. Moreover, the two-layer

network in the CPC model provide a novel framework to understand of how learning can involve multiple

processes that follow different temporal dynamics, a phenomenon widely observed in cognitive tasks. For

instance, value learning has been hypothesize to reflect the joint operation of a fast, volatile process and

a slow, stable process[70]. While these processes are typically viewed as operating in parallel, the CPC model

offers an example of how a hierarchical framework might prove more parsimonious.

**Methods**

<u>Cerebellar Population Coding (CPC) model</u>

Here we extend the classic Marr-Albus model, focusing on how learning is modulated when the

environment is variable. A foundational idea for our model is inspired by a recent work showing how PCs

in the oculomotor cerebellar cortex are simultaneously tuned to both movement direction and the error

that is associated with that movement [22,23].

To examine the implications of these tuning properties for cerebellar-dependent learning, we incorporate

PC tuning into a learning model. Specifically, we formalize the teaching signal, the complex spike (CS)

activity of a PC with a preferred direction of $i$ ($0 \leq i < \pi$) in response to a movement error $e$ (Fig 1d) as:

$$[1]\ CS_i^n = VM(\theta^e, i, s)F(\rho^e)$$

where $VM(\,i, s)$ is the probability density function of a simplified circular (von Mises) distribution with a

mean of $i$ and standard deviation of $s$. $\theta^e$ and $\rho^e$ refer to the direction and the size of $e$, respectively, and

$n$ is the trial number. Given we only applied one error size across all experiments, $F(\rho^e)$ is not relevant

and set as 1 here[55,57]. The form of this non-linear relationship is not relevant in the experiments with a

fixed perturbation; for those experiments, the exponent was set to 1.

Following the Marr-Albus model, the occurrence of a CS suppresses the strength of the parallel fiber input

synapse ($w$) through long-term depression (LTD):

$$[2]\ w_i^{n+1} = -lCS_i^n + f(w_o - w_i^n) + w_i^n$$

where $l$ ($l > 0$) and $f$ ($0 < f < 1$) are the learning and forgetting rates, respectively, and $w_o$ is the

baseline synaptic strength. Since the level of single spike (SS) activity will be greatest for cells coding a

movement direction opposite to the error, the modulation of synaptic strength will drive the next

movement in a direction that corrects for the observed error.

The preceding paragraph describes how parallel fiber synapses onto PCs are modified. A second

prominent site of plasticity is at deep cerebellar nuclei[25,35]. Importantly, PC and DCN neurons are

organized such that they share the same tuning direction for movement[40]. We posit that learning at the

DCN is gated by learning at the cerebellar cortex. Specifically, LTD at parallel fiber-PC (PF-PC) synapses will

reduce inhibitory PC input to the DCN, resulting in long term potentiation (LTP) at the mossy fiber-DCN

synapses ($m$) (Fig 1e):

$$[3]\ m_i^{n+1} = (w_{\mathrm{o}} - w_i^n)\text{ß}(m_{\max} - m_i^n) + \alpha(m_{\mathrm{o}} - m_i^n) + m_i^n$$

where ß and $\alpha$ are the learning rate and the forgetting rate of the DCN input synapse, respectively. The

parameters $m_{\mathrm{o}}$ and $m_{\max}$ represent baseline and maximal synaptic strength, respectively. The latter

constraint is based on empirical results showing that implicit adaptation saturates independent of the

error size.

In this initial version of the CPC model, we have assumed that the PC layer is dominated by LTD, and the

DCN layer is dominated by LTP. Classically, PC-layer learning has emphasized LTD[1,2,76,77], including recent

evidence from rodent work showing LTD during upper limb reach adaptation[3]. There is a dearth of

evidence concerning the mechanisms of learning in the DCN. As such, we assumed that learning here

follows a simple Hebbian process, one consistent with LTP. Importantly, our assumptions regarding LTD

and LTP are not critical computationally. The results would be similar if we assumed the reverse or had a

mixture of LTD and LTP at each layer. Indeed, a more realistic model should incorporate some degree of

bidirectionality given that LTP is also observed in the PC layer[23,60]. For example, an error signal at 0 could

result in units with a preferred direction at π becoming strengthened by LTP[22], presumably because of

their co-activation with a shared parallel fiber input. We expect a model with bi-directional modulation

would have greater flexibility. However, for the present purposes, the predictions would be largely

unchanged and thus, we opted to go with a simple dichotomy.

Considering the two sites of plasticity, DCN activity on a repeated trial following a movement error can be

formalized as:

$$[4]\ DCN_i^{n+1} \propto m_i^{n+1} - \gamma w_i^{n+1}$$

where $\gamma$ is a scaling factor. The output of the population of DCN neurons will correspond to the change in

movement direction in response to an error, a signal that can be used to adjust the movement. This can

be expressed as: (Fig 1f):

$$[5]\ \boldsymbol{h}^{n+1} = -\varepsilon \sum_i \boldsymbol{v}_i DCN_i^{n+1}$$

where $\boldsymbol{h}^n$ is a vector representing the hand angle on trial n, $\boldsymbol{v}_i$ is a vector representing the tuning direction

of unit $i$, and $\varepsilon$ is a scaling factor to transfer the neural activity into hand angle.

We note that we do not specify, for the purposes of this paper, whether the cerebellum is best viewed as

a forward or inverse model. The current implementation of the CPC model most resembles an inverse

model given that the system modifies the motor commends based on the error. However, the model could

be reframed as a forward model with the output a prediction that is fed into an (cerebellar or

extracerebellar) inverse model that refines the motor commands

<u>Parameterization of the CPC Model</u>

In our simulations of PC and DCN neurons, we modeled 1000 units for each layer and set the standard

deviation of the tuning function (*s*) to 0.2π. The results of most simulations were not sensitive to these

two parameters. While anatomical studies show considerable convergence from the PC layer to the DCN,

for simplicity, we opted to impose a one-to-one connection between the PC and DCN.

We used an empirical approach to estimate the learning and forgetting rate for PF-PC synapses, using the

data from Exp 4 in which +/- 30° clamps were presented with a 50% switching probability. To measure

single trial learning, we calculate the change of hand angle between trial n and trial n-1, flipping the sign

when the clamp on trial n-1 was negative. To measure single trial forgetting, we calculate the change of

hand angle between trial n and trial n-1, flipping the sign when the clamp on trial n-2 was negative.

PF-PC forgetting ($f$) is the ratio of single-trial forgetting and single-trial-learning. By definition, retention

rate is 1- ($f$). We applied the same method to measure the retention for all variable designs and this gave

us an $f$ around 0.5. Model simulations indicate that this method can precisely estimate retention when

the perturbation is random. In all of the analyses, we excluded the first 50 trials since learning at this early

stage is influenced by both PC and DCN. For comparing the learning rate between early and late training

in a by variable design, we employed the same general approach but limited the analysis to the first 50

trials to estimate early learning (Fig S8).

The baseline and maximal strength of MF-DCN synapses can be set to arbitrary values: We used 1 and

1.85 for $m_o$ and $m_{max}$, respectively. We measured the retention rate of the MF-DCN synapse ($\alpha$)

empirically using the data from the no-feedback washout phase in Exp 1:

651
$$[6]\ \alpha = \sqrt[10]{mean(\frac{y^{n+10}}{y^n})}$$

where $y^n$ is the hand angle in trial n. The first 20 trials in the washout phase were excluded since they

may be contaminated by a volatile process.

The learning rate of the PC ($l$) and DCN (ß) and the scaling factors ($\gamma, \varepsilon$) were jointly fitted from the

learning block in Exp 1 and the single trial learning in Exp 4. This results in a set of parameters: $l$ = .05, $f$

= .018, ß = 2, $\alpha$ = .5, $\gamma$ = 0.15, $\varepsilon$ = 130. These parameters were fixed in the simulations of all the other

experiments. The only exception is Exp 9, where we set the PF-PC retention rate for the long ITI conditions

($f'$) to be 0.3, based on the empirically observed value in the variable design of Exp 8.

<u>Revised CPC model</u>

The results of Exp 8 led us to develop a post-hoc variant in which the output of the cerebellum modulates

the input, an idea that is consistent with cerebellar anatomy and physiology[63,78]. The basic version of the

CPC model predicts that learning in a long ITI condition will reach a lower asymptote compared to a short

ITI condition. This occurs because the contribution of the volatile process is suppressed in the long ITI

condition. However, the results of Exp 9 showed that, with a sufficient number of trials, learning in the

long ITI condition eventually reaches the same asymptote as in the short ITI condition. This observation

led us to revise the model by adding an inhibitory pathway from the DCN to the inferior olive[63,78].

We assume that the output of the DCN integrates the activation of directionally tuned units and that this

signal serves as a generic inhibitory signal to the inferior olive. We implemented this generic suppression

by subtracting a common value from the activation of cells tuned to all error directions in the inferior olive

(IO):

$$[7] IO_i = 1 - \omega * \sum_i dDCN_i^n$$

$$[8] \text{ if } IO_i > 0: cs'_i = IO_i * cs_i;$$

$$\text{otherwise: } cs'_i = 0$$

where $\omega$ represents the strength of suppression. Given the assumption that $\omega$ decreases across time, we

used separate parameter values of $\omega$ for the long and short ITI conditions $\sum_i dDCN_i^n$ is the sum of the

change of all NCD units relative to their baseline activities. $cs'_i$ is the corrected CS activation value after

taking the DCN-IO pathway into the consideration and replaces the $cs_i$ term in EQ [1-5]. The retention

rates of the volatile and stable processes ($f, \alpha$) in the revised CPC model were set as in the basic two-layer

model. The other parameters ($l$, ß, $\varepsilon, \omega$) were jointly fitted from two data sets, the learning block in Exp 1

and the variable condition in Exp 8. The parameter set is as follow: $l$ = .1, $f$ = .018, ß = 2, $\alpha$ = .5, $\gamma$ = .2, $\varepsilon$ =

210, $\omega(short)$ = 2.5, $\omega(long)$ = 0.

<u>Alternative Models for Comparison</u>

*Variants of the CPC Model*

To help clarify the importance of a two-layer model, we describe two variants of the CPC model. First, we

implemented a single-layer version of the CPC model by modifying Eq 4 to:

$$[9]\ DCN_i^{n+1} = m_i^{n+1}$$

In this version, the output of the system is solely determined by the strength of the MF-DCN.

Second, we implemented a model in which the volatile and the stable processes operate in parallel

(PARALLEL) rather than hierarchical as in the CPC model. Since the stable process is insensitive to ITI, we

estimate the MF-DCN synapse ($m$) by simulations using a short ITI. The simulated value was then used in

simulations of the long ITI condition. For the volatile process, the strength of the PF-PC synapse ($w$) was

measured separately for the two ITI conditions.

*State-space model*

We employed a standard version of a state-space model[21,79]:

$[10]\ x(n+1) = a*x(n) + b(e,n)e(n) + \varepsilon_x(n)$

where $x$ is the internal estimate of the motor state (i.e., the hand movement required to compensate for

the perturbation), $a$ is the retention factor, $e(n)$ is the size of the perturbation in trial $n$, $b$ is the error

sensitivity for a given error size, and $\varepsilon_x$ represents planning noise.

The actual motor response on trial $n$ is given as:

$[11]\ y(n) = x(n) + \varepsilon_y(n)$

where $y$ is the reaching direction relative to the target, determined by $x(n)$ and motor noise, $\varepsilon_y$ .

For the dual state-space model, we added a second, slower learning process ($xs$) with a different retention

rate ($as$) and learning rate ($bs$),

$[12]\ xs(n+1) = as*xs(n) + bs(e,n)e(n) + \varepsilon_{xs}(n)$

Where $as > a$ and $bs < b$. As such, Eq [11] can be written as:

$[13]\ y(n) = x(n) + xs(n) + \varepsilon_y(n)$

*Memory-of-Error model (MoE)*

The Memory-of-Error model describes how the learning rate in the state-space model is modulated by

experience. In the MoE model, error sensitivity (*b*) is set to an initial value that is modulated by errors that

are experienced during training. Specifically, *b(e,n)* will increase if the error on trial n+1 shares the same

sign and *b(e, n)* will decrease if the error on trial n+1 is of the opposite sign. This is formalized as:

$[14]\ b(e(n), n+1) = \alpha * (b(e(n), n+1) - b0) + b0 + \beta * sign(e(n) * e(n+1))$

where $\beta$ and $\alpha$ are the learning rate and retention rate of *b*, respectively. Since the error size is fixed at

30° in our experiments, we replace $b(e)$ with a single value $b$.

*Contextual Inference (COIN) model*

We simulated the Contextual Interference (COIN) using the code provided by Heald et al.[29], focusing on

its prediction with respect to savings and spontaneous recovery. We assumed that the introduction of a

perturbation (e.g., clamped feedback) defines a new context and, as such, leads to the establishment of a

new motor memory. Similarly, reversing the sign of the perturbation would define another context and

thus require establishment of another memory. We simulated the clamps as if they were contingent

rotations so that the learning can reach an asymptote. Before each movement, the output is determined

by averaging the state of different contexts weighted by the expected probabilities of the contexts.

Participants observed an error after each movement and update the state estimation.

<u>Behavioral Experiments</u>

*Participants*

A total of 451 participants (297 female, mean age = 28.0, SD = 5.3) were recruited through the website

prolific.co. After eliminating participants who failed to meet our performance criteria (2.8%, see below),

the analyses were based on data from 438 participants. Based on a survey included in a prescreening

questionnaire, the participants were right-handed with normal or corrected-to-normal vision. The

participants were paid based on a rate of $8/h. The protocol was approved by the Institutional Review

Board at the University of California, Berkeley. Informed consent was obtained from all participants.

*Apparatus*

All of the behavioral experiments were conducted online using a web-based experimental platform,

OnPoint[58], which is written in JavaScript and presented via Google Chrome. It is designed to operate on

any laptop computer. Visual stimuli were presented on the laptop monitor and movements were

produced on the computer trackpad. Data were collected and stored using Google Firebase.

*Clamp rotation task*

We applied clamp feedback in the experiments, under the assumption that learning in response to this

type of feedback is limited to implicit, cerebellar-dependent sensorimotor recalibration. To start each trial,

the participant moved the cursor to a white start circle (radius: 1% of the screen height) positioned in the

center of the screen. After 500ms, the target, a blue circle (radius: 1% of the screen height) appeared with

the radial distance set to 40% of the screen size. The target appears at -45°, a workspace location selected

because it exhibits minimal bias across participants[80]. The participant was instructed to produce a rapid,

out-and-back movement, attempting to intersect the target. If the movement time (from onset to time at

which movement amplitude reached the target) was longer than 500ms, the message 'Too Slow' was

presented on the screen for 500ms.

There were three types of feedback. On veridical feedback trials, the position of the cursor moved was

matched to the position of the hand, subject to the translation in reference frames (screen assumed to

be vertical, hand movement assumed to be horizontal) and scaling (trackpad space expanded to

encompass most of the screen). On clamped feedback trials, the cursor followed a fixed path. As with

veridical feedback, the radial location of the cursor was based on the radial extent of the participant's

hand. However, the angular position of the cursor was independent of the position of the hand, instead

determined relative to the position of the target. The clamp angle was set at 30° relative to the target

except for Exp 5 and 8 (see below). On no feedback trials, the cursor was blanked at movement onset.

On veridical and clamped feedback trials, after the amplitude of the movement reached the target

distance, the cursor was presented at the target distance for another 50ms then it disappeared. Target

disappeared after 200ms. The cursor was then reset to a random position on an invisible circle with a

radius equal to 10% of the target distance and the participant moved the cursor back to the start circle.

At the onset of the first block of trials involving perturbed feedback, the experiment was paused, and a

set of instructions were presented to describe the clamped feedback. The participant was informed that

the cursor would no longer be linked to their movement but rather would follow a fixed path on all trials.

The participant was instructed to always reach directly to the target, ignoring the cursor. These

instructions were then repeated twice to emphasize the atypical nature of the feedback. After the first 10

trials with clamped feedback, a new instruction screen appeared in which the participant was asked to

indicate where they were aiming on each trial. If the participant indicated they were reaching somewhere

other than the target, the experiment was terminated.

Each experiment started with two baseline blocks: First a no-feedback block of 10 trials and second, a

veridical feedback block of 10 trials. For experiments using a fixed design (direction and size of

perturbation remain constant), the direction of the clamp (counterclockwise, CCW; clockwise; CW) was

counterbalanced across participants.

*Experiment 1*

Exp 1 was designed to determine the parameters of the CPC model. There was a total of 180 trials. The

two baseline blocks were followed by a learning block of 100 trials with clamped feedback with learning

expected to reach an asymptotic level in response to a fixed perturbation. This was followed by a final no-

feedback block of 60 trials. 30 participants were recruited for Exp 1 (29 valid, 5 males, age: 27.4 ± 4.9

years).

*Experiment 2*

Exp 2 was designed to measure antegrade interference. The baseline and initial perturbation blocks were

as in Exp 1. For the final block (150 trials), the direction of the clamp was reversed (e.g., from 30° to -30°).

30 participants were recruited for Exp 6 (30 valid, 10 males, age: 30.3 ± 4.3 years).

*Experiment 3*

Exp 3 was designed to assess spontaneous recovery and savings in implicit adaptation. The baseline and

initial perturbation blocks were as in Exp 2. We then included a 15-trial block with the clamp reversed

under the assumption that this would be a sufficient number of trials to bring the hand angle back to

baseline. This was followed by no-feedback block (35 trials) to examine spontaneous recovery and then a

100-trial relearning block in which the clamp feedback was identical to that used in the first perturbation

block. 34 participants were recruited for Exp 3 (34 valid, 16 males, age: 22.7 ± 4.8 years).

*Experiment 4*

Exp 4 examined how the consistency of the perturbation influenced implicit adaptation. The first blocks

were identical to Exp 3, providing initial exposure to clamped feedback and then a reversed clamp to bring

the hand angle back to baseline. This was followed by a 300-trial block in which the clamp changed sign

in a probabilistic manner. The probability of a sign change was either 90%, 50%, and 12.5% in a between-

subject manipulation. The sequence of clamps was preset to ensure that clockwise and counterclockwise

occurred on 50% of the trials each across the 300 trials. The experiment ended with a relearning block in

which the initial perturbation was presented for 100 trials. 36/40/36 participants were recruited for 90%,

50%, and 12.5% conditions respectively (34/38/33 valid, 37 males, age: 28.6 ± 5.5 years).

*Experiment 5*

To estimate the learning rate and retention at top layer of the CPC model, the PF-PC synapse, we

employed a variable design in which the error size and direction varied across trials. After the two baseline

sections, participants completed a 540-trial random perturbation block. Here the clamp size ranged from

-135° to 135° in steps of 1°. The size/direction was determined at random with the constraint that each

clamp was selected once every 270 trials. 72 participants were recruited for Exp 5 (70 valid, 25 males, age:

26.2 ± 5.2 years).

*Experiment 6*

Exp 6 was designed to evaluate different models of asymptotic adaptation. A 10-trial feedback baseline

block was followed by a learning block of 100 trials with clamped feedback. We then alternated between

no-feedback and clamp feedback trials for 60 trials (half-wash phase). 40 participants were recruited for

Exp 6 (38 valid, 8 males, age: 30.7 ± 6.6 years).

*Experiment 7*

Experiment 7 was designed to measure the time course of retention during the initial washout phase.

After the two baseline blocks, the perturbation block consisted of 31 mini-blocks, each composed of 10

trials with clamped feedback and 10 trials without feedback (620 trials). 57 participants were recruited

for Exp 7 (57 valid, 12 males, age: 28.3 ± 5.4 years).

*Experiment 8*

To quantify the temporal dynamics of volatile processes, we used variable clamped feedback with

extended inter-trial intervals (ITI) in Exp 8. For the long ITI, the interval between the end of one trial and

the start of the next trial was 6 s, 7 s, or 8 s, randomized across trials. The message "wait" was displayed

on the monitor after each trial. Exp 8 included two baseline blocks and a 180-trial learning block in which

a 30° perturbation was randomly selected to be either clockwise or counterclockwise, subject to the

constraint that each direction occurred four times every 8 trials. For the short ITI condition, we used the

data from Exp 4 for the variable condition (0 s ITI). 28 participants were recruited for each condition (27

valid, 13 males, age: 28.1 ± 4.8 years).

*Experiment 9*

To understand how the volatile and stable learning processes are jointly modulated by time, we used a

fixed design in Exp 9. The design was similar to that employed in Exp 1 with one notable modification. We

included a 10-trial familiarization block following the two baseline blocks to demonstrate the clamp

feedback. The clamp size in the familiarization block varied from -90° to 90° across trials to show that the

cursor is unaffected by the direction of hand movement. To avoid the influence of pre-exposure to the

error signal on learning, the familiarization block utilized a different target (45°) from the other blocks

(315°). Two groups of participants performed the task with either long ITI (6-8s) or short ITI (0s). 26

participants were recruited for each condition (51 valid, 21 males, age: 26.8 ± 4.6 years).

<u>Data analyses</u>

Hand angle was calculated as the angle difference a line from the start position to the target and a line

from the start position to the hand position at the target radius. Positive values indicate hand angles in

the opposite direction of the perturbation, the direction one would expect due to adaptation. Trials with

a movement duration longer than 500 ms or an error larger than 70° were excluded from the analyses.

We excluded the entire data from participants who had less than 70% valid trials (2.8% participants).

Between-condition comparisons were performed with t-tests or ANOVAs. Learning and relearning were

compared by paired-t-test. For all the statistical tests, we confirmed that the data met the assumptions

of a Gaussian distribution and homoscedasticity.

<u>Data and Software Availability</u>

The data and code supporting this work are available at https://github.com/shion707/CPC.

**Author contributions**

T.W., R.B.I. contributed to the conceptual development of this project. T.W. collected the data, analyzed the data, prepared the figures, and wrote the initial draft of the paper, with both authors participated in the editing process.

**Acknowledgment**

This study is funded by the NIH (grants NS116883 and NS105839). We thank Chris Miall for helpful discussion.

**Competing interests**

RI is a co-founder with equity in Magnetic Tides, Inc.

**References**

1.  Ito, M. Mechanisms of motor learning in the cerebellum. *Brain Res.* **886**, 237–245 (2000).

2.  Albus, J. S. A theory of cerebellar function. *Math. Biosci.* **10**, 25–61 (1971).

3.  Calame, D. J., Becker, M. I. & Person, A. L. Cerebellar associative learning underlies skilled reach
adaptation. *Nat. Neurosci.* **26**, 1068–1079 (2023).

4.  Albergaria, C., Silva, N. T., Pritchett, D. L. & Carey, M. R. Locomotor activity modulates associative
learning in mouse cerebellum. *Nat. Neurosci.* **21**, 725–735 (2018).

5.  Yeo, C. H., Hardiman, M. J. & Glickstein, M. Classical conditioning of the nictitating membrane
response of the rabbit. I. Lesions of the cerebellar nuclei. *Exp. Brain Res.* **60**, 87–98 (1985).

6.  Bastian, A. J. Moving, sensing and learning with cerebellar damage. *Curr. Opin. Neurobiol.* **21**, 596–
601 (2011).

7.  Taylor, J. A. & Ivry, R. B. Cerebellar and prefrontal cortex contributions to adaptation, strategies, and
reinforcement learning. *Prog. Brain Res.* **210**, 217–253 (2014).

8.  Kawato, M. Internal models for motor control and trajectory planning. *Curr. Opin. Neurobiol.* **9**, 718–
727 (1999).

9.  Ebner, T. J. & Pasalar, S. Cerebellum predicts the future motor state. *Cerebellum* **7**, 583–588 (2008).

10. Morehead, J. R., Taylor, J. A., Parvin, D. E. & Ivry, R. B. Characteristics of Implicit Sensorimotor
Adaptation Revealed by Task-irrelevant Clamped Feedback. *J. Cogn. Neurosci.* **29**, 1061–1074 (2017).

11. Taylor, J. A., Klemfuss, N. M. & Ivry, R. B. An Explicit Strategy Prevails When the Cerebellum Fails to
Compute Movement Errors. *Cerebellum* **9**, 580–586 (12/2010).

12. Mazzoni, P. & Krakauer, J. W. An implicit plan overrides an explicit strategy during visuomotor
adaptation. *J. Neurosci.* **26**, 3642–3645 (2006).

13. Taylor, J. A., Krakauer, J. W. & Ivry, R. B. Explicit and Implicit Contributions to Learning in a
Sensorimotor Adaptation Task. *Journal of Neuroscience* **34**, 3023–3032 (2014).

14. Wang, T. & Taylor, J. A. Implicit adaptation to mirror reversal is in the correct coordinate system but the wrong direction. *J. Neurophysiol.* **126**, 1478–1489 (2021).

15. Wilterson, S. A. & Taylor, J. A. Implicit Visuomotor Adaptation Remains Limited after Several Days of Training. *eNeuro* **8**, (2021).

16. Avraham, G., Keizman, M. & Shmuelof, L. Environmental consistency modulation of error sensitivity during motor adaptation is explicitly controlled. *J. Neurophysiol.* **123**, 57–69 (2020).

17. Wang, T., Avraham, G., Tsay, J. S., Abram, S. J. & Ivry, R. B. Perturbation Variability Does Not Influence Implicit Sensorimotor Adaptation. *bioRxiv* 2023.01.27.525949 (2023) doi:10.1101/2023.01.27.525949.

18. Avraham, G., Morehead, J. R., Kim, H. E. & Ivry, R. B. Reexposure to a sensorimotor perturbation produces opposite effects on explicit and implicit learning processes. *PLoS Biol.* **19**, e3001147 (2021).

19. Denny-Brown, D. Conditioned reflexes: An investigation of the physiological activity of the cerebral cortex. *Nature* **121**, 662–664 (1928).

20. Ebbinghaus, H. Memory: a contribution to experimental psychology. *Ann. Neurosci.* **20**, 155–156 (2013).

21. Smith, M. A., Ghazizadeh, A. & Shadmehr, R. Interacting Adaptive Processes with Different Timescales Underlie Short-Term Motor Learning. *PLoS Biol.* **4**, e179 (2006).

22. Herzfeld, D. J., Kojima, Y., Soetedjo, R. & Shadmehr, R. Encoding of action by the Purkinje cells of the cerebellum. *Nature* **526**, 439–442 (2015).

23. Sedaghat-Nejad, E., Pi, J. S., Hage, P., Fakharian, M. A. & Shadmehr, R. Synchronous spiking of cerebellar Purkinje cells during control of movements. *Proceedings of the National Academy of Sciences* **119**, e2118954119 (2022).

24. Junker, M. *et al.* Learning from the past: A reverberation of past errors in the cerebellar climbing fiber signal. *PLoS Biol.* **16**, e2004344 (2018).

25. Carulli, D. *et al.* Cerebellar plasticity and associative memories are controlled by perineuronal nets. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 6855–6865 (2020).

26. Raymond, J. L., Lisberger, S. G. & Mauk, M. D. The cerebellum: a neuronal learning machine? *Science* **272**, 1126–1131 (1996).

27. Lisberger, S. G., Pavelko, T. A. & Broussard, D. M. Responses during eye movements of brain stem neurons that receive monosynaptic inhibition from the flocculus and ventral paraflocculus in monkeys. *J. Neurophysiol.* **72**, 909–927 (1994).

28. Lee, K. H. *et al.* Circuit mechanisms underlying motor memory formation in the cerebellum. *Neuron* **86**, 529–540 (2015).

29. Heald, J. B., Lengyel, M. & Wolpert, D. M. Contextual inference underlies the learning of sensorimotor repertoires. *Nature* **600**, 489–493 (2021).

30. Haruno, M., Wolpert, D. M. & Kawato, M. Hierarchical MOSAIC for movement generation. *Int. Congr. Ser.* **1250**, 575–590 (2003).

31. Collins, A. & Koechlin, E. Reasoning, learning, and creativity: frontal lobe function and human decision-making. *PLoS Biol.* **10**, e1001293 (2012).

32. Kawato, M. Feedback-Error-Learning Neural Network for Supervised Motor Learning. in *Advanced Neural Computers* (ed. Eckmiller, R.) 365–372 (North-Holland, Amsterdam, 1990).

33. Wolpert, D. M., Miall, R. C. & Kawato, M. Internal models in the cerebellum. *Trends Cogn. Sci.* **2**, 338–347 (1998).

34. Apps, R. & Garwicz, M. Anatomical and physiological foundations of cerebellar information processing. *Nat. Rev. Neurosci.* **6**, 297–311 (2005).

35. Moscato, L. *et al.* Long-Lasting Response Changes in Deep Cerebellar Nuclei in vivo Correlate With Low-Frequency Oscillations. *Front. Cell. Neurosci.* **13**, 84 (2019).

36.  Ito, M. Cerebellar control of the vestibulo-ocular reflex--around the flocculus hypothesis. *Annu. Rev.*

*Neurosci.* **5**, 275–296 (1982).

37.  Medina, J. F., Garcia, K. S., Nores, W. L., Taylor, N. M. & Mauk, M. D. Timing mechanisms in the

cerebellum: testing predictions of a large-scale computer simulation. *J. Neurosci.* **20**, 5516–5525

(2000).

38.  Garcia, K. S. & Mauk, M. D. Pharmacological analysis of cerebellar contributions to the timing and

expression of conditioned eyelid responses. *Neuropharmacology* **37**, 471–480 (1998).

39.  Herzfeld, D. J., Hall, N. J., Tringides, M. & Lisberger, S. G. Principles of operation of a cerebellar

learning circuit. *Elife* **9**, (2020).

40.  Shadmehr, R. Population coding in the cerebellum: a machine learning perspective. *J. Neurophysiol.*

**124**, 2022–2051 (2020).

41.  Pugh, J. R. & Raman, I. M. Mechanisms of potentiation of mossy fiber EPSCs in the cerebellar nuclei

by coincident synaptic excitation and inhibition. *J. Neurosci.* **28**, 10549–10560 (2008).

42.  Kim, H. E., Avraham, G. & Ivry, R. B. The Psychology of Reaching: Action Selection, Movement

Implementation, and Sensorimotor Learning. *Annu. Rev. Psychol.* **72**, 61–95 (2021).

43.  Avraham, G., Taylor, J. A., Breska, A., Ivry, R. B. & McDougle, S. D. Contextual effects in sensorimotor

adaptation adhere to associative learning rules. *Elife* **11**, e75801 (2022).

44.  Tsay, J. S., Parvin, D. E. & Ivry, R. B. Continuous reports of sensed hand position during sensorimotor

adaptation. *J. Neurophysiol.* **124**, 1122–1130 (2020).

45.  Morehead, J. R. & Smith, M. The magnitude of implicit sensorimotor adaptation is limited by

continuous forgetting. *Abstract. Advances in Motor Learning & Motor Control* (2017).

46.  Lerner, G. *et al.* The origins of anterograde interference in visuomotor adaptation. *Cereb. Cortex* **30**,

4000–4010 (2020).

47. Brashers-Krug, T., Shadmehr, R. & Bizzi, E. Consolidation in human motor memory. *Nature* **382**, 252–

255 (1996).

48. Leow, L.-A., Hammond, G. & de Rugy, A. Anodal motor cortex stimulation paired with movement

repetition increases anterograde interference but not savings. *Eur. J. Neurosci.* **40**, 3243–3252 (2014).

49. Sing, G. C. & Smith, M. A. Reduction in learning rates associated with anterograde interference results

from interactions between different timescales in motor adaptation. *PLoS Comput. Biol.* **6**, e1000893

(2010).

50. Heald, J. B., Lengyel, M. & Wolpert, D. M. Contextual inference in learning and memory. *Trends Cogn.*

*Sci.* (2022) doi:10.1016/j.tics.2022.10.004.

51. Ingram, J. N., Flanagan, J. R. & Wolpert, D. M. Context-dependent decay of motor memories during

skill acquisition. *Curr. Biol.* **23**, 1107–1112 (2013).

52. Herzfeld, D. J., Vaswani, P. A., Marko, M. K. & Shadmehr, R. A memory of errors in sensorimotor

learning. *Science* **345**, 1349–1353 (2014).

53. Kording, K. P., Tenenbaum, J. B. & Shadmehr, R. The dynamics of memory as a consequence of

optimal adaptation to a changing body. *Nat. Neurosci.* **10**, 779–786 (2007).

54. Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R. & Ivry, R. B. Invariant errors reveal limitations

in motor correction rather than constraints on error sensitivity. *Commun Biol* **1**, 19 (12/2018).

55. Wei, K. & Körding, K. Relevance of Error: What Drives Motor Adaptation? *J. Neurophysiol.* **101**, 655–

664 (02/2009).

56. Tsay, J. S. *et al.* The effect of visual uncertainty on implicit motor adaptation. *J. Neurophysiol.* **125**,

12–22 (2021).

57. Hutter, S. A. & Taylor, J. A. Relative sensitivity of explicit reaiming and implicit motor adaptation. *J.*

*Neurophysiol.* **120**, 2640–2648 (2018).

58.    Tsay, J. S., Ivry, R. B., Lee, A. & Avraham, G. Moving outside the lab: The viability of conducting

sensorimotor learning studies online. *Neurons, Behavior, Data analysis, and Theory* (2021)

doi:10.51628/001c.26985.

59.    Medina, J. F. & Lisberger, S. G. Links from complex spikes to local plasticity and motor learning in the

cerebellum of awake-behaving monkeys. *Nat. Neurosci.* **11**, 1185–1192 (2008).

60.    Herzfeld, D. J., Kojima, Y., Soetedjo, R. & Shadmehr, R. Encoding of error and learning to correct that

error by the Purkinje cells of the cerebellum. *Nat. Neurosci.* **21**, 736–743 (5/2018).

61.    Hadjiosif, A. M., Morehead, J. R. & Smith, M. A. A double dissociation between savings and long-term

memory in motor learning. *PLoS Biol.* **21**, e3001799 (2023).

62.    McDougle, S. D., Bond, K. M. & Taylor, J. A. Explicit and Implicit Processes Constitute the Fast and

Slow Processes of Sensorimotor Learning. *Journal of Neuroscience* **35**, 9568–9579 (七月 1 2015).

63.    De Zeeuw, C. I. & Ruigrok, T. J. Olivary projecting neurons in the nucleus of Darkschewitsch in the cat

receive excitatory monosynaptic input from the cerebellar nuclei. *Brain Res.* **653**, 345–350 (1994).

64.    Turecek, J. & Regehr, W. G. Cerebellar and vestibular nuclear synapses in the inferior olive have

distinct release kinetics and neurotransmitters. *Elife* **9**, (2020).

65.    Bengtsson, F. & Hesslow, G. Cerebellar control of the inferior olive. *Cerebellum* **5**, 7–14 (2006).

66.    Tsay, J. S., Irving, C. & Ivry, R. B. Signatures of contextual interference in implicit sensorimotor

adaptation. *Proc. Biol. Sci.* **290**, 20222491 (2023).

67.    Collins, A. G. E. *The Tortoise and the Hare: Interactions between Reinforcement Learning and Working

Memory*. http://biorxiv.org/lookup/doi/10.1101/234724 (2017).

68.    Collins, A. G. E. & McDougle, S. D. Context is key for learning motor skills. *Nature* vol. 600 387–388

(2021).

69.    Herzfeld, D. J. & Shadmehr, R. Cerebellum estimates the sensory state of the body. *Trends in cognitive

sciences* vol. 18 66–67 (2014).

70. Gonzalez Castro, L. N., Hadjiosif, A. M., Hemphill, M. A. & Smith, M. A. Environmental consistency

determines the rate of motor adaptation. *Curr. Biol.* **24**, 1050–1061 (2014).

71. Negrello, M. *et al.* Quasiperiodic rhythms of the inferior olive. *PLoS Comput. Biol.* **15**, e1006475

(2019).

72. Loyola, S. *et al.* How inhibitory and excitatory inputs gate output of the inferior olive. *Elife* **12**, (2023).

73. Op de Beeck, H. P. & Baker, C. I. The neural basis of visual object learning. *Trends Cogn. Sci.* **14**, 22–

30 (2010).

74. Narain, D., Remington, E. D., Zeeuw, C. I. D. & Jazayeri, M. A cerebellar mechanism for learning prior

distributions of time intervals. *Nat. Commun.* **9**, 469 (2018).

75. Wang, T. *et al.* A unitary mechanism underlies adaptation to both local and global environmental

statistics in time perception. *PLoS Comput. Biol.* **19**, e1011116 (2023).

76. Ito, M. & Kano, M. Long-lasting depression of parallel fiber-Purkinje cell transmission induced by

conjunctive stimulation of parallel fibers and climbing fibers in the cerebellar cortex. *Neurosci. Lett.*

**33**, 253–258 (1982).

77. Marr, D. & Thach, W. T. A theory of cerebellar cortex. in *From the Retina to the Neocortex* 11–50

(Birkhäuser Boston, Boston, MA, 1991).

78. Best, A. R. & Regehr, W. G. Inhibitory regulation of electrically coupled neurons in the inferior olive

is mediated by asynchronous release of GABA. *Neuron* **62**, 555–565 (2009).

79. Shadmehr, R. & Mussa-Ivaldi, F. A. Adaptive representation of dynamics during learning of a motor

task. *J. Neurosci.* **14**, 3208‑3224 (1994).

80. Vindras, P., Desmurget, M., Prablanc, C. & Viviani, P. Pointing Errors Reflect Biases in the Perception

of the InitialHand Position. *J. Neurophysiol.* **79**, 3290–3294 (1998).

81. Shea, J. B. & Morgan, R. L. Contextual interference effects on the acquisition, retention, and transfer

of a motor skill. *J. Exp. Psychol. Hum. Learn.* **5**, 179–187 (1979).

82.  Tsay, J. S., Irving, C. & Ivry, R. B. Signatures of contextual interference in implicit sensorimotor

adaptation. *bioRxiv* (2022) doi:10.1101/2022.07.03.498608.

1039

1040 **Supplementary Information**

1041

1042 <u>Supplemental Result 1: The CPC model accounts for contextual interference.</u>

1043

The two-layer model provides an alternative explanation for another type of context-dependent learning,

contextual interference. The term is a bit of a misnomer since the phenomenon refers to the fact that,

while performance gains when training in multiple contexts is slower compared to training in a single

context, retention is better in the former[69,70]. As such, exposure to multiple contexts during training

actually enhances learning as measured by long-term gains. Interestingly, this phenomenon is not limited

to skill acquisition tasks but is also observed in studies of implicit adaptation[81] (Fig S9).

1050

In the revised CPC model, contextual interference occurs due to the parallel operation of volatile and

stable learning processes. With multiple targets (constituting multiple contexts), the rate of acquisition is

slower compared to a single target since learning from the volatile process decays between successive

reaches to a given target. However, early retention is higher since the contribution of the volatile process

is small. Thus, as with anterograde interference, contextual interference arises from the dynamics of the

system without postulating any representation of context.

1057

1058

1059

1060

1061
1062

1063 **Fig. S1 Predictions of the Dual State Space model for Exp 2-3.** The time course of the predicted hand

1064 angle as well as the underlying states of the fast and slow processes are shown.

1065

1066

1067

**Fig. S2 Effect of experience and error consistency on implicit adaptation. a)** Attenuation in relearning in

Exp 3. Adaptation was attenuated in response to re-exposure to a perturbation compared to the initial

exposure (t(33)=3.1, p=0.004) Data are averaged across each training phase. **b)** Spontaneous recovery was

not observed in Exp 3 during the no-feedback phase after washout. Hand angle over the first 5 trials of

the no-feedback phase (Early) is similar to hand angle over the last 5 trials (Late, t(33)=1.2, p=0.23). c)

Error consistency did not affect adaptation during initial learning and during relearning in Exp 4. A mixed

ANOVA showed a main effect of learning/relearning, (F(1,101)=37.7, p<0.001), similar to the antegrade

interference observed in Exp 6. There was no effect of error consistency (F(2,101)=0.18, p=0.84) or

interaction between phase and error consistency (F(2,101)=0.12, p=0.88). Box plots indicate median, max

and min values, and 25% and 75% quartiles.

1078

1079

1080 **Fig. S3 Predicted time course of stable and volatile processes in Exps 2-4 and 6.** The stable process is

1081 responsible for anterograde interference (a) and attenuation in relearning (b-c). The volatile process does

1082 not make a significant contribution to either phenomenon because of its low retention rate.

1083

1084

1085

1086 **Fig. S4 Retention increases during the initial washout trials. a)** To provide a stronger test of how the rate

1087 of retention changes (Exp 1), Exp 7 included mini-blocks (10 trials/mini-block) that alternated between

1088 clamp and no feedback trials. **B**) We estimated the change in retention rate over time by averaging by trial

1089 number across the no feedback blocks. Retention is relatively low in the first trials of the washout block

1090 and gradually rises (F(6,264)=4.64, p<0.001). The dark green curve shows the fit of the CPC model.

1091

1092

1093

**Fig. S5 Revised CPC with DCN-IO inhibitory pathway.** The original CPC predicts that the asymptote should

be lower in the long ITI condition compared to the short ITI condition because the latter includes a volatile

component. However, as shown in Fig 7, the asymptote is similar in the two ITI conditions. This

observation motivated a revision to the CPC model in which the DCN sends a inhibitory signal to the

inferior olive. **a)** Model schematic. DCN-IO inhibition suppresses the error signal to the DCN and cerebellar

cortex. This suppression is generic given that the output of the DCN integrates activation across

directionally tuned units. **b)** When the inter-trial-interval is short, the CS response is suppressed (top).

Note that the suppression is implemented by subtracting a common value to the IO and thus alters the

activation in PCs. On the next trial, SS activation is stronger in the long ITI condition since the PF-PC

synapse will have recovered during the ITI (middle). However, there are a subset of tuned elements that

in which SS activation is weaker in the long ITI condition (yellow arrows). This weaker activation induces

adaptation in DCN units tuned to the same direction (bottom). **c)** State of the volatile and stable processes

over the course of a fixed design under long and short ITI conditions. The change in the volatile process is

smaller in the long ITI condition due to forgetting. The stable process is also smaller in the long ITI

condition because SS activity at the preferred error direction will dominate learning. However, the long

ITI condition induces adaptation in neurons with sub-preferred error directions, resulting in larger

adaptation late in training.

1111

1112

1113  **Fig. S6 State-space models fails to explain the learning asymptote. a)** A single state motor cannot account

1114  for fast early learning and slow forgetting. **b)** The state-space model assumes the asymptote reflect a

1115  balance between learning and forgetting. As such the asymptote will drop to a half in the half washout

1116  phase of Exp 6. However, there is only a slight decrease in the asymptote during the half washout,

1117  consistent with the predictions of the CPC model rather than the dual SS model.

1118

1120

**Fig. S7 Contribution of stable and volatile processes in response to variable perturbations. a)** The stable process (top) contributes to learning during early training and has saturated by the 50th trial. The contribution of the volatile process (bottom) remains similar throughout training. **b)** Change in hand angle as a function of trial number when the size and direction of the perturbation varies across trials. The change of hand angle is larger in early training because the stable process has not saturated. **i)** As predicted by the two-process CPC model, when exposed to a variable perturbation, the Δhand is larger in early training compared to late training. Shaded areas and error bars indicate standard error.

1128

1129

1130

Fig. S8 Revised CPC model provides a good fit for the key results for all of the experiments. Dark green

line depicts model prediction. Error bars (c, g, h, i) and shaded areas (a, b, d, e) indicate standard error.

1131

1132

1133

1134

1135

**Fig. S9 Revised CPC model accounts for effect of number of target locations on adaptation. a)** In Tsay et

al.[82], participants were trained with either one target or three targets. In both conditions, participants

reached to a single target during the washout block. **b)** Learning functions for the target location probed

during washout. The 3-target condition showed slower learning but a larger aftereffect. Adding more

targets is effectively akin to imposing a long ITI since successive reaches to a given target are separated

by reaches to the other two locations; thus, there is more forgetting but stronger retention due to reduced

contribution of volatile process. Shaded area in b indicates standard error. Dash lines indicate the

predictions of the Revised CPC model.