Variations in bacterial community at Chlorophyll Maximum (C-Max) depths along the

west coast of India due to seasonal changes in the primary productivity

Short title: Bacterial dynamics in Chlorophyll Maximum depths

Ashutosh Shankar Parab[1, 2], Mayukhmita Prithunandan Ghose[1] Cathrine Sumathi Manohar[1]*

[1]Biological Oceanography Division, CSIR- National Institute of Oceanography, Dona Paula,

Goa, 403004. India.

[2]School of Earth, Ocean and Atmospheric Sciences, Goa University, Taleigao Plateau, Goa,

403206. India.

*Corresponding author e-mail: cathrine@nio.org, ORCID: 0000-0003-0178-7784

Abstract

An increase in primary productivity is recorded annually during the southwest monsoon season along the west coast of India, an important upwelling zone. The influence of the seasonal variations in the *in situ* primary productivity on the bacterial dynamics and community structure was explored during the non-monsoon and productive monsoon seasons. In the monsoon season, shallower mixed layer depth, increased nutrient concentration and a significant, positive correlation of bacterial carbon ($p < 0.01$) with primary productivity was observed. Bacterial diversity was assessed in the chlorophyll maxima depths during both seasons based on next-generation, metagenomic analysis. In the non-monsoon season, genera such as *Idiomarina*, *Salinimonas*, *Marinobacter* of Proteobacteria and *Bacillus* and *Lactobacillus* of Firmicutes were dominant. These major bacterial genera are shown through canonical correspondence analysis to play an important ecological role. They could be responsible for the increased heterotrophic activity recorded through predicted functional gene profiles in this season. In the monsoon season, increased abundance in the autotrophic Cyanobacteria community and its photosynthetic activity was recorded in the gene profiles. Higher diversity of heterotrophic phyla such as Bacteroidetes, Actinobacteria and a few Candidatus phyla and changes in the diversity of Proteobacteria with a representation of *Alteromonas*, SAR86 clade and OM60 (NOR5) was observed. These results highlight the bacterial dynamics associated with seasonal variations in primary productivity along the west coast of India.

Keywords: Bacterial diversity, community structure, functional activity, metagenomics, next-generation sequencing, monsoon, non-monsoon, west coast of India.

1. Introduction

Upwelling plays a crucial role in the biogeochemical processes, as it provides essential nutrients for phytoplankton growth, which in turn supports the marine food chain (Capone et al., 2013). Along the west coast of India (WCI), upwelling occurs annually from June to September, during the southwest monsoon season (Smitha et al., 2008; Gupta et al., 2016). The upwelling in this region is driven by the strong, persistent winds that blow parallel to the coast, which create a coastal upwelling front (Schott et al., 2009). These winds push the surface waters away from the coast, allowing the colder, nutrient-rich waters from the deeper ocean to rise to the surface (Vinayachandran et al., 2021; Gupta et al., 2021). This brings in nutrient-rich water to the surface that is abundant in nutrients such as nitrates and phosphates, which leads to an increase in phytoplankton productivity. Subsequently, this results in the generation of substantial amounts of organic matter along the WCI during the southwest monsoon season (Kumar et al., 2000; Ramaiah et al., 1996; Habeebrehman et al., 2011). Bacteria, which are referred to as the "Unseen Majority" play crucial roles in the biogeochemical cycling of organic matter and are therefore fundamental for the operation of marine ecosystems (Whitman et al., 1998). The heterotrophic bacteria through the "microbial loop" break down the particulate organic matter into the dissolved organic matter that sustains the marine food web and nutrient cycling (Pomeroy et al., 2007). Thus providing necessary nutrients for the primary production of phytoplankton, which forms the basis of the aquatic food web (Wiggert et al., 2005; Azam and Malfatti, 2007). Evaluating the processes that influence the structure of bacterial communities on spatial and temporal scales is essential for understanding the recycling of organic matter in marine ecosystems (Danovaro et al., 2017).

Bacteria play critical roles in regulating global nutrient cycles and primary production

(Baltar et al., 2015). Therefore, any changes in their abundance or activity as well as the

surrounding environmental changes can have significant effects on the functioning of marine

ecosystems. Initial studies of the bacterial community analysis from major upwelling regions

across the globe, was carried out based on conventional cloning and PCR-based methods, has

studied specific bacterial groups and diversity in the various upwelling regions have shown

seasonal variations in dominant bacterial communities, including Alphaproteobacteria,

Gammaproteobacteria, Bacteroidetes and certain Planctomycetes (Kuypers et al., 2005; Cai et

al., 2007; Alonso-Gutiérrez et al., 2009). This signified their involvement in biogeochemical

cycling and highlighted the impact of environmental variations on the bacterial diversity.

However, studies carried out based on recent Next-generation sequencing (NGS) technologies

have provided valuable advancements in understanding the bacterial communities in upwelling

regions, allowing in-depth examination of the bacterial dynamics (Jing et al., 2013; Bergen et

al., 2015; Thiele et al., 2019; Liu et al., 2022). These studies have found bacterial communities

including Proteobacteria, Cyanobacteria, Bacteroidota, Firmicutes, Actinobacteria and certain

previously unidentified Candidatus phyla prevalent in diverse upwelling regions, highlighting

their ecological significance. NGS technologies have significantly enhanced the understanding

of the active members within bacterial communities, revealing the ecological significance of

rare phylotypes such as SAR11, Alteromonadales and Vibrionales in their response to

environmental changes (Gutiérrez-Barral et al., 2021; Mena et al., 2022). Additionally, the

impact of upwelling on microbial community composition has also been observed at C-Max

depths (Gong et al., 2022). Collectively, these studies underscore the variability and complexity

of bacterial community structure and abundance within upwelling zones shaped by distinct

environmental factors and seasons.

Studies on bacterial communities along the WCI have primarily focused on specific

regions, such as the oxygen minimum zones (Bandekar et al., 2018; Fernandes et al., 2020;

Lincy and Manohar, 2021), the mud bank regions (Anas et al., 2021; Vijayan et al., 2022) and

estuarine habitats (Khandeparker et al., 2017; Kuchi et al., 2021). Studies based on culturable

approaches have shown the influence of monsoonal upwelling and higher productivity on the

bacterial community structure and functional diversity in the estuarine, coastal and open ocean

waters along the WCI (Anas et al., 2021; Vijayan et al., 2022; Parab et al., 2022). Our prior

investigation, on the culturable bacterial diversity along the WCI, showed higher bacterial

abundance during the monsoon season and with the highest colony-forming units (CFUs) from

the C-Max depths (Parab et al., 2022). The C-Max, is a prominent feature of tropical marine

environments occurring between 20-100 m water depth with maximum primary productivity,

due to the optimal physicochemical conditions for the growth of phytoplankton (Cullen, 2015).

These C-Max depths have very high a high availability of organic substrate for bacteria.

However, there are limited research from the upwelling regions from the WCI, on the impact

of varying primary productivity on the bacterial communities from C-Max zones utilising high-

throughput sequencing techniques. To address this knowledge gap, this study aimed to

comprehensively analyse the impact of increased productivity and surrounding environmental

factors on the bacterial dynamics along one of the most productive zone WCI, during non-

monsoon and monsoon seasons, using high-throughput sequencing of 16S rRNA amplicons.

This approach allows an in-depth understanding of the regional bacterial community structure

and its potential seasonal variations. This study provides the first detailed analysis of the

bacterial community and functional diversity associated with seasonal changes in the primary

productivity along the WCI.

2. Methodology

2.1 Study location, sample collection and assessment of physicochemical parameters

Sampling was conducted during oceangraphic cruises onboard RV Sindhu Sankalp during the non-monsoon season in February 2019, Cruise # SSK-125 and in the southwest monsoon season in September 2019, Cruise # SSK-131 along the WCI. Samples were collected between 8°N and 16°N and 72°N to 76°E, at three stations: off Goa, Mangalore and Trivandrum (Fig. S1). At each station, samples were collected from a coastal location at 30 m water depth and an off-shore location at 600 m depth. The sampling locations in the coastal and off-shore regions of Goa are designated as G30 and G600 respectively. Similar to this, sampling locations off Mangalore and Trivandrum stations are designated as M30, M600 and T30, T600 respectively (Fig. S1). At each of the six sampling locations, the conductivity-temperature-depth (CTD) underwater profiler, Sea-Bird, USA, model: SBE-911 plus with sensor accuracy of $\pm$ 0.0003 S/m, $\pm$ 0.001°C and $\pm$ 0.015 % for conductivity, temperature and depth respectively was employed to quantify the physicochemical characteristics. The water temperature, density, salinity, dissolved oxygen (DO) and fluorescence, which represents the concentration of chlorophyll in the water column was measured. In all coastal locations (30 m contour), samples were collected from the (i) surface at about 2 m below the sea surface, (ii) the depth at which maximum concentration of chlorophyll was recorded (C-Max) and (iii) bottom depth at about 3 m above the seafloor. In the off-shore region along the 600 m contour, samples were collected from (i) surface, (ii) C-Max, (iii) 100 m, (iv) 200 m (v) 400 m and (vi) bottom depth. The depth of C-Max was defined based on the readings obtained from the fluorescence sensor during the downward haul of the CTD underwater profiler and samples

were collected at desired depths during the upward haul. Water samples collected from all the

depths were immediately sub-sampled and appropriately stored for subsequent examination.

Nutrient analysis was carried out in an automated continuous flow analyzer (SKALAR 5000,

Netherlands) to measure the levels of nitrate, nitrite, phosphate and silicate, present in water

samples.

2.2. Estimation of chlorophyll and primary productivity

Chlorophyll-a (Chl-a) concentrations in the water column at discreet depths were

determined fluorometrically following a standard protocol of the Joint Global Ocean Flux

Study (JGOFS, UNESCO 1994). For this, 1 L of seawater sample from each depth was filtered

using a 0.4 µm, 47 mm GF/F filter (Whatman, USA) in triplicates. Chl-a was extracted from

the filters using 10 mL of 90 % acetone through overnight incubation at 20°C in the dark. After

the incubation, the samples were vortexed and centrifuged to remove cellular debris. The clear

supernatant was collected and used for the estimation of chl-a concentration on a Trilogy

Laboratory Fluorometer (Turner Designs, USA). The water column chl-a concentration from

the euphotic zone was determined through the integration of chl-a values from surface to 100

m depth and were expressed as mg m^{-2}. Gross primary productivity (PP) was assessed in water

samples from coastal and off-shore locations in both seasons based on a ^{14}C-labelled method

using labeled inorganic C, $NaH^{14}CO_3$ following the standard protocol of the JGOFS (UNESCO

1994). For this, water samples were collected and incubated without any pre-filtration in three

light and one dark, 250 mL polycarbonate bottle (Nalgene, Germany), from all the sampling

depths of the coastal (3 depths) and surface, C-Max, 100 and 200 m depths from the off-shore

locations. Each bottle was supplemented with one ampoule of $NaH^{14}CO_3$, obtained from the

Board of Radiation and Isotope Technology in Mumbai, with a specific activity of 185 kBq.

The bottles were then incubated at the respective water depths using an *in situ* mooring system

for 12 daylight hours. At the end of the incubation, water samples were filtered using GF/F 47

mm diameter, 0.7 m pore size filters (Whatman, USA) and stored in scintillation vials in the

dark till further analysis. For estimation of the radioactivity, filter papers were dissolved

overnight using concentrated HCl followed by incubation in recommended volume of liquid

scintillation cocktail (Sisco Research Laboratory, Mumbai, India). The levels of radioactivity

were assessed on a scintillation counter (PerkinElmer Wallac 1409 DSA, USA) and the values

were expressed as mg C m^{-3} d^{-1}. To determine depth-integrated PP, productivity measurements

up to 100 m depth were taken and integrated to obtain water column productivity and expressed

as mg C m^{-2} d^{-1}.

2.3. Estimation of bacterial productivity and dynamics

To estimate the rate of bacterial productivity (BP) at coastal and off-shore stations, water

samples from all the sampling depths were collected during both seasons and subjected to the

^{3}H-thymidine incorporation method as outlined in the JGOFS protocols (UNESCO 1994).

Following the protocol, water samples (20 mL) in triplicates from each depth were incubated

in the dark with ^{3}H-thymidine at final concentration of aboute 0.6 nM (specific activity 18 Ci

mmole^{-1}, Bhabha Atomic Research Center, Mumbai) for 2 hours at ambient temperature. The

incubation was halted by adding filtered formaldehyde and the samples were filtered using 0.22

µm cellulose acetate, 25 mm diameter filters (Millipore, India). The filters were washed with

cold trichloroacetic acid (10 % w/v) followed by ethanol (70 % v/v) and were stored in

scintillation vials for further analysis. The filter papers were soaked overnight in ethyl acetate

and then a 4 mL of liquid scintillation cocktail was added and radioactivity was quantified using a scintillation counter (PerkinElmer Wallac 1409 DSA, USA). BP was calculated utilizing an average oceanic conversion rate of 2.17×10^{18} cells mol^{-1} of thymidine incorporated (Ducklow in 1993) and the results were expressed as mg C m^{-3} d^{-1}.

Bacterial abundance was estimated using epifluorescence microscopy (Porter and Feig 1980). For this, 20 mL water subsamples from each depth were preserved with buffered formaldehyde to achieve a 2 % final concentration. Triplicate aliquots of the preserved samples, each measuring 2 - 5 mL, were stained with 4'6-diamidino-2-phenylindole (DAPI) to reach a final concentration of 1 μg mL $^{-1}$. These samples were then filtered using black polycarbonate membrane filters with a pore size of 0.22 μm, (Millipore, U.S.A.). The bacterial counts were carried out using an epifluorescence microscope (Olympus BX-51, Japan) and expressed in terms of the number of bacterial cells L^{-1}. The bacterial counts were also converted to bacterial carbon (BC) using the conversion factor of 20 fg cell $^{-1}$ (Ducklow et al., 1995) and reported as mg C m^{-3}. The total plates counts (TPC) of heterotrophic bacteria present in the water samples were also determined based on the spread plate technique on Zobell Marine Agar (ZMA) expressed as colony-forming units (CFUs mL^{-1}) (Parab et al., 2022).

2.4. Bacterial community and functional diversity analysis

2.4.1. Extraction of metagenomic DNA, library preparation and Illumina sequencing

For extraction of metagenomic DNA, 20 L of water samples from C-Max depths at each location were filtered through 0.22 mm Durapore filters (Millipore, USA) using a Peristaltic pump (Millipore, USA) on board the research vessel. The filters were immediately stored in

cryovials at -80°C in a DNA storage buffer containing 0.75M sucrose, 40 mM EDTA and 50

mM Tris (pH 8.3). DNA extraction was carried out using a FastDNA^{TM} SPIN kit (MP

Biomedical, USA) according to the manufacturer's protocol. The quality of the metagenomics

DNA extracted was tested and a sufficient quantity of DNA was utilized for library preparation,

which targeted the 16S rRNA gene V3-V4 region using specific proprietary primers (16S

amplicon PCR forward primer: 5' TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCC

TACGGGNGGCWGCAG 3' and 16S amplicon PCR reverse primer: 5' GTCTCGTGGGCTC

GGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC 3'). Illumina PCR

sequencing adapter sequence used were, Index 1 Read 5′ CAAGCAGAAGACGGCATA

CGAGAT[i7]GTCTCGTGGGCTCGG and Index 2 Read 5′ AATGATACGGCGACC

ACCGAGATCTACAC [i5] TCGTCGGCAGCGTC. (Qu et al., 2018; Bora et al., 2022)The

16S rRNA gene libraries generated were sequenced on NovaSeq 6000 v1.5 Illumina

sequencing platform (Illumina Inc., USA) at the National Institute of Biomedical Genomics,

West Bengal, India. The sequences of the amplicons obtained from the C-Max depths from all

the locations during both seasons were processed with a standard pipeline using the QIIME2

tool (Bolyen et al., 2019). The raw reads were initially quality filtered and primers were

removed using Cutadapt v4.2. After quality filtration, the ASV file was generated using the

DADA2 tool v3.16 and taxonomic profiling of the operational taxonomic units (OTU) was

carried out against the V3-V4 classifier.

2.4.1. Functional diversity analysis

Functional diversity of bacterial communities was derived from marker gene sequences

using the PICRUSt2 software, version 2.4.1. These acquired gene profiles were subsequently

categorized into KEGG pathways (Douglas et al. 2020). The PICRUSt tool facilitated the

prediction of potential functionalities from 16S rRNA sequence data. Employing the KEGG

database, closed reference OTU picking was performed against the Greengenes taxonomy. The

predicted functional profiles with gene profiles further grouped into KEGG pathways,

following to PICRUSt protocols. Additionally, to measure the variance between reference

genomes and the inferred metagenome, NSTI (Nearest Sequenced Taxon Index) scores were

determined.

2.5. Data analysis

Statistical analyses were performed using SPSS, to investigate the significance of the

distribution between physicochemical and biological characteristics in the two seasons using a

single-factor analysis of variance (ANOVA). Pearson's correlation analysis was employed to

interpret relationships within the physicochemical and biological parameters between the two

seasons, at a significance level of 5 % ($P < 0.05$). Alpha diversity indices and Beta diversity

between two seasons at OUT level were calculated using Non-metric Multidimensional Scaling

Analysis (NMDS) based on Bray–Curtis distance using the R package Phyloseq v1.22.3

(McMurdie and Holmes, 2013). A significance in the distribution of bacterial community

between non-monsoon and monsoon season were estimated using STAMP bioinformatics

software v2.1.3 (Parks et al., 2014). Heatmap of bacterial communities at a class level was

plotted using hierarchical clustering analysis based on the Euclidean similarity index in R

package Pheatmap v1.0.12. The R packages: ggplots2 and were used for the visualization of

the data. All these analyses were performed in R studio v3.5.1 (R Core Team, 2015). Bacterial

community structure at the genera level, along with environmental variables of the non-

monsoon and monsoon season, were analysed based on canonical correspondence analysis

(CCA) using PAST software v3.23 (Hammer et al., 2006).

3. Results

3.1. Physicochemical parameters

The physicochemical parameters varied significantly between the non-monsoon and

monsoon seasons. During the non-monsoon season, the temperature in the water column

remained constant at coastal stations. In the monsoon season, the temperature decreased by 2

to 4 °C from the surface to the bottom of the water column at coastal stations. In the off-shore

stations, the thermocline depth was observed between 80 and 100 m during the non-monsoon

season and it was shallower at 25 to 50 m, during the monsoon season (Fig. 1). During the non-

monsoon season, halocline depth along the off-shore stations was observed at 80 ± 20 m, which

decreased to 25 to 50 m during the monsoon season similar to the thermocline depth (Fig. 1).

The influence of salinity on the water column characteristicts was also studied based on BVF

from both the seasons, the influence of reduced salinity on the BVF was evident during the

monsoon season (Fig. S2). The average sea surface DO concentration at the coastal and off-

shore locations was 4 ± 2 mg L^{-1} during the non-monsoon and monsoon season. In the coastal

stations at 30 m, DO values were consistent throughout the water column during the non-

monsoon season and in the monsoon season, DO values dropped to < 1 mg L^{-1} at mid-depth

and nearly reached anoxic conditions at the bottom depths. In the off-shore stations (600 m

depth) DO concentration reached close to anoxic conditions at the mixed layer depth (MLD)

at about 80 -100 m in the non-monsoon and monsoon seasons (Fig. 1).

Fluorescence values recorded during the non-monsoon season throughout the water

column was an average ~ 0.4 ± 0.3 mg m^{-3} and it reached a maximum of around 1 mg m^{-3} at

the C-Max depth. In the monsoon season, the average value of fluorescence in the water column

was ~ 0.9 ± 1 mg m^{-3} and it reached a maximum of ~ 6 mg m^{-3} at the C-Max depth (Fig. 1). C-

Max depths in the water column varied during both seasons, which were about 15 m and 55 m

during the non-monsoon season in the coastal and off-shore stations respectively. In the

monsoon season, C-Max in coastal and off-shore stations was observed at approximately 10 m

and 30 m respectively. Higher nutrient concentration was observed in the monsoon season in

both the coastal and off-shore stations. The nitrate concentration, measured at all stations

during the non-monsoon season, had an average of around 2.55 ± 2.47 µM, which increased to

~ 17.37 ± 16.14 µM in the monsoon season (Fig. 2). The average nitrite concentration along

all stations was ~ 0.16 ± 0.2 µM during the non-monsoon season and increased in the monsoon

season to ~ 0.53 ± 0.85 µM. The average phosphate concentration in the water column during

the non-monsoon and monsoon seasons was ~ 3 ± 2.8 µM and 0.92 ± 0.73 µM respectively

(Fig. 2). The average silicate values detected during the non-monsoon and monsoon seasons

were ~10.83 ± 7.30 µM and ~18.93 µM, respectively.

3.2. Primary productivity and bacterial dynamics

Chl-a concentration estimated fluorometrically from discreet water depths was ranging

from 0.002 to 0.16 mg m^{-3} in the non-monsoon and monsoon season it was estimated to be

0.003 to 1.05 mg m^{-3} (Table S1) Column integrated chl-a in the euphotic zone calculated for

coastal and off-shore stations was observed to be 5.64 ± 2 mg m^{-2} in the non-monsoon and it

increased to 11.42 ± 6 mg m^{-2} during the monsoon seasons (Fig. 3a). The PP in the water

column during the non-monsoon season varied between 1.20 to 145.38 mg C m^{-3} d^{-1}, while

during the monsoon season, it ranged from 29.62 to 266.24 mg C m^{-3} d^{-1} (Table S1). The

column-integrated PP in the euphotic zone was 800.31 ± 538 mg C m^{-2} during the non-monsoon

season and the PP estimated during the monsoon season was higher and estimated to be

3355.60 ± 1608 mg C m^{-2} (Fig. 3b). This indicated an upwelling-induced increase in the

primary productivity along the WCI during the monsoon season.

The bacterial activity in both seasons was estimated by quantifying the bacterial

abundance and productivity. The estimated BP did not vary much between the seasons which

was about 10 ± 2 mg C m^{-3} d^{-1} in the coastal and off-shore stations (Table 1). Bacterial counts

were determined microscopically and were expressed as bacterial carbon, which was estimated

to be 209.88 ± 47 mg C m^{-3} during the non-monsoon season and it increased to an average of

1017.62 ± 132 mg C m^{-3} in monsoon season (Table 1).

TPC was enumerated to estimate the viable bacterial counts in the water column from

all the stations. An average of 4162.30 CFU mL^{-1} in the non-monsoon season was recorded

and it increased to 11392.59 CFU mL^{-1} during the monsoon season (Fig. 4). The highest

bacterial CFUs were obtained from the C-Max depths from almost all the locations sampled

along the WCI (Fig. 4). An increasing trend of the bacterial carbon biomass and CFUs was in

accordance with the higher primary productivity observed during the monsoon season.

Statistical analysis based on the ANOVA test was carried out to evaluate the significant

changes in different parameters during both seasons (Table 2). Estimated environmental

parameters including DO, fluorescence, nitrate, nitrite, phosphate, silicate, chl-a, PP, TPC and

BC showed significant variations between seasons ($p < 0.05$). Correlation between

environmental variables was also determined using Pearson's correlation matrix for the physicochemical and biological variables measured from water samples during non-monsoon and monsoon seasons (Table 3). A significant correlation was observed between the different environmental variables estimated during both seasons. Temperature and DO had a significant positive correlation with fluorescence. *In situ* fluorescence concentration determined showed a significant positive correlation with PP and BC. Temperature and DO showed a significant negative correlation with nitrate and silicate. Among the nutrients, nitrate and silicate were positively correlated and Chl-a and PP showed a positive correlation with TPC and BC respectively (Table 3). These results indicate temperature has a significant impact on the fluorescence values which is a measure of the phytoplankton abundance. A significant positive correlation between the primary productivity variables and bacterial abundance and counts shows the influence of the former on the latter in the study area.

3.4. Alpha and Beta diversity of bacteria from the C-Max along the WCI

The bacterial diversity from the stations along the WCI during the non-monsoon and monsoon season was assessed from the C-Max depths based on metagenomic analysis, using next-generation Illumina sequencing techniques. Sequence statistics show that the number of reads generated was nearly similar during both seasons, however, OTUs generated after denoising the sequence reads were higher in the monsoon season (Table S2). The rarefaction curves, based on observable features, were curved toward the saturation plateau at the 97 % cut-off value, indicating that the sampling sizes were enough for the community analysis (Fig. S3). The average OTU abundance in the non-monsoon season was 646 OTUs and it increased in the monsoon season, to an average of 2286 OTUs. In comparison to other stations, higher

OTU abundance was seen in the Trivandrum coastal and off-shore locations during the

monsoon season (Table S2). For both non-monsoon and monsoon seasons, various alpha

diversity indices for diversity and richness were calculated. Shannon, Chao1 and ACE indices

were significantly higher during the monsoon season. The coastal stations showed considerably

higher Shannon, Chao-1 and ACE diversity indices than the off-shore stations (Fig. 5a).

Statistical analysis of the diversity indices calculated shows significant seasonal differences

(Table S3).

Beta diversity analysis of bacterial communities during non-monsoon and monsoon

seasons was performed based on the NMDS method. NMDS ordination differentiates the

samples into abundance-weighted community compositions based on the Bray-Curtis distance.

The bacterial communities were grouped based on the seasons and locations and NMDS

analysis showed that sampling locations during the non-monsoon season were divided into two

apparent groups (Fig. 5b). The bacterial communities from the non-monsoon season of the Goa

and Mangalore locations clustered together and a similar grouping was evident in the

Trivandrum coastal and off-shore communities. A similar trend was observed in the monsoon

season, where the Trivandrum station exhibited a different level of diversity than the Goa and

Mangalore coastal and off-shore stations (Fig. 5b).

3. 5. Bacterial community structure at C-Max along the WCI

Bacterial OTUs generated in Illumina sequencing were classified based on the Silva

V3-V4 classifier for taxonomic profiling. For classification in this study, the sklearn classifier

within QIIME was utilized, which operated on a confidence score range of 0.7 to 1. Based on

taxonomic annotation, OTUs generated during both seasons belonged to 36 bacterial phyla in

total, of which 26 phyla were present during the non-monsoon season and the number of phyla increased during the monsoon season to 35 (Fig. 6a). In the non-monsoon season, 18 of the 26 phyla belonged to well-defined (Based on List of Prokaryotic names with Standing in Nomenclature (LPSN), 7 of it were of candidatus status and 1 unknown or uncharacterized phyla were identified. During the monsoon season, the number of well-defined phyla increased to 22, candidatus to 12 and one unknown phyla were reported (Fig. 6a). The dominant phyla observed in the C-Max depth in the non-monsoon season were Proteobacteria (59 %) and Firmicutes (33 %), while in the monsoon season, it was Cyanobacteria (49 %) followed by Proteobacteria (32 %) (Fig. 6b). The proportion of Cyanobacteria communities increased significantly from 3 % in the non-monsoon season to 49 % during the monsoon season. Though the relative abundance of phylum Proteobacteria decreased during the monsoon season, it remained the second most dominant phyla. The relative abundance of phylum Firmicutes decreased considerably from 33 to 1 % during monsoon season. Bacteroidetes, Actinobacteria, Planctomycetes and Verrucomicrobia were also among the top 10 phyla observed during both the seasons and their abundance increased by two to three folds during the monsoon season. The temporal differences observed in dominant bacterial communities during non-monsoon and monsoon seasons were statistically significant ($p < 0.001$) (Fig. 6c). Among the well-defined phyla, only one phylum, Halanaerobiaeota was specific to the non-monsoon season.

However, about 5 bacterial phyla including Nitrospirae, Lentisphaerae, Kiritimatiellaeota, Fibrobacteres and Fusobacteria were specific only to the monsoon season. There was a marginal increase in the relative abundance of six bacterial phyla, namely Nitrospinae, Spirochaetes, Gemmatimonadetes, Chlamydiae, Acidobacteria and Chloroflexi, during the monsoon season, despite their low abundance levels of less than 0.01 %. The relative abundance of a few phyla, Dianococcus-Thermus, Deferribacters and Epsilonbacteroeota

showed a decreasing trend during the monsoon season. However, the diversity was higher in

the monsoon with an increase in well-defined and candidatus phyla, which accounted for about

7 and 12 phyla from the non-monsoon and monsoon seasons respectively. The candidatus phyla

including Ca. Patescibacteria, Ca. Dependencies, Ca. Schekmanbacteria, Ca. Latescibacteria,

Ca. WPS-2, Ca. Dadabacteria, Ca. Marinimicrobia was found during both seasons. Most of the

candidatus phyla that were commonly identified in both seasons also showed an increase in

their relative abundance during the monsoon season. Ca. Marinimicrobia and Ca.

Patescibacteria had a significantly (p < 0.001) higher contribution in diversity, ranging from

0.5 to 1 %, during the monsoon season (Fig. 6c). Candidatus phyla including Ca. Zixibacteria,

Ca. PAUC34f, Ca. BRC1, Ca. Margulisbacteria and Ca. Hydrogenedentes were exclusively

present during the monsoon season (Fig. 6b). These results suggest that the monsoon season

had a higher diversity of both defined and Candidatus phyla (Fig. 6).

Distribution of the bacterial OTUs at the class level was also investigated using

hierarchical clustering of the heatmap, based on the Euclidian similarity index for the top 5

phyla (Fig. S4a). Within the Proteobacteria phylum, the bacterial class Gammaproteobacteria

was dominant during both seasons, followed by Alphaproteobacteria and Deltaproteobacteria.

Although Gammaproteobacteria was dominant during both seasons, during the monsoon

season, their relative abundance reduced from 48 to 17 % and there was a marginal increase in

the abundance of Alphaproteobacteria and Deltaproteobacteria. The monsoon season was

marked by a higher abundance of phylum Cyanobacteria, which was represented by class

Melainabacteria, Oxyphotobacteria and Sericytochromatia (Fig. S4a). The diversity of phylum

Firmicutes decreased during the monsoon season and distinct differences in the class level were

also observed with the representation of class Bacilli and Clostridia during the non-monsoon

and Erysipelotrichia and Negativicutes during the monsoon season.

The dominant classes of phyla that increased during the monsoon season are

represented by Acidimicrobiia, Actinobacteria, Coriobacteriia, Nitriliruptoria, Rubrobacteria,

Thermoleophilia and an unknown class of Actinobacteria; and class Bacteroidia, Ignavibacteria

and Rhodothermia of phylum Bacteroidetes (Fig. S4a). The other predominant representatives

were classes Planctomycetacia and Verrucomicrobiae of phylum Planctomycetes and

Verrucomicrobia respectively in both the seasons (Fig. S4b). A few classes belong to the phyla

Planctomycetes and Candidatus phyla Marinimicrobia (SAR406 clade), namely Ca.

Patescibacteria and Ca. Margulisbacteria were exclusively present during the monsoon season.

A significant difference in the distribution of the bacterial classes was observed between both

seasons, with higher diversity in the monsoon season (Fig. S4b). To further investigate the

community structure of bacterial populations, bacterial communities were categorized at order

and family levels (Table S4). During the non-monsoon season, the most abundant bacterial

family was Moraxellaceae, belonging to the order Pseudomonadales of Gammaproteobacteria.

During the monsoon season, order Synechococcales of the family Cyanobiaceae belonging to

phylum Cyanobacteria was the predominant autotrophic community. The most abundant

heterotrophic bacterial community at order level in this season was Flavobacteriales from

phylum Bacteroidetes and SAR86 clade from Gammaproteobacteria (Table S4).

3.6. Correlation between environmental variables and bacterial communities

The influence of environmental and primary productivity parameters on bacterial

community structure at the genera level was evaluated for both the non-monsoon and monsoon

seasons using CCA (Fig. 7). For the analysis, bacterial genera with a relative abundance of 0.5

% and greater were plotted along with temperature, salinity, DO, fluorescence, chl-a and PP

measured during both seasons. Based on the CCA ordination results, the environmental factors

considered in this study accounted for 94.21 % of the variation in the community during the

non-monsoon season. In this season, the bacterial genera *Cyanobium PCC-6307*, belonging to

Cyanobacteria; *Bacillus* from phylum Firmicutes, showed a significant correlation with

temperature, DO and chl-a. The primary productivity parameters fluorescence and PP had a

major influence on the bacterial genera largely belonging to Proteobacteria, including,

Idiomarina, *Alcanivorax*, *Alteromonas*, *Salinimonas*, *Marinobacter*, *Acinetobacter* from

Gammaproteobacteria; Alphaproteobacteria including *Rhodobacteraceae_uncultured,*

Sphingomonadaceae_uncultured and *Sulfitobacter* (Fig. 7a). One of the genera *Lactobacillus*

from phylum Firmicutes was also positively associated with fluorescence concentration during

the non-monsoon season. Environmental variables account for 87.01 % of the variation in the

bacterial community during the monsoon season. PP and fluorescence were the most significant

contributors to the community variation during this season (Fig. 7b). Majority of bacterial

genera belonging to Proteobacteria including Ectothiorhodospiraceae_uncultured, SAR86

clade, sOM60 (NOR5) clade from class Gammaproteobacteria, SAR11 clade_Clade I,

Rhodobacteraceae_uncultured, AEGEAN-169 marine group, *Euzebya* from class

Alphaproteobacteria were highly correlated with the primary productivity parameters during

the monsoon season. PP and fluorescence had also shown a positive influence on bacterial taxa

from phylum Bacteroidetes including NS4 and NS5 marine groups and Candidatus

Marinimicrobia (SAR406 clade) during the monsoon season. The bacterial genera Sva0996

marine group Ca. *Actinomarina*, from phylum Actinobacteria; OCS116 clade,

*Sphingomonadaceae_*uncultured from Gammaproteobacteria and *Alteromonas* from

Alphaproteobacteria, was strongly correlated to temperature (Fig. 7b).

Bacterial genera *Synechococcus CC9902*, *Cyanobium PCC-6307* and *Chloroplast* from

Cyanobacteria were associated with the DO and chl-a concentrations in the C-Max depth.

Overall, the CCA results suggested that DO and chl-a influenced the distribution of the

Cyanobacterial community. The primary productivity parameters fluorescence and PP are seen

to influence the bacterial populations belonging to Proteobacteria which included a large

majority of the Gammaproteobacteria and candidatus taxa. This shows that the variations in

the primary productivity associated with upwelling events have a significant impact on the

bacterial community composition in WCI.

3.7. Functional diversity at C-Max along the WCI

The functional profiles of the bacterial communities showed that the predicted

pathways belonged to a variety of vital classes of functions like metabolism, processing of

genetic and environmental information and pathways for cellular processes during both seasons

(Fig. 8a). Significant variations in the abundance of functional genes were also evident between

the non-monsoon and monsoon seasons. Further investigations into seasonal changes in the

gene profiles show that, during the non-monsoon season, there was a significant increase in the

abundance of metabolic pathways related to carbohydrate metabolism, lipid and amino acid

metabolism, glycan biosynthesis and xenobiotic degradation. Pathways related to energy

metabolism including, carbon fixation by prokaryotes, photosynthesis and membrane transport

were abundant in the monsoon season (Fig. 8b and c). Functional profiling of the pathways

related to the cellular processes revealed that signalling pathways, biofilm formation and

pathways related to cellular prokaryotic communities were abundant during the non-monsoon

season. Pathways related to cell growth and death, cell motility and prokaryotic quorum sensing

were uniformly distributed in both seasons (Fig. 8c). Based on the functional gene profiles, it

is evident that the non-monsoon season was associated with an increased representation of

pathways related to heterotrophic metabolism, however during monsoon season autotrophic

activities were higher.

4. Discussion

4.1. Influence of seasonal primary productivity on the bacterial dynamics

510   Seasonal upwelling occurs along the WCI during the southwest monsoon season which

leads to increased phytoplankton productivity and accumulation of organic matter in this region

(Smitha et al., 2008; Gupta et al., 2021). The availability of organic matter resulting from

phytoplankton blooms during upwelling fosters the proliferation of various bacterial

communities (Rocke et al., 2020). In our study, we investigated the impact of seasonal changes

in primary productivity along the WCI on the bacterial dynamics to understand the differences

in the bacterial communities and their functional profiles during the non-monsoon and

monsoon season. The assessment of water column physicochemical parameters shows

upwelling signatures, characterized by shallow MLD and increased nutrients ($p < 0.01$)

associated with higher concentration of chl-a ($p < 0.05$) (Table 2) in the subsurface waters

during the southwest monsoon season. This season also exhibited a significant increase in

fluorescence ($p < 0.05$) values, indicating higher primary productivity compared to the non-

monsoon season (Table 2). Throughout the monsoon season, nitrate ($p < 0.01$) and phosphate

($p < 0.01$) levels in the water column were significantly high and this causes an increase in

phytoplankton productivity. *In situ* estimation of PP, as well as chl-a concentration from the

study region, also indicate the presence of higher phytoplankton productivity during monsoon

season. The organic matter produced by primary producers drives the microbial carbon pump

and contributes to the overall energy flow and nutrient cycling (Jiao et al., 2010; Buchan et al.,

2014). Increased organic load also causes a decline in the DO levels and a significant increase

in nitrite concentration ($P < 0.01$). This was observed during the monsoon season, which may

be attributed to denitrification activities of the bacterial community, which are known to occur

along the WCI (Naqvi et al., 2010; Gonsalves et al., 2011). High primary productivity results

in the generation of organic matter, leading to an increase in microbial activity for sustaining

the food web and ecosystem functioning (Habeebrehman et al., 2008; Zingone et al., 2019;

Rixen et al. 2019). Studies have suggested that high primary production is typically promoted

by elevated nutrient levels associated with recently upwelled shelf waters, which are usually

accompanied by a surge in bacterial abundance (Gauns et al., 2005; Lamont et al., 2014; Reji

et al., 2020).

Results from this study region on the bacterial dynamics associated with changes in the

primary productivity along the WCI showed an increase in viable bacterial counts during the

monsoon season in the C-Max depths (Fig. 4). This suggests strong dependence of the bacterial

community on organic matter derived from phytoplankton (Molina et al., 2020). Bacterial

carbon estimated based on abundance value increased significantly during monsoon season and

showed a positive correlation with primary productivity measurement (Table 1). These findings

align with previous research conducted in the Arabian Sea, which also indicates that the higher

levels of phytoplankton growth during the monsoon season have an impact on the bacterial

abundance and culturable bacterial communities (Ramaiah et al., 2005; Parab et al., 2022).

These studies on the bacterial dynamics in upwelling regions have also shown that a correlation

between bacterial productivity and primary productivity exists, underlining the crucial role

played by bacteria. However, the characterization of the relationship between seasonal

variations in primary productivity levels and the structure of bacterial communities using

metagenomics tools along the WCI remains unexplored.

4.2. Bacterial community structure and functional profiles at C-Max depths in the non-

monsoon season

The intricate relationship between primary production and bacterial communities has

garnered substantial attention, as bacteria are essential players in carbon cycling and organic

matter recycling within the marine ecosystem. Hence, in this study, we employed metagenomic

approaches to investigate the bacterial diversity associated with seasonal changes in primary

production along the WCI in the C-Max depth. Our results revealed substantial variations in

bacterial diversity and community structure in response to seasonal changes in primary

production. Higher OTU richness and diversity (H) were observed at all the stations sampled

during the monsoon season. Temporal differences in alpha diversity were also evident between

the two seasons and within the locations studied along the WCI (Fig. 5).

Metagenomcc analysis of the bacaterial diversity showed that, during the non-monsoon

season, only 26 of the total 36 phyla were represented and the dominance of Proteobacteria and

Firmicutes followed by Bacteroidetes and Actinobacteria was observed. Within Proteobacteria,

Gammaproteobacteria were the most dominant class represented by major genera such as

Psychrobacter, *Idiomarina* and *Alcanivorax,* followed by a small proportion of

Alphaproteobacteria and Deltaproteobacteria (Fig. 6). Gammaproteobacteria are reported from

various marine ecosystems and the observed increase in their relative abundance can be

attributed to their remarkable ability to adapt and thrive under conditions of limited organic

matter availability (Cho and Giovannoni, 2004). Additionally, these bacteria possess the

capacity to utilize alternative sources of energy for growth (Cho et al., 2007; Chen et al., 2021).

Moreover, Gammaproteobacteria exhibit a unique capability of efficiently utilizing organic

matter through the secretion of extracellular hydrolysis enzymes, enabling them to access and

metabolize complex substrates present in the environment (Spring et al., 2015; Liu and Liu,

2020). Firmicutes were the second dominant phyla in the non-monsoon season represented by

major genera *Bacillus* and *Lactobacillus* and their total abundance decreased significantly in

the monsoon. They are known to be metabolically versatile and can adapt to different

environmental conditions (Siezen and van Hylckama Vlieg, 2011; Kadnikov et al., 2020). The

dominance of this group during the non-monsoon season could be due to their ability to

efficiently utilize and grow in the presence of limited organic matter. These bacteria have

evolved strategies to thrive well in these conditions by employing various mechanisms such as

sporulation, biofilm formation and metabolic versatility (Kadnikov et al., 2020).

During the non-monsoon season, the presence of bacterial phyla such as Actinobacteria,

Bacteroidetes, Planctomycetes, Verrucomicrobia and various candidatus phyla was observed

to be in minor proportions (Fig. 6b). Actinobacteria and Bacteroidetes are known to encompass

diverse microbial taxa with versatile metabolic capabilities (Bunse et al., 2016). However, their

reduced abundance during the non-monsoon season suggests a potential limitation in their

ability to thrive under nutrient-limited conditions (Ho et al., 2017). These bacterial groups may

rely on higher nutrient availability, which is typically associated with periods of increased

primary production, such as during the monsoon season. Furthermore, the presence of various

candidate phyla in low percentages further supports the notion that these bacterial groups may

struggle to flourish under conditions of limited nutrient availability. The minor representation

of these candidate phyla during the non-monsoon season could suggest that they possess

specialized ecological niches or physiological traits that restrict their growth and proliferation

in low-nutrient environments (Yilmaz et al., 2016). This finding exhibits interesting

implications regarding the adaptability of minor bacterial groups to conditions of low

productivity during this season.

To understand the influence of environmental variables and primary productivity on the

bacterial communities, CCA was carried out, which showed two significant correlations in the

non-monsoon season. This analysis showed, that in this relatively less productive season, the

dominant bacterial genera from Gammaproteobacteria and Alphaproteobacteria exhibited a

significant correlation with the fluorescence concentration. The dominant bacterial genera,

from the phylum Cyanobacteria, Firmicutes, Gammaproteobacteria and Bacteroidetes obtained

from this season, exhibited a significant correlation with temperature, dissolved oxygen (DO)

and chl-a levels (Fig. 7a). Environmental variables such as temperature influence the metabolic

rates and growth of these organisms, with optimal ranges facilitating their proliferation

(Mohanty et al., 2022). These metabolic capabilities of the bacterial communities enable them

to thrive in diverse environments and outcompete other bacterial taxa. The physicochemical

factors play an important role in shaping the structure of marine microbial communities and

cause variations at spatial scales (Wang et al., 2019). Recent studies have also shown that

variations in the community composition of marine bacteria as well as patterns of worldwide

bacterial (16S) diversity and community structure are strongly correlated with temperature and

DO (Sun et al., 2015; James et al., 2022).

Seasonal variations in primary productivity exert a significant influence on the

predicted functional profiles of the bacterial communities. The functional diversity of the

bacterial assemblages is known to vary, with evidence suggesting that the upwelling seasons

may exhibit higher levels of functional diversity than other periods. This indicates that during

the upwelling season, the metabolism of microbial communities is particularly active in breaking down complex organic compounds such as carbohydrates, proteins and polysaccharides (Caroppo et al., 2022). However, our previous study based on culturable bacterial populations found that functional activities in the water column during the monsoon season were reduced comparatively than in the non-monsoon season (Parab et al., 2022). Similarly, this study based on metagenomics approaches and predictive gene profiling showed that the heterotrophic metabolic activities were higher during the non-monsoon season (Fig. 8). The abundance of genes involved in membrane transport, carbohydrate metabolism, xenobiotic degradation and amino acid metabolism shows that metabolic exchange and nutrient transformation through extracellular heterotrophic activities were significantly higher during non-monsoon could be due to limited nutrient availability. Genes related to two-component signal transduction systems were also significantly higher during the non-monsoon season when productivity was low. Studies have also reported that bacteria can sense and adapt to diverse environments, stressors and growth conditions and also respond to stimuli like low nutrient and quorum signals (Mitrophanov and Groisman, 2008; Sebastián et al., 2013; Prüß, 2017). Limited nutrient availability can indeed increase heterotrophic activities in bacterial communities. When nutrients are scarce, heterotrophic bacteria like Alphaproteobacteria SAR11 and Flavobacteria are known to adapt by maximizing their resource utilization and metabolic efficiency (Sebastián et al., 2013, 2016). Bacteria may exhibit increased enzymatic activity and cellular activity such as quorum sensing and biofilm formation allowing them to efficiently break down available organic matter and obtain essential nutrients for growth (Lever et al., 2015 Prüß, 2017), which was also evident from our study (Fig. 8c). An increase in functional genes of pathways associated with heterotrophic metabolic activities were found to be more prevalent during the non-monsoon season. This can be linked to the contribution of

the dominant Gammaproteobacteria and Firmicutes communities. Their higher representation

during the non-monsoon season suggests their active involvement in heterotrophic metabolic

activities during the non-monsoon season.

4.3. Bacterial community structure and functional profiles at C-Max depths in the

monsoon season

During the monsoon season, the bacterial diversity was higher with representatives

from 35 phyla belonging to various defined and candidatus phyla. The dominant bacterial

groups during this season were from the autotrophic bacterial phyla, the Cyanobacteria which

increased by three folds during the monsoon season. The significant increase in the abundance

of Cyanobacteria from 3 % during the non-monsoon season to 49 % during the monsoon season

implies that this group of bacteria may have a competitive advantage over other bacterial

groups during the upwelling season. Cyanobacteria are known to form blooms in nutrient-rich

waters and thus increase in their abundance as compared to non-monsoon season (James et al.,

2022). During both seasons, *Synechococcus* and *Cyanobium* were the dominant genera of

Cyanobacteria observed. *Synechococcus* is considered a significant Cyanobacterial genus due

to its important ecological role in oceanic upwelling regions, specifically in carbon cycling and

trophic networks (Dore et al., 2022). Though there was an increase in the abundance and

diversity of bacterial communities in the monsoon season; there was a decrease in the

abundance (from 59 to 32 %) and community shift in the diversity of the most prevalent

heterotrophic bacterial group Proteobacteria. The decrease in the abundance of this major group

was obtained largely due to the reduction in the class Gammaproteobacteria. There was also a

change in the diversity within this group which was represented by *Alteromonas*, SAR86 clade

and OM60 (NOR5) in this season. These genera could be of ecological importance in the

upwelling regions as this class of bacteria is reported from regions of high primary productivity

(González et al., 2000; Alonso-Sáez et al., 2007). Within this phyla, Alphaproteobacteria and

Deltaproteobacteria abundance also increased significantly during the monsoon season (Fig.

S4). Alphaproteobacteria have been identified as particle-degrading microorganisms, which

suggests their potential significance as key contributors to the biogeochemical cycling of

carbon and other nutrients during upwelling (Bachmann et al., 2018). Deltaproteobacterial

abundance has also been shown to be higher in upwelling regions. They exchange nutrients

with Alphaproteobacterial and other bacterial communities through mutualism interactions,

providing inorganic compounds in exchange for organic carbon compounds and thus play

important roles in linking the nutrient cycles (Sheik et al., 2013; Aldunate et al., 2018; Sun et

al., 2022).

The abundance of Bacteroidetes, another prominent phylum observed in this study and

increases significantly during the monsoon season and has been commonly found associated

in upwelling systems with algal blooms (Alonso et al., 2006; Schattenhofer et al., 2009). They

are also important players in the breakdown of particulate organic materials in the water

column (Kirchman, 2002; Pinhassi et al., 2004; Bergen et al., 2015). It was found that

Actinobacteria is a prevalent and widely distributed phylum and its abundance increased from

2 % during the non-monsoon season to 6 % during the monsoon season. The diversity of these

phyla is reported from C-Max depths in other upwelling regions (Morris et al., 2012; Montes

et al., 2020); which are possibly increased in numbers due to the accumulation of

phytoplankton biomass at these depths. Other dominant phyla observed were Verrucomicrobia

and Planctomycetes; known to be frequently prevalent and numerous in marine bacterial

assemblages (Freitas et al., 2012). Thus, during upwelling events, Cyanobacteria and

Proteobacteria tend to dominate, indicating their ecological significance, while a significant
increase in other bacterial groups during upwelling, such as Actinobacteria, Bacteroidetes,
Verrucomicrobia and Planctomycetes contribute to organic matter recycling (Rocke et al.,
2020; Sun et al., 2020; Reji et al., 2020). In addition to the well-defined phyla observed in the
study region, the presence of various candidatus phyla was also detected. Notably, during the
monsoon season characterized by high primary productivity, candidatus phyla including Ca.
Marinimicrobia, Ca. Patescibacteria and Ca. Dadabacteria exhibited a significant increase in
their relative abundance. Their potential roles in the decomposition and utilization of the
abundant organic matter generated during this period highlight their ecological significance in
nutrient cycling and carbon metabolism within the study region (Mestre et al., 2018; Frank et
al., 2016). Further investigaions into the functional potential and ecological contributions of
these candidatus phyla are warranted to elucidate their specific roles in microbial community
dynamics and ecosystem functioning during the monsoon season.

Based on CCA analysis, an association of Cyanobacterial genera including *Cyanobium
and Synecoccoccus* with temperature, DO and chl-a concentrations during monsoon season
was evident, this highlights their sensitivity to environmental conditions (Fig. 7b). Dissolved
oxygen availability is crucial for the survival of aerobic bacteria, including Cyanobacteria
(Berg et al., 2022). The structure of the bacterial assemblages in the marine environment is
subject to temporal and regional change due to oceanographic phenomena like upwelling
(Alonso-Gutiérrez et al., 2009) and also due to environmental factors induced by upwelling
including physicochemical changes and organic matter generation. Physicochemical factors
such as temperature and DO availability play a crucial role in shaping the diversity, abundance
and activity of bacterial communities in the marine ecosystem, particularly in upwelling
regions (Sun et al., 2022). A positive correlation of fluorescence and PP with the majority of

bacterial communities including Gammaproteobacteria SAR66, OM60 (NOR5), Bacteroidetes NS4 and NS5 marine groups, Actinobacterial genera Sva0996 marine groups, Ca. *Actinomarina*, Ca. *Marinimicrobia* and other candidatus groups during the monsoon season were observed. These bacterial groups have previously been shown to be associated with high-productivity environments and their presence may reflect an increase in nutrient availability due to upwelling (Bachmann et al., 2018; Reji et al., 2020; James et al., 2022). This surge in the abundance of candidatus phyla can be attributed to their adaptation to thrive in environments rich in organic matter derived from primary productivity (Mestre et al., 2018; Frank et al., 2016). The enhanced availability of nutrients and labile carbon sources during the productive monsoon season likely stimulates the growth and metabolic activities of these bacterial phyla (Hug et al., 2015; Reji et al., 2020; Graham and Tully, 2021). Seasonal variations in primary productivity have been recognized as a crucial driver of microbial community composition and abundance. As primary productivity variations occur, bacterial communities respond by adjusting their composition and functional capacities to adapt to changing environmental conditions (Nguyen et al., 2020). The relationship between primary productivity-derived organic matter and bacterial diversity existed at both local and global scales (Phoma et al., 2021). Understanding the influence of seasonal primary productivity variations on increased bacterial diversity is essential for unraveling the intricate relationships between primary producers and bacterial populations, as well as their roles in nutrient cycling and ecosystem functioning. The correlation of bacterial communities with fluorescence and PP indicated that changes in these parameters in the seawater column could impact the abundance and composition of bacterial community variations. Studies had shown that the interactions between phytoplankton and accompanying bacteria could have important effects on both communities (Jasti et al., 2005; Rooney-Varga et al., 2005).

Predictive functional profiling showed autotrophic activities including, prokaryotic carbon fixation activities and cellular catabolism increased during the monsoon season. Autotrophic activities, as well as the abundance of prokaryotic carbon fixation pathways, could be due to the significant increase in Cyanobacterial populations during the upwelling season, which may have impacted the results of our metagenomics comparisons. Studies suggest that the differences in functional profiles observed between different seasons using metagenomics approaches may be influenced by the high abundance of genes associated with photosynthesis from Cyanobacteria during the upwelling season (Ward et al., 2017; Matsumoto et al., 2021). Upwelling circumstances are linked to an increase in energy metabolic processes (Coelho-Souza et al., 2012; Gregoracci et al., 2015; Parvathi et al., 2019), such as prokaryotic carbon fixation, nitrogen metabolism and methane metabolism, which was also evident from our study (Fig.8a). Understanding the metabolic interactions between primary producers, such as photoautotrophic species like phytoplanktons and Cyanobacteria and heterotrophic organisms, such as bacteria and archaea, is essential for understanding how ecosystems work (Caroppo et al., 2022). The results of our study indicate that upwelling has the potential to increase the metabolic capacity of the majority of the bacterial community, particularly in the areas of prokaryotic carbon fixation and various cellular catabolic processes (Fig. 8b). Compounds such as dissolved organic matter can enhance the growth of bacterial species such as Roseobacter, Oceanospirillales and Flavobacteriales (Osterholz et al., 2016), leading to alterations in the microbial community and functional diversity. Overall during the monsoon season, the dominant pathways related to autotrophic carbon fixation and cellular processing were abundant. The dominance of these pathways during monsoonal upwelling season indicates that bacterial communities are adapting to the changes in nutrient availability and environmental conditions during monsoon season and are prioritizing biomass production and maintenance of

cellular metabolism as key functions. This observation underscores the significant impact of

primary productivity variations, particularly those associated with upwelling events, on the

composition of bacterial communities within the WCI region.

5. Conclusion

This study presents a detailed account of the bacterial community structure in C-Max

depths during the non-monsoon and monsoon seasons along WCI. This study has improved the

understanding of microbial functioning along the WCI, an important upwelling zone and a

significant carbon sequestration region. Major implications from this study is based on the

observed taxonomic and functional diversity patterns, combined with physicochemical factors,

due to primary productivity variations and relationships among bacterial communities. This

study using Next-generation sequencing methods, has provided a deeper understanding of the

community structure and diversity associated with seasonal changes in primary productivity,

along the WCI. The level of taxonomic resolution provided by Illumina sequencing allowed us

to build upon previous research and uncover that, seasonal gradients of the bacterial taxonomic

makeup, revealing spatiotemporal variations at genera level, that was previously unexplored

from this region. Additionally, Illumina sequencing has enabled us to understand the presence

of candidatus phylotypes which are metabolically active, which are not characterised by

conventional culture-based methods. The seasonal transitions in WCI, and the dynamics of the

bacterial community composition are most likely the outcome of increased primary

productivity and environmental factors. Further studies are required to understand the basis of

microbial mechanisms connected to organic matter recycling and export to the deeper depths

of this region. This study has improved the understanding of the seasonal changes in microbial

community structure along the WCI in the Arabian Sea.

Funding

This research did not receive any specific grant from funding agencies in the public,

commercial, or not-for-profit sectors.

Declaration of competing interest

The authors declare no conflicts of interest.

Data availability

Data will be made available on request. The metagenomics data generated in this study have

been deposited in the National Centre for Biotechnology Information (NCBI) Sequence Read

Archive (SRA) under the bioproject number PRJNA873935. The data can be accessed from

the following link: [https://www.ncbi.nlm.nih.gov/bioproject/PRJNA873935].

Acknowledgments

research fellowship, Grant No. F.16-6 (Dec.2016)/2017 (NET). This is NIO contribution

number xxxx.

Appendix A. Supplementary data

Appendix B. Supplementary file

References

Aldunate, M., De La Iglesia, R., Bertagnolli, A.D., Ulloa, O., 2018. Oxygen modulates

bacterial community composition in the coastal upwelling waters off central Chile. Deep

Sea Research Part II: Topical Studies in Oceanography 156, 68–79.

https://doi.org/10.1016/j.dsr2.2018.02.001

Alonso, C., Pernthaler, J., 2006. Roseobacter and SAR11 dominate microbial glucose uptake

in coastal North Sea waters. Environ Microbiol 8, 2022–2030.

https://doi.org/10.1111/j.1462-2920.2006.01082.x

Alonso-Gutiérrez, J., Lekunberri, I., Teira, E., Gasol, J.M., Figueras, A., Novoa, B., 2009.

Bacterioplankton composition of the coastal upwelling system of 'Ría de Vigo', NW Spain.

FEMS Microbiology Ecology 70, 493–505. https://doi.org/10.1111/j.1574-

6941.2009.00766.x

Alonso-Sáez, L., Arístegui, J., Pinhassi, J., Gómez-Consarnau, L., González, J., Vaqué, D.,

Agustí, S., Gasol, J., 2007. Bacterial assemblage structure and carbon metabolism along a

productivity gradient in the NE Atlantic Ocean. Aquat. Microb. Ecol. 46, 43–53.

https://doi.org/10.3354/ame046043

Alonso-Sáez, L., Gasol, J.M., 2007. Seasonal Variations in the Contributions of Different

Bacterial Groups to the Uptake of Low-Molecular-Weight Compounds in Northwestern

Mediterranean Coastal Waters. Appl Environ Microbiol 73, 3528–3535.

https://doi.org/10.1128/AEM.02627-06

Anas, A., Chekidhenkuzhiyil, J., Nilayangod, C., Krishna, K., Raju, G.T., 2021. Prevalence of

obligate and facultative anaerobic bacteria in the mudbank along the southwest coast of

India. Regional Studies in Marine Science 42, 101660.

https://doi.org/10.1016/j.rsma.2021.101660

Anas, A., V.A., S., C., J., T.R., G., Mathew, D., Krishna, K., Nair, S., K.R., M., K.V., J., 2018.

Upwelling induced changes in the abundance and community structure of archaea and

bacteria in a recurring mud bank along the southwest coast of India. Regional Studies in

Marine Science 18, 113–121. https://doi.org/10.1016/j.rsma.2018.01.006

Azam, F., Malfatti, F., 2007. Microbial structuring of marine ecosystems. Nat Rev Microbiol

5, 782–791. https://doi.org/10.1038/nrmicro1747

Bachmann, J., Heimbach, T., Hassenrück, C., Kopprio, G.A., Iversen, M.H., Grossart, H.P.,

Gärdes, A., 2018. Environmental Drivers of Free-Living vs. Particle-Attached Bacterial

Community Composition in the Mauritania Upwelling System. Front. Microbiol. 9, 2836.

https://doi.org/10.3389/fmicb.2018.02836

Baltar, F., Arístegui, J., Gasol, J., Hernández-León, S., Herndl, G., 2007. Strong coastocean

and surfacedepth gradients in prokaryotic assemblage structure and activity in a coastal

transition zone region. Aquat. Microb. Ecol. 50, 63–74. https://doi.org/10.3354/ame01156

Bandekar, M., Ramaiah, N., Meena, R.M., 2018. Diversity and abundance of denitrifying and

anammox bacteria from the Arabian Sea oxygen minimum zone. Deep Sea Research Part

II: Topical Studies in Oceanography 156, 19–26.

https://doi.org/10.1016/j.dsr2.2018.08.008

Berg, J.S., Ahmerkamp, S., Pjevac, P., Hausmann, B., Milucka, J., Kuypers, M.M.M., 2022.

How low can they go? Aerobic respiration by microorganisms under apparent anoxia.

FEMS Microbiology Reviews 46, fuac006. https://doi.org/10.1093/femsre/fuac006

Bergen, B., Herlemann, D.P.R., Jürgens, K., 2015. Zonation of bacterioplankton communities

along aging upwelled water in the northern Benguela upwelling. Front. Microbiol. 6.

https://doi.org/10.3389/fmicb.2015.00621

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., et al.

Reproducible, interactive, scalable and extensible microbiome data science using QIIME

2. Nat Biotechnol 37, 852–857. https://doi.org/10.1038/s41587-019-0209-9

Bora, S.S., Dullah, S., Dey, K.K., Hazarika, D.J., Sarmah, U., Sharma, D., Goswami, G., Singh,

N.R., Barooah, M., 2022. Additive-induced pH determines bacterial community

composition and metabolome in traditional mustard seed fermented products. Front.

Sustain. Food Syst. 6, 1006573. https://doi.org/10.3389/fsufs.2022.1006573

Buchan, A., LeCleir, G.R., Gulvik, C.A., González, J.M., 2014. Master recyclers: features and

functions of bacteria associated with phytoplankton blooms. Nat Rev Microbiol 12, 686–

698. https://doi.org/10.1038/nrmicro3326

Bunse, C., Bertos-Fortis, M., Sassenhagen, I., Sildever, S., Sjöqvist, C., Godhe, A., Gross, S.,

Kremp, A., Lips, I., Lundholm, N., Rengefors, K., Sefbom, J., Pinhassi, J., Legrand, C.,

2016. Spatio-Temporal Interdependence of Bacteria and Phytoplankton during a Baltic Sea

Spring Bloom. Front. Microbiol. 7. https://doi.org/10.3389/fmicb.2016.00517

Cai, Y.-M., Ning, X.-R., Liu, C.-G., Hao, Q., 2007. Distribution Pattern of Photosynthetic

Picoplankton and Heterotrophic Bacteria in the Northern South China Sea. J Integrative

Plant Biology 49, 282–298. https://doi.org/10.1111/j.1744-7909.2007.00347.x

Capone, D.G., Hutchins, D.A., 2013. Microbial biogeochemistry of coastal upwelling regimes

in a changing ocean. Nature Geosci 6, 711–717. https://doi.org/10.1038/ngeo1916

Caroppo, C., Azzaro, F., Bergamasco, A., Caruso, G., Decembrini, F., 2022. Phytoplankton

and Bacterial Communities' Patterns in a Highly Dynamic Ecosystem (Central

Mediterranean Sea). Water 14, 2057. https://doi.org/10.3390/w14132057

Chen, S., He, Y.-B., Xie, Z.-X., Kong, L.-F., Yan, K.-Q., Li, D.-X., Wu, P.-F., Zheng, R.-W.,

Peng, L., Chen, J.-W., Lin, L., Liu, S.-Q., Fan, G.-Y., Wang, D.-Z., 2021. Metaproteomics

reveals nutrient availability shaping distinct microbial community and metabolic niche in

the nutrient-depleted and replete layers of an oligotrophic euphotic zone. Science of The

Total Environment 774, 145123. https://doi.org/10.1016/j.scitotenv.2021.145123

Cho, J.-C., Giovannoni, S.J., 2004. Cultivation and Growth Characteristics of a Diverse Group

of Oligotrophic Marine *Gammaproteobacteria*. Appl Environ Microbiol 70, 432–440.

https://doi.org/10.1128/AEM.70.1.432-440.2004

Cho, J.-C., Stapels, M.D., Morris, R.M., Vergin, K.L., Schwalbach, M.S., Givan, S.A.,

Barofsky, D.F., Giovannoni, S.J., 2007. Polyphyletic photosynthetic reaction centre genes

in oligotrophic marine Gammaproteobacteria. Environ Microbiol 9, 1456–1463.

https://doi.org/10.1111/j.1462-2920.2007.01264.x

Coelho-Souza, S.A., López, M.S., Guimarães, J.R.D., Coutinho, R., Candella, R.N., 2012.

Biophysical interactions in the Cabo Frio upwelling system, southeastern Brazil. Brazilian

Journal   of   Oceanography   60,   353–365.   https://doi.org/10.1590/S1679-

87592012000300008

Cullen, J.J., 2015. Subsurface Chlorophyll Maximum Layers: Enduring Enigma or Mystery

Solved? Annu. Rev. Mar. Sci. 7, 207–239. https://doi.org/10.1146/annurev-marine-

010213-135111

Danovaro, R., Corinaldesi, C., Dell'Anno, A., Rastelli, E., 2017. Potential impact of global

climate change on benthic deep-sea microbes. FEMS Microbiology Letters 364.

https://doi.org/10.1093/femsle/fnx214

Doré, H., Leconte, J., Guyet, U., Breton, S., Farrant, G.K., Demory, D., Ratin, M., Hoebeke,

M., Corre, E., Pitt, F.D., Ostrowski, M., Scanlan, D.J., Partensky, F., Six, C., Garczarek,

L., 2022. Global Phylogeography of Marine *Synechococcus* in Coastal Areas Reveals

Strong Community Shifts. mSystems 7, e00656-22.

https://doi.org/10.1128/msystems.00656-22

Douglas, G.M., Maffei, V.J., Zaneveld, J.R., Yurgel, S.N., Brown, J.R., Taylor, C.M.,

Huttenhower, C., Langille, M.G.I., 2020. PICRUSt2 for prediction of metagenome

functions. Nat Biotechnol 38, 685–688. https://doi.org/10.1038/s41587-020-0548-6

Du, J., Xiao, K., Li, L., Ding, X., Liu, H., Lu, Y., Zhou, S., 2013. Temporal and Spatial

Diversity of Bacterial Communities in Coastal Waters of the South China Sea. PLoS ONE

8, e66968. https://doi.org/10.1371/journal.pone.0066968

Ducklow, H.W., Quinby, H.L., Carlson, C.A., 1995. Bacterioplankton dynamics in the

equatorial Pacific during the 1992 El Niño. Deep Sea Research Part II: Topical Studies in

Oceanography 42, 621–638. https://doi.org/10.1016/0967-0645(95)00022-I

Fernandes, G.L., Shenoy, B.D., Damare, S.R., 2020. Diversity of Bacterial Community in the

Oxygen Minimum Zones of Arabian Sea and Bay of Bengal as Deduced by Illumina

Sequencing. Front. Microbiol. 10, 3153. https://doi.org/10.3389/fmicb.2019.03153

Fernandes, V., Bogati, K., 2022. Analysis of Bacteria–Phytoplankton relationships at three

discrete locations in the Eastern Arabian Sea during winter. Continental Shelf Research

243, 104751. https://doi.org/10.1016/j.csr.2022.104751

Frank, A.H., Garcia, J.A.L., Herndl, G.J., Reinthaler, T., 2016. Connectivity between surface

and deep waters determines prokaryotic diversity in the North Atlantic Deep Water.

Environmental Microbiology 18, 2052–2063. https://doi.org/10.1111/1462-2920.13237

Freitas, S., Hatosy, S., Fuhrman, J.A., Huse, S.M., Mark Welch, D.B., Sogin, M.L., Martiny,

A.C., 2012. Global distribution and diversity of marine Verrucomicrobia. ISME J 6, 1499–

1505. https://doi.org/10.1038/ismej.2012.3

Gauns, M., Madhupratap, M., Ramaiah, N., Jyothibabu, R., Fernandes, V., Paul, J.T., Prasanna

Kumar, S., 2005. Comparative accounts of biological productivity characteristics and

estimates of carbon fluxes in the Arabian Sea and the Bay of Bengal. Deep Sea Research

Part II: Topical Studies in Oceanography 52, 2003–2017.

https://doi.org/10.1016/j.dsr2.2005.05.009

Gomes, J., Khandeparker, R., Bandekar, M., Meena, R.M., Ramaiah, N., 2018. Quantitative

analyses of denitrifying bacterial diversity from a seasonally hypoxic monsoon governed

tropical coastal region. Deep Sea Research Part II: Topical Studies in Oceanography 156,

34–43. https://doi.org/10.1016/j.dsr2.2017.12.012

Gong, F., Ji, Q., Li, G., Yin, K., Gong, J., 2022. Diversity and distribution of bacterioplankton

in the coastal upwelling waters off Hainan Island, China. Acta Oceanol. Sin. 41, 76–85.

https://doi.org/10.1007/s13131-021-1807-1

Gonsalves, M.-J., Paropkari, A.L., Fernandes, C.E.G., Loka Bharathi, P.A., Krishnakumari, L.,

Fernando, V., Nampoothiri, G.E., 2011. Predominance of anaerobic bacterial community

over aerobic community contribute to intensify 'oxygen minimum zone' in the eastern

Arabian Sea. Continental Shelf Research 31, 1224–1235.

https://doi.org/10.1016/j.csr.2011.04.011

González, J.M., Moran, M.A., 1997. Numerical dominance of a group of marine bacteria in the

alpha-subclass of the class Proteobacteria in coastal seawater. Appl Environ Microbiol 63,

4237–4242. https://doi.org/10.1128/aem.63.11.4237-4242.1997

González, J.M., Simó, R., Massana, R., Covert, J.S., Casamayor, E.O., Pedrós-Alió, C., Moran,

M.A., 2000. Bacterial Community Structure Associated with a

Dimethylsulfoniopropionate-Producing North Atlantic Algal Bloom. Appl Environ

Microbiol 66, 4237–4246. https://doi.org/10.1128/AEM.66.10.4237-4246.2000

Graham, E.D., Tully, B.J., 2021. Marine Dadabacteria exhibit genome streamlining and

phototrophy-driven niche partitioning. ISME J 15, 1248–1256.

https://doi.org/10.1038/s41396-020-00834-5

Gregoracci, G.B., Soares, A.C.D.S., Miranda, M.D., Coutinho, R., Thompson, F.L., 2015.

Insights into the Microbial and Viral Dynamics of a Coastal Downwelling-Upwelling

Transition. PLoS ONE 10, e0137090. https://doi.org/10.1371/journal.pone.0137090

Gupta, G.V.M., Jyothibabu, R., Ramu, C.V., Yudhistir Reddy, A., Balachandran, K.K.,

Sudheesh, V., Kumar, S., Chari, N.V.H.K., Bepari, K.F., Marathe, P.H., Bikram Reddy, B.,

Vijayan, A.K., 2021. The world's largest coastal deoxygenation zone is not

anthropogenically driven. Environ. Res. Lett. 16, 054009. https://doi.org/10.1088/1748-

9326/abe9eb

Gupta, G.V.M., Sudheesh, V., Sudharma, K.V., Saravanane, N., Dhanya, V., Dhanya, K.R.,

Lakshmi, G., Sudhakar, M., Naqvi, S.W.A., 2016. Evolution to decay of upwelling and

associated biogeochemistry over the southeastern Arabian Sea shelf. J. Geophys. Res.

Biogeosci. 121, 159–175. https://doi.org/10.1002/2015JG003163

Gutiérrez-Barral, A., Teira, E., Hernández-Ruiz, M., Fernández, E., 2021. Response of

prokaryote community composition to riverine and atmospheric nutrients in a coastal

embayment: Role of organic matter on Vibrionales. Estuarine, Coastal and Shelf Science

251, 107196. https://doi.org/10.1016/j.ecss.2021.107196

Habeebrehman, H., Prabhakaran, M.P., Jacob, J., Sabu, P., Jayalakshmi, K.J., Achuthankutty,

C.T., Revichandran, C., 2008. Variability in biological responses influenced by upwelling

events in the Eastern Arabian Sea. Journal of Marine Systems 74, 545–560.

https://doi.org/10.1016/j.jmarsys.2008.04.002

Ho, A., Lonardo, D.P.D., Bodelier, P.L.E., 2017. Revisiting life strategy concepts in

environmental microbial ecology. Microbiology Ecology fix006.

https://doi.org/10.1093/femsec/fix006

Hug, L.A., Thomas, B.C., Sharon, I., Brown, C.T., Sharma, R., Hettich, R.L., Wilkins, M.J.,

Williams, K.H., Singh, A., Banfield, J.F., 2016. Critical biogeochemical functions in the

subsurface are associated with bacteria from new phyla and little studied lineages: N- and

C-cycling organisms in the subsurface. Environ Microbiol 18, 159–173.

https://doi.org/10.1111/1462-2920.12930

James, C.C., Barton, A.D., Allen, L.Z., Lampe, R.H., Rabines, A., Schulberg, A., Zheng, H.,

Goericke, R., Goodwin, K.D., Allen, A.E., 2022. Influence of nutrient supply on plankton

microbiome biodiversity and distribution in a coastal upwelling region. Nat Commun 13,

2448. https://doi.org/10.1038/s41467-022-30139-4

Jasti, S., Sieracki, M.E., Poulton, N.J., Giewat, M.W., Rooney-Varga, J.N., 2005. Phylogenetic

Diversity and Specificity of Bacteria Closely Associated with *Alexandrium* spp. and Other

Phytoplankton. Appl Environ Microbiol 71, 3483–3494.

https://doi.org/10.1128/AEM.71.7.3483-3494.2005

Jiao, N., Herndl, G.J., Hansell, D.A., Benner, R., Kattner, G., Wilhelm, S.W., Kirchman, D.L.,

Weinbauer, M.G., Luo, T., Chen, F., Azam, F., 2010. Microbial production of recalcitrant

dissolved organic matter: long-term carbon storage in the global ocean. Nat Rev Microbiol

8, 593–599. https://doi.org/10.1038/nrmicro2386

Jing, H., Xia, X., Suzuki, K., Liu, H., 2013. Vertical Profiles of Bacteria in the Tropical and

Subarctic Oceans Revealed by Pyrosequencing. PLoS ONE 8, e79423.

https://doi.org/10.1371/journal.pone.0079423

Kadnikov, V.V., Mardanov, A.V., Beletsky, A.V., Karnachuk, O.V., Ravin, N.V., 2020.

Microbial Life in the Deep Subsurface Aquifer Illuminated by Metagenomics. Front.

Microbiol. 11, 572252. https://doi.org/10.3389/fmicb.2020.572252

Khandeparker, L., Eswaran, R., Gardade, L., Kuchi, N., Mapari, K., Naik, S.D., Anil, A.C.,

2016. Elucidation of the tidal influence on bacterial populations in a monsoon influenced

estuary through simultaneous observations. Environ Monit Assess 189, 41.

https://doi.org/10.1007/s10661-016-5687-3

Khandeparker, L., Kuchi, N., Kale, D., Anil, A.C., 2017. Microbial community structure of

surface sediments from a tropical estuarine environment using next generation sequencing.

Ecological Indicators 74, 172–181. https://doi.org/10.1016/j.ecolind.2016.11.023

Kirchman, D.L., 2002. The ecology of Cytophaga–Flavobacteria in aquatic environments.

FEMS Microbiology Ecology 39, 91–100. https://doi.org/10.1111/j.1574-

6941.2002.tb00910.x

Kuchi, N., Khandeparker, L., Anil, A.C., 2021. Response of the bacterial metagenome in port

environments to changing environmental conditions. Marine Pollution Bulletin 172,

112869. https://doi.org/10.1016/j.marpolbul.2021.112869

Kumar, S.P., Madhupratap, M., Kumar, M.D., Gauns, M., Muraleedharan, P.M., Sarma,

V.V.S.S., De Souza, S.N., 2000. Physical control of primary productivity on a seasonal

scale in central and eastern Arabian Sea. J Earth Syst Sci 109, 433–441.

https://doi.org/10.1007/BF02708331

Kumar, S.P., Ramaiah, N., Gauns, M., Sarma, V.V.S.S., Muraleedharan, P.M., Raghukumar,

S., Dileep Kumar, M., Madhupratap, M., 2001. Physical forcing of biological productivity

in the Northern Arabian Sea during the Northeast Monsoon. Deep Sea Research Part II:

Topical Studies in Oceanography 48, 1115–1126. https://doi.org/10.1016/S0967-

0645(00)00133-8

Kuypers, M.M.M., Lavik, G., Woebken, D., Schmid, M., Fuchs, B.M., Amann, R., Jørgensen,

B.B., Jetten, M.S.M., 2005. Massive nitrogen loss from the Benguela upwelling system

through anaerobic ammonium oxidation. Proc. Natl. Acad. Sci. U.S.A. 102, 6478–6483.

https://doi.org/10.1073/pnas.0502088102

Lamont, T., Barlow, R.G., Kyewalyanga, M.S., 2014. Physical drivers of phytoplankton

production in the southern Benguela upwelling system. Deep Sea Research Part I:

Oceanographic Research Papers 90, 1–16. https://doi.org/10.1016/j.dsr.2014.03.003

Lever, M.A., Rogers, K.L., Lloyd, K.G., Overmann, J., Schink, B., Thauer, R.K., Hoehler,

T.M., Jørgensen, B.B., 2015. Life under extreme energy limitation: a synthesis of

laboratory- and field-based investigations. FEMS Microbiology Reviews 39, 688–728.

https://doi.org/10.1093/femsre/fuv020

Li, Y., Sun, L.-L., Sun, M.-L., Su, H.-N., Zhang, X.-Y., Xie, B.-B., Chen, X.-L., Zhang, Y.-Z.,

Qin, Q.-L., 2018. Vertical and horizontal biogeographic patterns and major factors

affecting bacterial communities in the open South China Sea. Sci Rep 8, 8800.

https://doi.org/10.1038/s41598-018-27191-w

Li, Y.-Y., Chen, X.-H., Xie, Z.-X., Li, D.-X., Wu, P.-F., Kong, L.-F., Lin, L., Kao, S.-J., Wang, D.-Z., 2018. Bacterial Diversity and Nitrogen Utilization Strategies in the Upper Layer of the Northwestern Pacific Ocean. Frontiers in Microbiology 9.

Lincy, J., Manohar, C.S., 2021. 16s rRNA and Hydrazine Gene-Based Profiling of the Candidatus Scalindua Community from the Arabian Sea Hypoxic Sediments. Current Science 120, 684. https://doi.org/10.18520/cs/v120/i4/684-693

Liu, S., Liu, Z., 2020. Distinct capabilities of different Gammaproteobacterial strains on utilizing small peptides in seawater. Sci Rep 10, 464. https://doi.org/10.1038/s41598-019-57189-x

Liu, X., Xie, N., Li, J., Bai, M., Sen, B., Wang, G., 2022. Potential Contribution of Coastal Upwelling to Carbon Sink through Interaction between Cyanobacteria and Microbial Eukaryotes. Water 14, 3097. https://doi.org/10.3390/w14193097

Matsumoto, K., Sakami, T., Watanabe, T., Taniuchi, Y., Kuwata, A., Kakehi, S., Engkong, T., Igarashi, Y., Kinoshita, S., Asakawa, S., Hattori, M., Watabe, S., Ishino, Y., Kobayashi, T., Gojobori, T., Ikeo, K., 2021. Metagenomic analysis provides functional insights into seasonal change of a non-cyanobacterial prokaryotic community in temperate coastal waters. PLoS ONE 16, e0257862. https://doi.org/10.1371/journal.pone.0257862

McMurdie, P.J., Holmes, S., 2013. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. PLoS ONE 8, e61217. https://doi.org/10.1371/journal.pone.0061217

Mena, C., Reglero, P., Balbín, R., Martín, M., Santiago, R., Sintes, E., 2022. Dynamics of actively dividing prokaryotes in the western Mediterranean Sea. Sci Rep 12, 2064. https://doi.org/10.1038/s41598-022-06120-y

Mestre, M., Ruiz-González, C., Logares, R., Duarte, C.M., Gasol, J.M., Sala, M.M., 2018.

Sinking particles promote vertical connectivity in the ocean microbiome. Proc. Natl. Acad.

Sci. U.S.A. 115. https://doi.org/10.1073/pnas.1802470115

Mitrophanov, A.Y., Groisman, E.A., 2008. Signal integration in bacterial two-component

regulatory systems. Genes Dev. 22, 2601–2611. https://doi.org/10.1101/gad.1700308

Mohanty, B., Majedi, S.M., Pavagadhi, S., Te, S.H., Boo, C.Y., Gin, K.Y.-H., Swarup, S.,

2022. Effects of Light and Temperature on the Metabolic Profiling of Two Habitat-

Dependent Bloom-Forming Cyanobacteria. Metabolites 12, 406.

https://doi.org/10.3390/metabo12050406

Molina, V., Fernández, C., 2020. Bacterioplankton response to nitrogen and dissolved organic

matter produced from salmon mucus. MicrobiologyOpen 9.

https://doi.org/10.1002/mbo3.1132

Montes, T., Guerrero-Feijóo, E., Moreira-Coello, V., Bode, A., Ruiz-Villarreal, M., Mouriño-

Carballido, B., Varela, M.M., 2020. Vertical zonation of bacterial assemblages attributed

to physical stratification during the summer relaxation of the coastal upwelling off Galicia

(NW Spain). Estuarine, Coastal and Shelf Science 245, 106791.

https://doi.org/10.1016/j.ecss.2020.106791

Morris, R.M., Frazar, C.D., Carlson, C.A., 2012. Basin-scale patterns in the abundance of

SAR11 subclades, marine Actinobacteria (OM1), members of the Roseobacter clade and

OCS116 in the South Atlantic: Bacterioplankton communities in the South Atlantic.

Environmental Microbiology 14, 1133–1144. https://doi.org/10.1111/j.1462-

2920.2011.02694.x

Naqvi, S.W.A., Bange, H.W., Farías, L., Monteiro, P.M.S., Scranton, M.I., Zhang, J., 2010.

Marine hypoxia/anoxia as a source of CH_4 and N_2O. Biogeosciences 7, 2159–2190.

https://doi.org/10.5194/bg-7-2159-2010

Nguyen, J., Lara-Gutiérrez, J., Stocker, R., 2021. Environmental fluctuations and their effects

on microbial communities, populations and individuals. FEMS Microbiology Reviews 45,

fuaa068. https://doi.org/10.1093/femsre/fuaa068

Osterholz, H., Singer, G., Wemheuer, B., Daniel, R., Simon, M., Niggemann, J., Dittmar, T.,

2016. Deciphering associations between dissolved organic molecules and bacterial

communities in a pelagic marine system. ISME J 10, 1717–1730.

https://doi.org/10.1038/ismej.2015.231

Parab, A.S., Jagtap, A.S., Meena, R.M., Manohar, C.S., 2022. Bacterial dynamics along the

west coast of India during the non-monsoon and monsoon season. Continental Shelf

Research 251, 104876. https://doi.org/10.1016/j.csr.2022.104876

Parks, D.H., Tyson, G.W., Hugenholtz, P., Beiko, R.G., 2014. STAMP: statistical analysis of

taxonomic and functional profiles. Bioinformatics 30, 3123–3124.

https://doi.org/10.1093/bioinformatics/btu494

Parvathi, A., Jasna, V., Aswathy, V.K., Nathan, V.K., Aparna, S., Balachandran, K.K., 2019.

Microbial diversity in a coastal environment with co-existing upwelling and mud-banks

along the south west coast of India. Mol Biol Rep 46, 3113–3127.

https://doi.org/10.1007/s11033-019-04766-y

Phoma, B.S., Makhalanyane, T.P., 2021. Depth-Dependent Variables Shape Community

Structure and Functionality in the Prince Edward Islands. Microb Ecol 81, 396–409.

https://doi.org/10.1007/s00248-020-01589-4

Pinhassi, J., Sala, M.M., Havskum, H., Peters, F., Guadayol, Ò., Malits, A., Marrasé, C., 2004.

Changes in Bacterioplankton Composition under Different Phytoplankton Regimens. Appl

Environ Microbiol 70, 6753–6766. https://doi.org/10.1128/AEM.70.11.6753-6766.2004

Pomeroy, L., leB. Williams, P., Azam, F., Hobbie, J., 2007. The Microbial Loop. Oceanog. 20,

28–33. https://doi.org/10.5670/oceanog.2007.45

Porter, K.G., Feig, Y.S., 1980. The use of DAPI for identifying and counting aquatic

microflora1. Limnol. Oceanogr. 25, 943–948. https://doi.org/10.4319/lo.1980.25.5.0943

Prüß, B.M., 2017. Involvement of Two-Component Signaling on Bacterial Motility and

Biofilm Development. J Bacteriol 199. https://doi.org/10.1128/JB.00259-17

Ramaiah, N., Raghukumar, S., Gauns, M., 1996. Bacterial abundance and production in the

central and eastern Arabian Sea. Current Science 71, 878–882.

Ramaiah, N., Raghukumar, S., Mangesh, G., Madhupratap, M., 2005. Seasonal variations in

carbon biomass of bacteria, thraustochytrids and microzooplankton in the Northern

Arabian Sea. Deep Sea Research Part II: Topical Studies in Oceanography 52, 1910–1921.

https://doi.org/10.1016/j.dsr2.2005.05.004

Reji, L., Tolar, B.B., Chavez, F.P., Francis, C.A., 2020. Depth-Differentiation and Seasonality

of Planktonic Microbial Assemblages in the Monterey Bay Upwelling System. Front.

Microbiol. 11, 1075. https://doi.org/10.3389/fmicb.2020.01075

Rixen, T., Gaye, B., Emeis, K.-C., Ramaswamy, V., 2019. The ballast effect of lithogenic

matter and its influences on the carbon fluxes in the Indian Ocean. Biogeosciences 16, 485–

503. https://doi.org/10.5194/bg-16-485-2019

Rocke, E., Cheung, S., Gebe, Z., Dames, N.R., Liu, H., Moloney, C.L., 2020. Marine Microbial

Community Composition During the Upwelling Season in the Southern Benguela. Front.

Mar. Sci. 7, 255. https://doi.org/10.3389/fmars.2020.00255

Rooney-Varga, J.N., Giewat, M.W., Savin, M.C., Sood, S., LeGresley, M., Martin, J.L., 2005.

Links between Phytoplankton and Bacterial Community Dynamics in a Coastal Marine

Environment. Microb Ecol 49, 163–175. https://doi.org/10.1007/s00248-003-1057-0

Sachithanandam, V., Saravanane, N., Chandrasekar, K., Karthick, P., Lalitha, P., Sai

Elangovan, S., Sudhakar, M., 2020. Microbial diversity from the continental shelf regions

of the Eastern Arabian Sea: A metagenomic approach. Saudi Journal of Biological Sciences

27, 2065–2075. https://doi.org/10.1016/j.sjbs.2020.06.011

Sarma, V.V.S.S., 2002. An evaluation of physical and biogeochemical processes regulating

perennial suboxic conditions in the water column of the Arabian Sea: perennial existence

of suboxic zone in the arabian sea. Global Biogeochem. Cycles *6, 29-1-29–11.

https://doi.org/10.1029/2001GB001461

Schattenhofer, M., Fuchs, B.M., Amann, R., Zubkov, M.V., Tarran, G.A., Pernthaler, J., 2009.

Latitudinal distribution of prokaryotic picoplankton populations in the Atlantic Ocean.

Environmental   Microbiology   11,   2078–2093.   https://doi.org/10.1111/j.1462-

2920.2009.01929.x

Schott, F.A., Xie, S.-P., McCreary, J.P., 2009. Indian Ocean circulation and climate variability.

Rev. Geophys. 47, RG1002. https://doi.org/10.1029/2007RG000245

Sebastián, M., Gasol, J.M., 2013. Heterogeneity in the nutrient limitation of different

bacterioplankton groups in the Eastern Mediterranean Sea. ISME J 7, 1665–1668.

https://doi.org/10.1038/ismej.2013.42

Sebastián, M., Smith, A.F., González, J.M., Fredricks, H.F., Van Mooy, B., Koblížek, M.,

Brandsma, J., Koster, G., Mestre, M., Mostajir, B., Pitta, P., Postle, A.D., Sánchez, P.,

Gasol, J.M., Scanlan, D.J., Chen, Y., 2016. Lipid remodelling is a widespread strategy in

marine heterotrophic bacteria upon phosphorus deficiency. ISME J 10, 968–978.

https://doi.org/10.1038/ismej.2015.172

Sheik, C.S., Jain, S., Dick, G.J., 2014. Metabolic flexibility of enigmatic SAR 324 revealed

through metagenomics and metatranscriptomics. Environmental Microbiology 16, 304–

317. https://doi.org/10.1111/1462-2920.12165

Siezen, R.J., Van Hylckama Vlieg, J.E., 2011. Genomic diversity and versatility of

Lactobacillus plantarum, a natural metabolic engineer. Microb Cell Fact 10, S3.

https://doi.org/10.1186/1475-2859-10-S1-S3

Smitha, B.R., Sanjeevan, V.N., Vimalkumar, K.G., Revichandran, C., 2008. On the Upwelling

off the Southern Tip and along the West Coast of India. Journal of Coastal Research 24,

95–102.

Spring, S., Scheuner, C., Göker, M., Klenk, H.-P., 2015. A taxonomic framework for emerging

groups of ecologically important marine gammaproteobacteria based on the reconstruction

of evolutionary relationships using genome-scale data. Front. Microbiol. 6.

https://doi.org/10.3389/fmicb.2015.00281

Sun, F.-L., Wang, Y.-S., Wu, M.-L., Sun, C.-C., Cheng, H., 2015. Spatial and vertical

distribution of bacterial community in the northern South China Sea. Ecotoxicology 24,

1478–1485. https://doi.org/10.1007/s10646-015-1472-2

Sun, P., Wang, Y., Huang, X., Huang, B., Wang, L., 2022. Water masses and their associated

temperature and cross-domain biotic factors co-shape upwelling microbial communities.

Water Research 215, 118274. https://doi.org/10.1016/j.watres.2022.118274

Taylor, A.G., Landry, M.R., Selph, K.E., Wokuluk, J.J., 2015. Temporal and spatial patterns

of microbial community biomass and composition in the Southern California Current

Ecosystem. Deep Sea Research Part II: Topical Studies in Oceanography 112, 117–128.

https://doi.org/10.1016/j.dsr2.2014.02.006

Teira, E., Aranguren-Gassis, M., González, J., Martínez-García, S., Pérez, P., Serret, P., 2009.

Influence of allochthonous matter on microbial community structure and function in an

upwelling system off the northwest Iberian Peninsula. Aquat. Microb. Ecol. 55, 81–93.

https://doi.org/10.3354/ame01283

Thiele, S., Basse, A., Becker, J.W., Lipski, A., Iversen, M.H., Mollenhauer, G., 2019.

Microbial communities in the nepheloid layers and hypoxic zones of the Canary Current

upwelling system. MicrobiologyOpen 8, e00705. https://doi.org/10.1002/mbo3.705

Tinta, T., Vojvoda, J., Mozetič, P., Talaber, I., Vodopivec, M., Malfatti, F., Turk, V., 2015.

Bacterial community shift is induced by dynamic environmental parameters in a changing

coastal ecosystem (northern Adriatic, northeastern Mediterranean Sea) - a 2-year time-

series study: Bacterial community shift in a dynamic coastal ecosystem. Environ Microbiol

17, 3581–3596. https://doi.org/10.1111/1462-2920.12519

UNESCO, 1994 UNESCO Protocols for the Joint Global Ocean Flux Study (JGOFS) Core

Measurements, IOC Manuals and Guides, vol. 29, UNESCO, Paris (1994), p. 170

Vijayan, J., Ammini, P., Nathan, V.K., 2022. Diversity pattern of marine culturable

heterotrophic bacteria in a region with coexisting upwelling and mud banks in the

southeastern Arabian Sea. Environ Sci Pollut Res 29, 3967–3982.

https://doi.org/10.1007/s11356-021-15772-8

Vinayachandran, P.N.M., Masumoto, Y., Roberts, M.J., Huggett, J.A., Halo, I., Chatterjee, A.,

Amol, P., Gupta, G.V.M., Singh, A., Mukherjee, A., Prakash, S., Beckley, L.E., Raes, E.J.,

Hood, R., 2021. Reviews and syntheses: Physical and biogeochemical processes associated

with upwelling in the Indian Ocean. Biogeosciences 18, 5967–6029.

https://doi.org/10.5194/bg-18-5967-2021

Wang, Z., Juarez, D.L., Pan, J., Blinebry, S.K., Gronniger, J., Clark, J.S., Johnson, Z.I., Hunt,

D.E., 2019. Microbial communities across nearshore to offshore coastal transects are

primarily shaped by distance and temperature. Environ Microbiol 21, 3862–3872.

https://doi.org/10.1111/1462-2920.14734

Ward, C.S., Yung, C.-M., Davis, K.M., Blinebry, S.K., Williams, T.C., Johnson, Z.I., Hunt,

D.E., 2017. Annual community patterns are driven by seasonal switching between closely

related marine bacteria. ISME J 11, 1412–1422. https://doi.org/10.1038/ismej.2017.4

Whitman, W.B., Coleman, D.C., Wiebe, W.J., 1998. Prokaryotes: The unseen majority. Proc.

Natl. Acad. Sci. U.S.A. 95, 6578–6583. https://doi.org/10.1073/pnas.95.12.6578

Wiggert, J.D., Hood, R.R., Banse, K., Kindle, J.C., 2005. Monsoon-driven biogeochemical

processes in the Arabian Sea. Progress in Oceanography 65, 176–213.

https://doi.org/10.1016/j.pocean.2005.03.008

Yilmaz, P., Yarza, P., Rapp, J.Z., Glöckner, F.O., 2016. Expanding the World of Marine

Bacterial and Archaeal Clades. Front. Microbiol. 6.

https://doi.org/10.3389/fmicb.2015.01524

Zingone, A., D'Alelio, D., Mazzocchi, M.G., Montresor, M., Sarno, D., Team, L.-M., 2019.

Time series and beyond: multifaceted plankton research at a marine Mediterranean LTER

site. Nature Conservation 34, 273–310.

https://doi.org/10.3897/natureconservation.34.30789

Qu, W., Lin, D., Zhang, Z., Di, W., Gao, B., Zeng, R., 2018. Metagenomics Investigation of

Agarlytic Genes and Genomes in Mangrove Sediments in China: A Potential Repertory for

Table 1

Average bacterial productivity (BP), bacterial abundance (BA) and bacterial carbon biomass (BC) in the water column of the coastal and off-shore stations during non-monsoon and monsoon season.

Stations	BP (mg C m^{-3} d^{-1})	BA (Cells $\times$ 10^9 L^{-1})	BC (mg C m^{-3})
Non-monsoon season			
G30	4.30 ± 5.19	10.64 ± 2.68	212.87 ± 53.52
G600	2.95 ± 1.33	11.31 ± 4.37	226.11 ± 87.48
M30	3.79 ± 2.72	13.65 ± 0.78	272.99 ± 15.65
M600	25.92 ± 10.99	6.36 ± 2.06	127.23 ± 41.14
T30	6.75 ± 0.93	10.97 ± 0.09	219.30 ± 1.73
T600	17.80 ± 9.53	10.04 ± 4.10	200.78 ± 82
Monsoon season			
G30	13.89 ± 0.79	52.94 ± 14.27	2657.00 ± 1436.22
G600	7.34 ± 2.26	57.09 ± 40.05	1682.00 ± 293.67
M30	7.13 ± 0.44	55.51 ± 10.33	1923.33 ± 158.31
M600	7.91 ± 2.04	41.25 ± 11.19	1580.00 ± 394.08
T30	14.51 ± 2.35	54.45 ± 26.72	2226.67 ± 156.86
T600	10.93 ± 2.27	44.05 ± 26.75	1756.50 ± 399.50

Table 2

Analysis of variance (ANOVA: single factor) assessing the seasonal difference in the physicochemical and biological parameters: temperature, salinity, dissolved oxygen (DO), fluorescence, nutrients, chlorophyll-a (chl-a), primary productivity (PP), bacterial productivity (BP), bacterial abundance (BA), bacterial carbon (BC) and total plates counts (TPC) between non-monsoon and monsoon season.

	df	F	F crit	P-value
Temperature	1.00	0.57	4.03	0.45
Salinity	1.00	0.02	4.03	0.88
DO	1.00	8.68	4.03	0.001**
Fluorescence	1.00	4.58	4.03	0.04*
Nitrate	1.00	22.34	4.03	< 0.001**
Nitrite	1.00	4.86	4.03	0.03*
Phosphate	1.00	14.39	4.03	< 0.001**
Silicate	1.00	9.14	4.03	< 0.001**
Chl-a	1.00	5.41	4.04	0.02*
PP	1.00	11.45	4.08	< 0.001**
BP	1.00	0.96	4.03	0.33
BA	1.00	70.05	4.03	< 0.001**
BC	1.00	70.05	4.03	< 0.001**
TPC	1.00	3.78	4.03	0.06*

Df: degree of freedom, F value greater than F critical value indicates statistical significance.

*: Significant at 0.1%., **: Significant at 1%.

Table 3

Pearson's correlation matrix of physicochemical and biological parameters, temperature (T), salinity (S), dissolved oxygen (DO), fluorescence (F), nutrients, Chlorophyll-a (Chl-a), primary production (PP), bacterial production (BP), bacterial carbon biomass (BC) and total plates counts (TPC) during non-monsoon and monsoon season.

	T	S	DO	F	Nitrate	Nitrite	Phosphate	Silicate	Chl-a	PP	BP
T	1										
S	-0.06	1									
DO	**.80**	-0.16	1								
F	**.43**	**-.40**	**.39**	1							
Nitrate	**-.72**	-0.01	**-.58**	-0.30	1						
Nitrite	-0.03	-0.01	-0.26	-0.05	-0.05	1					
Phosphate	0.01	0.31	0.16	-0.25	-0.14	-0.05	1				
Silicate	**-.59**	0.08	**-.45**	-0.35	**.76**	0.22	0.24	1			
Chl-a	-0.06	0.14	0.04	-0.02	0.15	-0.12	-0.10	0.11	1		
PP	0.13	**-.37**	0.003	**.72**	0.04	0.05	**-.44**	-0.03	0.24	1	
BP	-0.013	0.23	-0.03	-0.03	-0.18	-0.26	0.25	-0.16	0.04	-0.12	1
BC	-0.01	-0.30	-0.17	**.58**	0.24	0.17	**-.42**	0.13	0.14	**.90**	-0.23
TPC	0.00	-0.01	0.04	-0.11	-0.02	0.07	-0.17	0.03	**.55**	0.26	-0.04

**Correlation is significant at the 0.01 level (2-tailed). *correlation is significant at the 0.05 level (2-tailed).

Fig. 1. Profiles of physiochemical variables temperature (T), salinity (S), dissolved oxygen (DO) and fluorescence (F) in the water column of the coastal and off-shore stations during non-monsoon and monsoon season along the west coast of India.

Fig. 2. Profiles of nitrate, nitrite, phosphate and silicate in the water column of the coastal and off-shore stations during non-monsoon and monsoon season along the west coast of India.

Fig. 3. Column integrated (a) chlorophyll-a (Chl-a) concentration and (b) primary productivity (PP) in the water column of the coastal and off-shore stations during non-monsoon and monsoon season along the west coast of India.

83 **Fig. 4.** Total plate counts (TPC) of culturable bacteria in the water column of the coastal and

84 off-shore stations during non-monsoon and monsoon season along the west coast of India.

(a)

Fig. 5. (a) Alpha diversity measure and (b) Non-metric multidimensional scaling (NMDS) plot (axis defines 2D space that allows the best spatial representation of sample distance, based on Bray–Curtis distance with stress = 0.08) depicting variations in the bacterial operational taxonomic units (OTUs) in the coastal and off-shore stations during monsoon and non-monsoon season along the west coast of India.

Fig. 6. (a) Distribution of total bacterial phyla as Defined, Candidatus and Unknown taxa, (b) Relative abundance of bacterial phyla (number in parentheses indicates the relative abundance in the non-monsoon and monsoon season) and (c) Distribution of bacterial phyla showing significant statistical difference during the non-monsoon and monsoon season.

Fig. 7. Canonical correspondence analysis (CCA) ordination plot of dominant bacterial genera and environmental variables temperature, salinity, dissolved oxygen (DO), fluorescence, chlorophyll-a (chl-a), primary productivity (PP) and nutrients during (a) non-monsoon and (b) monsoon season.

(a)

235 **(b)**

Fig. 8. Heatmap showing functional pathways during the non-monsoon and monsoon season, (a) predicted based on KEGG database (all categories), (b) depicts pathways related to metabolism and (c) pathways related to genetic, environmental information processes and cellular processing.

1231   Carbohydrate-Active   Enzymes.   Front.   Microbiol.   9,   1864.

1232   https://doi.org/10.3389/fmicb.2018.01864

1233

1234

1235

1236

1237

1238