It's a Trap! Escape from an ancient, ancestral sex chromosome system and implication of *Foxl2* as a primary sex determining gene in a lizard (Anguimorpha; Shinisauridae)

Brendan J. Pinto[1,2,3,*], Stuart V. Nielsen[4,5], Kathryn A. Sullivan[3,6], Ashmika Behere[6], Shannon E. Keating[6], Mona van Schingen-Khan[7], Truong Quang Nguyen[8,9], Thomas Ziegler[10,11], Jennifer Pramuk, Melissa A. Wilson[1,2,12], Tony Gamble[3,4,13,*]

1. School of Life Sciences, Arizona State University, Tempe, AZ USA

2. Center for Evolution and Medicine, Arizona State University, Tempe, AZ USA

3. Department of Zoology, Milwaukee Public Museum, Milwaukee, WI USA

4. Department of Biological Sciences, Museum of Life Sciences, Louisiana State University-Shreveport, Shreveport, LA USA

5. Florida Museum of Natural History, University of Florida, Gainesville, FL USA

6. Department of Biological Sciences, Marquette University, Milwaukee WI USA

7. Federal Agency for Nature Conservation, CITES Scientific Authority, Konstantinstraße 110, 53179 Bonn, Germany

8. Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet Road, Hanoi 10072, Vietnam

9. Graduate University of Science and Technology, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi 10072, Vietnam

10. Cologne Zoo, Riehler Straße 173, 50735 Cologne, Germany

11. Department of Biology, Institute of Zoology, University of Cologne, Zülpicher Straße 47b, 50674 Cologne, Germany

12. Center for Mechanisms of Evolution, Biodesign Institute, Tempe, AZ USA

13. Bell Museum of Natural History, University of Minnesota, St Paul, MN USA

*Correspondence:
brendanjohnpinto@gmail.com
tgamble@geckoevolution.org

Abstract

Although sex determination is ubiquitous in vertebrates, mechanisms of sex determination vary from environmentally- to genetically-influenced. Genetic sex determination is typically accomplished with sex chromosomes via the presence/absence of a specific allele at a genetic locus, which initiates the sex determination signaling network. In vertebrates, well-studied groups like mammals and birds maintain conserved sex chromosome systems (XY and ZW, respectively), while sex chromosomes in many other clades may not be conserved across long evolutionary timeframes. Among squamate reptiles, for example, some groups are relatively stable, like iguanids (Pleurodonta), caenophidian snakes (Caenophidia), and lacertids (Lacertidae) while others are highly variable in their modes of sex determination, such as geckos (Gekkota) and the clade containing chameleons and dragon lizards (Acrodonta). One group inferred to have an evolutionarily stable mode of sex determination is the Anguimorpha, a clade of charismatic taxa, including monitor lizards (*Varanus*), beaded lizards (*Heloderma*), glass lizards (*Ophisaurus*), and the crocodile lizard (*Shinisaurus crocodilurus*). The most recent common ancestor to extant anguimorphs possessed a ZW system that has been retained across the clade. However, determination of sex chromosomes in some anguimorph species has remained elusive including in the endangered, monotypic family of crocodile lizards (Shinisauridae). Here, we use a combination of whole genome re-sequencing and RADseq to demonstrate that *S. crocodilurus* has replaced the ancestral anguimorph ZW system on their LG7 chromosome with a novel ZW system on LG3. The syntenic region that corresponds with this linkage group in chicken is chromosome 9, and this is the first documented use of this syntenic block as a sex chromosome in amniotes. Additionally, this ~1Mb region harbors approximately 10 genes, including a duplication of the well-characterized transcription factor, *Foxl2*—a critical locus for the maintenance of sexual differentiation in vertebrates, and thus a putative primary sex determining gene for *S. crocodilurus*.

Introduction

The evolution of sex determination in vertebrates is impressive in its ability to combine a highly conserved developmental network that can be initiated by quite distinct molecular mechanisms in different species (Bachtrog et al. 2014; Graves, 2008). In vertebrates, sex is commonly determined via either environmental and/or genetic cues at critical points in development. In vertebrate groups that use genetic mechanisms, the most common mechanism is sex chromosomes; either a male or female heterogametic system where the male or female inherits the sex-limited (Y or W) chromosome, respectively (Bachtrog et al. 2014; Gamble et al. 2015). Sex chromosomes have been traditionally identified by comparing male and female karyotypes under the light microscope. The presence of morphological differences between the X and Y (or Z and W) chromosomes (i.e. heteromorphic sex chromosomes) identify a species' sex chromosome system (Stevens, 1905; Bull, 1983). However, many species possess sex chromosomes that cannot be identified via light microscopy because the X and Y (or Z and W) are not morphologically distinguishable from each other (i.e. homomorphic sex chromosomes). Other methods must be employed, such as advanced cytogenetic techniques or high-throughput DNA sequencing technologies, to identify sex chromosome systems in these taxa (Gamble and Zarkower, 2014; Pinto et al. 2022).

Squamate reptiles (lizards and snakes) demonstrate high variability in modes of sex determination: where some clades have conserved, often heteromorphic, sex chromosomes, while others display extraordinary lability in their modes of sex determination and a high incidence of homomorphic sex chromosomes (Gamble et al. 2015, Kratochvíl et al. 2021; Augstenová et al. 2021a). One hypothesis of sex chromosome evolution is that ancient, degenerated sex chromosome systems may act as an "evolutionary trap", where the existence of highly differentiated (i.e. heteromorphic) sex chromosomes preclude transitions to other sex-determining systems (Bull 1983; Bull and Charnov, 1985; Pokorná and Kratochvíl, 2009). The stability of old sex chromosome systems in mammals, birds, caenophidian snakes, and others, provides anecdotal support for this hypothesis (Bull and Charnov, 1985;

Pokorná and Kratochvíl, 2009; Gamble et al. 2015). As more and more sex chromosome transitions are identified, it remains unclear whether all sex chromosomes are destined to become traps. And examples of taxa transitioning away from ancient, degenerated sex chromosome systems are rare (Acosta et al. 2019; Nielsen et al. 2019; Rovatsos et al. 2019a). Previous phylogenetic studies have supported the trap hypothesis in squamates (Pokorná and Kratochvíl, 2009; Gamble et al. 2015), but also suffered from incomplete taxonomic sampling, which likely biased the conclusions. In other words, testing this hypothesis is contingent upon having sufficient data necessary to identify a transition away from a sex chromosome system which typically requires a reference genome to coordinate linkage groups (which are rare in squamates; Pinto et al. 2023), genome-scale data from both sexes (e.g. Vicoso et al. 2013; Gamble et al. 2015; Pinto et al. 2022), and a robust phylogenetic hypothesis to establish relationships within the focal taxa (Nielsen et al. 2019). Thus, the burden of proof is higher for identifying escapes from these putative "evolutionary traps," which may be partially responsible for the dearth of examples.

The sex chromosomes of the infraorder Anguimorpha (lizards including monitor lizards, Gila monsters, alligator lizards, and their allies) have been a topic of interest for many years, likely resulting from the paucity of genetic and cytogenetic data for this group. In recent years, advanced cytogenetic techniques (FISH) have facilitated karyotypic analysis and identification of ZW sex chromosomes in the Gila monster (*Heloderma suspectum*; Pokorná et al. 2014) and Komodo dragon (*Varanus komodoensis*; Pokorná et al. 2016) using leading to expanded interest in studying chromosome evolution in this enigmatic group. More recently, RNAseq and qPCR analysis, in conjunction with draft genomes of these same two anguimorph species (Gila monster; Webster et al. 2023, and Komodo dragon; Lind et al. 2019), have provided some additional insights into this system (Rovatsos et al. 2019b). Namely, the homology of the ZW systems in the anguimorph genera *Abronia*, *Heloderma*, and *Varanus* (Rovatsos et al. 2019b; Webster et al. 2023). The presence of a ZW sex chromosome on the same linkage group—syntenic with chromosome 28 in the chicken genome—in these three genera, spanning the phylogenetic breadth of extant

Anguimorpha, is strong evidence that this is the ancestral sex chromosome system in the clade. Ancient sex chromosome systems, like those ancestral to anguimorphs (115–180 million years old), fit the criteria that should render them as an evolutionary trap (Pokorná and Kratochvíl, 2009; Rovatsos et al. 2019b). However, the sex chromosomes of many anguimorph taxa remain unknown, including the monotypic family Shinisauridae, which is nested deep within the anguimorph phylogeny (Figure 1).

The crocodile lizard (*Shinisaurus crocodilurus*) is the sole living member of the family Shinisauridae and native to small disjunct regions of southeastern China and northern Vietnam (Le and Ziegler, 2003; Huang et al. 2008; Nguyen et al. 2014). It is one of the rarest lizard species in the world and is listed as Endangered in the IUCN Red List (Nguyen et al. 2014). Due to anthropogenic pressures, populations have experienced severe declines in recent years with less than 1000 individuals in the wild in China and less than 100 adults in Vietnam (Huang et al. 2008; van Schingen et al. 2014a). They are semi-aquatic habitat specialists and depend upon clean streams in broadleaf evergreen forest (Ning et al. 2006; van Schingen et al. 2016a) and their restricted ecological niche is predicted to all but disappear due to climate change by the end of this century (Li et al. 2013; van Schingen et al. 2014a; see also van Schingen-Khan et al. 2022). Habitat destruction threatens remaining populations, as well as being collected for food and the international pet trade (Huang et al. 2008; van Schingen et al. 2014b; van Schingen et al. 2016a). Although still recognized as a single species, there exist several conservation units, with *S. crocodilurus vietnamensis* currently only known from Vietnam and the nominate subspecies from China consisting of several distinct lineages (van Schingen et al. 2016b; Ngo et al. 2020; Nguyen et al. 2022). Crocodile lizards do not have a clear sexual dimorphism. While morphological traits, such as coloration or body morphometry may provide an indication of the sex, a robust method to reliably identify the sex of individuals is lacking (van Schingen et al. 2016b). Relevant to the present study, examination of male and female *S. crocodilurus* karyotypes have revealed no heteromorphic sex chromosomes (Zhang et al. 1996; Augstenová et al. 2021b). To identify sex chromosomes in this species, we analyzed whole-genome re-sequencing data for more than 50 individual crocodile lizards (held animals originating

from China; Xie et al. 2022) using whole-genome re-sequencing and, in parallel, restriction site-associated DNA sequencing (RADseq) data for 13 additional individuals (held animals originating from both China and Vietnam) to show that the sex determining system in *S. crocodilurus* is a novel ZW system that has eluded previous analyses, at least in part, due to the small size (<1Mb) of its sex determining region (SDR).

Figure 1: Summary of current anguimorph sex chromosome knowledge summarized from Rovatsos et al. (2019) indicated by blue and green tips/branches); information identified in this study indicated by red tips/branches and what remains unknown across the phylogeny indicated by black tips/branches. Phylogeny from TimeTree using a representative species from each clade (Kumar et al. 2017) and visualized using Figtree [v1.4.4] (http://tree.bio.ed.ac.uk/software/figtree/). "Gg" stands for chicken (*Gallus gallus*) linkage group.

Methods

WGS analysis

We downloaded whole genome resequencing reads from NCBI SRA for multiple male and female individuals (see *Data Availability* for accessions). We constructed a Snakemake [v6.10.0] (Mölder et al. 2021) workflow in an isolated conda environment [v4.11.0] (https://docs.anaconda.com/) containing relevant packages: BBmap [v38.93] (Bushnell, 2014), FastQC [v0.11.9] (Andrews, 2010), Freebayes [v1.3.5] (Garrison and Marth, 2012), GFF utilities [v0.10.1] (Pertea and Pertea, 2020), Minimap2 [v2.22] (Li, 2018), Mosdepth [v0.3.2] (Pedersen and Quinlan, 2018), MultiQC [v1.11] (Ewels et al. 2016), Parallel [v20211022] (Tange, 2018), pixy [v1.2.5.beta1] (Korunes and Samuk, 2021), Sambamba [v0.8.1] (Tarasov et al. 2015), Samtools [v1.12] (Li et al. 2009), seqkit [v0.11.0] (Shen et al. 2016), STACKS [v2.6.0] (Catchen et al. 2013), Trim Galore! [v0.6.7] (https://doi.org/10.5281/zenodo.5127899), and vcftools [v0.1.15] (Danecek et al. 2011). To process the raw sequencing data, we trimmed adapters and low-quality regions using Trim Galore!, then removed PCR duplicates using BBmap. Quality assessment using FastQC and MultiQC was conducted at each step and we subsequently removed samples with fewer than 5 million PE reads after filtering PCR duplicates. The final dataset possessed 28 male and 23 female individuals. We proceeded to map reads for each individual to the female reference genome (Xie et al. 2022) with minimap2 and calculated read depth and read mapping statistics using mosdepth and samtools, respectively. Then generated an all-sites vcf file with freebayes-parallel. Lastly, we calculated Weir and Cockerham (1984) F_{ST} and nucleotide diversity statistics using pixy at 500kb resolution and, for LG3 only, also at 100kb resolution.

RADseq analysis

In a similar fashion, we analyzed a modest dataset of restriction site-associated DNA sequencing (RADseq) for four males and six females (Supplemental Table 1). We followed a modified protocol from Etter et al. (2011) as outlined in Gamble et al. (2015). Libraries were size-selected into 300- to 600-bp fragments using magnetic beads in a PEG/NaCl buffer (Rohland and Reich, 2012), then pooled and sequenced using paired-

end 150-bp reads on an Illumina® HiSeqX at Psomagen®. Similarly to the WGS data, raw illumina reads were trimmed using Trim Galore!, PCR duplicates removed using BBmap, and mapped to the reference genome using minimap2 with quality assessment using FastQC and MultiQC at each step. Diverging from the WGS pipeline, we processed the RADseq data herein using the `refmap.pl` script included in the STACKS suite and calculated Weir and Cockerham (1984) F_{ST} using VCFtools.

Validation of the putative ZW system in Shinisaurus crocodilurus

We generated a dataset of '*in silico* poolseq' reads by subsampling each read file to 10 million reads (20 million per sample) using seqkit and combined into male and female pools. We analyzed the pools using Pooled Sequencing Analyses for Sex Signal [PSASS; v3.1.0] (https://doi.org/10.5281/zenodo.3702337). We then generated PCR primers targeting the annotated version of *Foxl2*'s second exon [FOXL2-ex2-F2 5' – CAGAGCTCGTCCCATTCACTT – 3' and FOXL2-ex2-R2 5' – GAGAGATGTACCACCGGGAG – 3'] and sequenced the resultant amplicon using Sanger sequencing (Psomagen).

Genome Annotation

We used previously lifted over annotations (Pinto et al. 2023; https://doi.org/10.6084/m9.figshare.20201099.v1) from the draft genome of a male *S. crocodilurus* (Gao et al. 2017) to the new, unannotated female reference genome (Xie et al. 2022; GCA_021292165.1). We pulled coding transcripts from the genome using GFF Utilities. We used the 10 genes within the putative ~1Mb SDR on LG3 to perform a high-stringency tBLASTx query (Altschul et al. 1990) to the chicken genome on Ensembl (Howe et al. 2020) with a word size of 3, maximum of 10 hits, e-value cutoff of 1e^{-50}, using BLOSUM62 scoring matrix. These queries received hits on 7 of the 10 total genes (Table 1).

Results

Across WGS experiments, read mapping efficiency ranged from 80.60% (for SRR5019740) to 99.40% (for SRR14583318). In RADseq experiments, read mapping efficiency ranged from 96.03% to 98.67%. Across both experiments, only one F_{ST} peak appeared in both datasets (Supplemental Figure 1). The poor resolution of the RADseq dataset, compared with the whole genome resequencing data, likely results from the small number of individuals used and the relatively small size of the non-recombining portion of the sex chromosome. Comparing M/F F_{ST} values does not necessarily diagnose which sex is heterogametic (i.e. XY vs. ZW). Therefore, we composed a dataset of '*in silico* poolseq' reads to identify an excess of female-associated SNPs aligning to the previously identified region of high M/F F_{ST} (Supplemental Figure 2). Taken together, these data suggest that *S. crocodilurus* possesses a female heterogametic system (ZW) with an SDR located in a ~900kb region on LG3.

Upon further investigation of the SDR, we identified a total of 10 genes annotated within this region. To better characterize these genes, we BLAST-ed each to the chicken genome. We recovered high-quality BLAST hits for seven of the 10 annotated *S. crocodilurus* SDR genes in chicken (Table 1). Six out of the seven queries hit genes located on chicken chromosome 9, while the other landed on a chicken chromosome 30 (Table 1). In our poolseq analysis, one of these genes possessed half the read depth in females relative to males and, upon closer inspection, we identified a duplicated, unannotated copy of that gene Forkhead Box L2 (*Foxl2*) also located within the SDR. We included this *Foxl2* copy in a BLAST search against chicken, where it was again identified as a *Foxl2* homolog (Table 1). We also BLAST-ed *Foxl2* to the earlier male *S. crocodilurus* draft genome (Gao et al. 2017) and found only a single copy of *Foxl2* in this genome matching one copy in the updated reference genome with 100% sequence identity, consistent with both (1) the duplicated version being W-specific and (2) the female reference being chimeric for Z and W alleles (Xie et al. 2022). Lastly, we generated a gene tree using *Foxl2* copies from across reptiles to confirm its duplicated origination was within Shinisauridae (Supplemental Figure 3). Thus, in the chimeric female reference genome, this putative W-linked *Foxl2* copy was located approximately

70kb upstream of the annotated Z-linked copy of *Foxl2* on the other side of an assembly gap.

Our WGS data used in poolseq and other analyses were restricted to only individuals from Chinese populations to reduce the influence of population-specific demographic processes (Xie et al. 2022). To include the less-numerous Vietnam samples, we generated PCR primers for a segment of *Foxl2*'s second exon and Sanger sequenced multiple females (Vietnam) and males (China and Vietnam) (Supplemental Figure 4). We identified one SNP in the female Vietnamese samples in this region and tested its association with sex using Fisher's exact test (p-value = 0.0047***). Thus, the ZW SDR containing *Foxl2* appears to be conserved between populations of *S. crocodilurus* from both China and Vietnam.

Table 1: Top tBLASTx hits in chicken for each of the genes present in the *Shinisaurus* 900kb-SDR. The duplicated *Foxl2* copy is dubbed ENSGALP00000033127-W.

Shinisaurus Gene ID	Chicken Gene Id	Location (Chicken)	E-value
ENSACAP00000003394-D1	ENSGALG00000026187	9:6115461-6115832	1.42E-95
ENSGALP00000008531-D1	*RBP1*	9:6110534-6110713	7.54E-54
ENSACAP00000003392-D1		***No hits.***	
ENSACAP00000003371-D1	ENSGALG00000034575	30:1402166-1402369	2.51E-78
ENSACAP00000003355-D1	ENSGALG00000005357	9:6041607-6041822	0
ENSACAP00000003221-D1	ENSGALG00000005367	9:6037018-6037164	1.03E-80
ENSGALP00000033127-W	***FOXL2W***	**9:5875297-5875587**	**2.44E-76**

ENSGALP00000033127-D1	*FOXL2*	**9:5875297-5875587**	**6.78E-77**
ENSACAP00000003172-D1	*PIK3CB*	9:5800560-5800790	0
ENSGALP00000040175-D1		*No hits.*	
ENSACAP00000002765-D1		*No hits.*	

Discussion

Escaping the "Evolutionary Trap"

An open question within sex chromosome evolution is whether ancient, degenerated sex chromosomes act as evolutionary traps (Pokorná and Kratochvíl, 2009; Nielsen et al. 2019; Pinto et al. 2023). The most recent common ancestor of extant anguimorphs is thought to have possessed a ZW system on the linkage group syntenic with chicken chromosome 28, which is located on the distal region of LG7 in in the *S. crocodilurus* reference genome (Rovatsos et al. 2019b; Webster et al. 2023). The sex determining region (SDR) in *S. crocodilurus* is located on LG3, a region syntenic with chicken chromosome 9. To our knowledge, this is the first demonstration in a tetrapod of the syntenic region of chicken chromosome 9 being recruited in a sex determining role (Kratochvíl et al. 2021), lending further support to the idea that all chromosomes will likely be recruited into a sex determining role given thorough enough phylogenetic sampling (Graves and Peichel, 2010; Hodgkin, 2002; O'Meally et al. 2012; Pinto et al. 2022).

It is clear from these genomic data that *S. crocodilurus* possesses a distinct sex chromosome system from all other known anguimorphs. Unlike the case of Corytophanidae and other pleurodonts, where phylogenetic relationships among taxa were inconclusive (Nielsen et al. 2019), the relationship of *S. crocodilurus* to all other anguimorphs is far less divisive. Indeed, *S. crocodilurus* is well-supported as nested within Anguimorpha—either sister to Varanidae as a member of the "Paleoanguimorpha" (Burbrink et al. 2020) or as sister to a clade containing Varanidae and Lanthanotidae (Singhal et al. 2021), depending on taxonomic sampling. Thus, assuming the hypothesis that an ancient origin of the ZW sex chromosome system possessed by extant *Varanus*, *Heloderma*, and *Abronia* is correct, then *S. crocodilurus* has successfully escaped the evolutionary trap of their ancestral, degenerated sex chromosome system—a system nearly as ancient as those systems found in both mammals and birds (Rovatsos et al. 2019b; Webster et al. 2023). It is worth noting that there remains another putative escape from the ancestral anguimorph sex chromosome system in *Anguis* that has yet to be explored further (Rovatsos et al. 2019b) and more

recent phylogenetic work has implicated that Corytophanidae is likely nested somewhere within other pleurodonts, rather than being sister to all other species (Burbrink et al. 2020; Singhal et al. 2021). This suggests that there are a minimum of two evolutionary escapes within Toxicofera (snakes, iguanians, and anguimorphs)—and perhaps even two within the infraorder Anguimorpha alone.

Figure 2: Identification of the ZW sex chromosome system in *Shinisaurus crocodilurus*. (A) Whole genome F_{ST} scan with a clear peak in a ~1Mb region on LG3. The square block on LG7 is syntenic with the sex-determining region in *Varanus* and *Heloderma* (Webster et al. 2023). (B) Isolation and magnification of LG3 F_{ST} peak. (C) Modest increase in male, relative to female, nucleotide diversity and (D) decrease in male/female read depth in the region corresponding to the F_{ST} peak on LG3.

Primary Sex Determination in Shinisauridae

In many vertebrate groups where the primary sex determiner (PSD) is known, a relatively short list of commonly-recruited PSDs have been identified (i.e. the 'usual

suspects'; Adolfi et al. 2021; Dor et al. 2019; Herpin and Schartl, 2015). Indeed, the same genes, or paralogs, have been independently co-opted to function as the PSD in many taxa, including *Sox3* in placental mammals and some medaka (members of the *Oryzias celebensis* and *O. javanicus* groups); *Amh* in tilapia, northern pike, and potentially other anguimorphs (Li et al. 2015; Myosho et al. 2015; Pan et al. 2019; Rovatsos et al. 2019; Webster et al. 2023; and see Pan et al. 2021 for recent review); and *Dmrt1* in birds, a frog (*Xenopus laevis*), tongue sole, and other medaka fish (members of the *Oryzias latipes* group) (Chen et al. 2014; Ioannidis et al. 2021; Matsuda et al. 2002; Nanda et al. 2002; Smith et al. 2009).

This is the first time Forkhead Box L2 (*Foxl2*) has been implicated as a PSD in a vertebrate, although it has been predicted to be one (Ma et al. 2022). *Foxl2* is a direct transcriptional activator of aromatase, involved in development of the ovaries, and its loss in mice during embryogenesis leads to abnormal ovarian development and infertility (Fleming et al. 2010; Pannetier et al. 2006; Schmidt et al. 2004; Uda et al. 2004). After primary sex determination and sexual development has concluded, *Dmrt1* and *Foxl2* antagonize each other transcriptionally in gonadal tissue, where sustained *Dmrt1* and *Foxl2* expression is required for adult maintenance of testis and ovary tissue, respectively (Garcia-Ortiz et al. 2009; Matson et al. 2011; Uhlenhaut et al. 2009). Importantly, *Dmrt1* has been recruited to act as a primary sex determining gene in multiple taxa, while *Foxl2* has remained mysteriously absent from this list—with the singular putative exception being recently described in some species of bivalve mollusks (Han et al. 2022). Thus, the identification of both *Foxl2* and a duplicated *Foxl2* copy in the W-limited region of the *Shinisaurus* genome confirms the expanded list of the "usual suspects" that can act as the PSD in vertebrates.

Pragmatically, the identification of a novel ZW system in *S. crocodilurus* may present an important juncture in the conservation efforts of this endangered lizard species, that are urgently needed (Nguyen et al. 2014). Body morphometrics in mature specimens may provide an indication of the sex, i.e. males tend to have a relatively larger head, relative to abdomen length than females (van Schingen et al. 2016b).

However, definitive sexually dimorphic characters are lacking in the species, especially in hatchlings, juveniles, and subadults. Therefore, a molecular genetic sex test could assist in well-managed captive breeding efforts in this species (Ziegler et al. 2019). This is vital as it's estimated only ~1,000 individuals remain in the wild populations in China and Vietnam during the last census (Huang et al. 2008; van Schingen et al. 2016a), while loss of remaining habitats as well as poaching is considered ongoing.

In conclusion, using a combination of sequencing and validation techniques we identified the elusive ZW system in the endangered crocodile lizard, *Shinisaurus crocodilurus*. This ZW system is located on LG3 and, although interpretation inherits strong reference bias, the SDR appears to be <1Mb in size and contains approximately 10 genes. One of these genes, *Foxl2*, possesses a duplicated copy and is important in ovarian development and fertility in vertebrates. Because of its sequence conservation (either strictly age-related or via gene conversion) and proximity to the original Z copy of *Foxl2*, we hypothesize that if *Foxl2* is the PSD in this system, it's likely a gene dosage-dependent mechanism, where ZW females possess three copies of *Foxl2* instead of the two copies of ZZ males. Our proposed mechanism would essentially be the inverse of *Dmrt1* sex determination in birds, where a lack of *Dmrt1* on the W decreases *DMRT1* expression in females, allowing for *Foxl2* to proceed with ovarian development (Ioannidis et al. 2021; Smith et al. 2009). Here, extra gene copies of *Foxl2* increase *FOXL2* expression to initiate ovarian development in the developing gonad. This information may play a vital role in conservation efforts of this species and this information should be adopted in ongoing captive breeding work (Ziegler et al. 2019).

Data Availability:

The data used in this study is available on NCBI, SRA accessions for WGS data are: SRR14583317, SRR14583321, SRR14583324-26, SRR14583330, SRR14583333, SRR14583340-49, SRR14583351, SRR14583353-54, SRR14583356, SRR14583360-66, SRR5019733-45, SRR14583318-20, SRR14583322-23, SRR14583331, SRR14583334-39, SRR14583346, SRR14583350, SRR14583352, SRR14583355, SRR14583357-59. RADseq and HiC data generated in this study are available on SRA under BioProject PRJNA975696 (SAMN35342808-17).

Code used in the WGS analysis is available on GitHub: https://github.com/DrPintoThe2nd/Shinisaurus_ZW.

Acknowledgements:

The authors would like to acknowledge Research Computing at Arizona State University for providing high-performance computing and storage resources that have contributed to the research results reported within this paper (http://www.researchcomputing.asu.edu). We thank Anna Rauhaus (Cologne Zoo) for her help with the application and preparation of tissue sending and the Woodland Park Zoo for their respective assistance. Many thanks CITES Management Authority of Vietnam for issuing permits (CITES permits No. 13VN1246N/CT-KL and 16VN0920N/CT-KL).This work was funded by the Morris Animal Foundation (Study grant D19ZO-021) for their generous funding of this project (T.G.) and also supported by the National Institute of General Medical Sciences (NIGMS) of the National Institutes of Health grant R35GM124827 (M.A.W.).

References

Adolfi, M. C., Herpin, A., & Schartl, M. (2021). The replaceable master of sex determination: bottom-up hypothesis revisited. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 376(1832):20200090.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol.* 215: 403-410.

Andrews S. 2010. FastQC: A quality control tool for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Augstenová, B., Pensabene, E., Veselý, M., Kratochvíl, L., Rovatsos, M. 2021a. Are geckos special in sex determination? Independently evolved differentiated ZZ/ZW sex chromosomes in carphodactylid geckos. *Genome Biology and Evolution*, 13(7), evab119.

Augstenová, B., Pensabene, E., Kratochvíl, L., Rovatsos, M. 2021b. Cytogenetic evidence for sex chromosomes and karyotype evolution in anguimorphan lizards. *Cells*, 10(7), 1612.

Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman TL, Hahn MW, Kitano J, Mayrose I, Ming R, Perrin N, Ross L, Valenzuela N, Vamosi JC. 2014. Sex determination: Why so many ways of doing it? *PLoS Biol.* 12: e1001899.

Bull, JJ. 1983. Evolution of Sex Determining Mechanisms. Benjamin Cummings: Menlo Park, CA.

Bull J, Charnov E. 1985. On irreversible evolution. *Evolution* 39, 1149–1155.

Burbrink, F. T., Grazziotin, F. G., Pyron, R. A., Cundall, D., Donnellan, S., Irish, F., Keogh, J. S., Kraus, F., Murphy, R. W., Noonan, B., Raxworthy, C. J., Ruane, S., Lemmon, A. R., Lemmon, E. M., Zaher, H. (2019). Interrogating Genomic-Scale Data for Squamata (Lizards, Snakes, and Amphisbaenians) Shows no Support for Key Traditional Morphological Relationships. Systematic Biology, 69(3), 502–520.

Bushnell B. 2014. BBMap: a fast, accurate, splice-aware aligner (No. LBNL-7065E). Lawrence Berkeley National Lab (LBNL), Berkeley, CA (United States).

Catchen J, Hohenlohe P, Bassham S, Amores A, Cresko W. 2013. Stacks: an analysis tool set for population genomics. *Mol Ecol.* 22: 3124-3140.

Chen, S., Zhang, G., Shao, C., Huang, Q., Liu, G., Zhang, P., Song, W., An, N., Chalopin, D., Volff, J.-N., Hong, Y., Li, Q., Sha, Z., Zhou, H., Xie, M., Yu, Q., Liu, Y., Xiang, H., Wang, N., … Wang, J. (2014). Whole-genome sequence of a flatfish provides insights into ZW sex chromosome evolution and adaptation to a benthic lifestyle. *Nature Genetics*. 46(3):253–260.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics*. 27: 2156-2158.

Dor L, Shirak A, Kohn YY, Gur T, Weller JI, Zilberg D, Seroussi E, Ron M. 2019. Mapping of the sex determining region on linkage group 12 of guppy (*Poecilia reticulata*). *G3*. 9: 3867-3875.

Etter P, Bassham S, Hohenlohe P, Johnson E, Cresko W. 2011. SNP discovery and genotyping for evolutionary genetics using RAD sequencing. In: Orgogozo V, Rockman M, editors. Molecular methods for evolutionary genetics. New York: Springer. p. 157-178.

Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*. 32: 3047-3048.

Fleming, N. I., Knower, K. C., Lazarus, K. A., Fuller, P. J., Simpson, E. R., & Clyne, C. D. (2010). Aromatase Is a Direct Target of *FOXL2*: C134W in Granulosa Cell Tumors via a Single Highly Conserved Binding Site in the Ovarian Specific Promoter. *PLoS ONE*. 5(12):e14389.

Gamble T, Coryell J, Ezaz T, Lynch J, Scantlebury D, Zarkower D. 2015. Restriction site-associated DNA sequencing (RAD-seq) reveals an extraordinary number of transitions among gecko sex-determining systems. *Mol Biol Evol*. 32: 1296-1309.

Gamble T, Castoe TA, Nielsen SV, Banks JL, Card DC, Schield DR, Schuett GW, Booth W. 2017. The discovery of XY sex chromosomes in a *Boa* and *Python*. *Curr Biol*. 27: 2148-2153.

Gamble T, Zarkower D. 2014. Identification of sex-specific molecular markers using restriction site-associated DNA sequencing. *Molecular Ecology Resources*. 14:902–913.

Gao J, Li Q, Wang Z, Zhou Y, Martelli P, Li F, ... Zhang G. 2017. Sequencing, de novo assembling, and annotating the genome of the endangered Chinese crocodile lizard *Shinisaurus crocodilurus*. *GigaScience*. 6: gix041.

Garcia-Ortiz JE, Pelosi E, Omari S, Nedorezov T, Piao Y, Karmazin J, ... Ottolenghi C. 2009. *Foxl2* functions in sex determination and histogenesis throughout mouse ovary development. *BMC Dev Biol*. 9: 36.

Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing. *arXiv*. 1207.3907. https://arxiv.org/abs/1207.3907

Graves J. 2008. Weird animal genomes and the evolution of vertebrate sex and sex chromosomes. *Ann Rev Genet*. 42: 565-586.

Graves J, Peichel C. 2010. Are homologies in vertebrate sex determination due to shared ancestry or to limited options? *Genome Biol*. 11: 205. doi:10.1186/gb-2010-11-4-205

Han, W., Liu, L., Wang, J., Wei, H., Li, Y., Zhang, L., ... Wang, S. (2022). Ancient homomorphy of molluscan sex chromosomes sustained by reversible sex-biased genes and sex determiner translocation. *Nat Ecol Evol*. 1-16.

Herpin A, Schartl M. 2015. Plasticity of gene- regulatory networks controlling sex determination: of masters, slaves, usual suspects, newcomers, and usurpators. *EMBO Rep*. 16:1260-1274.

Hodgkin J. 2002. Exploring the envelope: systematic alteration in the sex-determination system of the nematode *Caenorhabditis elegans*. *Genetics*. 162: 767-780.

Howe KL, Achuthan P, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Azov AG, Bennett R, Bhai J, Billis K, Boddu S, Charkhchi M, Cummins C, Da Rin Fioretto L, Davidson C, Dodiya K, El Houdaigui B, Fatima R, … Flicek P. 2020. Ensembl 2021. *Nucleic Acids Res*. 49: 884-891. doi:10.1093/nar/gkaa942

Huang C.M., Yu H., Wu Z.J., Li Y.B., Wei F.W., Gong M.H. (2008). Population and conservation strategies for the Chinese crocodile lizard (*S. crocodilurus*) in China. *Animal Biodiversity and Conservation*. 31(2):63-70.

Ioannidis, J., Taylor, G., Zhao, D., Liu, L., Idoko-Akoh, A., Gong, D., Lovell-Badge, R., Guioli, S., McGrew, M. J., & Clinton, M. (2021). Primary sex determination in birds depends on DMRT1 dosage, but gonadal sex does not determine adult secondary sex characteristics. *PNAS*. 118(10).

Korunes KL, Samuk K. 2021. pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of missing data. *Mol Ecol Res*. 21: 1359-1368.

Kratochvíl, L., T. Gamble, and M. Rovatsos. 2021. Sex chromosome evolution among amniotes: Is the origin of sex chromosomes non-random? Philosophical Transactions of the Royal Society B: Biological Sciences 376:20200108.

Kumar S, Stecher G, Suleski M, Hedges SB. (2017). TimeTree: a resource for timelines, timetrees, and divergence times. *Mol Biol Evol*. 34:1812–1819

Le K.Q., Ziegler T. (2003). First record of the Chinese crocodile lizard from outside of China: report on a population of *S. crocodilurus* Ahl, 1930 from North-Eastern Vietnam. *Hamadryad*. 27(2):193-199.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, ... Durbin R. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics*. 25:2078-2079.

Li H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*. 34:3094-3100.

Li, M., Sun, Y., Zhao, J., Shi, H., Zeng, S., Ye, K., Jiang, D., Zhou, L., Sun, L., Tao, W., Nagahama, Y., Kocher, T. D., & Wang, D. (2015). A Tandem Duplicate of Anti-Müllerian Hormone with a Missense SNP on the Y Chromosome Is Essential for Male Sex Determination in Nile Tilapia, *Oreochromis niloticus*. *PLoS Genetics*. 11(11):e1005678.

Li, X., Tian, H., Wang, Y., Li, R., Song, Z., Zhang, F., ... & Li, D. (2013). Vulnerability of 208 endemic or endangered species in China to the effects of climate change. *Regional Environmental Change*, *13*, 843-852.

Ma, X., Liu, F., Chen, Q., Sun, W., Shen, J., Wu, K., Zheng, Z., Huang, J., Chen, J., Qian, G., & Ge, C. (2022). *Foxl2* is required for the initiation of the female pathway in a temperature-dependent sex determination system in *Trachemys scripta*. *Development*. 149(13).

Matson, C. K., Murphy, M. W., Sarver, A. L., Griswold, M. D., Bardwell, V. J., & Zarkower, D. (2011). *DMRT1* prevents female reprogramming in the postnatal mammalian testis. *Nature*. 476(7358), 101–104.

Matson, C. K., Zarkower, D. (2012). Sex and the singular DM domain: insights into sexual regulation, evolution and plasticity. *Nature Reviews Genetics*. 13(3):163-174.

Matsuda, M., Nagahama, Y., Shinomiya, A., Sato, T., Matsuda, C., Kobayashi, T., Morrey, C. E., Shibata, N., Asakawa, S., Shimizu, N., Hori, H., Hamaguchi, S., & Sakaizumi, M. (2002). DMY is a Y-specific DM-domain gene required for male development in the medaka fish. *Nature*. 417(6888):559–563.

Mölder F, Jablonski KP, Letcher B, Hall MB, Tomkins-Tinch CH, Sochat V, Forster J, Lee S, Twardziok SO, Kanitz A, Wilm A, Holtgrewe M, Rahmann S, Nahnsen S, Köster J. 2021. Sustainable data analysis with Snakemake. *F1000Res*. 10:33.

Nanda, I., Kondo, M., Hornung, U., Asakawa, S., Winkler, C., Shimizu, A., Shan, Z., Haaf, T., Shimizu, N., Shima, A., Schmid, M., & Schartl, M. (2002). A duplicated

copy of *DMRT1* in the sex-determining region of the Y chromosome of the medaka, *Oryzias latipes*. *PNAS*. 99(18):11778–11783.

Ngo, H. T., Nguyen, T. T., Le, M. D., van Schingen-Khan, M., Nguyen, T. Q., Rauhaus, R., Vences, M. & T. Ziegler (2020): Genetic screening of captive crocodile lizards (*Shinisaurus crocodilurus*) in Europe. – Der Zoologische Garten 88: 17-30.

Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2014). IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and Evolution*. 32(1), 268–274.

Nguyen TQ, Hamilton P, Ziegler T. 2014. *Shinisaurus crocodilurus*. *The IUCN Red List of Threatened Species*. doi:10.2305/IUCN.UK.2014-1.RLTS.T57287221A57287235.en. Accessed on 04 March 2022.

Nguyen, T. T., Ngo, H. T., Ha, Q. Q., Nguyen, T. Q., Le, T. Q., Nguyen, S. H., Pham, C. T., Ziegler, T., van Schingen-Khan, M. & M. D. Le (2022): Molecular phylogenetic analyses and ecological niche modeling provide new insights into threats to the endangered Crocodile Lizard (*Shinisaurus crocodilurus*). - Frontiers of Biogeography 2022, 14.1, e54779

Nielsen SV, Guzmán-Mendez IA, Gamble T, Blumer M, Pinto BJ, Kratochvíl L, Rovatsos M. 2019. Escaping the evolutionary trap? Sex chromosome turnover in basilisks and related lizards (Corytophanidae: Squamata). *Biol Lett*. 15: 20190498.

Ning J., Huang C., Yu H., Dai D., Wu Z., Zhong Y. (2006). Summer habitat characteristics of the Chinese Crocodile Lizard (*Shinisaurus crocodilurus*) in the Loukeng Nature Reserve, Guangdong. *Zoological Research*. 27:419-426.

O'Meally D, Ezaz T, Georges A, Sarre SD, Graves JAM. 2012. Are some chromosomes particularly good at sex? Insights from amniotes. *Chromosome Res*. 20: 7-19.

Pan Q, Feron R, Yano A, Guyomard R, Jouanno E, Vigouroux E, Wen M, Busnel JM, Bobe J, Concordet JP, … Guiguen Y. 2019. Identification of the master sex determining gene in Northern pike (*Esox lucius*) reveals restricted sex chromosome differentiation. *PLoS Genetics*. 15: e1008013. doi:10.1371/journal.pgen.1008013

Pan Q, Kay T, Depincé A, Adolfi M, Schartl M, Guiguen Y, Herpin A. 2021. Evolution of master sex determiners: TGF-β signalling pathways at regulatory crossroads. *Phil Trans Roy Soc B*. 376:20200091.

Pannetier, M., Fabre, S., Batista, F., Kocer, A., Renault, L., Jolivet, G., Mandon-Pépin, B., Cotinot, C., Veitia, R., & Pailhoux, E. (2006). FOXL2 activates P450 aromatase gene transcription: towards a better characterization of the early steps of mammalian ovarian development. *Journal of Molecular Endocrinology*. 36(3):399–413.

Pedersen BS, Quinlan AR. 2018. Mosdepth: Quick coverage calculation for genomes and exomes. *Bioinformatics*. 34: 867-868.

Pertea G, Pertea M. 2020. GFF utilities: GffRead and GffCompare. *F1000Research*. 9.

Pinto BJ, Keating SE, Nielsen SV, Scantlebury DP, Daza JD, Gamble T. (2022). Chromosome-level genome assembly reveals dynamic sex chromosomes in Neotropical leaf-litter geckos (Sphaerodactylidae: *Sphaerodactylus*). *Journal of Heredity*. 113(3): 272-287.

Pinto, B. J., Gamble, T., Smith, C. H., Wilson, M. A. (2023). A lizard is never late: squamate genomics as a recent catalyst for understanding sex chromosome and microchromosome evolution. *Journal of Heredity. In press.*

Pokorná, M., Kratochvíl, L. (2009). Phylogeny of sex-determining mechanisms in squamate reptiles: are sex chromosomes an evolutionary trap? *Zoological Journal of the Linnean Society*, 156(1), 168–183.

Rohland N, Reich D. 2012. Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture. *Genome Res.* 22: 939-46.

Rovatsos, M., Farkačová, K., Altmanová, M., Johnson Pokorná, M., Kratochvíl, L. (2019a). The rise and fall of differentiated sex chromosomes in geckos. *Molecular Ecology*, 28(12), 3042–3052.

Rovatsos M, Rehák I, Velenský P, Kratochvíl L. 2019b. Shared Ancient Sex Chromosomes in Varanids, Beaded Lizards, and Alligator Lizards. *Mol Biol Evol.* 36: 1113-20.

Schmidt, D., Ovitt, C. E., Anlag, K., Fehsenfeld, S., Gredsted, L., Treier, A.-C., & Treier, M. (2004). The murine winged-helix transcription factor *Foxl2* is required for granulosa cell differentiation and ovary maintenance. *Development.* 131(4):933–942.

Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A cross-platform and ultrafast toolkit for fasta/q file manipulation. *PLoS One.* 11: e0163962. doi:10.1371/journal.pone.0163962

Shumate A, Salzberg SL. 2021. Liftoff: Accurate mapping of gene annotations. *Bioinformatics.* 12: 1639-1643.

Singhal, S., Colston, T. J., Grundler, M. R., Smith, S. A., Costa, G. C., Colli, G. R., ... Rabosky, D. L. (2021). Congruence and conflict in the higher-level phylogenetics of squamate reptiles: an expanded phylogenomic perspective. *Systematic Biology*, 70(3), 542-557.

Smith, C. A., Roeszler, K. N., Ohnesorg, T., Cummins, D. M., Farlie, P. G., Doran, T. J., & Sinclair, A. H. (2009). The avian Z-linked gene DMRT1 is required for male sex determination in the chicken. *Nature.* 461(7261):267-271.

Stevens NM. 1905. A study of the germ cells of *Aphis rosae* and *Aphis oenotherae. J Exp Zool.* 2: 313-333.

Tange O. 2018. Gnu parallel 2018. pp. 112. doi:10.5281/zenodo.1146014

Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. 2015. Sambamba: fast processing of NGS alignment formats. *Bioinformatics.* 31: 2032-2034.

Uda, M., Ottolenghi, C., Crisponi, L., Garcia, J. E., Deiana, M., Kimber, W., Forabosco, A., Cao, A., Schlessinger, D., & Pilia, G. (2004). *Foxl2* disruption causes mouse ovarian failure by pervasive blockage of follicle development. *Human Molecular Genetics.* 13(11):1171–1181.

Uetz P, Freed P, Aguilar R, Hošek J. (eds.) 2021. The Reptile Database, http://www.reptile-database.org, accessed 02/18/2022.

Uhlenhaut NH, Jakob S, Anlag K, Eisenberger T, Sekido R, Kress J, Treier AC, Klugmann C, Klasen C, Holter NI, Riethmacher D, Schütz G, Cooney AJ, Lovell-Badge R, Treier M. 2009. Somatic sex reprogramming of adult ovaries to testes by *FOXL2* ablation. *Cell.* 139: 1130-1142.

van Schingen M., Ihlow F., Nguyen T.Q., Ziegler T., Bonkowski M., Wu Z., Rödder D. (2014a): Potential distribution and effectiveness of the protected area

network for the crocodile lizard, *Shinisaurus crocodilurus* (Reptilia: Squamata: Sauria). *Salamandra*. 50(2):71-76.

van Schingen M., Pham C.T., Thi H.A., Bernardes M., Hecht V., Nguyen T.Q., Bonkowski M., Ziegler T. (2014b). Current status of the crocodile lizard *Shinisaurus crocodilurus* Ahl, 1930 in Vietnam with implications for conservation measures. *Revue suisse de Zoologie*. 121(3):1-15.

van Schingen M., Ha Q.Q., Pham C.T., Le T.Q., Nguyen T.Q., Bonkowski M., Ziegler T. (2016a). Discovery of a new crocodile lizard population in Vietnam: Population trends, future prognoses and identification of key habitats for conservation. *Revue suisse de Zoologie*. 123(2):241-251.

van Schingen, M., Duc Le, M., Thi Ngo, H., The Pham, C., Quy Ha, Q., Quang Nguyen, T., Ziegler, T. (2016b). Is there more than one Crocodile Lizard? An Integrative Taxonomic Approach Reveals Vietnamese and Chinese Shinisaurus crocodilurus Represent Separate Conservation and Taxonomic Units. *Der Zoologische Garten*. 85(5):240-260.

van Schingen-Khan, M., Barthel, L. M. F., Pham, D. T. K., Pham, C. T., Nguyen, T. Q., Ziegler, T. & M. Bonkowski (2022): Will climatic changes affect the Vietnamese crocodile lizard? Seasonal variation in microclimate and activity pattern of *Shinisaurus crocodilurus vietnamensis*. Amphibia-Reptilia.

Vicoso, B., Emerson, J. J., Zektser, Y., Mahajan, S., Bachtrog, D. (2013). Comparative Sex Chromosome Genomics in Snakes: Differentiation, Evolutionary Strata, and Lack of Global Dosage Compensation. PLoS Biology, 11(8), e1001643.

Webster TH, Vannan A, Pinto BJ, Denbrock G, Morales M, Dolby GA, Fiddes IT, DeNardo DF, Wilson MA. 2023. Complete dosage compensation without balance between the sexes in the ZZ/ZW Gila monster (*Heloderma suspectum*) revealed by *de novo* genome assembly. *In review*.

Weir BS, Cockerham CC. 1984. Estimating F-statistics for the analysis of population structure. *Evolution*. 38: 1358-1370.

Xie HX, Liang XX, Chen ZQ, Li WM, Mi CR, Li M, ... Du WG. 2022. Ancient demographics determine the effectiveness of genetic purging in endangered lizards. *Mol Biol Evol*. 39: msab359.

Zhang, Y., R. Ban, C. Wu, and C. Zhijian. 1996. Studies on Ultrastructure and Karyotypes of Crocodilian Lizard. China Guanxi Teachers University Press, pp.77.

Ziegler, T., Van Schingen, M., Rauhaus, A., Dang, P. H., Pham, D. T. K., Pham, C. T. & T. Q. Nguyen (2019): New insights into the habitat use and husbandry of crocodile lizards (Reptilia: Shinisauridae) including the conception of new facilities for Vietnamese crocodile lizards *Shinisaurus crocodilurus vietnamensis* in Vietnam and Germany. - International Zoo Yearbook 53: 250–269.