It's a Trap?! Escape from an ancient, ancestral sex chromosome system
and implication of *Foxl2* as the putative primary sex determining
gene in a lizard (Anguimorpha; Shinisauridae)

Brendan J. Pinto[1,2,3,*], Stuart V. Nielsen[4,5], Kathryn A. Sullivan[3,6], Ashmika Behere[6], Shannon E. Keating[6], Mona van Schingen-Khan[7], Truong Quang Nguyen[8,9], Thomas Ziegler[10,11], Jennifer Pramuk, Melissa A. Wilson[1,2,12], Tony Gamble[3,6,13,*]

1. School of Life Sciences, Arizona State University, Tempe, AZ USA

2. Center for Evolution and Medicine, Arizona State University, Tempe, AZ USA

3. Department of Zoology, Milwaukee Public Museum, Milwaukee, WI USA

4. Department of Biological Sciences, Museum of Life Sciences, Louisiana State University-Shreveport, Shreveport, LA USA

5. Florida Museum of Natural History, University of Florida, Gainesville, FL USA

6. Department of Biological Sciences, Marquette University, Milwaukee WI USA

7. Federal Agency for Nature Conservation, CITES Scientific Authority, Konstantinstraße 110, 53179 Bonn, Germany

8. Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet Road, Hanoi 10072, Vietnam

9. Graduate University of Science and Technology, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi 10072, Vietnam

10. Cologne Zoo, Riehler Straße 173, 50735 Cologne, Germany

11. Department of Biology, Institute of Zoology, University of Cologne, Zülpicher Straße 47b, 50674 Cologne, Germany

12. Center for Mechanisms of Evolution, Biodesign Institute, Tempe, AZ USA

13. Bell Museum of Natural History, University of Minnesota, St Paul, MN USA

*Correspondence:

brendanjohnpinto@gmail.com

tgamble@geckoevolution.org

Abstract

Although sex determination is ubiquitous in vertebrates, mechanisms of sex

determination vary from environmentally- to genetically-influenced. In vertebrates,

genetic sex determination is typically accomplished with sex chromosomes. Groups like

mammals maintain conserved sex chromosome systems, while sex chromosomes in

most vertebrate clades aren't conserved across similar evolutionary timescales. One

group inferred to have an evolutionarily stable mode of sex determination is

Anguimorpha, a clade of charismatic taxa including: monitor lizards, Gila monsters, and

crocodile lizards. The common ancestor of extant anguimorphs possessed a ZW

system that has been retained across the clade. However, the sex chromosome system

in the endangered, monotypic family of crocodile lizards (Shinisauridae) has remained

elusive. Here, we analyze genomic data to demonstrate that *Shinisaurus* has replaced

the ancestral anguimorph ZW system on LG7 chromosome with a novel ZW system on

LG3. The linkage group LG3 corresponds to chromosome 9 in chicken, and this is the

first documented use of this syntenic block as a sex chromosome in amniotes.

Additionally, this ~1Mb region harbors approximately 10 genes, including a duplication

of the sex-determining transcription factor, *Foxl2*—critical for the determination and

maintenance of sexual differentiation in vertebrates, and thus a putative primary sex

determining gene for *Shinisaurus*.

Introduction

The evolution of sex determination in vertebrates is impressive in its ability to

combine a highly conserved developmental network that can be initiated by quite

distinct molecular mechanisms in different species (Bachtrog et al. 2014; Graves, 2008).

In vertebrates, sex is commonly determined via either environmental and/or genetic

cues at critical points in development. In vertebrate groups that use genetic

mechanisms, the most common mechanism is sex chromosomes; either a male or

female heterogametic system where the male or female inherits the sex-limited (Y or W)

chromosome, respectively (Bachtrog et al. 2014; Gamble et al. 2015). Sex

chromosomes have been traditionally identified by comparing male and female

karyotypes under the light microscope. The presence of morphological differences

between the X and Y (or Z and W) chromosomes (i.e. heteromorphic sex

chromosomes) identify a species' sex chromosome system (Stevens, 1905; Bull, 1983).

However, many species possess sex chromosomes that cannot be identified via light

microscopy because the X and Y (or Z and W) are not morphologically distinguishable

from each other (i.e. homomorphic sex chromosomes). Other methods must be

employed, such as advanced cytogenetic techniques or high-throughput DNA

sequencing technologies, to identify sex chromosome systems in these taxa (Gamble

and Zarkower, 2014; Gamble et al. 2017; Pinto et al. 2022).

Squamate reptiles (lizards and snakes) demonstrate high variability in modes of

sex determination: where some clades have conserved, often heteromorphic, sex

chromosomes, while others display extraordinary lability in their modes of sex

determination and a high incidence of homomorphic sex chromosomes (Gamble et al.

2015, Kratochvíl et al. 2021; Augstenová et al. 2021a). One hypothesis of sex

chromosome evolution is that ancient, degenerated sex chromosome systems may act

as an "evolutionary trap", where the existence of highly differentiated (i.e.

heteromorphic) sex chromosomes preclude transitions to other sex-determining

systems (Bull 1983; Bull and Charnov, 1985; Pokorná and Kratochvíl, 2009). The

stability of old sex chromosome systems in mammals, birds, caenophidian snakes, and

others, provides anecdotal support for this hypothesis (Bull and Charnov, 1985;

Pokorná and Kratochvíl, 2009; Gamble et al. 2015). As more and more sex

chromosome transitions are identified, it remains unclear whether all ancient sex

chromosome systems are destined to become traps, but examples of taxa transitioning

away from ancient, degenerated sex chromosome systems are rare in amniotes (Acosta

et al. 2019; Nielsen et al. 2019; Rovatsos et al. 2019a). Previous phylogenetic studies

have supported the trap hypothesis in squamates (Pokorná and Kratochvíl, 2009;

Gamble et al. 2015), but also suffered from incomplete taxonomic sampling, which

might have biased the conclusions. In other words, testing this hypothesis is contingent

upon having sufficient data necessary to identify transitions away from an ancient sex

chromosome system, which typically requires (1) a reference genome to coordinate

linkage groups (which are rare in squamates; Pinto et al. 2023), (2) genome-scale data

from both sexes (e.g. Vicoso et al. 2013; Gamble et al. 2015; Pinto et al. 2022), and (3)

a robust phylogenetic hypothesis to establish relationships within the focal taxa (Nielsen

et al. 2019). Thus, the burden of proof is higher for identifying escapees from these

ancient sex chromosome systems, which may be responsible for the dearth of

examples and the previous lack of conclusive examinations of the evolutionary trap

hypothesis. Future identification of additional escapees will permit more conclusive

analyses of whether or not ancient sex chromosome systems truly act as evolutionary

traps across a broader phylogenetic scale.

The sex chromosomes of the infraorder Anguimorpha (lizards including monitor

lizards, Gila monsters, alligator lizards, and their allies) have long been a topic of

interest, likely resulting from the paucity of genetic and cytogenetic data for this group.

In recent years, advanced cytogenetic techniques (FISH) have facilitated karyotypic

analysis and identification of ZW sex chromosomes in the Gila monster (*Heloderma*

suspectum; Pokorná et al. 2014) and Komodo dragon (*Varanus komodoensis*; Pokorná

et al. 2016) leading to expanded interest in studying chromosome evolution in this

enigmatic group. More recently, RNAseq and qPCR analysis, in conjunction with draft

genomes of these same two anguimorph species (Gila monster; Webster et al. 2023,

and Komodo dragon; Lind et al. 2019), have provided some additional insights into this

system (Rovatsos et al. 2019b). Namely, the homology of the heteromorphic ZW

systems in the anguimorph genera *Abronia*, *Heloderma*, and *Varanus* (Rovatsos et al.

2019b; Webster et al. 2023). The presence of a ZW sex chromosome on the same

linkage group—syntenic with chromosome 28 in the chicken genome—in these three

genera, spanning the phylogenetic breadth of extant Anguimorpha, is strong evidence

that this is the ancestral sex chromosome system in the clade. Ancient sex

chromosome systems, like those ancestral to anguimorphs (115–180 million years old),

fit the criteria that should render them as an evolutionary trap (Pokorná and Kratochvíl,

2009; Rovatsos et al. 2019b). However, the sex chromosomes of many anguimorph

taxa remain unknown, including the monotypic family Shinisauridae, which is nested

within the anguimorph phylogeny (Figure 1).

The crocodile lizard (*Shinisaurus crocodilurus*) is the sole living member of the

family Shinisauridae and native to small disjunct regions of southeastern China and

northern Vietnam (Le and Ziegler, 2003; Huang et al. 2008; Nguyen et al. 2014). It is

one of the rarest lizard species in the world and is listed as Endangered in the IUCN

Red List (Nguyen et al. 2014). Due to anthropogenic pressures, populations have

experienced severe declines in recent years with less than 1000 individuals in the wild

in China and less than 100 adults in Vietnam (Huang et al. 2008; van Schingen et al.

2014a). They are semi-aquatic habitat specialists and depend upon clean streams in

broadleaf evergreen forest (Ning et al. 2006; van Schingen et al. 2016a) and their

restricted ecological niche is predicted to all but disappear due to climate change by the

end of this century (Li et al. 2013; van Schingen et al. 2014a; see also van Schingen-

Khan et al. 2022). Habitat destruction threatens remaining populations, as well as

overcollection for food and the international pet trade (Huang et al. 2008; van Schingen

et al. 2014b; van Schingen et al. 2016a). Although still recognized as a single species,

there exist multiple conservation units, with *S. crocodilurus vietnamensis* from Vietnam

and the nominal subspecies from China consisting of several distinct lineages (van

Schingen et al. 2016b; Ngo et al. 2020; Nguyen et al. 2022). Crocodile lizards do not

have a clear sexual dimorphism. While morphological traits, such as coloration or body

morphometry, may provide some indication of the sex, it remains difficult for most

people to identify the sex of individuals (van Schingen et al. 2016b). Relevant to the

present study, examination of male and female *S. crocodilurus* karyotypes have

revealed no heteromorphic sex chromosomes (Zhang et al. 1996; Augstenová et al.

2021b). To identify sex chromosomes in this species, we analyzed whole-genome re-

sequencing data for approximately 50 sexed, individual crocodile lizards (Xie et al.

2022) using whole-genome re-sequencing to show that the sex determining system in

S. crocodilurus is a novel ZW system that has eluded previous analyses, at least in part,

due to the small size (<1Mb) of its sex determining region (SDR).

Methods

WGS analysis

We downloaded low-coverage whole genome Illumina resequencing (WGS) reads from NCBI SRA for multiple male and female individuals (see *Data Availability* for accessions). We constructed a Snakemake [v6.10.0] (Mölder et al. 2021) workflow in an isolated conda environment [v4.11.0] (https://docs.anaconda.com/) containing relevant packages: BBmap [v38.93] (Bushnell, 2014), FastQC [v0.11.9] (Andrews, 2010), Freebayes [v1.3.5] (Garrison and Marth, 2012), GFF utilities [v0.10.1] (Pertea and Pertea, 2020), Minimap2 [v2.22] (Li, 2018), Mosdepth [v0.3.2] (Pedersen and Quinlan, 2018), MultiQC [v1.11] (Ewels et al. 2016), Parallel [v20211022] (Tange, 2018), pixy [v1.2.5.beta1] (Korunes and Samuk, 2021), RTGTools [v3.12.1] (Cleary et al. 2015), Sambamba [v0.8.1] (Tarasov et al. 2015), Samtools [v1.12] (Li et al. 2009), seqkit [v0.11.0] (Shen et al. 2016), STACKS [v2.6.0] (Catchen et al. 2013), and Trim Galore! [v0.6.7] (https://doi.org/10.5281/zenodo.5127899). To process the raw sequencing data, we trimmed adapters and low-quality regions using Trim Galore!, then removed PCR duplicates using BBmap. Quality assessment using FastQC and MultiQC was conducted at each step, and we subsequently removed samples with fewer than 5 million PE reads after filtering PCR duplicates. The final WGS dataset possessed 50 sexed samples (27 male and 23 female individuals) sourced from China and Vietnam. We proceeded to map reads for each individual to the female reference genome (Xie et al. 2022) with minimap2 and calculated read depth and read mapping statistics using mosdepth and samtools, respectively. Then, we generated an all-sites VCF file with freebayes-parallel. Lastly, we calculated Weir and Cockerham (1984) F_{ST} between males and females and nucleotide diversity statistics using pixy at 500kb resolution and, for LG3 only, also at 100kb resolution.

Validation of the putative ZW system in Shinisaurus crocodilurus

Male vs. female F_{ST} values are agnostic to which sex is heterogametic (i.e. XY vs. ZW). Therefore, we generated a dataset of '*in silico* poolseq' reads by subsampling each WGS sample to 10 million paired reads (20 million total reads per sample) using seqkit and combined into male and female pools. We analyzed the pools using Pooled

Sequencing Analyses for Sex Signal [PSASS; v3.1.0]

(https://doi.org/10.5281/zenodo.3702337). We then generated PCR primers targeting

the annotated version of *Foxl2*'s second exon [FOXL2-ex2-F2 5' –

CAGAGCTCGTCCCATTCACTT – 3' and FOXL2-ex2-R2 5' –

GAGAGATGTACCACCGGGAG – 3'] and sequenced the resultant amplicon using

Sanger sequencing (Psomagen). Individuals used in Sanger sequencing are detailed in

Supplemental Table 1.

Genome Annotation

160   We used previously lifted over annotations (Pinto et al. 2023;

https://doi.org/10.6084/m9.figshare.20201099.v1) via Liftoff [v1.6.3] (Shumate and

Salzberg, 2021) from the draft genome of a male *S. crocodilurus* (Gao et al. 2017) to

the new, unannotated female reference genome (Xie et al. 2022; GCA_021292165.1).

We pulled coding transcripts from the genome using GFF Utilities. We used the 10

genes within the putative ~1Mb SDR on LG3 to perform a high-stringency tBLASTx

query (Altschul et al. 1990) to the chicken genome on Ensembl (Howe et al. 2020) with

a word size of 3, maximum of 10 hits, e-value cutoff of $1e^{-50}$, using BLOSUM62 scoring

matrix. These queries received hits on 7 of the 10 total genes (Table 1).

Results

Across WGS experiments, read mapping efficiency ranged from 80.60% (for

SRR5019740) to 99.40% (for SRR14583318). After variant calling, the WGS dataset

contained 6,202,005 biallelic variants (see Data Availability section for additional VCF

statistics). We identified a region of high F_{ST} between males and females on linkage

group 3 (LG3; Figure 2), however, comparing M/F F_{ST} values does not necessarily

diagnose which sex is heterogametic (i.e. XY vs. ZW). Therefore, we composed a

dataset of '*in silico* poolseq' reads to identify an excess of female-associated SNPs

aligning to the previously identified region of high M/F F_{ST} (Supplemental Figure 1).

Taken together, these data suggest that *S. crocodilurus* possesses a female

heterogametic system (ZW) with an SDR located in a ~900kb region on LG3.

Upon further investigation of the SDR, we identified a total of 10 genes annotated

within this region of high F_{ST} and an excess of female-specific SNPs. To better

characterize these genes, we BLAST-ed each to the chicken genome. We recovered

high-quality BLAST hits for seven of the 10 annotated *S. crocodilurus* SDR genes in

chicken (Table 1). Six out of the seven queries hit genes located on chicken

chromosome 9, while the other landed on a chicken chromosome 30 (Table 1). In our

poolseq analysis, one of these genes possessed half the read depth in females relative

to males (Supplemental Table 2) and, upon closer inspection, we identified a duplicated,

unannotated copy of that gene Forkhead Box L2 (*Foxl2*), located approximately 70kb

upstream—with 99% sequence identity, also located within the putative SDR. We

included this *Foxl2* copy in a BLAST search against chicken, where it was again

identified as a *Foxl2* homolog (Table 1). We also BLAST-ed *Foxl2* to the earlier male *S.*

crocodilurus draft genome (Gao et al. 2017) and found only a single copy of *Foxl2* in

this genome matching one copy in the updated reference genome with 100% sequence

identity, consistent with both (1) the duplicated version being W-specific and (2) the

female reference being chimeric for Z and W alleles (Xie et al. 2022). Lastly, we

generated a gene tree using *Foxl2* copies from across reptiles to confirm its duplicated

origination was within Shinisauridae (Supplemental Figure 2). Thus, in the chimeric

female reference genome, this putative W-linked *Foxl2* copy was located approximately

70kb upstream of the annotated Z-linked copy of *Foxl2* on the other side of an assembly

gap.

The WGS data used in the *in silico* PoolSeq analysis were restricted to only

individuals from Chinese populations to reduce the influence of population-specific

demographic processes (Xie et al. 2022). To include the less-numerous Vietnam

samples, we generated PCR primers for a segment of *Foxl2*'s second exon and Sanger

sequenced multiple females (Vietnam) and males (China and Vietnam) (Supplemental

Figure 3). We identified one SNP in the female Vietnamese samples in this region and

tested its association with sex using Fisher's exact test (p-value = 0.0003***). Thus, the

ZW SDR containing *Foxl2* appears to be conserved between populations of *S.*

crocodilurus from both China and Vietnam.

Discussion

Escaping the "Evolutionary Trap"

An open question within sex chromosome evolution is whether ancient, degenerated sex chromosomes act as evolutionary traps (Pokorná and Kratochvíl, 2009; Nielsen et al. 2019; Pinto et al. 2023). The most recent common ancestor of extant anguimorphs is thought to have possessed a ZW system on the linkage group syntenic with chicken chromosome 28, which is located on the distal region of LG7 in in the *S. crocodilurus* reference genome (Rovatsos et al. 2019b; Webster et al. 2023). The sex determining region (SDR) in *S. crocodilurus* is located on LG3, a region syntenic with chicken chromosome 9. Of note, however, at present it is difficult to assess the precise genomic coordinates and gene content of the SDR due to the chimeric nature of the reference genome assembly. To our knowledge, this is the first demonstration in a tetrapod of the syntenic region of chicken chromosome 9 being recruited in a sex determining role (Kratochvíl et al. 2021), lending further support to the idea that all chromosomes will likely be recruited into a sex determining role given thorough enough phylogenetic sampling (Graves and Peichel, 2010; Hodgkin, 2002; O'Meally et al. 2012; Pinto et al. 2022).

It is clear from these genomic data that *S. crocodilurus* possesses a distinct sex chromosome system from all other known anguimorphs. Unlike the case of Corytophanidae and other pleurodonts, where phylogenetic relationships among taxa were inconclusive (Nielsen et al. 2019), the relationship of *S. crocodilurus* to all other anguimorphs is far less divisive. Indeed, *S. crocodilurus* is well-supported as nested within Anguimorpha—either sister to Varanidae as a member of the "Paleoanguimorpha" (Burbrink et al. 2020) or as sister to a clade containing Varanidae and Lanthanotidae (Singhal et al. 2021), depending on taxonomic sampling. Thus, assuming the hypothesis that an ancient origin of the ZW sex chromosome system possessed by extant *Varanus*, *Heloderma*, and *Abronia* is correct, then *S. crocodilurus* has successfully escaped the evolutionary trap of their ancestral, degenerated sex chromosome system—a system nearly as ancient as those systems found in both mammals and birds (Rovatsos et al. 2019b; Webster et al. 2023). It is worth noting that

there remains another putative escape from the ancestral anguimorph sex chromosome

system in *Anguis* that has yet to be explored further (Rovatsos et al. 2019b) and more

recent phylogenetic work has implicated that Corytophanidae is likely nested

somewhere within other pleurodonts, rather than being sister to all other species

(Burbrink et al. 2020; Singhal et al. 2021). This suggests that there are a minimum of

two evolutionary escapes within Toxicofera (snakes, iguanians, and anguimorphs)—and

perhaps even two within the infraorder Anguimorpha alone.

Primary Sex Determination in Shinisauridae

In many vertebrate groups where the primary sex determiner (PSD) is known, a

relatively short list of commonly-recruited PSDs have been identified (i.e. the 'usual

suspects'; Adolfi et al. 2021; Dor et al. 2019; Herpin and Schartl, 2015). Indeed, the

same genes, or their paralogs, have been independently co-opted to function as the

PSD in many taxa, examples including *Sox3* in placental mammals and some medaka

(members of the *Oryzias celebensis* and *O. javanicus* groups); *Amh* in tilapia, northern

pike, and potentially other anguimorphs (Li et al. 2015; Myosho et al. 2015; Pan et al.

2019; Rovatsos et al. 2019b; Webster et al. 2023; and see Pan et al. 2021 for recent

review); and *Dmrt1* in birds, a frog (*Xenopus laevis*), tongue sole, and other medaka fish

(members of the *Oryzias latipes* group) (Chen et al. 2014; Ioannidis et al. 2021;

Matsuda et al. 2002; Nanda et al. 2002; Smith et al. 2009). This is the first time

Forkhead Box L2 (*Foxl2*) has been implicated as a PSD in a vertebrate, although it has

been predicted to be one (e.g. Ma et al. 2022).

The transcription factor, *Foxl2*, is a direct transcriptional activator of aromatase,

involved in development of the ovaries and its loss in mice during embryogenesis leads

to abnormal ovarian development and infertility (Fleming et al. 2010; Pannetier et al.

2006; Schmidt et al. 2004; Uda et al. 2004). After primary sex determination and sexual

development have concluded, *Dmrt1* and *Foxl2* antagonize each other transcriptionally

in gonadal tissue, where sustained *Dmrt1* and *Foxl2* expression is required for adult

maintenance of testis and ovary tissue, respectively (Garcia-Ortiz et al. 2009; Matson et

al. 2011; Uhlenhaut et al. 2009). Indeed, *Foxl2* also behaves in a dose-dependent

manner in some turtle species where its overexpression at the embryonic stage can

induce male-to-female sex reversal in ZZ soft-shelled turtles (*Pelodiscus sinensis*) and

female differentiation in male-temperature-incubated red-eared sliders (*Trachemys*

scripta) (Jin et al. 2022; Ma et al. 2022). Importantly, *Dmrt1* has been recruited to act as

a primary sex determining gene in multiple taxa (Matson and Zarkower, 2012), while

Foxl2 has remained mysteriously absent from this list—with the singular putative

exception being recently described in some species of bivalve mollusks (Han et al.

2022). Thus, the identification of both *Foxl2* and a duplicated *Foxl2* copy in the W-

limited region of the *Shinisaurus* genome supports the expanded list of the "usual

suspects" that might act as the PSD in vertebrates.

Pragmatically, the identification of a novel ZW system in *S. crocodilurus* may

present an important juncture in the conservation efforts of this endangered lizard

species, that are urgently needed (Nguyen et al. 2014). Body morphometrics in mature

specimens may provide an indication of the sex, i.e. males tend to have a relatively

larger head, relative to abdomen length than females (van Schingen et al. 2016b).

However, definitive sexually dimorphic characters are lacking in the species, especially

in hatchlings, juveniles, and subadults. Therefore, a molecular genetic sex test could

assist in well-managed captive breeding efforts in this species (Ziegler et al. 2019). This

is vital as it's estimated only ~1,000 individuals remain in the wild populations in China

and Vietnam during the last census (Huang et al. 2008; van Schingen et al. 2016a),

while loss of remaining habitats and poaching are considered ongoing. This information

may play a vital role in conservation efforts of this species and should be incorporated

into ongoing captive breeding work (Ziegler et al. 2019).

In conclusion, using a combination of sequencing and validation techniques we

identified the elusive ZW system in the endangered crocodile lizard, *Shinisaurus*

crocodilurus. This ZW system is located on LG3 and, although interpretation inherits

strong reference bias (a chimeric ZW reference genome), the SDR appears to be <1Mb

in size and contains approximately 10 genes. One of these genes, *Foxl2*, possesses a

duplicated copy and is important in ovarian development and fertility in vertebrates.

Because of its sequence conservation (either strictly age-related or via gene

conversion) and possibly its proximity to the original Z copy of *Foxl2*, we hypothesize

that if *Foxl2* is the PSD in this system, it may be a gene dosage-dependent mechanism,

where ZW females possess three copies of *Foxl2* instead of the two copies of ZZ males.

This specific hypothesis assumes that the Z copy of *Foxl2* is retained in the

pseudoautosomal region of the W chromosome, however, phased Z and W sequences

are needed to provide additional support to this model. The hypothetical mechanism

would essentially be the inverse of the dose-dependent *Dmrt1* sex determination in

birds, where a lack of *Dmrt1* on the W decreases *DMRT1* expression in females,

allowing for *Foxl2* to proceed with ovarian development (Ioannidis et al. 2021; Smith et

al. 2009). Here, extra gene copies of *Foxl2* increase *FOXL2* expression to downregulate

Dmrt1 expression and initiate ovarian development in the developing gonad. Thus, we

provide a putative sex determining gene for the crocodile lizard (*Shinisaurus*

crocodilurus) and speculate as to its potential mechanism of action in this system.

Data Availability:

The WGS data used in this study is available on NCBI, SRA accessions for WGS

data are: SRR14583317, SRR14583321, SRR14583324-26, SRR14583330,

SRR14583333, SRR14583340-49, SRR14583351, SRR14583353-54, SRR14583356,

SRR14583360-66, SRR5019733-45, SRR14583318-20, SRR14583322-23,

SRR14583331, SRR14583334-39, SRR14583346, SRR14583350, SRR14583352,

SRR14583355, SRR14583357-59. Sequence data generated in this study are available

on SRA under BioProject PRJNA975696, detailed in Supplemental Table 1, and code,

including and VCF statistics and gene alignments, are available on GitHub:

https://github.com/DrPintoThe2nd/Shinisaurus_ZW.

Acknowledgements:

The authors would like to acknowledge Research Computing at Arizona State

University for providing high-performance computing and storage resources that have

contributed to the research results reported within this paper

(http://www.researchcomputing.asu.edu). We thank Anna Rauhaus (Cologne Zoo) for

her help with the application and preparation of tissue sending and the Woodland Park

Zoo for their respective assistance. Many thanks CITES Management Authority of

Vietnam for issuing permits (CITES permits No. 13VN1246N/CT-KL and

16VN0920N/CT-KL).This work was funded by the Morris Animal Foundation (Study

grant D19ZO-021) for their generous funding of this project (T.G.) and also supported

by the National Institute of General Medical Sciences (NIGMS) of the National Institutes

of Health grant R35GM124827 (M.A.W.).

References

References
Adolfi, M. C., Herpin, A., & Schartl, M. (2021). The replaceable master of sex
determination: bottom-up hypothesis revisited. *Philosophical Transactions of the Royal*
Society B: Biological Sciences. 376(1832):20200090.
Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local
alignment search tool. *J Mol Biol*. 215: 403-410.
Andrews S. 2010. FastQC: A quality control tool for high throughput sequence
data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.
Augstenová, B., Pensabene, E., Veselý, M., Kratochvíl, L., Rovatsos, M. 2021a.
Are geckos special in sex determination? Independently evolved differentiated ZZ/ZW
sex chromosomes in carphodactylid geckos. *Genome Biology and Evolution*, 13(7),
evab119.
Augstenová, B., Pensabene, E., Kratochvíl, L., Rovatsos, M. 2021b. Cytogenetic
evidence for sex chromosomes and karyotype evolution in anguimorphan lizards. *Cells*,
10(7), 1612.
Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman TL, Hahn
MW, Kitano J, Mayrose I, Ming R, Perrin N, Ross L, Valenzuela N, Vamosi JC. 2014.
Sex determination: Why so many ways of doing it? *PLoS Biol*. 12: e1001899.
Bull, JJ. 1983. Evolution of Sex Determining Mechanisms. Benjamin Cummings:
Menlo Park, CA.
Bull J, Charnov E. 1985. On irreversible evolution. *Evolution* 39, 1149–1155.
Burbrink, F. T., Grazziotin, F. G., Pyron, R. A., Cundall, D., Donnellan, S., Irish,
F., Keogh, J. S., Kraus, F., Murphy, R. W., Noonan, B., Raxworthy, C. J., Ruane, S.,
Lemmon, A. R., Lemmon, E. M., Zaher, H. (2019). Interrogating Genomic-Scale Data
for Squamata (Lizards, Snakes, and Amphisbaenians) Shows no Support for Key
Traditional Morphological Relationships. Systematic Biology, 69(3), 502–520.
Bushnell B. 2014. BBMap: a fast, accurate, splice-aware aligner (No. LBNL-
7065E). Lawrence Berkeley National Lab (LBNL), Berkeley, CA (United States).
Chen, S., Zhang, G., Shao, C., Huang, Q., Liu, G., Zhang, P., Song, W., An, N.,
Chalopin, D., Volff, J.-N., Hong, Y., Li, Q., Sha, Z., Zhou, H., Xie, M., Yu, Q., Liu, Y.,
Xiang, H., Wang, N., … Wang, J. (2014). Whole-genome sequence of a flatfish provides
insights into ZW sex chromosome evolution and adaptation to a benthic lifestyle. *Nature*
Genetics. 46(3):253–260.
Cleary JG, Braithwaite R, Gaastra K, Hilbush BS, Inglis S, Irvine SA, Francisco
M. (2015). Comparing variant call files for performance benchmarking of next-
generation sequencing variant calling pipelines. *BioRxiv*.
Dor L, Shirak A, Kohn YY, Gur T, Weller JI, Zilberg D, Seroussi E, Ron M. 2019.
Mapping of the sex determining region on linkage group 12 of guppy (*Poecilia*
reticulata). *G3*. 9: 3867-3875.
Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics*. 32: 3047-3048.
Fleming, N. I., Knower, K. C., Lazarus, K. A., Fuller, P. J., Simpson, E. R., &
Clyne, C. D. (2010). Aromatase Is a Direct Target of *FOXL2*: C134W in Granulosa Cell
Tumors via a Single Highly Conserved Binding Site in the Ovarian Specific Promoter.
PLoS ONE. 5(12):e14389.

Gamble T, Zarkower D. 2014. Identification of sex-specific molecular markers
using restriction site-associated DNA sequencing. *Molecular Ecology Resources.*
14:902–913.
Gamble T, Coryell J, Ezaz T, Lynch J, Scantlebury D, Zarkower D. 2015.
Restriction site-associated DNA sequencing (RAD-seq) reveals an extraordinary
number of transitions among gecko sex-determining systems. *Mol Biol Evol.* 32: 1296-
1309.
Gamble T, Castoe TA, Nielsen SV, Banks JL, Card DC, Schield DR, Schuett
GW, Booth W. 2017. The discovery of XY sex chromosomes in a *Boa* and *Python*. *Curr*
Biol. 27: 2148-2153.
Gao J, Li Q, Wang Z, Zhou Y, Martelli P, Li F, ... Zhang G. 2017. Sequencing, de
novo assembling, and annotating the genome of the endangered Chinese crocodile
lizard *Shinisaurus crocodilurus*. *GigaScience.* 6: gix041.
Garcia-Ortiz JE, Pelosi E, Omari S, Nedorezov T, Piao Y, Karmazin J, ...
Ottolenghi C. 2009. *Foxl2* functions in sex determination and histogenesis throughout
mouse ovary development. *BMC Dev Biol.* 9: 36.
Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read
sequencing. *arXiv*. 1207.3907. https://arxiv.org/abs/1207.3907
Graves J. 2008. Weird animal genomes and the evolution of vertebrate sex and
sex chromosomes. *Ann Rev Genet.* 42: 565-586.
Graves J, Peichel C. 2010. Are homologies in vertebrate sex determination due
to shared ancestry or to limited options? *Genome Biol.* 11: 205. doi:10.1186/gb-2010-
11-4-205
Han, W., Liu, L., Wang, J., Wei, H., Li, Y., Zhang, L., ... Wang, S. (2022). Ancient
homomorphy of molluscan sex chromosomes sustained by reversible sex-biased genes
and sex determiner translocation. *Nat Ecol Evol.* 1-16.
Herpin A, Schartl M. 2015. Plasticity of gene- regulatory networks controlling sex
determination: of masters, slaves, usual suspects, newcomers, and usurpators. *EMBO*
Rep. 16:1260-1274.
Hodgkin J. 2002. Exploring the envelope: systematic alteration in the sex-
determination system of the nematode *Caenorhabditis elegans*. *Genetics.* 162: 767-
780.
Howe KL, Achuthan P, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean
IM, Azov AG, Bennett R, Bhai J, Billis K, Boddu S, Charkhchi M, Cummins C, Da Rin
Fioretto L, Davidson C, Dodiya K, El Houdaigui B, Fatima R, … Flicek P. 2020. Ensembl
2021. *Nucleic Acids Res.* 49: 884-891. doi:10.1093/nar/gkaa942
Huang C.M., Yu H., Wu Z.J., Li Y.B., Wei F.W., Gong M.H. (2008). Population
and conservation strategies for the Chinese crocodile lizard (*S. crocodilurus*) in China.
Animal Biodiversity and Conservation. 31(2):63-70.
Ioannidis, J., Taylor, G., Zhao, D., Liu, L., Idoko-Akoh, A., Gong, D., Lovell-
Badge, R., Guioli, S., McGrew, M. J., & Clinton, M. (2021). Primary sex determination in
birds depends on DMRT1 dosage, but gonadal sex does not determine adult secondary
sex characteristics. *PNAS.* 118(10).
Jin, L., Sun, W., Bao, H., Liang, X., Li, P., Shi, S., Wang, Z., Qian, G., & Ge, C.
(2022). The forkhead factor *Foxl2* participates in the ovarian differentiation of Chinese
soft-shelled turtle *Pelodiscus sinensis*. *Developmental Biology*. 492:101–110.

Korunes KL, Samuk K. 2021. pixy: Unbiased estimation of nucleotide diversity
and divergence in the presence of missing data. *Mol Ecol Res.* 21: 1359-1368.
Kratochvíl, L., T. Gamble, and M. Rovatsos. 2021. Sex chromosome evolution
among amniotes: Is the origin of sex chromosomes non-random? Philosophical
Transactions of the Royal Society B: Biological Sciences 376:20200108.
Kumar S, Stecher G, Suleski M, Hedges SB. (2017). TimeTree: a resource for
timelines, timetrees, and divergence times. *Mol Biol Evol.* 34:1812–1819
Le K.Q., Ziegler T. (2003). First record of the Chinese crocodile lizard from
outside of China: report on a population of *S. crocodilurus* Ahl, 1930 from North-Eastern
Vietnam. *Hamadryad.* 27(2):193-199.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, ... Durbin R. (2009).
The sequence alignment/map format and SAMtools. *Bioinformatics.* 25:2078-2079.
Li H. (2018). Minimap2: pairwise alignment for nucleotide sequences.
Bioinformatics. 34:3094-3100.
Li, M., Sun, Y., Zhao, J., Shi, H., Zeng, S., Ye, K., Jiang, D., Zhou, L., Sun, L.,
Tao, W., Nagahama, Y., Kocher, T. D., & Wang, D. (2015). A Tandem Duplicate of Anti-
Müllerian Hormone with a Missense SNP on the Y Chromosome Is Essential for Male
Sex Determination in Nile Tilapia, *Oreochromis niloticus. PLoS Genetics.*
11(11):e1005678.
Li, X., Tian, H., Wang, Y., Li, R., Song, Z., Zhang, F., ... & Li, D. (2013).
Vulnerability of 208 endemic or endangered species in China to the effects of climate
change. *Regional Environmental Change*, *13*, 843-852.
Ma, X., Liu, F., Chen, Q., Sun, W., Shen, J., Wu, K., Zheng, Z., Huang, J., Chen,
J., Qian, G., & Ge, C. (2022). *Foxl2* is required for the initiation of the female pathway in
a temperature-dependent sex determination system in *Trachemys scripta.*
Development. 149(13).
Matson, C. K., Murphy, M. W., Sarver, A. L., Griswold, M. D., Bardwell, V. J., &
Zarkower, D. (2011). *DMRT1* prevents female reprogramming in the postnatal
mammalian testis. *Nature.* 476(7358), 101–104.
Matson, C. K., Zarkower, D. (2012). Sex and the singular DM domain: insights
into sexual regulation, evolution and plasticity. *Nature Reviews Genetics.* 13(3):163-
174.
Matsuda, M., Nagahama, Y., Shinomiya, A., Sato, T., Matsuda, C., Kobayashi,
T., Morrey, C. E., Shibata, N., Asakawa, S., Shimizu, N., Hori, H., Hamaguchi, S., &
Sakaizumi, M. (2002). DMY is a Y-specific DM-domain gene required for male
development in the medaka fish. *Nature.* 417(6888):559–563.
Mölder F, Jablonski KP, Letcher B, Hall MB, Tomkins-Tinch CH, Sochat V,
Forster J, Lee S, Twardziok SO, Kanitz A, Wilm A, Holtgrewe M, Rahmann S, Nahnsen
S, Köster J. 2021. Sustainable data analysis with Snakemake. *F1000Res.* 10:33.
Nanda, I., Kondo, M., Hornung, U., Asakawa, S., Winkler, C., Shimizu, A., Shan,
Z., Haaf, T., Shimizu, N., Shima, A., Schmid, M., & Schartl, M. (2002). A duplicated
copy of *DMRT1* in the sex-determining region of the Y chromosome of the medaka,
Oryzias latipes. PNAS. 99(18):11778–11783.
Ngo, H. T., Nguyen, T. T., Le, M. D., van Schingen-Khan, M., Nguyen, T. Q.,
Rauhaus, R., Vences, M. & T. Ziegler (2020): Genetic screening of captive crocodile
lizards (*Shinisaurus crocodilurus*) in Europe. – Der Zoologische Garten 88: 17-30.

Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2014). IQ-TREE:
A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood
Phylogenies. *Molecular Biology and Evolution*. 32(1), 268–274.
Nguyen TQ, Hamilton P, Ziegler T. (2014). *Shinisaurus crocodilurus. The IUCN*
Red List of Threatened Species. doi:10.2305/IUCN.UK.2014-
1.RLTS.T57287221A57287235.en. Accessed on 04 March 2022.
Nguyen, T. T., Ngo, H. T., Ha, Q. Q., Nguyen, T. Q., Le, T. Q., Nguyen, S. H.,
Pham, C. T., Ziegler, T., van Schingen-Khan, M. & M. D. Le (2022): Molecular
phylogenetic analyses and ecological niche modeling provide new insights into threats
to the endangered Crocodile Lizard (*Shinisaurus crocodilurus*). - Frontiers of
Biogeography 2022, 14.1, e54779
Nielsen SV, Guzmán-Mendez IA, Gamble T, Blumer M, Pinto BJ, Kratochvíl L,
Rovatsos M. 2019. Escaping the evolutionary trap? Sex chromosome turnover in
basilisks and related lizards (Corytophanidae: Squamata). *Biol Lett*. 15: 20190498.
Ning J., Huang C., Yu H., Dai D., Wu Z., Zhong Y. (2006). Summer habitat
characteristics of the Chinese Crocodile Lizard (*Shinisaurus crocodilurus*) in the
Loukeng Nature Reserve, Guangdong. *Zoological Research*. 27:419-426.
O'Meally D, Ezaz T, Georges A, Sarre SD, Graves JAM. 2012. Are some
chromosomes particularly good at sex? Insights from amniotes. *Chromosome Res*. 20:
7-19.
Pan Q, Feron R, Yano A, Guyomard R, Jouanno E, Vigouroux E, Wen M, Busnel
JM, Bobe J, Concordet JP, … Guiguen Y. 2019. Identification of the master sex
determining gene in Northern pike (*Esox lucius*) reveals restricted sex chromosome
differentiation. *PLoS Genetics*. 15: e1008013. doi:10.1371/journal.pgen.1008013
Pan Q, Kay T, Depincé A, Adolfi M, Schartl M, Guiguen Y, Herpin A. 2021.
Evolution of master sex determiners: TGF-β signalling pathways at regulatory
crossroads. *Phil Trans Roy Soc B*. 376:20200091.
Pannetier, M., Fabre, S., Batista, F., Kocer, A., Renault, L., Jolivet, G., Mandon-
Pépin, B., Cotinot, C., Veitia, R., & Pailhoux, E. (2006). FOXL2 activates P450
aromatase gene transcription: towards a better characterization of the early steps of
mammalian ovarian development. *Journal of Molecular Endocrinology*. 36(3):399–413.
Pedersen BS, Quinlan AR. 2018. Mosdepth: Quick coverage calculation for
genomes and exomes. *Bioinformatics*. 34: 867-868.
Pertea G, Pertea M. 2020. GFF utilities: GffRead and GffCompare.
F1000Research. 9.
Pinto BJ, Keating SE, Nielsen SV, Scantlebury DP, Daza JD, Gamble T. (2022).
Chromosome-level genome assembly reveals dynamic sex chromosomes in
Neotropical leaf-litter geckos (Sphaerodactylidae: *Sphaerodactylus*). *Journal of*
Heredity. 113(3): 272-287.
Pinto, B. J., Gamble, T., Smith, C. H., Wilson, M. A. (2023). A lizard is never late:
squamate genomics as a recent catalyst for understanding sex chromosome and
microchromosome evolution. *Journal of Heredity. In press.*
Pokorná, M., Kratochvíl, L. (2009). Phylogeny of sex-determining mechanisms in
squamate reptiles: are sex chromosomes an evolutionary trap? *Zoological Journal of*
the Linnean Society, 156(1), 168–183.

Rovatsos, M., Farkačová, K., Altmanová, M., Johnson Pokorná, M., Kratochvíl, L.
(2019a). The rise and fall of differentiated sex chromosomes in geckos. *Molecular*
Ecology, 28(12), 3042–3052.
Rovatsos M, Rehák I, Velenský P, Kratochvíl L. 2019b. Shared Ancient Sex
Chromosomes in Varanids, Beaded Lizards, and Alligator Lizards. *Mol Biol Evol*. 36:
1113-20.
Schmidt, D., Ovitt, C. E., Anlag, K., Fehsenfeld, S., Gredsted, L., Treier, A.-C., &
Treier, M. (2004). The murine winged-helix transcription factor *Foxl2* is required for
granulosa cell differentiation and ovary maintenance. *Development*. 131(4):933–942.
Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A cross-platform and ultrafast toolkit for
fasta/q file manipulation. *PLoS One*. 11: e0163962. doi:10.1371/journal.pone.0163962
Shumate A, Salzberg SL. 2021. Liftoff: Accurate mapping of gene annotations.
Bioinformatics. 12: 1639-1643.
Singhal, S., Colston, T. J., Grundler, M. R., Smith, S. A., Costa, G. C., Colli, G.
R., ... Rabosky, D. L. (2021). Congruence and conflict in the higher-level phylogenetics
of squamate reptiles: an expanded phylogenomic perspective. *Systematic Biology*,
70(3), 542-557.
Smith, C. A., Roeszler, K. N., Ohnesorg, T., Cummins, D. M., Farlie, P. G.,
Doran, T. J., & Sinclair, A. H. (2009). The avian Z-linked gene DMRT1 is required for
male sex determination in the chicken. *Nature*. *461*(7261):267-271.
Stevens NM. 1905. A study of the germ cells of *Aphis rosae* and *Aphis*
oenotherae. *J Exp Zool*. 2: 313-333.
Tange O. 2018. Gnu parallel 2018. pp. 112. doi:10.5281/zenodo.1146014
Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. 2015. Sambamba: fast
processing of NGS alignment formats. *Bioinformatics*. 31: 2032-2034.
Uda, M., Ottolenghi, C., Crisponi, L., Garcia, J. E., Deiana, M., Kimber, W.,
Forabosco, A., Cao, A., Schlessinger, D., & Pilia, G. (2004). *Foxl2* disruption causes
mouse ovarian failure by pervasive blockage of follicle development. *Human Molecular*
Genetics. 13(11):1171–1181.
Uhlenhaut NH, Jakob S, Anlag K, Eisenberger T, Sekido R, Kress J, Treier AC,
Klugmann C, Klasen C, Holter NI, Riethmacher D, Schütz G, Cooney AJ, Lovell-Badge
R, Treier M. 2009. Somatic sex reprogramming of adult ovaries to testes by *FOXL2*
ablation. *Cell*. 139: 1130-1142.
van Schingen M., Ihlow F., Nguyen T.Q., Ziegler T., Bonkowski M., Wu Z.,
Rödder D. (2014a): Potential distribution and effectiveness of the protected area
network for the crocodile lizard, *Shinisaurus crocodilurus* (Reptilia: Squamata: Sauria).
Salamandra. 50(2):71-76.
van Schingen M., Pham C.T., Thi H.A., Bernardes M., Hecht V., Nguyen T.Q.,
Bonkowski M., Ziegler T. (2014b). Current status of the crocodile lizard *Shinisaurus*
crocodilurus Ahl, 1930 in Vietnam with implications for conservation measures. *Revue*
suisse de Zoologie. 121(3):1-15.
van Schingen M., Ha Q.Q., Pham C.T., Le T.Q., Nguyen T.Q., Bonkowski M.,
Ziegler T. (2016a). Discovery of a new crocodile lizard population in Vietnam:
Population trends, future prognoses and identification of key habitats for conservation.
Revue suisse de Zoologie. 123(2):241-251.

van Schingen, M., Duc Le, M., Thi Ngo, H., The Pham, C., Quy Ha, Q., Quang
Nguyen, T., Ziegler, T. (2016b). Is there more than one Crocodile Lizard? An Integrative
Taxonomic Approach Reveals Vietnamese and Chinese Shinisaurus crocodilurus
Represent Separate Conservation and Taxonomic Units. *Der Zoologische Garten*.
85(5):240-260.
van Schingen-Khan, M., Barthel, L. M. F., Pham, D. T. K., Pham, C. T., Nguyen,
T. Q., Ziegler, T. & M. Bonkowski (2022): Will climatic changes affect the Vietnamese
crocodile lizard? Seasonal variation in microclimate and activity pattern of *Shinisaurus*
crocodilurus vietnamensis. Amphibia-Reptilia.
Vicoso, B., Emerson, J. J., Zektser, Y., Mahajan, S., Bachtrog, D. (2013).
Comparative Sex Chromosome Genomics in Snakes: Differentiation, Evolutionary
Strata, and Lack of Global Dosage Compensation. PLoS Biology, 11(8), e1001643.
Webster TH, Vannan A, Pinto BJ, Denbrock G, Morales M, Dolby GA, Fiddes IT,
DeNardo DF, Wilson MA. 2023. Complete dosage compensation without balance
between the sexes in the ZZ/ZW Gila monster (*Heloderma suspectum*) revealed by *de*
novo genome assembly. *In review*.
Weir BS, Cockerham CC. 1984. Estimating F-statistics for the analysis of
population structure. *Evolution*. 38: 1358-1370.
Xie HX, Liang XX, Chen ZQ, Li WM, Mi CR, Li M, ... Du WG. 2022. Ancient
demographics determine the effectiveness of genetic purging in endangered lizards.
Mol Biol Evol. 39: msab359.
Zhang, Y., R. Ban, C. Wu, and C. Zhijian. 1996. Studies on Ultrastructure and
Karyotypes of Crocodilian Lizard. China Guanxi Teachers University Press, pp.77.
Ziegler, T., Van Schingen, M., Rauhaus, A., Dang, P. H., Pham, D. T. K., Pham,
C. T. & T. Q. Nguyen (2019): New insights into the habitat use and husbandry of
crocodile lizards (Reptilia: Shinisauridae) including the conception of new facilities for
Vietnamese crocodile lizards *Shinisaurus crocodilurus vietnamensis* in Vietnam and
Germany. - International Zoo Yearbook 53: 250–269.
595

596 **Table(s) and Figures:**
597

Table 1: Top tBLASTx hits in chicken for the CDS of each gene present in the *Shinisaurus* 900kb-SDR. The duplicated *Foxl2* copy is dubbed ENSGALP00000033127-W.

Shinisaurus Gene ID	Chicken Gene Id	Location (Chicken)	E-value
ENSACAP00000003394-D1	ENSGALG00000026187	9:6115461-6115832	1.42E-95
ENSGALP00000008531-D1	*RBP1*	9:6110534-6110713	7.54E-54
ENSACAP00000003392-D1	**No hits.**		
ENSACAP00000003371-D1	ENSGALG00000034575	30:1402166-1402369	2.51E-78
ENSACAP00000003355-D1	ENSGALG00000005357	9:6041607-6041822	0
ENSACAP00000003221-D1	ENSGALG00000005367	9:6037018-6037164	1.03E-80
ENSGALP00000033127-W	***FOXL2W***	**9:5875297-5875587**	**2.44E-76**
ENSGALP00000033127-D1	***FOXL2***	**9:5875297-5875587**	**6.78E-77**
ENSACAP00000003172-D1	*PIK3CB*	9:5800560-5800790	0
ENSGALP00000040175-D1	**No hits.**		
ENSACAP00000002765-D1	**No hits.**		

598
599

Figure 1: Summary of current anguimorph sex chromosome knowledge summarized
from Rovatsos et al. (2019b) indicated by blue and green tips/branches); information
identified in this study indicated by red tips/branches and what remains unknown across
the phylogeny indicated by black tips/branches. Phylogeny from TimeTree using a
representative species from each clade (Kumar et al. 2017) and visualized using Figtree
[v1.4.4] (http://tree.bio.ed.ac.uk/software/figtree/). Of note, "pleurodonts" represents
"non-corytophanid pleurodonts" and "Gg" stands for chicken (*Gallus gallus*) linkage
group.
608
609
610
611
Figure 2: Identification of the ZW sex chromosome system in *Shinisaurus crocodilurus*.
(A) Whole genome F_{ST} scan with a clear peak in a ~1Mb region on LG3. The square
block on LG7 is syntenic with the sex-determining region in *Varanus* and *Heloderma*
(Webster et al. 2023). (B) Isolation and magnification of LG3 F_{ST} peak. (C) Modest
increase in male, relative to female, nucleotide diversity and (D) decrease in
male/female read depth in the region corresponding to the F_{ST} peak on LG3.

Iguania

Corytophanidae (XY, Gg17)

Pleurodonta (XY, Gg15)

Chamaeleo (XY, Gg15)

Pogona (ZW, Gg17/23)

Varanus (ZW, Gg28)

Lanthanotus

Shinisaurus (ZW, Gg9)

Pseudopus

Anguis (*Transition*)

Hyalosaurus

Dopasia

Anguimorpha

Abronia (ZW, Gg28)

Barisia

Gerrhonotus

Elgaria

Wetmorena

Diploglossus

Anniella

Heloderma (ZW, Gg28)

Xenosaurus

150   100   50   0 mya

Shinisaurus sex chromosome identification