Title:		Effects of sexual dimorphism and estrous cycle on murine *Clostridioides difficile* infection
Running title:		*Sex effects on murine C. difficile infections*
Authors:		Jacqueline R. Phan[1], Erica Bacab[1], McKenzie Washington[1], Maria Niamba[1], Dung M. Do[1], Tiffany V. Mata[1], Amber Consul[1], Liahm Blank[1,2], Stephanie Yang[1], Amber J. Howerton[3], Efren Heredia[1], Robert Soriano[3], Chandler Hassan[1], Chad L. Cross[4], and Ernesto Abel-Santos[1,5]*
Affiliation:		[1] Department of Chemistry and Biochemistry, University of Nevada, Las Vegas, Las Vegas, NV, 89154, USA
		[2] Kirk Kerkorian School of Medicine, University of Nevada, Las Vegas, Las Vegas, NV, 89106
		[3] Department of Physical and Life Sciences, Nevada State University, Henderson, NV 89002, USA
		[4] School of Public Health, University of Nevada, Las Vegas, Las Vegas, NV, 89154, USA
		[5] Nevada Institute of Personalized Medicine, University of Nevada, Las Vegas, Las Vegas, NV, 89154
Contact:		4505 Maryland Parkway
		Campus Box 4003

Las Vegas, NV 89154

Tel: (702) 895-2608

Fax: (702) 895-4072

Email: ernesto.abelsantos@unlv.edu

Summary

Clostridioides difficile infection (CDI) is responsible for the majority of identifiable

antibiotic-associated diarrhea. Women are more susceptible to CDI than men. In this study, we

show that female mice developed more severe CDI than males. Furthermore, females in estrus

developed only mild CDI 1-2 days later, while females in proestrus developed deadly disease.

Mirroring the delayed effect of the estrous cycle, pre-infection prolactin levels formed a complex

network with immunoglobulins and cytokines that affected early CDI severity one day after

challenge. Similarly, pre-infection progesterone and luteinizing hormone formed a network that

affected CDI two days after challenge. As expected, immune effectors early in the infection

formed a hormone-independent network that concurrently correlated with CDI severity.

Interestingly, early infection follicular stimulating hormone levels created a network that affected

the CDI recovery phase. In summary, murine sexual hormones affect CDI progression by

affecting the immune system both before and during disease progression.

Keywords: *Clostridioides difficile* infection, CDI, sex hormones, cytokines, chemokines,

immunoglobulins, sexual dimorphism

Background

Clostridioides difficile infection (CDI) is responsible for approximately 25% of all antibiotic-associated diarrhea [1, 2]. In the U.S., over 223,900 CDI cases occur annually, with a mortality rate of up to 6.5% with costs estimated to be $6.3 billion [3-5].

When adjusted for age, immune status, and antibiotic exposure, sex is also an important risk factor in human CDI. Contrary to other GI bacterial infections [6], women are more vulnerable than men to both primary CDI and CDI relapse [7-14]. Furthermore, CDI has been flagged as an emerging threat to pregnant and peripartum women [15-18]. Similarly, older women might be at higher risk of developing CDI due to permanent hormonal depletion after menopause.

An obvious factor underlying sexual dimorphism is the fluctuating hormone concentrations during the female ovulation cycle [19]. Sex hormones have strong effects on both human and murine metabolism [20] and can affect the immune and humoral responses [21-23], as well as susceptibility to bacterial infections [6]. Changes in sex hormone levels can also affect the gut microbiota [24-26], the main protective barrier against C. difficile colonization [27].

Similar to humans, female mice undergo hormonal changes during the estrous cycle [28]. Estradiol levels increase during the proestrus stage, followed by spikes in luteinizing hormone (LH) and follicular stimulating hormone (FSH) [29]. The estrus stage is dominated by rapid increase of prolactin (PRL), followed by a sharp decrease to trigger metestrus entry. The metestrus stage shows a sharp increase of progesterone, with a smaller simultaneous increase in estradiol. Finally, progesterone levels peak during the diestrus stage. Testosterone is produced uniformly throughout the estrous cycle, but its circulating concentration is 10-fold lower than in males [30]. Hence, each murine estrous cycle stage is characterized by a unique pattern of

hormone levels.

We have shown that steroidal sex hormones can interact with *C. difficile* spores and

modulate germination [31, 32]. Similarly, since spore germination is a necessary step for disease

onset, we have found that sterane analogs can inhibit this process and prevent CDI onset in

rodent models of disease [33-43]. During our search for CDI prophylactics, we serendipitously

observed that male mice tended to develop less severe CDI than their female counterparts.

In this study, we used a murine model to assess for sexual dimorphism during CDI. We

found that female mice developed more severe CDI signs than males when infected with three

different *C. difficile* strains. Sexual dimorphism did not disappear under nutritional conditions

known to trigger lethal murine CDI [44]. Within a normal infection, CDI sign severity in female

mice correlated with the estrous stages 1-2 days prior to maximal CDI scores. As a result of this

delayed effect, animals were mostly protected from CDI if they were in estrus 1-2 days before

reaching maximal signs. In contrast, animals developed deadly disease signs 1-2 days after

proestrus. Mirroring the delayed effect of the estrous cycle on CDI, we found that pre-infection

levels of PRL, immunoglobulin IgG2b, cytokine IL-1β, cytokine G-CSF, and chemokine KC

were the primary nodules of a complex network that correlated with CDI symptomatology on the

day after infection. Similarly, we found that the pre-infection levels of progesterone, LH,

immunoglobulin IgG1, chemokine eotaxin, and chemokine IP-10 were the primary network

nodules that affected CDI severity but with a 2-day delay. As expected, the levels of

immunoglobulins, cytokines, and chemokines during early infection formed a hormone-

independent network that concurrently correlated with CDI severity. Interestingly, early infection

levels of FSH, together with cytokine IL-1β, and chemokine KC were the main nodules of a

network that affect CDI severity 2-days later, during the recovery phase of the infection. In

summary, we show that murine female sexual hormones affect CDI progression, likely as a

function of immune system modulation both before infection and during disease progression.

Materials and Methods

Materials: *C. difficile* strains R20291 and 9001966 were donated by Professor Nigel

Minton at the University of Nottingham, United Kingdom. *C. difficile* strain 630 was obtained

from the American Type Culture Collection (ATCC, Manassas, VA). The synthetic bile salt

analogs, CAMSA and CAPA were prepared by Professor Steven M. Firestine at Wayne State

University in Detroit, Michigan. Standard Laboratory Rodent Diet was obtained from LabDiet

(St. Louis, MO, USA). Custom purified diet (Atkins-type diet) containing approximately 50%

protein and 25% fast was purchased from TestDiet (TestDiet Catalog # 18181230286). All *C.*

difficile strains were grown and maintained in a Coy Laboratories (Grass Lake, MI) vinyl

anaerobic chamber (5% H_2, 5% CO_2, 95% N_2).

Animals: All procedures involving animals in this study were performed in accordance

with the Guide for Care and Use of Laboratory Animals outlined by the National Institutes of

Health. The protocol was reviewed and approved by the Institutional Animal Care and Use

Committee (IACUC) at the University of Nevada, Las Vegas (Permit Number: R0914-297).

Weaned C57BL/6 mice were purchased from Charles River Laboratories (Wilmington, MA,

USA). Mice were housed in groups of five per cage at the University of Nevada, Las Vegas

animal care facility. Upon arrival at the facility, animals were allowed to acclimate for at least

one week prior to the start of experimentation. All post-challenge manipulations were performed

within a biosafety level 2 laminar flow hood.

C. difficile spore harvest and purification: *C. difficile* cells from frozen stocks were

streak plated in an anaerobic chamber onto BHI agar supplemented with 2% yeast extract, 0.1%

L-cysteine-HCl, and 0.05% sodium taurocholate (BHIS) to yield colonies (15). After 48 hours, a

single colony was inoculated into BHI broth supplemented with 0.5% yeast extract and incubated

for 48 hours. The inoculated broth was then spread plated onto multiple BHI agar plates prepared

as described above. Inoculated plates were incubated for 7 days at 37 °C.

Plates were then flooded with ice-cold deionized (DI) water. Cells and spores were

harvested by scraping bacterial colonies from the plates. The cell and spore mixtures were

pelleted via centrifugation at 8,000 ×g for 5 minutes, resuspended in DI water, and pelleted

again. This washing step was repeated twice more. The cell/spore mixture was then centrifuged

through a 20% to 50% HistoDenz gradient at 18,200 ×g for 30 minutes with no brake (22).

Under these conditions, spores pelleted at the bottom of the centrifuge tube, while cell debris

remains above the 20% HistoDenz layer. Pelleted spores were transferred to a clean centrifuge

tube and washed five times before storing in DI water at 4 °C. To determine spore purity,

selected samples were stained using the Schaeffer-Fulton endospore staining method (23) or

were visualized via phase-contrast microscopy. Spore preparations used were greater than 95%

pure. Spores were washed three times with DI water, heat shocked at 68°C for 30 minutes, then

washed three additional times before being resuspended in DI water prior to administration.

Estrous cycle tracking in mice: The estrous cycle tracking method performed in this

study was adapted from *McLean et al., 2012* [29]. Mice estrous stages were tracked for at least

one week prior to any manipulation to ensure that animals were cycling normally. Animals were

tracked daily for their estrous cycle stage via vaginal lavage at the same time each morning

throughout the duration of the study. First, excess urine is dabbed off with low-lint Kimwipes. A

micropipette was used to lavage 20 μL of PBS into the vaginal orifice of the animal. Water was

swooshed 3-4 times around the vaginal orifice until the suspension appeared mildly cloudy. The

20 μL sample was then transferred onto a clean glass microscope slide and left to air dry for at

least 1 hour. Animals were monitored to ensure there were no signs of vaginal infection.

Air-dried vaginal sample slides were dipped into a 2% crystal violet solution inside a

Coplin jar for 1 minute. Slides were then moved to a Coplin jar containing fresh DI water to

wash off excess stain. Slides were dabbed gently with clean wipes, allowed to air dry, and

examined using a compound light microscope at 100x and 400x magnification (**Figure S1**) [45].

Murine CDI model: The murine CDI model used in this study was adapted from Chen

et al. [42]. Mice were given three consecutive days of antibiotic cocktail containing kanamycin

(0.4 mg/ml), gentamycin (0.035 mg/ml), colistin (850 U/ml), metronidazole (0.215 mg/ml), and

vancomycin (0.045 mg/ml) *ad libitum* (14). Mice were then given DI water for the remainder of

the experiment. On the day prior to *C. difficile* challenge (day -1), mice were given an

intraperitoneal (IP) injection of 10 mg/kg clindamycin. On the day of infection (day 0),

experimental groups were challenged with 10^8 *C. difficile* spores via oral gavage.

Mice were observed for signs of CDI twice daily and disease severity was scored

according to a CDI sign rubric adapted from previously published work [36]. According to the

rubric, pink anogenital area, mild wet tail, and weight loss of 8-15% were each given a score of

1. Red anogenital area, lethargy/distress, increased diarrhea/soiled bedding, hunched posture, and

weight loss greater than 15% were each given a score of 2. Animals with a cumulative scored

less than 3 were indistinguishable from noninfected controls and were considered non-diseased.

Animals scoring 3–4 were considered to have mild CDI. Animals scoring 5–6 were considered to

have moderate CDI. Animals scoring greater than 6 were considered to have severe CDI and

were immediately culled.

Tracking the estrous cycle in the murine CDI model: To better visualize the diverse

range of murine CDI signs [46], the hypervirulent strain R20291 was employed for all estrous-

related studies. Estrous stages for each mouse were tracked every morning at the same time via

vaginal lavage and cytology. Mice that were not cycling or cycling irregularly were excluded

from the study. In the first round of the study (cohort 1), animals were sacrificed at random on

day 0 (prior to *C. difficile* challenge), day 1, day 2, and day 3 (post- *C. difficile* challenge) after

daily estrous stage sampling. Animals that were severely ill were immediately culled regardless

of day or estrous stage. Surviving animals on each day were analyzed for estrous stage and CDI

signs.

After initial analysis of data from the first experiments showed that estrous stages have a

delayed effect on CDI severity, the protocol was further modified in an attempt to specifically

track certain estrous stages and collect non-terminal blood samples on the days prior to expected

maximal CDI signs. In the second round (cohort 2), mice were pooled based on their predicted

estrous stage on the day of spore challenge. This prediction was estimated from the estrous stage

they were on at the start of antibiotic treatment five days prior to infection. For each estrous stage

grouping, animals were randomly selected for blood sampling via facial vein bleed on day 0 (day

of challenge) or day 1 (24 hours after challenge). Animals were monitored for CDI signs, as

above and sacrificed either one day or two days after blood samples were collected.

Blood sample processing for sex hormones, cytokines/chemokines, and

immunoglobulins: Approximately 100 μl of blood was collected from each mouse via facial

vein bleeds. The blood rested at room temperature for 30 minutes prior to spinning down to

collect serum. Serum was frozen at -20°C and then stored at -80°C until processing. Each serum

sample was divided into four aliquots.

The first aliquot was analyzed to determine the concentration of follicle-stimulating

hormone (FSH), luteinizing hormone (LH), and prolactin (PRL) using the Milliplex Mouse

Pituitary Magnetic Bean Panel from Millipore according to kit instructions on a Bio-Plex 200

system. The second aliquot was processed using Millipore's Milliplex Mouse

Cytokine/Chemokine Magnetic Bead Panel according to kit instructions to test for cytokines IL-

13, IL-10, IL-2, IP-10, IL-5, RANTES, KC, IL-15, IL-1α, IL-1β, IL-17, IL-6, TNF-α, G-CSF,

eotaxin, IFN-γ, and GM-CSF. The third aliquot was processed using Milliplex Mouse

Immunoglobulin Isotyping Magnetic Bead Panel from Millipore according to kit instructions to

test for IgM, IgG, IgA, IgE, and IgD.

The fourth aliquot was sent to The Metabolomics Innovation Centre (TMIC, Alberta,

Canada) for analysis of progesterone concentrations by HPLC-MS. Briefly, 10 μl of plasma was

extracted with 40 μl of acetonitrile containing 3.1 nM progesterone-d9 internal standard

(Cayman chemical, Ann Arbor, MI). The extraction mixture was sonicated in a water bath for 10

minutes, centrifuged at 21000 x g for one minute, and 40 μl of the supernatant was transferred to

the autosampler vials with glass inserts. LC-MRM-MS analysis was performed using a Nexera

XR UPLC system (Shimadzu, Japan) coupled to a Sciex QTrap 6500+ mass spectrometer

equipped with an IonDrive Turbo V electrospray ionization source. Chromatography was

performed using a Zorbax Eclipse Plus C18 column (2.1 × 150 mm, 1.8 μm, Agilent, USA) at a

flow rate of 0.2 mL/min of mobile phases 0.1% formic acid in water (A) and 0.1% formic acid in

acetonitrile (B). The LC gradient was 50-100% B / 0.0-5.0 min, 100-50% B / 5.0-5.1 min, 50% B

/ 5.1-8.0 min. The sample injection volume was 10 μl, the column oven was off, and the

autosampler temperature was 15°C. MRM-MS was performed using electrospray ionization in

the positive mode. 315.2/97.1 and 324.3/100.1 Q1/Q3 transitions were monitored for

progesterone and progesterone-d9, respectively (1). Dwell time was 50 ms, collision energy (CE)

32 V, declustering potential (DP) 50 V, entrance potential (EP) 10 V, collision cell exit potential

(CXP) 12 V, source temperature 450 ºC; ion source gas 1 40 psi; ion source gas 2 40 psi. The

collision gas was set to high, and the curtain gas was set to 40 psi. Data acquisition and analysis

was performed using Analyst 1.6.3 software (Sciex, Framingham, MA). Progesterone

concentrations were determined from chromatographic peak area ratios of the analyte and the

internal standard.

Statistical analyses: For the murine CDI model, severity of signs was visualized via box-

and-whisker plots with a minimum of three independent values ($n \geq 3$). One-way ANOVA or

one-tailed *t*-test were performed, as appropriate, using R (v. 4.4.3, R Core Team, 2025) to assess

differences between groups at every time point, with significance defined as $p < 0.05$. Significant

main-effect ANOVAs were analyzed using Scheffe's correction for *post hoc* comparisons

between all groups. Time-to-event data were analyzed using Kaplan-Meier survival analysis

models, with log-rank tests for curve comparisons. For assessment of associations among

variables, we utilized the robust Kendall's Tau correlation coefficient for bivariate comparisons

among hormone levels, cytokine levels, immunoglobulin levels, and CDI severity scores, with

significance defined as $p < 0.10$. Correlations were also calculated for combined samples and

individually by bleed day (day 0 or day 1).

Results

Female mice exhibit more severe CDI than male mice. We infected male and female

C57BL/6 mice with three different *C. difficile* strains [33]. One group of male and female mice

was infected with spores from *C. difficile* strain 9001966 (ribotype 106) (**Fig. 1A**). For both

sexes, mice started to develop signs of CDI on day 2 post-challenge (day 2). Male mice showed

varied CDI sign severities during days 2-4, but all animals recovered by day 5. Female mice, on

the other hand, showed more homogenous mild to moderate CDI symptomatology in days 2-3

and all animals recovered by day 4. Even though the CDI timeline was shorter in female mice,

these animals developed statistically more severe signs than their male counterpart at the peak of

infection (**Fig. 1A**).

A second group of male and female mice was infected with spores from *C. difficile* strain

630 (ribotype 012). Male mice infected with *C. difficile* strain 630 spores showed less individual

variability in CDI symptomatology and shorter infection progression compared to animals

infected with *C. difficile* strain 9001966. Male mice infected with *C. difficile* strain 630 did not

show any signs on day 1 after challenge but started to show CDI signs by day 2. Female mice

started to develop mild CDI signs one day after spore challenge (**Fig. 1B**). By day 2, female mice

showed a notable increase in CDI sign severity, followed by a return to mild CDI signs by day 3

and complete remission by day 5 post-challenge. Similar to female mice, CDI signs for male

mice started to resolve by day 3 and disappeared by day 5 post-challenge. In both male and

female mice infected with *C. difficile* strain 630, CDI sign severity peaked on day 2 post-

challenge. Notably, male mice showed mostly mild-to-moderate sign severity, while female mice

showed statistically more severe symptomatology (**Fig. 1B**).

A third group of male and female mice was infected with spores from hypervirulent *C.*

difficile strain R20291 (ribotype 027). Male mice challenged with *C. difficile* strain R20291

started to develop moderate CDI symptoms by day 1 post-spore challenge (**Fig. 1C**). Male mice

remained moderately symptomatic until day 4, when CDI signs started to resolve. Similar to

animals infected with *C. difficile* strain 9001966 and *C. difficile* strain 630, all male mice

infected with *C. difficile* strain R20291 recovered and showed no CDI signs by day 4.

In contrast, female mice challenged with *C. difficile* strain R20291 spores developed

statistically more severe CDI signs than male mice the day after spore challenge (**Fig. 1C**).

Indeed, while all males infected with *C. difficile* strain R20291 recovered from CDI, more than

half of the female cohort succumbed to the infection (**Fig. 2B**, p=0.049). Surviving female mice

started to slowly recover from CDI and showed no symptoms by day 4. Because animals

challenged with *C. difficile* strain R20291 developed the most severe CDI signs, we used this

hypervirulent strain in subsequent experiments.

A high-protein/high fat (Atkins-type) diet aggravates CDI in mice of both sexes. We

recently showed that an Atkins-type diet can exacerbate CDI severity in mice [36]. To determine

if sex affects CDI exacerbation by diet, male and female mice challenged with *C. difficile* strain

R20291 while on this high protein/high fat (Atkins-type) diet showed delayed onset of CDI signs

compared to animals fed a standard diet (**Fig. 2A**). Although female mice fed an Atkins-type diet

did not show statistically more severe CDI signs (**Fig. 2A**) or survival (**Fig. 2B**) than female

mice fed a standard diet, by day 4 post-challenge all female mice had developed severe CDI and

had to be culled.

In contrast, once the infection started, male mice fed an Atkins-type diet developed

statistically more severe CDI signs compared to challenged mice fed a standard diet (**Fig. 2A**).

On a standard diet, 100% of males survived compared to 40% of females (p=0.049). Indeed,

while all male mice on a standard diet survived CDI, 75% of male mice fed an Atkins-type diet

developed signs severe enough to require euthanasia (**Fig. 2B**). In fact, survival for male mice on

an Atkins-type diet was statistically worse than male mice fed a standard diet (p=0.025)

comparable to female mice fed a standard diet, but still better than females on the Atkins-type

diet (p=0.018).

Estrous stages have a delayed effect on CDI severity prior to infection. Using vaginal

smear cytology (**Figure S1**), we were able to monitor the estrous cycle of female mice. Animals

were binned (**Fig. 3**) into estrus stage (blue columns), metestrus stage (orange columns), diestrus

stage (green columns), and proestrus stage (magenta columns). Some animals could not be

unambiguously assigned to a specific estrous stage. These animals showed borderline cytology

readings and were thus binned into two new categories: late diestrus/early proestrus (pink

columns) and late proestrus/early estrus (yellow columns).

To determine the effect of the estrous cycle on murine CDI severity, a first set of female

mice in distinct stages of the estrous cycle were challenged with *C. difficile* strain R20291

spores. We continued to simultaneously monitor the estrous cycle and CDI sign progressions. As

the estrous cycle does not stop during the course of infection, animals could cycle across

different estrous stages while CDI signs are being monitored. As a first approach, we tried to

correlate daily CDI signs for each animal with their concurrent estrous stage, but we found no

statistical significance between groups (**Fig. 3A**).

Intriguingly, when daily CDI severity was plotted against the estrous stages of the prior

day, we found distinct daily patterns (**Fig. 3B**). The strongest statistical correlation between CDI

signs and prior day estrous stages was found on day 1 post-challenge. Animals that were in

estrus when challenged with *C. difficile* spores (day 0), showed no CDI signs the following day

(day 1). Compared to animals that were in estrus on day 0, animals that were in metestrus,

diestrus, and late diestrus/early proestrus during spore challenge developed statistically higher

CDI severity the following day. Finally, animals that were at proestrus at day 0 developed the

most severe CDI symptoms the day after challenge with more than half reaching the clinical

endpoint.

The effect of prior-day estrous stages on CDI sign severity started to dissipate by day 2

post-challenge. On day 2 post-challenge, animals that were in estrus on day 1 showed no CDI

symptoms. In contrast, animals that were in metestrus, diestrus, or late proestrus/early estrus on

day 1 post-challenge showed statistically higher CDI sign severity. Unfortunately, there was an

insufficient number of animals in the proestrus and the late diestrus/early proestrus groups on

day 1 for a complete statistical analysis of CD severity on day 2 post-challenge. Furthermore,

because of the high CDI-mediated mortality on days 1 and 2, there were insufficient numbers of

animals left for statistical analysis on CDI score on day 3 post-challenge vs estrous stages.

To determine if the prior-day estrous effect is independent of the infection time course,

we combined the data for each estrous stage regardless of the day they were determined. This

allowed us to increase the number of animals in each estrous stage for statistical analysis. As

expected, plotting CDI severity against their concurrent estrous stages showed no statistically

significant differences (**Fig. 4A**).

However, when CDI severity for each animal was plotted against their prior-day estrous

stage, a similar delayed pattern was observed as in **Fig. 3B**. Indeed, when animals were in estrus

at any day during infection, they showed few CDI signs the next day (**Fig. 4B**). Animals that

were in metestrus, diestrus, and diestrus/proestrus at any time showed more varied but similar

intermediate symptomatology one day later. In contrast, animals in proestrus at any time during

infection became extremely sick the next day and had to be culled. Any animals in

proestrus/estrus showed next-day CDI severity intermediate between proestrus and estrus.

Estrous stages are correlated with day of maximal CDI sign severity. During the

previous part of the study, only vaginal lavage samples were collected and examined to

determine the estrous stage of animals. To further investigate the delayed effect of the estrous

stages on CDI severity, a second cohort of mice were tracked and timed for blood collection and

sacrificed based on their predicted estrous stage on the day of spore challenge with hypervirulent

strain R20291. Instead of collecting terminal blood samples, facial vein blood samples were

taken 1-2 days prior to sacrifice. Contrary to expectations, mice did not develop maximal CDI

signs 1-day post-challenge as the previous cohort of mice (**Fig. 1A**). Instead, all mice developed

the most severe CDI signs 2 days after infection. Consequently, no correlation was observed

between estrous stage (via vaginal lavage and cytology) and day 1 post-challenge.

Since the maximum CDI signs for the second cohort did not develop until 2-days post-

challenge, the temporal effect of the estrous stage was also delayed (**Fig. 5**) compared to animals

in the first cohort (**Fig. 4B**). Indeed, second cohort animals that were in the estrus stage showed

mild CDI signs two days later. In contrast, second cohort animals in the proestrus stage

developed more severe CDI signs two days later.

Biomarkers obtained before infection (D0) and during early infection (D1). Pre-

infection blood samples (D0, obtained hours before spore challenge) and early infection blood

samples (D1, one day after spore challenge) were obtained from female mice to test for the effect

of hormones, cytokines, and immunoglobulins on CDI progression. As expected, pre-infection

(D0) hormone levels varied between individuals since animals were at different estrous stages

when they were challenged with *C. difficile* spores on day 0. Interestingly, we also found inter-

individual variabilities for pre-infection levels of cytokines and immunoglobulins, even though

blood was collected before spore challenge. In contrast, early infection (D1) cytokine and

immunoglobulin levels were more varied, as expected for animals progressing through active

infection and showing different sign severities.

Discussion

Few attempts have been made to understand how sex affects CDI onset and severity. To

assess the clinical importance of sex on human CDI, we did an extensive PUBMED search for

primary articles relating to CDI risks. We found that most studies (n=87) did not stratify for sex.

Of the studies that assessed sex as a variable (n=24), most linked CDI to other comorbidities.

These studies tended to have small sample sizes, making it difficult to determine if sex was

associated with CDI or rather to the underlying comorbidity. However, when CDI was the

primary diagnosis, multiple large cohort studies showed that women are at higher risk than men

of developing symptoms [7, 8, 10, 11, 47]. Indeed, a multicenter retrospective cohort study of

over 1 million patients at 150 US hospitals showed that the odds of CDI in women was 1.2 times

as likely as in men [47]. Although the correlation between sex and human CDI is clear, testing

for susceptibility in women is cost prohibitive and impractical. Yet patient outcomes could be

dramatically improved if clinicians could adapt antibiotic treatments to match specific hormone

levels.

In this study, the murine model recapitulates many of the differences in CDI outcome

observed between men and women [7, 8, 10, 11, 47]. For instances, during our investigations

into the factors affecting CDI progression and prophylaxis [33, 40, 41], we serendipitously found

that female mice developed more severe CDI signs than males.

To further test for sexual dimorphism in the murine model of CDI, we infected mice with

three different *C. difficile* strains. These strains represent different ribotypes and were selected

because they show distinct timelines for CDI onset. *C. difficile* strain 9001966 (ribotype 106) is a

clinical strain from the United Kingdom and shows delayed onset of murine CDI. *C. difficile*

strain 630 (ribotype 012) is a type strain [48, 49] that we have extensively tested for its capacity

to induce CDI symptoms in mice [40, 41]. Finally, *C. difficile* strain R20291 (ribotype 027) is a

hypervirulent strain that causes earlier infection onset and more severe murine CDI signs than

other strains [33].

When animals were infected with any of these strains, female mice developed statistically

more severe CDI than their male counterpart. The timing of these differences varied, but in every

case, the most significant differences in CDI severity coincided with the peak of disease severity

in their respective infection timeline. Because animals challenged with hypervirulent *C. difficile*

strain R20291 showed the strongest CDI signs, we continued to use this strain for all subsequent

experiments.

In contrast to hamsters [38, 50], murine CDI seldom tends to be lethal [33, 40, 41].

However, we recently showed that an Atkins-type (high-protein/high-fat) diet can greatly

increase murine CDI sign severity duration and mortality [36]. Here, we determined if the

Atkins-type diet affected CDI severity between male and female mice. When animals were fed a

diet known to exacerbate CDI signs, mortality of male mice increased significantly. Even then,

male mice showed less severe signs than females fed the same diet. This shows that the sexual

dimorphism of murine CDI is still present under conditions where CDI is severe.

The most obvious difference between male and female mice is the female estrous cycle.

However, intraday correlation between estrous cycle and CDI severity showed no statistical

differences at any time during the infection. In contrast, estrous stages seem to have a 1 day to 2

days delayed effect on CDI severity.

The interplay between estrous stage cycling and CDI progression is quite complex. For

CDI caused by *C. difficile* strain R20291, infection starts and reaches a maximum of severity on

days 1 or 2 following spore challenge. Surviving animals completely recover by days 4-5 post-

challenge. In contrast, the estrous cycle can take between 3-6 days to be completed. Furthermore,

the duration of each estrous stage is variable, with proestrus and metestrus lasting as little as 8

hours and diestrus lasting as long as 72 hours [51]. This was further complicated by the

observation that estrous stages showed a 1–2-day delayed effect on CDI severity. Thus, binning

animals into different estrous stages for each day reduced our statistical power for some of the

estrous stages. Nevertheless, our data clearly shows that animals challenged while in the estrus

stage were largely protected from CDI. In contrast, animals challenged while in the proestrus

stage developed deadly CDI.

The contrast between proestrus-mediated and estrus-mediated CDI susceptibility could

not be starker. While CDI is most deadly following proestrus, estrus offers strong CDI protection

to animals on the typical day of maximal disease severity whether1 day or 2 days after estrous

staging. Furthermore, metestrus and diestrus correlate with more heterogeneous CDI

symptomatology on days of maximal onset.

An obvious factor underlying CDI sexual dimorphism is the fluctuating hormone levels

during the female ovulation cycle [19]. In general, CDI-exacerbating proestrus is characterized

by large increases in estradiol, progesterone, LH, and FSH levels and low concentrations of

prolactin [29]. These spikes in hormone levels signal entry into the CDI-protective estrus. At this

point, the high concentrations of circulating estradiol, progesterone, LH, and FSH characteristic

of proestrus are severely diminished followed by an increase of serum PRL. The metestrus and

diestrus stages tend to have hormonal levels that fall in between those of proestrus and estrus

[29].

Correlation analysis of pre-infection (D0) biomarkers form two, almost separate cluster.

In fact, cytokine KC is the only connection between the two clusters. Cluster 1 (**Fig. 6**, top) is

centered around pituitary hormones FSH, PRL, and LH. Cluster 2 (**Fig. 6**, bottom), on the other

hand, is centered around steroidal hormone progesterone.

From cluster 1, pre-infection (D0) levels of hormone LH, hormone FSH, and cytokine IL-

10 form a positively correlated network that negatively influences CDI severity during the

following day (day 1 after spore challenge) through PRL and cytokine IL-1β. Only cytokine KC

positively correlate with CDI sign on day 1 post-challenge while simultaneously negatively

correlating with IL-1β and IgA levels and positively correlating with IL-6.

From cluster 2, pre-infection (D0) levels of hormone progesterone, cytokine IL-1α,

immunoglobulin IgG1, immunoglobulin IgG3, immunoglobulin IgM, and immunoglobulin

IgG2b form a complex positively correlated group that negatively affect CDI severity on the

following day through immunoglobulin IgG2b. From the same group, IgM and IgG3 negatively

correlate with cytokine G-CSF. In turn, G-CSF negatively influences CDI score on day 1.

Mirroring the temporal delay between estrous stages and CDI severity, pre infection (D0)

levels of PRL, FSH, LH, cytokine IP-10, and cytokine eotaxin form a group that positively

correlate with CDI severity two days post-challenge. The same network negatively affect CDI

severity two days post-challenge trough LH

The two-day delayed effect is also observed in cluster 2. Pre infection (D0) levels of the

progesterone/immunoglobulin group negatively influences CDI severity two days post-challenge

through IgG1 and progesterone. As expected after spore challenge, early infection (D1)

immunoglobulin, cytokine, and chemokine levels create a single complex network that correlate

CDI severity on the same day (**Fig. 7**). Early infection levels of LH and FSH positively influence

same day CDI sign severity through IFNγ and KC, respectively.

We also observed a temporal effect between early infection biomarkers and CDI

recovery, two days later. Indeed, early infection (D1) levels of KC and IL-1β show strong

positive correlation to CDI severity on day 3 post-challenge. Central to these effects, early

infection (D1) FSH levels shows strong negative correlation with early infection (D1) levels of

KC and IL-1β. FSH also shows negative correlation with CDI severity on day 3 post-challenge.

We have previously shown that *C. difficile* spores germinate between 6- and 9-hours

post-challenge in the intestines of mice. Potentially, the strong association between day 0 estrous

stages, pre-infection biomarker levels, and CDI severity on days 1-2 could be due to estrous

cycle and hormone fluctuations influencing the initial stages of CDI onset. On the other hand,

delayed effects of the estrous cycle and sex hormones seem to continue throughout CDI

progression into infection recovery. This suggests that estrous stages can increase or decrease

female mice susceptibility to CDI both at the early and late stages of the infection course.

The negative correlation of pre-infection PRL (D0) with early CDI signs mirrors the

protective effect of the estrus cycle. However, the fact that pre-infection (D0) levels of

progesterone and LH show a time-delayed, negative correlation to CDI severity is more difficult

to explain. It should be also noted that the boundaries between estrous stages are arbitrary and

dependent on cellular response to hormonal changes. In contrast, changes in sex hormone levels

follow a time continuum throughout the estrous cycle. For example, even though progesterone is

expected to fall during estrus, entry into the stage happens when progesterone levels are at its

maximum. Hence, small variations in sample collection timing, could result in uncoupling

between expected sex hormone levels and identified estrous stages. Indeed, murine sex hormone

fluctuations have been shown to follow strain-specific patterns (REF) and even show significant

variabilities between individuals on the same assigned estrous stages. These temporal

uncertainties could partially explain the paradoxical nature of sex hormone effects on CDI

severity.

Conclusions

Human CDI is not diagnosed unless symptoms are severe enough to warrant clinical intervention. In that sense, CDI diagnosis serves as a surrogate for symptoms severity in humans. Similar to other infection models, murine CDI provides an idealized and controlled (albeit imperfect) model to study disease onset, severity, and progression. The murine CDI model presents symptoms progression similar to human CDI and animals respond to the same treatments [52, 53]. We have found that murine CDI follows a well-ordered progression of disease signs. Based on these observations, we have implemented a rubric for CDI severity development [33, 41].

Little is known regarding sex hormones, their role in intestinal infection, and the adaptation of the immune response in the GI tract. The delayed effect of the estrous stage and sex hormones on CDI severity might be due to the time needed for these hormones to affect host physiological (e.g., immune system, intestinal microbiota, steroidal hormone accumulation in the gut) responses during infection.

During pregnancy in humans and mice, progesterone stimulates the expression of IL-4 and IL-10 and favors the activation of Th2 cytokines while suppressing the major Th1 cytokines, IL-12p70, IL-2, IFN-y, IL-18 [54-56]. However, little is known about the role of progesterone in intestinal infections. A recent study showed that exposure to progesterone resulted in decreased *Listeria monocytogenes* burden in cultured intestinal cells [57].

PRL is both a hormone and a cytokine with roles in reproduction, lactation, and stress regulation [58]. PRL has been shown to have immune modulating effects. In this study, pre-infection PRL levels affect early CDI progression. This is consistent with typical chronological presentation of hormones in the murine estrous cycle with PRL levels rising approximately one

day (estrus stage) prior to elevation of progesterone levels (metestrus-diestrus stage). However,

pre-infection PRL does not seem to correlate with any cytokine, chemokine, or immunoglobulin

that that affect early CDI progression. Thus, the mechanism by which PRL affect CDI onset is

not clear.

Of all the immunoregulatory compounds tested, pre-infection PRL only correlated with

IP-10 levels. Similar to what we observed in this murine model, a strong direct correlation

between severe CDI and upregulation of IP-10 has been observed in humans [59]. Hence, pre-

infection PRL spike during murine protective-estrus stage, may cause a decrease in pre-infection

IP-10 production resulting in less severe disease two days later. In our study progesterone did not

show any significant direct relationships with any chemokines and cytokines. Nevertheless,

animals with higher levels of pre-infection progesterone were protected from CDI onset two days

later. In contrast to the lack of effect on cytokines, pre-infection progesterone forms a positive

correlated group with almost all pre-infection immunoglobulins tested. Although progesterone's

effect on immunoglobulins is not well understood, progesterone's immunomodulatory

capabilities suggest that it can potentially influence immunoglobulin productions. Indeed,

murine, human, and nonhuman primate, cervicovaginal IgG subtypes are highest during times of

increased progesterone [60].

We observed a similar positive correlation between pre-infection progesterone levels and

pre-infection IgG subtypes. In turn, both IgG1 and IgG3 showed correlations with cytokines and

chemokines. IgG1 positively correlated with IL-1a, a dual role cytokine involved in homeostasis

and the immune response [61] and negatively correlated with proinflammatory cytokine

RANTES. In turn, IgG3 negatively correlated with proinflammatory cytokines G-CSF and

RANTES.

Stimulated macrophages and other immune cells produce G-CSF and RANTES in

response to infection. Both are important as chemoattractants and aid in maturation and

activation of immune cells [62, 63] In fact, *C. difficile* has been shown to initiate a strong G-CSF

and RANTES expression in the mice intestinal tract which may be used as markers for severity

of disease [64, 65]. The direct and indirect effects of progesterone on immunoglobulins and

cytokines could account for protection from CDI in mice.

Previous work in the Abel-Santos laboratory has shown that progesterone can inhibit *C.*

difficile spore germination *in vitro*. Interestingly, sex steroidal hormones are also present at low

concentrations in the intestinal tract and show a 1-day accumulation delay compared to serum

[32], a timeline reminiscent to the delayed effect of progesterone on murine CDI. Due to the

heterogenous environment of the intestinal lumen, there could be localized areas of higher

hormone concentrations where they could interact with *C. difficile* spores. Whether progesterone

affects CDI indirectly by modulating the immune system or directly by inhabiting spore

germination, needs to be elucidated.

The gonadotropin LH is typically regarded as a reproductive hormone but studies have

shown a correlation to intestinal cells where LH binds to receptors expressed in the intestine and

slows gastric motility [66]. To our knowledge, the role of LH in intestinal infections has not been

reported. Similar to progesterone, we found a negative correlation between pre-infection LH

levels and CDI severity two days later. Pre-infection LH formed a positive correlated group with

the other two pituitary hormones, PRL and FSH. Furthermore, like PRL, LH also negative

correlated with pre-infection IP-10 while also positively correlating with eotaxin. Pre-infection

IP-10 and eotaxin levels strongly positively correlate to CDI severity two days later. IP-10 has

been reported as a potential biomarker for severity of disease including CDI [59, 67]. Eotaxin is

a chemokine that recruits eosinophils to areas of inflammation and has been shown to play a role

in the onset of the murine estrus cycle [68]. similar to IP-10, higher eotaxin levels have been

associated with more severe CDI in humans [69]. Eotaxin eosinophil recruitment has also been

shown to be necessary to prevent severe tissue damage in CDI [69].

After infection onset, the effect of sex hormones on disease progression seems to weaken.

This might be due to the complex immune response that occurs after infection onset. At this

point, the robust feedback look of the immune system would become more important than the

more subtly effects of sex hormones.

Nevertheless, we found a potential PRL/FSH effect on CDI recovery. Early infection

PRL levels positively correlated with IL-1β. Meanwhile, early infection FSH negatively

correlated with both IL-1β and KC. In a mutual feedback loop, IL-1β can stimulate the secretion

of PRL and PRL can increase the secretion of IL-1β through stimulated differentiation of T cells

[58, 70]. Although IL-1β is typically considered a pro-inflammatory signal stimulated by

infection, IL-1β has been shown to be complex and also play a role in commensal microbiota

regulation and maintenance [71]. We are likely seeing this early infection loop playing a role in

CDI recovery. Little is known regarding the effect of FSH on cytokine and chemokine

production in the intestine of humans or mice. In fact, receptors for FSH are not highly expressed

in the intestine [72]. In osteoclast differentiation FSH has also been shown to induce the

expression of IL-1β and IL-6 from monocytes [73, 74]. Hence, In the murine intestine, both pre-

infection and early infection FSH may have a role in the upregulation of IL-1β and IL-10 to aid

in tissue maintenance and repair. These complex interactions between hormones,

immunoglobulins and cytokines/chemokines warrants more study to better understand the onset

and progression of CDI in mice and humans. Approaching *C. difficile* infection by studying the

hypothalamic-pituitary-gonadal (HPG) axis, allows us to determine how sex hormones and the

estrus cycle affect the severity and progression of CDI. Many cells of the immune system contain

receptors for sex hormones which can activate cytokines, showing a connection between immune

modulation and sex hormones. The full host immune response initiated by infection with *C.*

difficile spores is yet to be fully elucidated. Many reports have identified cytokine and

chemokine correlations between expression and severity of disease with similarities and

differences in these reports suggesting dual roles in pathogenicity and protection. There is likely

a very complex crosstalk between the systemic host immune response, the local immune cells

and intestinal microbiota resulting in a fine line between infection clearance and tissue damage.

Together, these data suggests that sex hormones could affect CDI symptomatology

directly by modulating spore germination in the intestine and/or indirectly by globally changing

the host's gut microbiome and/or immune status. The effect of individual hormones on CDI

severity needs to be established but provides an open door into explorations on the relationship

between sex hormones, the gut, and CDI.

Acknowledgments

Acknowledgments

This work was supported by the National Institute of Allergy and Infectious Diseases at

the National Institute of Health [grants numbers R15AI164346, R16AI175022] to EAS, and by

the National Institute of General Medical Sciences at the National Institute of Health [grant

number P20GM12132] to NIPM.

Figures and Figure Legends

Figure 1. *Effect of sex on CDI severity.* Box-and-whisker plots of CDI sign severity in female

mice (pink bars) and male mice (blue bars) challenged with *C. difficile* spores from (A) strain

9001966, (B) strain 630, and (C) strain R20291. Signs of severity were determined daily based

on the CDI scoring rubric reported previously [36]. Animals with a score of <3 were

indistinguishable from noninfected controls and were considered non-diseased. Animals with a

score of 3 to 4 were considered to have mild CDI. Animals with a score of 5 to 6 were

considered to have moderate CDI. Animals with a score of >6 were considered to have severe

CDI and were immediately culled. Dashed line represents the clinical end point. Horizontal bold

lines represent median values, asterisks represent mean values, bars represent interquartile

ranges, error bars (whiskers) represent maximum and minimum values, and dots represent inner

point values. Daily statistical differences between males and females were determined by one-

tailed *t*-test.

Figure 2. *Effect of Atkins-style diet on mice CDI survival.* (A) Box-and-whisker plots of CDI

sign severity and (B) Kaplan-Meier survival plots for male mice fed a standard diet (solid blue),

female mice fed a standard diet (solid pink), male mice fed an Atkins-type diet (blue crosshatch),

and female mice fed an Atkins-type diet (pink crosshatch) challenged with *C. difficile* spores

from strain R20291 [36]. For panel A, horizontal bold lines represent median values, asterisks

represent mean values, bars represent interquartile ranges, error bars (whiskers) represent

maximum and minimum values, and dots represent inner point values. Single-factor ANOVA

was performed at every time point to assess differences among the sign severity means of the

different groups. ANOVA results with *P* values of <0.05 were analyzed *post hoc* using the

Scheffe's test for pairwise comparison between daily groups. Columns that are labeled with

different letters are statistically different. For panel B, statistical survival comparisons of

different groups were performed via log rank test.

Figure 3. *Time-dependent effect of estrous cycle on daily CDI severity.* Box-and-whisker plots of

daily CDI sign severity plotted against (A) their concurrent estrous stages and (B) their prior-day

estrous stages. Animals in estrus (blue bars), metestrus (orange bars), diestrus (green bars), late

diestrus/early proestrus (pink bars), proestrus (magenta bars), and late proestrus/early estrus

(yellow bars) were challenged with *C. difficile* spores from strain R20291 on day 0. CDI sign

severity and estrous cycle stages were determined daily. Signs of severity were determined based

on the CDI scoring rubric reported previously [36]. Horizontal bold lines represent median

values, asterisks represent mean values, bars represent interquartile ranges, error bars (whiskers)

represent maximum and minimum values, and dots represent inner point values. Single-factor

ANOVA was performed at every time point to assess differences among the sign severity means

of the different estrous stages. ANOVA results with *P* values of <0.05 were analyzed *post*

hoc using the Scheffe's test for pairwise comparison between daily groups. Columns that are

labeled with different letters are statistically different.

Figure 4. *Time-independent effect of estrous cycle on CDI severity.* Cumulative box-and-whisker

plots of CDI sign severity caused by *C. difficile* strain R20291 plotted against (A) their

concurrent estrous stages, and (B) their prior-day estrous stages. Animals that were at estrus

(blue bars), metestrus (orange bars), diestrus (green bars), late diestrus/early proestrus (pink

bars), proestrus (magenta bars), and late proestrus/early estrus (yellow bars) at any day were

pooled together. CDI sign severity and estrous cycle stages were determined daily. Signs of

severity were determined based on the CDI scoring rubric reported previously [36]. Horizontal

bold lines represent median values, asterisks represent mean values, bars represent interquartile

ranges, error bars (whiskers) represent maximum and minimum values, and dots represent inner

point values. Single-factor ANOVA was performed to assess differences among the sign severity

means of the different estrous stages. ANOVA results with P values of <0.05 were analyzed *post*

hoc using the Schiffer test for pairwise comparison between all groups. Columns that are labeled

with different letters are statistically different.

Figure 5. *CDI severity score and estrous stage 48 prior.* were scored daily for 5-7 days

following infection, and estrous stages were simultaneously tracked daily by microscopic

examination of cells collected by vaginal lavage. The maximum CDI score for each mouse was

determined and compared to its estrous stage 48 hours prior to the maximum score. Single-factor

ANOVA was performed to assess differences among the sign severity means of the different

estrous stages. ANOVA results with P values of <0.05 were analyzed *post hoc* using the Schiffer

test for pairwise comparison between all groups. Columns that are labeled with different letters

are statistically different.

Figure 6. *Biomarker networks before infection.* Serum was obtained from female mice before

challenge (D0) with *C. difficile* spores and individually analyzed for concentrations of sex

hormones, cytokines, and immunoglobulins. Animals were scored for CDI sign severity on days

1-5 post-challenge. Days with correlations between biomarkers and CDI severity are highlighted

(days 1 and 2 post-challenge). Biomarker levels and CDI scores were correlated using Kendall-

tau. Positive correlations are represented with green lines. Negative correlations are represented

with red lines. Line thickness and hues represent the strengths of the correlations. Solid lines

represent statistically significant correlations with $p \leq 0.05$. Dashed lines represent statistically

significant correlations with $0.051 \leq p \leq 0.10$.

Figure 7. *Biomarker networks during early infection.* Serum was obtained from female mice the

day after challenge (D1) with *C. difficile* spores and individually analyzed for concentrations of

sex hormones, cytokines, and immunoglobulins. Animals were scored for CDI sign severity on

days 1-5 post-challenge. Days with correlations between biomarkers and CDI severity are

highlighted (days 1 and 3 post-challenge). Biomarker levels and CDI scores were correlated

using Kendall-tau approaches. Positive correlations are represented with green lines. Negative

correlations are represented with red lines. Line thickness and hues represent the strengths of the

correlations. Solid lines represent statistically significant correlations with $p \leq 0.05$. Dashed lines

represent statistically significant correlations with $0.051 \leq p \leq 0.10$.

References

1.   Dicks LMT, Mikkelsen LS, Brandsborg E, Marcotte H. *Clostridium difficile*, the Difficult

"Kloster" Fuelled by Antibiotics. Current Microbiology. 2019;76(6):774–82; doi:

10.1007/s00284-018-1543-8.

2.   CDC: Antibiotic Resistance Threats in the United States 2019. In. Edited by Services

USDoHaH. Atlanta, GA; 2019.

3.   Zhang S, Palazuelos-Munoz S, Balsells EM, Nair H, Chit A, Kyaw MH. Cost of hospital

management of Clostridium difficile infection in United States—a meta-analysis and

modelling study. J BMC Infectious Diseases. 2016;16(1):447; doi: 10.1186/s12879-016-

1786-6.

4.   Ghose C. *Clostridium difficile* Infection in the Twenty-First Century. Emerging microbes

& Infections. 2013;2(9):e62–e; doi: 10.1038/emi.2013.62.

5.   Olsen MA, Young-Xu Y, Stwalley D, Kelly CP, Gerding DN, Saeed MJ, et al. The

Burden of *Clostridium difficile* Infection: Estimates of the Incidence of CDI from U.S.

Administrative Databases. BMC Infect Dis. 2016;16:177–; doi: 10.1186/s12879-016-

1501-7.

6.   Vázquez-Martínez ER, García-Gómez E, Camacho-Arroyo I, González-Pedrajo B.

Sexual dimorphism in bacterial infections. Biology of Sex Differences. 2018;9(1):27; doi:

10.1186/s13293-018-0187-5.

7.   Lucado J GCEA: Clostridium difficile Infections (CDI) in Hospital Stays, 2009. In.

Edited by (US) AfHRaQ. Rockville (MD); 2012.

8.   Elixhauser A SC, Gould C. : Readmissions following Hospitalizations with Clostridium

difficile Infections, 2009. In. Edited by (US) AfHRaQ. Rockville (MD); 2012.

9.   Saffouri G, Gupta A, Loftus EV, Baddour LM, Pardi DS, Khanna S. The incidence and

outcomes from Clostridium difficile infection in hospitalized adults with inflammatory

bowel disease. Scandinavian Journal of Gastroenterology. 2017;52(11):1240–7; doi:

10.1080/00365521.2017.1362466.

10.   Choi H-Y, Park S-Y, Kim Y-A, Yoon T-Y, Choi J-M, Choe B-K, et al. The epidemiology

and economic burden of Clostridium difficile infection in Korea. BioMed research

international. 2015;2015:510386–; doi: 10.1155/2015/510386.

11.   King A, Mullish BH, Williams HRT, Aylin P. Comparative epidemiology of Clostridium

difficile infection: England and the USA. International Journal for Quality in Health

Care. 2017;29(6):785–91; doi: 10.1093/intqhc/mzx120.

12.   Garey KW, Dao-Tran TK, Jiang ZD, Price MP, Gentry LO, DuPont HL. A clinical risk

index for Clostridium difficile infection in hospitalised patients receiving broad-spectrum

antibiotics. Journal of Hospital Infection. 2008;70(2):142–7; doi:

https://doi.org/10.1016/j.jhin.2008.06.026.

13.   Johnson S. Recurrent Clostridium difficile infection: A review of risk factors, treatments,

and outcomes. Journal of Infection. 2009;58(6):403–10; doi:

https://doi.org/10.1016/j.jinf.2009.03.010.

14.   Garey KW, Sethi S, Yadav Y, DuPont HL. Meta-analysis to assess risk factors for

recurrent Clostridium difficile infection. Journal of Hospital Infection. 2008;70(4):298–

304; doi: https://doi.org/10.1016/j.jhin.2008.08.012.

15.   Rouphael NG, O'Donnell JA, Bhatnagar J, Lewis F, Polgreen PM, Beekmann S, et al.

Clostridium difficile–associated diarrhea: an emerging threat to pregnant women.

American Journal of Obstetrics and Gynecology. 2008;198(6):635.e1–.e6; doi:

https://doi.org/10.1016/j.ajog.2008.01.062.

16.   Cózar-Llistó A, Ramos-Martinez A, Cobo J. Clostridium difficile Infection in Special

High-Risk Populations. Infectious diseases and therapy. 2016;5(3):253–69; doi:

10.1007/s40121-016-0124-z.

17.   Meda M, Virgincar N, Gentry V, Walker A, Macdonald N, Hooper M, et al. Clostridium

difficile infection in pregnant and postpartum women in 2 hospitals and a review of

literature. American Journal of Infection Control. 2019;47(1):e7–e14; doi:

https://doi.org/10.1016/j.ajic.2018.06.001.

18.   Garey KW, Jiang Z-D, Yadav Y, Mullins B, Wong K, Dupont HL. Peripartum

Clostridium difficile infection: case series and review of the literature. American Journal

of Obstetrics and Gynecology. 2008;199(4):332–7; doi:

https://doi.org/10.1016/j.ajog.2008.05.001.

19.   Draper CF, Duisters K, Weger B, Chakrabarti A, Harms AC, Brennan L, et al. Menstrual

cycle rhythmicity: metabolic patterns in healthy women. Sci Rep. 2018;8(1):14568; doi:

10.1038/s41598-018-32647-0.

20.   Comitato R, Saba A, Turrini A, Arganini C, Virgili F. Sex hormones and macronutrient

metabolism. Crit Rev Food Sci Nutr. 2015;55(2):227–41; doi:

10.1080/10408398.2011.651177.

21.   Taneja V. Sex Hormones Determine Immune Response. Front Immunol. 2018;9:1931–;

doi: 10.3389/fimmu.2018.01931.

22.   Klein SL. The effects of hormones on sex differences in infection: from genes to

718   behavior. Neuroscience & Biobehavioral Reviews. 2000;24(6):627–38; doi:

719   https://doi.org/10.1016/S0149-7634(00)00027-0.

720   23.   Shah NM, Lai PF, Imami N, Johnson MR. Progesterone-Related Immune Modulation of

721   Pregnancy and Labor. Frontiers in Endocrinology. 2019;10(198); doi:

722   10.3389/fendo.2019.00198.

723   24.   Mayneris-Perxachs J, Arnoriaga-Rodríguez M, Luque-Córdoba D, Priego-Capote F,

724   Pérez-Brocal V, Moya A, et al. Gut microbiota steroid sexual dimorphism and its impact

725   on gonadal steroids: influences of obesity and menopausal status. Microbiome.

726   2020;8(1):136; doi: 10.1186/s40168-020-00913-x.

727   25.   Colldén H, Landin A, Wallenius V, Elebring E, Fändriks L, Nilsson ME, et al. The gut

728   microbiota is a major regulator of androgen metabolism in intestinal contents. American

729   Journal of Physiology-Endocrinology and Metabolism. 2019;317(6):E1182–E92; doi:

730   10.1152/ajpendo.00338.2019.

731   26.   Chen KL, Madak-Erdogan Z. Estrogen and Microbiota Crosstalk: Should We Pay

732   Attention? Trends in Endocrinology & Metabolism. 2016;27(11):752–5; doi:

733   https://doi.org/10.1016/j.tem.2016.08.001.

734   27.   Shin J-H, Park Y-H, Sim M, Kim S-A, Joung H, Shin D-M. Serum level of sex steroid

735   hormone is associated with diversity and profiles of human gut microbiome. Research in

736   Microbiology. 2019; doi: https://doi.org/10.1016/j.resmic.2019.03.003.

737   28.   Byers SL, Wiles MV, Dunn SL, Taft RA. Mouse Estrous Cycle Identification Tool and

738   Images. PLOS ONE. 2012;7(4):e35538; doi: 10.1371/journal.pone.0035538.

739   29.   McLean AC, Valenzuela N, Fai S, Bennett SAL. Performing Vaginal Lavage, Crystal

740   Violet Staining, and Vaginal Cytological Evaluation for Mouse Estrous Cycle Staging

741   Identification. JoVE. 2012;(67):e4389; doi: doi:10.3791/4389.

742   30.   Nilsson ME, Vandenput L, Tivesten Å, Norlén AK, Lagerquist MK, Windahl SH, et al.

743   Measurement of a Comprehensive Sex Steroid Profile in Rodent Serum by High-

744   Sensitive Gas Chromatography-Tandem Mass Spectrometry. Endocrinology.

745   2015;156(7):2492–502; doi: 10.1210/en.2014-1890.

746   31.   Liggins M, Ramirez N, Magnuson N, Abel-Santos E. Progesterone analogs influence

747   germination of *Clostridium sordellii* and *Clostridium difficile* spores *in vitro*. Journal of

748   Bacteriology. 2011;193(11):2776–83; doi: 10.1128/jb.00058-11.

749   32.   Adlercreutz H, Martin F. Biliary excretion and intestinal metabolism of progesterone and

750   estrogens in man. Journal of Steroid Biochemistry. 1980;13(2):231–44; doi:

751   https://doi.org/10.1016/0022-4731(80)90196-X.

752   33.   Phan JR, Do DM, Truong MC, Ngo C, Phan JH, Sharma SK, et al. An Aniline-

753   Substituted Bile Salt Analog Protects both Mice and Hamsters from Multiple

754   *Clostridioides difficile* Strains. Antimicrobial Agents and Chemotherapy.

755   2022;66(1):e01435–21; doi: doi:10.1128/AAC.01435-21.

756   34.   Yip C, Okada NC, Howerton A, Amei A, Abel-Santos E. Pharmacokinetics of Camsa, a

757   Potential Prophylactic Compound Against *Clostridioides difficile* Infections. Biochemical

758   Pharmacology. 2021;183:114314; doi: https://doi.org/10.1016/j.bcp.2020.114314.

759   35.   Sharma SK, Yip C, Simon MP, Phan J, Abel-Santos E, Firestine SM. Studies on the

760   Importance of the 7α-, and 12α- hydroxyl groups of N-Aryl-3α,7α,12α-trihydroxy-5β-

761   cholan-24-amides on their Antigermination Activity Against a Hypervirulent Strain of

762   *Clostridioides (Clostridium) difficile*. Bioorganic & Medicinal Chemistry.

763   2021;52:116503; doi: https://doi.org/10.1016/j.bmc.2021.116503.

764   36.   Mefferd CC, Bhute SS, Phan JR, Villarama JV, Do DM, Alarcia S, et al. A High-

765   Fat/High-Protein, Atkins-Type Diet Exacerbates *Clostridioides (Clostridium) difficile*

766   Infection in Mice, whereas a High-Carbohydrate Diet Protects. mSystems.

767   2020;5(1):e00765–19; doi: 10.1128/mSystems.00765-19.

768   37.   Yip C, Hassan C, Phan J, Abel-Santos E. Modulators of Bacterial Spore Germination. In:

769   Mora-Escobedo R, Martinez C, Reynoso R, editors. Germination: Types, Process and

770   Effects. New York, NY: Nova Science Publisher; 2019.

771   38.   Howerton A, Seymour CO, Murugapiran SK, Liao Z, Phan JR, Estrada A, et al. Effect of

772   the Synthetic Bile Salt Analog CamSA on the Hamster Model of *Clostridium difficile*

773   Infection. Antimicrobial Agents and Chemotherapy. 2018;62(10): e02251–17.

774   39.   Yip C, Phan J, Abel-Santos E. Treatment of *Clostridium difficile* Infections. In: Firestine

775   SM, Lister T, editors. Antibiotic Drug Discovery: New Targets and Molecular Entities.

776   London, UK: The Royal Society of Chemistry; 2017. p. 1–19

777   40.   Howerton A, Patra M, Abel-Santos E. Fate of Ingested *Clostridium difficile* Spores in

778   Mice. PloS one. 2013;8(8):e72620–e; doi: 10.1371/journal.pone.0072620.

779   41.   Howerton A, Patra M, Abel-Santos E. A New Strategy for the Prevention of *Clostridium*

780   *difficile* Infection. Journal of Infectious Diseases. 2013;207(10):1498–504; doi:

781   10.1093/infdis/jit068.

782   42.   Chen X, Katchar K, Goldsmith JD, Nanthakumar N, Cheknis A, Gerding DN, et al. A

783   Mouse Model of Clostridium difficile-Associated Disease. Gastroenterology.

784   2008;135(6):1984–92.

785   43.   Sun X, Wang H, Zhang Y, Chen K, Davis B, Feng H. Mouse Relapse Model of

786   Clostridium difficile Infection. Infection and Immunity. 2011;79(7):2856–64; doi:

787   10.1128/iai.01336-10.

788   44.   Mefferd CC, Bhute SS, Phan JR, Villarama JV, Do DM, Alarcia S, et al. A High-

789   Fat/High-Protein, Atkins-Type Diet Exacerbates Clostridioides (Clostridium) difficile

790   Infection in Mice, whereas a High-Carbohydrate Diet Protects. mSystems. 2020;5(1);

791   doi: 10.1128/mSystems.00765-19.

792   45.   Cora MC, Kooistra L, Travlos G. Vaginal Cytology of the Laboratory Rat and Mouse:

793   Review and Criteria for the Staging of the Estrous Cycle Using Stained Vaginal Smears.

794   Toxicol Pathol. 2015;43(6):776–93; doi: 10.1177/0192623315570339.

795   46.   Phan JR, Do DM, Truong MC, Ngo C, Phan JH, Sharma SK, et al. An Aniline-

796   Substituted Bile Salt Analog Protects both Mice and Hamsters from Multiple

797   Clostridioides difficile Strains. Antimicrob Agents Chemother. 2022;66(1):e0143521;

798   doi: 10.1128/aac.01435-21.

799   47.   Watson T, Hickok J, Fraker S, Korwek K, Poland RE, Septimus E. Evaluating the Risk

800   Factors for Hospital-Onset Clostridium difficile Infections in a Large Healthcare System.

801   Clin Infect Dis. 2018;66(12):1957–9; doi: 10.1093/cid/cix1112.

802   48.   Dannheim H, Riedel T, Neumann-Schaal M, Bunk B, Schober I, Spröer C, et al. Manual

803   curation and reannotation of the genomes of Clostridium difficile 630Δerm and C.

804   difficile 630. Journal of Medical Microbiology. 2017;66(3):286–93; doi:

805   https://doi.org/10.1099/jmm.0.000427.

806   49.   Sebaihia M, Wren BW, Mullany P, Fairweather NF, Minton N, Stabler R, et al. The

807   Multidrug-Resistant Human Pathogen *Clostridium difficile* Has A Highly Mobile, Mosaic

808   Genome. Nature Genetics. 2006;38(7):779–86.

809   50.   Bhute SS, Mefferd CC, Phan JR, Ahmed M, Fox-King AE, Alarcia S, et al. A High-

810   Carbohydrate Diet Prolongs Dysbiosis and Clostridioides difficile Carriage and Increases

811   Delayed Mortality in a Hamster Model of Infection. Microbiol Spectr. 2022:e0180421;

812   doi: 10.1128/spectrum.01804-21.

813   51.   Ajayi AF, Akhigbe RE. Staging of the estrous cycle and induction of estrus in

814   experimental rodents: an update. Fertil Res Pract. 2020;6:5; doi: 10.1186/s40738-020-

815   00074-3.

816   52.   Warn P, Thommes P, Sattar A, Corbett D, Flattery A, Zhang Z, et al. Disease Progression

817   and Resolution in Rodent Models of Clostridium difficile Infection and Impact of

818   Antitoxin Antibodies and Vancomycin. 2016;60(11):6471–82; doi: 10.1128/AAC.00974-

819   16 %J Antimicrobial Agents and Chemotherapy.

820   53.   Hutton ML, Mackin KE, Chakravorty A, Lyras D. Small animal models for the study of

821   Clostridium difficile disease pathogenesis. FEMS Microbiology Letters.

822   2014;352(2):140–9; doi: 10.1111/1574-6968.12367.

823   54.   Thiele K, Hierweger AM, Riquelme JIA, Solano ME, Lydon JP, Arck PC. Impaired

824   Progesterone-Responsiveness of CD11c(+) Dendritic Cells Affects the Generation of

825   CD4(+) Regulatory T Cells and Is Associated With Intrauterine Growth Restriction in

826   Mice. Front Endocrinol (Lausanne). 2019;10:96; doi: 10.3389/fendo.2019.00096.

827   55.   Piccinni MP, Giudizi MG, Biagiotti R, Beloni L, Giannarini L, Sampognaro S, et al.

828   Progesterone favors the development of human T helper cells producing Th2-type

cytokines and promotes both IL-4 production and membrane CD30 expression in

established Th1 cell clones. J Immunol. 1995;155(1):128–33.

56.   Raghupathy R, Szekeres-Bartho J. Progesterone: A Unique Hormone with

Immunomodulatory Roles in Pregnancy. Int J Mol Sci. 2022;23(3); doi:

10.3390/ijms23031333.

57.   Hugon AM, Golos TG. Listeria monocytogenes infection in intestinal epithelial Caco-2

cells with exposure to progesterone and estradiol-17beta. PLoS One.

2025;20(3):e0320631; doi: 10.1371/journal.pone.0320631.

58.   Borba VV, Zandman-Goddard G, Shoenfeld Y. Prolactin and Autoimmunity. Front

Immunol. 2018;9:73; doi: 10.3389/fimmu.2018.00073.

59.   Wu X, Lai X, Tu H, Zou H, Cao J. Elevated serum CXCL10 in patients with Clostridium

difficile infection are associated with disease severity. Int Immunopharmacol.

2019;72:92–7; doi: 10.1016/j.intimp.2019.03.033.

60.   Collins MK, McCutcheon CR, Petroff MG. Impact of Estrogen and Progesterone on

Immune Cells and Host-Pathogen Interactions in the Lower Female Reproductive Tract. J

Immunol. 2022;209(8):1437–49; doi: 10.4049/jimmunol.2200454.

61.   Galozzi P, Bindoli S, Doria A, Sfriso P. The revisited role of interleukin-1 alpha and beta

in autoimmune and inflammatory disorders and in comorbidities. Autoimmun Rev.

2021;20(4):102785; doi: 10.1016/j.autrev.2021.102785.

62.   Bajrami B, Zhu H, Kwak HJ, Mondal S, Hou Q, Geng G, et al. G-CSF maintains

controlled neutrophil mobilization during acute inflammation by negatively regulating

CXCR2 signaling. J Exp Med. 2016;213(10):1999–2018; doi: 10.1084/jem.20160393.

63.   Ajuebor MN, Hogaboam CM, Kunkel SL, Proudfoot AE, Wallace JL. The chemokine

852   RANTES is a crucial mediator of the progression from acute to chronic colitis in the rat. J

853   Immunol. 2001;166(1):552–8; doi: 10.4049/jimmunol.166.1.552.

854   64.   Jose S, Madan R. Neutrophil-mediated inflammation in the pathogenesis of Clostridium

855   difficile infections. Anaerobe. 2016;41:85–90; doi: 10.1016/j.anaerobe.2016.04.001.

856   65.   Morteau O, Castagliuolo I, Mykoniatis A, Zacks J, Wlk M, Lu B, et al. Genetic

857   deficiency in the chemokine receptor CCR1 protects against acute Clostridium difficile

858   toxin A enteritis in mice. Gastroenterology. 2002;122(3):725–33; doi:

859   10.1053/gast.2002.31873.

860   66.   Hammar O, Veress B, Montgomery A, Ohlsson B. Expression of Luteinizing Hormone

861   Receptor in the Gastrointestinal Tract in Patients with and without Dysmotility. Drug

862   Target Insights. 2012;6:13–8; doi: 10.33393/dti.2012.2403.

863   67.   Cisoń-Jurek S, Czajka-Francuz P, Francuz T, Owczarek AJ, Szymczak B, Wojnar J, et al.

864   The Value of Prolactin, a Panel of Cytokines, and the Soluble Human Epidermal Growth

865   Factor Receptor 2 in the Prediction of Rapid Progression and Shorter Survival during

866   Palliative Chemotherapy of Colorectal Cancer Patients. Biomedicines. 2023;11(7); doi:

867   10.3390/biomedicines11072014.

868   68.   Gouon-Evans V, Pollard JW. Eotaxin is required for eosinophil homing into the stroma

869   of the pubertal and cycling uterus. Endocrinology. 2001;142(10):4515–21; doi:

870   10.1210/endo.142.10.8459.

871   69.   Rao K, Erb-Downward JR, Walk ST, Micic D, Falkowski N, Santhosh K, et al. The

872   systemic inflammatory response to Clostridium difficile infection. PLoS One.

873   2014;9(3):e92578; doi: 10.1371/journal.pone.0092578.

874   70.   Pereira Suarez AL, López-Rincón G, Martínez Neri PA, Estrada-Chávez C. Prolactin in

875   inflammatory response. Adv Exp Med Biol. 2015;846:243–64; doi: 10.1007/978-3-319-

876   12114-7_11.

877   71.   Narros-Fernández P, Chomanahalli Basavarajappa S, Walsh PT. Interleukin-1 family

878   cytokines at the crossroads of microbiome regulation in barrier health and disease. Febs j.

879   2024;291(9):1849–69; doi: 10.1111/febs.16888.

880   72.   Mao L, Wang L, Bennett S, Xu J, Zou J. Effects of follicle-stimulating hormone on fat

881   metabolism and cognitive impairment in women during menopause. Front Physiol.

882   2022;13:1043237; doi: 10.3389/fphys.2022.1043237.

883   73.   Iqbal J, Sun L, Kumar TR, Blair HC, Zaidi M. Follicle-stimulating hormone stimulates

884   TNF production from immune cells to enhance osteoblast and osteoclast formation. Proc

885   Natl Acad Sci U S A. 2006;103(40):14925–30; doi: 10.1073/pnas.0606805103.

886   74.   Crawford ED, Schally AV, Pinthus JH, Block NL, Rick FG, Garnick MB, et al. The

887   potential role of follicle-stimulating hormone in the cardiovascular, metabolic, skeletal,

888   and cognitive effects associated with androgen deprivation therapy. Urol Oncol.

889   2017;35(5):183–91; doi: 10.1016/j.urolonc.2017.01.025.