Regulation of potassium homeostasis in *Caulobacter crescentus*

Alex Quintero-Yanes[1+], Madeline Collignon[1], Loïc Léger[1], Aurélie Mayard[1], Régis Hallez[1,2,$,#]

[1]Bacterial Cell cycle & Development (BCcD), Biology of Microorganisms Research Unit (URBM), Namur Research Institute for Life Science (NARILIS), University of Namur, Namur (5000), Belgium.

[2]WELBIO, University of Namur, Namur (5000), Belgium

[+] ORCID ID: 0000-0002-8350-0881

[$] ORCID ID: 0000-0003-1175-8565

[#]Corresponding author: Tel: +32 81 724 244; E-mail: regis.hallez@unamur.be

Short title: Potassium homeostasis in *C. crescentus*.

Key words: Potassium transport, two-component system, Kup, KdpE, KdpD

Abstract

Potassium (K^+) is an essential physiological element determining membrane potential, intracellular pH, osmotic/turgor pressure, and protein synthesis in cells. Nevertheless, K^+ homeostasis remains poorly studied in bacteria. Here we describe the regulation of potassium uptake systems in the oligotrophic α-proteobacterium *Caulobacter crescentus* known as a model for asymmetric cell division. We show that *C. crescentus* can grow in concentrations from the micromolar to the millimolar range by essentially using two K^+ transporters to maintain potassium homeostasis, the low affinity Kup and the high affinity Kdp uptake systems. When K^+ is not limiting, we found that the *kup* gene is essential while *kdp* inactivation does not impact the growth. In contrast, *kdp* becomes critical but not essential and *kup* dispensable for growth in K^+-limited environments. However, in the absence of *kdp*, mutations in *kup* were selected to improve growth in K^+-depleted conditions, likely by improving the affinity of Kup for K^+. In addition, mutations in the KdpDE two-component system, which regulates *kdpABC* expression, suggest that the inner membrane sensor regulatory component KdpD works as a kinase in early stages of growth and as a phosphatase to regulate transition into stationary phase. Our data also show that KdpE is not only phosphorylated by KdpD but also by another non-cognate histidine kinase. On top of this, we determined the KdpE-dependent and independent K^+ transcriptome as well as the direct targets of KdpE. Together, our work illustrates how an oligotrophic bacterium responds to fluctuation in K^+ availability.

Introduction

Potassium (K$^+$) is the most abundant monovalent cation within living cells (Danchin & Nikel, 2019, Epstein, 2003) and it is essential to set the membrane potential and intracellular pH (Bakker & Mangerich, 1981, Booth, 1999, Epstein, 2003, Ochrombel *et al.*, 2011). As K$^+$ also emerges as an important regulator of bacterial physiology in view of its role in diverse processes (Castaneda-Garcia *et al.*, 2011, Dominguez-Ferreras *et al.*, 2009, Feng *et al.*, 2022, Gries *et al.*, 2013, Humphries *et al.*, 2017, Liu *et al.*, 2013, MacGilvary *et al.*, 2019, Prindle *et al.*, 2015, Quintero-Yanes *et al.*, 2019, Rasmussen, 2023, Su *et al.*, 2009, Valente & Xavier, 2016) including biofilm formation, chemotaxis, cell to cell communication or virulence, it is therefore expected that K$^+$ transport is tightly regulated.

In bacteria, the KdpDE two-component system (TCS) responds to potassium depletion by regulating the expression of the cognate high-affinity K$^+$ transporter KdpFABC (Pedersen *et al.*, 2019). This inner membrane complex consists of four subunits all encoded in the same operon, often together with the *kdpDE* genes (Ali *et al.*, 2017, Dani *et al.*, 2017). In *Escherichia coli*, KdpD was shown to act as a dual sensor for K$^+$, which independently regulates both enzymatic activities. Whereas the kinase activity was inhibited by extracellular K$^+$, phosphatase activity was stimulated by intracellular K$^+$ (Schramke *et al.*, 2016). This ultimately determines the phosphorylation status of the response regulator KdpE and thus provides a tight regulation of *kdpFABC* expression and as well as a robust homeostasis in fluctuating environments (Schramke *et al.*, 2016).

Other K$^+$ transporters have been described in detail in Gram-negative bacteria, such as the inner membrane low-affinity Trk and Kup systems for uptake, and the KefB efflux

pump. The Trk potassium uptake system is highly conserved in bacteria and works in

a multi-subunit complex comprising the TrkG/H, TrkA and TrkE proteins. TrkH (an extra

copy known as TrkG is found in *E. coli*) is the trans-membrane transporter unit, which

interacts with the NAD^+- and NADH-binding peripheral protein TrkA and the ATP-

binding protein TrkE in the cytoplasmic leaflet to control extracellular potassium uptake

when high extracellular concentrations of K^+ are available (Bossemeyer *et al.*, 1989a,

Dosch *et al.*, 1991, Harms *et al.*, 2001). On the other hand, Kup is a single protein,

constitutively expressed and thought to function as K^+/Na^+ symporter (Dosch *et al.*,

1991, Schleyer & Bakker, 1993, Zakharyan & Trchounian, 2001).

Recently, it was reported that glutathione controls cell division via negative regulation

of KefB in the Gram-negative oligotrophic α-proteobacterium *Caulobacter crescentus*

(3). Indeed, mutants unable to synthesize glutathione led to cell filamentation and

decreased intracellular K^+ levels. Interestingly, these phenotypes were suppressed by

mutations in the KefB efflux pump, suggesting that K^+ homeostasis is crucial for

bacterial cell cycle regulation. *C. crescentus* has been extensively used as a model for

cell cycle studies since it divides asymmetrically to give birth to a larger sessile stalked

cell and a smaller flagellated swarmer cell (1). DNA replication and cell division take

place only in sessile stalked cells whereas the swarmer cells remain in a non-

replicative but chemotactically active and motile phase able to explore new

environments. Once swarmer cells find a suitable – resourceful – habitat for

reproduction, they differentiate into stalked cells and concomitantly start DNA

replication. In contrast, upon deprivation of environmental resources, the swarmer cells

do not differentiate into replicative stalked cells.

In this study, we characterized the response of *Caulobacter* cells grown in environments containing what we defined as limiting, abundant and excessive K^+ concentrations. *In silico* analyses revealed that *C. crescentus* does not have Trk system components but encodes a putative transporter with a cytoplasmic TrkA-like domain. Moreover, in addition to the KefB and KefC efflux pumps, we found genes predicted to code for a Kup and a KdpABC transporters as well as a KdpDE TCS. We generated mutants of these predicted K^+ transporters and assessed their fitness in limiting and abundant K^+ environments. We also characterized the KdpDE system and determined its regulon by ChIP-seq and RNA-seq.

Results

Growth of *C. crescentus* at different K^+ concentrations

To primarily assess the impact of K^+ on *C. crescentus* fitness, the wild-type (WT) strain was grown in K^+-free minimal media (M2G-K) supplemented with K^+ at different concentrations using either KCl or a combination of K_2HPO_4 and KH_2PO_4 (K_2HPO_4 + KH_2PO_4) like the one used in standard M2G minimal media (**Fig. 1**). First, when phosphate salts were used as K^+ source, we observed that compared to M2G, which contains 7 mM K^+, growth started to be significantly impaired at K^+ concentrations below or equal to 0.025 mM and above or equal to 7 mM (**Fig. 1AB**). Second, by using KCl, we found different K^+ concentrations critical for growth since the WT already failed to grow at 0.1 mM but supports growth up to 25 mM (**Fig. 1CD** and **Fig. S1A**). Notwithstanding these differences likely due to the PO_4^{3-} and Cl^- counter-anions, the growth of the WT at K^+ concentrations ranging from 0.5 mM to 5 mM was indistinguishable from the one in M2G, that is 7 mM (**Fig. 1**). Therefore, based on these

data, we will use in this study potassium phosphate salts at 0.025 mM and 0.5 mM

respectively as limiting and abundant concentrations while KCl will be used at 50 mM

as excessive concentration.

K$^+$ transport and regulation systems in *C. crescentus*

By using *in silico* analyses, we found in the genome of *C. crescentus* NA1000

(NC_011916.1) several genes encoding potential K$^+$ uptake (*kup* and *kdpABCDE*) or

efflux (*kefB*, *kefC* and *kefG*) systems (**Fig. 2**). Although Trk orthologs were not found,

a putative transporter (CCNA_01688) containing a cytoplasmic TrkA-like domain was

identified (**Fig. 2**). Based on the function of the Kdp, Kef, Kup and Trk systems in other

bacteria (Bossemeyer *et al.*, 1989b, Cao *et al.*, 2011, Epstein, 2003, MacGilvary *et al.*,

2019, Roosild *et al.*, 2010), we predicted that transport of K$^+$ in *C. crescentus* occurs

via CCNA_01688, KefBG, KefC and Kup when potassium is abundant in the

environment (**Fig. 2A**) whereas the Kdp system might be inactive (**Fig. 2B**) (Heermann

& Jung, 2010, Roe *et al.*, 2000). In contrast, K$^+$-depleted conditions should activate the

KdpDE TCS and trigger expression of the high affinity KdpABC transporter (**Fig. 2C**)

(Laimins *et al.*, 1981).

To test these predictions, we first constructed knock-out (KO) mutants for all the K$^+$

transport and regulatory systems described above. We successfully inactivated all

these genes except *kup*, suggesting it might be essential at least at the K$^+$

concentration found in the complex media PYE used to construct the KO strains. Note

that a Δ*kefB* and Δ*kdpABCDE* (hereafter referred to as Δ*kdp*) were respectively used

as KefBG and Kdp inactive mutants (**Fig. 2C**). Then, we measured the growth for the

mutants in M2G-K supplemented with limiting (0.025 mM), abundant (0.5 mM) or

excessive (50 mM) K$^+$ concentrations (**Fig. 3**). In limiting K$^+$ conditions, only Δ*kdp* had

a strong growth delay (**Fig. 3A**) whereas all the KO mutants grew as good as the WT,

both in minimal and complex media supplemented with abundant K$^+$ concentrations

(**Fig. 3B** and **Fig. S1B**). However, and as far as we know in contrast to what is

described in other bacteria, a *kdp* inactive mutant could still grow in such limiting

conditions, suggesting that another transporter is used in K$^+$-depletion conditions. At

excessive K$^+$ concentration, we found that (i) Δ*CCNA_01688* had a slight growth delay;

(ii) Δ*kdp* had a lower plateau and (iii) Δ*kefC* barely grew in comparison to WT (**Fig.**

3C). Interestingly, the growth delay of Δ*CCNA_01688* and the lower plateau of Δ*kdp*

were not observed in complex PYE media supplemented with an excess of K$^+$ whereas

Δ*kefC*, like in minimal medium, did not grow (**Fig. S1C**).

K$^+$-dependent essential genome

Transposon mutagenesis coupled to next-generation sequencing (Tn-seq) were used

to determine a core set of genes important for adaptation and response to abundant

or limiting K$^+$ condition. Isolated colonies harbouring transposon insertions were grown

on and collected from solid agar plates. We used M2G-K with 1 % agar concentration

to lower as much as possible K$^+$ traces from the agar in our experiment. Indeed, even

at 1% agar, the K$^+$ traces were sufficient to allow WT cells to grow without exogenous

source of K$^+$ (on M2G-K, **Fig. 4A**). In contrast, at least 0.025 mM K$^+$ had to be

supplemented to M2G-K to allow growth of Δ*kdp* while at 0.5 mM K$^+$, both the mutant

and the WT grew similarly to the growth observed on regular M2G which contains 7

mM K$^+$ (**Fig. 4A**). Therefore, we decided to use M2G-K without exogenous K$^+$ and M2G

plates respectively as the limiting and abundant K$^+$ conditions for the Tn-seq (**Fig. 4A**).

We did calculations on the ratio of transposon (Tn) insertions per base pair (# Tn

insertions / bp) for internal 80% of each ORF in each condition (M2G and M2G-K)

(**Table S1**). As in studies performed in other bacteria, we analysed the frequency of Tn

insertions and observed a bimodal distribution that allowed us visualizing two clusters

of genes in each condition, one with low density of Tn insertions (considered as

essential genes) and the other one with a higher density (considered as non-essential

genes) (**Figure S2A)**. Based on this frequency of Tn insertions, we used Ward's

clustering analysis (Curtis & Brun, 2014) and defined three fitness cost categories, that

is (i) essential, (ii) high fitness cost (HFC) and (iii) non-essential. For the M2G library,

genes with values of Tn insertions / bp $\leq$ 0.01333333 were considered as essential,

0.01333779 $\leq$ and $\leq$ 0.04063018 as HFC, and $\geq$ 0.04081632 as non-essential. For the

M2G-K library, genes values of $\leq$ 0.0047112 were considered as essential, 0.0047909

$\leq$ and $\leq$ 0.0211178 as HFC, and $\geq$ 0.0215792 as non-essential (**Figure 4B** and **Table**

S1). Among the 4186 annotated genes in *C. crescentus*, the percentage of essential

genes in both media (M2G and M2G-K) was similar (~12 %), whilst the percentage of

HFC genes was lower (~7 %) in M2G-K than in M2G (~13 %). Hence, the percentage

of non-essential genes was higher in M2G-K (~ 81 %) than in M2G-K (~75 %). The

shift from the HFC to the non-essential category in *C. crescentus* cells grown in M2G-

K could be explained by the reduction of osmotic stress in this medium. Indeed, we

showed recently that M2G generates a cell envelope stress for *C. crescentus* due to

the high content of monovalent cations (Quintero-Yanes *et al.*, 2022).

Genes with known or predicted biological functions were grouped according to the

classification of cluster of orthologues groups (COG) (**Fig S2B**). This allowed to

highlight biological functions that are enriched in abundant and limited K$^+$ conditions.

Overall, the COG enrichment in the three fitness categories (Essential– E, High Fitness

Cost– HFC and Non-Essential– N-E) was essentially similar in both M2G and M2G-K

media. The C group "energy production and conversion" genes was the most abundant

for all the fitness categories in M2G-K, as well as for E and N-E genes in M2G, while

for HFC it was the second most abundant, after the J group "translation, ribosomal

structure and biogenesis".

By comparing the gene pool in each category for each media, we identified a set of

537 genes that changed their fitness cost from M2G to M2G-K and vice versa (**Fig. 4C**

and **Table S2**), including 153 genes which became exclusively essential (28.5 %), 153

HFC (28.5 %) and 231 non-essential (43 %) in M2G-K. First, we observed that the TCS

genes *chvGI*, known to be HFC in M2G due to its response to hyperosmotic stress

(Quintero-Yanes *et al.*, 2022), became N-E in M2G-K, thereby supporting that M2G-K

is a hypo-osmotic environment compared to M2G. Second, all the *kdp* genes

categorised as N-E genes in M2G became essential in M2G-K except *kdpD* which

became HFC. In contrast, *kup* moved from the HFC category in M2G to N-E one in

M2G-K. This confirms the importance of the Kdp and Kup systems repectively in

limiting and abundant K$^+$ conditions. Considering the conservation of the *kdp* system

in bacteria, it is intriguing that *kdpD* – encoding the HK component – did not have the

same fitness cost in M2G-K than the genes coding for the transporter (*kdpABC*) and

the response regulator (*kdpE*) components. While this supports that KdpE likely

regulates, directly, the expression of *kdpABC* genes, it suggests that KdpD and KdpE

have different regulatory effects on growth in K$^+$-depleted environments. Finally, we

did not observed changes in fitness for the other predicted potassium transport

systems since all of them were categorised as N-E genes in both M2G and M2G-K,

further supporting that CCNA_01688, KefBG and KefC are not important for growth in

abundant and limiting K$^+$ conditions.

Kup becomes dispensable in limiting K$^+$ conditions

Tn-seq data suggest that inactivation of *kup* gene could be facilitated at limiting K$^+$

concentrations (**Fig. 4B**, **Tables S1** and **S2**). Kup has been described in different

bacteria as the major K$^+$ transporter, essential in hyperosmotic conditions using

sucrose as osmotic stressor (de Oliveira *et al.*, 2017, Trchounian & Kobayashi, 1999).

Moreover, in a previous study, we found that the TCS ChvGI in *C. crescentus* positively

controls the expression of *kup* in hyperosmotic conditions (Quintero-Yanes *et al.*,

2022). In agreement with this, we could not obtain a Δ*kup* strain by using a markerless,

SacB-dependent recombination approach on complex media (PYE) plates

supplemented with 3% sucrose for the counterselection. Since addition of sucrose

considerably increases the osmolality of the media (Hocking *et al.*, 2012), we aimed to

construct a Δ*kup* mutant by lowering osmolytes in the media, that is on M2G-Na-K

(lacking both Na$_2$HPO$_4$ and KH$_2$PO$_4$) plates supplemented with 0.5 % sucrose. In these

conditions, we successfully generated Δ*kup* mutant candidates that were then tested

for growth at different K$^+$ regimes. Interestingly, we observed that the Δ*kup* mutant

grew in limiting K$^+$ concentration (M2G-K supplemented with 0.025 mM K$^+$), although

at a slower growth rate but with a higher plateau compared to the WT (**Fig. 5A**). As

expected from the Tn-seq data, we found that Δ*kup* failed to grow when exposed to

abundant and excessive K$^+$ concentrations (**Fig. 5A**). We also confirmed that Kup is

essential in hypertonic conditions since the mutant did not grow in limiting K$^+$ conditions

when 3% sucrose were added. (**Fig. 5A**). Furthermore, we also observed that

overexpression of *kup* in WT is detrimental for growth in limiting K$^+$ conditions (**Fig.**

5B). Altogether, these data indicate that Kup allows potassium uptake in abundant and

excessive K$^+$ conditions but can be detrimental for unknown reasons at limiting K$^+$

concentrations.

While characterizing the growth of Δ*kdp* cells on M2G-K plates (**Fig. 4A**), we

inadvertently selected growth suppressors. These suppressors improved the growth of

Δ*kdp* in limiting K$^+$ conditions (**Fig. 5C**). Whole genome sequencing analyses of several

of these suppressors allowed the identification of three mutations in *kup*: *kup*$_{A87P}$,

kup$_{G253S}$ and *kup*$_{S456R}$. Altogether, these data indicate that Kup allows potassium

uptake in abundant and excessive K$^+$ conditions but unexpectedly also works at low –

limiting – K$^+$ concentrations, and that single point mutations in Kup likely improve its

affinity for K$^+$.

K$^+$- and KdpE-dependent regulons

To further characterise the response of *C. crescentus* to K$^+$-depleted conditions, we

performed RNA-seq on RNA samples extracted from cells grown in minimal media in

K$^+$ limiting or abundant conditions. In our experiment, we identified a total of 594 genes

– 385 upregulated and 209 downregulated – whose expression was impacted upon K$^+$

limitation with at least a 2-fold change (FC ≥ 2) and a false-discovery rate (FDR) lower

than 5% (P*adj* ≤ 0.05) (**Fig. 6A**). As expected, this experiment confirmed that the

expression of *kdp* genes was strongly induced in the K$^+$-depleted medium. Among the

other upregulated genes, we found genes coding for the phosphate starvation protein

PhoH (CCNA_02727), a xylose isomerase family protein (CCNA_01701), a TonB-

dependent receptor (TBDR, CCNA_01859), an AraC family transcription regulator

(CCNA_01858) and hypothetical proteins (CCNA_03313 and CCNA_03970) (**Fig. 6A**

and **Table S3A**). Among the top downregulated candidates, we found genes

expressing a small RNA (CCNA_R0158), a hemin receptor (CCNA_02277), a putative

transporter (CCNA_03022), a protein with a calcium binding EF Hand binding domain

(CCNA_02274) and the chaperons GroES (CCNA_00722) and GroEL (CCNA_00721)

(**Fig. 6A** and **Table S3B**).

As our results suggest that the Kdp system is important for growth upon K^+ depletion,

we also performed RNA-seq experiments with a $\Delta kdpE$ mutant grown in M2G-K

supplemented with limiting K^+ concentrations. Then, we compared the RNA enrichment

in $\Delta kdpE$ *vs* WT to determine which genes are affected by KdpE in a K^+-depleted

condition using FC $\geq$ 1.5 and $P_{adj} \leq 0.05$ as parameters. Compared to WT, we found

83 upregulated and 50 downregulated genes in $\Delta kdpE$ grown in the K^+-depleted media

(**Fig. 6B** and **Tables S4AB**). Among the 50 downregulated genes in $\Delta kdpE$ (**Table**

S4B), 72% (36/50) were also found in the regulon of the WT starved for K^+, 19

upregulated (including *kdp* genes; **Table S3A**) and 17 downregulated (**Table S3B**).

This suggests that KdpE is strictly required to activate the expression of the 19 genes

in a K^+-depleted environment. On the contrary, the 17 other genes that are repressed

in WT cells upon K^+ depletion are even further downregulated in the absence of KdpE,

suggesting that KdpE limits the repression of these genes. The expression of the last

14 genes was insensitive to K^+ concentration while downregulated in $\Delta kdpE$ (**Tables**

S4AB), suggesting that they are positively regulated by KdpE but independently of the

K^+ concentration. Among the 83 upregulated genes in $\Delta kdpE$ (**Table S4A**), only ~36%

(30/83) were also found in the regulon of the WT starved for K^+, 12 downregulated

(including *groEL* and *groES* genes; **Table S3B**) and 18 upregulated (**Table S3A**). This

indicates KdpE is required to repress the first 12 genes upon K^+ depletion while it limits

the activation of the other 18 genes in a K^+-depleted environment. The remaining 53

upregulated genes in $\Delta kdpE$ (**Tables S4AB**) were insensitive to K^+ availability,

suggesting that they are likely repressed by KdpE but independently of the K^+

concentration.

Surprisingly, the vast majority (528 genes; 88.9 %) of the 594 genes whose expression

changed in the WT grown in K^+ limiting conditions seems to be KdpE-independent,

suggesting that other regulators, yet to be discovered, are activated upon K^+ depletion.

An alternative explanation might be that $\Delta kdpE$ cells exposed to limiting K^+

concentration suffer from a general stress response, which ultimately leads to global

transcriptional changes. This could also explain why half of the 163 genes (67/133)

found in the $\Delta kdpE$ regulon were not identified in the WT regulon.

In order to unveil genes whose expression is directly regulated by KdpE, we identified

the DNA regions bound by KdpE using ChIP-seq. For that, we first constructed strains

in which *kdpE* was expressed under the xylose inducible promoter (P_{xylX}::*kdpE*) in a

$\Delta kdpE$ $\Delta xylX$ background. Inactivating *xylX* avoids xylose consumption by *C.*

crescentus as a carbon source thereby allowing constant expression of *kdpE* upon

induction. Expression of *kdpE* from P_{xylX} efficiently complemented the growth of both

strains in the presence of xylose since $\Delta kdpE$ $\Delta xylX$ P_{xylX}::*kdpE* cells grew similarly to

the WT or $\Delta xylX$ cells whereas $\Delta kdpDE$ $\Delta xylX$ P_{xylX}::*kdpE* cells grew at a higher plateau

(**Fig. S3A**) like $\Delta kdpD$ cells (see below). Moreover, using polyclonal anti-KdpE

antibodies, KdpE was easily detected by western blot in both strains although KdpE

protein levels were lower in $\Delta kdpDE$ $\Delta xylX$ than in $\Delta kdpE$ $\Delta xylX$ (**Fig. S3B**), suggesting

that KdpD has a global positive impact on KdpE levels. It is noteworthy that KdpE was

undetectable in WT strain, indicating that P$_{xylX}$::*kdpE* leads to KdpE overexpression

upon xylose induction.

A total of 110 and 146 DNA binding sites were respectively detected in the Δ*kdpE*

Δ*xylX* P$_{xylX}$::*kdpE* and Δ*kdpDE* Δ*xylX* P$_{xylX}$::*kdpE* strains grown in K$^+$ limiting conditions

(**Fig. 6C**; **Tables S5** and **S6**). About a third of the DNA regions bound by KdpE

identified in both strains – 40 out of 110 (36.4%) found in Δ*kdpE* Δ*xylX* P$_{xylX}$::*kdpE* and

46 out of 146 (31.5%) found Δ*kdpDE* Δ*xylX* P$_{xylX}$::*kdpE* – are located in the close vicinity

of genes whose expression showed a FC ≥ 1.5 in RNA-seq experiments (highlighted

in green in **Tables S5** and **S6**), supporting that these regions are indeed KdpE-binding

sites.

The highest numbers of reads were found in the *kdp* promoter region, not only in Δ*kdpE*

Δ*xylX* P$_{xylX}$::*kdpE* cells but more surprisingly also in Δ*kdpDE* Δ*xylX* P$_{xylX}$::*kdpE* cells.

Even more surprising is the fact that the peak for Δ*kdpDE* Δ*xylX* P$_{xylX}$::*kdpE* was higher

than in Δ*kdpE* Δ*xylX* P$_{xyl}$::*kdpE* (**Fig. 6C**). In fact, this is true for most of if not all the

regulated genes, but the difference was particularly important for the *kdp* promoter

(**Fig. 6D**), suggesting that KdpE could be hyperphosphorylated in the absence of KdpD

and therefore that KdpD mainly acts as a negative regulator of KdpE~P. Moreover, it

indicates that, although P$_{kdp}$ is the main target of KdpE, multiple other genes are

potential direct targets of KdpE. Analysis of the top 50 sequences identified in the ChIP-

seq with the Δ*kdpE* Δ*xylX* P$_{xylX}$::*kdpE* strain using MEME allowed the identification of

the conserved DNA motif TCGAMRACGCSMKC likely bound by KdpE (**Fig. 6E**). For

the peak in the vicinity of the *kdpABCDE* operon (top hit of the analysis), the KdpE

motif was located 113 bp upstream of the *kdpA* start codon, within the gene

CCNA_01662 (**Fig. 6E**).

C. crescentus **KdpD regulates KdpE both positively and negatively**

To further assess the role of the *C. crescentus* Kdp system in K^+-limiting conditions,

we first constructed KO mutants of the transport complex ($\Delta kdpABC$) and the TCS

($\Delta kdpDE$) genes. In comparison to the WT, both $\Delta kdpABC$ and $\Delta kdpDE$ mutants had

a growth delay at limiting but not at abundant K^+ concentration (**Fig. 7A** and **Fig. S4A**).

Moreover, both mutants grew similarly, suggesting that the *kdpABC* is not expressed

in the absence of the TCS, as described in other bacteria.

To study the HK *kdpD* and RR *kdpE* genes independently of each other, we

constructed single KO mutants, $\Delta kdpD$ and $\Delta kdpE$. Deletion of *kdpE* resulted in a

similar growth delay than the ones displayed by $\Delta kdpABC$, $\Delta kdpDE$ and $\Delta kdpABCDE$

(**Fig. 7A** and **Fig. S4B**). On the contrary, $\Delta kdpD$ behave differently since compared to

WT, it had a slight but reproducible growth delay in exponential phase while it later

reached a higher plateau in stationary phase (**Fig. 7A**). This peculiar growth behaviour

was complemented by expressing *kdpD* from the native *kdp* promoter ($P_{kdp}::kdpD$) on

a replicative and low-copy number plasmid in a $\Delta kdpD$ background (**Fig. 7B**).

Complementation of the $\Delta kdpE$ mutant was achieved by expressing *kdpE* from the

xylose inducible promoter ($P_{xylX}::kdpE$) either in a $\Delta kdpE$ or a $\Delta kdpDE$ background.

While the growth defect of $\Delta kdpE$ was fully complemented, even in the absence of the

xylose inducer, expressing back *kdpE* in $\Delta kdpDE$ cells led to the same growth

behaviour than the single $\Delta kdpD$ mutant, that is a slight growth delay in exponential

phase and a higher plateau in stationary phase (**Fig. 7C** and **Fig. S4C**). In addition,

the fact that $\Delta kdpE$ is epistatic on $\Delta kdpD$, since $\Delta kdpDE$ grew as poor as $\Delta kdpE$ or

$\Delta kdpABC$ (**Fig. 7A**), shows KdpE is entirely responsible for the $\Delta kdpD$ phenotypes.

Together, these data support that both components of the TCS are important for WT

growth in K$^+$-depleted environments and suggest that KdpD and KdpE have

antagonistic roles for growth in such conditions. Kdp regulatory proteins KdpD and

KdpE were further analysed by identifying domains and their organisation. Comparison

of Pfam features highlighted that the KdpD sequence of *C. crescentus* lacks the

cytoplasmic GAF domain known to be linked to the DHp domain in *E. coli* [103]. Also,

multiple sequence alignments enabled to locate the conserved catalytic –

phosphorylatable – residues in both the HK (KdpD) and the RR (KdpE), that is

respectively the histidine 670 (His670) and the aspartate 56 (Asp56). To better

understand the KdpDE regulatory mechanism, we constructed *kdpD and kdpE*

catalytic point mutants and assessed their impact on growth in abundant and limiting

K$^+$ conditions. In agreement with what we observed with the KO mutants, all the

catalytic mutants grew as good as the WT in abundant K$^+$ conditions (**Fig. S4D**). At

limiting K$^+$ concentrations however, the phospho-ablative mutant of KdpE (*kdpE$_{D56A}$*)

had a strong growth delay, similar to the one displayed by $\Delta kdpE$ (**Fig. 7D** and **Fig.**

S4E). In contrast, the phospho-mimetic mutant (*kdpE$_{D56E}$*) grew similarly to the WT up

to the late exponential phase, to finally reach a higher final OD comparable to the one

observed with $\Delta kdpD$ grown in same conditions (**Fig. 7D** and **Fig. S4E**). As expected,

a KdpD catalytically dead mutant (*kdpD$_{H670N}$*) phenocopied $\Delta kdpD$ in limited K$^+$ (**Fig.**

7D and **Fig. S4E**). Since *kdpE$_{D56A}$* cells had a much stronger growth delay than $\Delta kdpD$

or *kdpD$_{H670N}$*, it indicates that KdpE is still phosphorylated despite the absence of KdpD

kinase activity. Thus, KdpE is very likely phosphorylated by another – non-cognate –

HK. On the other hand, the fact that $kdpE_{D56E}$ cells grew slightly but reproducibly better

than $\Delta kdpD$ and $kdpD_{H670N}$ suggests that KdpD also phosphorylates KdpE, at least

during the early exponential phase of growth. Thus, KdpD might (i) phosphorylate

KdpE in early exponential phase, with at least another HK, and (ii) mainly act as a

phosphatase on KdpE~P in late exponential phase, likely to downregulate the

expression of kdp and restrict growth in K^+-limiting conditions.

To test this hypothesis, we constructed a KdpD kinase-deficient but phosphatase

active mutant (K^-/P^+) by replacing a conserved phenylalanine residue by a leucine

residue ($kdpD_{F832L}$) in a way reminiscent to what was described for another HK

encoding gene, $pleC_{F778L}$ (Matroule et $al.$, 2004). Interestingly, we found that unlike the

$kdpD_{H670N}$, $kdpD_{F832L}$ cells failed to reach a higher plateau in stationary phase (**Fig. 7D**

and **Fig. S4E**), strongly supporting our hypothesis that KdpD primarily works as a

phosphatase in late exponential phase. Moreover, the K^-/P^+ mutant $kdpD_{F832L}$ grew

better than $kdpE_{D56A}$ and $\Delta kdpE$ mutants in low K^+ conditions (**Fig. 7D** and **Fig. S4E**),

supporting that there is indeed at least another HK that phosphorylates KdpE. Finally,

the fact that the $kdpD_{F832L}$ mutant had a growth delay compared to WT suggests that

KdpD also acts as a kinase for KdpE in early exponential phase (**Fig. 7D** and **Fig.**

S4E). However, we cannot rule out the possibility that the phosphatase activity of

$KdpD_{F832L}$ is impacted by the mutation. For instance, it might be stronger than the WT

thereby interfering with the phosphorylation levels of KdpE in early exponential phase.

Nevertheless, our results altogether indicate that (i) KdpD works both as a kinase and

phosphatase to control KdpE phosphorylation (KdpE~P) levels depending on the

growth phase and K^+ availability and that (ii) KdpE can be phosphorylated by (an)other

non-cognate HK(s) for regulating K^+ transport in K^+-depleted environments.

KdpD-dependent activation of the *kdp* promoter in low K⁺ conditions

In order to study the Kdp system at the transcriptional level, a plasmid carrying a P$_{kdp}$::*lacZ* reporter fusion was introduced in the different mutant strains, cells were grown in limiting and abundant K⁺ concentrations and β-galactosidase activity was measured. Consistent with the idea that the Kdp system is a high-affinity potassium transporter, P$_{kdp}$ basal activity in WT cells was barely detectable (~300 Miller Units) at abundant K⁺ concentration while it was strongly induced in WT (~9.000 Miller Units) or in Δ*kdpABC* cells (~10.500 Miller Units) upon K⁺ depletion (**Fig. 8A**). In contrast, P$_{kdp}$ activity was completely annihilated in Δ*kdpE* and *kdpE$_{D56A}$* cells (**Fig. 8A**), indicating that phosphorylated KdpE (KdpE~P) is strictly required for *kdp* expression both in limiting and abundant K⁺ conditions. In support of that, the basal activity of P$_{kdp}$ at abundant K⁺ concentration was 10 times higher (~3.000 Miller Units) in *kdpE$_{D56E}$* cells than in WT cells (**Fig. 8A**). Interestingly, P$_{kdp}$ was still active at low K⁺ concentration in a Δ*kdpD*, *kdpD$_{H670N}$,* and *kdpD$_{F832L}$* background (~4.600, ~4.200, ~4.700 Miller Units, respectively), but not to the same extent than in WT cells (~9.000 Miller Units) (**Fig. 8A**). In addition, the basal P$_{kdp}$ activity at abundant K⁺ concentration was completely abolished in the strains expressing the K⁻/P⁺ *kdpD$_{F832L}$* variant, suggesting that the KdpD phosphatase activity is counterbalanced by its kinase activity on KdpE in K⁺-replenished conditions. In support of that, compared to WT cells grown in abundat K⁺ condition, the basal P$_{kdp}$ activity was higher when both the hydrolase and kinase activities were inactivated (K⁻/P⁻), that is in Δ*kdpD* cells and to a lesser extent in *kdpD$_{F832L}$* cells.

Then, we followed the expression of the P_{kdp}::*lacZ* fusion in WT and *kdp* mutant cells

along the growth at limiting K$^+$ concentration. Despite $\Delta kdpABC$, $\Delta kdpE$ and $kdpE_{D56A}$

all poorly grew in limiting K$^+$ conditions, P_{kdp} was active in $\Delta kdpABC$ cells to a level

comparable to the WT one while P_{kdp} activity was almost undetectable in both *kdpE*

mutants (**Fig. 8B**). These data suggest that (i) the Kdp transporter (KdpABC) is not

required for *kdp* expression while (ii) the response regulator is, in contrast,

indispensable. In support of this, we found the phospho-mimetic mutant of *kdpE*

($kdpE_{D56E}$) displayed a relatively constant but high activity of P_{kdp} alongside the growth

curve (**Fig. 8B**).

The *kdpD* mutants had a different behaviour. First, compared to WT, the activity of P_{kdp}

was $\Delta kdpD$ cells was lower during early exponential phase but higher in stationary

phase (**Fig. 8C**). On the contrary, *kdp* expression was significantly lower both in

$kdpD_{H670N}$ (K$^-$/P$^-$) and in $kdpD_{F832L}$ (K$^-$/P$^+$) during early exponential phase. However,

P_{kdp} activity strongly increased only in $kdpD_{F832L}$ cells when reaching the late

exponential. Altogether, these results support our hypothesis that (i) KdpD plays dual

antagonistic activities on KdpE, as a kinase in early exponential phase and a

phosphatase in late exponential phase of growth, and that (ii) KdpE can be

phosphorylated by another non-cognate HK all along the growth.

Discussion

As the most abundant monovalent cation used by living cells to drive many cellular

processes, regulating potassium (K$^+$) homeostasis is critical for survival. The

oligotrophic α-proteobacterium *Caulobacter crescentus*, widely used as a cell cycle

model, is cyclically subjected to nutrients deprivation in its natural environments. In

contrast to carbon, nitrogen or phosphate, the response of *C. crescentus* to K⁺

depletion has never been characterised. Here, we first determined the range of K⁺

concentrations at which *C. crescentus* supports growth. We also identified all the genes

predicted to code for proteins involved the transport, the sensing and the regulation of

K⁺ homeostasis and tested their growth behaviour at low (limiting), optimal (abundant)

and high (excessive) K⁺ concentrations. Then, we defined the set of genes that (i)

become essential and/or (ii) are induced upon K⁺-environmental depletion. This

allowed us to identify the low affinity and the high affinity K⁺ transporters, respectively

Kup and KdpABC, as critical components that maintain K⁺ homeostasis in *C.*

crescentus. Finally, we deeply characterized the role played by the two-component

(TCS) regulatory system KdpDE in the response to K⁺ depletion.

By analysing the Tn-Seq data, we found that the COG categories – biological functions

– for which the highest number of genes became essential (EG) or high fitness cost

(HFCG) upon K⁺ limitation (M2G *vs* M2G-K) were (i) C "energy production and

conversion" (~1% increase of EG and ~6 % increase of HFCG) and (ii) J "Translation,

ribosomal structure and biogenesis" (~2% increase of EG and ~3 % increase of HFCG)

(**Table S2**). In fact, it is not so surprising to highlight these two categories as sensitive

to K⁺ availability. Indeed, ATP concentration and respiration state are two well-known

parameters that drive K⁺ transport across the membrane. For instance, the electron

transport chain determines the membrane potential, which in turn will influence the

distribution of other ions across like K⁺ that accumulate on the cytoplasmic face of the

membrane. This might therefore explain why Tn-insertions in genes coding for subunits

of the respiratory chain or the ATP synthase are counter-selected when K⁺ become

limiting. On top of this, active transport of K^+ across the membrane by the high-affinity

transporter consumes a lot of energy provided by ATP, so that decreasing ATP

concentration likely has a negative impact on K^+ import. On the other hand, K^+ cations

are critical for maintaining the structure and sustaining the function of ribosomes. Not

only K^+ is required to stabilize the messenger RNA, the transfer RNAs, the ribosomal

RNAs and the ribosomal proteins within the ribosomes, but also to increase

proofreading by accommodating the right aminoacyl-tRNA in the ribosomal A-site

(Grason *et al.*, 2008, Rozov *et al.*, 2019) . Additionally, many GTPases involved in

translation display a K^+-dependent activation of GTP hydrolysis (Danchin & Nikel,

2019).Thus, the impact K^+ has on translation could explain the lower Tn insertion

density observed in rRNA, tRNA and genes coding for ribosomal proteins in cells grown

in limiting on K^+ conditions.

Several physiological processes have been described to be K^+-dependent. Besides

being required by numerous enzymes related to basic metabolic functions (Danchin &

Nikel, 2019, Gohara & Di Cera, 2016), such as for example the pyruvate kinase

(Epstein, 2003), K^+ modulates the activity of enzymes involved in bacterial cell cycle.

For instance, in *E. coli*, K^+ has been shown to stimulate the polymerisation of the cell

division protein FtsZ (Ahijado-Guzman *et al.*, 2013), the activity of the DNA polymerase

(Klenow & Henningsen, 1969). Moreover, the DNA supercoiling activity of the DNA

gyrase was described to be triggered by K^+ in both *B. subtilis* (Gubaev & Klostermeier,

2012) and *Micrococcus luteus* (Liu & Wang, 1978). It was also described as critical for

the chaperonin activity of GroEL/GroES (Grason *et al.*, 2008) or DnaK (Feifel *et al.*,

1996). Interestingly, we observed that *groEL* and *groES* became essential upon K^+

depletion, thereby suggesting that some K+-dependent mechanisms are conserved in
bacteria.

KdpDE is the major, if not the only bacterial system described in the literature to
respond to limiting K+ availability. Although KdpE was shown in *E. coli* to regulate
expression of the anti-sigma factor Rsd and the ribosome modulation factor Rmf at low
K+ concentrations, thereby allowing KdpE to globally control transcription and
translation in limiting K+, other regulatory systems independent of KdpDE have not
been yet described in bacteria. The K+-dependent regulon of *C. crescentus* allowed us
to identify almost 600 genes whose expression changes upon K+ limitation.
Surprisingly, only a small proportion of this regulon depends directly or indirectly on
the response regulator KdpE, suggesting that there are unknown transcriptional
regulators sensitive to K+ availability.

The analysis of the *C. crescentus* genome allowed us to identify several genes and
systems predicted to transport K+. By systematically characterizing the growth profile
of the corresponding mutants in different K+ regimes, we found that the high affinity
Kdp system primarily serves to import K+ at limiting K+ concentrations, like in many
other bacterial species. Nonetheless, Kdp is unlikely the sole K+ transport system
involved in this context. Despite its growth defect, the Δ*kdpABC* mutant was still able
to grow at low K+ concentration. Since complete depletion of K+ led to a growth arrest
in *C. crescentus*, this implies that K+ is imported by additional system(s), at least in
limiting K+ conditions. Interestingly, we found that the low affinity Kup transporter
became dispensable in such limiting K+ conditions. However, mutations in Kup have
been identified to improve the growth the Δ*kdp* mutant on M2G-K plates (**Fig. 5D**),
strongly suggesting that Kup participates to K+ import when it is poorly available.

To test whether Kup is at work at low K^+ conditions, it might be interesting to try to

combine Δkdp and Δkup mutations together. Nonetheless, K^+ could be transported in

such limiting conditions by other unknown transporters. Several putative transporters

were identified as essential and high fitness cost in our Tn-seq analysis (i.e

CCNA_01159 and CCNA_00494) and up-regulated in the RNA-seq analysis (i.e

CCNA_00339, CCNA_01852, CCNA_01587, CCNA_01588 CCNA_02570,

CCNA_02571, …). It is possible that one or several of them assist the activity of

KdpABC in limiting K^+.

As in *E. coli*, the expression of *kdp* operon in *C. crescentus* is activated upon K^+

starvation by the KdpDE TCS (Polarek *et al.*, 1992). However, we found surprising

differences since in the absence of the histidine kinase (HK) KdpD, (i) *kdp* is still

expressed, although to a lesser extent than in WT cells and (ii) the *kdp* promoter (P_{kdp})

is more active in stationary phase. Our genetic dissection of the catalytic activity of

KdpDE with point mutants in *kdpE* (phospho-mimetic *kdpE$_{D56E}$* and phospho-ablative

kdpE$_{D56A}$) and KdpD (K^-/P^- *kdpD$_{H670N}$* and K^-/P^+ *kdpD$_{F832L}$*) strongly suggest that (i)

KdpE can be phosphorylated by another – non-cognate – HK, at least in the absence

of KdpD and that (ii) KdpD switches mainly to a phosphatase in late exponential phase

of growth.

In *E. coli,* deletion of the *kdpD* results in a prominent growth arrest, of more than 20

hours, before restarting growth thanks to KdpE phosphorylation by PhoR. However, it

never surpassed the WT plateau as seen with *C. crescentus $\Delta kdpD$, kdpD$_{H670N}$* and

kdpE$_{D56E}$ mutants. Thus, the higher plateau is due to the loss of KdpD phosphatase

activity. The KdpD phosphatase activity has be shown to be stimulated by intracellular

K^+ intracellular levels (Schramke *et al.*, 2016). Four amino acids located in the second

periplasmic loop were identified as an extracellular K^+-specific recognition site. Hence,

substitution of those four amino acids by alanine resulted in a strain unable to sense

extracellular K^+. In contrast, the K^+-recognition site for intracellular sensing remains to

be identified. However, although specific residues could not be highlighted, the C-

terminal cytoplasmic domain of KdpD was described to display a K^+-dependent

phosphatase activity (Cann, 2007, Dani *et al.*, 2017). Although this has yet to be

experimentally investigated, authors suggested that the cytoplasmic GAF domain of

KdpD may be responsible for intracellular K^+ sensing. Most GAF domains are involved

in regulatory functions through binding to cyclic nucleotides or other small ligands

including ions (Cann, 2007, Heikaus *et al.*, 2009). Interestingly, we found that KdpD

from *C. crescentus* lacks a GAF domain. Therefore, it is possible that such a

connection between intracellular sensing and phosphatase activity does not exist in *C.*

crescentus, which would explain why KdpD mainly acts as a phosphatase.

As mentioned above, in *E. coli*, the HK PhoR can crosstalk with the RR KdpE, which

allows a $\Delta kdpD$ mutant to grow in a low K^+ environment after a long lag phase

(Schramke *et al.*, 2017) . Since PhoR is conserved in *C. crescentus*, it constitutes a

good candidate for the non-cognate HK phosphorylating KdpE. To test this hypothesis,

we could measure growth behaviour as well as P$_{kdp}$ activity in a strain in which both

kdpD and *phoR* were inactivated. PhoR is a sensor kinase regulating the

phosphorylation levels and activity of the RR PhoB, which drives the Pho regulon, that

is the entire set of genes whose expression relies on phosphate availability.

Interestingly, a crosstalk between KdpD and PhoB was also described, which means

that in the absence of PhoR, PhoB can be phosphorylated by KdpD in low phosphate

conditions (Schramke *et al.*, 2017) . The double crosstalk suggests there is tight

connection between K$^+$ and phosphate homeostasis. Since both TCS – KdpDE and

PhoBR – are conserved in *C. crescentus*, it could be worth testing if the crosstalk exists

in both directions as well.

Interestingly, we observed that Kup activity varies in excessive concentrations

depending on the media. This indicates that Kup activity is regulated in toxic K$^+$

concentrations depending on environmental resources and/or metabolic states. Also,

suppressor mutations that bypass the growth defect of ∆*kdp* cells in limiting K$^+$

conditions have been mapped in *kup* without impacting the growth when K$^+$ was

available. A plausible explanation would be that mutations found in Kup increase its

affinity for K$^+$, thereby increasing K$^+$ transport to compensate the loss of Kdp, and that

the efflux systems (KefB and KefC) limits the intracellular K$^+$ concentration in abundant

K$^+$ conditions. Studies in other bacteria indicate Kup is a single membrane protein

showing moderate affinity for K$^+$ (Rhoads *et al.*, 1976, Sato *et al.*, 2014). In

Lactococcus lactis, the second messenger c-di-AMP binds to Kup homologues, in

addition to the Trk homologue, to inhibit K$^+$ transport (Quintana *et al.*, 2019). It is

therefore possible that Kup activity is regulated by intracellular signalling molecules to

coordinate K$^+$ uptake with environmental conditions.

KdpD homologues have also been reported to bind to c-di-AMP in other Gram-positive

bacteria (Dutta *et al.*, 2021, Moscoso *et al.*, 2016). High levels of c-di-AMP inhibit the

KdpD-dependent upregulation of the *kdp* operon in *Staphylococcus aureus* (Moscoso

et al., 2016). On top of this, genes encoding K$^+$ transporters are also regulated by a c-

di-AMP-dependent riboswitch. The 5' leader region (LR) of the *B. subtilis kimA*

transcript can adopt two mutually exclusive conformations, one favouring transcription

readthrough while the other promoting transcription termination (Nelson *et al.*, 2013).

Upon direct binding of c-di-AMP, the *kimA* 5' LR preferentially forms a structure

exposing a predicted intrinsic transcription terminator, thereby inhibiting transcription

elongation of *kimA*. Interestingly, a study showed that transcription levels of *kimA* were

higher in low K^+ conditions (0.1 mM KCl) than at higher K^+ concentrations (5 mM KCl)

(Gundlach *et al.*, 2017). Indeed, the high-affinity K^+ transporter KimA was exclusively

detected in low K^+ conditions. This K^+-dependent control of *kimA* expression is

suggested to be achieved through a modulation of c-di-AMP intracellular levels.

Supporting this hypothesis, concentrations of c-di-AMP were increased by two-fold in

high K^+ conditions. More than just increasing in concentration, c-di-AMP becomes

indispensable in high K^+ conditions since a strain lacking c-di-AMP synthesising genes

is viable at low but not at high K^+ concentrations. *C. crescentus* produces c-di-GMP,

rather than c-di-AMP, to control cell development and division. If we speculate that c-

di-GMP levels fluctuate reminiscently to that of c-di-AMP, K^+-depletion would result in

low c-di-GMP levels thereby preventing progression in cell cycle. Although it remains

unclear how K^+ limitation leads to a decrease in c-di-AMP levels, it has been shown

that the enzymes producing c-di-AMP in *B. subtilis* were less abundant in low K^+

conditions (Gundlach *et al.*, 2017). Evaluating the growth behaviour of cells unable to

synthesize c-di-GMP (Abel *et al.*, 2013) in excessive or limiting K^+ conditions could be

an initial step to assess if such regulation is conserved.

The nitrogen-related phosphotransferase component EIIA[Ntr] has been shown in *E. coli*

to regulate not only K^+ transport, by modulating KdpD kinase activity (Luttmann *et al.*,

2015, Mork-Morkenstein *et al.*, 2017, Zimmer *et al.*, 2008) but also phosphate

starvation response, by modulating PhoR activity (Luttmann *et al.*, 2012). For instance,

in limiting K^+ conditions (< 5 mM), the unphosphorylated form of EIIA[Ntr] was shown to

directly interact with KdpD and stimulate its kinase activity, thereby promoting

phosphorylation of KdpE and the subsequent increase of *kdp* genes expression

(Luttmann *et al.*, 2009)*. This EIIA^Ntr-based control of KdpD has also been reported in

the plant symbionts *Rhizobium leguminosarum* (Untiet *et al.*, 2013) and *Sinorhizobium*

fredii (Feng *et al.*, 2022), two α-proteobacteria closely related to *C. crescentus*. It was

later discovered that EIIA^Ntr regulates in fact sigma factor selectivity by modulating K^+

levels. Indeed, the competition between σ^{70} and σ^S for interaction with the RNA

polymerase (RNAP) is influenced by K^+ intracellular levels (Lee *et al.*, 2010).

Depending on the phosphorylation state of EIIA^Ntr, the resulting K^+ intracellular levels

preferentially favour the binding of either σ^{70} or σ^S to the core RNAP (Lee *et al.*, 2010).

Interestingly, PTS^Ntr components regulate the levels of another signalling molecule –

(p)ppGpp – that connects cell division with environmental and metabolic cues in *C.*

crescentus (Hallez *et al.*, 2017, Ronneau *et al.*, 2016, Shyp *et al.*, 2021). Knowing that

(p)ppGpp also influences gene expression by regulating sigma factor competition

(Jishage *et al.*, 2002), it is therefore tempting to speculate that EIIA^Ntr and/or (p)ppGpp

participate to the K^+ homeostasis in *C. crescentus*, even in other alphaproteobacteria.

We also generated strains in which either *kefB* or *kefC* – both coding for efflux pumps

– was inactivated. We observed that deletion of *kefC* ($\Delta kefC$) dramatically impaired

growth in excess of K^+, while the *kefB* mutant ($\Delta kefB$) behaved like the WT strain,

suggesting that KefB cannot support KefC functions. Recently, it was reported that

glutathione controls cell division via negative regulation of the potassium (K^+) efflux

pump KefB (Hartl *et al.*, 2020). Indeed, cell filamentation and decreased intracellular

K^+ levels were observed in mutants unable to synthesize glutathione ($\Delta gshB$),

suggesting that K^+ homeostasis is crucial for bacterial cell cycle regulation.

Interestingly, mutations that suppressed Δ*gshB* phenotypes were found in *kefB* and

kefG, but not in *kefC*. This indicated that, glutathione regulation in *C. crescentus* occurs

mainly through *kefB*. Therefore, it is likely that the glutathione-dependent negative

regulation on KefB still occurred in Δ*kefC* cells, which would explain why the single

kefC knock-out mutant failed to grow when K$^+$ was in excess whereas the single Δ*kefC*

grew.

Methods

Bacterial strains and growth conditions.

All *Escherichia coli* strains used in this thesis were grown aerobically in Luria-Bertani

(LB) broth (Sigma). *E. coli* Top10 was used for cloning purpose while *E. coli* MT607

was used for triparental matings. Thermo-competent cells were used for transformation

of *E. coli*. All *Caulobacter crescentus* strains used are derived from the synchronizable

wild-type strain NA1000. The traditional synthetic media supplemented with glucose

(M2G) was modified by replacing potassium salts by their equivalent sodium salts to

obtain a medium without any K$^+$ (K$_0$) to which the desired K$^+$ concentrations could be

added. The source of K$^+$ consists in both KH_2PO_4 and K_2HPO_4 mixed to an equivalent

ratio in terms of K$^+$ concentration. For each experiment, cells were first grown at 30 °C;

in Peptone Yeast Extract (PYE), then adapted overnight at 30 °C in K$_0$ media

supplemented with 0.5 mM K$^+$, washed twice in K$_0$ media and finally grown at 30 °C in

K$_0$ media supplemented with the desired K$^+$ concentration. For growth curve

measurements, cultures were diluted to a final OD$_{660}$ of 0.02 in a 96-well plate. Growth

was monitored during 48 h with continuous shaking at 30 °C, in an automated plate

reader (Bioscreen C, Lab Systems) measuring OD_{660} every 10 min. For *E. coli*,

antibiotics were used at the following concentrations (μg/ml; in liquid/solid medium):

chloramphenicol (20/30), kanamycin (30/50), oxytetracycline (12.5/12.5),

spectinomycin (100/100), streptomycin (50/100) while for *C. crescentus*, media were

supplemented with kanamycin (5/20), oxytetracycline (2.5/2.5), nalidixic acid (20) when

appropriate. Genes under the control of the inducible xylose promoter (P_{xylX}) were

induced with 0.1 % xylose. Plasmid delivery into *C. crescentus* was achieved by either

bi- or tri-parental mating using *E. coli* S17-1 and *E. coli* MT607 as helper strains,

respectively. In-frame deletions were created by using the pNPTS138-derivative

plasmids as follows. Integration of the plasmids in the *C. crescentus* genome after

single homologous recombination were selected on PYE plates supplemented with

kanamycin. Three independent recombinant clones were inoculated in PYE medium

without kanamycin and incubated overnight at 30 °C. Then, dilutions were spread on

PYE plates supplemented with 3% sucrose and incubated at 30 °C. Single colonies

were picked and transferred onto PYE plates with and without kanamycin. Finally, to

discriminate between mutated and wild-type loci, kanamycin-sensitive clones were

tested by PCR on colony using locus-specific oligonucleotides.

Spotting assays

Ten-fold serial dilutions (in M2G-K) were prepared in 96-well plates from 5 ml cultures

in standard universal tubes grown overnight at 30 °C in the corresponding media. Cells

(5μl) were then spotted on plates, incubated at 30 °C for two-to-three days and pictures

were taken.

β-galactosidase assays

β-galactosidase assays were performed as already described in (Quintero-Yanes *et*

al., 2022). Briefly, cultures were either grown to stationary phase directly in a 96-wells

plate or grown in flasks from which samples were withdrawn throughout growth and

placed at -80 °C in between time points. Permeabilization of cells was performed by

incubating cells with lysis buffer (20 mg/ml polymyxin B, β-mercaptoethanol) for 30 min

at 28 °C. Then, 50 µl of 4 mg/ml O-nitrophenyl- β-D-galactopyranoside (ONPG) were

added. Lysis and ONPG solutions were prepared using Z buffer as a solvent (60 mM

Na_2HPO_4, 40 mM NaH_2PO_4, 10 mM KCl, 1 mM $MgSO_4$, pH 7.0). Both OD_{420} and OD_{550}

were measured every minute for 30 min at 30 °C using a fluorimeter (SpectraMax®,

Molecular Devices). Miller Units were calculated using the following formula: M.U. =

$\frac{[OD420 - (1.75 \, x \, OD550)]x \, 1000}{OD660 x \, t \, x \, v}$ where t is the reaction time in min v is the volume of culture

used in ml. Then, ONPG hydrolysis was measured at 30 °C for 30 min. The activity of

the β-galactosidase expressed in miller units (MU) was calculated using the following

equation: MU = $(OD_{420} \times 1{,}000) / [OD_{660} \times t \times v]$ where "t" is the time of the reaction

(min), and "v" is the volume of cultures used in the assays (ml). Experimental values

were the average of three independent experiments.

Protein purification and production of polyclonal antibodies

To immunize rabbits for production of polyclonal antibodies, an *E. coli* BL21 (DE3)

pLysS strain carrying plasmid pET-28a-*kdpE* was grown in LB medium supplemented

with kanamycin and chloramphenicol until an OD600 of 0.5. After addition of IPTG to

a final concentration of 1 mM, the culture was incubated at 18 °C for 18 hrs. Cells were

then harvested by centrifugation for 30 min at 5,000 x g, 4 °C. The pellet was

resuspended in 20 ml of binding buffer (20 mM Tris-HCl pH 8.0, 500 mM NaCl, 10%

glycerol, 10 mM $MgCl_2$, 12.5 mM Imidazole) supplemented with complete EDTA-free

protease cocktail inhibitor (Roche), 400 mg lysozyme (Sigma) and 10 mg DNaseI

(Roche) and incubated for 30 min on ice. Cells were then lysed by sonication. After

centrifugation at 12,000 rpm for 30 min at 4°C, the lysate was loaded on a Ni-NTA

column and incubated 1 hr at 4 °C with end-over-end agitation. The column was then

washed with 5 ml binding buffer, 3 ml Wash1 buffer (binding buffer with 25 mM

imidazole), 3 ml Wash2 buffer (binding buffer with 50 mM imidazole), 3 ml Wash3

buffer (binding buffer with 75 mM imidazole). Proteins bound to the column were eluted

with 3 ml Elution buffer (binding buffer with 100 mM imidazole) and aliquoted in 300 μl

fractions. All the fractions containing the protein of interest (checked by Coomassie

blue staining) were pooled and dialyzed in Dialysis buffer (20 mM Tris pH 8, 500 mM

NaCl, 10% glycerol). Purified KdpE was used to immunize rabbits (CER Groupe,

Marloie, Belgium).

Tn-seq analysis.

A mini-Tn5 was introduced in *C. crescentus* NA1000 WT strain by bi-parental mating.

Briefly, overnight cultures of recipient and donor strains (grown in LB supplemented

with kanamycin) were mixed in 95:5 ratio to a final volume of 1 ml. Cells were harvested

by centrifugation at 9,000 rpm (7,600 x g) at room temperature for 2 min. The

supernatant was removed, and 1 ml of fresh PYE medium was added to wash the

pellet. A second centrifugation step was done to remove again the supernatant.

Thereafter, the pellet was resuspended in 50 μl and spotted on PYE agar and

incubated at 30 ° C for 4h. Cells were collected, resuspended in 10 ml M2G or M2G-K

and washed twice with the same volume. Dilutions (10^{-6}) were plated on M2G or M2G-

K plates supplemented with aztreonam (5μg l^{-1}) and kanamycin (5μg l^{-1}) and incubated

at 30 ° C for 5 days. Then, at least 3 x10^5 colonies were collected in M2G or M2G-K

supplemented with 10% glycerol and stored at -80 °C. Genomic DNA was then

extracted following the *Nucleospin Tissue Kit* protocol (Macherey-Nagel) and

resuspended in 50 µl elution buffer (5 mM Tris-HCl [pH 8.5]). DNA sequencing was

performed using an Illumina NextSeq (paired-end 2x75) instrument (Fasteris, Geneva,

Switzerland). Reads corresponding to the mini-Tn5 insertion sites were first filtered

with the 5'-GGTTGAGATGTGTATAAGAGACAG sequence before being processed as

described in (Coppine *et al.*, 2020). Briefly, filtered reads were mapped on the genome

of *C. crescentus* NA1000 (GenBank accession number NC_011916.1) and converted

to Sequence Alignment/Map (SAM) format using the Burrows-Wheeler Aligner (BWA)

and SAMtools, respectively, from the Sourceforge server (https://sourceforge.net/).

Next, the number of reads overlapping each genomic position was computed using

custom Python scripts. The total number of reads for internal 80% of each ORF in each

condition were then computed using custom Python scripts. For comparing strains, the

total number of reads was normalized by the ratio of the number of reads between the

two strains. Ward's clustering analysis was performed to define the fitness categories

as previously reported.

Chromatin immunoprecipitation followed by deep sequencing (ChIP-Seq assay)

ChIP-Seq was performed as already described in (Coppine *et al.,* 2020). Briefly, 80 ml

of mid-log-phase cells (OD_{660} of 0.6) were cross-linked in 1% formaldehyde and 10

mM sodium phosphate (pH 7.6) at room temperature (RT) for 10 min and then for 30

min on ice. Cross-linking was stopped by addition of 125 mM glycine and incubated for

5 min on ice. Cells were washed twice in phosphate buffer solution (PBS; 137 mM

NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, pH 7.4) resuspended in 450 µl

TES buffer (10 mM Tris-HCl [pH 7.5], 1 mM EDTA, and 100 mM NaCl), and lysed with

2 μl of Ready-lyse lysozyme solution for 5 min at RT. Protease inhibitors (Roche) were

added, and the mixture was incubated for 10 min. Then, 550 μl of ChIP buffer (1.1%

Triton X-100, 1.2 mM EDTA, 16.7 mM Tris-HCl [pH 8.1], 167 mM NaCl, protease

inhibitors) were added to the lysate and incubated at 37 °C for 10 min before sonication

(2 x 8 bursts of 30 sec on ice using a Diagenode Bioruptor) to shear DNA fragments

to a length of 300 to 500 bp. Lysate was cleared by centrifugation for 10 min at 12,500

rpm at 4 °C, and protein content was assessed by measuring the OD_{280}. Then, 7.5 mg

of proteins was diluted in ChIP buffer supplemented with 0.01% SDS and precleared

for 1 hr at 4 °C with 50 μl of SureBeads Protein A Magnetic Beads (BioRad) and 100

μg bovine serum albumin (BSA). One microliter of polyclonal anti-KdpE antibodies was

added to the supernatant before overnight incubation at 4 °C under gentle agitation.

Next, 80 μl of BSA presaturated protein A-agarose beads were added to the solution

and incubated for 2 hrs at 4 °C with rotation, washed once with low-salt buffer (0.1%

SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl [pH 8.1], 150 mM NaCl), once

with high-salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl [pH

8.1], 500 mM NaCl), once with LiCl buffer (0.25 M LiCl, 1% NP-40, 1% deoxycholate,

1 mM EDTA, 10 mM Tris-HCl [pH 8.1]), and once with TE buffer (10 mM Tris-HCl [pH

8.1] 1 mM EDTA) at 4 °C, followed by a second wash with TE buffer at RT. The DNA-

protein complexes were eluted twice in 250 μl freshly prepared elution buffer (0.1 M

NaHCO3, 1% SDS). NaCl was added at a concentration of 300 mM to the combined

eluates (500 μl) before overnight incubation at 65 °C to reverse the cross-link. The

samples were treated with 20 μg of proteinase K in 40mM EDTA and 40 mM Tris-HCl

(pH 6.5) for 2 h at 45 °C. DNA was extracted using a Nucleospin PCR cleanup kit

(Macherey-Nagel) and resuspended in 50 μl elution buffer (5 mM Tris-HCl [pH 8.5]).

DNA sequencing was performed using an Illumina NextSeq 550 (paired-end 2x75)

instrument (Seqalis). NGS data were analysed as described in (Coppine *et al.*, 2020).

RNA-Seq

WT and Δ*kdpE* cells were grown overnight to $OD_{660} \sim 0.3$ in M2G-K 0,025 mM K^+ and

WT in M2G-K 0,5 mM K^+. Thereafter, total RNA was extracted with RNeasy® Protect

Bacteria Kit from Qiagen and following manufacturer's instructions. The quantity and

quality (A260/A280 ration) of RNA was determined with a Thermo Scientific™

Nanodrop ™ One Microvolume UV-Vis Spectrophotometer. RNA-Seq TTRNA libraries

were prepared according to the manufacturer's instructions and sequenced with

Illumina NovaSeq 6000 (paired-end 2x100) instrument (Seqalis). NGS data were

analysed using Galaxy (https://usegalaxy.org) (Quintero-Yanes *et al.*, 2022). Briefly,

FastQC was used to evaluate the quality of the reads; HISAT2 was used to map the

reads onto the NA1000 reference genome (NC_011916.1) and generate bam files;

featureCounts was used to generate counts tables using bam files and DESeq2 was

used to determine differentially expressed genes. The Volcano plot was generated

using GraphPad Prism 9 software.

MEME analysis

Sequences corresponding to the peaks founds in ChIP-seq were analysed for a KdpE

conserved DNA binding motif using MEME (Quintero-Yanes *et al.*, 2022) (**Table S6**).

As searching parameters, the classical motif discovery mode, zoops motif site

distribution, 10 minimum motif width and 20 maximum motif width, 2 as maximum

number of motifs and a p-value<0.001 as cut-off were used. The motif with highest site

count was selected.

Statistical analyses

All the statistical analyses were performed using GraphPad Prism 9 software. A *P* value of <0.05 was considered as significant.

Data availability

ChIP-Seq and RNA-Seq data have been deposited to the Gene Expression Omnibus (GEO) repository with the accession number XXX.

References

Abel, S., Bucher, T., Nicollier, M., Hug, I., Kaever, V., Abel Zur Wiesch, P., and Jenal, U. (2013) Bi-modal distribution of the second messenger c-di-GMP controls cell fate and asymmetry during the caulobacter cell cycle. *PLoS Genet* **9**: e1003744.

Ahijado-Guzman, R., Alfonso, C., Reija, B., Salvarelli, E., Mingorance, J., Zorrilla, S., Monterroso, B., and Rivas, G. (2013) Control by potassium of the size distribution of Escherichia coli FtsZ polymers is independent of GTPase activity. *J Biol Chem* **288**: 27358-27365.

Ali, M.K., Li, X., Tang, Q., Liu, X., Chen, F., Xiao, J., Ali, M., Chou, S.H., and He, J. (2017) Regulation of Inducible Potassium Transporter KdpFABC by the KdpD/KdpE Two-Component System in Mycobacterium smegmatis. *Front Microbiol* **8**: 570.

Bakker, E.P., and Mangerich, W.E. (1981) Interconversion of components of the bacterial proton motive force by electrogenic potassium transport. *J Bacteriol* **147**: 820-826.

Booth, I.R. (1999) The regulation of intracellular pH in bacteria. *Novartis Found Symp* **221**: 19-28; discussions 28-37.

Bossemeyer, D., Borchard, A., Dosch, D.C., Helmer, G.C., Epstein, W., Booth, I.R., and Bakker, E.P. (1989a) K+-transport protein TrkA of Escherichia coli is a peripheral membrane protein that requires other trk gene products for attachment to the cytoplasmic membrane. *J Biol Chem* **264**: 16403-16410.

Bossemeyer, D., Schlosser, A., and Bakker, E.P. (1989b) Specific cesium transport via the Escherichia coli Kup (TrkD) K+ uptake system. *J Bacteriol* **171**: 2219-2221.

Cann, M. (2007) A subset of GAF domains are evolutionarily conserved sodium sensors. *Mol Microbiol* **64**: 461-472.

Cao, Y., Jin, X., Huang, H., Derebe, M.G., Levin, E.J., Kabaleeswaran, V., Pan, Y., Punta, M., Love, J., Weng, J., Quick, M., Ye, S., Kloss, B., Bruni, R., Martinez-Hackert, E., Hendrickson, W.A., Rost, B., Javitch, J.A., Rajashankar, K.R., Jiang, Y., and Zhou, M. (2011) Crystal structure of a potassium ion transporter, TrkH. *Nature* **471**: 336-340.

Castaneda-Garcia, A., Do, T.T., and Blazquez, J. (2011) The K+ uptake regulator TrkA controls membrane potential, pH homeostasis and multidrug susceptibility in Mycobacterium smegmatis. *J Antimicrob Chemother* **66**: 1489-1498.

Coppine, J., Kaczmarczyk, A., Petit, K., Brochier, T., Jenal, U., and Hallez, R. (2020) Regulation of Bacterial Cell Cycle Progression by Redundant Phosphatases. *J Bacteriol* **202**.

Curtis, P.D., and Brun, Y.V. (2014) Identification of essential alphaproteobacterial genes
reveals operational variability in conserved developmental and cell cycle systems. *Mol*
Microbiol **93**: 713-735.

Danchin, A., and Nikel, P.I. (2019) Why Nature Chose Potassium. *J Mol Evol* **87**: 271-288.

Dani, P., Ujaoney, A.K., Apte, S.K., and Basu, B. (2017) Regulation of potassium dependent
ATPase (kdp) operon of Deinococcus radiodurans. *PLoS One* **12**: e0188998.

de Oliveira, M.V., Intorne, A.C., Vespoli, L.S., Andrade, L.F., Pereira, L.M., Rangel, P.L., and de
Souza Filho, G.A. (2017) Essential role of K(+) uptake permease (Kup) for resistance to
sucrose-induced stress in Gluconacetobacter diazotrophicus PAl 5. *Environ Microbiol*
Rep **9**: 85-90.

Dominguez-Ferreras, A., Munoz, S., Olivares, J., Soto, M.J., and Sanjuan, J. (2009) Role of
potassium uptake systems in Sinorhizobium meliloti osmoadaptation and symbiotic
performance. *J Bacteriol* **191**: 2133-2143.

Dosch, D.C., Helmer, G.L., Sutton, S.H., Salvacion, F.F., and Epstein, W. (1991) Genetic analysis
of potassium transport loci in Escherichia coli: evidence for three constitutive systems
mediating uptake potassium. *J Bacteriol* **173**: 687-696.

Dutta, A., Batish, M., and Parashar, V. (2021) Structural basis of KdpD histidine kinase binding
to the second messenger c-di-AMP. *J Biol Chem* **296**: 100771.

Epstein, W. (2003) The roles and regulation of potassium in bacteria. *Prog Nucleic Acid Res*
Mol Biol **75**: 293-320.

Feifel, B., Sandmeier, E., Schonfeld, H.J., and Christen, P. (1996) Potassium ions and the
molecular-chaperone activity of DnaK. *Eur J Biochem* **237**: 318-321.

Feng, X.Y., Tian, Y., Cui, W.J., Li, Y.Z., Wang, D., Liu, Y., Jiao, J., Chen, W.X., and Tian, C.F. (2022)
The PTS(Ntr)-KdpDE-KdpFABC Pathway Contributes to Low Potassium Stress
Adaptation and Competitive Nodulation of Sinorhizobium fredii. *mBio* **13**: e0372121.

Gohara, D.W., and Di Cera, E. (2016) Molecular Mechanisms of Enzyme Activation by
Monovalent Cations. *J Biol Chem* **291**: 20840-20848.

Grason, J.P., Gresham, J.S., Widjaja, L., Wehri, S.C., and Lorimer, G.H. (2008) Setting the
chaperonin timer: the effects of K+ and substrate protein on ATP hydrolysis. *Proc Natl*
Acad Sci U S A **105**: 17334-17338.

Gries, C.M., Bose, J.L., Nuxoll, A.S., Fey, P.D., and Bayles, K.W. (2013) The Ktr potassium
transport system in Staphylococcus aureus and its role in cell physiology, antimicrobial
resistance and pathogenesis. *Mol Microbiol* **89**: 760-773.

Gubaev, A., and Klostermeier, D. (2012) Potassium ions are required for nucleotide-induced
closure of gyrase N-gate. *J Biol Chem* **287**: 10916-10921.

Gundlach, J., Herzberg, C., Kaever, V., Gunka, K., Hoffmann, T., Weiss, M., Gibhardt, J.,
Thurmer, A., Hertel, D., Daniel, R., Bremer, E., Commichau, F.M., and Stulke, J. (2017)
Control of potassium homeostasis is an essential function of the second messenger
cyclic di-AMP in Bacillus subtilis. *Sci Signal* **10**.

Hallez, R., Delaby, M., Sanselicio, S., and Viollier, P.H. (2017) Hit the right spots: cell cycle
control by phosphorylated guanosines in alphaproteobacteria. *Nat Rev Microbiol* **15**:
137-148.

Harms, C., Domoto, Y., Celik, C., Rahe, E., Stumpe, S., Schmid, R., Nakamura, T., and Bakker,
E.P. (2001) Identification of the ABC protein SapD as the subunit that confers ATP
dependence to the K+-uptake systems Trk(H) and Trk(G) from Escherichia coli K-12.
Microbiology (Reading) **147**: 2991-3003.

Hartl, J., Kiefer, P., Kaczmarczyk, A., Mittelviefhaus, M., Meyer, F., Vonderach, T., Hattendorf,
B., Jenal, U., and Vorholt, J.A. (2020) Untargeted metabolomics links glutathione to
bacterial cell cycle progression. *Nat Metab* **2**: 153-166.
Heermann, R., and Jung, K. (2010) The complexity of the 'simple' two-component system
KdpD/KdpE in Escherichia coli. *FEMS Microbiol Lett* **304**: 97-106.
Heikaus, C.C., Pandit, J., and Klevit, R.E. (2009) Cyclic nucleotide binding GAF domains from
phosphodiesterases: structural and mechanistic insights. *Structure* **17**: 1551-1557.
Hocking, J., Priyadarshini, R., Takacs, C.N., Costa, T., Dye, N.A., Shapiro, L., Vollmer, W., and
Jacobs-Wagner, C. (2012) Osmolality-dependent relocation of penicillin-binding
protein PBP2 to the division site in Caulobacter crescentus. *J Bacteriol* **194**: 3116-3127.
Humphries, J., Xiong, L., Liu, J., Prindle, A., Yuan, F., Arjes, H.A., Tsimring, L., and Suel, G.M.
(2017) Species-Independent Attraction to Biofilms through Electrical Signaling. *Cell*
168: 200-209 e212.
Jishage, M., Kvint, K., Shingler, V., and Nystrom, T. (2002) Regulation of sigma factor
competition by the alarmone ppGpp. *Genes Dev* **16**: 1260-1270.
Karp, P.D., Billington, R., Caspi, R., Fulcher, C.A., Latendresse, M., Kothari, A., Keseler, I.M.,
Krummenacker, M., Midford, P.E., Ong, Q., Ong, W.K., Paley, S.M., and Subhraveti, P.
(2019) The BioCyc collection of microbial genomes and metabolic pathways. *Brief*
Bioinform **20**: 1085-1093.
Klenow, H., and Henningsen, I. (1969) Effect of monovalent cations on the activity of the DNA
polymerase of Escherichia coli B. *Eur J Biochem* **9**: 133-141.
Laimins, L.A., Rhoads, D.B., and Epstein, W. (1981) Osmotic control of kdp operon expression
in Escherichia coli. *Proc Natl Acad Sci U S A* **78**: 464-468.
Lee, C.R., Cho, S.H., Kim, H.J., Kim, M., Peterkofsky, A., and Seok, Y.J. (2010) Potassium
mediates Escherichia coli enzyme IIA(Ntr) -dependent regulation of sigma factor
selectivity. *Mol Microbiol* **78**: 1468-1483.
Liu, L.F., and Wang, J.C. (1978) Micrococcus luteus DNA gyrase: active components and a
model for its supercoiling of DNA. *Proc Natl Acad Sci U S A* **75**: 2098-2102.
Liu, Y., Ho, K.K., Su, J., Gong, H., Chang, A.C., and Lu, S. (2013) Potassium transport of
Salmonella is important for type III secretion and pathogenesis. *Microbiology*
(Reading) **159**: 1705-1719.
Luttmann, D., Gopel, Y., and Gorke, B. (2012) The phosphotransferase protein EIIA(Ntr)
modulates the phosphate starvation response through interaction with histidine
kinase PhoR in Escherichia coli. *Mol Microbiol* **86**: 96-110.
Luttmann, D., Gopel, Y., and Gorke, B. (2015) Cross-Talk between the Canonical and the
Nitrogen-Related Phosphotransferase Systems Modulates Synthesis of the KdpFABC
Potassium Transporter in Escherichia coli. *J Mol Microbiol Biotechnol* **25**: 168-177.
Luttmann, D., Heermann, R., Zimmer, B., Hillmann, A., Rampp, I.S., Jung, K., and Gorke, B.
(2009) Stimulation of the potassium sensor KdpD kinase activity by interaction with
the phosphotransferase protein IIA(Ntr) in Escherichia coli. *Mol Microbiol* **72**: 978-994.
MacGilvary, N.J., Kevorkian, Y.L., and Tan, S. (2019) Potassium response and homeostasis in
Mycobacterium tuberculosis modulates environmental adaptation and is important
for host colonization. *PLoS Pathog* **15**: e1007591.
Matroule, J.Y., Lam, H., Burnette, D.T., and Jacobs-Wagner, C. (2004) Cytokinesis monitoring
during development; rapid pole-to-pole shuttling of a signaling protein by localized
kinase and phosphatase in Caulobacter. *Cell* **118**: 579-590.

Mork-Morkenstein, M., Heermann, R., Gopel, Y., Jung, K., and Gorke, B. (2017) Non-canonical
activation of histidine kinase KdpD by phosphotransferase protein PtsN through
interaction with the transmitter domain. *Mol Microbiol* **106**: 54-73.
Moscoso, J.A., Schramke, H., Zhang, Y., Tosi, T., Dehbi, A., Jung, K., and Grundling, A. (2016)
Binding of Cyclic Di-AMP to the Staphylococcus aureus Sensor Kinase KdpD Occurs via
the Universal Stress Protein Domain and Downregulates the Expression of the Kdp
Potassium Transporter. *J Bacteriol* **198**: 98-110.
Nelson, J.W., Sudarsan, N., Furukawa, K., Weinberg, Z., Wang, J.X., and Breaker, R.R. (2013)
Riboswitches in eubacteria sense the second messenger c-di-AMP. *Nat Chem Biol* **9**:
834-839.
Ochrombel, I., Ott, L., Kramer, R., Burkovski, A., and Marin, K. (2011) Impact of improved
potassium accumulation on pH homeostasis, membrane potential adjustment and
survival of Corynebacterium glutamicum. *Biochim Biophys Acta* **1807**: 444-450.
Pedersen, B.P., Stokes, D.L., and Apell, H.J. (2019) The KdpFABC complex - K(+) transport
against all odds. *Mol Membr Biol* **35**: 21-38.
Polarek, J.W., Williams, G., and Epstein, W. (1992) The products of the kdpDE operon are
required for expression of the Kdp ATPase of Escherichia coli. *J Bacteriol* **174**: 2145-
2151.
Prindle, A., Liu, J., Asally, M., Ly, S., Garcia-Ojalvo, J., and Suel, G.M. (2015) Ion channels enable
electrical communication in bacterial communities. *Nature* **527**: 59-63.
Quintana, I.M., Gibhardt, J., Turdiev, A., Hammer, E., Commichau, F.M., Lee, V.T., Magni, C.,
and Stulke, J. (2019) The KupA and KupB Proteins of Lactococcus lactis IL1403 Are
Novel c-di-AMP Receptor Proteins Responsible for Potassium Uptake. *J Bacteriol* **201**.
Quintero-Yanes, A., Mayard, A., and Hallez, R. (2022) The two-component system ChvGI
maintains cell envelope homeostasis in Caulobacter crescentus. *PLoS Genet* **18**:
e1010465.
Quintero-Yanes, A., Monson, R.E., and Salmond, G.P.C. (2019) Environmental potassium
regulates bacterial flotation, antibiotic production and turgor pressure in Serratia
through the TrkH transporter. *Environ Microbiol* **21**: 2499-2510.
Rasmussen, T. (2023) The Potassium Efflux System Kef: Bacterial Protection against Toxic
Electrophilic Compounds. *Membranes (Basel)* **13**.
Rhoads, D.B., Waters, F.B., and Epstein, W. (1976) Cation transport in Escherichia coli. VIII.
Potassium transport mutants. *J Gen Physiol* **67**: 325-341.
Roe, A.J., McLaggan, D., O'Byrne, C.P., and Booth, I.R. (2000) Rapid inactivation of the
Escherichia coli Kdp K+ uptake system by high potassium concentrations. *Mol
Microbiol* **35**: 1235-1243.
Ronneau, S., Petit, K., De Bolle, X., and Hallez, R. (2016) Phosphotransferase-dependent
accumulation of (p)ppGpp in response to glutamine deprivation in Caulobacter
crescentus. *Nat Commun* **7**: 11423.
Roosild, T.P., Castronovo, S., Healy, J., Miller, S., Pliotas, C., Rasmussen, T., Bartlett, W.,
Conway, S.J., and Booth, I.R. (2010) Mechanism of ligand-gated potassium efflux in
bacterial pathogens. *Proc Natl Acad Sci U S A* **107**: 19784-19789.
Rozov, A., Khusainov, I., El Omari, K., Duman, R., Mykhaylyk, V., Yusupov, M., Westhof, E.,
Wagner, A., and Yusupova, G. (2019) Importance of potassium ions for ribosome
structure and function revealed by long-wavelength X-ray diffraction. *Nat Commun* **10**:
2519.

Sato, Y., Nanatani, K., Hamamoto, S., Shimizu, M., Takahashi, M., Tabuchi-Kobayashi, M.,
Mizutani, A., Schroeder, J.I., Souma, S., and Uozumi, N. (2014) Defining membrane
spanning domains and crucial membrane-localized acidic amino acid residues for K(+)
transport of a Kup/HAK/KT-type Escherichia coli potassium transporter. *J Biochem* **155**:
315-323.
Schleyer, M., and Bakker, E.P. (1993) Nucleotide sequence and 3'-end deletion studies indicate
that the K(+)-uptake protein kup from Escherichia coli is composed of a hydrophobic
core linked to a large and partially essential hydrophilic C terminus. *J Bacteriol* **175**:
6925-6931.
Schramke, H., Laermann, V., Tegetmeyer, H.E., Brachmann, A., Jung, K., and Altendorf, K.
(2017) Revisiting regulation of potassium homeostasis in Escherichia coli: the
connection to phosphate limitation. *Microbiologyopen* **6**.
Schramke, H., Tostevin, F., Heermann, R., Gerland, U., and Jung, K. (2016) A Dual-Sensing
Receptor Confers Robust Cellular Homeostasis. *Cell Rep* **16**: 213-221.
Shyp, V., Dubey, B.N., Bohm, R., Hartl, J., Nesper, J., Vorholt, J.A., Hiller, S., Schirmer, T., and
Jenal, U. (2021) Reciprocal growth control by competitive binding of nucleotide second
messengers to a metabolic switch in Caulobacter crescentus. *Nat Microbiol* **6**: 59-72.
Su, J., Gong, H., Lai, J., Main, A., and Lu, S. (2009) The potassium transporter Trk and external
potassium modulate Salmonella enterica protein secretion and virulence. *Infect*
Immun **77**: 667-675.
Trchounian, A., and Kobayashi, H. (1999) Kup is the major K+ uptake system in Escherichia coli
upon hyper-osmotic stress at a low pH. *FEBS Lett* **447**: 144-148.
UniProt, C. (2023) UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Res*
51: D523-D531.
Untiet, V., Karunakaran, R., Kramer, M., Poole, P., Priefer, U., and Prell, J. (2013) ABC transport
is inactivated by the PTS(Ntr) under potassium limitation in Rhizobium leguminosarum
3841. *PLoS One* **8**: e64682.
Valente, R.S., and Xavier, K.B. (2016) The Trk Potassium Transporter Is Required for RsmB-
Mediated Activation of Virulence in the Phytopathogen Pectobacterium wasabiae. *J*
Bacteriol **198**: 248-255.
Zakharyan, E., and Trchounian, A. (2001) K+ influx by Kup in Escherichia coli is accompanied
by a decrease in H+ efflux. *FEMS Microbiol Lett* **204**: 61-64.
Zimmer, B., Hillmann, A., and Gorke, B. (2008) Requirements for the phosphorylation of the
Escherichia coli EIIANtr protein in vivo. *FEMS Microbiol Lett* **286**: 96-102.

1007

1008

1009

1010

1011

 Figures

Figure 1. Impact of K⁺ on *C. crescentus* growth. Growth of WT in M2G-K supplemented with different K^+ concentrations using K_2HPO_4 + KH_2PO_4 (A-B) and KCl (C-D) as K^+ source. Data represent average, n=3, and error bars= ±SD. Other concentrations tested are presented in Fig. S1.

1018

Figure 2. K⁺ transport and response systems in Gram-negative bacteria.
Schematic diagram of K⁺-related proteins predicted to work at high (A-B) or low (C) K⁺
concentrations. Coloured proteins correspond to the ones encoded in *C. crescentus*
genome: CCNA_01688, Kup (CCNA_00130), KefB (CCNA_00204), KefG
(CCNA_00205), KefC (CCNA_03611), KdpDE (CCNA_01666-CCNA_01667)

1024 KdpFABC (CCNA_01662-CCNA_01665). The genes were identified using Biocyc

1025 collection of microbial genomes (Karp *et al.,* 2019) and structures of the proteins

1026 obtained using predictions with alphafold from Uniprot (UniProt, 2023). Outer

1027 Membrane (OM), peptidoglycan (PG), inner membrane (IM).

1028

1029 **Figure 3. Deletion of K+-related genes impact *C. crescentus* growth.** Growth of

1030 WT, ΔkefB, ΔkefC, ΔCCNA_01688 and ΔkdpABCDE mutants in minimal media M2G-

1031 K supplemented with (A) limiting (0.025 mM), (B) abundant (0.5 mM) and (C) excessive

1032 (50 mM) K+ concentrations. Data represent average, n=3, and error bars= ±SD.

1033

Figure 4. Tn-seq and RNA-seq profile of *C. crescentus* cells in limiting K⁺

conditions. (A) Growth of WT and Δ*kdp* cells in M2G-K plates without or

supplemented with K+. (B) Number of Transposon-insertions (#Tn-insertions) against

the length (#bp) of 4186 genes in M2G and M2G-K media. Non-essential genes are

highlighted in grey, high-fitness cost genes in light blue, and essential in dark blue. (C)

Representation of genes that changed in their fitness-cost categories according to

analysis done in M2G and M2G-K (B). (B-C) The percentage of genes in each category

is indicated following the colour code.

1042

Figure 5. Impact of deletion and overexpression of the *kup* gene. (A) Growth of

WT and Δ*kup* mutant in minimal media M2G-K supplemented with limiting (0.025 mM),

abundant (0.5 mM), excessive (50 mM) K+ concentrations or limiting (0.025 mM) K+

concentrations and 3% sucrose. (B) Growth of WT expressing *kup* under xylose

induction. (C) Growth of WT, Δ*kdp* and Δ*kdp* suppressors (Δ*kdp*^{SUP}) in minimal media

M2G-K supplemented with limiting (0.025 mM) or abundant (0.5 mM). Data represent

average, n=3, and error bars= ±SD.

1050

Figure 6. K$^+$ and KdpE-dependent regulon. (A-B) Volcano plots representing the

relation between the log$_2$ fold change (FC) and -log False Discovery Rate (FDR) on

gene expression between (A) WT in limiting *vs* abundant K$^+$ condition, and (B) Δ*kdpE*

vs WT in limiting K$^+$ condition. Genes identified are presented as dots. Significant

down- and up-regulated genes are presented as blue (A) and purple (B) dots, while

genes with no significant alterations are presented as black dots. Top 10 down and

upregulated genes are highlighted for each analysis. (C) Genome-wide occupancy of

KdpE on the chromosome of *C. crescentus* determined by ChIP-seq on Δ*kdpE*

P$_{xylX}$::*kdpE* Δ*xylX* and Δ*kdpDE* P$_{xylX}$::*kdpE* Δ*xylX*. The x-axis represents the coordinates

on the genome (Mb), the y-axis shows the normalized ChIP-Seq read abundance in

reads. The promoter region of *kdp* is highlighted. (D) Occupancy of KdpE on the *kdp*

promoter Δ*kdpE* P$_{xylX}$::*kdpE* Δ*xylX* and Δ*kdpDE* P$_{xylX}$::*kdpE* Δ*xylX*. (E) WebLogo of

predicted KdpE consensus sequence obtained with MEME and identification (with a *p-*

value 7.71 x 10^{-5}) of the DNA binding sequence upstream of *kdpA.* Arrows indicate

gene orientation for expression.

1066

Figure 7. KdpD downregulates KdpE to control growth in low K⁺. (A) Growth of WT, ΔkdpABC, ΔkdpD, ΔkdpE and ΔkdpDE. (B) Growth of WT and ΔkdpD carrying either pMR10 empty vectors (EV) or pMR10 P$_{kdp}$::kdpD. (C) Growth of WT, ΔkdpE, ΔkdpDE expressing kdpE from the xylX locus (P$_{xylX}$::kdpE). (D) Growth of kdpD and kdpE catalytic point mutants. Growth was done in minimal media M2G-K supplemented with limiting (0.025 mM) K⁺ concentrations (A-D), media was additionally supplemented tetracycline for plasmid selection (B) or with 0.01 % xylose

1074   for the induction of *kdpE* expression (C). Data represent average, n=3, and error bars=

1075   ±SD.

A

B

C

1076

Figure 8. Regulation of *kdp* expression (A) β-Galactosidase activities of WT and *kdp*

mutants carrying a transcriptional P$_{kdp}$::*lacZ* fusion grown in M2G-K supplemented with

limiting (0.025 mM) or abundant (0.5 mM). β-Galactosidase activities of WT and *kdp*

mutants carrying a transcriptional P$_{kdp}$::*lacZ* fusion along the growth in M2G-K

supplemented with limiting (0.025 mM). Statistical analyses were carried out via single

factor ANOVA. Data represent average, n=3, and error bars= ±SD.

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093