# A seascape approach to the microbial biogeography of mesophotic and

# deep-sea octocorals

Samuel A. Vohsen [1*] & Santiago Herrera [1#]

[1] Department of Biological Sciences, Lehigh University, Bethlehem, PA

[*] Corresponding Author: sav419@lehigh.edu

[#] Co-corresponding Author: santiago.herrera@lehigh.edu

**Competing Interests Statement**

The authors declare no conflicts of interest

**Abstract**:

Coral-associated microbiomes vary greatly between coral colonies and sampling
locations with functional consequences on the coral host. However, the full extent of variability
across the ranges of most coral species remains unknown, especially in octocorals and species
from mesophotic and deep-sea habitats. Here we characterized the microbiomes of four octocoral
species from mesophotic and deep-sea habitats in the northern Gulf of Mexico, *Muricea pendula,*
*Swiftia exserta, Callogorgia delta,* and *Paramuricea biscaya* using 16S metabarcoding. We
tested for microbiome differentiation between and within species, examining the influence of the
coral's genotype and environmental factors that vary with depth (53-2224 m) and geographic
location (over 680 m). Coral microbiomes were often dominated by a single amplicon sequence
variant – ASV (>50% relative abundance) whose abundance varied across the range of the host
coral. These included corallicolid apicomplexans, *Endozoicomonas*, members of the Mollicutes,
and the BD1-7 clade of Spongiibacteraceae. Coral species, depth, and geographic location
significantly affected microbial community composition, richness, and evenness, as well as the
abundance of individual microbes. Differences in bottom temperature and surface primary
productivity could explain part of the variation associated with depth and geographic location,
respectively. However the coral genotype was confounded with depth so its effect could not be
dissected. Altogether, this work demonstrates that the microbiomes of corals vary substantially
across their ranges in ways that may have functional consequences, identifies important
ecological drivers in mesophotic and deep-sea corals, and can inform restoration efforts.

**Introduction**

Shallow-water, reef-building corals are well-known for associating with microalgae in the family Symbiodinaceae. Still, all corals, including those in the deep sea, form associations with diverse groups of bacteria[1], archaea[2, 3], unicellular eukaryotes[4], and viruses[5], collectively referred to as the coral microbiome. The coral microbiome resides throughout the coral, including skeleton[6, 7], tissues[8, 9], gastrovascular cavity[10], and even the mucus[1, 11]. Many members of the microbiome have proven important to the health of corals, either as pathogens[12] or providing sources of nutrition[13, 14].

Common coral associates include eukaryotic apicomplexans of the family Corallicolidae[15-19] and the bacterial genera *Mycoplasma*[20-25] and *Endozoicomonas*[1, 26-28]. These taxa are among the most-studied coral-associated microbes and exhibit many indications of a close relationship with their coral hosts. *Endozoicomonas* form dense aggregates in coral tissue[9, 29]. They are often dominant in individual microbiomes[29, 30] of soft corals of the subclass Octocorallia[27, 28, 31], hard corals of the order Scleractinia [1], and black corals of the order Antipatharia[32]. *Endozoicomonas* isolated from corals possess capabilities relevant to the coral host such as the ability to produce vitamins[33, 34], degrade dimethylsulfoniopropionate[35], cycle phosphorus[8], and degrade chitin[30]. *Endozoicomonas* responds to coral tissue extracts by activating genes involved in initiating symbiosis[33]. Similarly, corallicolid apicomplexans and *Mycoplasma* are commonly found in many coral species and reside within or on coral tissue[16, 24, 36]. Most apicomplexans[37] and *Mycoplasma* spp.[38] are parasites or pathogens. However, those found in corals have not yet been associated with evident pathologies or reduced fitness[39]. While these common coral associates have been the focus of extensive research, we have a limited understanding of the diversity of coral- microbe associations.

The compositions of coral microbiomes can vary substantially between and within coral species and across geographic locations[26, 40, 41]. The genotype of the host coral underlies some of this variation[40, 42, 43] as well as environmental factors such as water temperature[44], habitat type[45], and pollution levels[44, 46]. These factors can differ across the ranges of coral species, yet the scope of most studies is limited to a narrow selection of a species' known geographic or depth range. Further, most studies of microbiome variability have focused on shallow water corals, mainly scleractinians, and to a lesser degree on octocorals and antipatharian black corals[32, 41, 47]. In summary, the extent of coral microbiome variability is likely underestimated.

Variations in coral microbiomes can have functional implications. In shallow-water scleractinian species, certain microbial assemblages are associated with increased thermal tolerance[48-50]. Microbiome variation could be significant for mesophotic and deep-sea corals (those living deeper than 50 meters) because they often have wider depth and geographic ranges than shallow-water species[51].

Mesophotic and deep-sea corals support diverse and abundant biological communities in Earth's largest biome[52-56]. Most of these corals are heterotrophic because they live beyond the reach of sunlight sufficient for photosynthesis. Like shallow-water corals, mesophotic and deep-sea corals face human-induced environmental changes such as ocean warming, acidification, pollution, and ecosystem destruction [57-60]. In the mesophotic zone and deep sea, octocorals comprise approximately 70% of all coral species[61]. A comprehensive characterization of microbiome variation in octocorals across their geographic ranges is needed to understand the diversity and ecological roles of coral microbiomes in the mesophotic zone and the deep sea.

Mesophotic and deep-sea octocorals occur across a wide depth range down to at least 8,000m
and worldwide along continental margins and seamounts[51, 62]. Light intensity, oxygen
concentration, temperature, pressure, pH, salinity, food availability, and habitat structure vary
with depth and geography. All these factors combine to strongly influence the ecology and
evolution of marine animals[63-66]. Thus, factors that vary with depth and geographic location
likely drive microbiome variation in mesophotic and deep-sea octocorals.

Here we test the hypotheses that the microbiomes of mesophotic and deep-sea octocorals
vary significantly across their ranges and in correlation with environmental factors and host
genotype. We characterized the microbiomes of the mesophotic octocorals *Muricea pendula* and
*Swiftia exserta,* and the deep-sea octocorals *Callogorgia delta* and *Paramuricea biscaya* across
their known depth and geographic ranges in the northern Gulf of Mexico (**Fig. 1**). All of these
corals are dominant foundation species in their respective communities. At least three were
impacted by the Deepwater Horizon oil spill[57, 58, 67, 68] and are the focus of restoration efforts.
Understanding the geographic variation of their associated microbes is essential to understanding
the ecology of coral ecosystems and can also inform restoration plans.

**Methods**

*Sample Collection*

Octocorals were sampled between 2009 and 2019 from multiple sites across their ranges
in the northern Gulf of Mexico using Remotely Operated Vehicles (ROVs) or the human-
occupied vehicle Alvin, and placed in temperature-insulated containers during multiple research
cruises. Specimens of the mesophotic octocoral *Muricea pendula,* n = 140, were collected from
14 sites spanning depths between 53-117 m, and geographic distances between 34 and 690
kilometers. Specimens of the co-occurring mesophotic octocoral *Swiftia exserta,* n = 100, were
collected from 10 sites spanning depths between 62-100 m, and geographic distances between 13
and 600 kilometers. Specimens of the upper-bathyal octocoral *Callogorgia delta*, n = 60, were
collected from 6 sites spanning depths between 439-852 m and geographic distances between 14
and 350 kilometers. Finally, specimens of the lower-bathyal octocoral *Paramuricea biscaya*, n =
60, were collected from 6 sites spanning depths between 1370-2224 m, and geographic distances
between 14 and 655 kilometers. Upon ROV recovery, subsamples from each colony were frozen
with liquid nitrogen and/or in >95% ethanol and/or RNAlater. Details, including research cruise,
time of collection, and preservation method for each sample, can be found in **Supplementary**
**Table A**.

*DNA purification*

Whole DNA was purified from *ca.* ten coral colonies from each sampling site using a
modified salting-out procedure (dx.doi.org/10.17504/protocols.io.bypypvpw). In short, about
3mm$^3$ of coral tissue was digested in a lysis buffer with proteinase K and beta-mercaptoethanol,
protein and cellular debris were precipitated with 7.5M ammonium acetate, and DNA was
pelleted in 50% isopropanol, washed with 70% ethanol, and then resuspended in TE buffer. To
standardize DNA concentrations and reduce the concentration of PCR inhibitors, DNA extract
concentrations were measured on a Qubit 4.0 fluorometer with a qubit Broad Range assay and
then diluted with TE buffer to 2ng/µl for *Swiftia exserta*, 20ng/µL for *Muricea pendula*, 10ng/µL
for *Callogorgia delta*, and 10-40ng/µL for *Paramuricea biscaya*. DNA extracts of *C. delta* and
*P. biscaya* that were less concentrated than these thresholds were not diluted. See
**Supplementary Methods** for details.

*Microbial metabarcoding*

120
The hypervariable V1-V2 regions of the bacterial 16S rRNA gene were amplified
following Vohsen et al.[15] using the universal primers 27F and 355R (see **Supplementary**
**Methods** for details). PCR products were sent to the University of Illinois Chicago for library
preparation and sequencing[69] on an Illumina Miseq platform with 250bp paired-end sequencing.
Raw sequence data is available on the Sequence Read Archive (SRA) under BioProject ID
PRJNA875098.

Raw sequence data were imported into *qiime2*[70] ver2021.2 and processed following
Vohsen et al.[15] to create an amplicon sequence variant (ASV) table. In brief, reads were quality
filtered, joined, checked for chimeras, and ASVs were determined with deblur and classified
using a naïve-bayes classifier trained on the SILVA small subunit (SSU) ribosomal RNA
database (Ref NR 99 release 132)[71]. Details can be found in the **Supplementary Methods**.

*Coral genotyping*

To investigate the influence of population structures of the host, Restriction site-
Associated DNA sequence (RADseq) data were obtained from a subset of individuals as
described by Bracco et al.[64] and Galaska et al.[65] In brief, DNA was sent to Floragenex Inc.
(Eugene, OR) RAD for library preparation (PstI) and sequencing on an Illumina HiSeq platform.
Reads were demultiplexed and quality filtered using the *process_radtags* program in Stacks
v2.1[72]. Read clustering and single nucleotide polymorphism (SNP) calling were performed with
DeNovoGBS v4.0.1[73] for *C. delta* and *P. biscaya*, and with the gstacks v2.1[72] after mapping to a
genome assembly for *S. exserta* and *M. pendula*. We excluded SNP loci that: (1) had more than
10% of missing data, (2) had a minor allele frequency smaller than 0.01, or (3) were not biallelic.
We also excluded individuals if they had missing data in >10% of the SNP loci for *S. exserta* and
*M. pendula* and >30% for *C. delta* and *P. biscaya* or were identified as clones with R package
poppr v2.8.6[74]. To reduce the dimensionality of the genetic data, a principal component analysis
(PCA) was performed on the SNP data of each species separately (*S. exserta,* n = 54, 23,869
SNPs; *M. pendula,* n = 100, 9,611 SNPs; *C. delta*, n= 51, 14,013 SNPs, and *P. biscaya,* n = 58,
7,293 SNPs) using the R package ade4 v1.7[75]. The PCA axes were reduced to two dimensions
using the uniform manifold approximation and projection method[76] using the R package *umap*
v0.2.8. The two-dimensional coordinates were then used in statistical analyses to represent the
genotype of each coral.

*Environmental Variables*

Publicly available satellite data and model-derived values for the Gulf of Mexico were
obtained to evaluate the influence of environmental variables on microbial communities. The
satellite-derived estimates of surface primary productivity (mg/m$^2$d), chlorophyll concentration
(mg/m$^3$), inorganic suspended particulate matter concentration (g/m$^3$), and the particulate back-
scattering coefficient at 443nm (m$^{-1}$) were obtained from the Copernicus-GlobColour Global
Ocean Colour biogeochemical dataset. Model-derived values were obtained for the depth of the
surface mixed layer (m), bottom seawater temperature (°C), bottom seawater oxygen
concentration (mL/L), bottom seawater salinity, and bottom current magnitude (m/s) from the
HYbrid Coordinate Ocean Model (HYCOM). Both datasets comprised monthly means from Jan
2011 to Dec 2018 at 4km grid resolution. The mean and standard deviation of monthly means for
each variable were extracted at the sampling location of each coral colony. Only the mean
bottom seawater temperature and surface primary productivity were retained for statistical

analyses. Other variables were excluded from the analyses because they were strongly correlated,
or not substantially variable across sites, or were not justified by *a priori* hypotheses. See
**Supplementary Methods** for further details.

*Statistical Analyses*

Rarefied microbial ASV richness was calculated for each coral colony using the *rarefy*
function in the R package *vegan*[77] v2.5.7 using the minimum library size as the sampling depth
according to Hurlbert[78]. Pielou's evenness was calculated for each individual using the function
*evenness* in the R package *asbio*[79] v1.7 without including zeros.  Microbial community richness
and evenness were compared between coral species and across sites using paired two-tailed
Wilcoxon rank-sum tests and a Bonferroni-adjusted α-value with the function *wilcox.test* in the
base R package *stats* v4.1.2. In addition, the effects of depth and longitude on richness and
evenness were evaluated for each species with a linear model using depth and longitude as
predictors and richness or evenness as response variables independently.

Distance-based redundancy analysis[80] (dbRDA) was performed to evaluate the influence
of several factors on microbial community compositions including host species, host genotype,
depth, geography, bottom temperature, and surface primary productivity. Longitude was used to
test for regional scale patterns across geography while sampling site was used as a finer scale
variable. Models were constructed using the R command *dbrda* in the package *vegan* with Bray-
Curtis dissimilarities. The significance of the models and their predictor variables were tested
with the function *anova.cc*a using 9,999 permutations. A series of dbRDA models were
constructed and evaluated to test several hypotheses regarding the influence of each variable.
The details of each test are provided in **Supplementary Table F**. In addition, the diversity
among *Endozoicomonas* and corallicolids associated with *S. exserta* and *M. pendula* were
individually analyzed by applying the same dbRDA analyses described above. Relative
abundances were re-calculated for all ASVs classified as *Endozoicomonas* or corallicolids as
their relative abundance among each of those taxa. Coral samples without any representatives
from the taxon of focus were removed.

Individual ASVs whose abundance differed along depth, longitude, and across sites were
identified using Analysis of Compositions of Microbiomes with Bias Correction (ANCOM-
BC)[81] using the R packages *ancombc* v1.4.0 and *phyloseq* 1.38.0. Two separate ANCOM-BC
analyses were conducted on each species individually: 1) using both depth and longitude as
explanatory variables and 2) a global analysis using sampling sites as groups. ASVs were
excluded from the analysis if they were present in fewer than five individual corals (half the
colonies sampled at a single site). FDR was controlled using the Benjamini-Hochberg method.
Since the global analysis using sampling sites as groups depends on which site was used as the
reference, the analysis was repeated using each site as the reference. Only ASVs with a
maximum q-value below α=0.05 in all permutations were considered significantly different. In
general, results were consistent regardless of the reference site. However, a few sites had much
higher variation in microbial communities, resulting in many more significantly differential
ASVs when used as the reference site. These analyses were repeated on the family level instead
of ASVs to inform functional differences.

## Results

*Microbial communities are distinct among octocoral species*

Each coral species had a distinct microbial community in terms of composition and diversity (**Fig. 2A,B**; $R^2$ 43%, p≤$10^{-4}$). The microbiomes of *M. pendula* and *P. biscaya* had higher ASV richness and Pielou's evenness than *S. exserta* and *C. delta* (**Fig. 2C,D**; p≤5.4x$10^{-6}$). The differences in richness and evenness between species appeared to be driven by the prevalence of a few ASVs that dominated the microbial community of individual colonies (**Fig. 2B**). The microbiomes of all four coral species were dominated by at least one ASV that represented more than 50% of the microbial community in at least one coral colony. A higher percentage of *S. exserta* (94%) and *C. delta* (85%) colonies hosted a dominant ASV compared to *P. biscaya* (70%) and *M. pendula* (38%). *S. exserta* hosted only two dominant ASVs: one classified as BD1-7 clade (Spongiibacteraceae) and another classified as *Mycoplasma*. Both *C. delta* and *P. biscaya* hosted three dominant ASVs. All three dominant ASVs in *C. delta* were recently described[36] as novel members of the class Mollicutes: two *Ca. Oceanoplasma* and one *Ca. Thalassoplasma*. The dominant ASVs in *P. biscaya* were classified as *Mycoplasma*, *Endozoicomonas,* and *Pseudoalteromonas*. *M. pendula* had the greatest diversity of dominant ASVs including two *Mycoplasma*, two *Marinoplasma*, an *Endozoicomonas*, an unclassified alphaproteobacterium, and a corallicolid apicomplexan (**Fig. 2B**). Dominant ASVs belonging to the same taxon were given short identifiers (ex: M1, M2, S2) to distinguish which coral species they were associated with (first letter of coral genus) and the rank of abundance (1 being most abundant).

*Factors shaping microbial community composition*

The microbial communities of all coral species differed significantly between sites. Differences between sites accounted for 33 to 59% of the variation within species (**Supp. Table F**) and were driven by significant trends across both depth (p≤$10^{-4}$) and longitude (p≤0.0213), which could simultaneously account for 11 to 17% and 3 to 12% of the variation, respectively, depending on the coral species. The only exception was *C. delta,* whose microbial community composition did not significantly correlate with longitude. After accounting for depth and longitude, the sampling site was still associated with a significant (p≤0.012) and substantial amount of additional variation (14-48%) for each coral species.

Several RDAs were conducted with and without depth and longitude to investigate whether bottom temperature, primary productivity, and host genotype contributed to the trends with depth and longitude. While excluding depth, host genotype was significantly correlated with microbial community composition in all species (p≤0.0186) except *S. exserta* as was bottom temperature (p≤0.0261) except in *P. biscaya* in which it was not tested due to low variation between sites (4.3-4.8°C). Similarly, surface primary productivity was a significant predictor (p≤0.0114) in all species except *S. exserta*. After including all variables, depth and longitude still explained a significant amount of variation (p≤0.036) except depth in *C. delta*.

*Microbiome patterns with depth*

Depth was correlated with microbiome diversity in various ways. In *Muricea pendula,* both richness (**Fig. 2E**; p=6x$10^{-6}$) and evenness (**Fig. 2F**; p=0.012) decreased significantly with depth. The decrease in diversity in *M. pendula* was great enough that the average richness among colonies from the deepest sites was comparable to that observed in *S. exserta* and *C. delta*.

Conversely, both richness and evenness in *S. exserta* increased with depth (p<6x10$^{-6}$), as did
richness in *C. delta* (p<0.038).

The relative abundances of multiple microbial taxa strongly correlated with depth
including many dominant taxa (**Fig. 3A-D**). In *M. pendula*, *Endozoicomonas* ASV M1 increased
in abundance with depth and so did the total abundance of all *Endozoicomonas* ASVs. Similarly,
the abundance of the dominant corallicolid ASV M1 increased with depth in both *M. pendula*
and *S. exserta*. This was true for the total abundance of all corallicolid ASVs as well (**Fig. 3B,**
**Fig. S1A**). In contrast, other dominant ASVs decreased in abundance with depth: BD1-7 ASV
S1 in *S. exserta*, *Ca. Marinoplasma* ASV M1 in *M. pendula,* and both *Ca. Oceanoplasma* ASV
C1 and *Ca. Thalassoplasma* C1 in *C. delta*.

In addition, many other taxa that comprised a smaller proportion of the microbial
community differed with depth (**Fig. 3E**). *M. pendula* hosted most of these and the majority
decreased in abundance with depth. The most abundant of these non-dominant depth-decreasing
taxa included photosynthetic groups such as Cyanobiaceae and chloroplasts but also ASVs in the
families Spirochaetaceae, Helicobacteraceae, Flavobacteriaceae, Francisellaceae,
Burkholderiaceae, and unclassified bacteria that clustered with Spirochaetia and
Epsilonproteobacteria. Together with the dominant *Ca. Marinoplasma*, these taxa comprised a
mean of 70% of the microbial community among colonies from the shallowest site (Sonnier
Bank, 55 m) and decreased down to a mean total of 3% at the deepest site (Bright Bank, 116 m).
Although many of these taxa were also detected in *S. exserta,* their abundances were either not
significantly correlated with depth or increased with depth. In *C. delta,* the abundances of two
sulfur-oxidizing taxa increased with depth: SUP05 and members of the novel family
Thiobarbaceae.

In addition to microbial taxa differing in abundance with depth, their ASV compositions
also differed. For example, in *S. exserta,* the composition of corallicolid ASVs differed across
depth. (**Fig. S1K**, p≤10$^{-4}$). Notably, *M. pendula* and *S. exserta* each hosted a unique plastotype at
the deep site Elvers Bank (96 m). Similarly, the composition of *Endozoicomonas* ASVs in *M.*
*pendula* differed with depth (**Fig. 3F**; p≤10$^{-4}$). Similar transitions of ASVs within microbial taxa
were also prevalent among the associates of *C. delta*. *Ca. Oceanoplasma* ASV C1 dominated the
microbial community of most *C. delta* colonies from most sites except the deepest two sites
(GC249 and GC290), where the closely related *Ca. Oceanoplasma* ASV C2 dominated some
colonies (**Fig. 2B**).  This pattern was also seen among *Endozoicomonas* and *Shewanella*
associates of *C. delta* (**Fig. 3G,H**). A single common *Endozoicomonas* ASV was present in most
colonies from all sites (dark gray), while other ASVs were mostly restricted between three depth
zones. More strikingly, *Shewanella* communities between colonies from two depth zones
(<637m and >794m) were completely non-overlapping, failing to share a single ASV.

*Microbiome patterns with geography*
Both richness and evenness were higher in eastern than western sites for both *M. pendula*
and *S. exserta* (p<0.0221). Multiple microbes exhibited differences in abundance across the
geographic range of their host coral on a regional scale along the east-west axis (**Fig. 4A-F**). *M.*
*pendula* colonies from eastern sites had higher abundances of microbes that may be free-living
such as ASVs classified as *Alteromonas, Pseudoalteromonas, Pseudomonas, Marinobacter,*
*Ruegeria, Caedibacter,* Sphingomonadaceae, and photosynthetic taxa such as Cyanobiaceae but
also microbes that may be symbiotic such as individual *Endozoicomonas* ASVs, corallicolids,

BD1-7, Spirochaetaceae, and the dominant *Mycoplasma* ASV M2. The mean total relative abundance of all ASVs that were more abundant in the east ranged between 0.8% and 26%, excluding the dominant *Mycoplasma* ASV M2 and dominant corallicolid plastotype. Some of these same ASVs and other closely related taxa were also more abundant in *S. exserta* in the east including *Pseudomonas, Pseudoalteromonas, Alteromonas, Ruegeria, Endozoicomonas,* Cyanobiaceae, BD1-7, and Spirochaetaceae (**Supplementary File 1**).

Other ASVs from the same microbial taxa were more abundant in *M. pendula* from western sites such as some classified as Cyanobiaceae, *Mycoplasma,* and *Endozoicomonas* (**Fig. 4C,D**). In addition, ASVs from other taxa were abundant in the west such as several *Ca. Marinoplasma* ASVs and photosynthetic taxa including several chloroplasts. Notably, two ASVs classified as *Prochloron* comprised up to a total of over 50% of the microbial community in some corals from adjacent sites in the west (**Fig. 4C**). Conversely, only two ASVs associated with *S. exserta* were significantly more abundant in the west including an unclassified gamma-proteobacterium and the dominant *Mycoplasma* ASV S1 which was completely absent from all samples from the three easternmost sites (**Fig. 4F**).

*Remaining Site Scale Geographic variation*

Many microbial taxa were not correlated with depth nor longitude but still differed in abundance across sites. In *P. biscaya,* there were no consistent patterns with depth nor longitude in the abundance of individual ASVs. However, there were substantial differences between sites (**Fig. 4G**). For instance, SUP05 had a high abundance in colonies from GC852, *Endozoicomonas* ASV P1 was abundant at DC673 and KC405, *Pseudoalteromonas* ASV P1 was abundant at MC297, and *Mycoplasma* ASV P1 was abundant at most sites but ranged from a mean of 0.1% to 74% across sites.

Microbes whose abundance differed between sites followed two general patterns. First, some microbes were present in most colonies from most sites but differed consistently in abundance across sites, such as the dominant *Mycoplasma* ASVs in *P. biscaya* (**Fig. 4G**)*,* and ASVs classified as Gracilibacteria in *C. delta*. Second, many other microbes were only detected in colonies from a few sites or even a single site. This included a corallicolid only detected in *C. delta* colonies from VK826, GC234, and MC885 and another corallicolid only detected in colonies from GC234 (**Fig. S2G,H**). Other examples involved abundant ASVs such as three close relatives of *Endozoicomonas* in *M. pendula* at McNeil bank averaging a total relative abundance of 26% (**Fig. S2E**). Less abundant examples include potential parasites within Rickettsiales in all coral species.

## Discussion:

*Differences Between Coral Species*

Coral species was the strongest factor differentiating microbial communities. Species differed markedly in which microbes they associated with and also the diversity of their microbial communities. This finding is consistent with previous coral studies[1, 27, 41, 82, 83]. All dominant ASVs were only abundant (over X%) in one coral species except some corallicolids which were found in both *S. exserta* and *M. pendula* where they totaled X% and Y% on average. The microbial taxa that dominated some coral species belonged to different phyla, such as BD1-7 in *S. exserta* and *Ca. Oceanoplasma* in *C. delta,* suggesting that their metabolic functions differ between coral species. However, some taxa such as *Endozoicomonas* and members of the

Mollicutes were associates of multiple species in this study and other studies suggesting that they
may serve common functions across a wide range of coral species. The differences in community
structure observed between these four coral species may reflect their unique ecologies.
Differences in diversity have been proposed to reflect different ecological strategies[84]. Some
corals may be microbial generalists that foster more diverse and flexible microbial communities,
like those of *M. pendula* and *P. biscaya,* to respond to changing environmental conditions, while
other corals associate with less diverse, more stable, and specialized communities like *S. exserta*
and *C. delta*[83, 84] Alternatively, *S. exserta* and *C. delta* may have higher absolute abundances of
their dominant associates reducing the richness and evenness of their communities. Measuring
absolute microbial abundances would aid the interpretation of differences in richness and
evenness between species.

*The Influence of Depth*

Depth strongly structured the microbial communities of all coral species investigated
which exhibited differences in composition, diversity, and the abundance of specific microbial
ASVs and families. The number and variety of taxon-specific patterns with depth reflect the
likelihood that depth structures microbial communities through multiple mechanisms.

For instance, some variation in microbial communities may be due to temperature.
Temperature decreases with ocean depth down to ~1500m in the Gulf of Mexico.  Depth and
temperature were correlated with community composition in all species shallower than this
depth. Other studies have shown that shallow-water, scleractinian corals from locations with
different temperature regimes host different microbial communities in both the Caribbean and
the Red seas[48, 85]. Specifically, the stress of high temperatures has been shown to shape the
microbial communities associated with corals[86], especially if they bleach[87-90]. Thus, for the coral
species in our study, the microbiomes of colonies may differ across depth in correlation with
differing thermal regimes because their host corals experience different frequencies or intensities
of thermally induced stress. While temperature may be the underlying cause of some of the
variation with depth, it does not appear to be the only factor since depth correlated with
additional variation after accounting for temperature.

The genetic divergence of coral populations may further contribute to differentiation in
microbial communities across depths. In the scleractinian *Pocillopora damicornis,* genetically
distinct coral populations have been shown to harbor distinct microbial communities[43]. In our
study, the genotype of coral hosts was significantly correlated with microbial community
structure, but only when the correlates of depth and temperature were ignored. While the
potential influence of host genotype is difficult to distinguish from correlates with depth, the
turnover of the *Shewanella* community is consistent with the population genetic structuring of *C.*
*delta*[63, 64]. Coral colonies from the deepest two sites were genetically distinct and hosted unique
*Shewanella* communities. Thus, some of the differentiation in microbial communities across
depth may be caused by genetic differentiation of the coral host.

Some depth patterns may reflect habitat type. For instance, members of the SUP05
cluster and the family Thiobarbaceae increased in abundance with depth in *C. delta*. These
microbes are sulfur-oxidizers associated with chemosymbiotic fauna[91] and may reflect the
greater prevalence of seeps along depth in the northern Gulf of Mexico. One of these microbes,
SUP05, has been shown to correlate with markers of chemoautotrophic nutrition in corals[92] and
thus may reflect an increase in chemoautotrophic food sources and a decrease in the abundance

of marine snow with depth. Other patterns with depth appear to be driven by different causes.
For instance, the diversity of microbial associates of *M. pendula* decreased continuously with
depth (Fig. 2E,F). The decrease in the abundance of bacterioplankton with depth could account
for some of this. For example, the abundance of likely free-living bacteria found in *M. pendula*,
such as *Serratia* and *Achromobacter*, decreased with depth. Further, potentially photosynthetic
taxa associated with *M. pendula* predictably decreased with depth, likely reflecting decreasing
light levels.

Altogether, depth structures microbial communities through multiple mechanisms,
including changes in free-living microbial communities, decreasing light levels, differentiation of
host corals, and transitions in habitat types.

Many microbial associates that may be symbiotic differed in abundance with depth, such
as *Spirochaeta 2, Ca. Marinoplasma, Endozoicomonas, Mycoplasma, Ca. Oceanoplasma, Ca.*
*Thalassoplasma,* and corallicolid apicomplexans. The reasons for this are unclear because many
of these microbe's interactions with their coral hosts are not well characterized. Interestingly,
corallicolids increased in abundance with depth in both *M. pendula* and *S. exserta,* but no depth
pattern was found in corallicolids that infect *C. delta*. This is in contrast to previous studies that
found a decrease in abundance along depth within the range of a shallow-water coral and across
a depth range of 2 kilometers[15, 93]. These inconsistencies may be explained if corallicolids utilize
intermediate host species like some other apicomplexans and if the distributions of these
potential intermediate host species differ with depth in various ways. Some of the closest
relatives of corallicolids are found in fish[94-96], opening the possibility that corallicolids may use
fish as intermediate hosts and are limited by their depth ranges.

*Variation Across Geography*

In addition to variation across depth, we found that the microbial communities of all four
species differed geographically. Corals from different sites differed strongly in microbial
communities in all species. but nearby sites showed similarities creating regional-scale
geographic variation.

*Regional scale geographic variation*

Regional scale variation was identified in both mesophotic coral species. Both *M.*
*pendula* and *S. exserta* had higher diversity near the mouth of the Mississippi River, and many
likely free-living microbial taxa were detected at higher abundances, such as *Alteromonas*,
*Pseudoalteromonas*, *Pseudomonas*, and photosynthetic taxa. Their increased abundance may
reflect the influence of the Mississippi River on bacterioplankton in the region. Further, there are
marked longitudinal differences in surface primary productivity in the Northern Gulf of Mexico,
higher in the area of influence of the Mississippi. Microbial communities of other shallow-water
corals seem to be significantly influenced by surface primary productivity[97]. This is consistent
with our observation that microbial communities of octocorals were significantly correlated with
surface primary productivity.

However, taxa that are symbiotic, like *Endozoicomonas,* Mollicutes, and corallicolids
also differed in abundance regionally consistent with other studies that found geographic
variation in *Endozoicomonas* and *Mycoplasma* associates in shallow water corals[21, 26]. Some of
the regional patterns in these potentially symbiotic taxa may result from limited dispersal since

some were restricted to nearby sites like *Mycoplasma* ASV S1, which was completely absent in
all corals from the three easternmost sites.

*Site scale geographic variation*

For every coral species, microbial communities differed strongly between sites after
accounting for depth and geographic region. This is likely due to differences in local conditions
between sites that influence microbial community structure. For instance, the high abundance of
the SUP05 cluster in *P. biscaya* at GC852 may reflect the presence of seeps nearby[98] (< 450 m).
Similarly, the abundance of *Pseudoalteromonas* in *P. biscaya* at the Mississippi Canyon sites
may reflect the lasting impact of the Deepwater Horizon oil spill[58, 68, 99]. Consistent deviations
from depth trends at certain sites suggest that other factors in addition to depth play a role in
determining the abundance of certain groups, such as the total corallicolid abundance in *M.*
*pendula* at Roughtongue Reef which was higher than the trend with depth (Fig 3B). Conversely,
corals from some subsets of sites displayed similar microbial communities despite being far apart
and inhabiting different depths. For instance, *P. biscaya* from both KC405 and DC673 were
dominated by *Endozoicomonas* ASV P1, but it comprised no more than 0.12% in any colony
from any other site. These sites are the farthest apart and do not have overlapping depths.
Interestingly, corals from these two populations are closely related[65] suggesting an influence of
the coral host's genotype on microbiome composition.

Implications for restoration efforts

Characterizations of coral microbiomes based on samples from a single site or even a few
may miss substantial variation across a species' range. This is important to consider in
restoration plans that utilize collections from different locations. For example, a restoration plan
could use this information to avoid inadvertently transmitting parasites to new sites through
transplants or to avoid rearing infected and non-infected corals together in aquaria. Examples of
geographic variation in potential parasites discovered in this work include corallicolids in *C.*
*delta* at VK826, MC885, and GC234 and the unique corallicolids in *M. pendula* and *S. exserta* at
Elvers Bank.

Conclusion:
This work characterizes the microbial communities of four dominant coral species in the
Gulf of Mexico and demonstrates substantial variability across their ranges. These results
provide promising targets to study the interactions between corals and their associated microbes.
They also reveal substantial variability associated with depth and geography that could only be
uncovered through extensive sampling. The numerous patterns associated with depth
demonstrate that in addition to its ecological and evolutionary effects on corals, it also strongly
structures their microbial communities.

Acknowledgements:

This research was funded by the NOAA's National Centers for Coastal Ocean Science,
Competitive Research Program, the Office of Ocean Exploration and Research, and the
RESTORE Science Program under awards NA18NOS4780166 and NA17NOS4510096 to
Santiago Herrera at Lehigh University. Santiago Herrera was also supported by the National

Competing Interests:

The authors declare no competing interests.

Data Availability Statement:

16S amplicon sequence data were submitted to the Sequence Read Archive (SRA) under
Bioproject number PRJNA875098.

Figures:

Fig. 1: Sampling sites and their environmental and coral genotypic makeup. (A) In situ mages of
the octocoral species in this study. (B) Geographic map displaying sampling locations in the
Northern Gulf of Mexico (BOEM bathymetry). The mean seafloor depth of all samples from a
site is reported next to each site name in meters. Sites are colored based on which coral species
were sampled: light gray = *Muricea pendula,* red = *Swiftia exserta,* blue = *Callogorgia delta,*
yellow = *Paramuricea biscaya.* Images of *C. delta* and *P. biscaya* courtesy of the ECOGIG
consortium. (C) Primary productivity of each sampling site displayed across longitude and
colored by coral species sampled. (D) Bottom seawater temperature for each sampling site
displayed across depth for each coral species. (E) UMAP projections of the genetic background
of coral colonies based on RADseq data. Each point represents the genotype of a coral colony
and is colored by depth in meters.

Fig. 2: Characterization of coral microbiomes. A) Non-metric multidimensional scaling plot of
microbiome compositions using Bray-Curtis dissimilarities of ASV proportions. B) Barplots of
dominant ASVs in each coral species. Each horizontal bar is an individual coral colony. Colonies
are grouped by site which are sorted by depth. ASVs that comprised 50% or more in any sample
are designated with different colors. All other ASVs are collapsed and shown in light gray. C)
Rarefied ASV richness and D) Pielou's evenness between different coral species. MP = *Muricea
pendula,* SE = *Swiftia exserta,* CD = *Callogorgia delta,* PB = *Paramuricea biscaya.* E) Rarefied
ASV richness and F) Pielou's evenness in *Muricea pendula* with depth.

Fig. 3: Examples of depth patterns in coral microbiome composition. The relative abundances along depth in *Muricea pendula* of A) the sum of all *Endozoicomonas* ASVs, B) the sum of all corallicolid plastotypes, D) *Ca. Marinoplasma* ASV M1, E) the sum of all microbial families that decreased significantly with depth. F) NMDS ordination of *Endozoicomonas* communities in *M. pendula* showing patterns with depth. C) Relative abundance of BD1-7 ASV S1 with depth in *Swiftia exserta.* The community composition of G) *Endozoicomonas* and H) *Shewanella* communities in *Callogorgia delta.* Each horizontal bar is a single coral colony whose composition reflects the relative abundances of each ASV when restricted to only *Endozoicomonas* or *Shewanella* ASVs. Colors denote different ASVs. Colonies are arranged by site in order of increasing mean depth.

Fig. 4: Examples of geographic patterns in coral microbiome composition. Barplots of the
relative abundance of microbial ASVs from *Muricea pendula* that differed significantly across
longitude and were associated with A) eastern and C) western areas of its range in the Northern
Gulf of Mexico. ASVs are colored by family or other taxonomic level detailed in the legend.
Each coral colony is represented by a vertical bar, and colonies are grouped by site sorted from
westernmost to easternmost starting from the left. The total relative abundances of these ASVs
are shown in B) and D) plotted against longitude to better reflect relative distances between sites.
The relative abundance of E) *Mycoplasma* ASV M1 in *M. pendula* and F) *Mycoplasma* ASV S1
in *S. exserta* along longitude. Samples where zero reads were detected are represented with open
circles. G) Barplots showing the five most abundant ASVs that differed between sites in *P.*
*biscaya.*

References:

1.     Pollock, F.J., McMinds, R., Smith, S., Bourne, D.G., Willis, B.L., Medina, M., Thurber,
520           R.V. & Zaneveld, J.R. Coral-associated bacteria demonstrate phylosymbiosis and
521           cophylogeny. *Nature Communications* **9**, 4921 (2018).
522    2.     Siboni, N., Ben☐Dov, E., Sivan, A. & Kushmaro, A. Global distribution and diversity of
523           coral☐associated Archaea and their possible role in the coral holobiont nitrogen cycle.
524           *Environmental Microbiology* **10**, 2979-2990 (2008).
525    3.     Wegley, L., Yu, Y., Breitbart, M., Casas, V., Kline, D.I. & Rohwer, F. Coral-associated
526           archaea. *Marine Ecology Progress Series* **273**, 89-96 (2004).
527    4.     Ainsworth, T., Fordyce, A. & Camp, E. The other microeukaryotes of the coral reef
528           microbiome. *Trends in Microbiology* **25**, 980-991 (2017).
529    5.     Thurber, R.V., Payet, J.P., Thurber, A.R. & Correa, A. Virus–host interactions and their
530           roles in coral reef health and disease. *Nature Reviews Microbiology* **15**, 205-216 (2017).
531    6.     Yang, S.H., Lee, S.T., Huang, C.R., Tseng, C.H., Chiang, P.W., Chen, C.P., Chen, H.J. &
532           Tang, S.L. Prevalence of potential nitrogen☐fixing, green sulfur bacteria in the skeleton
533           of reef☐building coral I sopora palifera. *Limnology and oceanography* **61**, 1078-1086
534           (2016).
535    7.     Ricci, F., Rossetto Marcelino, V., Blackall, L.L., Kühl, M., Medina, M. & Verbruggen,
536           H. Beneath the surface: community assembly and functions of the coral skeleton
537           microbiome. *Microbiome* **7**, 1-10 (2019).
538    8.     Wada, N., Hsu, M.-T., Tandon, K., Hsiao, S.S.-Y., Chen, H.-J., Chen, Y.-H., Chiang, P.-
539           W., Yu, S.-P., Lu, C.-Y. & Chiou, Y.-J. High-resolution spatial and genomic
540           characterization of coral-associated microbial aggregates in the coral Stylophora
541           pistillata. *Science Advances* **8**, eabo2431 (2022).
542    9.     La Rivière, M., Garel, M. & Bally, M. Localization of endobacteria in the gastrodermis
543           of a Mediterranean gorgonian coral, *Paramuricea clavata*, using fluorescence in situ
544           hybridization. *Marine Biology* **163**, 206 (2016).
545    10.    Shapiro, O.H., Kramarsky-Winter, E., Gavish, A.R., Stocker, R. & Vardi, A. A coral-on-
546           a-chip microfluidic platform enabling live-imaging microscopy of reef-building corals.
547           *Nature communications* **7**, 1-10 (2016).
548    11.    Kim, B.R. Regulation of microbial populations by coral surface mucus and mucus-
549           associated bacteria. *Marine Ecology Progress Series* **322**, 1-14 (2006).
550    12.    Klinges, G., Maher, R.L., Vega Thurber, R.L. & Muller, E.M. Parasitic 'Candidatus
551           Aquarickettsia rohweri' is a marker of disease susceptibility in Acropora cervicornis but
552           is lost during thermal stress. *Environmental Microbiology* **22**, 5341-5355 (2020).
553    13.    Lesser, M.P., Falcón, L.I., Rodrí-guez-Román, A., Enr-íquez, S., Hoegh-Guldberg, O. &
554           Iglesias-Prieto, R. Nitrogen fixation by symbiotic cyanobacteria provides a source of
555           nitrogen for the scleractinian coral Montastraea cavernosa. *Marine Ecology Progress*
556           *Series* **346**, 143-152 (2007).
557    14.    Lesser, M.P., Mazel, C.H., Gorbunov, M.Y. & Falkowski, P.G. Discovery of Symbiotic
558           Nitrogen-Fixing Cyanobacteria in Corals. *Science* **305**, 997-1000 (2004).
559    15.    Vohsen, S.A., Anderson, K.E., Gade, A.M., Gruber-Vodicka, H.R., Dannenberg, R.P.,
560           Osman, E.O., Dubilier, N., Fisher, C.R. & Baums, I.B. Deep-sea corals provide new

insight into the ecology, evolution, and the role of plastids in widespread apicomplexan symbionts of anthozoans. *Microbiome* **8**, 34 (2020).

16. Kwong, W.K., del Campo, J., Mathur, V., Vermeij, M.J.A. & Keeling, P.J. A widespread coral-infecting apicomplexan with chlorophyll biosynthesis genes. *Nature* **568**, 103-107 (2019).

17. Mathur, V., del Campo, J., Kolisko, M. & Keeling, P.J. Global diversity and distribution of close relatives of apicomplexan parasites. *Environmental Microbiology* **20**, 2824-2833 (2018).

18. Kirk, N.L., Ritson-Williams, R., Coffroth, M.A., Miller, M.W., Fogarty, N.D. & Santos, S.R. Tracking Transmission of Apicomplexan Symbionts in Diverse Caribbean Corals. *Plos One* **8**, 10 (2013).

19. Kirk, N.L., Thornhill, D.J., Kemp, D.W., Fitt, W.K. & Santos, S.R. Ubiquitous associations and a peak fall prevalence between apicomplexan symbionts and reef corals in Florida and the Bahamas. *Coral Reefs* **32**, 847-858 (2013).

20. Weiler, B.A., Verhoeven, J.T.P. & Dufour, S.C. Bacterial Communities in Tissues and Surficial Mucus of the Cold-Water Coral *Paragorgia arborea*. *Frontiers in Marine Science* **5** (2018).

21. Woo, S., Yang, S.-H., Chen, H.-J., Tseng, Y.-F., Hwang, S.-J., De Palmas, S., Denis, V., Imahara, Y., Iwase, F., Yum, S. & Tang, S.-L. Geographical variations in bacterial communities associated with soft coral *Scleronephthya gracillimum*. *PLOS ONE* **12**, e0183663 (2017).

22. Holm, J.B. & Heidelberg, K.B. Microbiomes of Muricea californica and M. fruticosa: Comparative Analyses of Two Co-occurring Eastern Pacific Octocorals. *Frontiers in Microbiology* **7** (2016).

23. Kellogg, C.A., Lisle, J.T. & Galkiewicz, J.P. Culture-Independent Characterization of Bacterial Communities Associated with the Cold-Water Coral *Lophelia pertusa* in the Northeastern Gulf of Mexico. *Applied and Environmental Microbiology* **75**, 2294-2303 (2009).

24. Neulinger, S.C., Gärtner, A., Järnegren, J., Ludvigsen, M., Lochte, K. & Dullo, W.-C. Tissue-Associated "*Candidatus* Mycoplasma corallicola" and Filamentous Bacteria on the Cold-Water Coral *Lophelia pertusa* (Scleractinia). *Applied and Environmental Microbiology* **75**, 1437-1444 (2009).

25. Gray, M.A., Stone, R.P., McLaughlin, M.R. & Kellogg, C.A. Microbial consortia of gorgonian corals from the Aleutian islands. *FEMS Microbiology Ecology* **76**, 109-120 (2011).

26. Neave, M.J., Rachmawati, R., Xun, L., Michell, C.T., Bourne, D.G., Apprill, A. & Voolstra, C.R. Differential specificity between closely related corals and abundant Endozoicomonas endosymbionts across global scales. *The ISME Journal* **11**, 186-200 (2017).

27. van de Water, J.A.J.M., Voolstra, C.R., Rottier, C., Cocito, S., Peirano, A., Allemand, D. & Ferrier-Pagès, C. Seasonal Stability in the Microbiomes of Temperate Gorgonians and the Red Coral *Corallium rubrum* Across the Mediterranean Sea. *Microbial Ecology* **75**, 274-288 (2018).

28. van de Water, J.A.J.M., Melkonian, R., Voolstra, C.R., Junca, H., Beraud, E., Allemand, D. & Ferrier-Pagès, C. Comparative Assessment of Mediterranean Gorgonian-Associated

Microbial Communities Reveals Conserved Core and Locally Variant Bacteria.
*Microbial Ecology* **73**, 466-478 (2017).

29.   Bayer, T., Neave, M.J., Alsheikh-Hussain, A., Aranda, M., Yum, L.K., Mincer, T.,
Hughen, K., Apprill, A. & Voolstra, C.R. The microbiome of the Red Sea coral
Stylophora pistillata is dominated by tissue-associated Endozoicomonas bacteria. *Applied*
*and environmental microbiology* **79**, 4759-4762 (2013).

30.   Keller-Costa, T., Kozma, L., Silva, S.G., Toscan, R., Gonçalves, J., Lago-Lestón, A.,
Kyrpides, N.C., Rocha, U.N. & Costa, R. Evidence for cross-feeding, metabolic
specialization, and niche partitioning in the octocoral holobiont. (2022).

31.   McCauley, E.P., Haltli, B., Correa, H. & Kerr, R.G. Spatial and temporal investigation of
the microbiome of the Caribbean octocoral *Erythropodium caribaeorum*. *FEMS*
*Microbiology Ecology* **92** (2016).

32.   Dannenberg, R.P. Characterization and Oil Response of the Deep Sea Coral-associated
Microbiome. (2015).

33.   Pogoreutz, C., Oakley, C.A., Rädecker, N., Cárdenas, A., Perna, G., Xiang, N., Peng, L.,
Davy, S.K., Ngugi, D.K. & Voolstra, C.R. Coral holobiont cues prime Endozoicomonas
for a symbiotic lifestyle. *The ISME Journal*, 1-13 (2022).

34.   Neave, M.J., Michell, C.T., Apprill, A. & Voolstra, C.R. *Endozoicomonas* genomes
reveal functional adaptation and plasticity in bacterial strains symbiotically associated
with diverse marine hosts. *Scientific Reports* **7**, 40579 (2017).

35.   Tandon, K., Lu, C.-Y., Chiang, P.-W., Wada, N., Yang, S.-H., Chan, Y.-F., Chen, P.-Y.,
Chang, H.-Y., Chiou, Y.-J. & Chou, M.-S. Comparative genomics: dominant coral-
bacterium Endozoicomonas acroporae metabolizes dimethylsulfoniopropionate (DMSP).
*The ISME journal* **14**, 1290-1303 (2020).

36.   Vohsen, S.A., Gruber-Vodicka, H.R., Meyer, M., Sadowski, M., Dubilier, N., Fisher,
C.R. & Baums, I.B. The deep-sea coral, Callogorgia delta, associates with bacteria
belonging to a novel marine branch of the Mollicutes. *bioRxiv* (2022).

37.   Morrison, D.A. Evolution of the Apicomplexa: where are we now? *Trends in*
*parasitology* **25**, 375-382 (2009).

38.   Razin, S. & Hayflick, L. Highlights of mycoplasma research—an historical perspective.
*Biologicals* **38**, 183-190 (2010).

39.   Toller, W.W., Rowan, R. & Knowlton, N. Genetic evidence for a protozoan (phylum
Apicomplexa) associated with corals of the *Montastraea annularis* species complex.
*Coral Reefs* **21**, 143-146 (2002).

40.   Dubé, C.E., Ziegler, M., Mercière, A., Boissin, E., Planes, S., Bourmaud, C.A.F. &
Voolstra, C.R. Naturally occurring fire coral clones demonstrate a genetic and
environmental basis of microbiome composition. *Nature Communications* **12**, 6402
(2021).

41.   van de Water, J.A.J.M., Coppari, M., Enrichetti, F., Ferrier-Pagès, C. & Bo, M. Local
Conditions Influence the Prokaryotic Communities Associated With the Mesophotic
Black Coral Antipathella subpinnata. *Frontiers in Microbiology* **11** (2020).

42.   Durante, M.K., Baums, I.B., Williams, D.E., Vohsen, S. & Kemp, D.W. What drives
phenotypic divergence among coral clonemates of Acropora palmata? *Molecular Ecology*
**28**, 3208-3224 (2019).

43.  van Oppen, M.J.H., Bongaerts, P., Frade, P., Peplow, Lesa M., Boyd, S.E., Nim, H.T. &
Bay, L.K. Adaptation to reef habitats through selection on the coral animal and its
associated microbiome. *Molecular Ecology* **27**, 2956-2971 (2018).
44.  Wang, L., Shantz, A.A., Payet, J.P., Sharpton, T.J., Foster, A., Burkepile, D.E. &
Thurber, R.V. Corals and Their Microbiomes Are Differentially Affected by Exposure to
Elevated Nutrients and a Natural Thermal Anomaly. *Frontiers in Marine Science* **5**, 16
(2018).
45.  Roder, C., Bayer, T., Aranda, M., Kruse, M. & Voolstra, C.R. Microbiome structure of
the fungid coral Ctenactis echinata aligns with environmental differences. *Molecular
Ecology* **24**, 3501-3511 (2015).
46.  Klinges, J.G., Patel, S.H., Duke, W.C., Muller, E.M. & Vega Thurber, R.L. Phosphate
enrichment induces increased dominance of the parasite Aquarickettsia in the coral
Acropora cervicornis. *FEMS Microbiology Ecology*, fiac013 (2022).
47.  van de Water, J.A.J.M., Allemand, D. & Ferrier-Pagès, C. Host-microbe interactions in
octocoral holobionts - recent advances and perspectives. *Microbiome* **6**, 64 (2018).
48.  Osman, E.O., Suggett, D.J., Voolstra, C.R., Pettay, D.T., Clark, D.R., Pogoreutz, C.,
Sampayo, E.M., Warner, M.E. & Smith, D.J. Coral microbiome composition along the
northern Red Sea suggests high plasticity of bacterial and specificity of endosymbiotic
dinoflagellate communities. *Microbiome* **8**, 1-16 (2020).
49.  Osman, E.O., Smith, D.J., Ziegler, M., Kürten, B., Conrad, C., El☐Haddad, K.M.,
Voolstra, C.R. & Suggett, D.J. Thermal refugia against coral bleaching throughout the
northern Red Sea. *Glob. Change Biol.* **24**, e474-e484 (2018).
50.  Rosado, P.M., Leite, D.C.A., Duarte, G.A.S., Chaloub, R.M., Jospin, G., Nunes da
Rocha, U., P. Saraiva, J., Dini-Andreote, F., Eisen, J.A., Bourne, D.G. & Peixoto, R.S.
Marine probiotics: increasing coral resistance to bleaching through microbiome
manipulation. *The ISME Journal* **13**, 921-936 (2019).
51.  Roberts, J.M., Wheeler, A.J., Freiwald, A. & Cairns, S.D. *Cold-Water Corals The
Biology and Geology of Deep-Sea Coral Habitats*. (Cambridge University Press, New
York; 2009).
52.  Jensen, A. & Frederiksen, R. The fauna associated with the bank-forming deepwater
coral Lophelia pertusa (Scleractinaria) on the Faroe shelf. *Sarsia* **77**, 53-69 (1992).
53.  Buhl-Mortensen, L. & Mortensen, P.B. Distribution and diversity of species associated
with deep-sea gorgonian corals off Atlantic Canada. *Cold-water corals and ecosystems*,
849-879 (2005).
54.  Sinniger, F., Ballantine, D., Bejarano, I., Colin, P., Pochon, X., Pomponi, S., Puglise, K.,
Pyle, R., Reaka, M. & Spalding, H. Biodiversity of mesophotic coral ecosystems.
*Mesophotic coral ecosystems—a lifeboat for coral reefs*, 50-62 (2016).
55.  Pinheiro, H., Goodbody-Gringley, G., Jessup, M., Shepherd, B., Chequer, A. & Rocha, L.
Upper and lower mesophotic coral reef fish communities evaluated by underwater visual
censuses in two Caribbean locations. *Coral Reefs* **35**, 139-151 (2016).
56.  Henderson, M.J., Huff, D.D. & Yoklavich, M.M. Deep-sea coral and sponge taxa
increase demersal fish diversity and the probability of fish presence. *Frontiers in Marine
Science* **7**, 593844 (2020).
57.  Silva, M., Etnoyer, P.J. & MacDonald, I.R. Coral injuries observed at mesophotic reefs
after the Deepwater Horizon oil discharge. *Deep Sea Research Part II: Topical Studies in
Oceanography* **129**, 96-107 (2016).

58.  White, H.K., Hsing, P.Y., Cho, W., Shank, T.M., Cordes, E.E., Quattrini, A.M., Nelson,
R.K., Camilli, R., Demopoulos, A.W.J., German, C.R., Brooks, J.M., Roberts, H.H.,
Shedd, W., Reddy, C.M. & Fisher, C.R. Impact of the Deepwater Horizon oil spill on a
deep-water coral community in the Gulf of Mexico. *Proc. Natl. Acad. Sci. U. S. A.* **109**,
20303-20308 (2012).
59.  Guinotte, J.M., Orr, J., Cairns, S., Freiwald, A., Morgan, L. & George, R. Will
human☐induced changes in seawater chemistry alter the distribution of deep☐sea
scleractinian corals? *Frontiers in Ecology and the Environment* **4**, 141-146 (2006).
60.  Hoegh-Guldberg, O., Poloczanska, E.S., Skirving, W. & Dove, S. Coral reef ecosystems
under climate change and ocean acidification. *Frontiers in Marine Science* **4**, 158 (2017).
61.  Cairns, S.D. Deep-water corals: an overview with special reference to diversity and
distribution of deep-water scleractinian corals. *Bulletin of marine Science* **81**, 311-322
(2007).
62.  Yesson, C., Taylor, M.L., Tittensor, D.P., Davies, A.J., Guinotte, J., Baco, A., Black, J.,
Hall-Spencer, J.M. & Rogers, A.D. Global habitat suitability of cold-water octocorals.
*Journal of Biogeography* **39**, 1278-1292 (2012).
63.  Quattrini, A.M., Baums, I.B., Shank, T.M., Morrison, C.L. & Cordes, E.E. Testing the
depth-differentiation hypothesis in a deepwater octocoral. *Proceedings of the Royal*
*Society B: Biological Sciences* **282**, 20150008 (2015).
64.  Bracco, A., Liu, G., Galaska, M.P., Quattrini, A.M. & Herrera, S. Integrating physical
circulation models and genetic approaches to investigate population connectivity in deep-
sea corals. *Journal of Marine Systems* **198**, 103189 (2019).
65.  Galaska, M.P., Liu, G., West, D., Erickson, K., Quattrini, A.M., Bracco, A. & Herrera, S.
Seascape genomics reveals metapopulation connectivity network of Paramuricea biscaya
in the northern Gulf of Mexico. *Frontiers in Marine Science*, 1790 (2021).
66.  Rex, M.A. & Etter, R.J. *Deep-sea biodiversity: pattern and scale*. (Harvard University
Press, 2010).
67.  Etnoyer, P.J., Wickes, L.N., Silva, M., Dubick, J., Balthis, L., Salgado, E. & MacDonald,
I.R. Decline in condition of gorgonian octocorals on mesophotic reefs in the northern
Gulf of Mexico: before and after the Deepwater Horizon oil spill. *Coral Reefs* **35**, 77-90
(2016).
68.  Fisher, C.R., Hsing, P.-Y., Kaiser, C.L., Yoerger, D.R., Roberts, H.H., Shedd, W.W.,
Cordes, E.E., Shank, T.M., Berlet, S.P., Saunders, M.G., Larcom, E.A. & Brooks, J.M.
Footprint of *Deepwater Horizon* blowout impact to deep-water coral communities.
*Proceedings of the National Academy of Sciences* **111**, 11744-11749 (2014).
69.  Naqib, A., Poggi, S., Wang, W., Hyde, M., Kunstman, K. & Green, S.J. Making and
sequencing heavily multiplexed, high-throughput 16S ribosomal RNA gene amplicon
libraries using a flexible, two-stage PCR protocol, in *Gene expression analysis* 149-169
(Springer, 2018).
70.  Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C., Al-Ghalith, G.A.,
Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K.,
Brejnrod, A., Brislawn, C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodríguez, A.M.,
Chase, J., Cope, E., Da Silva, R., Dorrestein, P.C., Douglas, G.M., Durall, D.M.,
Duvallet, C., Edwardson, C.F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J.M.,
Gibson, D.L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H.,
Huttenhower, C., Huttley, G., Janssen, S., Jarmusch, A.K., Jiang, L., Kaehler, B., Kang,

K.B., Keefe, C.R., Keim, P., Kelley, S.T., Knights, D., Koester, I., Kosciolek, T., Kreps, J., Langille, M.G.I., Lee, J., Ley, R., Liu, Y.-X., Loftfield, E., Lozupone, C., Maher, M., Marotz, C., Martin, B.D., McDonald, D., McIver, L.J., Melnik, A.V., Metcalf, J.L., Morgan, S.C., Morton, J., Naimey, A.T., Navas-Molina, J.A., Nothias, L.F., Orchanian, S.B., Pearson, T., Peoples, S.L., Petras, D., Preuss, M.L., Pruesse, E., Rasmussen, L.B., Rivers, A., Robeson, I.I.M.S., Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A., Sinha, R., Song, S.J., Spear, J.R., Swafford, A.D., Thompson, L.R., Torres, P.J., Trinh, P., Tripathi, A., Turnbaugh, P.J., Ul-Hasan, S., van der Hooft, J.J.J., Vargas, F., Vázquez-Baeza, Y., Vogtmann, E., von Hippel, M., Walters, W., Wan, Y., Wang, M., Warren, J., Weber, K.C., Williamson, C.H.D., Willis, A.D., Xu, Z.Z., Zaneveld, J.R., Zhang, Y., Zhu, Q., Knight, R. & Caporaso, J.G. QIIME 2: Reproducible, interactive, scalable, and extensible microbiome data science. *PeerJ Preprints* **6**, e27295v27292 (2018).

71.  Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. & Glöckner, F.O. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic acids research* **41**, D590-D596 (2012).

72.  Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A. & Cresko, W.A. Stacks: an analysis tool set for population genomics. *Molecular ecology* **22**, 3124-3140 (2013).

73.  Parra-Salazar, A., Gomez, J., Lozano-Arce, D., Reyes-Herrera, P.H. & Duitama, J. Robust and efficient software for reference-free genomic diversity analysis of GBS data on diploid and polyploid species. *bioRxiv*, 2020.2011. 2028.402131 (2020).

74.  Kamvar, Z.N., Tabima, J.F. & Grünwald, N.J. Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**, e281 (2014).

75.  Dray, S. & Dufour, A.-B. The ade4 package: implementing the duality diagram for ecologists. *Journal of statistical software* **22**, 1-20 (2007).

76.  McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and projection for dimension reduction. *arXiv preprint arXiv:1802.03426* (2018).

77.  Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R., Simpson, G.L., Solymos, P., Stevens, M.H.H. & Wagner, H. Package 'vegan'. *Community ecology package, version* **2**, 1-295 (2013).

78.  Hurlbert, S.H. The Nonconcept of Species Diversity: A Critique and Alternative Parameters. *Ecology* **52**, 577-586 (1971).

79.  Aho, K.A. *Foundational and applied statistics for biologists using R*. (CRC Press, 2013).

80.  Legendre, P. & Anderson, M.J. Distance□based redundancy analysis: testing multispecies responses in multifactorial ecological experiments. *Ecological monographs* **69**, 1-24 (1999).

81.  Lin, H. & Peddada, S.D. Analysis of compositions of microbiomes with bias correction. *Nature Communications* **11**, 3514 (2020).

82.  Haydon, T.D., Suggett, D.J., Siboni, N., Kahlke, T., Camp, E.F. & Seymour, J.R. Temporal variation in the microbiome of tropical and temperate octocorals. *Microbial Ecology* **83**, 1073-1087 (2022).

83.  Ziegler, M., Grupstra, C.G.B., Barreto, M.M., Eaton, M., BaOmar, J., Zubier, K., Al-Sofyani, A., Turki, A.J., Ormond, R. & Voolstra, C.R. Coral bacterial community structure responds to environmental change in a host-specific manner. *Nature Communications* **10**, 3092 (2019).

84.   Voolstra, C.R. & Ziegler, M. Adapting with Microbial Help: Microbiome Flexibility
Facilitates Rapid Responses to Environmental Change. *BioEssays* **42**, 2000004 (2020).
85.   Lima, L.F., Weissman, M., Reed, M., Papudeshi, B., Alker, A.T., Morris, M.M.,
Edwards, R.A., de Putron, S.J., Vaidya, N.K. & Dinsdale, E.A. Modeling of the coral
microbiome: the influence of temperature and microbial network. *MBio* **11**, e02691-
02619 (2020).
86.   Lee, S.T.M., Davy, S.K., Tang, S.-L., Fan, T.-Y. & Kench, P.S. Successive shifts in the
microbial community of the surface mucus layer and tissues of the coral Acropora
muricata under thermal stress. *FEMS Microbiology Ecology* **91** (2015).
87.   Ziegler, M., Seneca, F.O., Yum, L.K., Palumbi, S.R. & Voolstra, C.R. Bacterial
community dynamics are linked to patterns of coral heat tolerance. *Nature
Communications* **8**, 14213 (2017).
88.   Pootakham, W., Mhuantong, W., Yoocha, T., Putchim, L., Jomchai, N., Sonthirod, C.,
Naktang, C., Kongkachana, W. & Tangphatsornruang, S. Heat☐induced shift in coral
microbiome reveals several members of the Rhodobacteraceae family as indicator species
for thermal stress in Porites lutea. *MicrobiologyOpen* **8**, e935 (2019).
89.   Bourne, D., Iida, Y., Uthicke, S. & Smith-Keune, C. Changes in coral-associated
microbial communities during a bleaching event. *The ISME journal* **2**, 350-363 (2008).
90.   Grottoli, A.G., Dalcin Martins, P., Wilkins, M.J., Johnston, M.D., Warner, M.E., Cai, W.-
J., Melman, T.F., Hoadley, K.D., Pettay, D.T. & Levas, S. Coral physiology and
microbiome dynamics under combined warming and ocean acidification. *PloS one* **13**,
e0191156 (2018).
91.   Assié, A., Leisch, N., Meier, D.V., Gruber-Vodicka, H., Tegetmeyer, H.E., Meyerdierks,
A., Kleiner, M., Hinzke, T., Joye, S., Saxton, M., Dubilier, N. & Petersen, J.M.
Horizontal acquisition of a patchwork Calvin cycle by symbiotic and free-living
Campylobacterota (formerly Epsilonproteobacteria). *The ISME Journal* **14**, 104-122
(2020).
92.   Vohsen, S.A., Gruber-Vodicka, H.R., Osman, E.O., Saxton, M.A., Joye, S.B., Dubilier,
N., Fisher, C.R. & Baums, I.B. Deep-sea corals near cold seeps associate with
chemoautotrophic bacteria that are related to the symbionts of cold seep and
hydrothermal vent mussels. *bioRxiv*, 2020.2002.2027.968453 (2020).
93.   Klaus, J.S., Janse, I., Heikoop, J.M., Sanford, R.A. & Fouke, B.W. Coral microbial
communities, zooxanthellae and mucus along gradients of seawater depth and coastal
pollution. *Environmental Microbiology* **9**, 1291-1305 (2007).
94.   Sikkel, P.C., Cook, C.A., Renoux, L.P., Bennett, C.L., Tuttle, L.J. & Smit, N.J. The
distribution and host-association of a haemoparasite of damselfishes (Pomacentridae)
from the eastern Caribbean based on a combination of morphology and 18S rDNA
sequences. *International Journal for Parasitology: Parasites and Wildlife* **7**, 213-220
(2018).
95.   Janouškovec, J., Tikhonenkov, D.V., Burki, F., Howe, A.T., Kolisko, M., Mylnikov, A.P.
& Keeling, P.J. Factors mediating plastid dependency and the origins of parasitism in
apicomplexans and their close relatives. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 10200-10207
(2015).
96.   Janouškovec, J., Horák, A., Barott, K.L., Rohwer, F.L. & Keeling, P.J. Global analysis of
plastid diversity reveals apicomplexan-related lineages in coral reefs. *Current Biology* **22**,
R518-R519 (2012).

97.    Glasl, B., Bourne, D.G., Frade, P.R., Thomas, T., Schaffelke, B. & Webster, N.S.
834            Microbial indicators of environmental perturbations in coral reef ecosystems.
835            *Microbiome* **7**, 1-13 (2019).
836    98.    Feng, D., Cordes, E.E., Roberts, H.H. & Fisher, C.R. A comparative study of authigenic
837            carbonates from mussel and tubeworm environments: Implications for discriminating the
838            effects of tubeworms. *Deep Sea Research Part I: Oceanographic Research Papers* **75**,
839            110-118 (2013).
840    99.    Girard, F. & Fisher, C.R. Long-term impact of the Deepwater Horizon oil spill on deep-
841            sea corals detected after seven years of monitoring. *Biological Conservation* **225**, 117-
842            127 (2018).