Defense-related callose synthase *PMR4* promotes root hair callose deposition and adaptation to phosphate deficiency in *Arabidopsis thaliana*

Kentaro Okada[1,2], Koei Yachi[1], Tan Anh Nhi Nguyen[1], Satomi Kanno[3], Shigetaka

Yasuda[1], Haruna Tadai[1], Chika Tateda[1], Tae-Hong Lee[1], Uyen Nguyen[4], Kanako Inoue[1],

Shunsuke Miyashima[1], Kei Hiruma[1], Kyoko Miwa[5], Takaki Maekawa[4, 6], Michitaka

Notaguchi[2], Yusuke Saijo[1]

[1]Graduate School of Science and Technology, Nara Institute of Science and Technology,

Takayama-cho, Ikoma 630-0192, Japan

[2]Bioscience and Biotechnology Center, Nagoya University, Furo-cho, Chikusa-ku,

Nagoya 464-8601, Japan.

[3]Institute for Advanced Research, Nagoya University, Furo-cho, Chikusa, Nagoya

464-8601, Japan

[4]Institute for Plant Sciences, University of Cologne, Cologne D-50674, Germany

[5]Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810,

Japan

[6]CEPLAS Cluster of Excellence on Plant Sciences at the University of Cologne,

Cologne D-50674, Germany

Author for correspondence: Yusuke Saijo

ORCID ID: 0000-0001-8985-9695

Tel: +81-743-72-5690

Email: saijo@bs.naist.jp

Summary

- Plants acquire phosphorus (P) primarily as inorganic phosphate (Pi) from the soil. Under Pi deficiency, plants induce an array of physiological and morphological responses, termed phosphate starvation response (PSR), thereby increasing Pi acquisition and use efficiency. However, the mechanisms underlying PSR remain poorly understood.

- Here, we report that deposition of a β-1,3-glucan polymer called callose is induced in *Arabidopsis thaliana* root hairs under Pi deficiency, in a manner independent of PSR-regulating *PHR1/PHL1* transcription factors and *LPR1/LPR2* ferroxidases.

- Genetic studies revealed *PMR4* (*GSL5*) callose synthase being required for the callose deposition in Pi-depleted root hairs. Loss of *PMR4* also affects Pi mobilization, plant growth and anthocyanin accumulation under low Pi conditions, indicating the significance of PMR4 in both PHR1/PHL1- and LPR1/LPR2-mediated responses to Pi deficiency. The defects are not affected by simultaneous disruption of *SID2*, mediating defense-associated salicylic acid (SA) biosynthesis, unlike SA-dependent powdery mildew resistance previously described in *pmr4*.

- Grafting experiments and characterization of plants expressing *PMR4* specifically in root hair cells suggest that PMR4 in the cell type contributes to shoot growth under Pi deficiency. Our findings thus illuminate an important role for *PMR4* in plant adaptation to Pi deficiency.

Introduction

As an essential macronutrient, plants acquire and distribute P from the rhizosphere to the whole tissues, primarily in the form of inorganic phosphate (Pi). To maintain Pi homeostasis under Pi deficiency, plants induce an array of adaptive responses, *e.g.* increasing Pi uptake, remobilization and utilization, termed Pi starvation response (PSR), and involve beneficial microbes in mutualistic associations (Péret *et al*., 2011; Chien *et al*., 2018; Isidra-Arellano *et al*., 2021). PSR is accompanied by extensive transcriptional reprogramming, mainly through the transcription factor *PHR1* and related *PHL1*, which leads to increases in Pi transporter expression, root hair formation and anthocyanin accumulation (Rubio *et al*., 2001, Bustos *et al*., 2010). *PHR1/PHL1* mediate systemic coordination of these responses according to the internal Pi availability. In addition, Pi depletion leads to remodeling root system architecture, a decrease in primary root growth while increasing lateral root formation and growth, through the ferroxidases *LPR1* and *LPR2* (Svistoonoff *et al*., 2007; Péret *et al*., 2014; Balzergue *et al*., 2017). *LPR1/LPR2* are required for arresting primary root growth when the external Pi levels are reduced surrounding the root tips (Svistoonoff *et al*., 2007). Recent studies illuminated an intimate relationship between PSR and biotic interactions, but the molecular basis for their coordination remains poorly understood (Chan *et al*., 2021).

Callose deposition on the plasma membrane through callose synthases *Glucan Synthase Likes* (*GSLs*) reinforces barrier functions of the cell wall and restricts plasmodesmatal permeability, in plant development and biotic/abiotic stress responses (Jacobs *et al*., 2003; Nishimura *et al*., 2003; Dong *et al*., 2005; Enns *et al*., 2005; Nishikawa *et al*., 2005; Chowdhury *et al*., 2014; Ellinger & Voigt, 2014; Sager & Lee,

2014; Wu *et al*., 2018). In *Arabidopsis thaliana* (hereafter Arabidopsis)*,* callose

deposition is induced in the root phloem in response to excess iron, thereby reducing

plasmodesmatal permeability (O'Lexy *et al*., 2018). Of 12 Arabidopsis *GSLs*, *GSL2*

(*CALS5*) and *GSL5* (*PMR4, CALS12*) respectively contribute to basal accumulation and

iron-induced deposition of callose in the root phloem (O'Lexy *et al*., 2018). Callose also

accumulates in the stem cell niche (SCN) surrounding the quiescent center cells, thereby

inhibiting cell-to-cell movement of the transcription factor SHORT ROOT (SHR)

during PSR. This serves to suppress cell division and elongation in the root apical

meristem and elongation zone, respectively, through the *LPR1/LPR2* pathway (Müller

et al., 2015; Mora-Macías *et al*., 2017). *GSL5* (*PMR4*) also mediates callose deposition

in response to pathogen challenge and microbe/damage-associated molecular patterns

(MAMPs/DAMPs), and promotes fungal penetration resistance (Wang *et al*., 2021).

However, the loss of *GSL5* function results in strong resistance to powdery mildew

fungi through tightened SA-based defenses, demonstrated by *gsl5* mutants revealed as

powdery mildew resistance4 (*pmr4*) (Vogel & Somerville, 2000; Nishimura *et al*.,

2003). This resistance is thought to represent a backup defense activated when *GSL5*

(*PMR4*)-mediated callose deposition, a key output of MAMP/DAMP-triggered defense

signaling, is disrupted. Despite increasing knowledge, our understanding is still limited

for the possible involvement and significance of callose deposition in biotic and abiotic

stress responses.

Here, we report that callose deposition is induced in root hairs under Pi deficiency, in

a manner independent of *PHR1/PHL1* and *LPR1/LPR2*. Our genetic studies revealed

that the Pi depletion-induced callose deposition is impaired in *pmr4* (*gsl5*) root hairs.

The loss of *PMR4*-mediated callose deposition is associated with reduced plant growth

as well as exaggerated stress symptoms under Pi deficiency, in addition with enhanced

powdery mildew resistance. The results indicate *PMR4*-mediated callose deposition in

root hairs as an important step in plant adaptation to Pi-limiting conditions.

Materials and Methods

Plant Materials and growth conditions

The *Arabidopsis thaliana* accessions used were Col-0 and Ws-2, including the mutants

examined (Table S1) and the wile type used was Col-0 unless otherwise stated. We

generated ethyl methanesulfonate (EMS)-mutagenesis (M2) seeds for screening in both

Col-0 and Ler backgrounds. All seeds were sterilized using 6% sodium hypochlorite

and 0.2% Trion X-100 for 15 min, and then kept at 4°C for 2 d before sowing a

half-strength Murashige and Skoog (MS) medium with 25 mM sucrose and 0.8% Agar

(WAKO, Japan). If not stated otherwise, the normal Pi media contained 625 µM

KH_2PO_4, whereas the low Pi media contained 0 µM, 10 µM, 50 µM or 150 µM KH_2PO_4

as indicated. The low N and low Fe media used contained 0 µM N and 0 µM Fe,

respectively. Plants were grown in a growth chamber under 12 h: 12 h, light: dark

cycles at 22°C. Six-day (d)-old seedlings were transferred to the indicated Pi liquid

media or agar media with 1% Difco Agar, Bacteriological (BD, Franklin Lakes, NJ,

USA).

Construction and plant transformation

The genomic DNA sequences of the *POWDERY MILDEW RESISTANCE 4* (*PMR4*,

AT4G03550) loci, together with a regulatory DNA sequence for root hair specific

expression (below), were cloned into the Gateway binary vector pGWB504 (Nakagawa

et al., 2007; primer information Table S2) to express a C-terminally GFP-fused PMR4

under its native regulatory DNA sequence. The root hair-specific promoter sequence

used was derived from *LRR-EXTENSIN 1* (*LRX1*, AT1G12040) (Baumberger *et al.*,

2003). Transgenic lines were generated by Agrobacterium-mediated transformation of

pmr4-1 with the DNA construct and subsequently selected on hygromycin resistance

and DNA genotyping.

Aniline blue and PI staining

Callose staining with aniline blue was conducted as described in Schenk & Schikora

(2015). Plants were examined at 3 d after the transfer of 5-d-old seedlings to the

indicated Pi, N or Fe liquid media (Figs. 1a, c). Plant tissues were fixed in a solution

containing acetic acid and ethanol (1:3 v/v) overnight. The fixation solution was

exchanged to a 150 mM K_2HPO_4 solution before callose staining with 0.1% aniline blue

in the K_2HPO_4 solution for 2 h. The data presented are representative of three

independent experiments (Figs 1a, c, 2a, 3a), two independent experiments (Figs 1b, 6d,

S1, S2) and one experiment (Figs 7a, S3). To localize the root meristematic regions, the

primary roots were stained with 20 µg/ml propidium iodide (PI). Callose or PI signals

were examined under microscopy using a fluorescence microscope DMI 6000B (Leica

Microsystems, Wetzlar, Germany) and a confocal microscope SP8 (Leica Microsystems,

Wetzlar, Germany).

Identification of mutants altered in the root hair callose deposition

We examined callose deposition in *gsl* mutants (Table S1) after aniline blue staining of

the roots transferred to the low Pi liquid media (50 µM) for 3 d. In the EMS mutagenic

screening, we subjected a region ~5 cm from the root tip to aniline blue staining, after

cutting out of the M2 individuals transferred to the Pi -depleted liquid media (0 μM) for

3 d. This enabled to keep the plant alive for subsequent characterization and seed

propagation. The *PMR4* genomic region was sequenced in the selected mutant

candidates.

Inoculation analyses with powdery mildew fungi

Inoculation of *Golovinomyces orontii* and determination of fungal versus plant DNA

ratio were conducted essentially by following Weßling & Panstruga (2012).

Eighteen-d-old plants grown in 10 h: 14 h, light: dark cycles at 22°C were inoculated

with *G. orontii* spores at approximately 750 spores cm^{-2}. The inoculated leaves were

harvested and frozen in liquid nitrogen before determining fungal biomass at 10 d after

inoculation. Genomic DNA was extracted essentially by following Brouwer *et al*.

(2001). Quantitative real-time PCR was performed with primers (Table S2) and

Brilliant Sybr Green QPCR Reagent Kit (Stratagene, Waldbronn, Germany). The ratio

of *G. orontii* to Arabidopsis genomic DNA was calculated using the ΔΔCt method

(Pfaffl, 2001). Data are representative of two independent experiments (Figs. 2c, d).

Determination of root hair density, primary root length and root area

Six-day-old seedlings were transferred and further grown in the low (50 μM) or normal

Pi media (625 μM) for 6 d, and then root hairs, primary root and root system area were

examined under stereomicroscope M80 (Leica Microsystems, Wetzlar, Germany). The

number of root hairs was determined in a region 2.0-3.0 mm from the root tip using

Image J software (National Institutes of Health, Bethesda, MD, USA). Data are

161 representative of similar results from three independent experiments (Figs. 3c, d) and

162 two independent experiments (Fig. S4). The primary root length and convex hull area

163 were determined using ImageJ software and Root nav software (Pound *et al.*, 2013).

164 Data are representative of three independent experiments (Figs. 3e-h). For growth under

165 different B conditions, plants were grown in MGRL solid media containing 1% sucrose

166 and 1% gellan gum (WAKO, Japan). B concentrations were adjusted with boric acid.

167 Plants were grown under 16 h: 8 h, light: dark cycles at 22°C. The data presented are

168 from one independent experiment (Fig. S7).

170 **Determination of shoot fresh weight, anthocyanin and Pi contents**

171 In plant growth assays on media, six-day-old seedlings were transferred to the indicated

172 media (10 μM, 50 μM or 625 μM Pi) and then further grown for the indicated days.

173 Shoot fresh weight (SFW) was determined 16 d after transfer to the indicated media (50

174 μM or 625 μM Pi; Fig. 3i, j). In plant growth assays with *pLRX1::PMR4* lines, SFW

175 was determined 10 d after the transfer to the low Pi media (10 μM：Fig. 6a, c). In plant

176 growth assay on soil, organic black soil (Maruishi Engei, Osaka, Japan) and Ezo sand

177 (Kobayashi Sangyo, Kita-Nagoya, Japan) mixed at 1:1 (v/v) were used as low Pi soil,

178 and Sakata-Super-Mix-A (Sakata Seed, Kyoto, Japan) and Vermiculite G20 (Maruishi

179 Engei, Osaka, Japan) mixed at 1:1 (v/v) were used as normal Pi soil (Fig. S5). SFW of

180 seedlings grown on soil was determined 14 d after the transfer of 10-d-old seedlings to

181 the low and normal Pi soils. Data are representative of two independent experiments

182 (Figs 3i, j, S5) and one independent experiment (Fig. 7b, c).

183 In the determination of anthocyanin content, six-day-old seedlings were transferred to

184 and further grown on low Pi media (10 μM) for 10 d (Fig. 3k, l). The shoots were

subject to the procedure described in Mita (1997). In brief, a HCl: methanol mixture at

1:99 (v/v) was added to the ground tissues with 1 ml per sample, and then the resultant

homogenates were incubated at 4°C overnight. A_{530} and A_{657} of the supernatants were

determined with a spectrometer UV-1800 (Shimadzu, Kyoto, Japan). Anthocyanin

content was calculated following the formula: $(A_{530} - 0.25 \times A_{657}) \times$ fresh weight

$(FW)^{-1}$. Data are representative of two independent experiments with the same

conclusions (Fig. 3k, l).

In the determination of Pi content, six-d-old seedlings were transferred to and further

grown on the low Pi (150 μM) or normal Pi (625 μM) media for 9 d (Fig. 4a-d). Pi

content was determined by following Ames (1966). In brief, the extraction buffer (10

mM Tris, 1 mM EDTA, 100 mM NaCl, 1 mM phenylmethylsulfonyl fluoride pH8.0)

was added with 10 μl per sample mg FW. The homogenate 100 μl was mixed with 1%

glacial acetic acid 900 μl, and then incubated at 42°C for 30 min. The supernatant was

recovered to be incubated in a solution (0.35% NH_4MoO_4, 0.86 N H_2SO_4, and 1.4%

ascorbic acid) at 42°C for 30 min. A_{820} of the aliquot was measured under Eppendorf

BioSpectrometer® basic (Eppendorf, Hamburg, Germany). Then, Pi content was

calculated by a P calibration curve prepared with a P standard solution. Data are

representative of three independent experiments (Fig. 4a, b) and two independent

experiments (Fig. 4c, d) with the same conclusions.

Scanning electron microscope

The maturation zone of primary roots 10 d after transfer of 5-d-old seedlings to low Pi

(50 μM) liquid media was fixed in half-Karnovsky's fixative overnight at 4°C, and then

postfixed in 1% osmium tetroxide for 1 h at room temperatures (Karnovsky, 1965). The

specimens were dehydrated in an ethanol series (30%, 50%, 70%, 80%, and 90%, and

100% at three times) for 10 min each at RT, and then the specimens were dipped in

t-butyl alcohol and dried with a vacuum evaporator. They were coated with vapor of 1%

osmium tetroxide and then observed with a scanning electron microscope (JIB-4601F,

JEOL, Tokyo, Japan).

Inductively coupled plasma-mass spectrometer (ICP-MS) analyses

Five-day-old seedlings were transferred to and further grown on the growth media (10

μM or 625 μM KH_2PO_4) for 4 d (Figs S6, S7). Shoots and roots were harvested and

dried at 60°C. After determining the dry weight, the dried tissues were digested with 2

ml nitric acid and 0.5 ml hydrogen peroxide (Wako Pure Chemicals) at 110°C. Digested

samples were then dissociated with 5 ml of 2% (w/v) nitric acid and subjected to

inductively coupled plasma mass spectrometry (NexION 2000, Perkin-Elmer). The

contents of P, K, Mg, Ca, Mn, Fe, Zn, Mo and B were determined. The data were

obtained from one experiment (Figs 5, S6).

Determination of ^{32}P uptake rate and translocation ratio

Seven-day-old seedlings were transferred to and further grown on the growth media (50

or 625 μM KH_2PO_4) for 10 d. The seedlings were exposed to the liquid media

containing ^{32}P-labeled Pi ($^{32}PO_4^{3-}$ 20 kBq / ml) for 30 min. The plant tissues were fixed

to an Imaging Plate (GE Healthcare co., Chicago, USA) for 3 h and then scanned by

Typhoon FLA 9000 (GE Healthcare co., Chicago, USA). ^{32}P uptake rate was calculated

based on a calibration curve for ^{32}P. For determining the Pi translocation rate into shoots,

the seedlings were exposed to the Pi-containing liquid media containing ^{32}P-labeled Pi

for 10 min, and then transferred to non-labelled agar media for 120 min. The

translocation rate of Pi was determined by calculating ^{32}P ratio of the shoot to the whole

seedlings. The data presented are representative of two independent experiments with

the same conclusions (Fig. 4f, g).

Grafting experiments

Grafting of WT and *pmr4* was performed with a supportive micro-device as described

in Tsutsui *et al*. (2020). Four-day-old seedlings grown on a 1/2 MS medium containing

0.5% sucrose and 1% agar under constant light at 22°C were subject to grafting. Both

hypocotyls of stock and scion were cut horizontally with a knife and assembled on the

device using forceps. The grafted plants were grown in a 1/2 MS medium containing

0.5% sucrose and 2% agar at 27°C for 6 d. In callose staining, the grafted plants

transferred and grown on low Pi media (50 µM) under constant light at 22°C for 9 d

were examined. In plant growth assays, the grafted plants transferred and grown on low

Pi media (150 µM) under constant light at 22°C for 14 d were examined. The data

presented are representative of two independent experiments with the same conclusions

(Fig. 6).

Results

Callose deposition occurs in root hairs under phosphate deficiency, independently

of *PHR1/PHL1* and *LPR1/LPR2*

Under low Pi conditions (Pi at 0 or 50 µM), we noticed callose deposition in the

epidermal cells, particularly in root hairs of the differentiation zone (Fig. 1a, b; O'Lexy

et al., 2018). However, callose deposition was not detected in root hairs under the tested

N, Fe or Fe/Pi deficient conditions (Fig. 1a). The results indicate that the callose

deposition is specific to Pi deficiency and is dependent on Fe. The absence of callose

staining in the epidermis of root hair-less *rhd6* mutants (Fig. 1c) validated that callose

deposition occurs in root hairs under low Pi. In contrast to previous study (Müller *et al*.,

2015), we did not detect callose deposition in the root SCN and cortex cells of the

elongation zone under our experimental conditions (Figs. 1a, c).

We next assessed whether *PHR1/PHL1* and *LPR1/LPR2* play a role in the root hair

callose deposition. Although *LPR1* and *LPR2* contribute to the callose deposition in

SCN (Müller *et al*., 2015), the root hair callose deposition was not affected in *lpr1 lpr2*

(Fig. 1c). Furthermore, wild type (WT)-like callose deposition was retained in root hairs

of *phr1 phl1 lpr1 lpr2* under low Pi conditions (Fig. 1c). The results indicate that callose

deposition is induced in root hairs under Pi deficiency, independently of these PSR

regulators and callose deposition in the root SCN.

***PMR4* callose synthase contributes to callose deposition under Pi deficiency**

To unravel genetic requirements for the callose deposition in root hairs under Pi

deficiency, we examined *glucan synthase like* (*gsl*) mutants for alternations in callose

deposition under phosphate deficiency. Of the tested *gsl* mutants, *gsl5* (hereafter, *pmr4*)

mutants specifically lacked callose deposition in root hairs (Fig. S1). We also screened a

total of 18,550 ethyl methane sulfonate (EMS)-mutagenized individuals in Col-0 and

Ler backgrounds. This revealed two mutants in the Col-0 background, designated

<u>*callose alteration under Pi starvation 1 and 2*</u> (*caps1 and caps2*, Fig. 2a). *caps1* and

caps2 mutants were severely defective in callose deposition of root hairs under low Pi

conditions (Fig. 2a). DNA Sequencing on the *PMR4* genomic region revealed a DNA

substitution within the coding region, resulting in a non-synonimus amino acid

substitution, in both *caps1* and *caps*2 alleles (Fig. 2b). The substitution in *caps*1

converts Tryptophan to Stop codon at the 260^{th} amino acid (PMR4^{W260*}), resulting in

PMR4 truncation, consistent with the severely impaired callose deposition in *caps1* as

well as *pmr4-1* (PMR4^{W687*}). In *caps2*, glutamic acid is substituted with lysine at the

1181^{st} amino acid (PMR4^{E1181K}), located within the cytosolic side of a predicted

glucan-synthase domain (residue 575 to 1558, orange-marked domain, Fig. 2b). The

results suggest that dysfunction of PMR4 fails to induce root hair callose deposition

under Pi deficiency in these *caps* alleles. We validated that leaf callose deposition in

response to the bacterial MAMP flg22, an output in pattern-triggered immunity (PTI)

dependent on *PMR4*, is abolished in these *caps* alleles (Fig. S2). Moreover, we showed

strong resistance to powdery mildew fungi in these *caps* plants (Fig. 2c, d), as

previously described in *pmr4* (Vogel & Somerville, 2000; Nishimura *et al*., 2003). The

results confirm that PMR4 functions are impaired in both *caps* alleles.

***PMR4* influences root development traits induced under Pi deficiency**

By using *pmr4-1* as a representative mutant allele, we further investigated the

functional significance of *PMR4* under Pi deficiency. The loss of *PMR4* leads to an

increase in salicylic acid (SA) biosynthesis and defense under biotic stress (Vogel &

Somerville, 2000; Nishimura *et al*., 2003), in a manner requiring *SID2 isochorismate*

synthase that mediates a critical step in SA biosynthesis during defense activation

(Wildermuth *et al*., 2001). Given the negative impact of SA defense activation on plant

growth (Mateo *et al*., 2006; Zhang *et al*., 2010), we assessed the possible SA

dependence of the observed *pmr4* phenotypes, by examining genetic interactions

between *pmr4* and *sid2*. Callose deposition was induced in *sid2* root hairs but not

induced in *pmr4 sid2* under low Pi conditions (Fig. 3a), demonstrating that

SID2-mediated SA biosynthesis is not required for *PMR4*-mediated callose deposition

in root hairs. The results also exclude over-activation of SA defense as the cause of the

callose defects in *pmr4* root hairs.

In Arabidopsis, root hair cells specifically occur at the junction of two underlying

cortical cells, whereas non-hair cells occur in the other epidermal cells (Dolan *et al.*,

1994; Schiefelbein *et al*., 2009). Root hair density was increased under low Pi

conditions compared to normal Pi conditions, and the increase was exaggerated in *pmr4*

as well as *pmr4 sid2*, without ectopic root hair formation (Figs 3b-d, S3a). Likewise,

root hair length was also increased in *pmr4* as well as *pmr4 sid2* under both Pi

conditions (Fig. 3c, d). The results suggest that *PMR4* is not required for the

differentiation or outgrowth of root hair cells, but that *PMR4* contributes to restricting

root hair formation and elongation, in the presence or absence of *SID2*. Conversely,

primary root length was reduced in *pmr4* compared with WT in both low and normal Pi

conditions (Fig. 3b, e). We then determined the ratio of the primary root length under

normal Pi versus low Pi as an indicator for the primary root growth inhibition under Pi

deficiency. The ratio (normal Pi/low Pi) was increased in *pmr4* compared with WT,

indicating that primary root growth inhibition, a Pi deficiency-characteristic response, is

exaggerated in *pmr4* (Fig. 3f). We also observed a decrease in the root meristem region

in *pmr4* compared with WT under Pi deficiency, suggesting that the meristem

exhaustion coincides with the growth inhibition in the primary root of *pmr4* (Fig. S3b).

The primary root growth inhibition under low Pi was alleviated in *sid2*, but it was again

exaggerated in *pmr4 sid2* compared to *sid2* (Fig. 3e, f), indicating that the loss of *PMR4*

enhances the Pi deficiency response, with or without *SID2*.

Since the primary root growth inhibition under Pi deficiency relies on *LPR1* and

LPR2 (Svistoonoff *et al*., 2007; Müller *et al*., 2015), we examined the possible

relationship between *LPR1/LPR2* and *PMR4* in the regulation of primary root growth.

The primary root length was reduced in *pmr4 lpr1 lpr2* compared with *lpr1 lpr2* under

both Pi conditions, whereas it was increased in *pmr4 lpr1 lpr2* compared with *pmr4*

specifically under low Pi (Figs. S4a, b). The results indicate that *PMR4* serves to sustain

the primary root growth irrespective of the Pi availability or *LPR1/LPR2* activity,

whereas the *LPR1/LPR2* inhibition is specific under low Pi with or without *PMR4*. The

ratio of the primary root length under normal Pi versus low Pi was lowered to a similar

extent in *lpr1 lpr2* and *pmr4 lpr1 lpr2* compared with WT (Fig. S4c), despite the

decrease in the primary root length by *pmr4* (Fig. S4b), indicating that *lpr1 lpr2* is

epistatic to *pmr4*.

We next determined the convex hull root area as a proxy of overall root system

growth, which was correlated with Pi availability (Fig. 3g). It was reduced in *pmr4* and

sid2 compared with WT under both normal and low Pi conditions, albeit to a lesser

extent in *sid2* than *pmr4* under low Pi (Fig. 3g), suggesting that *PMR4* and *SID2* both

contribute to expanding the root area. Notably, only in the presence of *SID2*, the root

area was reduced under both Pi conditions whereas its ratio under normal Pi versus low

Pi was increased, in *pmr4* relative to WT (Fig. 3g, h). The results suggest that *PMR4*

promotion of the root system expansion relies on *SID2*. Conversely, the root area was

greater in *pmr4 sid2* compared to *pmr4* under low Pi (Fig. 3g), suggesting that *SID2*

negatively impacts root expansion specifically in the absence of *PMR4* under Pi

deficiency. There results thus suggest that *PMR4* directs *SID2*-mediated SA

biosynthesis to root growth expansion in Pi-depleted plants, consistent with an

uncoupling of SA induction from defense activation under Pi deficiency (Gulabani *et al.*

2021).

***PMR4* serves to sustain shoot growth and restrict anthocyanin accumulation under**

Pi deficiency.

We next examined whether *PMR4* influences shoot growth, by determining shoot

fresh weight (FW) under Pi deficiency and sufficiency. Shoot FW was lowered in *pmr4*

compared with WT, and it was not even partially restored by the simultaneous loss of

SID2 in *pmr4 sid2* under low-Pi media or soil (Fig. 3i-j and S5). Together with the root

phenotypes described above, the results exclude *SID2* as the major cause of the growth

inhibition in *pmr4*. The results suggest that *PMR4* positively influences shoot growth in

the presence of *SID2*. Although high SA accumulation was previously described to

reduce plant growth (Miura *et al.*, 2010), we did not detect negative impacts of *SID2* on

shoot growth under low Pi conditions (Fig. 3j). *SID2* even positively influenced shoot

growth under the normal Pi conditions, indicated by the reduced shoot biomass in *sid2*

compared to WT (Fig. 3j).

Anthocyanin content was increased in *pmr4* compared with WT, in the presence or

absence of *SID2* (Fig. 3k, l). Although SA was described to enhance anthocyanin

accumulation under Pi deficiency (Morcillo *et al.*, 2019), *sid2* was largely

indistinguishable from WT in the anthocyanin contents under our low Pi conditions (Fig.

3k, l). The results imply that PMR4-mediated Pi acquisition and allocation serves to

alleviate Pi starvation causing anthocyanin accumulation under Pi deficiency in a

manner independent of *SID2*.

***PMR4* promotes the acquisition and root-to-shoot translocation of Pi**

The *PMR4* effects on shoot and root growth prompted us to examine its possible role

in Pi acquisition from the rhizosphere and Pi mobilization within the plant. We found

that shoot Pi concentrations were greatly reduced in *pmr4* compared with WT, without

alterations in root Pi concentrations, under Pi deficiency (Fig. 4a, b). This indicates that

PMR4 contributes to shoot Pi accumulation in Pi-depleted plants. It was described that

PHR1/PHL1 promotes shoot Pi accumulation (Wang *et al.*, 2018) and that *PHO1* Pi

transporter mediates Pi mobilization from roots to shoots through the root xylem vessels

(Stefanovic *et al.*, 2007). We validated that Pi contents were lowered in the shoots but

increased in the roots, in *phr1 phl1* and *pho1* compared with WT under low Pi (Fig. 4c).

A decrease in shoot Pi contents was also observed in *phr1 phl1* and *pho1* under normal

Pi (Fig. 4d). Since root hairs contribute to Pi accumulation under Pi deficiency (Tanaka

et al., 2014; Robertson-Albertyn *et al.*, 2017; Holz *et al.*, 2018), we next determined Pi

contents in a root hair less mutant, *rhd6* (Fig. 4c, d). As seen in *pmr4, phr1 phl1* and

pho1, Pi contents were lowered in the shoots but not in the roots of *rhd6* compared to

WT, specifically under Pi deficiency (Fig. 4c, d). In sum, all these mutants displayed a

decrease in shoot Pi contents without affecting or reducing root Pi contents under Pi

deficiency (Fig. 4e). We inferred from these results that *PMR4*-mediated callose

deposition in root hairs contributes to shoot Pi accumulation, possibly through

increasing the uptake or root-to-shoot translocation of Pi. To test the hypothesis, we

determined the rates of Pi uptake and Pi translocation by applying and tracing

radio-labelled P. The rate of ^{32}P uptake per time and the ratio of ^{32}P in the shoot to

whole seedling, standing for Pi translocation rate, were both lowered in *pmr4* and *pmr4*

sid2 compared to WT and *sid2,* respectively, under low Pi (Figs. 4f, g). In the presence

of *SID2*, *PMR4* contribution to the two proxies was significant only under low Pi (Figs.

4f, g). In the absence of *SID2*, however, *PMR4* contribution to P uptake was apparent

under normal Pi as well (Fig. 4f). Moreover, the ^{32}P uptake rate was reduced in *sid2*

compared to WT, whereas *pmr4 sid2* was not distinguishable from *pmr4* under low Pi

(Fig. 4f), suggesting that SID2 positively influences Pi uptake under Pi deficiency in the

presence of PMR4. These results suggest that PMR4 promotes Pi uptake and

root-to-shoot Pi translocation under Pi deficiency and that this is in part achieved

through diverting SID2 to promote these processes.

We further assessed the possible *PMR4* contributions to the acquisition and tissue

mobilization of Pi and other elements, by inductively coupled plasma mass

spectrometry (ICP-MS) analyses for P, K, Mg, Ca, Mn, Fe, Zn, Mo and B contents in

Pi-depleted and Pi-supplied plants. Under Pi sufficiency, although the contents of these

elements were not distinguishable between WT and *pmr4* roots, the contents of K and

the rest were increased and lowered, respectively, in *pmr4* compared to WT shoots (Fig.

S6). Under low Pi, Mn, Zn and Mo contents were greater in *pmr4* compared with WT

roots, whereas the contents of all the tested elements except K were lowered in *pmr4*

compared with WT shoots (Fig. 5). The results validated the role for *PMR4* in shoot P

accumulation without affecting root P contents under both Pi conditions (Figs. 4a, 4b, 5,

S6). Additionally, the results suggest that *PMR4* also positively influences shoot

accumulation of Mg, Ca, Mn, Fe, Zn, Mo and B (Figs. 5, S6). WT-like and increased K

accumulation in *pmr4* shoots compared to WT under Pi deficiency and sufficiency,

respectively, show that the root-to-shoot transport is not entirely reduced for all

elements in *pmr4*, pointing to a certain degree of element specificity in *PMR4*

dependence (Figs 5, S6).

The role of *PMR4* in the shoot accumulation of different elements prompted us to

examine whether *PMR4* influences plant responses to different nutrient deficiency. We

examined *pmr4* phenotypes under B scarcity, as B scarcity induces inhibition of the

primary root and transcriptional reprogramming related to ROS and defense in roots

(Miwa *et al*., 2013; Das & Purkait, 2020). Although the primary root length was

reduced in *pmr4* compared to WT over the tested range of B conditions, its ratio under

B sufficiency (100 µM) versus the examined low B conditions was not affected in *pmr4*

(Fig. S7). The results point to specific contribution of *PMR4* to plant growth under Pi

deficiency.

Root *PMR4* contributes to shoot growth under Pi deficiency

We next examined the possible significance of the root *PMR4* pool in promoting

shoot growth, through micrografting between WT and *pmr4* plants. We validated that

the micrografted plants of WT/WT (shoot/root genotypes) and *pmr4/pmr4* showed the

corresponding WT and *pmr4* phenotypes in root growth and SFW (Figs. 6a-c). The root

hair callose was detected only when *PMR4* was present in the roots, although the callose

signals were not as strong as in intact plants, possibly reflecting the stresses imposed by

the grafting procedure (Fig. 6d). Remarkably, SFW was greatly reduced in WT*/pmr4*

whereas it was less affected in *pmr4/*WT (Fig. 6b), pointing to a critical role for a root

PMR4 pool in shoot growth under Pi deficiency. However, the root growth was partially

restored in WT/*pmr4* (Fig. 6c), pointing to a role for a shoot *PMR4* pool in sustaining

the primary root growth in Pi-depleted plants.

To further examine the role of a PMR4 pool in root hairs, we generated transgenic

plants expressing *PMR4* under the control of a root hair-specific *LRX1* promoter

(*pLRX1*, Baumberger *et al.*, 2003) in the *pmr4-1* background (Figs. 7a, b).

pLRX1::PMR4 complemented *pmr4* in terms of the callose deposition in root hairs and

shoot growth under low Pi conditions (Figs. 7a-c). By contrast, the primary root growth

remained suppressed in the *pLRX1::PMR4* transgenic lines as well as in *pmr4* (Figs. 7b,

d). The results indicate that PMR4 in root hairs is sufficient to restore shoot growth,

possibly uncoupled from the primary root growth regulation, under Pi deficiency.

Discussion

Our studies on callose deposition in root hairs induced under Pi deficiency led to the

discovery of *PMR4* as a key regulator of plant adaptation to Pi deficiency (Fig. 8).

PMR4 mediates defense-induced callose deposition in pattern triggered immunity and

pre-invasion fungal resistance (Jacobs *et al.*, 2003; Chowdhury *et al.*, 2014), and that its

loss is linked to enhanced SA-based pathogen resistance (Nishimura *et al.*, 2003).

Therefore, *PMR4* seems to play a critical role in an array of biotic and abiotic stress

responses, possibly through callose deposition at different sites.

The present study suggests that the root hair callose contributes to root Pi uptake and

shoot Pi accumulation under Pi deficiency. Importantly, the loss of *PMR4* affects both

PHR1-dependent and *LPR1*-dependent processes (Figs 3, 4) (Bates & Lynch 1996;

Thibaud *et al.*, 2010), which have been genetically uncoupled in PSR (Wang *et al.*,

2010; Balzergue *et al.*, 2017). Pi accumulation and anthocyanin accumulation are

suppressed in *phr1 phl1* (Rubio *et al.*, 2001; Bustos *et al.*, 2010; Wang *et al.*, 2018). The

shoot Pi contents, and plant biomass are also reduced in *pmr4*, implying that *PMR4*

positively influences *PHR1/PHL1*-mediated Pi acquisition and plant growth under Pi

deficiency. However, over-accumulation of anthocyanin in *pmr4* (Fig. 3k, l) in contrast

to *phr1 phl1* (Rubio *et al.*, 2001; Bustos *et al.*, 2010; Wang *et al.*, 2018), suggests that

PMR4 contribution is specific to the former two outputs of *PHR1/PHL1* and that its loss

results in exaggeration of Pi starvation symptom. The loss of *PMR4* also strengthens

inhibition of the primary root growth under Pi deficiency, only in the presence of

LPR1/LPR2 (Fig. S4). The LPR1 pathway mediates callose deposition at the root SCN

to inhibit the cell-to-cell movement of the transcription factor SHR, thereby suppressing

the cell division and subsequent root elongation (Müller *et al.*, 2015; Mora-Macías *et al.*,

2017). *PMR4*-mediated primary root growth retention is prominent only in the presence

of *LPR1/LPR2*, whereas *LPR1/LPR2*-mediated growth inhibition is significant even in

the absence of *PMR4* under low Pi (Fig. S4). Our results suggest a role for PMR4 in

antagonizing LPR1/LPR2-dependent root growth inhibition.

The failure of root hair PMR4 to restore primary root growth (Figs. 7b, d) implies the

involvement of PMR4 from another cell type(s) than root hairs in this regulation. This is

consistent with the partial and full retention of shoot Pi accumulation in *rhd6* lacking

root hairs under low and normal Pi conditions, respectively (Fig. 4c-e). Although our

callose detection was limited to root hairs, it is conceivable that PMR4-mediated callose

deposition in different cell types serves to sustain the primary root growth against

LPR1/LPR2-mediated attenuation under Pi deficiency. The mechanisms underlying

antagonistic interactions between *PMR4* and *LPR1/LPR2* require further studies

including the identification of the SCN callose synthase.

Shoot Pi accumulation depends on root Pi uptake and subsequent Pi translocation.

PHT1;1 and *PHT1;4* provide two major Pi transporters in roots, contributing greatly to

Pi acquisition under Pi deficiency (Shin *et al*., 2004). *PHO1* mediates Pi translocation

from roots to shoots through Pi loading into the root xylem (Poirier *et al.,* 1991;

Hamburger *et al*., 2002; Liu *et al*., 2012). Notably, the loss of *PMR4* results in a

decrease in the shoot Pi contents without affecting the root Pi contents (Figs 4a, 4b, 5).

Likewise, the loss of *PHR1/PHL1* and *PHO1* also lowers shoot Pi contents without

reducing root Pi contents (Fig. 4c, d; Liu *et al*., 2012; Wang *et al*., 2018). In *pht1;1*

pht1;4, a decrease of Pi uptake in roots is associated with that of Pi translocation to the

shoots and shoot Pi accumulation (Shin *et al*., 2004). An intimate linkage between root

uptake and root-to-shoot translocation is also seen for K^+ (Nieves-Cordones *et al*., 2019).

It is thus conceivable that the lowered Pi accumulation in *pmr4* shoots is caused by a

decrease in the rates of the root Pi uptake and the root-to-shoot Pi translocation,

possibly through the compromised root hair function in the absence of *PMR4*. Indeed,

^{32}P tracer analyses indicate that *PMR4* promotes both root Pi uptake and root-to-shoot

Pi translocation, without affecting the root Pi contents (Figs. 4a, b, f, g). This is

consistent with the view that the root-to-shoot Pi translocation is driven by the Pi uptake

in the roots (Shin *et al*., 2004). Leaf transpiration affects root-to-shoot solute

translocation (Nieves-Cordones *et al*., 2019), but it might be ignorable under high

humidity in the closed plate assays. Notably, *PMR4* also contributes to the accumulation

of different elements except K in shoots under Pi deficiency (Figs. 5, S6). This might

largely attribute to secondary effects of *PMR4*-dependent root growth and expansion

(Figs 3e, g), but which do not explain the separate regulation of K (Figs. 5, S6). In this

respect, the disruption of multiple *PHT1* members (*PHT1;1/1;2/1;3* silencing in the

phf1 pht1;4 mutant background, designated mut5) results in a decrease in Na as well as

P contents compared to WT (Ayadi *et al.*, 2015). Cross-regulation among Fe, Zn and Pi

homeostasis relies on *PHR1* (Briat *et al.*, 2015). Consistently, our data highlight further

coordination among a wide spectrum of elements in their acquisition and shoot

accumulation through *PMR4* (Figs 5, S6).

Root hairs increase the root surface area to promote root anchorage, nutrient

acquisition, and metabolite exudation, thereby facilitating the interactions with the soil

environments (Datta *et al.*, 2011; Poole *et al.*, 2018; Robe *et al.*, 2021; Vissenberg *et al.*,

2020; Han *et al.,* 2023). The elongation and increased densities of root hairs in the

absence of PMR4-mediated callose deposition therein (Fig. 3c, d) can be inferred that

the root hairs formed in *pmr4* are less functional in Pi acquisition or Pi allocation to the

other tissues, thereby leading to further formation of root hairs. Our findings point to a

positive role for *PMR4* in the root hair functions under low Pi.

Additionally, root hairs also provide a key interface to soil-borne microbes and are

intimately associated with root defense regulation. Root hair formation and elongation

are facilitated by jasmonate (JA) (Zhu *et al.*, 2006), which mediates the expression of

key transcription factors for root hair differentiation and the de-repression of RHD6

from the interference by JAZ proteins (Han *et al*, 2020). In beneficial interactions with

the epiphytic bacterium *Pseudomonas simiae* WCS417, activation of defense-related

genes is increased in hair cell lineages compared to non-hair cell lineages (Verbon *et al.*,

2023). The endogenous elicitor-active Pep peptides induce root hair formation, and Pep

receptor signaling from the root hairs can activate defense responses without root

growth inhibition (Okada *et al.*, 2021). These findings suggest a specific capacity of

root hair cells in root immunity. How these properties of root hairs are influenced by

PMR4-mediated callose deposition under Pi deficiency also represents an interesting

topic for future studies.

Acknowledgement

We thank the Arabidopsis Biological Research Centre for published plant materials,

Yaichi Kawakatsu for advice on micrografting, and Michiko Tsukamoto and Mie

Matsubara for technical assistance. This work was supported in part by the grants from

Mitsubishi Foundation (YS), the Japan Science and Technology Agency (JST,

JPMJPR13B6 and JPMJSC1702 (YS) and JPMJPR16Q7 (KH)), and MEXT of Japan

18H02467 and 21H02507 (YS).

Author contribution

YS conceived the study. KO and YS designed the experiments. KO, KY, TANN, SK, SY,

HT, CT, THL, UN, KI, KM, TM and MN developed the methods, performed the

experiments, and/or analyzed the data. SM, MN and KH advised on the experiments. KO

and YS wrote the manuscript with contributions from the other authors.

References

Ames BN. **1966**. [10] Assay of inorganic phosphate, total phosphate and phosphatases.

Methods in Enzymology **8**: 115–118.

Ayadi A, David P, Arrighi JF, Chiarenza S, Thibaud MC, Nussaume L, Marin E.

2015. Reducing the genetic redundancy of arabidopsis phosphate transporter1

transporters to study phosphate uptake and signaling. *Plant Physiology* **167**:

1511–1526.

Balzergue C, Dartevelle T, Godon C, Laugier E, Meisrimler C, Teulon JM, Creff A,

Bissler M, Brouchoud C, Hagège A, *et al*. 2017. Low phosphate activates

STOP1-ALMT1 to rapidly inhibit root cell elongation. *Nat Commun.* doi:

10.1038/ncomms15300.

Baumberger N, Ringli C, Keller B. 2001. The chimeric leucine-rich repeat/extensin

cell wall protein LRX1 is required for root hair morphogenesis in *Arabidopsis*

thaliana. *Genes Dev*. **15**:1128-39. doi: 10.1101/gad.200201.

Bates TR, Lynch JP. **1996**. Stimulation of root hair elongation in *Arabidopsis thaliana*

by low phosphorus availability. *Plant, Cell and Environment* **19**: 529–538.

Briat JF, Rouached H, Tissot N, Gaymard F, Dubos C. 2015. Integration of P, S, Fe,

and Zn nutrition signals in Arabidopsis thaliana: potential involvement of

PHOSPHATE STARVATION RESPONSE 1 (PHR1). *Front Plant Sci.***6**:290.

doi: 10.3389/fpls.2015.00290.

Brouwer M, Lievens B, Van Hemelrijck W, Van Den Ackerveken G, Cammue

BPA, Thomma BPHJ. **2003**. Quantification of disease progression of several

microbial pathogens on *Arabidopsis thaliana* using real-time fluorescence PCR.

FEMS Microbiology Letters **228**: 241–248.

Bustos R, Castrillo G, Linhares F, Puga MI, Rubio V, Pérez-Pérez J, Solano R,

Leyva A, Paz-Ares J. **2010**. A central regulatory system largely controls

transcriptional activation and repression responses to phosphate starvation in

Arabidopsis. *PLoS Genetics* **6**: e1001102.

Chan C, Liao YY, Chiou TJ. 2021. The Impact of Phosphorus on Plant Immunity.

Plant and Cell Physiology **62**: 582–589

Chien PS, Chiang CP, Leong SJ, Chiou TJ. **2018**. Sensing and Signaling of

Phosphate Starvation: From Local to Long Distance. *Plant and Cell Physiology*

59: 1714–1722.

Chowdhury J, Henderson M, Schweizer P, Burton RA, Fincher GB, Little A. **2014**.

Differential accumulation of callose, arabinoxylan and cellulose in

nonpenetrated versus penetrated papillae on leaves of barley infected with

Blumeria graminis f. sp. *hordei*. *New Phytologist* **204**: 650–660.

Datta S, Kim CM, Pernas M, Pires ND, Proust H, Tam T, Vijayakumar P, Dolan L.

2011. Root hairs: Development, growth and evolution at the plant-soil interface.

Plant and Soil **346**: 1–14.

Das AK, Purkait A. 2020. Boron Dynamics in Soil: Classification, Sources, Factors,

Fractions, and Kinetics. *Commun. Soil Sci. Plant Anal.* **51**: 2778–2790.

Dolan L, Duckett CM, Grierson C, Linstead P, Schneider K, Lawson E, Dean C,

Poethig S, Roberts K. **1994**. Clonal relationships and cell patterning in the

root epidermis of Arabidopsis. *Development* **120**: 2465–2474.

Dong X, Hong Z, Sivaramakrishnan M, Mahfouz M, Verma DPS. **2005**. Callose

synthase (CalS5) is required for exine formation during microgametogenesis

and for pollen viability in Arabidopsis. *Plant Journal* **42**: 315–328.

Ellinger D, Voigt CA. **2014**. Callose biosynthesis in Arabidopsis with a focus on

pathogen response: What we have learned within the last decade. *Annals of*

Botany **114**: 1349–1358.

Enns LC, Kanaoka MM, Torii KU, Comai L, Okada K, Cleland RE. **2005.** Two

callose synthases, GSL1 and GSL5, play an essential and redundant role in

plant and pollen development and in fertility. *Plant Molecular Biology* **58**:

333–349.

Gulabani H, Goswami K, Walia Y, Roy A, Noor JJ, Ingole KD, Kasera M, Laha D,

Giehl RFH, Schaaf G, Bhattacharjee S. 2021. Arabidopsis inositol

polyphosphate kinases IPK1 and ITPK1 modulate crosstalk between

SA-dependent immunity and phosphate-starvation responses. *Plant Cell Rep.*

41:347-363.

Karnovsky MJ. 1965. A formaldehyde-glutaraldehyde fixative of high osmolality for

use in electron microscopy. *J Cell Biol* **27**: 137-138A.

Hamburger D, Rezzonico E, Petétot JMDC, Somerville C, Poirier Y. **2002**.

Identification and characterization of the Arabidopsis *PHO1* gene involved in

phosphate loading to the xylem. *Plant Cell* **14**: 889–902.

Han G, Wei X, Dong X, Wang C, Sui N, Guo J, Yuan F, Gong Z, Li X, Zhang Y, *et*

al. **2020**. Arabidopsis ZINC FINGER PROTEIN1 Acts Downstream of GL2 to

Repress Root Hair Initiation and Elongation by Directly Suppressing bHLH

Genes. *The Plant cell* **32**: 206–225.

Han X, Kui M, He K, Yang M, Du J, Jiang Y, Hu Y. **2023**. Jasmonate-regulated root

growth inhibition and root hair elongation. *Journal of Experimental Botany* **74**:

1176–1185.

Holz M, Zarebanadkouki M, Kuzyakov Y, Pausch J, Carminati A. **2018**. Root hairs

increase rhizosphere extension and carbon input to soil. *Annals of Botany* **121**:

61–69.

Isidra-Arellano MC, Delaux PM, Valdés-López O. 2021. The Phosphate Starvation

Response System: Its Role in the Regulation of Plant–Microbe Interactions.

Plant Cell Physiol **62**: 392–400

Jacobs AK, Lipka V, Burton RA, Panstruga R, Strizhov N, Schulze-Lefert P,

Fincher GB. 2003. An Arabidopsis Callose Synthase, GSL5, Is Required for

Wound and Papillary Callose Formation. *Plant Cell* **15**: 2503–2513.

Liu TY, Huang TK, Tseng CY, Lai YS, Lin SI, Lin WY, Chen JW, Chioua TJ.

2012. PHO2-dependent degradation of PHO1 modulates phosphate

homeostasis in *Arabidopsis*. *Plant Cell* **24**: 2168–2183.

Mateo A, Funck D, Mühlenbock P, Kular B, Mullineaux PM, Karpinski S. 2006.

Controlled levels of salicylic acid are required for optimal photosynthesis and

redox homeostasis. *J Exp Bot* **57**: 1795–1807

Mita S, Murano N, Akaike M, Nakamura K. 1997. Mutants of *Arabidopsis thaliana*

with pleiotropic effects on the expression on the gene for β-amylase and on the

accumulation that are inducible by sugars. *Plant Journal* **97**:1087-1093.

Miura K, Lee J, Miura T, Hasegawa PM. 2010. *SIZ1*controls cell growth and plant

development in Arabidopsis through salicylic acid. *Plant Cell Physiol* **51**:

103–113

Miwa K, Wakuta S, Takada S, Ide K, Takano J, Naito S, Omori H, Matsunaga T,

Fujiwara T. 2013. Roles of BOR2, a boron exporter, in cross linking of

rhamnogalacturonan II and root elongation under boron limitation in

Arabidopsis. *Plant Physiology* **163**: 1699–1709.

Mora-Macías J, Ojeda-Rivera JO, Gutiérrez-Alanís D, Yong-Villalobos L,

Oropeza-Aburto A, Raya-González J, Jiménez-Domínguez G,

Chávez-Calvillo G, Rellán-Álvarez R, Herrera-Estrell L. **2017**.

Malate-dependent Fe accumulation is a critical checkpoint in the root

developmental response to low phosphate. *Proceedings of the National*

Academy of Sciences of the United States of America **114**: E3563–E3572.

Morcillo RJ, Singh SK, He D, An G, Vílchez JI, Tang K, Yuan F, Sun Y, Shao C,

Zhang S, ***et al.*** **2020**. Rhizobacterium-derived diacetyl modulates plant

immunity in a phosphate-dependent manner. *The EMBO Journal* **39**: 1–15.

Müller J, Toev T, Heisters M, Teller J, Moore KL, Hause G, Dinesh DC,

Bürstenbinder K, Abel S. **2015**. Iron-Dependent Callose Deposition Adjusts

Root Meristem Maintenance to Phosphate Availability. *Developmental Cell* **33**:

216–230.

Nakagawa T, Suzuki T, Murata S, Nakamura S, Hino T, Maeo K, Tabata R,

Kawai T, Tanaka K, Niwa Y et al. 2007. Improved gateway binary vectors:

High-performance vectors for creation of fusion constructs in transgenic

analysis of plants. Bioscience, Biotechnology and Biochemistry 71:

2095–2100.

Nieves-Cordones M, Lara A, Ródenas R, Amo J, Rivero RM, Martínez V, Rubio F.

2019. Modulation of K+ translocation by AKT1 and AtHAK5 in Arabidopsis

plants. *Plant Cell and Environment* **42**: 2357–2371.

Nishikawa SI, Zinkl GM, Swanson RJ, Maruyama D, Preuss D. **2005**. Callose

(β-1,3 glucan) is essential for Arabidopsis pollen wall patterning, but not tube

growth. *BMC Plant Biology* **5**: 1–9.

Nishimura MT, Stein M, Hou BH, Vogel JP, Edwards H, Somerville SC. **2003**.

Loss of a callose synthase results in salicylic acid-dependent disease resistance.

Science **301**: 969–972.

Okada K, Kubota Y, Hirase T, Otani K, Goh T, Hiruma K, Saijo Y. **2021**.

Uncoupling root hair formation and defence activation from growth inhibition

in response to damage-associated Pep peptides in Arabidopsis thaliana. *New*

Phytologist **229**: 2844–2858.

O'Lexy R, Kasai K, Clark N, Fujiwara T, Sozzani R, Gallagher KL. **2018**.

Exposure to heavy metal stress triggers changes in plasmodesmatal

permeability via deposition and breakdown of callose. *Journal of Experimental*

Botany **69**: 3715–3728.

Omasits U, Ahrens CH, Müller S, Wollscheid B. **2014**. Protter: Interactive protein

feature visualization and integration with experimental proteomic data.

Bioinformatics **30**: 884–886.

Péret B, Clément M, Nussaume L, Desnos T. **2011**. Root developmental adaptation to

phosphate starvation: Better safe than sorry. *Trends in Plant Science* **16**:

442–450.

Péret B, Desnos T, Jost R, Kanno S, Berkowitz O, Nussaume L. **2014**. Root

architecture responses: In search of phosphate. *Plant Physiol* **166**: 1713–1723.

Pfaffl. **2001**. A new mathematical model for relative quantification in real-time

RT-PCR. *Nucleic Acids Research* **29**:2003-2007.

Poirier Y, Thoma S, Somerville C, Schiefelbein J. **1991**. A mutant of Arabidopsis

deficient in xylem loading of phosphate. *Plant Physiology* **97**: 1087–1093.

Poole J, Day CJ, Von Itzstein M, Paton JC, Jennings MP. **2018**. Glycointeractions in

bacterial pathogenesis. *Nature Reviews Microbiology* **16**: 440–452.

Pound MP, French AP, Atkinson JA, Wells DM, Bennett MJ, Pridmore T. **2013**.

RootNav: Navigating images of complex root architectures. *Plant Physiology*

162: 1802–1814.

Robe K, Conejero G, Gao F, Lefebvre-Legendre L, Sylvestre-Gonon E, Rofidal V,

Hem S, Rouhier N, Barberon M, Hecker A, *et al.* **2021**. Coumarin

accumulation and trafficking in Arabidopsis thaliana: a complex and dynamic

process. *New Phytologist* **229**: 2062–2079.

Robertson-Albertyn S, Terrazas RA, Balbirnie K, Blank M, Janiak A, Szarejko I,

Chmielewska B, Karcz J, Morris J, Hedley PE, *et al.* **2017**. Root hair

mutations displace the barley rhizosphere microbiota. *Frontiers in Plant*

Science **8**: 1–15.

Rubio V, Linhares F, Solano R, Martín AC, Iglesias J, Leyva A, Paz-Ares J. **2001**.

A conserved MYB transcription factor involved in phosphate starvation

signaling both in vascular plants and in unicellular algae. *Genes and*

Development **15**: 2122–2133.

Sager R, Lee JY. **2014**. Plasmodesmata in integrated cell signalling: Insights from

development and environmental signals and stresses. *Journal of Experimental*

Botany **65**: 6337–6358.

Schenk ST, Schikora A. **2015**. http://www.bio-protocol.org/e1429 Vol 5, Iss 6, Mar 20,

2015. **5**: 3–6.

Schiefelbein J, Kwak SH, Wieckowski Y, Barron C, Bruex A. **2009**. The gene

regulatory network for root epidermal cell-type pattern formation in

Arabidopsis. *Journal of Experimental Botany* **60**: 1515–1521.

Shin H, Shin HS, Dewbre GR, Harrison MJ. **2004**. Phosphate transport in

Arabidopsis: Pht1;1 and Pht1;4 play a major role in phosphate acquisition from

both low- and high-phosphate environments. *Plant Journal* **39**: 629–642.

Stefanovic A, Ribot C, Rouached H, Wang Y, Chong J, Belbahri L, Delessert S,

Poirier Y. **2007**. Members of the PHO1 gene family show limited functional

redundancy in phosphate transfer to the shoot, and are regulated by phosphate

deficiency via distinct pathways. *Plant Journal* **50**: 982–994.

Sun L, Song L, Zhang Y, Zheng Z, Liu D. **2016**. Arabidopsis PHL2 and PHR1 act

redundantly as the key components of the central regulatory system controlling

transcriptional responses to phosphate starvation. *Plant Physiology* **170**:

499–514.

Svistoonoff S, Creff A, Reymond M, Sigoillot-Claude C, Ricaud L, Blanchet A,

Nussaume L, Desnos T. **2007**. Root tip contact with low-phosphate media

reprograms plant root architecture. *Nat Genet* **39**: 792–796

Tanaka N, Kato M, Tomioka R, Kurata R, Fukao Y, Aoyama T, Maeshima M.

2014. Characteristics of a root hair-less line of *Arabidopsis thaliana* under

physiological stresses. *Journal of Experimental Botany* **65**: 1497–1512.

Thibaud MC, Arrighi JF, Bayle V, Chiarenza S, Creff A, Bustos R, Paz-Ares J,

Poirier Y, Nussaume L. **2010**. Dissection of local and systemic transcriptional

responses to phosphate starvation in Arabidopsis. *Plant Journal* **64**: 775–789.

Verbon EH, Liberman LM, Zhou J, Yin J, Pieterse CMJ, Benfey N, Stringlis IA,

Jonge R De. **2023**. Cell type-specific transcriptomics reveals that root hairs

and endodermal barriers play important roles in beneficial

plant-rhizobacterium-interactions. *MOLECULAR PLANT* doi:

https://doi.org/10.1016/j.molp.2023.06.001.

Vissenberg K, Claeijs N, Balcerowicz D, Schoenaers S. **2020**. Hormonal regulation of

root hair growth and responses to the environment in Arabidopsis. *Journal of*

Experimental Botany **71**: 2412–2427.

Vogel J, Somerville S. 2000. Isolation and characterization of powdery

mildew-resistant *Arabidopsis* mutants. *PNAS* **97**: 1897–1902.

Wang X, Du G, Wang X, Meng Y, Li Y, Wu P, Yi K. **2010**. The function of LPR1 is

controlled by an element in the promoter and is independent of SUMO E3

ligase SIZ1 in response to Low Pi stress in *Arabidopsis thaliana*. *Plant and*

Cell Physiology **51**: 380–394.

Wang Z, Zheng Z, Song L, Liu D. **2018**. Functional characterization of Arabidopsis

PHL4 in plant response to phosphate starvation. *Frontiers in Plant Science* **9**:

1–19.

Wang Y, Li X, Fan B, Zhu C, Chen Z. 2021. Regulation and function of

defense-related callose deposition in plants. *Int J Mol Sci* **22**: 1–15.

Weßling R, Panstruga R. **2012**. Rapid quantification of plant-powdery mildew

interactions by qPCR and conidiospore counts. *Plant Methods* **8**.

Wildermuth MC, Dewdney J, Wu G, Ausubel FM. **2002**. Erratum: corrigendum:

Isochorismate synthase is required to synthesize salicylic acid for plant defense.

Nature **417**: 571–571.

Wu SW, Kumar R, Iswanto ABB, Kim JY. **2018**. Callose balancing at

plasmodesmata. *Journal of Experimental Botany* **69**: 5325–5339.

Zhang X, Chen S, Mou Z. 2010. Nuclear localization of NPR1 is required for

regulation of salicylate tolerance, isochorismate synthase 1 expression and

salicylate accumulation in Arabidopsis. *J Plant Physiol* **167**: 144–148

Zhu C, Gan L, Shen Z, Xia K. **2006**. Interactions between jasmonates and ethylene in

the regulation of root hair development in Arabidopsis. *Journal of*

Experimental Botany **57**: 1299–1308.

Figure Legends

Fig. 1 Callose deposition is induced in root hairs under phosphate deficiency,

independently of *PHR1, PHL1, LPR1* and *LPR2*.

Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis*

thaliana seedlings to low Pi (P 50 µM), low N (N 0 µM), low Fe (Fe 0 µM), low Fe low

Pi (Fe 0 µM, P 0 µM) and normal Pi (P 625 µM) liquid media. (a, c, left) Bright field

(BF) images. (a, c, right) Aniline blue staining images. White arrowheads indicate root

hair tips. (b) Aniline blue and propidium iodide (PI) staining of root hairs under low Pi

and normal Pi conditions. (c) The wild-type (WT) used for *rhd6* was Ws-2. Magnified

images of the upper panels (a, c, lower panel). Bars, 500 µm (a, c, upper panel), 50 µm

(a, c lower panel) and 25 µm (b).

Fig. 2 Callose synthase *PMR4* is required for callose induction in root hairs under

phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis*

thaliana plants to low Pi (0 µM) liquid media. Left, bright filed (BF) images. Right,

aniline blue staining images. Two *caps* mutants are defective in inducing root hair

callose under phosphate starvation. Lower panels, magnified images of upper panels.

White arrowheads indicate root hair tips. Bars, 500 µm (upper panel) and 50 µm (lower

panel). (b) Positions of mutated sites in the *PMR4* locus of *caps* mutants, illustrated

with Protter (Omasits *et al.*, 2014). An orange region corresponds to the catalytic

domain of PMR4. Asterisks indicate stop codons. (c) Leaves 10 d after inoculation with

Golovinomyces orontii. Bar, 2 cm. (d) Ratios of *G. orontii* genomic DNA to

Arabidopsis genomic DNA were determined by qPCR with the primers for Plasma

membrane ATPase 1 (Go_V1_Conting3757) and RNA-binding family protein

(AT3G21215), respectively. Data show means ± SD (n=3). Asterisks (** $P < 0.01$)

indicate statistically significant differences relative to WT according to Student's *t*-test.

Fig. 3 *PMR4* modulates root and shoot growth under phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis*

seedlings to low Pi (50 µM) liquid media. Left, bright filed (BF) images. Right, aniline

blue staining images. Lower panels, magnified images of upper panels. White

arrowheads indicate root hair tips. Bars, 100 µm (upper panel) and 50 µm (lower panel).

(b) Six-d-old seedlings were exposed to low Pi (50 µM) or normal Pi (625 µM) media

for 6 d. Bar, 10 mm. (c) The primary roots 6 d after transfer of 6-d-old seedling to low

Pi (50 µM) and normal Pi media. Bar, 1 mm. (d) Root hair density and root hair length

in (c). (e) Primary root length in (b). (f) Ratio of the primary root length under normal

Pi relative to low Pi of (e). (g) Convex hull area of the roots in (b). (h) Ratio of convex

hull area under normal Pi relative to low Pi of (g). (i) Six-d-old seedlings were exposed

to low Pi (50 µM) or normal Pi for 16 d. Bar, 10 mm. (j) SFW in (i). (k) Six-d-old

seedlings were exposed to low Pi (10 µM) media for 10 d. Bar, 10 mm. (l) Anthocyanin

contents in (k). (d-h, j, l) Data show means ± SD (n = 7~30). Statistical significance was

assessed by one-way ANOVA and Tukey honestly significant different (HSD) test for

each medium dataset. Different letters indicate significant differences ($P < 0.05$). NS,

not significant.

Fig. 4 *PMR4* contributes to shoot growth and Pi accumulation under phosphate

deficiency.

(a-d) Pi contents per root or shoot fresh weight (FW) of 6-d-old *Arabidopsis* seedlings

exposed to low Pi (150 µM) or normal Pi (625 µM) for 9 d. The WT used for *rhd6* was

Ws-2 (c,d). (e) Illustration of Pi accumulation in the indicated genotypes, referred to

(a-d). (f) ^{32}P uptake rate determined in the whole seedlings exposed to the indicated

media (low Pi and normal Pi, 50 and 625 µM Pi, respectively) supplemented with

^{32}P-labelled Pi for 30 min. (g) Ratio of ^{32}P in the shoots relative to the whole plants after

pulse-chase labeling. Seedlings grown on low and normal Pi media (50 and 625 µM,

respectively) were exposed to ^{32}P-labeled Pi for 10 min, and then transferred to

non-labelled Pi media for 120 min (pulse chase). The ^{32}P ratio of the shoots to the whole

seedlings was calculated for the translocation rate of Pi. Data show means ± SD (n =

3~4) in (a-d, f-g). Statistical significance was assessed by one-way ANOVA and Tukey

HSD test for each medium dataset (a-b, f-g). Different letters indicate significant

differences ($P < 0.05$). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate statistically

significant differences relative to WT according to Student's *t*-test (c, d). NS, not

significant.

Fig. 5 Plant element contents in shoot and root under low phosphate conditions.

Element contents (P, K, Mg, Ca, Mn, Fe, Zn, Mo and B) per root or shoot dry weight

(DW) of 5-d-old *Arabidopsis thaliana* seedlings exposed to low Pi (10 μM) for 4 d.

Data show means ± SD (n=4~8). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate

statistically significant differences relative to WT according to Student's *t*-test. NS, not

significant. ND, no data.

Fig. 6 Root *PMR4* contributes to shoot growth under phosphate deficiency.

(a) Reciprocal micrografting of *Arabidopsis* WT and *pmr4* mutants. Images were

obtained at 14 d after transfer to low Pi (150 μM) media. Bar, 10 mm. (b) Shoot fresh

weight (SFW) of (a). (c) The primary root length of (a). (d) Aniline blue staining of

grafted-seedlings at 9 d after transfer to low Pi (50 μM). (d, left) Bright filed (BF) images.

(d, right) Aniline blue staining images. (d, lower panel) Magnified images of (d, upper

panel). White arrows indicate root hair tips. Bar, 100 μm (c, upper panel) or 20 μm (c,

lower panel). (b, c) Data show means ± SD (n=19-26). Statistical significance was

assessed by one-way ANOVA and Tukey HSD test for each medium dataset. Different

letters indicate significant differences ($P < 0.05$).

Fig. 7 PMR4 in root hairs contributes to shoot growth under phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis*

seedlings to low Pi (10 μM). White arrowheads indicate root hair chips. Bars, 100 μm

(upper panel) and 50 µm (lower panel). (b) Five-d-old seedlings were exposed to low Pi

(50 µM) for 10 d. Bar, 10 mm. (c) SFW of (b). (d) Primary root length of (b). (c, d)

Data show means ± SD (n = 22-48). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate

statistically significant differences relative to *pmr4* according to Student's *t*-test. NS, not

significant.

Fig. 8 Model for PMR4-mediated root hair callose deposition in plant adaptation

to Pi deficiency

PMR4 induces callose deposition in root hairs under phosphate deficiency, and thus

contributes to shoot Pi accumulation and growth through enhancing Pi uptake and

root-to-shoot transfer. This contribution alleviates phosphate starvation symptoms, such

as root system area reduction and anthocyanin accumulation.

Supporting information

Fig. S1 Callose deposition in *gsl* mutants under phosphate deficiency

Fig. S2 flg22-induced callose deposition is impaired in the *caps* and *pmr4* alleles.

Fig. S3 Meristematic region is reduced in *pmr4* mutants under phosphate

deficiency.

Fig. S4 *PMR4* restricts *LPR1/LPR2*-mediated inhibition of the primary root growth

in response to low Pi.

Fig. S5 Plant growth is lowered in *pmr4* on soil.

Fig. S6 Plant element contents in shoot and root under phosphate sufficiency.

Fig. S7 Primary root growth inhibition under B deficiency.

Table S1 *Arabidopsis thaliana* mutants used in this study

Table S2 DNA Primers used in DNA vector construction and qPCR analyses

Fig.1 Callose deposition is induced in root hairs under phosphate deficiency, independently of *PHR1*, *PHL1*, *LPR1* and *LPR2*.

Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis thaliana* seedlings to low Pi (P 50 µM), low N (N 0 µM), low Fe (Fe 0 µM), low Fe low Pi (Fe 0 µM, P 0 µM) and normal Pi (P 625 µM) liquid media. (a, c, left) Bright field (BF) images. (a, c, right) Aniline blue staining images. White arrowheads indicate root hair tips. (b) Aniline blue and propidium iodide (PI) staining of root hairs under low Pi and normal Pi conditions. (c) The wild-type (WT) used for *rhd6* was Ws-2. Magnified images of the upper panels (a, c, lower panel). Bars, 500 µm (a, c, upper panel), 50 µm (a, c lower panel) and 25 µm (b).

Fig. 2. Callose synthase *PMR4* is required for callose induction in root hairs under phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis thaliana* plants to low Pi (0 µM) liquid media. Left, bright filed (BF) images. Right, aniline blue staining images. Two *caps* mutants are defective in inducing root hair *callose under phosphate starvation.* Lower panels, magnified images of upper panels. White arrowheads indicate root hair tips. Bars, 500 µm (upper panel) and 50 µm (lower panel). (b) Positions of mutated sites in the *PMR4* locus of *caps* mutants, illustrated with Protter (Omasits *et al.*, 2014). An orange region corresponds to the catalytic domain of PMR4. Asterisks indicate stop codons. (c) Leaves 10 d after inoculation with *Golovinomyces orontii*. Bar, 2 cm. (d) Ratios of *G. orontii* genomic DNA to Arabidopsis genomic DNA were determined by qPCR with the primers for Plasma membrane ATPase 1 (Go_V1_Conting3757) and RNA-binding family protein (AT3G21215), respectively. Data show means ± SD (n=3). Asterisks (** $P < 0.01$) indicate statistically significant differences relative to WT according to Student's *t*-test.

Fig. 3 *PMR4* modulates root and shoot growth under phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis* seedlings to low Pi (50 µM) liquid media. Left, bright filed (BF) images. Right, aniline blue staining images. Lower panels, magnified images of upper panels. White arrowheads indicate root hair tips. Bars, 100 µm (upper panel) and 50 µm (lower panel). (b) Six-d-old seedlings were exposed to low Pi (50 µM) or normal Pi (625 µM) media for 6 d. Bar, 10 mm. (c) The primary roots 6 d after transfer of 6-d-old seedling to low Pi (50 µM) and normal Pi media. Bar, 1 mm. (d) Root hair density and root hair length in (c). (e) Primary root length in (b). (f) Ratio of the primary root length under normal Pi relative to low Pi of (e). (g) Convex hull area of the roots in (b). (h) Ratio of convex hull area under normal Pi relative to low Pi of (g). (i) Six-d-old seedlings were exposed to low Pi (50 µM) or normal Pi for 16 d. Bar, 10 mm. (j) SFW in (i). (k) Six-d-old seedlings were exposed to low Pi (10 µM) media for 10 d. Bar, 10 mm. (l) Anthocyanin contents in (k). (d-h, j, l) Data show means ± SD (n = 7~30). Statistical significance was assessed by one-way ANOVA and Tukey honestly significant different (HSD) test for each medium dataset. Different letters indicate significant differences ($P < 0.05$). NS, not significant.

Fig. 4 *PMR4* contributes to shoot growth and Pi accumulation under phosphate deficiency

(a-d) Pi contents per root or shoot fresh weight (FW) of 6-d-old *Arabidopsis* seedlings exposed to low Pi (150 μM) or normal Pi (625 μM) for 9 d. The WT used for *rhd6* was Ws-2 (c,d). (e) Illustration of Pi accumulation in the indicated genotypes, referred to (a-d). (f) ^{32}P uptake rate determined in the whole seedlings exposed to the indicated media (low Pi and normal Pi, 50 and 625 μM Pi, respectively) supplemented with ^{32}P-labelled Pi for 30 min. (g) Ratio of ^{32}P in the shoots relative to the whole plants after pulse-chase labeling. Seedlings grown on low and normal Pi media (50 and 625 μM, respectively) were exposed to ^{32}P-labeled Pi for 10 min, and then transferred to non-labelled Pi media for 120 min (pulse chase). The ^{32}P ratio of the shoots to the whole seedlings was calculated for the translocation rate of Pi. Data show means ± SD (n = 3~4) in (a-d, f-g). Statistical significance was assessed by one-way ANOVA and Tukey HSD test for each medium dataset (a-b, f-g). Different letters indicate significant differences ($P < 0.05$). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate statistically significant differences relative to WT according to Student's *t*-test (c, d). NS, not significant.

Fig. 5 Plant element contents in shoot and root under low phosphate conditions.

Element contents (P, K, Mg, Ca, Mn, Fe, Zn, Mo and B) per root or shoot dry weight (DW) of 5-d-old *Arabidopsis thaliana* seedlings exposed to low Pi (10 μM) for 4 d. Data show means ± SD (n=4~8). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate statistically significant differences relative to WT according to Student's *t*-test. NS, not significant. ND, no data.

Fig. 6 Root *PMR4* contributes to shoot growth under phosphate deficiency.

(a) Reciprocal micrografting of *Arabidopsis* WT and *pmr4* mutants. Images were obtained at 14 d after transfer to low Pi (150 µM) media. Bar, 10 mm. (b) Shoot fresh weight (SFW) of (a). (c) The primary root length of (a). (d) Aniline blue staining of grafted-seedlings at 9 d after transfer to low Pi (50 µM). (d, left) Bright filed (BF) images. (d, right) Aniline blue staining images. (d, lower panel) Magnified images of (d, upper panel). White arrows indicate root hair tips. Bar, 100 µm (c, upper panel) or 20 µm (c, lower panel). (b, c) Data show means ± SD (n=19-26). Statistical significance was assessed by one-way ANOVA and Tukey HSD test for each medium dataset. Different letters indicate significant differences ($P < 0.05$).

Fig. 7 PMR4 in root hairs contributes to shoot growth under phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis* seedlings to low Pi (10 μM). White arrowheads indicate root hair chips. Bars, 100 μm (upper panel) and 50 μm (lower panel). (b) Five-d-old seedlings were exposed to low Pi (50 μM) for 10 d. Bar, 10 mm. (c) SFW of (b). (d) Primary root length of (b). (c, d) Data show means ± SD (n = 22-48). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate statistically significant differences relative to *pmr4* according to Student's *t*-test. NS, not significant.

Fig. 8 Model for PMR4-mediated root hair callose deposition in plant adaptation to Pi deficiency

PMR4 induces callose deposition in root hairs under phosphate deficiency, and thus contributes to shoot Pi accumulation and growth through enhancing Pi uptake and root-to-shoot transfer. This contribution alleviates phosphate starvation symptoms, such as root system area reduction and anthocyanin accumulation.