1 Defense-related callose synthase *PMR4* promotes root hair callose deposition and

2 adaptation to phosphate deficiency in *Arabidopsis thaliana*

Kentaro Okada[1,2], Koei Yachi[1], Tan Anh Nhi Nguyen[1], Satomi Kanno[3], Shigetaka

Yasuda[1], Haruna Tadai[1], Chika Tateda[1], Tae-Hong Lee[1], Uyen Nguyen[4], Kanako Inoue[1],

Natsuki Tsuchida[1], Taiga Ishihara[1], Shunsuke Miyashima[1], Kei Hiruma[1], Kyoko Miwa[5],

Takaki Maekawa[4, 6], Michitaka Notaguchi[2], Yusuke Saijo[1]

[1]Graduate School of Science and Technology, Nara Institute of Science and Technology,

Takayama-cho, Ikoma 630-0192, Japan

[2]Bioscience and Biotechnology Center, Nagoya University, Furo-cho, Chikusa-ku,

Nagoya 464-8601, Japan.

[3]Institute for Advanced Research, Nagoya University, Furo-cho, Chikusa, Nagoya

464-8601, Japan

[4]Institute for Plant Sciences, University of Cologne, Cologne D-50674, Germany

[5]Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810,

Japan

[6]CEPLAS Cluster of Excellence on Plant Sciences at the University of Cologne,

Cologne D-50674, Germany

Author for correspondence: Yusuke Saijo

ORCID ID: 0000-0001-8985-9695

Tel: +81-743-72-5690

Email: saijo@bs.naist.jp

Summary

Plants acquire phosphorus (P) primarily as inorganic phosphate (Pi) from the soil.

Under Pi deficiency, plants induce an array of physiological and morphological

responses, termed phosphate starvation response (PSR), thereby increasing Pi

acquisition and use efficiency. However, the mechanisms by which plants adapt to Pi

deficiency remain to be elucidated. Here, we report that deposition of a β-1,3-glucan

polymer called callose is induced in *Arabidopsis thaliana* root hairs under Pi deficiency,

in a manner independent of PSR-regulating *PHR1/PHL1* transcription factors and

LPR1/LPR2 ferroxidases. Genetic studies revealed *PMR4* (*GSL5*) callose synthase

being required for the callose deposition in Pi-depleted root hairs. Loss of *PMR4* also

reduces Pi acquisition in shoots and plant growth under low Pi conditions. The defects

are not recovered by simultaneous disruption of *SID2*, mediating defense-associated

salicylic acid (SA) biosynthesis, excluding SA defense activation from the cause of the

observed *pmr4* phenotypes. Grafting experiments and characterization of plants

expressing *PMR4* specifically in root hair cells suggest that a PMR4 pool in the cell

type contributes to shoot growth under Pi deficiency. Our findings thus suggest an

important role for *PMR4* in plant adaptation to Pi deficiency.

Significance statement

We reveal that PMR4 callose synthase mediates callose deposition in root hairs under

phosphate (Pi) deficiency, without requiring Pi starvation response regulators

PHR1/PHL1 or *LPR1/LPR2*. The loss of the callose deposition is accompanied by

decreases in Pi acquisition and plant growth in *pmr4*. Root hair cell-specific *PMR4*

expression restores callose deposition in root hairs and shoot growth under Pi deficiency,

indicating a critical role for root hair callose in plant adaptation to Pi deficiency.

Keywords: root hair, cell wall, Glucan Synthase Likes (GSLs), phosphate acquisition,

phosphate transport, plant nutrition, nutrient deficiency, salicylic acid (SA), pattern

triggered immunity

Introduction

As an essential macronutrient, plants acquire and distribute P from the rhizosphere to

the whole tissues, primarily in the form of inorganic phosphate (Pi). To maintain Pi

homeostasis under Pi deficiency, plants induce an array of adaptive responses, *e.g.*

increasing Pi uptake, remobilization and utilization, termed Pi starvation response

(PSR), and involve beneficial microbes in mutualistic associations (Péret et al., 2011;

Chien et al., 2018; Isidra-Arellano et al., 2021). PSR is accompanied by extensive

transcriptional reprogramming, mainly through the transcription factor *PHR1* and

related *PHL1*, which leads to increases in Pi transporter expression, root hair formation

and anthocyanin accumulation (Rubio et al., 2001, Bustos et al., 2010). *PHR1/PHL1*

mediate systemic coordination of these responses according to the internal Pi

availability. In addition, Pi depletion leads to remodeling root system architecture, a

decrease in primary root growth while increasing lateral root formation and growth,

through the ferroxidases *LPR1* and *LPR2* (Svistoonoff et al., 2007; Péret et al., 2014;

Balzergue et al., 2017). *LPR1/LPR2* are required for arresting primary root growth when

the external Pi levels are reduced surrounding the root tips (Svistoonoff et al., 2007).

Recent studies illuminated an intimate relationship between PSR and biotic interactions,

but the molecular basis for their coordination remains poorly understood (Chan et al.,

2021).

Callose deposition on the plasma membrane through callose synthases *Glucan*

Synthase Likes (*GSLs*) reinforces barrier functions of the cell wall and restricts

plasmodesmatal permeability, in plant development and biotic/abiotic stress responses

(Jacobs et al., 2003; Nishimura et al., 2003; Dong et al., 2005; Enns et al., 2005;

Nishikawa et al., 2005; Chowdhury et al., 2014; Ellinger & Voigt, 2014; Sager & Lee,

2014; Wu et al., 2018). In *Arabidopsis thaliana* (hereafter Arabidopsis)*,* callose

deposition is induced in the root phloem in response to excess iron, thereby reducing

plasmodesmatal permeability (O'Lexy et al., 2018). Of 12 Arabidopsis *GSLs*, *GSL2*

(*CALS5*) and *GSL5* (*PMR4, CALS12*) respectively contribute to basal accumulation and

iron-induced deposition of callose in the root phloem (O'Lexy et al., 2018). Callose also

accumulates in the stem cell niche (SCN) surrounding the quiescent center cells, thereby

inhibiting cell-to-cell movement of the transcription factor SHORT ROOT (SHR)

during PSR. This serves to suppress cell division and elongation in the root apical

meristem and elongation zone, respectively, through the *LPR1/LPR2* pathway (Müller

et al., 2015; Mora-Macías et al., 2017). *GSL5* (*PMR4*) also mediates callose deposition

in response to pathogen challenge and microbe/damage-associated molecular patterns

(MAMPs/DAMPs), and promotes fungal penetration resistance (Wang et al., 2021).

However, the loss of *GSL5* function results in strong resistance to powdery mildew

fungi through tightened SA-based defenses, demonstrated by *gsl5* mutants revealed as

powdery mildew resistance4 (*pmr4*) (Vogel & Somerville, 2000; Nishimura et al., 2003).

This resistance is thought to represent a backup defense activated when *GSL5*

(*PMR4*)-mediated callose deposition, a key output of MAMP/DAMP-triggered defense

signaling, is disrupted.

Although the significance of *PMR4* has been genetically shown, our knowledge is

very limited for the mechanisms that involve and regulate PRM4 in callose deposition

during defense responses (Wang et al., 2021). *PMR4/GSL5* is expressed ubiquitously in

unwounded *Arabidopsis thaliana* plants (Enns et al., 2005). Inoculation with

Plectosphaerella cucumerina fungi results in a slight increase in *GSL5* expression

(García-Andrade et al., 2011). Downy mildew *Hyaloperonospora* and the application of

SA and flg22 induce *GSL5* expression (Dong et al., 2008; Keppler et al., 2018).

Whether and if so how GSL5 expression or activity is regulated according to the

nutrient status remains poorly understood.

Here, we report that callose deposition is induced in root hairs under Pi deficiency, in

a manner independent of *PHR1/PHL1* and *LPR1/LPR2*. Our genetic studies revealed

that the Pi depletion-induced callose deposition is impaired in *pmr4* (*gsl5*) root hairs.

The loss of *PMR4*-mediated callose deposition is associated with reduced plant growth

as well as exaggerated stress symptoms under Pi deficiency, in addition with enhanced

powdery mildew resistance. The results indicate *PMR4*-mediated callose deposition in

root hairs as an important step in plant adaptation to Pi-limiting conditions.

Results

Callose deposition occurs in root hairs under phosphate deficiency, independently

of *PHR1/PHL1* and *LPR1/LPR2*

Under low Pi conditions (Pi at 0 or 50 µM), we noticed callose deposition in the

epidermal cells, particularly in root hairs of the differentiation zone (Figure 1a, b;

O'Lexy et al., 2018). However, callose deposition was not detected in root hairs under

the tested N, Fe or Fe/Pi deficient conditions (Figure 1a). The results indicate that the

callose deposition is specific to Pi deficiency and is dependent on Fe, consistent with

the interplay between Fe and Pi (Müller., 2015; Abel, 2017; Xue et al., 2022). The

absence of callose staining in the epidermis of root hair-less *rhd6* mutants (Figure 1c)

validated that callose deposition occurs in root hairs under low Pi. In contrast to

previous study (Müller et al., 2015), we did not detect callose deposition in the root

SCN and cortex cells of the elongation zone under our experimental conditions (Figure

1a, c).

We next assessed whether *PHR1/PHL1* and *LPR1/LPR2* play a role in the root hair

callose deposition. Unlike *LPR1*/*LPR2*-mediated callose deposition in SCN (Müller et

al., 2015), the root hair callose deposition was not affected in *lpr1 lpr2* (Figure 1c)

despite its Fe dependence (Figure 1a). Furthermore, wild type (WT)-like callose

deposition was retained in root hairs of *phr1 phl1 lpr1 lpr2* under low Pi conditions

(Figure 1c). The results indicate that callose deposition is induced in root hairs under Pi

deficiency, independently of these PSR regulators, and its separation from callose

deposition in the root SCN.

***PMR4* callose synthase contributes to callose deposition under Pi deficiency**

To unravel genetic requirements for the callose deposition in root hairs under Pi

deficiency, we examined *glucan synthase like* (*gsl*) mutants for alternations in callose

deposition under phosphate deficiency. Of the tested *gsl* mutants, *gsl5* (hereafter, *pmr4*)

mutants specifically lacked callose deposition in root hairs (Figure S1). We also

screened a total of 18,550 ethyl methane sulfonate (EMS)-mutagenized individuals in

Col-0 and Ler backgrounds. This revealed two mutants in the Col-0 background,

designated <u>*c*allose *a*lteration under *P*i *s*tarvation 1 and 2</u> (*caps1 and caps2*, Figure 2a).

caps1 and *caps2* mutants were severely defective in callose deposition of root hairs

under low Pi conditions (Figure 2a). DNA Sequencing on the *PMR4* genomic region

revealed a DNA substitution within the coding region, resulting in a non-synonymous

amino acid substitution, in both *caps1* and *caps*2 alleles (Figure 2b). The substitution in

*caps*1 converts Tryptophan to Stop codon at the 260[th] amino acid (PMR4^{W260*}),

resulting in PMR4 truncation, consistent with the severely impaired callose deposition

in *caps1* as well as *pmr4-1* (PMR4^{W687*}). In *caps2*, glutamic acid is substituted with

lysine at the 1181st amino acid (PMR4^{E1181K}), located within the cytosolic side of a

predicted glucan-synthase domain (residue 575 to 1558, orange-marked domain, Figure

2b). The results suggest that dysfunction of PMR4 fails to induce root hair callose

deposition under Pi deficiency in these *caps* alleles. We validated that leaf callose

deposition in response to the bacterial MAMP flg22, a *PMR4*-dependent output in

pattern-triggered immunity (PTI), is abolished in these *caps* alleles (Figure S2).

Moreover, we showed strong resistance to powdery mildew fungi in these *caps* plants

(Figure 2c), as previously described in *pmr4* (Vogel & Somerville, 2000; Nishimura et

al., 2003). The results confirm that PMR4 functions are impaired in both *caps* alleles.

***PMR4* influences root development traits induced under Pi deficiency**

By using *pmr4-1* as a representative mutant allele, we further investigated the

biological significance of *PMR4* under Pi deficiency. Recent studies noted that some of

Pi deficiency-induced root responses are limited to the plate-grown plants whose roots

are light irradiated (Zheng et al., 2019; Yeh et al., 2020; Gao et al., 2021). To test this

possibility, we examined callose deposition in Pi-deplete and Pi-replete plants under

darkness. At 3 d after the transfer of seedlings to the different Pi conditions under

darkness, we detected callose deposition in non-irradiated root hairs of WT but not

pmr4 plants specifically under low Pi, validating that PMR4-mediated, root hair callose

deposition is not a consequence of light exposure to the root (Figure S3a). Moreover,

we also detected callose deposition in the cotyledons of WT but not *pmr4* plants under

these conditions (Figure S3b). The results strengthen the physiological relevance of

PMR4-mediated callose deposition under low Pi.

The loss of *PMR4* leads to an increase in salicylic acid (SA) biosynthesis and defense

under biotic stress (Vogel & Somerville, 2000; Nishimura et al., 2003), in a manner

requiring *SID2 isochorismate synthase* that mediates a critical step in SA biosynthesis

during defense activation (Wildermuth et al., 2001). Given the negative impact of SA

defense activation on plant growth (Mateo et al., 2006; Zhang et al., 2010), we assessed

the possible SA dependence of the observed *pmr4* phenotypes, by examining genetic

interactions between *pmr4* and *sid2*. Callose deposition was induced in *sid2* root hairs

but not induced in *pmr4 sid2* under low Pi conditions (Figure 3a), demonstrating that

SID2-mediated SA biosynthesis is not required for *PMR4*-mediated callose deposition

in root hairs. The results also exclude over-activation of SA defense as the cause of the

callose defects in *pmr4* root hairs.

In Arabidopsis, root hair cells specifically occur at the junction of two underlying

cortical cells, whereas non-hair cells occur in the other epidermal cells (Dolan et al.,

1994; Schiefelbein et al., 2009). Root hair density was increased under low Pi

conditions compared to normal Pi conditions, and the increase was exaggerated in *pmr4*

as well as *pmr4 sid2*, without ectopic root hair formation (Figures 3b-d, S4a). Likewise,

root hair length was also increased in *pmr4* as well as *pmr4 sid2* under both Pi

conditions (Figure 3c, d). The results suggest that *PMR4* is not required for the

differentiation or outgrowth of root hair cells, but that *PMR4* contributes to restricting

root hair formation and elongation, in the presence or absence of *SID2*. Conversely,

primary root length was reduced in *pmr4* compared with WT in both low and normal Pi

conditions (Figure 3b, e). We then determined the ratio of the primary root length under

normal Pi versus low Pi as an indicator for the primary root growth inhibition under Pi

deficiency. The ratio (normal Pi/low Pi) was increased in *pmr4* compared with WT,

indicating that primary root growth inhibition, a Pi deficiency-characteristic response, is

exaggerated in *pmr4* (Figure 3f). We also observed a decrease in the root meristem

region in *pmr4* compared with WT under Pi deficiency, suggesting that the meristem

exhaustion coincides with the inhibition of primary root growth in *pmr4* (Figure S4b).

The primary root growth inhibition under low Pi was alleviated in *sid2*, but it was again

exaggerated in *pmr4 sid2* compared to *sid2* (Figure 3e, f), indicating that the loss of

PMR4 enhances the Pi deficiency response, with or without *SID2*.

We next tested whether and if so how *PMR4* affects the expression of Pi

deficiency-inducible genes. Quantitative RT-PCR analyses on WT and *pmr4* roots

revealed that the expression of Pi starvation-induced genes, *AT4* and *IPS1* long

non-coding RNAs (Franco-Zorrilla et al., 2007), *PHT1;5* Pi transporter (Nagarajan et al.,

2011) and *SPX1* Pi-binding suppressor of PHR1 (Puga et al., 2014), was not affected in

pmr4 under the normal (625 µM) and low Pi (50 µM and 10 µM) conditions (Figure S5).

The results are consistent with WT-like callose deposition in the root hairs of *phr1 phl1*

and *phr1 phl1 lpr1 lpr2* (Figure 1c) and indicate that *PMR4* is not required for the

expression of these PSR-related genes.

The primary root growth inhibition under Pi deficiency relies on *LPR1* and *LPR2*

(Svistoonoff et al., 2007; Müller et al., 2015). We examined the possible relationship

between *LPR1/LPR2* and *PMR4* in the regulation of primary root growth. The primary

root length was reduced in *pmr4 lpr1 lpr2* compared with *lpr1 lpr2* under both Pi

conditions, whereas it was increased in *pmr4 lpr1 lpr2* compared with *pmr4* specifically

under low Pi (Figure S6a, b). The results indicate that *PMR4* serves to sustain the

primary root growth irrespective of the Pi availability or *LPR1/LPR2* activity, whereas

the *LPR1/LPR2* inhibition is specific under low Pi with or without *PMR4*. The ratio of

the primary root length under normal Pi versus low Pi was lowered to a similar extent in

lpr1 lpr2 and *pmr4 lpr1 lpr2* compared with WT (Figure S6c), despite the decrease in

the primary root length of *pmr4* (Figure S6b), indicating that *lpr1 lpr2* is epistatic to

pmr4. Consistently, under Pi deficiency, *PMR4* expression was reduced in *lpr1 lpr2,* but

not in *phr1 phl1*, compared to WT roots (Figure S7). Together with the WT-like callose

deposition in *lpr1 lpr2* root hairs (Figure 1c), the data suggest that *PMR4* expression is

dependent on *LPR1/LPR2* but that the residual *PMR4* expression is sufficient for the

callose deposition in *lpr1 lpr2* root hairs.

We next determined the convex hull root area as a proxy of overall root system

growth, which was correlated with Pi availability (Figure 3g). It was reduced in *pmr4*

and *sid2* compared with WT under both normal and low Pi conditions, albeit to a lesser

extent in *sid2* than *pmr4* under low Pi (Figure 3g), suggesting that *PMR4* and *SID2* both

contribute to expanding the root area. The ratio of the root area under normal Pi versus

low Pi conditions was increased in *pmr4* relative to WT in a manner dependent on *SID2*

(Figure 3h), suggesting that *SID2* contributes to low Pi-induced root growth inhibition

in *pmr4*. Consistently, the root area was greater in *pmr4 sid2* compared to *pmr4* under

low Pi (Figure 3g). The results suggest that SID2 negatively impacts root expansion

specifically in the absence of PMR4 under Pi deficiency. The results also imply that

SID2-mediated SA biosynthesis promotes root growth expansion in Pi-depleted plants

when PMR4 is present. The latter seems to be consistent with a previously described

uncoupling of SA induction from defense activation under Pi deficiency (Gulabani et al.

2021).

PMR4 serves to sustain shoot growth and restrict anthocyanin accumulation under

Pi deficiency.

We next examined whether _PMR4_ influences shoot growth, by determining shoot

fresh weight (FW) under Pi deficiency and sufficiency. Shoot FW was lowered in _pmr4_

compared with WT, and it was not even partially restored by the simultaneous loss of

SID2 in _pmr4 sid2_ under low-Pi media or soil (Figures 3i-j and S8). Together with the

root phenotypes described above, the results exclude _SID2_ as the major cause of the

growth inhibition in _pmr4_. Although high SA accumulation was previously described to

reduce plant growth (Miura et al., 2010), we did not detect negative impacts of _SID2_ on

shoot growth under low Pi conditions (Figure 3j). _SID2_ even positively influenced shoot

growth under the normal Pi conditions, indicated by the reduced shoot biomass in _sid2_

compared to WT (Figure 3j).

Anthocyanin content was increased in _pmr4_ compared with WT, in the presence or

absence of _SID2_ (Figure 3k, l). Although SA was described to enhance anthocyanin

accumulation under Pi deficiency (Morcillo et al., 2019), _sid2_ was largely

indistinguishable from WT in the anthocyanin contents under our low Pi conditions

(Figure 3k, l). The results imply that PMR4-mediated Pi acquisition and allocation

serves to alleviate Pi starvation causing anthocyanin accumulation under Pi deficiency.

PMR4 promotes the acquisition and root-to-shoot translocation of Pi

The _PMR4_ effects on shoot and root growth prompted us to examine its possible role

in Pi acquisition from the rhizosphere and Pi mobilization within the plant. We found

that shoot Pi concentrations were greatly reduced in _pmr4_ compared with WT, without

alterations in root Pi concentrations, under Pi deficiency (Figure 4a, b). This indicates

that *PMR4* contributes to shoot Pi accumulation in Pi-depleted plants. It was described

that *PHR1/PHL1* promotes shoot Pi accumulation (Wang et al., 2018) and that *PHO1* Pi

transporter mediates Pi mobilization from roots to shoots through the root xylem vessels

(Stefanovic et al., 2007). We validated that Pi contents were lowered in the shoots but

increased in the roots, in *phr1 phl1* and *pho1* compared with WT under low Pi (Figure

5a). A decrease in shoot Pi contents was also observed in *phr1 phl1* and *pho1* under

normal Pi (Figure 5b). Since root hairs contribute to Pi accumulation under Pi

deficiency (Tanaka et al., 2014; Robertson-Albertyn et al., 2017; Holz et al., 2018), we

next determined Pi contents in a root hair less mutant, *rhd6*. As seen in *pmr4, phr1 phl1*

and *pho1*, Pi contents were lowered in the shoots but not in the roots of *rhd6* compared

to WT, specifically under Pi deficiency (Figure 6a, b). As illustrated in Figure 5c and 6c,

all these mutants displayed a decrease in shoot Pi contents without reducing root Pi

contents under Pi deficiency. We inferred from these results that *PMR4*-mediated

callose deposition in root hairs contributes to shoot Pi accumulation, possibly through

increasing the uptake or root-to-shoot translocation of Pi. To test the hypothesis, we

determined the rates of Pi uptake and Pi translocation by applying and tracing

radio-labelled P. The rate of ^{32}P uptake per time and the ratio of ^{32}P in the shoot to

whole seedling, standing for Pi translocation rate, were both lowered in *pmr4* and *pmr4*

sid2 compared to WT and *sid2,* respectively, under low Pi (Figure 5d, 6d). In the

presence or absence of *SID2*, *PMR4* contribution to the two proxies was significant only

under low Pi (Figure 5d, 6d). These results suggest that PMR4 promotes Pi uptake and

root-to-shoot Pi translocation under Pi deficiency. Moreover, the ^{32}P uptake rate was

reduced in *sid2* compared to WT, whereas *pmr4 sid2* was not distinguishable from *pmr4*

under low Pi (Figure 6d), suggesting that SID2 positively influences Pi uptake under Pi

deficiency in the presence of PMR4.

We further assessed the possible *PMR4* contributions to the acquisition and tissue

mobilization of Pi and other elements, by inductively coupled plasma mass

spectrometry (ICP-MS) analyses for P, K, Mg, Ca, Mn, Fe, Zn, Mo and B contents in

Pi-depleted and Pi-replete plants. Under Pi sufficiency, although the contents of these

elements were not distinguishable between WT and *pmr4* roots, the contents of K and

the rest were increased and lowered, respectively, in *pmr4* compared to WT shoots

(Figure S9). Under low Pi, Mn, Zn and Mo contents were greater in *pmr4* compared

with WT roots, whereas the contents of all the tested elements except K were lowered in

pmr4 compared with WT shoots (Figure 7). The results were consistent with

PMR4-mediated shoot Pi accumulation without affecting root Pi contents under low Pi

conditions (Figures 4a). Additionally, the results suggest that *PMR4* also positively

influences shoot accumulation of Mg, Ca, Mn, Fe, Zn, Mo and B (Figures 7, S9).

WT-like and increased K accumulation in *pmr4* shoots compared to WT under Pi

deficiency and sufficiency, respectively, show that the root-to-shoot transport is not

entirely reduced for all elements in *pmr4*, pointing to a certain degree of element

specificity in *PMR4* dependence (Figures 7, S9).

The role of *PMR4* in the shoot accumulation of different elements prompted us to

examine whether *PMR4* influences plant responses to different nutrient deficiency. We

examined *pmr4* phenotypes under B scarcity, as B scarcity induces inhibition of the

primary root and transcriptional reprogramming related to ROS and defense in roots

(Miwa et al., 2013; Das & Purkait, 2020). Although the primary root length was

reduced in *pmr4* compared to WT over the tested range of B conditions, its ratio under

B sufficiency (100 µM) versus the examined low B conditions was not affected in *pmr4*

(Figure S10). The results point to specific contribution of *PMR4* to plant growth under

Pi deficiency.

Root *PMR4* contributes to shoot growth under Pi deficiency

We next examined the possible significance of the root *PMR4* pool in promoting

shoot growth, through micrografting between WT and *pmr4* plants. We validated that

the micro-grafted plants of WT/WT (shoot/root genotypes) and *pmr4/pmr4* showed the

corresponding WT and *pmr4* phenotypes in root growth and SFW (Figure 8a-c). The

root hair callose was detected only when *PMR4* was present in the roots, although the

callose signals were not as strong as in intact plants, possibly reflecting the stresses

imposed by the grafting procedure (Figure 8d). Remarkably, SFW was greatly reduced

in WT*/pmr4* whereas it was less affected in *pmr4/*WT (Figure 8b), pointing to a critical

role for a root *PMR4* pool in shoot growth under Pi deficiency. However, the root

growth was partially restored in WT/*pmr4* (Figure 8c), pointing to a role for a shoot

PMR4 pool in sustaining the primary root growth in Pi-depleted plants.

To further examine the role of a PMR4 pool in root hairs, we generated transgenic

plants expressing *PMR4* under the control of a root hair-specific *LRX1* promoter

(*pLRX1*, Baumberger et al., 2003) in the *pmr4-1* background (Figure 9a, b).

pLRX1::PMR4 complemented *pmr4* in terms of the callose deposition in root hairs and

shoot growth under low Pi conditions (Figure 9a-c). By contrast, the primary root

growth remained suppressed in the *pLRX1::PMR4* transgenic lines as well as in *pmr4*

(Figure 9b, d). The results indicate that PMR4 in root hairs is sufficient to restore shoot

growth, but not the primary root growth, under Pi deficiency.

Discussion

Our studies on callose deposition in root hairs induced under Pi deficiency revealed

PMR4 as a positive regulator of plant adaptation to Pi deficiency (Figure 10). *PMR4*

mediates defense-induced callose deposition in pattern triggered immunity and

pre-invasion fungal resistance (Jacobs et al., 2003; Chowdhury et al., 2014), and that its

loss is linked to enhanced SA-based pathogen resistance (Nishimura et al., 2003).

Therefore, *PMR4* seems to play a critical role in an array of biotic and abiotic stress

responses, possibly through callose deposition at different sites.

The present study suggests that the root hair callose contributes to root Pi uptake and

shoot Pi accumulation under Pi deficiency. Importantly, the loss of *PMR4* affects both

PHR1-dependent and *LPR1*-dependent processes (Figures 3, 4) (Bates & Lynch 1996;

Thibaud et al., 2010), which have been genetically uncoupled in PSR (Wang et al.,

2010; Balzergue et al., 2017). Pi accumulation and anthocyanin accumulation are

suppressed in *phr1 phl1* (Rubio et al., 2001; Bustos et al., 2010; Wang et al., 2018).

Shoot Pi contents and plant biomass are also reduced in *pmr4*, implying that *PMR4*

positively influences *PHR1/PHL1*-mediated Pi acquisition and plant growth under

internal Pi deficiency. However, over-accumulation of anthocyanin in *pmr4* (Figure 3k,

l) in contrast to *phr1 phl1* (Rubio et al., 2001; Bustos et al., 2010; Wang et al., 2018),

suggests that *PMR4* contribution is specific to the former two outputs of *PHR1/PHL1*

and that its loss results in exaggeration of Pi starvation symptom. The loss of *PMR4*

also strengthens inhibition of the primary root growth under Pi deficiency, which is

pronounced in the presence of *LPR1/LPR2* (Figure S6), consistent with *LPR1/LPR2-*

-dependent *PMR4* expression (Figure S7). The LPR1 pathway mediates callose

deposition at the root SCN to inhibit the cell-to-cell movement of the transcription

factor SHR, thereby suppressing the cell division and subsequent root elongation

(Müller et al., 2015; Mora-Macías et al., 2017). Our results suggest a role for PMR4 in

antagonizing LPR1/LPR2-dependent root growth inhibition.

The failure of a root hair PMR4 pool to restore primary root growth (Figure 9b, d)

implies the involvement of PMR4 from another cell type(s) than root hairs in this

regulation. Consistently, shoot Pi accumulation is partially and essentially retained in

rhd6 under low and normal Pi conditions, respectively, despite the lack of root hairs

(Figure 6a-c). It is conceivable that PMR4-mediated callose deposition in different cell

types serves to sustain the primary root growth against LPR1/LPR2-mediated

attenuation under Pi deficiency. In line with this, we observed low Pi-induced,

PMR4-mediated callose deposition in cotyledons as well under darkness (Figure S3).

The mechanisms underlying antagonistic interactions between *PMR4* and *LPR1/LPR2*

require further studies including the full identification of critical callose deposition sites.

Shoot Pi accumulation depends on root Pi uptake and subsequent Pi translocation.

PHT1;1 and *PHT1;4* provide two major Pi transporters in roots, contributing greatly to

Pi acquisition under Pi deficiency (Shin et al., 2004). *PHO1* mediates Pi translocation

from roots to shoots through Pi loading into the root xylem (Poirier et al., 1991;

Hamburger et al., 2002; Liu et al., 2012). Notably, the loss of *PMR4* results in a

decrease in the shoot Pi contents without affecting the root Pi contents (Figures 4a-b, 7).

Likewise, the loss of *PHR1/PHL1* and *PHO1* also lowers shoot Pi contents without

reducing root Pi contents (Figure 5a, b; Liu et al., 2012; Wang et al., 2018). In *pht1;1*

pht1;4, a decrease of Pi uptake in roots is associated with that of Pi translocation to the

shoots and shoot Pi accumulation (Shin et al., 2004). An intimate linkage between root

uptake and root-to-shoot translocation is also seen for K^+ (Nieves-Cordones et al., 2019).

It is thus conceivable that the lowered Pi accumulation in *pmr4* shoots is caused by a

decrease in the rates of the root Pi uptake and the root-to-shoot Pi translocation,

possibly through the compromised root hair function in the absence of *PMR4*. Indeed,

^{32}P tracer analyses indicate that *PMR4* promotes both root Pi uptake and root-to-shoot

Pi translocation, without affecting the root Pi contents (Figures 4, 5d, 6d). This is

consistent with the view that the root-to-shoot Pi translocation is driven by the Pi uptake

in the roots (Shin et al., 2004). Leaf transpiration affects root-to-shoot solute

translocation (Nieves-Cordones et al., 2019), but it might be ignorable under

constitutive high humidity in the closed plate assays used in the present study. Notably,

PMR4 also contributes to the accumulation of different elements except K in shoots

under Pi deficiency (Figures 7, S9). This might largely attribute to secondary effects of

PMR4-dependent root growth and expansion (Figure 3e-h), but the separate regulation

of K (Figures 7, S9) cannot be explained by this model. Notably, the disruption of

multiple *PHT1* members (*PHT1;1/1;2/1;3* silencing in the *phf1 pht1;4* mutant

background, designated mut5) results in a decrease in Na as well as P contents

compared to WT (Ayadi et al., 2015). *PHR1* mediates cross-regulation among Fe, Zn

and Pi homeostasis (Briat et al., 2015). Further studies will be required to unravel the

mechanisms underlying the complex coordination of the acquisition and tissue

distribution among different elements and the observed role of PMR4 in the regulation

(Figures 7, S9).

Root hairs increase the root surface area to promote root anchorage, nutrient

acquisition, and metabolite exudation, thereby facilitating the interactions with the soil

environments (Datta et al., 2011; Poole et al., 2018; Vissenberg et al., 2020; Robe et al.,

2021; Han et al., 2023). The elongation and increased densities of root hairs in *pmr4*

(Figure 3c, d) can be inferred that the root hairs formed in the absence of callose

deposition are less functional in Pi acquisition or Pi allocation to the other tissues,

thereby leading to further formation of root hairs.

How does PMR4-mediated callose deposition contribute to the root hair functions

under low Pi? Our findings are reminiscent of the nutrient transport through the phloem

that requires callose deposition in sieve plates (Barratt et al., 2011, Xie & Hong, 2011).

The loss of callose from sieve plate pores reduces the phloem conductivity and thus the

phloem transport (Barratt et al., 2011). Therefore, it is tempting to speculate that the

root hair callose plays a role in cell wall remodeling to reinforce the root hairs for

effective transport of water and nutrients and for providing a driving force for their

root-to-shoot transport. The precise mechanisms by which callose deposited in the root

hairs promotes Pi acquisition require further investigation.

Root hairs also provide a key interface to soil-borne microbes and are intimately

associated with root defense regulation. Root hair formation and elongation are

facilitated by jasmonate (JA) (Zhu et al., 2006), which mediates the expression of key

transcription factors for root hair differentiation and the de-repression of RHD6 from

the interference by JAZ proteins (Han et al., 2020). In beneficial interactions with the

epiphytic bacterium *Pseudomonas simiae* WCS417, activation of defense-related genes

is increased in hair cell lineages compared to non-hair cell lineages (Verbon et al., 2023).

The endogenous elicitor-active Pep peptides induce root hair formation (Okada et al.,

2021; Jing et al., 2024), and Pep receptor signaling from the root hairs can activate

defense responses without root growth inhibition (Okada et al., 2021). These findings

suggest a specific capacity of root hair cells in root immunity. How these properties of

root hairs are influenced by *PMR4*-mediated callose deposition under Pi deficiency also

represents an interesting topic for future studies.

Experimental procedures

Plant materials and growth conditions

The *Arabidopsis thaliana* accessions used were Col-0 and Ws-2, including the mutants

examined (Table S1) and the wild type used was Col-0 unless otherwise stated. We

generated ethyl methanesulfonate (EMS)-mutagenesis (M2) seeds for screening in both

Col-0 and Ler backgrounds. All seeds were sterilized using 6% sodium hypochlorite

and 0.2% Trion X-100 for 15 min, and then kept at 4°C for 2 d before sowing a

half-strength Murashige and Skoog (MS) medium with 25 mM sucrose and 0.8% Agar

(WAKO, Japan). If not stated otherwise, the normal Pi media contained 625 μM

KH_2PO_4, whereas the low Pi media contained 0 μM, 10 μM, 50 μM or 150 μM KH_2PO_4

as indicated. The low N and low Fe media used contained 0 μM N and 0 μM Fe,

respectively. Plants were grown in a growth chamber under 12 h: 12 h, white light: dark

cycles at 22°C unless otherwise stated. The white light intensity was 80 μmol m^{-2} s^{-1}.

Six-day (d)-old seedlings were transferred to the indicated Pi liquid media or agar media

with 1% Difco Agar, Bacteriological (BD, Franklin Lakes, NJ, USA).

Construction and plant transformation

The genomic DNA sequences of the *POWDERY MILDEW RESISTANCE 4* (*PMR4*,

AT4G03550) loci, together with a regulatory DNA sequence for root hair specific

expression (below), were cloned into the Gateway binary vector pGWB504 (Nakagawa

et al., 2007; primer information Table S2) to express a C-terminally GFP-fused PMR4

under its native regulatory DNA sequence. The root hair-specific promoter sequence

used was derived from *LRR-EXTENSIN 1* (*LRX1*, AT1G12040) (Baumberger et al.,

2003). Transgenic lines were generated by Agrobacterium-mediated transformation of

pmr4-1 with the DNA construct and subsequently selected on hygromycin resistance

and DNA genotyping.

Aniline blue and PI staining

Callose staining with aniline blue was conducted as described in Schenk & Schikora

(2015). Plants were examined at 3 d after the transfer of 5-d-old seedlings to the

indicated Pi, N or Fe liquid media (Figure 1a, c). Plant tissues were fixed in a solution

containing acetic acid and ethanol (1:3 v/v) overnight. The fixation solution was

exchanged to a 150 mM K_2HPO_4 solution before callose staining with 0.1% aniline blue

in the K_2HPO_4 solution for 2 h. The data presented are representative of three

independent experiments (Figures 1a, 1c, 2a, 3a), two independent experiments (Figures

1b, 6d, S1, S2) and one experiment (Figures 7a, S3). To localize the root meristematic

regions, the primary roots were stained with 20 µg/ml propidium iodide (PI). Callose or

PI signals were examined under microscopy using a fluorescence microscope DMI

6000B (Leica Microsystems, Wetzlar, Germany) and a confocal microscope SP8 (Leica

Microsystems, Wetzlar, Germany).

Quantitative real-time PCR analysis

Three-day-old seedlings were exposed to low Pi or normal Pi media for 4 d. Total RNA

was then extracted from the root tissues by NucleoSpin® RNA Plant kit

(Machete-Nagel, Dueren, Germany). Total RNA was reverse-transcribed using the

PrimeScript RT reagent kit including a genomic DNA eraser (Perfect Real Time, Takara

Bio, Ohtsu, Japan). Quantitative PCR was performed with SYBR Green (Life

Technologies, Carlsbad, CA, USA) and the Thermal Cycler Dice® Real-Time System

(TakaraBio). Expression levels of the examined genes (Table S2) were normalized

relative to a reference gene, *ACTIN 2* (*ACT2*). PSR-inducible markers examined were

AT4 (At5g03545), *IPS1* (AT3g09922), *PHT1;5* (AT2g32830) and *SPX1* (AT5g20150).

Data are representative of two independent experiments

Identification of mutants altered in the root hair callose deposition

We examined callose deposition in *gsl* mutants (Table S1) after aniline blue staining of

the roots transferred to the low Pi liquid media (50 μM) for 3 d. In the EMS mutagenic

screening, we subjected a region ~5 cm from the root tip to aniline blue staining, after

cutting out of the M2 individuals transferred to the Pi -depleted liquid media (0 μM) for

3 d. This enabled to keep the plant alive for subsequent characterization and seed

propagation. The *PMR4* genomic region was sequenced in the selected mutant

candidates.

Inoculation analyses with powdery mildew fungi

Inoculation of *Golovinomyces orontii* and determination of fungal versus plant DNA

ratio were conducted essentially by following Weßling & Panstruga (2012).

Eighteen-d-old plants grown in 10 h: 14 h, light: dark cycles at 22°C were inoculated

with *G. orontii* spores at approximately 750 spores cm^{-2}. The inoculated leaves were

harvested and frozen in liquid nitrogen before determining fungal biomass at 10 d after

inoculation. Genomic DNA was extracted essentially by following Brouwer et al.

(2001). Quantitative real-time PCR was performed with primers (Table S2) and

Brilliant Sybr Green QPCR Reagent Kit (Stratagene, Waldbronn, Germany). The ratio

of *G. orontii* to Arabidopsis genomic DNA was calculated using the $\Delta\Delta$Ct method

(Pfaffl, 2001). Data are representative of two independent experiments (Figure 2c).

Determination of root hair density, primary root length and root area

Six-day-old seedlings were transferred and further grown in the low (50 µM) or normal

Pi media (625 µM) for 6 d, and then root hairs, primary root and root system area were

examined under stereomicroscope M80 (Leica Microsystems, Wetzlar, Germany). The

number of root hairs was determined in a region 2.0-3.0 mm from the root tip using

Image J software (National Institutes of Health, Bethesda, MD, USA). Data are

representative of similar results from three independent experiments (Figure 3c, d) and

two independent experiments (Figure S4). The primary root length and convex hull area

were determined using ImageJ software and Root nav software (Pound et al., 2013).

Data are representative of three independent experiments (Figure. 3e-h). For growth

under different Boron (B) conditions, plants were grown in MGRL solid media

containing 1% sucrose and 1% gellan gum (WAKO, Japan). B concentrations were

adjusted with boric acid. Plants were grown under 16 h: 8 h, light: dark cycles at 22°C.

The data presented are from one independent experiment (Figure S10).

Determination of shoot fresh weight, anthocyanin and Pi contents

In plant growth assays on media, six-day-old seedlings were transferred to the indicated

media (10 µM, 50 µM or 625 µM Pi) and then further grown for the indicated days.

Shoot fresh weight (SFW) was determined 16 d after transfer to the indicated media (50

μM or 625 μM Pi; Figure 3i, j). In plant growth assays with *pLRX1::PMR4* lines, SFW

was determined 10 d after the transfer to the low Pi media (10 μM ： Figure 9a, c). In

plant growth assay on soil, organic black soil (Maruishi Engei, Osaka, Japan) and Ezo

sand (Kobayashi Sangyo, Kita-Nagoya, Japan) mixed at 1:1 (v/v) were used as low Pi

soil, and Sakata-Super-Mix-A (Sakata Seed, Kyoto, Japan) and Vermiculite G20

(Maruishi Engei, Osaka, Japan) mixed at 1:1 (v/v) were used as normal Pi soil (Figure

S8). SFW of seedlings grown on soil was determined 14 d after the transfer of 10-d-old

seedlings to the low and normal Pi soils. Data are representative of two independent

experiments (Figures 3i, 3j, S5) and one independent experiment (Figure 9b, c).

In the determination of anthocyanin content, six-day-old seedlings were transferred to

and further grown on low Pi media (10 μM) for 10 d (Figure 3k, l). The shoots were

subject to the procedure described in Mita et al. (1997). In brief, a HCl: methanol

mixture at 1:99 (v/v) was added to the ground tissues with 1 ml per sample, and then the

resultant homogenates were incubated at 4°C overnight. A_{530} and A_{657} of the

supernatants were determined with a spectrometer UV-1800 (Shimadzu, Kyoto, Japan).

Anthocyanin content was calculated following the formula: $(A_{530} - 0.25 \times A_{657}) \times$ fresh

weight $(FW)^{-1}$. Data are representative of two independent experiments with the same

conclusions (Figure 3k, l).

In the determination of Pi content, six-d-old seedlings were transferred to and further

grown on the low Pi (150 μM) or normal Pi (625 μM) media for 9 d (Figures 4, 5a-b,

6a-b). Pi content was determined by following Ames (1966). In brief, the extraction

buffer (10 mM Tris, 1 mM EDTA, 100 mM NaCl, 1 mM phenylmethylsulfonyl fluoride

pH8.0) was added with 10 μl per sample mg FW. The homogenate 100 μl was mixed

with 1% glacial acetic acid 900 μl, and then incubated at 42°C for 30 min. The

supernatant was recovered to be incubated in a solution (0.35% NH_4MoO_4, 0.86 N

H_2SO_4, and 1.4% ascorbic acid) at 42°C for 30 min. A_{820} of the aliquot was measured

under Eppendorf BioSpectrometer® basic (Eppendorf, Hamburg, Germany). Then, Pi

content was calculated by a P calibration curve prepared with a P standard solution.

Data are representative of three independent experiments (Figure 4a, b) and two

independent experiments (Figures 5d, 6d) with the same conclusions.

Scanning electron microscope

The maturation zone of primary roots 10 d after transfer of 5-d-old seedlings to low Pi

(50 µM) liquid media was fixed in half-Karnovsky's fixative overnight at 4°C, and then

postfixed in 1% osmium tetroxide for 1 h at room temperatures (Karnovsky, 1965). The

specimens were dehydrated in an ethanol series (30%, 50%, 70%, 80%, and 90%, and

100% at three times) for 10 min each at RT, and then the specimens were dipped in

t-butyl alcohol and dried with a vacuum evaporator. They were coated with vapor of 1%

osmium tetroxide and then observed with a scanning electron microscope (JIB-4601F,

JEOL, Tokyo, Japan).

Inductively coupled plasma-mass spectrometer (ICP-MS) analyses

Five-day-old seedlings were transferred to and further grown on the growth media (10

µM or 625 µM KH_2PO_4) for 4 d (Figures 7, S9). Shoots and roots were harvested and

dried at 60°C. After determining the dry weight, the dried tissues were digested with 2

ml nitric acid and 0.5 ml hydrogen peroxide (Wako Pure Chemicals) at 110°C. Digested

samples were then dissociated with 5 ml of 2% (w/v) nitric acid and subjected to

inductively coupled plasma mass spectrometry (NexION 2000, Perkin-Elmer). The

contents of P, K, Mg, Ca, Mn, Fe, Zn, Mo and B were determined. The data were

obtained from one experiment (Figures 7, S9).

Determination of ^{32}P uptake rate and translocation ratio

Seven-day-old seedlings were transferred to and further grown on the growth media (50

or 625 μM KH$_2$PO$_4$) for 10 d. The seedlings were exposed to the liquid media

containing ^{32}P-labeled Pi (^{32}PO$_4$$^{3-}$ 20 kBq / ml) for 30 min. The plant tissues were fixed

to an Imaging Plate (GE Healthcare co., Chicago, USA) for 3 h and then scanned by

Typhoon FLA 9000 (GE Healthcare co., Chicago, USA). ^{32}P uptake rate was calculated

based on a calibration curve for ^{32}P. For determining the Pi translocation rate into shoots,

the seedlings were exposed to the Pi-containing liquid media containing ^{32}P-labeled Pi

for 10 min, and then transferred to non-labelled agar media for 120 min. The

translocation rate of Pi was determined by calculating ^{32}P ratio of the shoot to the whole

seedlings. The data presented are representative of two independent experiments with

the same conclusions (Figures 5d, 6d).

Grafting experiments

Grafting of WT and *pmr4* was performed with a supportive micro-device as described

in Tsutsui et al. (2020). Four-day-old seedlings grown on a 1/2 MS medium containing

0.5% sucrose and 1% agar under constant light at 22°C were subject to grafting. Both

hypocotyls of stock and scion were cut horizontally with a knife and assembled on the

device using forceps. The grafted plants were grown in a 1/2 MS medium containing

0.5% sucrose and 2% agar at 27°C for 6 d. In callose staining, the grafted plants

transferred and grown on low Pi media (50 μM) under constant light at 22°C for 9 d

were examined. In plant growth assays, the grafted plants transferred and grown on low

Pi media (150 µM) under constant light at 22°C for 14 d were examined. The data

presented are representative of two independent experiments with the same conclusions

(Figure 8).

Acknowledgements

We thank the Arabidopsis Biological Research Centre for published plant materials,

Yaichi Kawakatsu for advice on micrografting, and Michiko Tsukamoto and Mie

Matsubara for technical assistance. This work was supported in part by the grants from

Mitsubishi Foundation (YS), the Japan Science and Technology Agency (JST,

JPMJPR13B6 and JPMJSC1702 (YS) and JPMJPR16Q7 (KH)), and MEXT of Japan

18H02467 and 21H02507 (YS).

Author contributions

YS conceived the study. KO and YS designed the experiments. KO, KY, TANN, SK, SY,

HT, CT, THL, UN, KI, NT, TI, KM, TM and MN developed the methods, performed the

experiments, and/or analyzed the data. SM, MN and KH advised on the experiments. KO

and YS wrote the manuscript with contributions from the other authors.

Supporting Information

Figure S1. Callose deposition in *gsl* mutants under phosphate deficiency.

Figure S2. flg22-induced callose deposition is impaired in the *caps* and *pmr4* alleles.

Figure S3. Pi deficiency-induced callose deposition under dark conditions.

Figure S4. Root hair and meristem morphology in *pmr4* under Pi deficiency.

Figure S5. PSR marker gene expression in *pmr4* roots

Figure S6. *PMR4* restricts *LPR1/LPR2*-mediated inhibition of the primary root

growth under low Pi.

Figure S7. *PMR4* expression in *phr1 phl1* and *lpr1 lpr2* roots.

Figure S8. Plant growth is lowered in *pmr4* on soil.

Figure S9. Plant element contents in shoot and root under phosphate sufficiency.

Figure S10. Primary root growth inhibition under B deficiency.

Table S1. *Arabidopsis thaliana* mutants used in this study.

Table S2. DNA Primers used in DNA vector construction and qPCR analyses.

References

Abel, S. (2017). Phosphate scouting by root tips. *Current Opinion in Plant Biology,* **39**,

635   168–177.

636 **Ames, B.N.** (1966) Assay of inorganic phosphate, total phosphate and phosphatases.

637   *Methods Enzymol*, **8**, 115–118.

638 **Ayadi, A., David, P., Arrighi, J.F., Chiarenza, S., Thibaud, M.C., Nussaume, L.**

639   **and Marin, E.** (2015) Reducing the genetic redundancy of Arabidopsis phosphate

640   transporter1 transporters to study phosphate uptake and signaling. *Plant Physiol*,

641   **167**, 1511–1526.

642 **Balzergue, C., Dartevelle, T., Godon, C., et al.** (2017) Low phosphate activates

643   STOP1-ALMT1 to rapidly inhibit root cell elongation. *Nat Commun*, **8**.

644 **Bates, T.R. and Lynch, J.P.** (1996) Stimulation of root hair elongation in *Arabidopsis*

645   *thaliana* by low phosphorus availability. *Plant Cell Environ*, **19**, 529–538.

Baumberger, N., Ringli, C. and Keller, B. (2001) The chimeric leucine-rich

repeat/extensin cell wall protein LRX1 is required for root hair morphogenesis in

Arabidopsis thaliana. *Genes Dev*, **15**, 1128–1139.

Briat, J.F., Rouached, H., Tissot, N., Gaymard, F. and Dubos, C. (2015) Integration

of P, S, Fe, and Zn nutrition signals in *Arabidopsis thaliana*: Potential involvement

of PHOSPHATE STARVATION RESPONSE 1 (PHR1). *Front Plant Sci*, **6**.

Brouwer, M., Lievens, B., Hemelrijck, W. Van, Ackerveken, G. Van Den, Cammue,

B.P.A. and Thomma, B.P.H.J. (2003) Quantification of disease progression of

several microbial pathogens on *Arabidopsis thaliana* using real-time fluorescence

PCR. *FEMS Microbiol Lett*, **228**, 241–248.

Bustos, R., Castrillo, G., Linhares, F., Puga, M.I., Rubio, V., Pérez-Pérez, J.,

Solano, R., Leyva, A. and Paz-Ares, J. (2010) A central regulatory system

largely controls transcriptional activation and repression responses to phosphate

starvation in Arabidopsis. *PLoS Genet*, **6**.

Chan, C., Liao, Y.Y. and Chiou, T.J. (2021) The Impact of Phosphorus on Plant

Immunity. *Plant Cell Physiol*, **62**, 582–589.

Chien, P.S., Chiang, C.P., Leong, S.J. and Chiou, T.J. (2018) Sensing and Signaling

of Phosphate Starvation: From Local to Long Distance. *Plant Cell Physiol*, **59**,

1714–1722.

Chowdhury, J., Henderson, M., Schweizer, P., Burton, R.A., Fincher, G.B. and

Little, A. (2014) Differential accumulation of callose, arabinoxylan and cellulose

in nonpenetrated versus penetrated papillae on leaves of barley infected with

Blumeria graminis f. sp. *hordei*. *New Phytologist*, **204**, 650–660.

Das, A.K. and Purkait, A. (2020) Boron dynamics in soil: classification, sources,

factors, fractions, and kinetics. *Commun Soil Sci Plant Anal*, **51**, 2778–2790.

Datta, S., Kim, C.M., Pernas, M., Pires, N.D., Proust, H., Tam, T., Vijayakumar, P.

and Dolan, L. (2011) Root hairs: development, growth and evolution at the

plant-soil interface. *Plant Soil*, **346**, 1–14.

Dolan, L., Duckett, C.M., Grierson, C., Linstead, P., Schneider, K., Lawson, E.,

Dean, C., Poethig, S. and Roberts, K. (1994) Clonal relationships and cell

patterning in the root epidermis of *Arabidopsis*. *Development*, **120**, 2465–2474.

Dong, X., Hong, Z., Sivaramakrishnan, M., Mahfouz, M. and Verma, D.P.S. (2005)

Callose synthase (CalS5) is required for exine formation during

microgametogenesis and for pollen viability in Arabidopsis. *Plant Journal*, **42**,

315–328.

Dong, X., Hong, Z., Chatterjee, J., Kim, S., & Verma, D. P. S. (2008). Expression of

callose synthase genes and its connection with Npr1 signaling pathway during

pathogen infection. *Planta*, **229**, 87–98.

https://doi.org/10.1007/s00425-008-0812-3.

Ellinger, D. and Voigt, C.A. (2014) Callose biosynthesis in arabidopsis with a focus on

pathogen response: What we have learned within the last decade. *Ann Bot*, **114**,

1349–1358.

Enns, L.C., Kanaoka, M.M., Torii, K.U., Comai, L., Okada, K. and Cleland, R.E.

(2005) Two callose synthases, GSL1 and GSL5, play an essential and redundant

role in plant and pollen development and in fertility. *Plant Mol Biol*, **58**, 333–349.

Franco-Zorrilla, J. M., Valli, A., Todesco, M., Mateos, I., Puga, M. I.,

Rubio-Somoza, I., Leyva, A., Weigel, D., García, J. A., & Paz-Ares, J. (2007).

Target mimicry provides a new mechanism for regulation of microRNA activity.

Nature Genetics, **39**, 1033–1037.

Gao, Y. Q., Bu, L. H., Han, M. L., Wang, Y. L., Li, Z. Y., Liu, H. T., & Chao, D. Y.

(2021). Long-distance blue light signalling regulates phosphate deficiency-induced

primary root growth inhibition. *Molecular Plant*, **14**, 1539–1553.

García-Andrade, J., Ramírez, V., Flors, V., & Vera, P. (2011). Arabidopsis *ocp3*

mutant reveals a mechanism linking ABA and JA to pathogen-induced callose

deposition. *Plant Journal*, **67**, 783–794.

Gulabani, H., Goswami, K., Walia, Y., et al. (2021) Arabidopsis inositol

polyphosphate kinases IPK1 and ITPK1 modulate crosstalk between SA-dependent

immunity and phosphate-starvation responses. *Plant Cell Rep*, **41**, 347–363.

Hamburger, D., Rezzonico, E., Petétot, J.M.D.C., Somerville, C. and Poirier, Y.

(2002) Identification and characterization of the Arabidopsis PHO1 gene involved

in phosphate loading to the xylem. *Plant Cell*, **14**, 889–902.

Han, G., Wei, X., Dong, X., et al. (2020) Arabidopsis ZINC FINGER PROTEIN1 Acts

Downstream of GL2 to Repress Root Hair Initiation and Elongation by Directly

Suppressing bHLH Genes. *Plant Cell*, **32**, 206–225.

Han, X., Kui, M., He, K., Yang, M., Du, J., Jiang, Y. and Hu, Y. (2023)

Jasmonate-regulated root growth inhibition and root hair elongation. *J Exp Bot*, **74**,

1176–1185.

Holz, M., Zarebanadkouki, M., Kuzyakov, Y., Pausch, J. and Carminati, A. (2018)

Root hairs increase rhizosphere extension and carbon input to soil. *Ann Bot*, **121**,

61–69.

Isidra-Arellano, M.C., Delaux, P.M. and Vald°s-L°pez, O. (2021) The Phosphate

Starvation Response System: Its Role in the Regulation of Plant–Microbe

Interactions. *Plant Cell Physiol*, **62**, 392–400.

Jacobs, A.K., Lipka, V., Burton, R.A., Panstruga, R., Strizhov, N., Schulze-Lefert,

P. and Fincher, G.B. (2003) An Arabidopsis Callose Synthase, GSL5, Is Required

for Wound and Papillary Callose Formation. *Plant Cell*, **15**, 2503–2513.

Jing, Y., Zhao, F., Lai, K., et al. (2024) Plant elicitor Peptides regulate root hair

development in *Arabidopsis*. *Front Plant Sci*, **15**.

Karnovsky, M.J. (1964) A Formaldehyde-Glutaraldehyde Fixative of High Osmolality

for Use in Electron Microscopy. *The Journal of Cell Biology*, **27**, 1A-149A.

https://www.researchgate.net/publication/244955881.

Keppler, B. D., Song, J., Nyman, J., Voigt, C. A., & Bent, A. F. (2018).

3-aminobenzamide blocks MAMP-induced callose deposition independently of its

poly(ADPribosyl)ation inhibiting activity. *Frontiers in Plant Science*, **871**.

https://doi.org/10.3389/fpls.2018.01907.

Liu, T.Y., Huang, T.K., Tseng, C.Y., Lai, Y.S., Lin, S.I., Lin, W.Y., Chen, J.W. and

Chioua, T.J. (2012) PHO2-dependent degradation of PHO1 modulates phosphate

homeostasis in *Arabidopsis*. *Plant Cell*, **24**, 2168–2183.

Mateo, A., Funck, D., Mühlenbock, P., Kular, B., Mullineaux, P.M. and Karpinski,

S. (2006) Controlled levels of salicylic acid are required for optimal photosynthesis

and redox homeostasis. *J Exp Bot*, **57**, 1795–1807.

Mita, S., Murano N., Akaike M. and Nakamura K. (1997) Mutants of *Arabidopsis*

thaliana with pleiotropic effects on the expression of the gene for β-amylase and

on the accumulation of anthocyanin that are inducible by sugars. *Plant Journal*, **11**,

841-851.

Miura, K., Lee, J., Miura, T. and Hasegawa, P.M. (2010) SIZ1 controls cell growth

and plant development in Arabidopsis through salicylic acid. *Plant Cell Physiol*, **51**,

103–113.

Miwa, K., Wakuta, S., Takada, S., Ide, K., Takano, J., Naito, S., Omori, H.,

Matsunaga, T. and Fujiwara, T. (2013) Roles of BOR2, a boron exporter, in

cross linking of rhamnogalacturonan II and root elongation under boron limitation

in Arabidopsis. *Plant Physiol*, **163**, 1699–1709.

Mora-Macías, J., Ojeda-Rivera, J.O., Gutiérrez-Alanís, D., et al. (2017)

Malate-dependent Fe accumulation is a critical checkpoint in the root

developmental response to low phosphate. *Proc Natl Acad Sci USA*, **114**,

E3563–E3572. https://doi.org/10.1073/pnas.1701952114

Morcillo, R.J., Singh, S.K., He, D., et al. (2020) Rhizobacterium⎕derived diacetyl

modulates plant immunity in a phosphate⎕dependent manner. *EMBO J*, **39**, 1–15.

Müller, J., Toev, T., Heisters, M., Teller, J., Moore, K.L., Hause, G., Dinesh, D.C.,

Bürstenbinder, K. and Abel, S. (2015) Iron-Dependent Callose Deposition

Adjusts Root Meristem Maintenance to Phosphate Availability. *Dev Cell*, **33**,

216–230.

Nagarajan, V. K., Jain, A., Poling, M. D., Lewis, A. J., Raghothama, K. G., &

Smith, A. P. (2011). Arabidopsis Pht1;5 Mobilizes Phosphate between Source and

Sink Organs and Influences the Interaction between Phosphate Homeostasis and

Ethylene Signaling. *Plant Physiology*, **156**, 1149–1163.

Nakagawa, T., Suzuki, T., Murata, S., et al. (2007) Improved gateway binary vectors:

High-performance vectors for creation of fusion constructs in transgenic analysis

of plants. *Biosci Biotechnol Biochem*, **71**, 2095–2100.

Nieves-Cordones, M., Lara, A., Ródenas, R., Amo, J., Rivero, R.M., Martínez, V.

and Rubio, F. (2019) Modulation of K^+ translocation by AKT1 and AtHAK5 in

Arabidopsis plants. *Plant Cell Environ*, **42**, 2357–2371.

Nishikawa, S.I., Zinkl, G.M., Swanson, R.J., Maruyama, D. and Preuss, D. (2005)

Callose (β-1,3 glucan) is essential for Arabidopsis pollen wall patterning, but not

tube growth. *BMC Plant Biol*, **5**, 1–9.

Nishimura, M.T., Stein, M., Hou, B.H., Vogel, J.P., Edwards, H. and Somerville,

S.C. (2003) Loss of a callose synthase results in salicylic acid-dependent disease

resistance. *Science (1979)*, **301**, 969–972.

Okada, K., Kubota, Y., Hirase, T., Otani, K., Goh, T., Hiruma, K. and Saijo, Y.

(2021) Uncoupling root hair formation and defence activation from growth

inhibition in response to damage-associated Pep peptides in *Arabidopsis thaliana*.

New Phytologist, **229**, 2844–2858.

O'Lexy, R., Kasai, K., Clark, N., Fujiwara, T., Sozzani, R. and Gallagher, K.L.

(2018) Exposure to heavy metal stress triggers changes in plasmodesmatal

permeability via deposition and breakdown of callose. *J Exp Bot*, **69**, 3715–3728.

Omasits, U., Ahrens, C.H., Müller, S. and Wollscheid, B. (2014) Protter: Interactive

protein feature visualization and integration with experimental proteomic data.

Bioinformatics, **30**, 884–886.

Pasternak, T., Groot, E.P., Kazantsev, F. V., Teale, W., Omelyanchuk, N.,

Kovrizhnykh, V., Palme, K. and Mironova, V. V. (2019) Salicylic acid affects

root meristem patterning via auxin distribution in a concentration-dependent

manner. *Plant Physiol*, **180**, 1725–1739.

Paul Barratt, D. H., Kölling, K., Graf, A., Pike, M., Calder, G., Findlay, K.,

Zeeman, S. C., & Smith, A. M. (2011). Callose synthase GSL7 is necessary for

normal phloem transport and inflorescence growth in Arabidopsis. *Plant*

Physiology, **155**, 328–341.

Péret, B., Clément, M., Nussaume, L. and Desnos, T. (2011) Root developmental

adaptation to phosphate starvation: Better safe than sorry. *Trends Plant Sci*, **16**,

442–450.

Péret, B., Desnos, T., Jost, R., Kanno, S., Berkowitz, O. and Nussaume, L. (2014)

Root architecture responses: In search of phosphate. *Plant Physiol*, **166**,

1713–1723.

Pfaffl, M.W. (2001) A new mathematical model for relative quantification in real-time

RT-PCR. *Nucleic Acids Research*, **29**, 2003-2007

Poirier, Y., Thoma, S., Somerville, C. and Schiefelbein, J. (1991) A mutant of

Arabidopsis deficient in xylem loading of phosphate. *Plant Physiol*, **97**,

1087–1093.

Poole, J., Day, C.J., Itzstein, M. Von, Paton, J.C. and Jennings, M.P. (2018)

Glycointeractions in bacterial pathogenesis. *Nat Rev Microbiol*, **16**, 440–452.

http://dx.doi.org/10.1038/s41579-018-0007-2.

Pound, M.P., French, A.P., Atkinson, J.A., Wells, D.M., Bennett, M.J. and

Pridmore, T. (2013) RootNav: Navigating images of complex root architectures.

Plant Physiol, **162**, 1802–1814.

Puga, M. I., Mateos, I., Charukesi, R., Wang, Z., Franco-Zorrilla, J. M., De

Lorenzo, L., Irigoyen, M. L., Masiero, S., Bustos, R., Rodríguez, J., Leyva, A.,

Rubio, V., Sommer, H., & Paz-Ares, J. (2014). SPX1 is a phosphate-dependent

inhibitor of Phosphate Starvation Response 1 in Arabidopsis. *Proceedings of the*

National Academy of Sciences of the United States of America, **111**, 14947–14952.

Robe, K., Conejero, G., Gao, F., et al. (2021) Coumarin accumulation and trafficking

in *Arabidopsis thaliana*: a complex and dynamic process. *New Phytologist*, **229**,

2062–2079.

Robertson-Albertyn, S., Terrazas, R.A., Balbirnie, K., et al. (2017) Root hair

mutations displace the barley rhizosphere microbiota. *Front Plant Sci*, **8**, 1–15.

Rubio, V., Linhares, F., Solano, R., Martín, A.C., Iglesias, J., Leyva, A. and

Paz-Ares, J. (2001) A conserved MYB transcription factor involved in phosphate

starvation signaling both in vascular plants and in unicellular algae. *Genes Dev*, **15**,

2122–2133.

Sager, R. and Lee, J.Y. (2014) Plasmodesmata in integrated cell signalling: Insights

from development and environmental signals and stresses. *J Exp Bot*, **65**,

6337–6358.

Schenk, S.T. and Schikora, A. (2015) Staining of Callose Depositions in Root and

Leaf Tissues. *Bio-protocol*, **5**. http://www.bio-protocol.org/e1429

Schiefelbein, J., Kwak, S.H., Wieckowski, Y., Barron, C. and Bruex, A. (2009) The

gene regulatory network for root epidermal cell-type pattern formation in

Arabidopsis. *J Exp Bot*, **60**, 1515–1521.

Shin, H., Shin, H.S., Dewbre, G.R. and Harrison, M.J. (2004) Phosphate transport in

832   *Arabidopsis*: Pht1;1 and Pht1;4 play a major role in phosphate acquisition from

833   both low- and high-phosphate environments. *Plant Journal*, **39**, 629–642.

834   **Stefanovic, A., Ribot, C., Rouached, H., Wang, Y., Chong, J., Belbahri, L.,**

835   **Delessert, S. and Poirier, Y.** (2007) Members of the *PHO1* gene family show

836   limited functional redundancy in phosphate transfer to the shoot, and are regulated

837   by phosphate deficiency via distinct pathways. *Plant Journal*, **50**, 982–994.

838   **Sun, L., Song, L., Zhang, Y., Zheng, Z. and Liu, D.** (2016) Arabidopsis PHL2 and

839   PHR1 act redundantly as the key components of the central regulatory system

840   controlling transcriptional responses to phosphate starvation. *Plant Physiol*, **170**,

841   499–514.

842   **Svistoonoff, S., Creff, A., Reymond, M., Sigoillot-Claude, C., Ricaud, L., Blanchet,**

843   **A., Nussaume, L. and Desnos, T.** (2007) Root tip contact with low-phosphate

844   media reprograms plant root architecture. *Nat Genet*, **39**, 792–796.

845   **Tanaka, N., Kato, M., Tomioka, R., Kurata, R., Fukao, Y., Aoyama, T. and**

846   **Maeshima, M.** (2014) Characteristics of a root hair-less line of *Arabidopsis*

847   *thaliana* under physiological stresses. *J Exp Bot*, **65**, 1497–1512.

848   **Thibaud, M.C., Arrighi, J.F., Bayle, V., Chiarenza, S., Creff, A., Bustos, R.,**

849   **Paz-Ares, J., Poirier, Y. and Nussaume, L.** (2010) Dissection of local and

850   systemic transcriptional responses to phosphate starvation in Arabidopsis. *Plant*

851   *Journal*, **64**, 775–789.

852   **Verbon, E.H., Liberman, L.M., Zhou, J., Yin, J., Pieterse, C.M.J., Benfey, N.,**

853   **Stringlis, I.A. and Jonge, R. De** (2023) Cell-type-specific transcriptomics reveals

854   that root hairs and endodermal barriers play important roles in beneficial

plant-rhizobacterium interactions. *Mol Plant*, **16**, 1160-1177.

https://doi.org/10.1016/j.molp.2023.06.001.

Vissenberg, K., Claeijs, N., Balcerowicz, D. and Schoenaers, S. (2020) Hormonal

regulation of root hair growth and responses to the environment in Arabidopsis. *J*

Exp Bot, **71**, 2412–2427.

Vogel, J. and Somerville, S. (2000) Isolation and characterization of powdery

mildew-resistant *Arabidopsis* mutants. *Proc Natl Acad Sci USA*, **97**, 1897–1902.

https://doi.org/10.1073/pnas.030531997

Wang, Xuemin, Du, G., Wang, Xuming, Meng, Y., Li, Y., Wu, P. and Yi, K. (2010)

The function of LPR1 is controlled by an element in the promoter and is

independent of SUMO E3 ligase SIZ1 in response to Low Pi stress in *Arabidopsis*

thaliana. *Plant Cell Physiol*, **51**, 380–394.

Wang, Y., Li, X., Fan, B., Zhu, C. and Chen, Z. (2021) Regulation and function of

defense-related callose deposition in plants. *Int J Mol Sci*, **22**, 1–15.

Wang, Z., Zheng, Z., Song, L. and Liu, D. (2018) Functional characterization of

Arabidopsis PHL4 in plant response to phosphate starvation. *Front Plant Sci*, **9**,

1–19.

Weßling, R. and Panstruga, R. (2012) Rapid quantification of plant-powdery mildew

interactions by qPCR and conidiospore counts. *Plant Methods*, **8**.

Wildermuth, M.C., Dewdney, J., Wu, G. and Ausubel, F.M. (2002) Isochorismate

synthase is required to synthesize salicylic acid for plant defence. *Nature*, **417**,

571–571.

Wu, S.W., Kumar, R., Iswanto, A.B.B. and Kim, J.Y. (2018) Callose balancing at

plasmodesmata. *J Exp Bot*, **69**, 5325–5339.

Xie, B., & Hong, Z. (2011). Unplugging the callose plug from sieve pores. *Plant*

Signaling and Behavior, **6**, 491–493.

Yeh, C. M., Kobayashi, K., Fujii, S., Fukaki, H., Mitsuda, N., & Ohme-Takagi, M.

(2020). Blue Light Regulates Phosphate Deficiency-Dependent Primary Root

Growth Inhibition in Arabidopsis. *Frontiers in Plant Science*, **10**.

https://doi.org/10.3389/fpls.2019.01803.

Zhang, X., Chen, S. andMou, Z. (2010) Nuclear localization of NPR1 is required for

regulation of salicylate tolerance, isochorismate synthase 1 expression and

salicylate accumulation in *Arabidopsis*. *J Plant Physiol*, **167**, 144–148.

http://dx.doi.org/10.1016/j.jplph.2009.08.002.

Zheng, Z., Wang, Z., Wang, X., & Liu, D. (2019). Blue Light-Triggered Chemical

Reactions Underlie Phosphate Deficiency-Induced Inhibition of Root Elongation

of *Arabidopsis* Seedlings Grown in Petri Dishes. *Molecular Plant*, **12**, 1515–1523.

https://doi.org/10.1016/j.molp.2019.08.001.

Zhu, C., Gan, L., Shen, Z. and Xia, K. (2006) Interactions between jasmonates and

ethylene in the regulation of root hair development in *Arabidopsis*. *J Exp Bot*, **57**,

1299–1308.

Figure Legends

Figure 1. Callose deposition is induced in root hairs under phosphate deficiency,

independently of *PHR1, PHL1, LPR1* and *LPR2*.

Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis*

thaliana seedlings to low Pi (P 50 µM), low N (N 0 µM), low Fe (Fe 0 µM), low Fe low

Pi (Fe 0 µM, P 0 µM) and normal Pi (P 625 µM) liquid media. (a, c, left) Bright field

(BF) images. (a, c, right) Aniline blue staining images. White arrowheads indicate root

hair tips. (b) Aniline blue and propidium iodide (PI) staining of root hairs under low Pi

and normal Pi conditions. (c) The wild-type (WT) used for *rhd6* was Ws-2. Magnified

images of the upper panels (a, c, lower panel). Bars, 500 µm (a, c, upper panel), 50 µm

(a, c lower panel) and 25 µm (b).

Figure 2. Callose synthase *PMR4* is required for callose induction in root hairs

under phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis*

thaliana plants to low Pi (0 µM) liquid media. Left, bright filed (BF) images. Right,

aniline blue staining images. Two *caps* mutants are defective in inducing root hair

callose under phosphate starvation. Lower panels, magnified images of upper panels.

White arrowheads indicate root hair tips. Bars, 500 µm (upper panel) and 50 µm (lower

panel). (b) Positions of mutated sites in the *PMR4* locus of *caps* mutants, illustrated

with Protter (Omasits *et al*., 2014). An orange region corresponds to the catalytic

domain of PMR4. Asterisks indicate stop codons. (c) Ratios of *G. orontii* genomic DNA

to Arabidopsis genomic DNA were determined by qPCR with the primers for Plasma

membrane ATPase 1 (Go_V1_Conting3757) and RNA-binding family protein

(AT3G21215), respectively. Data show means ± SD (n=3). Asterisks (** $P < 0.01$)

indicate statistically significant differences relative to WT according to Student's *t*-test.

Figure 3. *PMR4* modulates root and shoot growth under phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis*

seedlings to low Pi (50 µM) liquid media. Left, bright filed (BF) images. Right, aniline

blue staining images. Lower panels, magnified images of upper panels. White

arrowheads indicate root hair tips. Bars, 100 µm (upper panel) and 50 µm (lower panel).

(b) Six-d-old seedlings were exposed to low Pi (50 µM) or normal Pi (625 µM) media

for 6 d. Bar, 10 mm. (c) The primary roots 6 d after transfer of 6-d-old seedling to low

Pi (50 µM) and normal Pi media. Bar, 1 mm. (d) Root hair density and root hair length

in (c). (e) Primary root length in (b). (f) Ratio of the primary root length under normal

Pi relative to low Pi of (e). (g) Convex hull area of the roots in (b). (h) Ratio of convex

hull area under normal Pi relative to low Pi of (g). (i) Six-d-old seedlings were exposed

to low Pi (50 µM) or normal Pi for 16 d. Bar, 10 mm. (j) SFW in (i). (k) Six-d-old

seedlings were exposed to low Pi (10 µM) media for 10 d. Bar, 10 mm. (l) Anthocyanin

contents in (k). (d-h, j, l) Data show means ± SD (n = 7~30). Statistical significance was

assessed by one-way ANOVA and Tukey honestly significant different (HSD) test for

each medium dataset. Different letters indicate significant differences ($P < 0.05$). NS,

not significant.

Figure 4. *PMR4* contributes to shoot Pi accumulation under phosphate deficiency.

(a-b) Pi contents per root or shoot fresh weight (FW) of 6-d-old *Arabidopsis* seedlings

exposed to low Pi (150 µM) or normal Pi (625 µM) for 9 d. Data show means ± SD (n =

3~4) in (a-b). Statistical significance was assessed by one-way ANOVA and Tukey

HSD test for each medium dataset (a-b). Different letters indicate significant differences

($P < 0.05$). NS, not significant.

Figure 5. *PMR4* contributes to Pi translocation under phosphate deficiency.

(a-b) Pi contents per root or shoot fresh weight (FW) of 6-d-old *Arabidopsis* seedlings

exposed to low Pi (150 µM) or normal Pi (625 µM) for 9 d. (c) Illustration of Pi

accumulation in the indicated genotypes, referred to (Figure 4a, 5a-b). (d) Ratio of ^{32}P

in the shoots relative to the whole plants after pulse-chase labeling. Seedlings grown on

low and normal Pi media (50 and 625 µM, respectively) were exposed to ^{32}P-labeled Pi

for 10 min, and then transferred to non-labelled Pi media for 120 min (pulse chase). The

^{32}P ratio of the shoots to the whole seedlings was calculated for the translocation rate of

Pi. Data show means ± SD (n = 3~4) in (a-b, d). Asterisks (** $P < 0.01$, * $P < 0.05$)

indicate statistically significant differences relative to WT according to Student's *t*-test

(a-b). Statistical significance was assessed by one-way ANOVA and Tukey HSD test

for each medium dataset (d). Different letters indicate significant differences ($P < 0.05$).

NS, not significant.

Figure 6. *PMR4* contributes to Pi uptake under phosphate deficiency.

(a-b) Pi contents per root or shoot fresh weight (FW) of 6-d-old *Arabidopsis* seedlings

exposed to low Pi (150 µM) or normal Pi (625 µM) for 9 d. The WT used for *rhd6* was

Ws-2 (c) Illustration of Pi accumulation in the indicated genotypes, referred to (Figure

4a, 6a-b). (d) ^{32}P uptake rate determined in the whole seedlings exposed to the indicated

media (low Pi and normal Pi, 50 and 625 µM Pi, respectively) supplemented with

^{32}P-labelled Pi for 30 min. Data show means ± SD (n = 3~4) in (a-b, d). Asterisks (** P

< 0.01, * $P < 0.05$) indicate statistically significant differences relative to WT according

to Student's *t*-test (a-b). Statistical significance was assessed by one-way ANOVA and

Tukey HSD test for each medium dataset (d). Different letters indicate significant

differences ($P < 0.05$). NS, not significant.

Figure 7. Plant element contents in shoot and root under low phosphate conditions.

Element contents (P, K, Mg, Ca, Mn, Fe, Zn, Mo and B) per root or shoot dry weight

(DW) of 5-d-old *Arabidopsis thaliana* seedlings exposed to low Pi (10 μM) for 4 d.

Data show means ± SD (n=4~8). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate

statistically significant differences relative to WT according to Student's *t*-test. NS, not

significant. ND, no data.

Figure 8. Root *PMR4* contributes to shoot growth under phosphate deficiency.

(a) Reciprocal micrografting of *Arabidopsis* WT and *pmr4* mutants. Images were

obtained at 14 d after transfer to low Pi (150 μM) media. Bar, 10 mm. (b) Shoot fresh

weight (SFW) of (a). (c) The primary root length of (a). (d) Aniline blue staining of

grafted-seedlings at 9 d after transfer to low Pi (50 μM). (d, left) Bright filed (BF) images.

(d, right) Aniline blue staining images. (d, lower panel) Magnified images of (d, upper

panel). White arrows indicate root hair tips. Bar, 100 μm (c, upper panel) or 20 μm (c,

lower panel). (b, c) Data show means ± SD (n=19-26). Statistical significance was

assessed by one-way ANOVA and Tukey HSD test for each medium dataset. Different

letters indicate significant differences ($P < 0.05$).

Figure 9. PMR4 in root hairs contributes to shoot growth under phosphate

deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis*

seedlings to low Pi (10 μM). White arrowheads indicate root hair chips. Bars, 100 μm

(upper panel) and 50 μm (lower panel). (b) Five-d-old seedlings were exposed to low Pi

998   (50 µM) for 10 d. Bar, 10 mm. (c) SFW of (b). (d) Primary root length of (b). (c, d)

999   Data show means ± SD (n = 22-48). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate

1000   statistically significant differences relative to *pmr4* according to Student's *t*-test. NS, not

1001   significant.

1002

Figure 10. Model for PMR4-mediated root hair callose deposition in plant

adaptation to Pi deficiency.

PMR4 induces callose deposition in root hairs under phosphate deficiency, and thus

contributes to shoot Pi accumulation and growth through enhancing Pi uptake and

root-to-shoot transfer. This contribution alleviates phosphate starvation symptoms, such

as root system area reduction and anthocyanin accumulation.

(a)

(b)

(c)

Figure 1. Callose deposition is induced in root hairs under phosphate deficiency, independently of *PHR1, PHL1, LPR1* and *LPR2*.

Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis thaliana* seedlings to low Pi (P 50 µM), low N (N 0 µM), low Fe (Fe 0 µM), low Fe low Pi (Fe 0 µM, P 0 µM) and normal Pi (P 625 µM) liquid media. (a, c, left) Bright field (BF) images. (a, c, right) Aniline blue staining images. White arrowheads indicate root hair tips. (b) Aniline blue and propidium iodide (PI) staining of root hairs under low Pi and normal Pi conditions. (c) The wild-type (WT) used for *rhd6* was Ws-2. Magnified images of the upper panels (a, c, lower panel). Bars, 500 µm (a, c, upper panel), 50 µm (a, c lower panel) and 25 µm (b).

Figure 2. Callose synthase *PMR4* is required for callose induction in root hairs under phosphate deficiency.
(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis thaliana* plants to low Pi (0 µM) liquid media. Left, bright filed (BF) images. Right, aniline blue staining images. Two *caps* mutants are defective in inducing root hair *callose under phosphate starvation.* Lower panels, magnified images of upper panels. White arrowheads indicate root hair tips. Bars, 500 µm (upper panel) and 50 µm (lower panel). (b) Positions of mutated sites in the *PMR4* locus of *caps* mutants, illustrated with Protter (Omasits *et al.*, 2014). An orange region corresponds to the catalytic domain of PMR4. Asterisks indicate stop codons. (c) Ratios of *G. orontii* genomic DNA to Arabidopsis genomic DNA were determined by qPCR with the primers for Plasma membrane ATPase 1 (Go_V1_Conting3757) and RNA-binding family protein (AT3G21215), respectively. Data show means ± SD (n=3). Asterisks (** *P* < 0.01) indicate statistically significant differences relative to WT according to Student's *t*-test.

Figure 3. *PMR4* modulates root and shoot growth under phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis* seedlings to low Pi (50 μM) liquid media. Left, bright filed (BF) images. Right, aniline blue staining images. Lower panels, magnified images of upper panels. White arrowheads indicate root hair tips. Bars, 100 μm (upper panel) and 50 μm (lower panel). (b) Six-d-old seedlings were exposed to low Pi (50 μM) or normal Pi (625 μM) media for 6 d. Bar, 10 mm. (c) The primary roots 6 d after transfer of 6-d-old seedling to low Pi (50 μM) and normal Pi media. Bar, 1 mm. (d) Root hair density and root hair length in (c). (e) Primary root length in (b). (f) Ratio of the primary root length under normal Pi relative to low Pi of (e). (g) Convex hull area of the roots in (b). (h) Ratio of convex hull area under normal Pi relative to low Pi of (g). (i) Six-d-old seedlings were exposed to low Pi (50 μM) or normal Pi for 16 d. Bar, 10 mm. (j) SFW in (i). (k) Six-d-old seedlings were exposed to low Pi (10 μM) media for 10 d. Bar, 10 mm. (l) Anthocyanin contents in (k). (d-h, j, l) Data show means ± SD (n = 7~30). Statistical significance was assessed by one-way ANOVA and Tukey honestly significant different (HSD) test for each medium dataset. Different letters indicate significant differences (*P* < 0.05). NS, not significant.

Figure 4. *PMR4* contributes to shoot Pi accumulation under phosphate deficiency.

(a-b) Pi contents per root or shoot fresh weight (FW) of 6-d-old *Arabidopsis* seedlings exposed to low Pi (150 µM) or normal Pi (625 µM) for 9 d. Data show means ± SD (n = 3~4) in (a-b). Statistical significance was assessed by one-way ANOVA and Tukey HSD test for each medium dataset (a-b). Different letters indicate significant differences ($P < 0.05$). NS, not significant.

Figure 5. *PMR4* contributes to Pi translocation under phosphate deficiency.

(a-b) Pi contents per root or shoot fresh weight (FW) of 6-d-old *Arabidopsis* seedlings exposed to low Pi (150 µM) or normal Pi (625 µM) for 9 d. (c) Illustration of Pi accumulation in the indicated genotypes, referred to (Figure 4a, 5a-b). (d) Ratio of ^{32}P in the shoots relative to the whole plants after pulse-chase labeling. Seedlings grown on low and normal Pi media (50 and 625 µM, respectively) were exposed to ^{32}P-labeled Pi for 10 min, and then transferred to non-labelled Pi media for 120 min (pulse chase). The ^{32}P ratio of the shoots to the whole seedlings was calculated for the translocation rate of Pi. Data show means ± SD (n = 3~4) in (a-b, d). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate statistically significant differences relative to WT according to Student's *t*-test (a-b). Statistical significance was assessed by one-way ANOVA and Tukey HSD test for each medium dataset (d). Different letters indicate significant differences ($P < 0.05$). NS, not significant.

Figure 6. *PMR4* contributes to Pi uptake under phosphate deficiency.

(a-b) Pi contents per root or shoot fresh weight (FW) of 6-d-old *Arabidopsis* seedlings exposed to low Pi (150 µM) or normal Pi (625 µM) for 9 d. The WT used for *rhd6* was Ws-2 (c) Illustration of Pi accumulation in the indicated genotypes, referred to (Figure 4a, 6a-b). (d) ^{32}P uptake rate determined in the whole seedlings exposed to the indicated media (low Pi and normal Pi, 50 and 625 µM Pi, respectively) supplemented with ^{32}P-labelled Pi for 30 min. Data show means ± SD (n = 3~4) in (a-b, d). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate statistically significant differences relative to WT according to Student's *t*-test (a-b). Statistical significance was assessed by one-way ANOVA and Tukey HSD test for each medium dataset (d). Different letters indicate significant differences ($P < 0.05$). NS, not significant.

Figure 7. Plant element contents in shoot and root under low phosphate conditions.

Element contents (P, K, Mg, Ca, Mn, Fe, Zn, Mo and B) per root or shoot dry weight (DW) of 5-d-old *Arabidopsis thaliana* seedlings exposed to low Pi (10 μM) for 4 d. Data show means ± SD (n=4~8). Asterisks (** $P < 0.01$, * $P < 0.05$) indicate statistically significant differences relative to WT according to Student's *t*-test. NS, not significant. ND, no data.

Figure 8. Root *PMR4* contributes to shoot growth under phosphate deficiency.

(a) Reciprocal micrografting of *Arabidopsis* WT and *pmr4* mutants. Images were obtained at 14 d after transfer to low Pi (150 μM) media. Bar, 10 mm. (b) Shoot fresh weight (SFW) of (a). (c) The primary root length of (a). (d) Aniline blue staining of grafted-seedlings at 9 d after transfer to low Pi (50 μM). (d, left) Bright filed (BF) images. (d, right) Aniline blue staining images. (d, lower panel) Magnified images of (d, upper panel). White arrows indicate root hair tips. Bar, 100 μm (c, upper panel) or 20 μm (c, lower panel). (b, c) Data show means ± SD (n=19-26). Statistical significance was assessed by one-way ANOVA and Tukey HSD test for each medium dataset. Different letters indicate significant differences (*P* < 0.05).

Figure 9. PMR4 in root hairs contributes to shoot growth under phosphate deficiency.

(a) Aniline blue staining of the primary roots 3 d after transfer of 5-d-old *Arabidopsis* seedlings to low Pi (10 µM). White arrowheads indicate root hair chips. Bars, 100 µm (upper panel) and 50 µm (lower panel). (b) Five-d-old seedlings were exposed to low Pi (50 µM) for 10 d. Bar, 10 mm. (c) SFW of (b). (d) Primary root length of (b). (c, d) Data show means ± SD (n = 22-48). Asterisks (** *P* < 0.01, * *P* < 0.05) indicate statistically significant differences relative to *pmr4* according to Student's *t*-test. NS, not significant.

Figure 10. Model for PMR4-mediated root hair callose deposition in plant adaptation to Pi deficiency.

PMR4 induces callose deposition in root hairs under phosphate deficiency, and thus contributes to shoot Pi accumulation and growth through enhancing Pi uptake and root-to-shoot transport. This contribution alleviates phosphate starvation symptoms, such as root system area reduction and anthocyanin accumulation.