Multicomponent *Ayurveda* formulation *Lodhrasavam* ameliorates steatosis and lipotoxicity in HepG2 cell model of NAFLD

Sania Kouser[1], Pranav Girish Banvi[1], Soumya Garawadmath[1], Subrahmanya Kumar[1], Chethala N Vishnuprasad[1*]

The University of Transdisciplinary Health Sciences and Technology (TDU), No.74/2, Jarakabande Kaval Post: Attur, Via Yelahanka, Bangalore – 560-106

*Correspondence: drcnvp@tdu.edu.in

Abstract:

Background: Non-alcoholic fatty liver disease (NAFLD) is a complex, multifactorial and multi-system disorder. It is one of the major contributors of liver disease worldwide. Among the many factors involved in the pathogenesis of NAFLD, free fatty acids (FFAs) such as palmitic acid induced lipotoxicity promotes steatosis, oxidative stress and insulin resistance that activate apoptotic cascades leading to tissue damage and inflammation. Since NAFLD is a multifactorial metabolic disorder, conventional target-based drug therapies have limited success. Therefore, the use of multicomponent ayurveda herbal formulations could be a promising alternative due to their multitargeted mechanisms of action. The present study investigates the effects and underlying mechanism of actions of an ayurveda formulation, *Lodhrasavam* (TDU-LS-1), in the in-vitro model of NAFLD.

Methods: Lipotoxicity was induced in HepG2 cells by treating the cells with 1mM palmitic acid for 24 hrs followed by drug (TDU-LS-1) treatment for another 24 hrs. The effect of TDU-LS-1 on lipotoxicity was evaluated by MTT assay. The effect of TDU-LS-1 on steatosis was studied by estimating intracellular triglycerides, lipid droplets formation and expression of genes involved in lipid metabolism. Further, to examine the antioxidant activity, DPPH scavenging assay was performed.

Results/discussion: TDU-LS-1 was found to increase the antioxidant activity in a concentration dependent manner with an IC50 of 16.45 μg GAE/ml. Palmitic acid induced lipotoxicity in HepG2 cells was reduced by lower concentrations of TDU-LS-1. Also, the results from triglyceride (TAG) assay, Oil-Red-O staining and BODIPY 493/503 confocal imaging suggest that TDU-LS-1 reduces the palmitate induced triglyceride deposition and lipid droplet

accumulation in HepG2 cells. Further, the qRT-PCR analysis of TDU-LS-1 at a concentration of 32 μg/GAE revealed that it modulates the expression of SREBP, FASN, SCD1, ACOX, and PPARγ that are relevant in hepatic lipid metabolism. Our results suggest that TDU-LS-1 can reduce *de novo* lipogenesis, peroxisomal lipid peroxidation as well as lipotoxicity in the *in vitro* palmitate-induced NAFLD model in HePG2 cells.

Keywords: NAFLD, *Lodhrasavam*, lipotoxicity, HepG2, steatosis

Introduction:

Non-alcoholic fatty liver disease (NAFLD) is a complex, multisystem disease, extending its effect on extrahepatic organs and regulatory pathways (Byrne and Targher, 2015). High-calorie diet and sedentary lifestyle are typical risk factors for NAFLD. However, many other factors including, but not limited to, type 2 diabetes, insulin resistance, obesity, hypertension, metabolic syndrome, genetic variations, alterations in gut permeability and microbiome are also playing a crucial role in its development and progression (Carr et al., 2016; Friedman et al., 2018; Younossi et al., 2018). It is one of the major contributors of liver disease worldwide with an estimated prevalence of 25% in the adult population (Younossi et al., 2019). In India its prevalence varies between 9% in rural and to 35% in urban populations (Duseja, 2010). When left untreated, NAFLD leads to deleterious liver physiology like steatohepatitis, fibrosis and cirrhosis that can further increase the risk of liver cancer and end-stage liver disease (Haas et al., 2016; Michelotti et al., 2013). Also, the renaming of NAFLD as Metabolic-Associated Fatty Liver Disease (MAFLD) by a consensus arrived by a panel of experts is the reflection of its complex etiopathology and patient heterogeneity as well as the multidirectional relationship with other metabolic disorders (Eslam et al., 2020).

Despite its high clinical and socioeconomic burden there are no approved therapeutics for NAFLD and the current clinical management strategies are largely centered around controlling the predisposing factors like diabetes and obesity (Mundi et al., 2020). One of the reasons for this is the inadequacy of conventional single target-based drugs to address the complex etiopathology of NAFLD as well as the limited knowledge we have on the systemic biology of this disease. Therefore, frontiers of biology started looking at innovative systemic strategies for addressing the multifactorial pathophysiology of NAFLD. This is where a transdisciplinary approach of integrating biomedicine's molecular perspective of NAFLD with its systemic

counterpart from traditional medicines like *Ayurveda* become relevant and can probably open up new opportunities for better holistic strategies for the prevention and management of NAFLD.

One of the characteristics of NAFLD is the accumulation of lipid droplets (LDs) in the hepatocytes leading to a condition referred to as hepatic steatosis, and it happens via flux of free fatty acids (FFAs) from adipose tissue due to increased lipolysis, *de novo lipogenesis* (DNL), excess dietary fat intake, decreased export of triglycerides and reduced beta-oxidation (Alkhouri et al., 2009; Schulze and McNiven, 2019). Although *Ayurveda* has an epistemologically and ontologically different understanding of NAFLD from biomedicine, both schools emphasize the involvement of metabolic imbalances and the role of diabetes (*Prameha*) and obesity (*Sthoulya*) as predisposing factors in NAFLD. Previous studies from our laboratory by Butala et al., 2017 reported the anti-diabetic and anti-obesity effects of *Lodhrasavam*, a polyherbal formulation in *Ayurveda* prescribed for obese-diabetic patients. According to Ayurvedic pharmacology principles, the key herbal ingredients of this formulation belong to the *Lodhradi-gana* group of plants, a group having *medo-hara* (anti-obesity) and *kapha-hara* properties and both are essential for the management of glucose and lipid homeostasis in the body. Considering the profound ant-obesity and anti-lipidemic properties of *Lodhrasavam* as well as its rich herbal-ingredient diversity the present study aimed at evaluating its potential effect on reducing lipotoxicity and steatosis in free fatty acid (FFAs) induced liver model.

Materials and methods

Chemicals and Reagents

HepG2 human hepatoma cell line (NCCS, Pune, India); Dulbecco's modified Eagle's medium (DMEM) (GIBCO, Grand Island, New York); fetal bovine serum (FBS) (GIBCO, Grand Island, New York); penicillin/streptomycin (GIBCO, Grand Island, New York); fatty acid free bovine serum albumin (BSA) (Genei, Bangalore, India); sodium palmitate (PA) (Sigma-Aldrich, St. Louis, MO, USA); 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (SRL, India); triglyceride kit (BeneSphera, Avantor, USA); 2,2-Diphenyl-1-picrylhydrazyl (DPPH) (Sigma-Aldrich, St. Louis, MO, USA); Ascorbic Acid (Sigma-Aldrich, St. Louis, MO, USA); Folin – Ciocalteu reagent (Sigma-Aldrich, St. Louis, MO, USA); gallic acid (Sigma-Aldrich, St. Louis, MO, USA); BODIPY 493/503 (Cayman chemical 25892); Hoechst 33342 (Cayman chemical 15547); Oil-Red-O stain (Sigma-Aldrich, St. Louis, MO, USA); trizol reagent (Takara

Bio, Shiga, Japan); PrimeScript 1st strand cDNA Synthesis Kit (Takara Bio, Shiga, Japan); PrimeScript RT-PCR Kit (Takara Bio, Shiga, Japan)

Determination of total tannin content

TDU-LS-1 was estimated for total tannins following standard Folin - Ciocalteu method (Ainsworth and Gillespie, 2007). Briefly, 10 μL of TDU-LS-1 with 40 μL of water, 50 μL FC reagent (previously diluted with water 1:10 v/v) and 100 μL of 3.5% Na_2CO_3 was incubated at room temperature for 30 min. A set of gallic acid standards (50, 25, 12.5, 6.25, 3.125 μg/mL) were prepared in the same manner. The absorbance was measured at 760 nm using plate reader (xMark Microplate Spectrophotometer, BioRad, USA). The experimental concentrations of test samples were expressed as μg of gallic acid equivalent tannin (GAE)/mL of sample.

Determination of DPPH radical scavenging activity

Free radical scavenging ability of different concentrations of TDU-LS-1 was tested by DPPH radical scavenging assay (Rahman et al., 2015). DPPH produces violet/purple color in ethanol solution and fades to shades of yellow color in the presence of antioxidants. A solution of 0.1 mM DPPH in ethanol was prepared, and 2.4 mL of this solution was mixed with 1.6 mL of TDU-LS-1 in ethanol to get a concentration gradient (128, 64, 32, 16, 8, 4, 2 μg GAE/ml). The reaction mixture was vortexed thoroughly and left in the dark at room temperature (RT) for 30 min. The absorbance of the mixture was measured spectrophotometrically at 517 nm. Ascorbic acid was used as reference. Percentage DPPH radical scavenging activity was calculated by the following equation: % DPPH radical scavenging activity= $\{(A_0 - A_1)/A_0\} \times 100$; where A_0 is the absorbance of the control, and A_1 is the absorbance of TDU-LS-1 or the standard. The % of inhibition was plotted against concentration, and from the graph IC_{50} was calculated. The experiment was performed in triplicates and repeated three times. Ascorbic acid Equivalent Antioxidant Capacity (AEAC) was calculated using the formula: IC_{50} (AA) /IC_{50} (sample). Where, IC_{50}(AA) is IC_{50} of standard (i.e., Ascorbic acid) and IC_{50}(sample) is IC_{50} of sample.

Cell Culture and establishment of palmitic acid induced NAFLD model

HepG2 cells were cultured in DMEM containing 10% FBS and 1% penicillin/streptomycin in a humidified atmosphere containing 5% CO2 at 37°C. The cells were seeded in multi-well plates and grown for 48 hrs. The medium was changed to DMEM (serum free) containing 5% fatty-

acid-free bovine serum albumin (BSA) with or without 1mM palmitic acid (PA) and grown for 24 h. Further, the cells were treated with various concentrations of TDU-LS-1 in the same BSA-medium and incubated for 24 h. The changes in lipid droplet formation were observed and quantified using various methods as described in the below sections.

Cytotoxicity assay

The cytotoxic effects of various concentrations of TDU-LS-1 in HepG2 cells were studied using MTT assay, which is based on the reduction of a tetrazolium salt by mitochondrial dehydrogenase in viable cells. HepG2 cells were seeded in 96-well plates at the density of 1.5×10^4 cells/well and cultured in 100μL of DMEM containing 10% FBS overnight. The medium was then discarded and was replaced with 128, 64, 32, 16, 8, 4, 2, 0 μg GAE/ml TDU-LS-1 diluted in the culture medium, followed by incubation for a further 24 h at 37˚C. 500 μg/ml of MTT working solution were then added to each well followed by incubation for 4 h at 37˚C. The violet formazan crystal in each well was dissolved in DMSO and the absorbance of each well was measured at 570 nm using a microplate reader (xMark Microplate Spectrophotometer, BioRad, USA). HepG2 cells were treated with 1mM PA in 5% BSA media for 24 h at 37˚C, followed by treatment with 128, 64, 32, 16, 8, 4, 0 μg GAE/ml TDU-LS-1 for 24 h at 37˚C. Further, to assess they cytotoxicity effect MTT assay was performed as described.

BODIPY 493/503 staining of lipid droplets

HepG2 cells were seeded in 35mm confocal dishes, treated with 1 mM PA for 24 h, and then cultured with 32 μg GAE/ml TDU-LS-1 for 24 h at 37˚C. Cells were stained with 10μM BODIPY 493/503 dye and 10μg/ml Hoechst dye and incubated for 30 mins at 37°C (Qiu and Simon, 2016). Cells were washed with PBS, visualized under the LSM880 confocal live cell imaging system, (Carls Ziess, Germany) and images were taken at 40x magnification.

Oil Red O (ORO) staining of intracellular lipid accumulation

HepG2 cells were seeded in six-well plates (1.2×10^6 cells/well), treated with 1 mM PA for 24 h, and then cultured with 32 μg GAE/ml TDU-LS-1 for 24 h at 37˚C. Cells were washed twice with PBS, and fixed with 4% formaldehyde for 30 mins. After washing, cells were stained for 20 min in freshly diluted ORO solution (0.5% ORO in isopropanol diluted to 3:2 with H_2O, and filtered) at 37 °C. After staining, the cells were thoroughly washed with double distilled water to remove

the unbound staining solution and images of cells stained with Oil Red were captured using phase contrast microscope (Olympus-IX- Olympus America Inc, USA). To quantify the ORO contents, isopropanol was added to each well and the optical density of the samples at a wavelength of 450 nm was measured using a microplate reader (xMark Microplate Spectrophotometer, BioRad, USA) (Mehlem et al., 2013).

Triglyceride (TG) estimation

To analyze the content of cellular triglycerides, the cells were washed with PBS, scraped into 250µl 0.1% PBST and pulse sonicated for 45 % amplitude for 45 seconds. The lysates were assayed for their total triglyceride content using assay kit and cellular protein was estimated using the Bradford method. The results were expressed as µg of triglyceride per mg of cellular protein.

Quantification real-time PCR

Total RNA was isolated from treated HepG2 cells using trizol reagent according to the manufacturer's instructions, and the absorbances of the extracted RNAs at 260 nm and 280 nm were determined spectrophotometrically (Nanodrop, Thermofisher Scientific). RNA samples with purity ratios (A260/A280) between 1.8 and 2.0 were used for synthesizing single-strand cDNA by means of PrimeScript 1st strand cDNA Synthesis Kit for quantitative real-time polymerase chain reaction (qPCR). Gene-specific primers were designed (Table 2) (Bioserve, Hyderabad, India) and qPCR was performed to detect the relative mRNA expression levels using PrimeScript RT-PCR Kit as described in the protocol. To determine the specificity of the amplification, melting curve analysis was performed for all final PCR products. Relative changes in gene expression were calculated and expressed as fold changes using the relative quantification ($\Delta\Delta$Ct) method. The relative abundance of each transcript was normalized to that of GAPDH. Each sample was run in triplicate.

Statistical analysis

All results are expressed as the means $\pm$ standard deviations (SDs). A one-way ANOVA test was performed to determine the significance of test samples compared to the controls and a value of $p < 0.05$ was considered as significant.

Results

TDU-LS-1 increases DPPH radical scavenging activity in a concentration dependent manner

The antioxidant effect of TDU-LS-1 was studied using DPPH assay and the results showed a concentration dependent increase in radical scavenging activity by the formulation. The highest concentration of 128 µg GAE/ml tested in the assay showed 90.40% ± 1.39% inhibition of DPPH assay whereas 15.33% ± 5.38% inhibition was observed at the lowest concentration of 2 µg GAE/ml (Fig-1). IC_{50} of TDU-LS-1 is 16.45 µg GAE/ml and IC_{50} of ascorbic acid is 24.51 µg/ml. AEAC (Ascorbic acid equivalent antioxidant capacity) was found to be 1.49 (Fig 1).

TDU-LS-1 reduces cytotoxicity of HepG2 cells in palmitic acid induced NAFLD model

Different concentrations of TDU-LS-1 were tested on HepG2 cells for assessing the cytotoxicity of the formulation. Results of 24 hr treatment showed that only the highest concentration (128 µg GAE/ml) used in the experiment caused any significant cytotoxicity in HepG2 cells. It showed 25.67% ± 10.48% viability whereas the lowest concentration tested (2 µg GAE/ml) showed 92.66% ± 9.43% viability after 24 h treatment (Fig-2A). An *in vitro* model of NAFLD was established in HepG2 cells by inducing the cells with palmitic acid for 24 hrs. HepG2 cells were treated with various concentrations of PA and observed for cytotoxicity effects. Cells treated with 1mM PA reduced the viability of cells to 37.64% ± 5.71% along with increasing lipid and triglyceride accumulation. In order to study the effect of TDU-LS-1 on protecting the PA-induced cytotoxicity of HepG2 cells. On the other hand, TDU-LS-1 treatment post 1mM palmitic acid induction was found to reverse the palmitic acid-induced cytotoxicity, especially at lower concentrations of the formulation like 4, 8, 16 and 32 µg GAE/ml. These concentrations of TDU-LS-1 increased the viability of PA treated cells to 67.46% ± 4.9%, 64.81% ± 5.06%, 60.69% ± 3.44% and 41.64% ± 3.70% respectively. Although 64 µg GAE/ml TDU-LS-1 was non-toxic in the MTT assay, this concentration was found to be ineffective in the palmitic acid treatment (Fig 2 A-B).

TDU-LS-1 reduces intracellular lipid droplet and triglyceride

The *in vitro* model of PA induced hepatic steatosis was used to investigate the efficacy of TDU-LS-1 against hepatic steatosis. BODIPY 493/503 staining showed a 1.72-fold increase in mean fluorescence intensity in 1mM PA treated cells when compared to untreated cells (Fig 3 E-F). Further, it was found that 32µg GAE/ml TDU-LS-1 significantly reduced the mean fluorescence

intensity by 0.72-fold. Similar to BODIPY staining, cells stained with ORO also showed that PA treatment for 24 h significantly induced significant intracellular lipid accumulation in HepG2 cells. The absorbance of the stained cells in the PA-treated model group was 1.2 times that of the cells in the control group. However, the intracellular lipid accumulation in 32μg GAE/ml TDU-LS-1 treated cells significantly reduced the lipid droplet accumulation by 1.14-fold in comparison to PA induced cells (Fig 3 A-D). Further, in the PA treated cells the TG levels increased 1.5-fold when compared to untreated cells. However, 32 μg GAE/ml TDU-LS-1 treated cells significantly reduced the TG levels by 1.10-fold in comparison to PA induced cells (Fig 3 G).

TDU-LS-1 attenuates lipogenic mRNA expressions induced by PA

To investigate how TDU-LS-1 ameliorated fat deposition in the liver, the expressions of genes associated with lipid metabolism in HepG2 cells were examined using qRT-PCR. The mRNA expression levels of *PPARγ, SREBP-1C, FASN*, *SCD1* and ACOX1 significantly increased in PA treated cells by 6.2-fold, 3.2-fold, 2.0-fold, 1.9-fold and 2.36-fold respectively in comparison to untreated cells. 32 μg GAE/ml TDU-LS-1 significantly reduced the expression of *PPARγ* (2.62-fold), *SREBP-1C* (2.21-fold), *FASN* (1.39-fold), *SCD1* (1.54-fold) and ACOX1 (1.75) when compared with PA treated cells (Fig 4A-C)

Discussion

In previous studies conducted in our lab *Lodhrasavam* has shown potential anti-diabetic and anti-adipogenic effects. It inhibited α-amylase and α-glucosidase activity by 90% and 78% respectively and reduced adipocyte differentiation significantly (Butala et al., 2017). It has not been explored for its anti-steatosis and antioxidant activities. In the present study we aimed at evaluating the effect of TDU-LS-1 on saturated fatty acid (palmitate) induced hepatic steatosis and lipotoxicity.

In a case study conducted by Sahu et al., 2022, it was found that *Ayurveda* formulations (*Avipattikara Churna, Punarnava Mandura, Sankha Bhasma, Tarunikusumakara Churna, Triphala Churna* and *Kutaki Churna*) have immense potential in treating grade 2 fatty liver disease. A similar effect was seen in a randomized placebo-controlled clinical trial of a multiherbal formulation, *Sharapunkhadi* powder on subjects with grade 1-3 fatty liver disease (Remya et al., 2020). Livogrit, a tri-herbal *Ayurvedic* medicine, was studied for its

hepatoprotective effect on methionine and cysteine deficient NASH models in both HepG2 spheroids and rat primary hepatocytes. It reduced fat accumulation and ROS production and increased lipolysis hence attenuating NASH (Balkrishna et al., 2022)

NAFLD is characterized by the accumulation of lipid droplets (LDs) in the hepatocytes, referred to as hepatic steatosis. It is considered as the "first-hit" in the development of NAFLD. LDs can accumulate in the liver via three mechanisms: a) flux of free fatty acids (FFAs) from adipose tissue due to increased lipolysis, increased *de novo lipogenesis* (DNL), and dietary fat intake; b) decreased export of triglycerides and c) reduced beta-oxidation. Elevated plasma free fatty acid (FFA) levels play an etiological role in the pathogenesis of NAFLD. In particular, the saturated fatty acid palmitate, which makes up 30%-40% of high plasma FFA concentration, is a major contributor to an increase in intrahepatic triglyceride (Cacicedo et al., 2005; Schroeder-Gloeckler et al., 2007). In the present study, 1 mM PA was used to induce cellular steatosis (Gómez-Lechón et al., 2007). We found that the accumulation of lipid droplets and the intracellular TG content significantly increased in FFA-treated HepG2 cells. TDU-LS-1 treatment of HepG2 cells exhibited an inhibitory effect on FFA-induced hepatic steatosis.

To investigate the mechanism of TDU-LS-1 inhibition of hepatic steatosis, we evaluated the expression of key genes involved in lipogenesis, including SREBP-1c, FASN and SCD-1 (Parlati et al., 2021). SREBP-1c is a master regulator of lipogenesis, known to transcriptionally regulate downstream genes involved in de novo lipogenesis, including those encoding ACC and FAS (Osborne, 2000). In a study by Motoyuki et al., 2007, it showed an increase in expression of SREBP-1c, ACC and FAS in NAFLD patients. It has been reported that SREBP-1c activation is associated with increased lipogenesis in NAFLD (Yan et al., 2014). In mice, HFD caused fat accumulation and induced SREBP-1c mRNA expression which further contributes to the progression of hepatic steatosis (Kim et al., 2010; Zhu et al., 2019). Li et al., 2011 showed that inhibiting SREBP-1c activity can exert an anti-hepatic steatosis effect in diet-induced obese mice. SCD-1 is an endoplasmic reticulum bound rate limiting enzyme that catalyzes the formation of mono-unsaturated fatty acids from saturated fatty acids (Paton and Ntambi, 2009) Liver specific knock-out of SCDI in mice fed with high-carbohydrate diet showed reduction in hepatic steatosis (Miyazaki et al., 2007) (Jiang et al., 2005) in their experiments using antisense oligonucleotide inhibitor of SCD1 showed inhibition of lipogenesis and promoted lipid oxidation in both primary mouse hepatocytes and HFD fed mice. In the current study, 24hr incubation with

1mM PA increased the expression of these genes. Following which TDU-LS-1 significantly downregulated the expression of SREBP-1c, FAS and SCD-1.

The hepatic PPARγ plays a putative role in the progression of fatty liver disease in the NAFLD patients. PPAR-γ is up-regulated in the liver of obese patients with NAFLD, and recently, the expression of PPAR-γ is considered as an additional reinforcing lipogenic signal, assisting SREBP-1c to trigger the development of hepatic steatosis (Pettinelli and Videla, 2011). We showed that PA increased PPAR-γ mRNA expression in HepG2 cells, which was significantly reduced by TDU-LS-1.

When there is an excess of fatty acids in hepatocytes, alternative pathways of fatty acid oxidation are activated, such as ß-oxidation in peroxisomes. The peroxisomal acylCoA oxidases ACOX is one of the first and rate-limiting enzymes of ß-oxidation pathways in peroxisomes (Reddy and Hashimoto, 2001). Expression of ACOX increase in NAFLD patients indicating that peroxisomal ß-oxidation is compensatively enhanced in NAFLD (Motoyuki et al., 2007). In our study we observed an increase in ACOX gene expression in PA treated cells, whereas in TDU-LS-1 treated cells a significant decrease in its expression was shown. TDU-LS-1 reduced the ß-oxidation in peroxisomes indicative of its role in lowering the excessive ROS production which occurs as a result of excessive oxidation of fatty acids. Further studies need to be performed to understand the effect of TDU-LS-1 in lowering oxidative stress induced by palmitic acid.

Conclusion

Excessive accumulation of lipid droplets in the hepatocytes is the primary hallmark of NAFLD. Therefore, reducing the amount of lipid droplets and inhibiting *de novo* lipogenesis (DNL) are considered as important treatment strategies for NAFLD. Our study provides experimental evidence for the anti-steatosis effect of TDU-LS-1. It exerts its effects through reducing the lipid droplet formation in hepatocytes and inhibiting key genes involved in DNL. In addition, radical scavenging activities of TDU-LS-1 makes it more beneficial in addressing the complexities of NAFLD pathophysiology. Further studies need to be conducted to understand the holistic mechanism of action of this complex multicomponent *Ayurveda* formulation for managing NAFLD.

Acknowledgements

The authors acknowledge ICMR for granting Sania Kouser the ICMR-SRF fellowship as well as the fund support received from JM Financial and Rural India Support Trust (RIST) for conducting experiments.

Conflict of interest

Authors declare no conflict of interest.

Figure legends

Figure 1: TDU-LS-1 increases DPPH radical scavenging activity in a concentration dependent manner. IC50 of TDU-LS-1 is 16.45 μg GAE/ml.

Figure 2: TDU-LS-1 reduces cytotoxicity of HepG2 cells in palmitic acid induced NAFLD model. A) In the 24 hr treatment of HepG2 cells the highest concentration of 128 μg GAE/ml showed 25.67% ± 10.48% viability while the lower concentrations did not significantly affect the viability of the cells. B) 1mM PA reduced the viability of cells to 37.64% ± 5.71% and treatment with various concentrations of TDU-LS-1 gradually increased the viability as the concentration decreased.

Figure 3: TDU-LS-1 inhibits intracellular lipid droplet and triglyceride accumulation. A-C) After ORO staining it was observed that 32 μg GAE/ml of TDU-LS-1 reduced the lipid droplets induced by 1mM PA; D) the intracellular lipid accumulation in 32 μg GAE/ml TDU-LS-1 treated cells significantly reduced by 1.14-fold in comparison to PA induced cells; E) Untreated cells (E1-E3); 1mM PA treated cells (E4-E6); 32 μg GAE/ml TDU-LS-1 treated cells (E7-E9). Live cells stained with BODIPY 493/503 (Green) and Hoechst dyes (Blue); F) 1.72-fold increase in mean fluorescence intensity in 1mM PA treated cells when compared to untreated cells and 32 μg GAE/ml TDU-LS-1 significantly reduced the mean fluorescence intensity by 0.72-fold was observed.

Figure 4: Treatment with 1mM PA increased the expression of the lipogeneic genes while 32 μg GAE/ml TDU-LS-1 significantly reduced their expression (A-B). 32 μg GAE/ml TDU-LS-1

treated cells significantly decreased the expression of ACOX gene while it increased the expression of CPT1A (C)

References

1. Ainsworth, E.A., Gillespie, K.M., 2007. Estimation of total phenolic content and other oxidation substrates in plant tissues using Folin-Ciocalteu reagent. Nat Protoc 2, 875–877. https://doi.org/10.1038/nprot.2007.102

2. Alkhouri, N., Dixon, L.J., Feldstein, A.E., 2009. Lipotoxicity in nonalcoholic fatty liver disease: Not all lipids are created equal. Expert Rev Gastroenterol Hepatol. https://doi.org/10.1586/egh.09.32

3. Askari, F., Rashidkhani, B., Hekmatdoost, A., 2014. Cinnamon may have therapeutic benefits on lipid profile, liver enzymes, insulin resistance, and high-sensitivity C-reactive protein in nonalcoholic fatty liver disease patients. Nutrition Research 34, 143–148. https://doi.org/10.1016/j.nutres.2013.11.005

4. Balkrishna, A., Gohel, V., Kumari, P., Manik, M., Bhattacharya, K., Dev, R., Varshney, A., 2022. Livogrit Prevents Methionine-Cystine Deficiency Induced Nonalcoholic Steatohepatitis by Modulation of Steatosis and Oxidative Stress in Human Hepatocyte-Derived Spheroid and in Primary Rat Hepatocytes. Bioengineered 13, 10811–10826. https://doi.org/10.1080/21655979.2022.2065789

5. Butala, M.A., Kukkupuni, S.K., Vishnuprasad, C.N., 2017. Ayurvedic anti-diabetic formulation Lodhrasavam inhibits alpha-amylase, alpha-glucosidase and suppresses adipogenic activity in vitro. J Ayurveda Integr Med 8, 145–151. https://doi.org/10.1016/j.jaim.2017.03.005

6. Byrne, C.D., Targher, G., 2015. NAFLD: A multisystem disease. J Hepatol. https://doi.org/10.1016/j.jhep.2014.12.012

7. Cacicedo, J.M., Benjachareowong, S., Chou, E., Ruderman, N.B., Ido, Y., 2005. Palmitate-Induced Apoptosis in Cultured Bovine Retinal Pericytes Roles of NAD(P)H Oxidase, Oxidant Stress, and Ceramide, Diabetes.

8. Carr, R.M., Oranu, A., Khungar, V., 2016. Nonalcoholic Fatty Liver Disease: Pathophysiology and Management. Gastroenterol Clin North Am. https://doi.org/10.1016/j.gtc.2016.07.003

9. Chou, Y.T., Li, C.H., Sun, Z.J., Shen, W.C., Yang, Y.C., Lu, F.H., Chang, C.J., Wu, J.S., 2021. A positive relationship between betel nut chewing and significant liver fibrosis in NAFLD subjects, but not in non-nafld ones. Nutrients 13, 1–12. https://doi.org/10.3390/nu13030914

10. Chou, Y.T., Sun, Z.J., Shen, W.C., Yang, Y.C., Lu, F.H., Chang, C.J., Li, C.Y., Wu, J.S., 2022. Cumulative Betel Quid Chewing and the Risk of Significant Liver Fibrosis in Subjects With and Without Metabolic Syndrome. Front Nutr 9. https://doi.org/10.3389/fnut.2022.765206

11. Daneshi-Maskooni, M., Keshavarz, S.A., Qorbani, M., Mansouri, S., Alavian, S.M., Badri-Fariman, M., Jazayeri-Tehrani, S.A., Sotoudeh, G., 2019. Green cardamom supplementation improves serum irisin, glucose indices, and lipid profiles in overweight or obese non-alcoholic fatty liver disease patients: A double-blind randomized placebo-controlled clinical trial. BMC Complement Altern Med 19. https://doi.org/10.1186/s12906-019-2465-0

12. Dhami-Shah, H., Vaidya, R., Talwadekar, M., Shaw, E., Udipi, S., Kolthur-Seetharam, U., Vaidya, A.D.B., 2021. Intervention by picroside II on FFAs induced lipid accumulation and lipotoxicity in HepG2 cells. J Ayurveda Integr Med 12, 465–473. https://doi.org/10.1016/j.jaim.2021.04.007

13. Dhami-Shah, H., Vaidya, R., Udipi, S., Raghavan, S., Abhijit, S., Mohan, V., Balasubramanyam, M., Vaidya, A., 2018. Picroside II attenuates fatty acid accumulation in HEPG2 cells via modulation of fatty acid uptake and synthesis. Clin Mol Hepatol 24, 77–87. https://doi.org/10.3350/cmh.2017.0039

14. Duseja, A., 2010. Nonalcoholic fatty liver disease in India - a lot done, yet more required! Indian Journal of Gastroenterology. https://doi.org/10.1007/s12664-010-0069-1

15. Eslam, M., Sanyal, A.J., George, J., Sanyal, A., Neuschwander-Tetri, B., Tiribelli, C., Kleiner, D.E., Brunt, E., Bugianesi, E., Yki-Järvinen, H., Grønbæk, H., Cortez-Pinto, H., Fan, J., Valenti, L., Abdelmalek, M., Romero-Gomez, M., Rinella, M., Arrese, M., Bedossa, P., Newsome, P.N., Anstee, Q.M., Jalan, R., Bataller, R., Loomba, R.,

Sookoian, S., Sarin, S.K., Harrison, S., Kawaguchi, T., Wong, V.W.S., Ratziu, V., Yilmaz, Y., Younossi, Z., 2020. MAFLD: A Consensus-Driven Proposed Nomenclature for Metabolic Associated Fatty Liver Disease. Gastroenterology 158, 1999-2014.e1. https://doi.org/10.1053/j.gastro.2019.11.312

16. Friedman, S.L., Neuschwander-Tetri, B.A., Rinella, M., Sanyal, A.J., 2018. Mechanisms of NAFLD development and therapeutic strategies. Nat Med. https://doi.org/10.1038/s41591-018-0104-9

17. Gómez-Lechón, M.J., Donato, M.T., Martínez-Romero, A., Jiménez, N., Castell, J.V., O'Connor, J.E., 2007. A human hepatocellular in vitro model to investigate steatosis. Chem Biol Interact 165, 106–116. https://doi.org/10.1016/j.cbi.2006.11.004

18. Haas, J.T., Francque, S., Staels, B., 2016. Pathophysiology and Mechanisms of Nonalcoholic Fatty Liver Disease. Annu Rev Physiol. https://doi.org/10.1146/annurev-physiol-021115-105331

19. Huang, C.Z., Tung, Y.T., Hsia, S.M., Wu, C.H., Yen, G.C., 2017. The hepatoprotective effect of Phyllanthus emblica L. fruit on high fat diet-induced non-alcoholic fatty liver disease (NAFLD) in SD rats. Food Funct 8, 842–850. https://doi.org/10.1039/c6fo01585a

20. Jiang, G., Li, Z., Liu, F., Ellsworth, K., Dallas-Yang, Q., Wu, M., Ronan, J., Esau, C., Murphy, C., Szalkowski, D., Bergeron, R., Doebber, T., Zhang, B.B., 2005. Prevention of obesity in mice by antisense oligonucleotide inhibitors of stearoyl-CoA desaturase-1. Journal of Clinical Investigation 115, 1030–1038. https://doi.org/10.1172/JCI200523962

21. Kim, Y.W., Kim, Y.M., Yang, Y.M., Kim, T.H., Hwang, S.J., Lee, J.R., Kim, S.C., Kim, S.G., 2010. Inhibition of SREBP-1c-mediated hepatic steatosis and oxidative stress by sauchinone, an AMPK-activating lignan in Saururus chinensis. Free Radic Biol Med 48, 567–578. https://doi.org/10.1016/j.freeradbiomed.2009.12.006

22. Koshy, S.M., Bobby, Z., Jacob, S.E., Ananthanarayanan, P.H., Sridhar, M.G., Paulose, D.T., 2015. Amla prevents fructose-induced hepatic steatosis in ovariectomized rats: Role of liver FXR and LXRα. Climacteric 18, 299–310. https://doi.org/10.3109/13697137.2014.933408

23. Li, B., Li, J., Hu, S., 2022. Cinnamon could improve hepatic steatosis caused by a high-fat diet via enhancing hepatic beta-oxidation and inhibiting hepatic lipogenesis, oxidative

damage, and inflammation in male rats. J Food Biochem 46. https://doi.org/10.1111/jfbc.14077

24. Li, Y., Xu, S., Mihaylova, M.M., Zheng, B., Hou, X., Jiang, B., Park, O., Luo, Z., Lefai, E., Shyy, J.Y.J., Gao, B., Wierzbicki, M., Verbeuren, T.J., Shaw, R.J., Cohen, R.A., Zang, M., 2011. AMPK phosphorylates and inhibits SREBP activity to attenuate hepatic steatosis and atherosclerosis in diet-induced insulin-resistant mice. Cell Metab 13, 376–388. https://doi.org/10.1016/j.cmet.2011.03.009

25. Liu, Y.T., Chen, H.W., Lii, C.K., Jhuang, J.H., Huang, C.S., Li, M.L., Yao, H.T., 2020. A diterpenoid, 14-deoxy-11, 12-didehydroandrographolide, in andrographis paniculata reduces steatohepatitis and liver injury in mice fed a high-fat and high-cholesterol diet. Nutrients 12. https://doi.org/10.3390/nu12020523

26. Luo, X., Zhang, B., Pan, Y., Gu, J., Tan, R., Gong, P., 2022. Phyllanthus emblica aqueous extract retards hepatic steatosis and fibrosis in NAFLD mice in association with the reshaping of intestinal microecology. Front Pharmacol 13. https://doi.org/10.3389/fphar.2022.893561

27. Mehlem, A., Hagberg, C.E., Muhl, L., Eriksson, U., Falkevall, A., 2013. Imaging of neutral lipids by oil red O for analyzing the metabolic status in health and disease. Nat Protoc 8, 1149–1154. https://doi.org/10.1038/nprot.2013.055

28. Michelotti, G.A., Machado, M. V., Diehl, A.M., 2013. NAFLD, NASH and liver cancer. Nat Rev Gastroenterol Hepatol. https://doi.org/10.1038/nrgastro.2013.183

29. Miyazaki, M., Flowers, M.T., Sampath, H., Chu, K., Otzelberger, C., Liu, X., Ntambi, J.M., 2007. Hepatic Stearoyl-CoA Desaturase-1 Deficiency Protects Mice from Carbohydrate-Induced Adiposity and Hepatic Steatosis. Cell Metab 6, 484–496. https://doi.org/10.1016/j.cmet.2007.10.014

30. Motoyuki, K., Munechika, E., Nobito, H., Masaki, K., Kazuhiro, K., Tsuyoshi, Y., Tatsuya, F., Masayoshi, Y., Ryoko, Y., Naohiko, H., Ryoichi, T., Makoto, N., 2007. Re-evaluation of fatty acid metabolism-related gene expression in nonalcoholic fatty liver disease.

31. Mundi, M.S., Velapati, S., Patel, J., Kellogg, T.A., Abu Dayyeh, B.K., Hurt, R.T., 2020. Evolution of NAFLD and Its Management. Nutrition in Clinical Practice. https://doi.org/10.1002/ncp.10449

32. Nitasha Bhat, G.M., Nayak, N., Vinodraj, K., Chandralekha, N., Mathai, P., Cherian, J., 2015. Comparison of the efficacy of cardamom (Elettaria cardamomum) with pioglitazone on dexamethasone-induced hepatic steatosis, dyslipidemia, and hyperglycemia in albino rats. J Adv Pharm Technol Res 6, 136–140. https://doi.org/10.4103/2231-4040.157981

33. Oh, G.S., Yoon, J., Lee, G.G., Kwak, J.H., Kim, S.W., 2015. The hexane fraction of Cyperus rotundus prevents non-alcoholic fatty liver disease through the inhibition of liver X receptor α-mediated activation of sterol regulatory element binding protein-1c. American Journal of Chinese Medicine 43, 477–494. https://doi.org/10.1142/S0192415X15500305

34. Osborne, T.F., 2000. Sterol regulatory element-binding proteins (SREBPS): Key regulators of nutritional homeostasis and insulin action. Journal of Biological Chemistry. https://doi.org/10.1074/jbc.R000017200

35. Parlati, L., Régnier, M., Guillou, H., Postic, C., 2021. New targets for NAFLD. JHEP Reports. https://doi.org/10.1016/j.jhepr.2021.100346

36. Paton, C.M., Ntambi, J.M., 2009. Biochemical and physiological function of stearoyl-CoA desaturase. Am J Physiol Endocrinol Metab 297, 28–37. https://doi.org/10.1152/ajpendo.90897.2008.-A

37. Pettinelli, P., Videla, L.A., 2011. Up-regulation of PPAR-γ mRNA expression in the liver of obese patients: An additional reinforcing lipogenic mechanism to SREBP-1c induction. Journal of Clinical Endocrinology and Metabolism 96, 1424–1430. https://doi.org/10.1210/jc.2010-2129

38. Qiu, B., Simon, M., 2016. BODIPY 493/503 Staining of Neutral Lipid Droplets for Microscopy and Quantification by Flow Cytometry. Bio Protoc 6. https://doi.org/10.21769/bioprotoc.1912

39. Rahman, M.M., Islam, M.B., Biswas, M., Khurshid Alam, A.H.M., 2015. In vitro antioxidant and free radical scavenging activity of different parts of Tabebuia pallida growing in Bangladesh. BMC Res Notes 8. https://doi.org/10.1186/s13104-015-1618-6

40. Ran, L.S., Wu, Y.Z., Gan, Y.W., Wang, H.L., Wu, L.J., Zheng, C.M., Ming, Y., Xiong, R., Li, Y.L., Lei, S.H., Wang, X., Lao, X.Q., Zhang, H.M., Wang, L., Chen, C., Zhao, C.Y., 2023. Andrographolide ameliorates hepatic steatosis by suppressing FATP2-

mediated fatty acid uptake in mice with nonalcoholic fatty liver disease. J Nat Med 77, 73–86. https://doi.org/10.1007/s11418-022-01647-w

41. Reddy, J.K., Hashimoto, T., 2001. Peroxisomal β-oxidation and peroxisome proliferator-activated receptor α: An Adaptive Metabolic System.

42. Remya, E., Goyal, M., Varsakiya, J., 2020. Efficacy of Sharapunkhadi powder (a polyherbal formulation) and lifestyle modification in the management of nonalcoholic fatty liver disease-A randomized placebo-controlled clinical trial. AYU (An international quarterly journal of research in Ayurveda) 41, 98. https://doi.org/10.4103/ayu.ayu_281_19

43. Sahu, A.K., Upadhyay, A., Bhakuni, H., Attanayake, A.M.H.S., Sharma, P., 2022. Effect of Ayurveda interventions in non-alcoholic grade II fatty liver associated with obesity – A case report. J Ayurveda Integr Med 13. https://doi.org/10.1016/j.jaim.2022.100605

44. Schroeder-Gloeckler, J.M., Rahman, S.M., Janssen, R.C., Qiao, L., Shao, J., Roper, M., Fischer, S.J., Lowe, E., Orlicky, D.J., McManaman, J.L., Palmer, C., Gitomer, W.L., Huang, W., O'Doherty, R.M., Becker, T.C., Klemm, D.J., Jensen, D.R., Pulawa, L.K., Eckel, R.H., Friedman, J.E., 2007. CCAAT/enhancer-binding protein β deletion reduces adiposity, hepatic steatosis, and diabetes in Leprdb/db mice. Journal of Biological Chemistry 282, 15717–15729. https://doi.org/10.1074/jbc.M701329200

45. Schulze, R.J., McNiven, M.A., 2019. Lipid Droplet Formation and Lipophagy in Fatty Liver Disease. Semin Liver Dis 39, 283–290. https://doi.org/10.1055/s-0039-1685524

46. Sharma, V.K., Prateeksha, Gupta, S.C., Singh, B.N., Rao, C. V., Barik, S.K., 2022. Cinnamomum verum-derived bioactives-functionalized gold nanoparticles for prevention of obesity through gut microbiota reshaping. Mater Today Bio 13. https://doi.org/10.1016/j.mtbio.2022.100204

47. Shetty, S.N., Mengi, S., Vaidya, R., Vaidya, A.D.B., 2010. A study of standardized extracts of Picrorhiza kurroa Royle ex Benth in experimental nonalcoholic fatty liver disease. J Ayurveda Integr Med 1, 203–210. https://doi.org/10.4103/0975-9476.72622

48. Toppo, E., Al-Dhabi, N.A., Sankar, C., Kumar, S.N., Buvanesvaragurunathan, K., Darvin, S.S., Stalin, A., Balakrishna, K., Ceasar, S.A., Pandikumar, P., Ignacimuthu, S., Sivasankaran, K., Agastian, P., 2021. Hepatoprotective effect of selected

isoandrographolide derivatives on steatotic HepG2 cells and High Fat Diet fed rats. Eur J Pharmacol 899. https://doi.org/10.1016/j.ejphar.2021.174056

49. Toppo, E., Darvin, S.S., Esakkimuthu, S., Nayak, M.K., Balakrishna, K., Sivasankaran, K., Pandikumar, P., Ignacimuthu, S., Al-Dhabi, N.A., 2017. Effect of two andrographolide derivatives on cellular and rodent models of non-alcoholic fatty liver disease. Biomedicine and Pharmacotherapy 95, 402–411. https://doi.org/10.1016/j.biopha.2017.08.071

50. Tung, Y.T., Huang, C.Z., Lin, J.H., Yen, G.C., 2018. Effect of Phyllanthus emblica L. fruit on methionine and choline-deficiency diet-induced nonalcoholic steatohepatitis. J Food Drug Anal 26, 1245–1252. https://doi.org/10.1016/j.jfda.2017.12.005

51. Vivekanandan, P., Gobianand, K., Priya, S., Vijayalakshmi, P., Karthikeyan, S., 2007. Protective effect of picroliv against hydrazine-induced hyperlipidemia and hepatic steatosis in rats. Drug Chem Toxicol 30, 241–252. https://doi.org/10.1080/01480540701375216

52. Yan, F., Wang, Q., Lu, M., Chen, W., Song, Y., Jing, F., Guan, Y., Wang, L., Lin, Y., Bo, T., Zhang, J., Wang, T., Xin, W., Yu, C., Guan, Q., Zhou, X., Gao, L., Xu, C., Zhao, J., 2014. Thyrotropin increases hepatic triglyceride content through upregulation of SREBP-1c activity. J Hepatol 61, 1358–1364. https://doi.org/10.1016/j.jhep.2014.06.037

53. Yi, S., Chen, K., Sakao, K., Ikenaga, M., Wang, Y., Hou, D.X., 2023. Assessment of Areca Nut Bioactivities in Western Diet-Induced Mice NAFLD Model. Nutrients 15. https://doi.org/10.3390/nu15102403

54. Younossi, Z., Anstee, Q.M., Marietti, M., Hardy, T., Henry, L., Eslam, M., George, J., Bugianesi, E., 2018. Global burden of NAFLD and NASH: Trends, predictions, risk factors and prevention. Nat Rev Gastroenterol Hepatol. https://doi.org/10.1038/nrgastro.2017.109

55. Younossi, Z., Tacke, F., Arrese, M., Chander Sharma, B., Mostafa, I., Bugianesi, E., Wai-Sun Wong, V., Yilmaz, Y., George, J., Fan, J., Vos, M.B., 2019. Global Perspectives on Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis. Hepatology. https://doi.org/10.1002/hep.30251

56. Zhu, X., Bian, H., Wang, L., Sun, X., Xu, X., Yan, H., Xia, M., Chang, X., Lu, Y., Li, Y., Xia, P., Li, X., Gao, X., 2019. Berberine attenuates nonalcoholic hepatic

steatosis through the AMPK-SREBP-1c-SCD1 pathway. Free Radic Biol Med 141, 192–204. https://doi.org/10.1016/j.freeradbiomed.2019.06.019

Table 1. List of plant ingredients present in *Lodhrasavam*

Sl. No.	Scientific name	Sanskrit name	Plant part used
1	*Symplocos cochinchinensis*	*Lodhra*	Stem Bark
2	*Chonemorpha fragrans*	*Murva*	Root
3	*Kaempferia galanga*	*Sati*	Rhizome
4	*Embelia ribes*	*Vella (Vidanga)*	Seed
5	*Rotheca serrata*	*Bharngi*	Roots
6	*Valeriana jatamansi*	*Nata (Tagara)*	Rhizome
7	*Capparis sepiaria L. syn. C. zeylanica L.*	*Nakha (Vyaghranakha)*	Root
8	*Cyperus esculentus*	*Plava (Musta)*	Root
9	*Wrightia antidysenterica*	*Kalinga (Kutaja)*	Seed
10	*Saussurea costus*	*Kushta*	Root
11	*Areca catechu L.*	*Kramuka (Puga)*	Seeds
12	*Callicarpa macrophylla Yahl.*	*Priyangu*	Inflorescence
13	*Aconitum heterophyllum Wall. Ex Royle*	*Ativisha*	Root
14	*Plumbago zeylanica L.*	*Agni (Citraka)*	Root
15	*Citrullus colocynthis*	*Vishaala (Rakta-indravaruni)*	Whole plant
16	*Cinnamomum verum J.Presl .*	*Lavanga (Tvak)*	Stem Bark
17	*Elettaria cardamomum (L.) Maton*	*Ela (Sukshma ela)*	Seeds
18	*Cinnamomum tamala*	*Patra (Tejapatra)*	Leaves
19	*Mesua ferrea L.*	*Nagakesara*	Stamens
20	*Swertia chirayita*	*Bhunimba*	Whole plant

21	*Neopicrorhiza scrophulariiflora*	*Katukurohini*	Root
22	*Cuminum cyminum*	*Yavaani*	Seed
23	*Inula racemosa C.B.Clarke*	*Pushkara*	Root
24	*Cyclea peltata*	*Pathaa*	Root
25	*Piper longum (wild var.)*	*Granthi*	Root
26	*Piper mullesua*	*Cavya*	Stem
27	*Terminalia chebula Retz.*	*Haritaki*	Fruit pulp
28	*Terminalia bellerica Gaertn.*	*Bibhitaka*	Fruit pulp
29	*Phyllanthus Emblica Roxb.*	*Amalaki*	Fruit pulp

Table 2: List of primers

NM ID	Gene	Primer	Primer sequence	Primer length	Tm	Product length
NM_002046.7	GAPDH	FP	GGCCTCCAAGGAGTAAGACC	20	59.9	147
		RP	AGGGGTCTACATGGCAACTG	20	60	
NM_001321096.3	SREBP	FP	TTCTCACCTCCCAGCTCTGT	20	59.99	98
		RP	CCCAAGCTGTAGCTCCTGTC	20	60.01	
NM_004104.5	FASN	FP	TTCCGAGATTCCATCCTACG	20	60.01	100
		RP	TGTCATCAAAGGTGCTCTCG	20	59.98	
NM_005063.5	SCD	FP	TTCTGGGTAGTCCCCTGTTG	20	59.96	115
		RP	ACTCTCTCCCCCGACAATTT	20	59.99	
NM_001876.4	CPTA1	FP	CATCATCACTGGCGTGTACC	20	59.99	81
		RP	TTGGCGTACATCGTTGTCAT	20	60.01	
NM_001185039.2	ACOX1	FP	CGTGAAACCGCTGAGTAACA	20	59.9	139
		RP	TTCAGACTGGTGCCTCACAG	20	60.02	
NM_001354666.3	PAARG	FP	AGGGACCCGAAATATGCTTT	20	56.8	184
		RP	GAGGGATCCGAAGTTTAGGG	20	57.09	