Ablation of oligodendrogenesis in adult mice alters brain microstructure and activity independently of behavioural deficits

Malte S. Kaller[1], Alberto Lazari[1], Yingshi Feng[1], Annette van der Toorn[2], Sebastian Rühling[1,3], Christopher W. Thomas[4], Takahiro Shimizu[5], David Bannerman[6], Vladyslav Vyazovskiy[4,7,8], William D. Richardson[5], Cassandra Sampaio-Baptista[1,9, #], Heidi Johansen-Berg[1, #]

1. Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United Kingdo; 2. Biomedical MR Imaging and Spectroscopy Group, Center for Image Sciences, University Medical Center Utrecht & Utrecht University, Utrecht, The Netherlands; 3. Department of Diagnostic and Interventional Neuroradiology, School of Medicine, Klinikum rechts der Isar, Technical University of Munich, Munich, Germany; 4. Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, United Kingdom; 5. University College London, The Wolfson Institute for Biomedical Research, London, United Kingdom; 6. Department of Experimental Psychology, University of Oxford, Oxford, UK; 7. Sir Jules Thorn Sleep and Circadian Neuroscience Institute, University of Oxford, Oxford, United Kingdom; 8. The Kavli Institute for Nanoscience Discovery, Oxford, United Kingdom; 9. School of Psychology and Neuroscience, University of Glasgow, Glasgow, United Kingdom.

 # Contributed equally

Abstract

Oligodendrocytes continue to differentiate from their precursor cells even in adulthood, a process that can be modulated by neuronal activity and experience. Yet, our understanding of the functional role of adult oligodendrogenesis remains limited. Previous work has indicated that conditional ablation of oligodendrogenesis in adult mice can lead to learning and memory deficits in a range of behavioural tasks. Our results, reported here, have replicated a key finding that learning to run on a complex wheel with unevenly spaced rungs is disrupted by ablation of oligodendrogenesis. However, using ex vivo MRI (MTR and DTI), we also found that ablating oligodendrogenesis by itself alters brain microstructure, independent of behavioural experience. Furthermore, in vivo EEG recording in behaviourally naïve mice with ablated oligodendrogenesis revealed altered brain activity in the form of increased EEG power density across a broad frequency range. Together, our data indicate that disrupting the formation of new oligodendrocytes directly alters brain microstructure and activity. This suggests a role for adult oligodendrogenesis in the maintenance of brain function and indicates that task-independent changes to brain structure and function might contribute to the learning and memory deficits associated with oligodendrogenesis ablation.

Introduction

Myelination of axons plays a critical role in the functioning of the vertebrate nervous system by increasing the transmission speed and energy efficiency of neural processing (Nave, 2010; Salzer and Zalc, 2016). Recent studies have demonstrated that myelin is more dynamic than initially thought. Myelination can be stimulated by artificially exciting neuronal activity (Cullen et al., 2021; Gibson et al., 2014; Mitew et al., 2018), indicating that myelin plasticity, in addition to synaptic modification, might be one way in which experience can shape brain structure and function (Bonetto et al., 2021; Kaller et al., 2017; Xin and Chan, 2020). Indeed, changes in myelination and white matter (WM) microstructure have been consistently associated with learning in humans (Lakhani et al., 2016; Scholz et al., 2009) and rodents (Bacmeister et al., 2022; Sampaio-Baptista et al., 2013). In addition, adaptive myelination is proposed to regulate homeostatic coordination and oscillatory self-organization in local and large-scale brain networks (Dubey et al., 2022; Noori et al., 2020; Pajevic et al., 2022, 2014; Talidou et al., 2022). Thus, deficient myelin plasticity might lead to alterations in myelination and neural network function that impair neurological function (Geraghty et al., 2019; Knowles et al., 2022). Yet, the questions of when, how, and to what extent adaptive myelination contributes to relevant changes in neural circuit function require further investigation.

Myelination can be adaptively modified through the formation of new oligodendrocytes (OLs), the myelin forming glial cells in the central nervous system (Bergles and Richardson, 2016; Foster et al., 2019). OLs continue to differentiate even in adulthood (Hill et al., 2018; Hughes et al., 2018; Rivers et al., 2008; Young et al., 2013) from populations of oligodendrocyte precursor cells (OPCs) that remain abundant, dynamic and widespread in the CNS throughout life (Dawson et al., 2003). Recently, multi-photon imaging studies in live behaving mice have demonstrated continuous formation of oligodendrocytes and changes in oligodendroglial cells dynamics in response to motor learning (Bacmeister et al., 2022; Hill et al., 2018; Hughes et al., 2018). However, the contribution of such continuous and adaptive oligodendrocyte lineage dynamics to the functioning of the nervous system is still not well understood.

Transgenic mouse lines that allow conditional ablation of oligodendrogenesis in early adulthood have provided information about the causal involvement of new oligodendrocyte formation in behaviour. The first use of this transgenic approach revealed that mice with ablated oligodendrogenesis have an impaired ability to learn to run at speed on a complex wheel, suggesting a deficit in motor skills learning (McKenzie et al., 2014; Xiao et al., 2016). Subsequent studies found that interrupting OL generation also leads to deficits in long-term consolidation of spatial and fearful memories (Pan et al., 2020; Steadman et al., 2020). Additionally, reduced opioid reward learning (Yalçın et al., 2022) and impaired training-induced improvements in spatial working memory have also been reported (Shimizu et al., 2023). However, if and how the disruption of oligodendrogenesis in adulthood affects the CNS, independent of experience in a specific task, remains understudied.

The current study set out to further probe our understanding of behavioural, anatomical and physiological

consequences of disrupting the formation of new OLs during adulthood. Our first aim was to replicate key

findings indicating a motor learning deficit in adult mice with ablated oligodendrogenesis on the complex

wheel task (McKenzie et al., 2014; Xiao et al., 2016). Our results replicated the finding that ability to perform

a complex wheel (CW) running task is perturbed within the first hours of testing in animals with disrupted

oligodendrocyte. Our second aim was to study how genetic ablation of OL differentiation affects brain

microstructure and activity in adult mice using *ex vivo* magnetic resonance imaging (MRI) and *in vivo* EEG

recording. We hypothesised that disruption of new oligodendrocyte formation alters the metrics of brain

microstructure sensitive to myelin and, as a consequence, might modify cortical network activity. We

investigated how learning to run the complex wheel affects brain microstructure in the presence and

absence of ongoing OL differentiation, probing the mechanism by which adaptive OL formation might

contribute to skill acquisition. We found alterations in both the microstructure and activity of the brain in

animals with disruptioned oligodendrogenesis, irrespective of animal's experience with the CW task.

Results

Deleting Myrf in OPCs leads to ablation of oligodendrogenesis

In the current study, we used a transgenic mouse model that allowed conditional deletion of myelin

regulatory factor (*Myrf*) in the resident PDGFRα-positive OPC population (*Pdgfra-CreER^{T2}:R26R-YFP: Myrf$^{(fl/fl)}$*;

referred to as 'P-Myrf' from here on; McKenzie et al., 2014). As *Myrf* encodes a transcription factor required

for oligodendrocyte differentiation (Emery et al., 2009), animals with both alleles of *Myrf* conditionally

deleted during adulthood (P-Myrf$^{(-/-)}$) have markedly reduced formation of new oligodendrocytes 6 weeks

after tamoxifen treatment, as compared to control animals with one functional allele (P-Myrf$^{(+/-)}$) (Fig. 1A-D).

Ablation of oligodendrogenesis leads to impaired performance on the complex wheel

Previous work has indicated that such conditional ablation of oligodendrogenesis in adult mice leads to

learning deficits in a range of different behavioural tasks (McKenzie et al., 2014; Pan et al., 2020; Steadman

et al., 2020). We aimed to re-examine the ability of such mice to learn to run on a complex running wheel

with irregularly spaced rungs (McKenzie et al., 2014; Xiao et al., 2016) (Illustrated in Fig. 2A).

A total of 124 animals (P-*Myrf*$^{(+/-)}$; n=59, 28 females; P-*Myrf*$^{(-/-)}$; n=65, 29 females) were tested on the

complex wheel (CW) across several experiments (see Methods). Animals received tamoxifen in early

adulthood (age range: P60-90) and behavioural performance was tested 3-6 weeks later (Fig. 2C).

To investigate whether ablation of *Myrf* in OPCs during adulthood impacts performance on the complex

running wheel, we tested for main effects of genotype, time and interactions on daily average and maximum

speed using a mixed-two way ANOVAs. Sex was included as an additional between-subject factor.

Fig. 1. **Deleting Myrf in OPCs significantly reduced the formation of new oligodendrocytes. (A)** Representative individual image of immunohistochemistry, split into its separate channels. DAPI was used to identify individual cell nuclei, CC1 was used to label mature oligodendrocytes, and YFP marked OPCs at the time of tamoxifen administration (Scale bar 50 um). **(B)** CC1+ and YFP+ cells were quantified and labelled as either (1) an OPC that remained YFP+|CC1-, or (2) a recently matured Oligodendrocyte that was YFP+|CC1+. **(C)** Quantification confirmed a significant reduction of the density of YFP+|CC1+ recently matured oligodendrocytes in the P-Myrf$^{(-/-)}$ animals 6 weeks after tamoxifen administration. **(D)** The ratio between recently matured Oligodendrocytes (YFP+|CC1+) and labelled OPCs (YFP+|CC1-) was significantly lower in the P-Myrf$^{(-/-)}$. Data presented as Mean±95%CI, Welch's two sample t-test used for statistical comparison. *** represents statistical significance of p<0.001. P-Myrf$^{(+/-)}$, n=12; P-Myrf$^{(-/-)}$, n=15.

Supplementary Table 2 contains comprehensive information on the statistical analysis methods utilized and

the corresponding results for each dataset.

A main effect of genotype for average speed (F(1,120)= 10.00, *p*=0.002, Fig. 2D) and maximum speed

((F(1,120)=11.92, p<0.001, Fig. 2E) over 8 days confirmed that overall performance metrics of P-*Myrf*$^{(-/-)}$

animals were lower than for their P-*Myrf*$^{(+/-)}$ siblings. Furthermore, P-*Myrf*$^{(-/-)}$ animals had impaired capacity

to improve their performance with practice over training days (mixed two-way ANOVA, time*genotype

interaction for average daily speed (F(2.74,328.5)=5.56, *p*=0.001, Fig. 2D) and maximum daily speed

F(2.57,308.9)=4.76, *p*=0.005, Fig. 2E). Females outperformed males in terms of running performance (main

effect of sex, average speed: F(1,120)= 13.1, p<0.001, maximum speed: F(1,120)= 17.5, p<0.001, Suppl. Fig.

1G) and spent more time on the complex running wheel (Suppl. Fig. 1H, Mann-Whitney U, *p*=0.014), yet no

statistically significant interaction between animals' sex and genotype was detected in any ANOVA test

presented. Running ability on a normal running wheel and general motor skills, as tested by RotaRod and

Balance Beam, were not impaired in the P-Myrf$^{(-/-)}$ animals (Suppl. Fig. 1A-F), indicating that performance differences between genotypes did not generalize to other running tasks or tests of motor skill.

Fig. 2. **Preventing differentiation of new OL leads to impaired performance on the complex wheel. (A)** *Complex running wheel, with an irregular rung pattern displayed beside the wheel.* **(B)** *Animals bred on a PDGF-Rα-Cre-ER^{T2} back-*

matic Illustration of experimental design. Between 3-6 weeks after tamoxifen treatment, animals were given free access
to the CW for 8 days. (D) Mean average speed and (E) maximum speed on the complex wheel of P-Myrf$^{(-/-)}$ was lower
across 8 days. A statistically significant effect of Genotype and for Genotype*time interaction was detected for both
speed metrics. (F, G) Raincloud plots of Average (F) and Maximum (G) speed across the 8 days of running. (H) Average
running speed on the CW during the first 4 hours. P-Myrf$^{(-/-)}$ had a significantly lower speed improvement over the first 4
hours (Time*Genotype interaction). (I) Time spent on the wheel within the first 4 hours showed large variability and was
lower within the P-Myrf$^{(-/-)}$ group (Mann-Whitney U, p=0.013). (J) P-Myrf$^{(-/-)}$ spent less time running on the complex
wheel than their P-Myrf$^{(+/-)}$ siblings over 8 days of testing. (K) Statistically significant log relationship between the aver-
age speed of animals on day 8 of testing and the time animals spend on the wheel during the 8 days. (L) Average run-
ning speed plotted against the first hour of time animals spent on the wheel. (M) Average speed within 10 mins time
period of wheel running after the first hour on the wheel. (N) Average running speed plotted against the time animals
spend on the wheel for the first 5 hours (P-Myrf$^{(+/-)}$; n=58 ; P-Myrf$^{(-/-)}$; n=64). (O) Average speed within 1 hour of wheel
running after 5 hours on the wheel. Data presented as Mean±95%CI or boxplots. Mixed-ANOVA (Main Factors: Geno-
type, Sex; Factor: Time) used for statistical comparison of performance over time comparison. Mann-Whitney U test
used for statistical comparison of two groups. Asterix indicate statistical significance (*p<0.05, **p<0.01, ***p<0.001).

The average running speed (Fig. 2F) and maximum running speed (Fig. 2G) across the 8 days of training was

lower in P-*Myrf*$^{(-/-)}$ (Mann-Whitney U: average speed, Fig. 2F, p=0.003; maximum speed, Fig. 2G, p=0.002),

further indicating a deficit in performance on the CW in animals with ablation of *Myrf* in OPCs during

adulthood. However, the distribution of performance values was not equal between genotype groups, with

higher variability and a bimodally shaped distribution especially prominent in the group of P-*Myrf*$^{(-/-)}$ animals

(Fig. 2F & G), a feature that was also reported by McKenzie et al. (2014).

Furthermore, the observed deficit on the CW was already detectable within the first 4 hours of exposure to

the wheel (Mixed ANOVA: Genotype*Time $F_{(4.83, 579.6)}$ = 2.63, p=0.025, Fig. 2H), replicating an early deficit

in complex wheel learning previously reported by Xiao et al (2016).

P-Myrf$^{(-/-)}$ animals spend less time running on the complex wheel

As running on the wheel was voluntary and animals were given unrestricted access to the wheels, the time

individual animals spent on the running wheel varied significantly. Indeed, within the first 4 hours of

exposure to the wheel, the time animals spent rotating the wheel ranged from about 5 minutes to over 2

hours (Fig. 2I). Interestingly, P-*Myrf*$^{(-/-)}$ animals spent significantly less time on the wheel within this 4 hour

period (Fig. 2I, Mann-Whitney U, p=0.013). A similar trend was observed over the 8 days of testing. Time

running on the wheel ranged from around 4 hours to about 64 hours (Fig. 2H) and P-*Myrf*$^{(-/-)}$ animals spent

less time running on the complex running wheel than P-*Myrf*$^{(-/+)}$ (Mann-Whitney U: p=0.014, Fig. 2H, also see

Suppl. Fig. 1I). Similar to other performance metrics across 8 days (Fig. 2F & G), the time animals spent on

the wheel had a wider and bimodal distribution in the P-*Myrf*$^{(-/-)}$ group (Fig. 2H).

Importantly, we found a statistically significant log-positive relationship between the time animals spent

running on the wheel and their average running speed on day 8 of testing (Fig. 2I, Spearman's rank

correlation, across genotype, rho = 0.800, p<0.001; P-*Myrf*$^{(+/-)}$, rho=0.627, p<0.001; P-*Myrf*$^{(-/-)}$, rho=0.871

p<0.001). This suggests that when considering running speed metrics over chronological time periods (e.g.

days, Fig. 2D & E) it might be important to account for variation in the time animals spent running on the

wheel, and therefore inter-individual variation in the time available for learning the task.

Deficit on complex running wheel detected within the first hours of running

One way to account for such variation is to compare running performance of animals against the time they

spent on the wheel, rather than chronological time. Hence, we analysed average running speed on the

complex wheel for the first hour animals spent running on the wheel, regardless of when that running

happened (Fig. 2 L & M). Using a mixed two-way ANOVA (main factors: genotype, sex; within subject factor:

time(10 min intervals)), we found a significant effect of time ($F_{(2.21, 265.3)}= 88.0$, $p<0.001$) and a significant

time*genotype interaction for average daily speed ($F_{(2.21, 265.3)}= 9.06$, $p<0.001$, Fig. 2J), indicating that P-

$Myrf^{(-/-)}$ animals showed less improvement in performance with practice over the first hour of running. The

average running speed achieved in a 10-minute interval after 1 hour on the wheel was significantly lower in

P-$Myrf^{(-/-)}$ animals (Fig. 2K, Mann-Whitney U: $p= 0.004$).

As this deficit is prominent within the first hour spent running on the complex wheel, we wondered how the

genotype differences changed over subsequent time animals spent on the wheel. Comparing performance

across the first 5 hours running on the wheel, P-$Myrf^{(-/-)}$ animals had a persistent lower average running

speed than P-$Myrf^{(+/-)}$ controls (Fig 2 L-O). A mixed two-way ANOVA (main factors: genotype, sex; within

subject factor: time) confirms a statistically significant effect of time (average speed: $F_{(2.2,259.9)}= 53.8$,

$p<0.001$) and for genotype (average speed: $F_{(1,118)}= 4.99$, $p=0.027$, Fig. 2N), yet not for an interaction

between time spent on the wheel and the animals' genotype that might indicate a deficit in learning ability

over this time period (average speed: $F_{(2.2,259.3)}= 1.22$, $p=0.29$). The running speeds in the 5th hour of

running on the wheel was different between genotypes (average speed: Mann-Whitney U: $p= 0.031$, Fig.

2O). This suggests the group differences that became apparent within the first hour on the wheel (Fig.

2L&M) persisted over subsequent hours of running.

Overall, these results replicate the main behavioural results obtained by McKenzie et al (2014) and Xiao et al

(2016), indicating an early deficit in learning on the CW in animals with ablation of *Myrf* in OPCs during

adulthood. However, unlike McKenzie et al. (2014), we found wide variation in the amount of time animals

spent engaging with the task, a factor that is important to take into account when interpreting behavioural

changes. Taking this variation into account by considering time spent on wheel rather than chronological

time, we found that ablation of oligodendrogenesis results in deficits in performance improvement over the

first hour of experience on the wheel. These deficits are maintained, but not exacerbated, over subsequent

hours of experience.

Probing task demands of the complex wheel task

To better understand the task demands of the wheel running task, we tested wild type (WT) mice on

different variants of the wheel. Switching to a complex wheel after running on the normal wheel leads to a

partial reduction in running speed in WT mice (Suppl. Fig. 5), that recovers after a day or two of further

training, indicating the task demand specific to the CW is mainly probed in the first couple of days of

exposure to the wheel.

To test whether mice learn a specific sequence of movements to master the complex wheel, we tested the

effects of changing the rung sequence after exposure to one variant of the complex wheel. This does not

lead to reduction in running speed in WT mice (Suppl. Fig. 6), suggesting that mice learn a general strategy

for running on irregularly spaced rungs, rather than a specific sequence of movements.

Taken together, these behavioural findings in WT mice suggest that learning to run at speed on a complex

wheel primarily involves adaptation to the irregular spaced rung positions, particularly in the early days of

testing.

Ablation of oligodendrogenesis alters brain microstructure independently of wheel running

Previous work has reported alterations in white matter microstructure, measured with MRI techniques, in

response to motor learning (Sampaio-Baptista et al., 2020, 2013). However, the underlying mechanism is

unclear, as these changes could relate to remodelling of pre-existing myelin and/or recruitment of new

oligodendrocytes. Furthermore, the contribution of new oligodendrocytes to MRI metrics has not been

directly tested.

In the current study we hypothesised that disruption of oligodendrogenesis in early adulthood affects brain

microstructure, as it interferes with ongoing cellular dynamics and *de novo* myelination. Further, we

hypothesised that learning to run on the CW affects brain microstructure. Finally, given evidence that

adaptive oligodendrogenesis contributes to learning to run on the CW, we hypothesised that effects of CW

running on brain microstructure should differ between knock-outs and controls.

We employed *ex vivo* MRI (see Methods) to compare P-Myrf$^{(-/-)}$ (n=26) and P-Myrf$^{(-/+)}$ (n=25) animals that

had free access for 12 days to either the complex wheel (CW, n=26) or to a fixed immovable CW in which

running was not possible (from here on referred to as 'fixed wheel', FW, n=25, Fig. 3A). This FW control

condition was used to keep all environmental factors, other than wheel running, similar between groups

(schematic illustration of experimental design, Fig. 3B).

Brain microstructure was assessed *ex vivo* using a 9.4 T horizontal bore MR scanner (Varian, Palo Alto, CA,

USA). Magnetisation transfer imaging and diffusion-weighted imaging were acquired. Three parameters

were extracted (see Suppl. Fig. 2): i) magnetisation transfer ratio (MTR), which allows indirect detection of

water bound to macromolecules, such as lipids and proteins, and is thus sensitive to myelin (Deloire-Grassin

et al. 2000); ii) fractional anisotropy (FA), which describes the anisotropy of diffusion of water molecules;

and iii) mean diffusivity (MD), which describes the rotationally invariant magnitude of water diffusion within

brain tissue. FA and MD are sensitive but

not specific to a number of white matter features, such as myelin, axon calibre, density and organisation,

among others (Lazari and Lipp, 2021; Sampaio-Baptista and Johansen-Berg, 2017; Zatorre et al., 2012).

For a region of interest (ROI) analysis, five brain areas hypothesised to be involved in learning a running

wheel task, or sensitive to exercise on the wheel, were selected: primary (M1) and secondary (M2) motor

cortex, primary sensory cortex (S1), hippocampus (Hipp) and an anterior region of the corpus callosum

(antCC), as illustrated in Fig. 3C. For exploratory purposes, we ran voxel-wise analyses to test for more local

effects in any brain area across the grey matter mask and white matter skeleton (Fig. 3F). Finally, to

investigate global changes, we tested for effects averaged across the whole grey matter (GM) and whole

white matter (WM) skeleton (Fig. 3G,H).

Effects of genotype: First, to investigate the effect of the genetic manipulation on brain microstructure

without considering behavioural experience of wheel running, we tested for effects of genotype in animals

that were only exposed to the fixed wheel (FW, our control condition, n=25). When testing across the four

grey matter regions of interest (ROIs) (M1, M2, S1, Hipp), P-Myrf$^{(-/-)}$(n=13) were found to have significantly

lower MTR, compared to P-Myrf$^{(+/-)}$ (n=12) (mixed ANOVA; main effect of genotype (F(1,17)=7.32, *p*=0.015).

Post hoc tests for individual ROIs (ANOVA, Bonferroni adjusted for number of ROIs) found a significant

reduction in MTR in M1 (Fig. 3D, *p*=0.02) and M2 (Fig. 3D, *p*=0.032). Additionally, for the white matter region

of the anterior corpus callosum (antCC), P-Myrf$^{(-/-)}$ had significantly lower MTR, compared to P-Myrf$^{(+/-)}$ (Fig.

3E, *p*=0.013). No statistically significant changes between groups were observed for diffusion-derived

metrics of MD and FA (Suppl. Fig. 3). No effects were found in the voxel-wise analysis.

Effects of wheel condition: Next, to investigate the effect that running the complex wheel for 12 days may

have on brain microstructure, we tested for effects of wheel condition (fixed vs complex wheel) in P-Myrf$^{(+/-)}$

control animals only (CW:n=13, FW: n=12). No statistically significant difference between the two running

wheel groups were detected in MTR, FA or MD in the ROI analysis (Suppl. Fig 3). A voxel-wise analysis

revealed an increase in MD in the left M1 grey matter region (Suppl. Fig. 4A & B) for P-Myrf$^{(+/-)}$ that ran the

complex wheel.

Fig. 3. *Ablation of oligodendrogenesis alters brain microstructure even in the absence of wheel running. (A) Animals used for ex vivo MRI imaging were exposed to either a complex wheel on which they can run, or an immovable complex wheel (Fixed Wheel, FW). (B) Illustration of experimental design. (C) Coronal and sagittal section of MTR image illustrating Regions of Interest (ROI) masks used for the ROI analysis. (D) For animals only exposed to the Fixed Wheel (FW, P-Myrf$^{(+/-)}$, n=12, , P-Myrf$^{(-/-)}$, n=13) , ROI-specific mean MTR revealed that the ablation of oligodenrogenesis in the absence of wheel running lead to statistically significant changes in brain microstructure in the Primary motor cortex (M1), Secondary motor cortex (M2) and the (E) anterior corpus callosum (antCC). (F) Mask for the Grey Matter (GM, red) and White Matter skeleton (WM, yellow) used for analysis. (G-H) MTR was reduced in (G) GM and the (H) WM skeleton for P-Myrf$^{(-/-)}$ animals compared to control animals, regardless of wheel condition (FW, P-Myrf$^{(+/-)}$, n=25, , P-Myrf$^{(-/-)}$, n=25). Diffusion-weighted parameters, such as FA and MD, did not reveal any statistically significant differences. See Suppl. Fig. 3 for more details. Data presented as Mean±95%CI or boxplots. M1:Primary Motor Cortex, M2:Secondary Motor Cortex, S1: Primary Somatosensory Cortex, antCC: Anterior Corpus Callosum. Asterix indicate statistical significance (*p<0.05).*

Interactions between genotype and wheel condition: Finally, to investigate potential interactions between

genetic manipulation and behavioural experience, we performed an ANOVA with genotype and wheel (CW &

FW) as factors on all samples. Similar to effects found when considering the FW groups alone, significant

main effects of genotype on MTR were found within specific grey matter and white matter ROIs (M1, M2, S1,

antCC; Supp. Fig. 3A-E), as well as across the whole grey matter mask (Fig. 3G; $F_{(1, 34)}=7.77$, *p*=0.019) and

the white matter skeleton (Fig. 3H; $F_{(1,34)}=4.73$, 0.037). , with P-Myrf$^{(-/-)}$ animals having significantly lower

MTR. However, no main effect of wheel or interactions between genotype and wheel on MTR were found.

No main effects or interactions were detected in FA or MD in the ROI analysis (Suppl. Fig. 3F-N). The voxel

wise analysis detected small clusters of changes in MTR between genotypes in cortical and midbrain regions

(Suppl. Fig. 4C), yet no interaction between genotype and wheel was found for any voxel-wise analyses.

Correlations between MRI and behavioural performance: A voxel-wise analysis across all animals in the CW

condition (n=26) tested for correlations between brain metrics and performance metrics (maximum speed &

total distance run over 8 days). No relationship was found for MTR or FA. We found a negative correlation

between distance run and MD in broad areas in cortical grey matter and basal ganglia, independent of

genotype (Suppl. Fig. 3D&E). Mice that ran longer distances had lower MD, indicating that water in these

areas is more restricted (regardless of direction), potentially indicating higher brain tissue density.

Taken together, these results indicate that the ablation of oligodendrogenesis in young adult mice for 5-6

weeks led to decreases in the myelin sensitive metric MTR, across grey matter and white matter regions.

However, we found no interactions between wheel running and genotype, suggesting no evidence for

changes in brain microstructure in response to wheel running that depend on oligodendrogenesis.

Ablation of oligodendrogenesis changes oscillatory brain activity in Myrf-cKO mice in vivo

Previous work has indicated that even subtle deficits in myelination can lead to altered neuronal activity and

synchronicity (Dubey et al., 2022; Gould et al., 2018; Kato et al., 2020), while modelling approaches indicate

a role of activity-dependent myelination in oscillatory self-organization and homeostatic coordination of

brain networks (Noori et al., 2020; Talidou et al., 2022).

As our histological and MRI data suggests altered myelination in P-Myrf$^{(-/-)}$ mice, we tested oscillatory brain

activity using cortical EEG recordings in freely-behaving P-Myrf mice, not exposed to running wheels. In each

animal, the electrodes were implanted in two cortical regions (see Fig. 4A) at the age of P110-120, 5-6 weeks

after tamoxifen treatment (Fig. 4B). Mice were individually housed in standard conditions. Data were

recorded over a 2-day period, 1-2 weeks after surgery (P-Myrf[(+/-)]: n=3, P-Myrf[(-/-)]: n=4) during spontaneous

sleep and waking in undisturbed animals.

Spectral analysis revealed a significant increase in EEG spectral power across broad frequency bands in P-

Myrf[(-/-)] animals in the frontal and occipital region during wakefulness (Fig. 4C-E), when compared to P-

Myrf[(+/-)] animals. Using a Mixed ANOVA (Main effect: genotype, within-subject factor: electrode location),

the genotype difference in total EEG power density in 1-30 Hz frequency range was statistically significant

across both cortical derivations during wakefulness (main effect for genotype: $F_{(1,5)}=10.4$, $p=0.023$). This

change in total spectral power induced by the conditional deletion of *Myrf in OPCs* was also observed during

other vigilance states such as REM sleep (Fig. 4F, Suppl. FigA&B; $F_{(1,5)}=12.7$, $p=0.016$) and nonREM sleep

(Fig. 4G, Suppl. Fig. C&D,$F_{(1,5)}=11.4$, $p=0.02$). The fraction of time animals spent in different vigilance states

did not differ between genotypes (Fig. 4H). The peak EEG power density during wake and REM sleep (0.25

hz resolution) in the Theta range (4-12 hz) was significantly different between genotype groups ($F_{(1,5)}=7.4$,

$p=0.041$, Suppl. Fig. 5E), yet the frequency of the peak was not (Suppl. Fig 5F). Altogether, these results

provide evidence for changes in brain electrophysiology induced by the ablation of oligodendrogenesis for 7-

8 weeks prior to EEG recording in freely moving and behaviourally naïve mice.

In summary, we replicated the previously reported behavioural phenotype of P-Myrf[(-/-)] showing reduced

capacity to improve performance over time on the complex wheel, which is prominent within the first hours

of running. We additionally found a high degree of variation in engagement with the wheel, which should be

taken into account when interpreting results. Further, our results suggest that ablation of

oligodendrogenesis alters brain microstructure, as detected by a reduction in MTR, and brain activity, as

indicated by increase in EEG mean spectral power, independently of behavioural experience on the complex

running wheel. This suggests ablation of oligodendrogenesis in P-Myrf[(-/-)] can itself lead to physiological and

structural alterations in the brain.

Fig. 4. **Ablation of oligodendrogenesis alters brain activity in the absence wheel running. (A)** *Schematic illustration of the location of screws implanted for in vivo EEG recording.* **(B)** *Schematic illustration of experimental design.* **(C-D)** *Mean group Power Spectral Density (PSD) by Frequency during wakefulness in* **(C)** *frontal channel and* **(D)** *occipital channels. Data presented as Mean±SED* **(E)** *The mean PSD across the 1-30 hz frequency bands was lower for animals with ablated oligodendrogenesis in the frontal and occipital channel during wakefulness. This genotype difference was found to be statistically significant across frontal and occipital channels (Mixed ANOVA) during wakefulness.* **(F-G)** *The effect was not specific to wakefulness, as statistically significant differences in mean PSD were also observed during* **(F)** *REM sleep and* **(G)** *non-REM sleep.* **(H)** *The fraction of time animals spent in different vigilance states. Recordings were undertaken in seven male adult mice (P-Myrf(-/-): n=4; P-Myrf(+/-): n=3). Data presented as Mean±SD. Asterix indicate statistical significance (*p<0.05).*

Discussion

This study investigated the consequences of disrupting OL differentiation during adulthood (Fig. 1) on behaviour (Fig. 2), brain structure (Fig. 3) and brain activity (Fig. 4). Our results replicated the key finding that ability to perform a complex wheel (CW) running task is disrupted by ablation of oligodendrogenesis (Fig. 2, McKenzie et al., 2014; Xiao et al., 2016). These group differences on the complex wheel were present within hours of running on the wheel. Additionally, using *ex vivo* MRI (Fig. 3) and *in vivo* EEG (Fig. 4), we found that the ablation of oligodendrogenesis altered brain microstructure and activity independently of the behavioural experience of running the complex wheel. Taken together, our findings indicate that inhibiting the formation of new oligodendrocytes in adult mice not only led to a decline in performance on the complex wheel but also brought about alterations in both the microstructure and activity of the brain, irrespective of animals' experience with the task. Whether and how these task-independent changes in brain microstructure and function contributed to the observed behavioural phenotype requires further investigation.

Replication of behavioural deficit on the complex wheel

We found large variability in the time animals spent engaging with the complex wheel task, which was in turn related to their overall performance (Fig. 2I). Accounting for this variability by analysing performance on the CW as a function of time animals spent on the CW (Fig. 2L-O) revealed that P-Myrf[(-/-)] animal's deficit on the wheel arises within the first hour of running (Fig. 2L-<).

However, it is difficult to clearly define the underlying deficit in P-Myrf[(-/-)] mice that leads to the observed deficit on the CW, as the task demands of the CW are multifaceted. . Running speed improvements over several days are observed on both the normal and complex wheel and can therefore reflect generic aspects of wheel exposure as well as specific demands of the complex wheel (Suppl. Fig 1D, Suppl. Fig. 6). Switching from a normal wheel to a complex wheel leads to a temporary reduction in running performance in WT mice (Suppl. Fig. 6, Hibbits et al., 2009), indicating there is some specific task demand of the irregular spaced rungs. Yet, changing the rung sequence does not affect performance, indicating that mice adapt a strategy to cope with irregular spaced rungs in general and do not learn a specific sequence (Suppl. Fig 6). Together, this indicates that mice need to find a new strategy (e.g. adapt their gait pattern, McKenzie et al., 2014) to cope with irregular rungs and be able to run and improve their performance on the CW.

A deficit within the first hour running the complex wheel suggests that a key task requirement impaired in P-Myrf[(-/-)] animals is probed by early attempts to perform the CW task, rather than by learning processes evolving over several hours or days. Although the CW has been used as a test of subtle deficits in complex motor *execution* (Hibbits et al., 2009; Liebetanz et al., 2007; Schalomon and Wahlsten, 2002), the fact that P-Myrf[(-/-)] mice already trained on the CW prior to tamoxifen treatment have no deficits when re-exposed to

the CW (McKenzie et a l., 2014), together with the absence of any execution deficit on other motor tasks

(Suppl. Fig. 1A-F, McKenzie et al., 2014), suggests that impairment that we and others have found in CW

performance reflects impairment in the acquisition of a new skill rather than skill execution or general

motor performance.

Task-independent changes to brain structure and function might contribute behavioural deficit

How an inability to form new oligodendrocytes can cause the observed learning phenotype of P-Myrf[(-/-)] mice

within just a few hours is unclear. Previous studies suggest that rapid production of *Enpp6*-expressing

immature oligodendrocytes can be observed within the first hours of exposure to the complex wheel,

potentially contributing to circuit functioning and neuronal metabolism (Xiao et al., 2016). Yet, our

understanding of how new oligodendrocytes contribute to skill learning and memory formation remains

incomplete, with apparently different timescales of involvement found with different learning paradigms.

For example, while early learning deficits have been found with complex wheel running, studies investigating

spatial learning and fear conditioning in mice with a conditional knockout of *Myrf* in OPCs found no deficit in

early memory formation, but a deficit of memory consolidation over days or weeks (Pan et al., 2020;

Steadman et al., 2020). Different aspects of myelin plasticity might occur over distinct timescales. For

example, a study investigating motor learning in the dextrous reaching task found modulation of existing

myelin in upper layers of M1 predominant in early stages of learning, while addition of new myelin sheaths

was more relevant post learning (Bacmeister et al., 2022).

An alternative explanation for the behavioural deficit is that the disruption of oligodendrogenesis for weeks

prior to behaviour testing in P-Myrf[(-/-)] caused subtle changes in brain microstructure (Fig. 3) and activity (Fig.

4) which then impacted on capacity to learn a new skill. New oligodendrocytes (OLs) continue to

differentiate throughout early adulthood (Hill et al., 2018; Hughes et al., 2018; Rivers et al., 2008; Young et

al., 2013) and interfering with such ongoing cellular dynamics can have unintended consequences, as

suggested by the current study (Fig. 3 & Fig. 4). Activity-regulated myelination, including oligodendrogenesis,

is proposed to play a role in homeostatic coordination and oscillatory self-organization in local and large-

scale brain networks (Dubey et al., 2022; Noori et al., 2020; Talidou et al., 2022) and mild impairments in

myelination have been linked to decreased motor learning by increasing asynchrony and spontaneity in

neural activity (Kato et al., 2020). Hence, it is possible that brain differences between genotype groups at the

start of behavioural testing, as well as a lack of activity-dependent up-regulation of oligodendrogenesis

triggered by exposure to the CW, might have combined to contribute to the observed behavioural

phenotype.

Changes in brain microstructure due to Myrf knock-out

When investigating the consequences of genetic ablation of OL differentiation on brain microstructure using

ex vivo MRI, we found significantly lower MTR in multiple brain regions (Fig. 3). We hypothesised that

disruption of oligodendrogenesis will affect brain microstructure, as it disrupts ongoing cellular dynamics

and myelin formation. As MTR is related to myelin content of tissue (Lazari and Lipp, 2021), lower MTR may

reflect reduction in myelination caused by the significant reduction in new oligodendrocyte formation (Fig.

1) induced for ~5-6 weeks prior to *ex vivo* MRI. However, estimating how much myelin content was added

during the timeframe of interest is difficult and requires further histological investigation. Metrics derived

from diffusion-weighted imaging (FA, MD), that are sensitive to tissue microstructure and less sensitive or

specific to myelin, were not significantly affected by the deletion of *Myrf* (Suppl. Fig. 3). The finding that

changes in MTR were widespread across WM and GM regions is consistent with the global ablation of OL

differentiation across the whole CNS.

We found no evidence for task-dependent changes in brain microstructure related to myelination when

investigating how motor learning on the complex wheel affected brain microstructure (in comparison to a

fixed wheel control condition). Specifically, we found no effect of complex wheel running on the myelin-

sensitive metric MTR (Fig. 3, Suppl. Fig. 3), regardless of whether animals had disrupted oligodendrogenesis.

We had hypothesised that motor learning should lead to changes in myelin sensitive metrics of brain

microstructure if associated with increase in myelination, as previously reported in the context of the skilled

reaching motor learning tasks in rats (Sampaio-Baptista et al., 2013). However, there are several important

methodological differences between these studies, including species (rat vs mouse) and training task. The

reaching task involves the development of fine motor movements in comparison to the gross motor

movements necessary to perform the CW task. Furthermore, MRI techniques employed here can only

measure cellular effects very indirectly and are less sensitive to small effects compared to histological

approaches employed by others, which showed increased oligodendrocytes differentiation in motor cortex

and subcortical WM due to CW running (McKenzie et al., 2014; Xiao et al., 2016). In addition, the fixed wheel

control condition we employed might have offered more environmental enrichment in comparison to home

cage controls that are typically used. Furthermore, it is possible that at the timepoint of our measurement,

immediately after 12 days of training on the complex wheel, experience-induced changes in myelination

might not have been prominent enough to cause detectable changes in MTR. Together, we did not find

evidence for learning-induced changes in myelin sensitive MRI measures of brain microstructure after

learning to run the complex wheel.

Electrophysiological changes cause by Myrf knockout

In addition to genotype differences in microstructure, deletion of *Myrf* in OPCs caused alterations in oscillatory brain activity (Fig. 4), as reflected in higher EEG power density across broad frequency bands. Increases in spectral power of LFPs have been reported in Plp1-KO models with mild myelin deficits (Gould et al., 2018) and cuprizone models of demyelination (Dubey et al., 2022). Associated computational modelling indicated that a decrease in conduction velocity, as a result of reduced myelination, can lead to an increase in spectral power (Gould et al., 2018). Additionally, adaptive myelination, disrupted in P-Myrf$^{(-/-)}$ animals, might regulate homeostatic coordination and oscillatory self-organization in local and large-scale brain networks (Dubey et al., 2022; Noori et al., 2020; Talidou et al., 2022). Hence, it is plausible that reduced myelination and disruption of adaptive myelination, caused by ablated oligodendrogenesis over a 6-7 week period, led to the observed increases in EEG power density. However, identification of the underlying mechanism driving such broad alterations in brain activity requires further investigation.

Summary and conclusion

In summary, the findings presented in this study suggest that ablation of oligodendrogenesis by conditional deletion of Myrf in PDGFa-positive OPCs leads to subtle but detectable physiological and structural differences in the brain. The previously detected, and here replicated, behavioural deficit on the complex running wheel might therefore partly arise as a result of these task-independent brain differences, in addition to impairments in oligodendrogenesis driven by learning.

The current findings raise many questions for future work. For example, the timescale over which these structural and physiological changes develop should be explored as we only sampled effects at a single timepoint. Further, it remains to be tested whether the effects found here hold for other transgenic models of disrupted oligodendrogenesis. While the use of such models has thus far been predominantly to study of effects on learning, future work could investigate whether and how continuous oligodendrogenesis plays a broader role in maintaining brain microstructure and function.

This study suggests that the implications of disrupting oligodendrogenesis in the adult mouse brain extend beyond specific task-related effects and should be taken into account when interpreting behavioural phenotypes and designing future experiments.

Acknowledgements

The authors thank Eveliina Hanski and Luke Baxter for feedback on previous version of the manuscript. The authors also want to thank Piergiorgo Salva and the entire Plasticity Group at the Wellcome Centre for Integrative Neuroimaging for constructive discussions and suggestions during the project.

Materials and methods

Animals

All work carried out conformed to UK Home Office legislation (Scientific Procedures Act 1986).

Mouse breeding and genetic background

Mice used in this study were originally generated in the Richardson laboratory (McKenzie et al., 2014), UCL by crossing $Pdgfra\text{-}CreER^{T2}$ transgenic mice (Rivers et al., 2008) with Cre-conditional reporter mice $Rosa26R\text{-}YFP$ ($R26R\text{-}YFP$) (Srinivas et al., 2001) to generate double-homozygous offspring. Cre and YFP reporter lines in the Richardson Laboratory were maintained separately by crossing (more than 10 generations) with C57B6 females (Charles River; Margate, UK). These $Pdgfra\text{-}CreER^{T2}:Rosa26R\text{-}YFP$ were crossed with $Myrf^{(fl/fl)}$ mice (Emery et al., 2009) to obtain $Pdgfra\text{-}CreER^{T2}: R26R: Myrf^{(+/fl)}$ offspring, which were sibling-mated to obtain $Myrf^{(fl/fl)}$ and $Myrf^{(+/fl)}$ littermates. These animals were finally crossed to generate cohorts of $Pdgfra\text{-}CreER^{T2}: R26R\text{-} YFP: Myrf^{(fl/fl)}$ and $Myrf^{(+/fl)}$ littermates for behavioural testing and further breeding. In the Richardson Laboratory, $Myrf^{(fl/fl)}$ mice were obtained originally (in 2010) on a mixed 129/CBA/C57B6 background.

Five breeding pairs of $Pdgfra\text{-}CreER^{T2}: R26R\text{-} YFP: Myrf^{(fl/fl)}$ and $Pdgfra\text{-}CreER^{T2}: R26R\text{-} YFP: Myrf^{(+/fl)}$ (referred to as P-Myrf $^{(fl/fl)}$and P-Myrf $^{(+/fl)}$) were sent to our laboratory from the Richardson Lab, UCL in 2016 and were used to breed animals for our study by crossing P-Myrf $^{(fl/fl)}$ with P-Myrf $^{(+/fl)}$ mates. Mice carried copies of $Pdgfra\text{-}CreER^{T2}$ on both alleles and were either $R26R\text{-}YFP^{(+/-)}$ or $R26R\text{-}YFP^{(+/+)}$. Two generations of animals were bred before behaviour testing started for the current studies and animals from a total of 4 consecutive generations were used for behaviour testing. Hence, P-Myrf$^{(fl/fl)}$ and P-Myrf $^{(+/fl)}$ were maintained by sibling crossing for 6 generations in our facilities.

For testing the normal and complex wheel running in wild-type mice, twenty C57BL/6JOlaHsd mice (8 males and 8 females) were purchased from Envigo, UK at age P60. The mice were given a one-week period to acclimate before commencing behavioural testing.

Mouse Genotyping

Genotyping for Cre-ER^{T2} (Cre 5'-TCG ATG CAA CGA GTG ATG AG and Cre 3'-TTC GGC TAT ACG TAA CAG GG; Product Size 481bp) and R26R-YFP (R26WTF1-AAA GTC GCT CTG AGT TGT TAT, R26WTR1-GGA GCG GGA GAA ATG GAT ATG, R26KOF1-GCG AAG AGT TTG TCC TCA ACC) was done by PCR amplification of genomic DNA. Myrf $^{(fl/fl)}$ mice were genotyped by PCR amplification of genomic DNA using primers within intron 7 flanking the first lox site (5'-AGGAGTGTTGTGGGAA-GTGG and 5'-CCCAGGCTGAAGATGGAATA), which gives a 281 bp product for the wild-type allele and a 489 bp product for the $loxP$-flanked allele (Emery et al., 2009).

 Tamoxifen administration

Tamoxifen (Sigma T5648) was dissolved at 40 mg/ml in corn oil on a shaker overnight at RT and by

subsequent sonicating at 21°C for one hour. P-Myrf [(+/-)] and P-Myrf [(-/-)] mice were generated by administering

tamoxifen (0.3 mg/g body weight in corn oil) by oral gavage for four consecutive days to *Pdgfra-*

CreER^{T2}:R26R- YFP:Myrf$^{(flox/flox)}$ and *Pdgfra-CreER^{T2}:R26R- YFP:Myrf$^{(+/flox)}$*. Animals were left for at least 2

weeks before experimental procedures. During, and for a week after tamoxifen treatment, nutritious jelly

and mash were provided for all mice to prevent excessive weight loss due to the adverse effects of the drug.

Behaviour testing of mice

Assessment of behavioural phenotype was conducted by the same researcher. Wherever possible, equal

numbers of animals of each sex were included in each experimental group. For experiments for which data

collection was not automated, researchers were blinded to the genotype of the animals during data

collection and analysis. All mice were maintained on a 12-hour artificial light-dark cycle (9:00-21:00). Food

and water were provided *ad libitum.*

Complex running wheel

Cages with running wheels were purchased from Lafayette Neuroscience (Scurry mouse mis-step wheel

Model 80821S). This voluntary running wheel cage comes equipped with a wheel (circumference = 0.389 m)

that allows individual rungs to be removed to create a complex wheel or rung configuration (Liebetanz et al.,

2007; McKenzie et al., 2014). The mice were provided with a "complex wheel" (CW) with 16 out of 38 rungs

removed to create a wheel with an irregular pattern of 22-rungs (Fig. 2A), a configuration that was adapted

from (McKenzie et al., 2014). For behaviour testing, mice were single caged with the running wheel, a paper

house filled with nesting material and wood chippings to cover the floor of the cage. Mice were accustomed

to the cage for 30 mins before being provided access to the running wheel by removing a dividing wall to the

compartment where the wheel was located. Mice were placed in the wheel cages up to 2 hours before the

start of the dark period. Mice ran spontaneously, without artificial reward. Food and water were available *ad*

libitum within the specialised running wheel cages.

Infrared sensors recorded wheel rotation, which allowed computer-controlled recording of wheel rotation as

a function of time for 20 cages simultaneously for 24h a day (The Scurry Activity Monitoring Software,

Lafayette Neuroscience, Model 86165). For data analysis, counts of full wheel rotations were exported into a

spreadsheet, divided into time bins of 10 seconds. Custom R scripts were used extract the following running

parameters from the baseline activity data: 1) distance travelled, which represents the total number of

wheel rotations × wheel circumference. 2) Active intervals, which represent the number of 10 second

intervals in which at least one full wheel rotation was recorded. 3) Average speed, which represents the sum

of the distance run divided by the number of active intervals. 4) Maximum speed, which was calculated by

determining the average speed in the 1% of fastest intervals run. Analysis of performance was conducted for

either chronological time intervals (length of time vector) or for intervals of time that animals spent running

on the wheel (length of vector containing only active intervals on the wheel).

Running speed of animals (distance/time) was conducted using time bins of 10 seconds. McKenzie *et al.*

(2014) calculated the average running speed by quantifying wheel spins during a 1 min time bin. Yet, in the

current study between 35-40% of continuous running bouts were found to be shorter than 1 minute long

(data not shown). That meant that average running speed was often not accurately estimated when using 1

min intervals to calculate speed. Hence, 10 sec bins were chosen for this study.

Complex running wheel and fixed wheel testing for ex vivo MRI experiment

A total of 24 animals were tested on the complex running wheel (P-Myrf[(+/-)]:n=12, 5 females; P-Myrf[(-/-)]:n=12,

4 females) in specific running wheel cages as described above. Additionally, 25 littermates were exposed to a

static fixed complex wheel that was not able to rotate (referred to as fixed wheel, FW) in cages with

otherwise identical set up (P-Myrf[(+/-)]:n=12, 4 females; P-Myrf[(-/-)]:n=13, 5 females). As the fixed wheel had the

same missing rungs pattern that allows animals to pass between rungs, animals could climb and interact

with the wheel as a kind of environmental enrichment yet could not run and turn the wheel. The experiment

consisted of 5 distinct batches of P-Myrf animals bred from the same set of 5 breeding pairs over

consecutive litters. Littermates were semi-randomly assigned to cages equipped with either a complex

wheel (CW) or a fixed wheel (FW) and were tested in the same environment and timeframe. Semi-random

assignment meant that animals were randomly assigned to a wheel type, while attempting to balance the

numbers of genotype, sex and littermates equally between experimental groups.

Accelerating RotaRod

Mice were tested on a standard RotaRod apparatus (Ugo Basile, Italy). Mice were tested for their ability to

stay on a rod 3 cm in dimeter (Suppl. Fig. 1 B). The apparatus allowed simultaneous testing of up to 5 mice,

yet a maximum of 3 mice were tested at the same time. Mice were habituated to the testing room for 30

mins before behaviour testing, by placing their cages into the room. Mice were trained for three trials on the

rotarod at a constant speed (4 rpm), 120 s each trial, 2 days before behaviour testing, at which speed all

mice stayed on the rod for the duration of the trial.

On the day of testing, each mouse was held by the tail and allowed to grasp the rotating rod. If it was still in

place after 10 s, then the rotation of the rod was accelerated by 20 rpm to a maximum speed of 40 rpm. The

time at which each mouse fell off the rod was recorded, starting from the moment of acceleration. If a

mouse managed to stay on the rod for longer than 150 s, the trial was stopped, and a time of 150 s was

recorded. Each mouse was tested 3 times on the same day. Animals were allowed to rest for 10 minutes
between trials.

Balance beam

The beam apparatus consists of 1 meter round smooth wooden beams of 15 mm or 7.5 mm diameter
positioned 50 cm above the table. A soft foam matt was placed beneath the beam to cushion any potential
fall. The beam rested on the edge of a wooden box on one side, at which animals were placed for testing. On
the other side, the beam rested on the edge of the animal's home cage (Suppl. Fig. 1C). The beam was
slightly angled (~10 degrees) so that animals had to walk uphill to reach their home cage. A lamp (with a 60-
watt light bulb) was used to shine light above the start point and serve as an aversive stimulus, and the light
inside the room was slightly dimmed. Mice were habituated to the testing room for 1 hour before behaviour
testing, by placing their cage inside the room. Three days before behaviour assessment on the balance
beam, mice were trained to cross the beam. Training was done by placing the mice on the beam (1.5 cm
diameter) opposite the home cage. During training mice were freely allowed to cross the beam and reach
their home cage. This procedure was repeated 3 times with a 1-min rest time between attempts. Training
was conducted as almost all mice stopped on the beam or tried to turn around in the middle during the first
exposure to the apparatus. Only after learning the location of their home cage during training do the mice
reliably cross the beam without stopping and turning around, which allows reliable and comparable data
collection. All mice successfully crossed the beams during training at least one time.

For testing, mice were placed on the beam and freely allowed to cross the beam. Mice were first tested on
the wider beam (1.5 cm diameter). After a 5-min resting time in the home cage, the beam was exchanged for
the narrow (0.75 cm diameter) beam and the same animal was tested again. All animals successfully crossed
the beams without falling. The apparatus was cleaned with ethanol (70%) before testing another animal.

Testing was recorded using an HD camera placed on the side of the beam opposite the home cage. Videos
were analysed by a blinded experimenter. Foot slips (defined as the foot coming off the top of the beam)
and the number of steps taken to cross the beam were quantified.

Perfusion of animals and tissue processing

Animals were deeply anaesthetized (Euthatal: 80mg/kg, i.p) with sodium pentobarbital and transcardially
perfused with 10 ml PBS, followed by 30 ml paraformaldehyde (4% w/v in 0.1 M phosphate buffer) with the
help of a perfusion pump (Flow rate: 4ml/min). After perfusion, animal skulls were dissected by removing
soft tissue from the outside of the skull (skin, muscle, eyes, etc.) and the lower jaws. Dissected skulls
containing the CNS tissue were stored overnight in 4% PFA at 4°C for 24h. Samples were then washed with
phosphate buffer saline (PBS) before being stored in PBS at 4°C. Samples were sent to the Center for Image

Sciences, UMCU, Utrecht, Netherlands for magnetic resonance imaging (MRI). After MR imaging, tissue samples were sent back to the University of Oxford, cryoprotected with 30% sucrose solution, embedded in OCT and stored at -80°C until processed for histological analysis.

Immunohistochemistry of free-floating brain sections

For Immunohistochemical analysis, coronal cryosections (30 µm) were cut (anterior to posterior) from the first section at which the anterior corpus callosum was visible and continuously connected from one hemisphere to the other. Brain sections were collected and washed in Tris-buffered Saline (TBS; Sigma) and processed as floating sections. All incubations and washes were performed on a shaker at 60 rpm. Sections were blocked in TBS containing 10% Normal Goat Serum (Sigma), 0.5% Triton-X (Sigma) and 1% BSA for 2 hours at room temperature. The primary antibodies used were anti-PDGF-Ra (rabbit, New England Biolabs, 1:400 dilution), anti-APC (clone CC1; mouse, Calbiochem, 1:200 dilution) and/or anti-GFP(YFP) (chicken, Avas, 1:500), which were diluted in TBS containing 10% of the blocking solution (see above) at 4°C. Samples were then incubated with corresponding fluorophore conjugated secondary antibodies (1:500, Alexa FluorO-488 or 546, Thermo Fischer Scientific), diluted in blocking solution (1:10) for 2h at room temperature. Samples were mounted and dried for ~10 mins before being sealed under a glass cover with Vectorshield anti-fade mounting medium (Vectorshield laboratories) and viewed by a Zeiss L subse SM-700 confocal microscope and ZEN software. Cells were counted in low-magnification photo-micrographs (x20 objective) of non-overlapping fields of coronal sections (320mmx320mm each) of the corpus callosum, between the dorsolateral corners of the lateral ventricles (6 fields per section, three sections per animal), as illustrated in Suppl. Fig. 6. The surface area of corpus callosum was calculated for each animal to determine cellular density. Image analysis was performed using Fuji/ImageJ (NIH).

Statistical analysis of histological and behavioural data

Criteria for excluding animals from complex wheel data analysis

Previous experiments reveal that a small subset of healthy untreated animals show unusually low engagement in voluntary wheel running. To prevent such outliners from skewing overall group performance, we set the following pre-determined exclusion criteria for analysis of complex wheel behaviour performance: animals were excluded from the analysis if: 1) they ran less than 1km in 8 Days (P-Myrf$^{(+/-)}$; n=4, females; P-Myrf$^{(-/-)}$; n=1), or 2) ran less than 100 meters per day for at least half of the days recorded (P-Myrf$^{(+/-)}$; n=2, females; P-Myrf$^{(-/-)}$; n=1). In total, that led to the exclusion of 8 animals based on these criteria (P-Myrf$^{(+/-)}$; n=6; P-Myrf$^{(-/-)}$; n=2).

Testing for the effect of covariates on complex wheel running

Before analysing the effect of genotype on behaviour, we assessed whether a number of different covariates

might influence our findings, in order to determine which covariates to include in subsequent analyses. First,

to test for a potential effect of the generations and sex on the performance on the CW, a mixed two-way

ANOVA (within subject factor: days (over 8 days of running); between subject factors: sex, generation) was

used to analyse daily running speed (average or maximum). This analysis revealed that sex, but not

generation, had a significant effect maximum running speeds (sex: $F(1,116)=22.68$, $p<0.001$) on the complex

wheel, with females consistently outperforming males (Suppl. Fig. 1G). Apart from consistently running

longer distances on the CW (total distance in m: female, M±SD=35505±22374; males, M±SD=16413±14426),

females also spent more time on the CW than males over a period of 8 consecutive days (Suppl. Fig. 1H).

While there was an expected difference in body weight between sexes, there was no difference in body

weight between P-Myrf$^{(-/-)}$ and P-Myrf$^{(+/-)}$ animals for either sex at the start of behaviour testing (body

weight in g: male-Myrf$^{(+/)}$, M±SD=24.1±3.4; male-Myrf$^{(-/-)}$, M±SD=24.3±2.4; female-Myrf$^{(+/-)}$,

M±SD=19.7.1±1.4, female-Myrf$^{(-/-)}$, M±SD=19.4±2.3).

The main analysis of behavioural performance on the CW combines data from several experiments. Hence,

the age of tamoxifen treatment (range: P58-93, M±SD=77.11±8.6), the age at behaviour testing (range: P92-

135, M±SD=109.9±7.9) and the days between tamoxifen treatment and behaviour testing (range: P21-43,

M±SD=32.8±7.5) varied between animals, experiments and generations (Suppl. Table 1). However, we found

no statistically significant correlation between the performance of the P-Myrf$^{(-/-)}$ on the CW (average running

speed) and any of the age parameters mentioned above in the compiled data set of all P-Myrf animals. Yet, it

is of note that our experiments were not specifically designed to compare the effect of these age parameters

on running performance.

In summary, of the potential covariates tested, only sex was found to be significantly associated with running

behaviour and so this factor was included as a fixed factor in all subsequent analyses of genotype differences

to control for the induced variability.

Statistical analysis method

Statistical analysis was performed using R (R Core Team, 2021), using he rstatix (0.7.0; Kassambara, 2021)

and the psycho (v0.6.2; Makowski, 2021) packages. The statistical tests used for a particular analysis are

mentioned during the data presentation in the result sections. Normality of the data and assumptions of

homogeneity of variance were tested (e.g. with Levene's Test) before using parametric statistical analysis. If

these assumptions were not met, non-parametric alternative tests were used for statistical analysis, or the

test parameters were adjusted to account for the lack of homogeneity of variances (e.g. using un-pooled

variances and a correction to the degrees of freedom for unpaired t-test) whenever possible.

Performance on the complex wheel was analysed using a mixed two-way ANOVAs. However, sample data

did not always meet the assumption of normality due to a tendency towards bimodal distribution in the

sample population. Despite this we proceeded to apply ANOVA as I) this statistical analysis shows a certain

robustness to type 1 error due to moderate violation of assumptions of normality at the range of sample

sizes used in this study (Arnau et al., 2013; Blanca et al., 2018, 2017), II) no ideal alternative non-parametric

statistical analysis exist for a mixed model ANOVA, and III) the same statistical analysis was used for the data

set that we are attempting to replicate. If the assumption of sphericity (Mauchly's Test) was violated in the

mixed two-way ANOVA, Greenhouse-Geisser procedures were used to estimate epsilon in order to correct

the degrees of freedom of the F-distribution. In this case, adjusted degrees of freedom are presented.

Asterisks are used to indicate statistical significance when appropriate: *=$p<0.05$, **=$p<0.01$ ***=$p<0.001$.

Details of statistical analysis approaches and results for each dataset are provided in Suppl. Table 2

MRI methods

MRI acquisition

Post-mortem high-resolution structural MRI was performed on a 9.4 T horizontal bore MR system (Varian,

Palo Alto, CA, USA) at the Center for Image Sciences, UMCU, Utrecht, Netherlands, equipped with a 6 cm ID

gradient insert with gradients up to 1 T/m. A custom-made solenoid coil with an internal diameter of 2.6 cm

was used for excitation and reception of the MR signal. Three perfusion-fixed brains were inserted with the

skulls intact in a custom-made holder and immersed in non-magnetic oil (Fomblin, Solvay Solexis).

Diffusion weighted imaging (DWI) was acquired with 3D eight shot spin-echo EPI sequence with the

following parameters: 60 diffusion encoding directions; 1 average per image; 5 images with no diffusion

weighting; TR/TE 500/38.6 ms, field of view: 24x20x20 mm; isotropic resolution of 0.125 mm ; δ/Δ 5.5/8.91

ms, diffusion gradient strength (half-sine shape) 73 G/cm, b = 3522 s/mm2; total acquisition time 12.3 hrs.

Magnetization transfer images were also acquired with a 3D eight shot spin-echo EPI sequence with a train

of saturating pulses in front of the image acquisition saturating frequencies 5 kHz, 10 kHz, and 100 kHz off

resonance. Four volumes per frequency offset were acquired (TR/TE 500/28ms; field of view: 24x20x20 mm;

resolution of 0.125 mm isotropic; total time 2.3 hrs).

A total of 71 animals were scanned for *ex vivo* MRI. Unfortunately, a subset of 20 animals were perfused by a

different experimenter using a different perfusion method. Due to complications with the perfusion method

and a significant deviation in measured MRI metrics from the remaining cohort, these 20 samples had to be

excluded from the analysis.

MRI data processing

All data were Fourier transformed and combined using home-written software in MatLab (Mathworks®) from the Dijkhuizen lab, UMCU, Utrecht, Netherlands. These scripts can be provided on request. DWI data were analysed with the FMRIB Diffusion Toolbox. Voxel-wise values of fractional anisotropy (FA) and mean diffusivity (MD) were estimated from the original DWI data using 'dtifit'.

For alignment of the images to the same space, individual brain images were digitally separated from the triplets in which they were acquired. For the optimal registration of images in grey and white matter, two different contrasts were combined (both weighted equally): i) FA and ii) and mean of all diffusion weighted directions. A study specific template of all subject images was then acquired using an automated image registration pipeline as described previously (Lerch et al., 2011, 2008). In short, this approach brings all scans into anatomical alignment in an automated and unbiased fashion using a combination of mni_autoreg tools (Collins et al., 1994) and Advanced Normalization Tools (ANTS) (Avants et al., 2008). FSL (for FMRIB Software Library) tools, version 6.0 (www.fmrib.ox.ac.uk/fsl) were used for linear (FLIRT) and nonlinear transformations (FNIRT) to the study specific template (Scripts can be provided on request). Transformations gained from this approach were applied to the individual modalities of interest (FA, MD, MTR).

White matter structures were analysed using a modified version of Tract-Based Spatial Statistics (TBSS; Smith et al., 2006). The skeleton for TBSS was thresholded at an FA value of 0.28 as to reliably contain major tracts in the mouse brain that can be accurately aligned across individuals. Finally, the FA values of the tract centers (i.e., maximum FA values) were projected onto the skeleton for each mouse brain and fed into statistical analysis. MD and MTR values for the same voxels were also projected into the skeleton.

MRI data analysis

For region of interest **(ROI)** analysis, an aggregate atlas combining 182 individually segmented structures (Dorr et al., 2008; Richards et al., 2011; Steadman et al., 2014; Ullmann et al., 2013) was registered to the unbiased consensus average of the current study (the atlas is available at [http://repo.mouseimag-ing.ca/repo/DSURQE_40micron_nifti/](http://repo.mouseimaging.ca/repo/DSURQE_40micron_nifti/)) and masks for the specific ROI (M1, M2, S1, Hippocampus) were extracted from the atlas. Mean values from each ROI for each subject were extracted for statistical analysis in R studio. Mixed ANOVAs were used for group comparison. where regions of Interest (ROIs) were within subject factors and genotype and/or wheel were between subject factors. As data was collected from 4 consecutive experimental groups, experimental group (n=4) was included as a fixed factor to control for introduced variability. Significant effects in within-subject factor (ROIs) were followed by a post-hoc simple comparison between groups (wheel or genotype), Bonferroni adjusted for number of ROIs. Statistical analysis was performed using R (R Core Team, 2021), using he rstatix (0.7.0; Kassambara, 2021). To check for outliers, mean values for each metric were extracted from the whole brain mask and checked for extreme outliers ('*identify_outliers*',

rstatix package, R). One animal (P-Myrf$^{(-/-)}$, CW) was identified as an extreme outlier and subsequently re-

moved from the ROI analysis. Statistical analysis including the outlier are provided in Suppl Table 2.

For voxel-wise analysis across the whole brain, nonparametric permutation testing with a cluster-forming

threshold of t > 2 and 5000 permutations were used to determine corrected p values. Clusters with a

corrected significance of p < 0.05 were deemed significant. We tested for differences between the genotype

groups (Myrf$^{(-/-)}$ vs Myrf$^{(+/-)}$, wheel type (complex wheel vs fixed wheel) and for a genotype*wheel type

interaction (Scripts and GLMs can be provided on request). We also tested for voxel-wise for correlations

between contrast and wheel performance (maximum speed, total distance).

Finally, to test for global effects, we calculate mean values for each MR metric across all voxels within the

GM mask or the WM skeleton.

EEG methods

Electrophysiology data collection

Chronic electrophysiological recordings were undertaken in seven male adult mice (P-Myrf $^{(-/-)}$: n=4; P-Myrf

$^{(+/-)}$: n=3). All mice received tamoxifen at age P75 and were implanted at ages P110-120, and the reported

recordings were collected at age P130. The animals were surgically implanted with a custom-made

headstage tethered to electrodes for the continuous recording of electroencephalogram (EEG) and

electromyogram (EMG). EEG screw electrodes (Fine Science Tools) were inserted into the skull. 5 electrodes

were implanted in total: above the right and left frontal cortex (primary motor area: anteroposterior +2 mm,

mediolateral +2 mm), above the right and left occipital cortex (primary visual area: anteroposterior +3.5 mm,

mediolateral +2.5 mm), and above the cerebellum (which served as a reference). As per Fisher *et al.*, 2016),

two custom-soldered stainless steel wires were inserted into the right and left nuchal muscles respectively,

for the recording of EMG. A schematic diagram of implantation locations is shown in Fig. 4A.

Immediately before surgery, analgesics were administered (subcutaneous injection of 1–2 mg/kg metacam

and 0.08 mg/kg vetergesic). Isoflurane was used for induction and maintenance of anesthesia throughout

the surgical procedure (4% and 1–2% respectively). After surgery, analgesics were given for at least 3 days

(1–2 mg/kg oral metacam) and animal wellbeing was closely monitored for 1-2 weeks, until the animal

returned to baseline conditions. All procedures were performed under a UK Home Office Project License and

conformed to the Animal Scientific Procedures Act 1986.

Two animals were implanted each day, with all surgeries taking place in the same week. Order of surgery

was randomised to ensure balanced allocation of genotype to time of day of surgery (i.e. morning surgery vs

afternoon surgery) and to time of week surgery (i.e. early in the week or late in the week), to avoid systemic

bias in implantations and in length of time elapsed between implantation and recording.

Electrophysiology data processing

Data acquisition was performed using the Multichannel Neurophysiology Recording System (TDT, Alachua FL,

USA). EEG/EMG data were filtered between 0.1–100 Hz, amplified (PZ5 NeuroDigitizer pre-amplifier, TDT

Alachua FL, USA) and stored on a local computer at a sampling rate of 256.9 Hz. EEG/EMG data were

resampled offline at a sampling rate of 256 Hz. Signal conversion was performed using custom-written

Matlab (The MathWorks Inc, Natick, Massachusetts, USA) scripts and was then transformed into European

Data Format (Fisher et al., 2016). Custom MATLAB scripts were used to estimate the power spectral density

using the multitaper method ("pmtm()" function , Signal Processing Toolbox, Matlab) for 4-s epochs f. A

0.25 Hz resolution was used for plotting of EEG power density across frequency bands. Total power within

the 1-30hz range across the frontal and occipital channel was calculated and compared using ANOVAs for

repeated measures (Within-Subject Factor: Location; Between Subject Factor: Genotype). Peak frequency

was defined as the frequency bin (0.25 Hz resolution) with the highest power. Power within specific

frequency bands was calculated and compared between genotype groups: delta, δ (0.5–3 Hz), theta, θ (4–

12 Hz), beta, β (12.5–25 Hz), and gamma, γ (30–80 Hz). Statistical analysis was performed using R (R Core

Team, 2021), using he rstatix (0.7.0; Kassambara, 2021).

Scoring and analysis of vigilance states

Recordings were quality-checked during acquisition. All data was manually scored by a blinded investigator

(S.R.) for vigilance states, as previously described in Fisher et al. (2016). Briefly, waking was defined based on

a low-amplitude fast-frequency EEG activity and high EMG amplitude, NREM sleep was characterised by the

predominance of slow waves and sleep spindles on the EEG and low EMG tone, while REM sleep was

characterised by strong theta EEG activity, especially in the occipital derivation and low EMG activity. To

validate scoring accuracy, a subset of the data was re-scored by another investigator (A.L.). Scoring quality

was further verified by a third investigator (V.V.). The channel of the left hemisphere was selected for group

comparison, unless a blinded investigator (V.V.) indicated that the quality of the data was significantly better

in the right hemisphere, in which case the right hemisphere channel was used.

References

References

Arnau, J., Bendayan, R., Blanca, M.J., Bono, R., 2013. The effect of skewness and kurtosis on the robustness
of linear mixed models. Behav. Res. Methods 45, 873–879. https://doi.org/10.3758/s13428-012-
0306-x
Bacmeister, C.M., Huang, R., Osso, L.A., Thornton, M.A., Conant, L., Chavez, A., Poleg-Polsky, A., Hughes,
E.G., 2022. Motor Learning Drives Dynamic Patterns of Intermittent Myelination on Learning-
activated Axons. https://doi.org/10.1101/2021.10.13.464319
Bergles, D.E., Richardson, W.D., 2016. Oligodendrocyte Development and Plasticity. Cold Spring Harb.
Perspect. Biol. 8, a020453. https://doi.org/10.1101/cshperspect.a020453
Blanca, M.J., Alarcón, R., Arnau, J., Bono, R., Bendayan, R., 2018. Effect of variance ratio on ANOVA
robustness: Might 1.5 be the limit? Behav. Res. Methods 50, 937–962.
https://doi.org/10.3758/s13428-017-0918-2
Blanca, M.J., Alarcón, R., Arnau, J., Bono, R., Bendayan, R., 2017. Non-normal data: Is ANOVA still a valid
option? Psicothema 29, 552–557. https://doi.org/10.7334/psicothema2016.383
Bonetto, G., Belin, D., Káradóttir, R.T., 2021. Myelin: A gatekeeper of activity-dependent circuit plasticity?
Science 374, eaba6905. https://doi.org/10.1126/science.aba6905
Cullen, C.L., Pepper, R.E., Clutterbuck, M.T., Pitman, K.A., Oorschot, V., Auderset, L., Tang, A.D., Ramm, G.,
Emery, B., Rodger, J., Jolivet, R.B., Young, K.M., 2021. Periaxonal and nodal plasticities modulate
action potential conduction in the adult mouse brain. Cell Rep. 34, 108641.
https://doi.org/10.1016/j.celrep.2020.108641
Dawson, M.R.L., Polito, A., Levine, J.M., Reynolds, R., 2003. NG2-expressing glial progenitor cells: an
abundant and widespread population of cycling cells in the adult rat CNS. Mol. Cell. Neurosci. 24,
476–488. https://doi.org/10.1016/s1044-7431(03)00210-0
Dorr, A.E., Lerch, J.P., Spring, S., Kabani, N., Henkelman, R.M., 2008. High resolution three-dimensional brain
atlas using an average magnetic resonance image of 40 adult C57Bl/6J mice. NeuroImage 42, 60–69.
https://doi.org/10.1016/j.neuroimage.2008.03.037
Dubey, M., Pascual-Garcia, M., Helmes, K., Wever, D.D., Hamada, M.S., Kushner, S.A., Kole, M.H.P., 2022.
Myelination synchronizes cortical oscillations by consolidating parvalbumin-mediated phasic
inhibition. eLife 11, e73827. https://doi.org/10.7554/eLife.73827
Emery, B., Agalliu, D., Cahoy, J.D., Watkins, T.A., Dugas, J.C., Mulinyawe, S.B., Ibrahim, A., Ligon, K.L.,
Rowitch, D.H., Barres, B.A., 2009. Myelin gene regulatory factor is a critical transcriptional regulator
required for CNS myelination. Cell 138, 172–185. https://doi.org/10.1016/j.cell.2009.04.031
Fisher, S.P., Cui, N., McKillop, L.E., Gemignani, J., Bannerman, D.M., Oliver, P.L., Peirson, S.N., Vyazovskiy,
V.V., 2016. Stereotypic wheel running decreases cortical activity in mice. Nat. Commun. 7, 13138.
https://doi.org/10.1038/ncomms13138
Foster, A.Y., Bujalka, H., Emery, B., 2019. Axoglial interactions in myelin plasticity: Evaluating the relationship
between neuronal activity and oligodendrocyte dynamics. Glia 67, 2038–2049.
https://doi.org/10.1002/glia.23629
Geraghty, A.C., Gibson, E.M., Ghanem, R.A., Greene, J.J., Ocampo, A., Goldstein, A.K., Ni, L., Yang, T., Marton,
R.M., Paşca, S.P., Greenberg, M.E., Longo, F.M., Monje, M., 2019. Loss of Adaptive Myelination
Contributes to Methotrexate Chemotherapy-Related Cognitive Impairment. Neuron 103, 250-
265.e8. https://doi.org/10.1016/j.neuron.2019.04.032
Gibson, E.M., Purger, D., Mount, C.W., Goldstein, A.K., Lin, G.L., Wood, L.S., Inema, I., Miller, S.E., Bieri, G.,
Zuchero, J.B., Barres, B.A., Woo, P.J., Vogel, H., Monje, M., 2014. Neuronal activity promotes
oligodendrogenesis and adaptive myelination in the mammalian brain. Science 344, 1252304.
https://doi.org/10.1126/science.1252304
Gould, E.A., Busquet, N., Shepherd, D., Dietz, R.M., Herson, P.S., Simoes de Souza, F.M., Li, A., George, N.M.,
Restrepo, D., Macklin, W.B., 2018. Mild myelin disruption elicits early alteration in behavior and
proliferation in the subventricular zone. eLife 7, e34783. https://doi.org/10.7554/eLife.34783

Hibbits, N., Pannu, R., John Wu, T., Armstrong, R.C., 2009. Cuprizone demyelination of the corpus callosum in
mice correlates with altered social interaction and impaired bilateral sensorimotor coordination.
ASN NEURO 1, e00013. https://doi.org/10.1042/AN20090032
Hill, R.A., Li, A.M., Grutzendler, J., 2018. Lifelong cortical myelin plasticity and age-related degeneration in
the live mammalian brain. Nat. Neurosci. 21, 683–695. https://doi.org/10.1038/s41593-018-0120-6
Hughes, E.G., Orthmann-Murphy, J.L., Langseth, A.J., Bergles, D.E., 2018. Myelin remodeling through
experience-dependent oligodendrogenesis in the adult somatosensory cortex. Nat. Neurosci. 21,
696–706. https://doi.org/10.1038/s41593-018-0121-5
Kaller, M.S., Lazari, A., Blanco-Duque, C., Sampaio-Baptista, C., Johansen-Berg, H., 2017. Myelin plasticity and
behaviour-connecting the dots. Curr. Opin. Neurobiol. 47, 86–92.
https://doi.org/10.1016/j.conb.2017.09.014
Kassambara, A., 2023. rstatix: Pipe-Friendly Framework for Basic Statistical Tests.
Kato, D., Wake, H., Lee, P.R., Tachibana, Y., Ono, R., Sugio, S., Tsuji, Y., Tanaka, Y.H., Tanaka, Y.R., Masamizu,
Y., Hira, R., Moorhouse, A.J., Tamamaki, N., Ikenaka, K., Matsukawa, N., Fields, R.D., Nabekura, J.,
Matsuzaki, M., 2020. Motor learning requires myelination to reduce asynchrony and spontaneity in
neural activity. Glia 68, 193–210. https://doi.org/10.1002/glia.23713
Knowles, J.K., Batra, A., Xu, H., Monje, M., 2022. Adaptive and maladaptive myelination in health and
disease. Nat. Rev. Neurol. 18, 735–746. https://doi.org/10.1038/s41582-022-00737-3
Lakhani, B., Borich, M.R., Jackson, J.N., Wadden, K.P., Peters, S., Villamayor, A., MacKay, A.L., Vavasour, I.M.,
Rauscher, A., Boyd, L.A., 2016. Motor Skill Acquisition Promotes Human Brain Myelin Plasticity.
Neural Plast. 2016, 7526135. https://doi.org/10.1155/2016/7526135
Lazari, A., Lipp, I., 2021. Can MRI measure myelin? Systematic review, qualitative assessment, and meta-
analysis of studies validating microstructural imaging with myelin histology. NeuroImage 230,
117744. https://doi.org/10.1016/j.neuroimage.2021.117744
Liebetanz, D., Baier, P.C., Paulus, W., Meuer, K., Bähr, M., Weishaupt, J.H., 2007. A highly sensitive
automated complex running wheel test to detect latent motor deficits in the mouse MPTP model of
Parkinson's disease. Exp. Neurol. 205, 207–213. https://doi.org/10.1016/j.expneurol.2007.01.030
McKenzie, I.A., Ohayon, D., Li, H., de Faria, J.P., Emery, B., Tohyama, K., Richardson, W.D., 2014. Motor skill
learning requires active central myelination. Science 346, 318–322.
https://doi.org/10.1126/science.1254960
Micheva, K.D., Wolman, D., Mensh, B.D., Pax, E., Buchanan, J., Smith, S.J., Bock, D.D., 2016. A large fraction
of neocortical myelin ensheathes axons of local inhibitory neurons. eLife 5, e15784.
https://doi.org/10.7554/eLife.15784
Mitew, S., Gobius, I., Fenlon, L.R., McDougall, S.J., Hawkes, D., Xing, Y.L., Bujalka, H., Gundlach, A.L., Richards,
L.J., Kilpatrick, T.J., Merson, T.D., Emery, B., 2018. Pharmacogenetic stimulation of neuronal activity
increases myelination in an axon-specific manner. Nat. Commun. 9, 306.
https://doi.org/10.1038/s41467-017-02719-2
Nave, K.-A., 2010. Myelination and support of axonal integrity by glia. Nature 468, 244–252.
https://doi.org/10.1038/nature09614
Noori, R., Park, D., Griffiths, J.D., Bells, S., Frankland, P.W., Mabbott, D., Lefebvre, J., 2020. Activity-
dependent myelination: A glial mechanism of oscillatory self-organization in large-scale brain
networks. Proc. Natl. Acad. Sci. 117, 13227–13237. https://doi.org/10.1073/pnas.1916646117
Pajevic, S., Basser, P.J., Fields, R.D., 2014. Role of myelin plasticity in oscillations and synchrony of neuronal
activity. Neuroscience 276, 135–147. https://doi.org/10.1016/j.neuroscience.2013.11.007
Pajevic, S., Plenz, D., Basser, P.J., Fields, R.D., 2022. Oligodendrocyte-mediated Myelin Plasticity and its role
in Neural Synchronization. https://doi.org/10.1101/2022.07.03.498570
Pan, S., Mayoral, S.R., Choi, H.S., Chan, J.R., Kheirbek, M.A., 2020. Preservation of a remote fear memory
requires new myelin formation. Nat. Neurosci. 23, 487–499. https://doi.org/10.1038/s41593-019-
0582-1
Richards, K., Watson, C., Buckley, R.F., Kurniawan, N.D., Yang, Z., Keller, M.D., Beare, R., Bartlett, P.F., Egan,
G.F., Galloway, G.J., Paxinos, G., Petrou, S., Reutens, D.C., 2011. Segmentation of the mouse

hippocampal formation in magnetic resonance images. NeuroImage 58, 732–740.
https://doi.org/10.1016/j.neuroimage.2011.06.025
Rivers, L.E., Young, K.M., Rizzi, M., Jamen, F., Psachoulia, K., Wade, A., Kessaris, N., Richardson, W.D., 2008.
PDGFRA/NG2 glia generate myelinating oligodendrocytes and piriform projection neurons in adult
mice. Nat. Neurosci. 11, 1392–1401. https://doi.org/10.1038/nn.2220
Salzer, J.L., Zalc, B., 2016. Myelination. Curr. Biol. CB 26, R971–R975.
https://doi.org/10.1016/j.cub.2016.07.074
Sampaio-Baptista, C., Johansen-Berg, H., 2017. White Matter Plasticity in the Adult Brain. Neuron 96, 1239–
1251. https://doi.org/10.1016/j.neuron.2017.11.026
Sampaio-Baptista, C., Khrapitchev, A.A., Foxley, S., Schlagheck, T., Scholz, J., Jbabdi, S., DeLuca, G.C., Miller,
K.L., Taylor, A., Thomas, N., Kleim, J., Sibson, N.R., Bannerman, D., Johansen-Berg, H., 2013. Motor
skill learning induces changes in white matter microstructure and myelination. J. Neurosci. Off. J.
Soc. Neurosci. 33, 19499–19503. https://doi.org/10.1523/JNEUROSCI.3048-13.2013
Sampaio-Baptista, C., Vallès, A., Khrapitchev, A.A., Akkermans, G., Winkler, A.M., Foxley, S., Sibson, N.R.,
Roberts, M., Miller, K., Diamond, M.E., Martens, G.J.M., De Weerd, P., Johansen-Berg, H., 2020.
White matter structure and myelin-related gene expression alterations with experience in adult rats.
Prog. Neurobiol. 187, 101770. https://doi.org/10.1016/j.pneurobio.2020.101770
Schalomon, P.M., Wahlsten, D., 2002. Wheel running behavior is impaired by both surgical section and
genetic absence of the mouse corpus callosum. Brain Res. Bull. 57, 27–33.
https://doi.org/10.1016/S0361-9230(01)00633-5
Scholz, J., Klein, M.C., Behrens, T.E.J., Johansen-Berg, H., 2009. Training induces changes in white-matter
architecture. Nat. Neurosci. 12, 1370–1371. https://doi.org/10.1038/nn.2412
Shimizu, T., Nayar, S.G., Swire, M., Jiang, Y., Grist, M., Bannerman, D.M., Johansen-Berg, H., Ogasawara, K.,
Tohyama, K., Richardson, W.D., 2023. Oligodendrocyte dynamics dictate individual performance
outcomes of working memory training in mice. https://doi.org/10.1101/2023.01.17.524381
Smith, S.M., Jenkinson, M., Johansen-Berg, H., Rueckert, D., Nichols, T.E., Mackay, C.E., Watkins, K.E.,
Ciccarelli, O., Cader, M.Z., Matthews, P.M., Behrens, T.E.J., 2006. Tract-based spatial statistics:
voxelwise analysis of multi-subject diffusion data. NeuroImage 31, 1487–1505.
https://doi.org/10.1016/j.neuroimage.2006.02.024
Srinivas, S., Watanabe, T., Lin, C.-S., William, C.M., Tanabe, Y., Jessell, T.M., Costantini, F., 2001. Cre reporter
strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. BMC Dev. Biol. 1, 4.
https://doi.org/10.1186/1471-213X-1-4
Steadman, P.E., Ellegood, J., Szulc, K.U., Turnbull, D.H., Joyner, A.L., Henkelman, R.M., Lerch, J.P., 2014.
Genetic effects on cerebellar structure across mouse models of autism using a magnetic resonance
imaging atlas. Autism Res. 7, 124–137. https://doi.org/10.1002/aur.1344
Steadman, P.E., Xia, F., Ahmed, M., Mocle, A.J., Penning, A.R.A., Geraghty, A.C., Steenland, H.W., Monje, M.,
Josselyn, S.A., Frankland, P.W., 2020. Disruption of Oligodendrogenesis Impairs Memory
Consolidation in Adult Mice. Neuron 105, 150-164.e6. https://doi.org/10.1016/j.neuron.2019.10.013
Talidou, A., Frankland, P.W., Mabbott, D., Lefebvre, J., 2022. Homeostatic coordination and up-regulation of
neural activity by activity-dependent myelination. Nat. Comput. Sci. 1–12.
https://doi.org/10.1038/s43588-022-00315-z
Ullmann, J.F.P., Watson, C., Janke, A.L., Kurniawan, N.D., Reutens, D.C., 2013. A segmentation protocol and
MRI atlas of the C57BL/6J mouse neocortex. NeuroImage 78, 196–203.
https://doi.org/10.1016/j.neuroimage.2013.04.008
Xiao, L., Ohayon, D., McKenzie, I.A., Sinclair-Wilson, A., Wright, J.L., Fudge, A.D., Emery, B., Li, H., Richardson,
W.D., 2016. Rapid production of new oligodendrocytes is required in the earliest stages of motor-
skill learning. Nat. Neurosci. 19, 1210–1217. https://doi.org/10.1038/nn.4351
Xin, W., Chan, J.R., 2020. Myelin plasticity: sculpting circuits in learning and memory. Nat. Rev. Neurosci. 21,
682–694. https://doi.org/10.1038/s41583-020-00379-8

Yalçın, B., Pomrenze, M.B., Malacon, K., Chau, I.J., Taylor, K.R., Ni, L., Contreras-Esquivel, D., Malenka, R.C.,
Monje, M., 2022. Myelin plasticity in ventral tegmental area is required for opioid reward.
https://doi.org/10.1101/2022.09.01.506263
Young, K.M., Psachoulia, K., Tripathi, R.B., Dunn, S.-J., Cossell, L., Attwell, D., Tohyama, K., Richardson, W.D.,
2013. Oligodendrocyte Dynamics in the Healthy Adult CNS: Evidence for Myelin Remodeling. Neuron
77, 873–885. https://doi.org/10.1016/j.neuron.2013.01.006
Zatorre, R.J., Fields, R.D., Johansen-Berg, H., 2012. Plasticity in gray and white: neuroimaging changes in
brain structure during learning. Nat. Neurosci. 15, 528–536. https://doi.org/10.1038/nn.3045
901

*Suppl. Fig. 1. **Control task for motor behaviour and supplementary data for the complex running wheel data. (A)** To*
test general performance on a running wheel, a cohort of P-Myrf(-/-) and P-Myrf(+/-) animals were tested on a normal
wheel which had none of the rungs removed. To test for general motor ability and coordination, a cohort of animals was
tested on the (B) RotaRod and (C) Balance Beam task. (D) Conditional knockout of Myrf did not affect performance on a
normal running wheel (P-Myrf(+/-); n=17; P-Myrf(-/-); n=18) , nor did it lead to a difference in performance on the (E)
RotaRod or (F) Balance Beam tasks (P-Myrf(+/-); n=16, P-Myrf(-/-); n=25) which may have indicated a more general mo-
tor skill deficit. (G-H) Sex difference in running speed and activity level. (G) Females (n=57) outperformed males (n=67)
on the complex running wheel in terms of running performance (Mixed ANOVA: Main effect of Sex; $F_{(1,120)}= 19.98$,
$p<0.001$). (H) Females also spent more time active on the wheel over 8 days Mann-Whitney U, p=0.014. (K) Active inter-
vals over 8 days (1 hour time bins) for P-Myrf(-/-) and P-Myrf(+/-) animals. While both Genotypes displayed comparable
day/night activity patterns, P-Myrf(-/-) had on average fewer active intervals on the CW during the night when com-
pared to P-Myrf(+/-) (Mixed ANOVA: Main effect of Genotype: $F_{(1,120)}=4.48$ p=036). Data presented as Mean ± 95%CI.
*Data presented as Mean±95%CI or boxplots. Asterix indicate statistical significance (*p<0.05, **p<0.01, ***p<0.001).*

*Suppl. Fig. 2. **Example images for different parameter contrasts extracted from ex vivo MRI for quantification of brain***
***microstructure.** Brain microstructure was assessed using a 9.4 T horizontal bore MR scanner (Varian, Palo Alto, CA, USA)*
*(**A**) Magnetisation transfer ratio (MTR) allows indirect detection of water bound to macromolecules, such as lipids and*
*proteins, and is thus sensitive to myelin (Deloire-Grassin et al. 2000). (**B**) Fractional anisotropy (FA) describes the anisot-*
*ropy of diffusion of water molecules. (**C**) Mean Diffusivity (MD), which describes the rotationally invariant magnitude of*
water diffusion within brain tissue. Images shown are the mean of all individual subject images analysed for this study.

*Suppl. Fig. 3. **Region of Interest (ROI) Analysis of brain microstructure, as measure by MTR, MD and FA. (A-E)** MTR*
values in grey matter ROIs. A Mixed ANOVA revealed a main effect of Genotype in MTR grey matter regions, post-hoc
*pairwise comparison (Bonferroni adjusted) revealed a significant Genotype effect in the **(A)** Primary motor cortex (M1),*
***(C)** Somatosensory Cortex (S1), **(D)** Hippocampus. **(E)** MTR in the anterior Corpus Callosum. **(F-J)** DWI-derived Mean Dif-*
*fusivity (MD) values in ROI, **(K)** whole brain mask and **(L)** white matter skeleton. **(M-N)** DWI-derived Fractional Anisot-*
*ropy (FA) values in **(M)** white matter skeleton and **(N)** anterior corpus callosum. Asterix indicate statistical significance*
(*p<0.05, **p<0.01).

*Suppl. Fig. 4. **Voxel-wise analysis of DWI-derived parameters revealed an effect wheel running in brain microstruc-***
ture. *Results from Voxel-wise analysis of MRI-derived metrics. (**A**) A cluster in the right primary motor cortex (M1) grey*
matter region had higher MD for P-Myrf$^{(-/+)}$ control animals that ran on the Complex Wheel (CW), when compared to P-
*Myrf$^{(-/+)}$ exposed to the fixed wheel (FW). (**B**) Mean MD extracted from cluster depicted in (**A**). (**C**) MTR was lower for P-*
*Myrf$^{(-/-)}$ animals in a few clusters located in cortical regions and the midbrain (superior colliculus). (**D**) MD across a dif-*
fuse area across the grey matter was correlated to the total distance ran on the complex running wheel by animals
*across both genotypes. (**E**) Mean MD extracted from the ROI depicted in (**D**) plotted against the total distance run on the*
complex wheel. Permutation test ('randomise', FSL) used for voxel-wise analysis. Spearman's test used for correlation
analysis.

Suppl. Fig. 5. **Difference in EEG recording between P-Myrf$^{(-/-)}$ and P-Myrf$^{(+/-)}$.** *EEG power density by frequency during rapid eye movement (REM) sleep in (**A**) frontal channel and (**B**) occipital channels. EEG power density by frequency during non-REM sleep (NREM) in (**C**) frontal channel and (**D**) occipital channels. (**E**) Amplitude and (**F**) location of peak theta power per vigilance state (Wake, REM) as measured in occipital channel. While peak amplitude was different between the genotype groups, there was no shift in the frequency of the peak. (**G**) Mean group EEG power density (PSD) per frequency bands, delta, δ (0.5–3 Hz), theta, ϑ (4–12 Hz), beta, β (12.5–25 Hz), and gamma, γ (30-60 Hz). Data presented as Mean±SEM.*

*Suppl. Fig. 6. **The effect of complex wheel sequence on running performance.** Wildtype mice (C58BL/6JOIaHsd, 8 females, 8 males, Age: P84) exposed to a normal running wheel (NW) show signifi-cant improvements in their running speed over 8 days (Main effect of Day: F(3.14, 47.1)=21.8, p<0.001), with females outperforming males (Sex*Day Interaction: F(1, 15)=34.6, p<0.001). Switching to complex running wheel (CW1) leads to a temporary reduction in running speed (Day 8 on NW vs Day 1 on CW1; Main effect of wheel: F(1, 14)=27.0, p<0.001) , indicating a specific task demand of the complex wheel. However, running speed on day 1 on CW1 is significantly higher than on day 1 on NW (Main effect of wheel: F(1, 14)=12.0, p=0.004), indicating some skill in wheel running was retained between NW and CW1. Switching the rung sequence of the complex running wheel (CW2) after 8 days has no effect on run-ning performance compared to last day of running on CW1. Mixed Two-way ANOVA used for statistical analysis. Data presented as Mean±SEM.*

*Suppl. Fig. 7. **Quantification of oligodendrocyte differentiation in the corpus Callosum** Representative image of a brain slices showing the anterior Corpus Callosum region used for quantification of YFP+ positive OPCs and CC1+ mature Oligodendrocytes. Six images were taken from each section (three sections per animal), as illustrated by the white frames in the image.*

*Suppl. Table 1. **Variability of animal's age at important experimental timepoints within different generations of Myrf-cKO animals that ran the complex wheel.*** *Age is presented in days after birth*

Generation	Number of Animals	Age at tamoxifen treatment	Age between tamoxifen treatment and CW	Age at beginning of CW testing	Experiments / Data
G1	17 P-Myrf$^{(-/+)}$ = 7 P-Myrf$^{(-/-)}$ = 10	Range=69-84 Mean=73.5 SD=4.1	Range=31-42 Mean=36.5 SD=5.5	Range=100-126 Mean=109.8 SD=6.8	Complex Wheel (8 days)
G2	19 P-Myrf$^{(-/+)}$ = 11 P-Myrf$^{(-/-)}$ = 8	Range=58-93 Mean=74.6 SD=13.9	Range=34-42 Mean=38.1 SD=4.1	Range=92-135 Mean=112.4 SD=17.1	Complex Wheel (8 days)
G3	39 P-Myrf$^{(-/+)}$ = 15 P-Myrf$^{(-/-)}$ = 24	Range=64-83 Mean=71.4 SD=5.9	Range=30-43 Mean=37.8 SD=5.9	Range=96-114 Mean=109.14 SD=4.8	Complex Wheel (8 days) Followed by: RotaRod & Balance Beam
G4	49 P-Myrf$^{(-/+)}$ = 26 P-Myrf$^{(-/-)}$ = 23	Range=79-88 Mean=83.8 SD=2.3	Range=21-30 Mean=25.4 SD=2.9	Range=102-116 Mean=109.1 SD=2.9	Complex Wheel (12 days) Ex vivo MRI Histology

*Suppl. Table 2: **Comprehensive information on the statistical analysis methods utilized and the corresponding results for each dataset.***

Figure	Data & Methods	Statistical results				
Fig. 1C	Welch's t-test. P-Myrf(+/-), n=12; P-Myrf(-/-), n=15.	YFP+ cells: t = 4.5627, df = 15.462, p-value = 0.000347 YFP+CC1+: t = 7.1621, df = 11.274, p-value = 1.61e-05 YFP+CC1-: t = 0.72468, df = 24.246, p-value = 0.4756				
Fig. 1D	Welch's t-test. P-Myrf(+/-), n=12; P-Myrf(-/-), n=15.	YFP+CC1+/YFP+CC1-: t = 10.199, df = 11.338, p-value = 4.706e-07				
Fig. 2D	Mixed Two-way ANOVA P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=65 (Female, n=29); Between subject factor: MYRFGenotype (2 levels), Sex (2 levels) Within subject factor: Time (Days, 8 levels)	**Effect**	**DFn**	**DFd**	**F**	**p**
		(Intercept)	1	120	900.615	<0.001
		MYRFGenotype	1	120	10	0.002
		Sex	1	120	13.081	<0.001
		Time	2.74	328.53	170.149	<0.001
		MYRFGenotype:Sex	1	120	3.041	0.084
		MYRFGenotype:Time	2.74	328.53	5.569	0.001
		Sex:Time	2.74	328.53	6.344	<0.001
Fig. 2E	Mixed Two-way ANOVA P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=65 (Female, n=29); Between subject factor: MYRFGenotype (2 levels), Sex (2 levels) Within subject factor: Time (Days, 8 levels)	**Effect**	**DFn**	**DFd**	**F**	**p**
		(Intercept)	1	120	1791.129	<0.001
		MYRFGenotype	1	120	11.922	<0.001
		Sex	1	120	17.481	<0.001
		Time	2.57	308.96	185.786	<0.001
		MYRFGenotype:Sex	1	120	1.867	0.174
		MYRFGenotype:Time	2.57	308.96	4.764	0.005
		Sex:Time	2.57	308.96	4.614	0.006
Fig. 2F	Mann–Whitney U test P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=65 (Female, n=29);	W = 2504, p-value = 0.003369				
Fig. 2G	Mann–Whitney U test P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=65 (Female, n=29);	W = 2536, p-value = 0.001988				
Fig. 2H	Mixed Two-way ANOVA P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=65 (Female, n=29); Between subject factor: MYRFGenotype (2 levels), Sex (2 levels) Within subject factor: Time (12 levels (20 mins intervals))	**Effect**	**DFn**	**DFd**	**F**	**p**
		MYRFGenotype	1	120	2.727	0.101
		Sex	1	120	0.753	0.387
		Time	4.83	579.63	62.791	<0.001
		MYRFGenotype:Sex	1	120	5.446	0.021
		MYRFGenotype:Time	4.83	579.63	2.627	0.025
		Sex:Time	4.83	579.63	3.088	0.01
Fig. 2I	Mann–Whitney U test P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=65 (Female, n=29);	W = 2413.5, p-value = 0.01317				
Fig. 2J	Mann–Whitney U test P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=65 (Female, n=29);	W = 2411, p-value = 0.01364				
Fig. 2K	Spearman's rank correlation rho:	Across Genotype: S = 63551, p-value <0.001, rho = 0.7999962 P-Myrf(+/−): S = 12760, p-value <0.001, rho 0.6271278 P-Myrf(-/-): S = 5874.6, p-value <0.001, rho 0.8716223				

		cor.test(a, b method=c("spearman")) P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=65 (Female, n=29);					
Fig. 2L	Mixed Two-way ANOVA P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=64 (Female, n=29); Between subject factor: MYRFGenotype (2 Levels), Sex (2 Levels) Within subject factor: Time (6 levels (10 mins intervals))	**Effect**	**DFn**	**DFd**	**F**	**p**	
		MYRFGenotype	1	120	2.848	0.094	
		Sex	1	120	0.058	0.81	
		Time	2.21	265.32	88.001	<0.001	
		MYRFGenotype:Sex	1	120	2.846	0.094	
		MYRFGenotype:Time	2.21	265.32	9.06	<0.001	
		Sex:Time	2.21	265.32	1.81	0.161	
Fig. 2M	Mann–Whitney U test P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=65 (Female, n=29);	W = 2490, p-value = 0.00421					
Fig. 2N	Mixed Two-way ANOVA P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=64 (Female, n=29); Between subject factor: MYRFGenotype (2 Levels), Sex (2 Levels) Within subject factor: Time (5 Intervals (1H intervals))	**Effect**	**DFn**	**DFd**	**F**	**p**	
		MYRFGenotype	1	118	4.998	0.027	
		Sex	1	118	3.09	0.081	
		Time	2.2	259.93	53.822	<0.001	
		MYRFGenotype:Sex	1	118	3.709	0.057	
		MYRFGenotype:Time	2.2	259.93	1.224	0.298	
		Sex:Time	2.2	259.93	3.045	0.044	
Fig. 2O	Mann–Whitney U test P-Myrf(+/-), n=58 (Female, n=28); P-Myrf(-/-), n=64 (Female, n=29);	W = 2278, p-value = 0.0307					
Fig. 3D	Mixed Two-way ANOVA P-Myrf(+/-), n=12 P-Myrf(-/-), n=13 Between subject factor: MYRFGenotype (2 Levels), Batch (4 Levels) Within subject factor: ROI (4 Levels)	**Effect**	**DFn**	**DFd**	**F**	**p**	
		MYRFGenotype	1	17	7.318	0.015	
		Batch	3	17	0.889	0.467	
		ROI	1.27	21.57	190.596	<0.001	
		MYRFGenotype:Batch	3	17	2.786	0.072	
		MYRFGenotype:ROI	1.27	21.57	2.084	0.161	
		Batch:ROI	3.81	21.57	3.034	0.041	
Fig. 3D	Two-way ANOVAs P-Myrf(+/-), n=12 P-Myrf(-/-), n=13 Between subject factor: MYRFGenotype (2 Levels), Batch (4 Levels)	**ROI**	**Effect**	**F**	**p**	**p (adjusted)**	
		MTRHipp	MYRFGenotype	0.969	0.339	1	
		MTRM1	MYRFGenotype	10.523	0.005	0.02	
		MTRM2	MYRFGenotype	9.029	0.008	0.032	
		MTRS1	MYRFGenotype	5.224	0.035	0.14	
Fig. 3E	Two-way ANOVA P-Myrf(+/-), n=12 P-Myrf(-/-), n=13 Between subject factor: MYRFGenotype (2 Levels), Batch (4 Levels)	**ROI**	**DFn**	**DFd**	**F**	**p**	
		MTRaCCWMskel	1	17	7.77	0.013	
Fig. 3G	Grey matter skeleton - MTR Two-way ANOVA P-Myrf(+/-), n=25 P-Myrf(-/-), n=25 Between subject factor: MYRFGenotype (2 Levels), Batch (4 Levels)	Effect	DFn	DFd	F	p	
		MYRFGenotype	1	34	6.04	0.019	
		Wheel	1	34	0.065	0.8	
		Batch	3	34	2.179	0.108	
		MYRFGenotype:Wheel	1	34	0.061	0.807	
		MYRFGenotype:Batch	3	34	1.117	0.356	

		Wheel:Batch	3	34	0.395	0.758
Fig. 3H	White matter skeleton - MTR Two-way ANOVA P-Myrf(+/-), n=25 P-Myrf(-/-), n=25 Between subject factor: MYRFGenotype (2 Levels), Wheel (2 Levels), Batch (4 Levels)	**Effect**	**DFn**	**DFd**	**F**	**p**
		MYRFGenotype	1	34	4.728	0.037
		Wheel	1	34	0.028	0.869
		Batch	3	34	1.563	0.216
		MYRFGenotype:Wheel	1	34	0.083	0.775
		MYRFGenotype:Batch	3	34	0.939	0.433
		Wheel:Batch	3	34	0.697	0.56
Fig. 4E	Mixed Two-way ANOVA P-Myrf(+/-), n=3 P-Myrf(-/-), n=4 Between subject factor: MYRFGenotype (2 Levels), Within subject factor: EEG location (2 Levels)	**Effect**	**DFn**	**DFd**	**F**	**p**
		Genotype	1	5	10.398	0.023
		EEG location	1	5	4.207	0.096
		Genotype: EEG location	1	5	0.779	0.418
Fig. 4F	Mixed Two-way ANOVA P-Myrf(+/-), n=3 P-Myrf(-/-), n=4 Between subject factor: MYRFGenotype (2 Levels), Within subject factor: EEG location (2 Levels)	**Effect**	**DFn**	**DFd**	**F**	**p**
		Genotype	1	5	12.751	0.016
		EEG location	1	5	0.155	0.71
		Genotype: EEG location	1	5	1.463	0.28
Fig. 4H	Mixed Two-way ANOVA P-Myrf(+/-), n=3 P-Myrf(-/-), n=4 Between subject factor: MYRFGenotype (2 Levels), Within subject factor: EEG location (2 Levels)	**Effect**	**DFn**	**DFd**	**F**	**p**
		Genotype	1	5	11.385	0.02
		EEG location	1	5	1.762	0.242
		Genotype:EEG location	1	5	1.071	0.348
Suppl. Fig. 1H	Mann–Whitney U test Males: n=67; Females: n=57;	W = 2411, p-value = 0.01364				

		Effect	**DFn**	**DFd**	**F**	**p**
Suppl. Fig 1I	Mixed Two-way ANOVA P-Myrf(+/-), n=59 (Female, n=28); P-Myrf(-/-), n=64 (Female, n=29); Between subject factor: MYRFGenotype (2 Levels), Sex (2 Levels) Within subject factor: Time (192 Intervals – 1h size))	MYRFGenotype	1	120	4.484	0.036
		Sex	1	120	30.276	<0.001
		Interval	191	22920	103.115	<0.001
		MYRFGenotype:Sex	1	120	0.604	0.438
		MYRFGenotype:Interval	191	22920	2.12	<0.001
		Sex:Interval	191	22920	8.859	<0.001
Suppl. Fig 3A-D	Mixed Two-way ANOVA P-Myrf(+/-), n=25 P-Myrf(-/-), n=25 Between subject factor: MYRFGenotype (2 Levels), Batch (4 Levels) Within subject factor: ROI (4 Levels)	**Effect**	**DFn**	**DFd**	**F**	**p**
		MYRFGenotype	1	34	10.464	0.003
		Wheel	1	34	0.196	0.661
		Batch	3	34	3.426	0.028
		ROI_GM_MTR	1.27	43.21	299.344	<0.001
		MYRFGenotype:Wheel	1	34	0.076	0.784
		MYRFGenotype:Batch	3	34	2.39	0.086
		Wheel:Batch	3	34	0.455	0.715
		MYRFGenotype:ROI_GM_MTR	1.27	43.21	0.196	0.719
		Wheel:ROI_GM_MTR	1.27	43.21	0.34	0.615

		ROI	**Effect**	**F**	**p**	**p (adjusted)**
Suppl. Fig 3A-D	Two-way ANOVAs P-Myrf(+/-), n=25 P-Myrf(-/-), n=25	MTRHipp	MYRFGenotype	6.142	0.018	0.072
		MTRM1	MYRFGenotype	7.814	0.008	0.032
		MTRM2	MYRFGenotype	4.816	0.035	0.14

	Between subject factor: MYRFGenotype (2 Levels), Batch (4 Levels)	MTRS1	MYRFGenotype	13.376	<0.001	0.0032
Suppl. Fig. 3E	White matter skeleton - MTR Two-way ANOVA P-Myrf(+/-), n=25 P-Myrf(-/-), n=25 Between subject factor: MYRFGenotype (2 Levels), Wheel (2 Levels), Batch (4 Levels)	**Effect**	**DFn**	**DFd**	**F**	**p**
		MYRFGenotype	1	34	6.574	0.015
		Wheel	1	34	0.1	0.754
		Batch	3	34	4.91	0.006
		MYRFGenotype:Wheel	1	34	0.621	0.436

.055