Affinity hierarchies and amphiphilic proteins underlie the co-assembly of nucleolar and heterochromatin condensates

Authors

Srivarsha Rajshekar[1], Omar Adame-Arana[2,3], Gaurav Bajpai[2], Kyle Lin[1], Serafin Colmenares[1], Samuel Safran[2], Gary H Karpen[1,4]

[1] Department of Molecular and Cell Biology, University of California Berkeley, Berkeley, USA

[2] Department of Chemical and Biological Physics, Weizmann Institute of Science, Rehovot, Israel

[3] Max Planck Institute for the Physics of Complex Systems, Dresden, Germany

[4] Division of Biological Sciences and the Environment, Lawrence Berkeley National Laboratory, Berkeley, USA

Correspondence: gkarpen@berkeley.edu

Abstract

Nucleoli are surrounded by Pericentromeric Heterochromatin (PCH), reflecting a close spatial association between the two largest biomolecular condensates in eukaryotic nuclei. This nuclear organizational feature is highly conserved and is disrupted in diseased states like senescence, however, the mechanisms driving PCH-nucleolar association are unclear. High-resolution live imaging during early Drosophila development revealed a highly dynamic process in which PCH and nucleolar formation is coordinated and interdependent. When nucleolus assembly was eliminated by deleting the ribosomal RNA genes (rDNA), PCH showed increased compaction and subsequent reorganization to a shell-like structure. In addition, in embryos lacking rDNA, some nucleolar proteins were redistributed into new bodies or 'neocondensates,' including enrichment in the core of the PCH shell. These observations, combined with physical modeling and simulations, suggested that nucleolar-PCH associations are mediated by a hierarchy of affinities between PCH, nucleoli, and 'amphiphilic' protein(s) that interact with both nucleolar and PCH components. This result was validated by demonstrating that the depletion of one candidate amphiphile, the nucleolar protein Pitchoune, significantly reduced PCH-nucleolar associations. Together, these results unveil a dynamic program for establishing nucleolar-PCH associations during animal development, demonstrate that nucleoli are required for normal PCH organization, and identify Pitchoune as an amphiphilic molecular link that promotes PCH-nucleolar associations. Finally, we propose that disrupting affinity hierarchies between interacting condensates can liberate molecules to form neocondensates or other aberrant structures that could contribute to cellular disease phenotypes.

Introduction

The eukaryotic nucleus is organized into different membrane-less compartments known as biomolecular condensates that assemble via multivalent interactions between their constituent molecules. Prominent examples include the nucleolus, nuclear speckles, and heterochromatin, essential for cellular viability and function. An individual condensate can concentrate several distinct macromolecules in cells, including structured and intrinsically disordered proteins and nucleic acids (Lyon et al. 2021; Shin & Brangwynne 2017). Condensates with the same components that nucleate at different cellular locations can coarsen by fusions into larger clusters, whereas condensates with distinct compositions do not fuse (Ditlev et al. 2018). Nevertheless, in the crowded environment of the nucleus, distinct condensates can display close, conserved associations to form higher-order complex structures (Fare et al. 2021). Although many studies have examined the formation and function of individual condensates, we know relatively little about the interplay between distinct interacting condensates *in vivo* with respect to how they form and impact each other's structure and function.

Two prominent nuclear condensates that are proximally located in a highly conserved manner are the Nucleolus and Pericentromeric Heterochromatin (PCH) (Bizhanova & Kaufman 2021; Padeken & Heun 2014). The nucleolus is a multi-layered biomolecular condensate that plays an essential role in the assembly of ribosomal subunits and forms by liquid-liquid phase separation of its constituent molecules (Brangwynne et al. 2011; Feric et al. 2016; Lafontaine et al. 2020). RNA-Polymerase I (RNA-Pol-I) transcribes tandemly-repeated ribosomal RNA genes (rDNA) to form the innermost layer (Fibrillar Center or FC), rRNA is processed by proteins such as Fibrillarin to generate the Dense Fibrillar Compartment (DFC), and ribosomal proteins assemble with processed rRNA to form ribosomes in the outermost layer, the Granular Compartment (GC) (Feric et al. 2016; Thiry & Lafontaine 2005). The different subcompartments recruit specific factors involved in rRNA transcription, processing, and ribosome assembly in a sequential manner to facilitate the vectorial expulsion of ribosomes through and out of the nucleolus (King et al. 2022; Riback et al. 2023). In most eukaryotic nuclei, the nucleolus is surrounded by PCH, a compact chromatin compartment that assembles on megabases of pericentromeric repeats, including tandemly repeated satellite DNA and transposable elements. While the nucleolus is the most transcriptionally active site in the nucleus, PCH is associated with transcriptional silencing and has essential roles in nuclear architecture and genome stability (Janssen et al. 2018). PCH is distinct from the inactive B compartment as defined by chromosome conformation capture experiments since Hi-C analyses typically exclude repetitive PCH sequences (Lieberman-Aiden

et al. 2009). Under the microscope, PCH can be visualized as chromatin regions enriched for the

AT-rich DNA dye DAPI, histone modifications di- and tri-methylation of histone H3 (H3K9me2/3),

and the cognate epigenetic reader Heterochromatin Protein 1 (HP1) (Politz et al. 2013). HP1 is a

multivalent protein with structured and disordered domains (Eissenberg & Elgin 2000) that

undergoes phase separation *in vitro* and preferentially partitions DNA and nucleosomes (Keenen

et al. 2020; Larson et al. 2017), while *in vivo,* it forms a liquid-like condensate nucleated by

H3K9me2/3 enriched chromatin (Strom et al. 2017; Tortora et al. 2023; Wang et al. 2019).

A simple explanation for why PCH is organized close to the nucleolus is that tandem arrays of

rDNA genes are positioned within pericentromeric satellite repeats on a subset of chromosomes

(Potapova & Gerton 2019). However, proximity mapping nucleolus-associated domains (NADs)

and IF-FISH revealed that PCH sequences from chromosomes without rDNA genes also

associate with the nucleolus (Németh et al. 2010; Van Koningsbruggen et al. 2010). Thus, cis-

proximity to rDNA is not necessary for PCH-nucleolar associations, and how PCH from all

chromosomes becomes positioned around the nucleolus is unknown. Loss of the major nucleolar

component nucleophosmin (NPM1) and its Drosophila homolog NLP in human and fly cell lines

results in disorganization of nucleolus-associated heterochromatin (Holmberg Olausson et al.

2014; Padeken et al. 2013). Conversely, loss of major heterochromatin factors like Su(var)3-9

and HP1 impaired rDNA and nucleolar integrity (Ballmer et al. 2023; Peng & Karpen 2007),

suggesting structural or functional relationships between the two domains. The proximal

positioning of rDNA to PCH sequences is likely functionally relevant since not all rDNA genes are

expressed, and a subset is silenced by heterochromatin formation (Srivastava et al. 2016). PCH

repeats are dissociated from the nucleolus in senescent cells, suggesting that PCH-nucleolar

association is linked to cellular health (Dillinger et al. 2017). Thus, the association of PCH and

nucleoli is a prominent and conserved feature of 3D nuclear organization, but we lack mechanistic

insight into how the two condensates co-assemble and affect each other's structure and function.

This study uses live imaging and genetic tools in Drosophila embryos and tissues to reveal highly

dynamic behaviors that characterize the *de novo* assembly of PCH around the nucleolus. During

the initial phases of formation, PCH and nucleolar compartments are spatially separated. After

independent growth and fusions, PCH first assembles at the nuclear edge and progressively

reorganizes around the nucleolus as the embryo develops. Mutational analyses that perturb

heterochromatin disrupt nucleolar shape and morphology. Reciprocally, embryos that lack rDNA

and, thus, a functional nucleolus, show dramatic reorganization of PCH characterized by

increased compaction followed by reorganization to a shell-like structure. *In vivo* experiments and

physical modeling revealed that a hierarchy of interaction strengths between PCH, nucleoli, and
'amphiphilic' protein(s) able to interact with both nucleolar and PCH components generate the
layered organization of nucleoli and PCH. This model led to identifying Pitchoune, a DEAD-box
RNA-Helicase protein, as a molecular linker responsible for PCH-nucleolar associations. Finally,
these results suggest that perturbed condensate associations and affinity hierarchies can
generate abnormal cellular phenotypes in disease states.

Results
De novo **establishment of PCH-nucleolar associations during early Drosophila**
embryogenesis is dynamic
Although PCH and nucleolar assembly dynamics have been studied independently in Drosophila
embryos (Falahati et al. 2016; Strom et al. 2017), they have not been analyzed simultaneously
during *de novo* establishment. To determine how PCH is assembled de novo around the
nucleolus, we performed high-resolution time-lapse imaging of early Drosophila embryos
expressing RFP-Fib and GFP-HP1a transgenes as markers for the nucleolus and PCH,
respectively. Drosophila embryos undergo 14 synchronized, syncytial nuclear divisions (S and M
phases only) before cellularizing at blastoderm (cycle 14) and thereafter undergoing cell divisions
(Farrell & O'Farrell 2014). Major chromatin features, including H3K9 methylation, are
progressively established during the syncytial cycles (Ahmad & Henikoff 2022; Seller et al. 2019).
PCH condensates first emerge in cycle 11 (Strom et al. 2017), while nucleoli first emerge in cycle
13, seeded by transcribing rRNA (Falahati et al. 2016), making cycle 13 the earliest time PCH
and nucleolar condensates appear in the same nuclei. Upon entry in cycle 13 interphase, HP1a
and Fibrillarin proteins are initially diffuse throughout the nucleus, then within ~8 mins, each
becomes enriched in multiple, distinct foci (**Figure 1A(i) and Supplementary Movie 1**). Small
PCH and nucleolar condensates remain separated throughout cycle 13, likely because growth is
limited by the short interphase (~15 mins) before dissolution in mitosis.

Cycle 14 begins like cycle 13 in that HP1a and Fibrillarin foci emerge soon after mitotic exit and
are initially separated. However, PCH and nucleolar condensates grow larger and undergo liquid-
like self-fusions due to the extended length of cycle 14 interphase (~90 mins) (Falahati et al. 2016;
Strom et al. 2017; Tortora et al. 2023) (**Figure 1A(ii) and Supplementary Movie 2**). By the end
of cycle 14, a small section of the HP1a domain is associated with Fibrillarin, corresponding to
the rDNA locus on the sex chromosomes (**Extended Figure 1A-B**). The majority of the HP1a

Figure 1: PCH reorganizes from the nuclear to the nucleolar edge during early Drosophila development. (**A**) Maximum intensity projections (xy view) of individual live nuclei of GFP-HP1a (magenta) and RFP-Fibrillarin (green) localization in Cycles 13-17 of Drosophila embryogenesis. The numbers on the top left corner of each image indicate time (in minutes) after mitotic exit. (**B**) Quantification of HP1a occupancy at the nucleolar edge during specific developmental cycles. "Early Cycle" refers to nuclei between 15-30 mins into the specified interphase, "Late Cycle" refers to nuclei between 50-70 mins. Data points = the mean, error bars = 95% confidence intervals around the mean. n>50 nuclei (from 5 embryos). at each time point. (**C**) Representative maximum intensity projections of PCH (GFP-HP1a, magenta) and nucleoli (RFP-Fibrillarin, green) in live epidermal nuclei in a late Drosophila embryo (Stage 16, ~14-16hr). The nucleus within the white dashed box is magnified on the right and presented in xy and xz views. White dashed lines indicate the nuclear boundary. (**D**) Schematic depicting the organization of the nucleolus and PCH in nuclei in early (top) and late (bottom) Drosophila embryos. 359bp and 1.686 repeats are major pericentromeric sequences on Chr X and Chr 2 & 3 respectively.

domain, including chromosomes 2 and 3, is distributed apically and juxtaposed to the nuclear

periphery (**Figure 1A(ii) and Extended Figure 1C**). This 'extended' conformation represents a

previously undescribed intermediate during the establishment of PCH-nucleolar associations.

Continued live and fixed imaging of HP1a and Fibrillarin in post-blastoderm, asynchronous cell
divisions revealed how the extended PCH configuration dynamically transitions into the canonical
surrounded form observed in later developmental stages. As for cycles 13 and 14, in each post-
blastoderm interphase (cycle 15 onwards), HP1a and Fibrillarin proteins are initially diffuse, then
nucleate into separate foci that grow and fuse to form the extended intermediate. The PCH
remains in the extended configuration and predominantly associates with the nuclear periphery
through the rest of cycle 15 (**Figure 1A(iii) and Supplementary Movie 3**) but transitions between
the extended and surrounded configurations during cycle 16 (**Figure 1A(iv) and Supplementary**
Movie 4). Finally, stable formation of the surrounded organization is observed ~15 min into cycle
17 interphase (**Figure 1A(v) and Supplementary Movie 5**), after PCH condensates quickly
transition through the extended configuration. Further analyses substantiate these observed live
PCH-nucleolar dynamics. First, the fraction of the nucleolar edge occupied by HP1a increases
between cycles 14 and 17 (**Figure 1B and** representative stills in **Extended Figure D**). Second,
H3K9me2 displays the extended conformation relative to Fibrillarin in fixed Stage 8/cycle 15
embryos and localizes to the nucleolar edge later in development (**Extended Figure 1E)**. Third,
immuno-FISH revealed that satellite repeats 1.686 on Chr 2 and 3 are located farther from the
nucleolus in cycle 15 (Stage 8; mean=1.28μm) and display significantly more proximal localization
in Stage 16 (mean=0.31μm) (**Extended Figure 1F-G)**.

Live imaging in late embryos (~12-16 hr, Embryonic Stage 14-16), larval tissues, and cultured S2
cells expressing fluorescently tagged transgenes for HP1a revealed that the canonical
surrounded conformation persists in later developmental stages (**Figure 1C and Extended**
Figure 2A). *Drosophila* nucleoli are also multi-layered as in other eukaryotes; the GC is marked
by Modulo (the fly ortholog for Nucleolin), Fibrillarin marks the DFC, and Rpl-135, a subunit of
RNA-Pol-I localizes within the Fibrillarin compartment (**Extended Figure 2B**). HP1a/PCH
organizes around the outermost nucleolar layer (GC) in late Drosophila embryos (**Extended**
Figure 2C). Together, we conclude that PCH-nucleolar associations are highly dynamic during
each interphase; PCH and nucleolar proteins are diffuse upon mitotic exit, then both condensates
undergo independent nucleation, growth, and fusions, and display cycle-specific spatiotemporal
differences while transitioning from the extended (nuclear periphery) to the surrounded (nucleolar
periphery) configurations (summarized in **Figure 1D, schematic)**.

Reduced levels of heterochromatin result in decondensed rDNA and enlarged nucleoli
Is the organization of PCH and nucleolar condensates interdependent? We investigated whether
PCH is required for proper nucleolus formation by removing one of the major H3K9

methyltransferase genes. Su(var)3-9[06/17] embryos (Schotta et al. 2002) generated from Su(var)3-

9[06/17] mothers are depleted for H3K9me2 in cycle 14 (**Extended Figure 3A**) and display a 1.5-fold

increase in nucleolar volume compared to wild-type controls (**Extended Figure 3B-C).** In addition,

combined immuno-FISH analysis revealed that the high-intensity rDNA block located at the

nucleolar edge in wild-type is reduced in Su(var)3-9[06/17] embryos, with corresponding increases

in the amount of dispersed rDNA within the nucleolus (**Extended Figure 3D**). Similarly, nucleolar

volume increased in cycle 14 after maternal knockdown of HP1a using RNAi (**Extended Figure

3E-F**). We conclude that PCH constrains nucleolar size and limits rDNA distribution within the

nucleolus (summarized in **Extended Figure 3G**). This is consistent with previous studies that

show HP1/H3K9me2/3 is enriched at silent rDNA in flies and mammalian cells (Fefelova et al.

2022; Santoro et al. 2002) and that HP1 maintains nucleolar structure by excluding repeats from

the nucleolus (Ballmer et al. 2023). Further research is required to determine if the absence of

H3K9me2/3 and HP1 alters nucleolar morphology by de-silencing rRNA transcription versus a

requirement for physical association with PCH condensates.

Embryos lacking nucleoli display abnormal PCH organization and neocondensate

formation

Next, we determined if nucleolar condensates impact PCH organization by imaging RFP-HP1a

and GFP-Fib in embryos that completely lack rDNA repeats (designated hereafter as -rDNA) due

to the presence of a rearranged X chromosome (C(1)DX/0, Valencia et al., 1949). No functional

nucleoli are formed in -rDNA embryos (Falahati et al. 2016), but they develop into late-stage

embryos due to maternal deposition of ribosomes. While no nucleolus is present, Fibrillarin forms

spherical structures (as previously reported by Falahati et al. 2016), which are located at a

significantly longer distance from the HP1a/PCH condensate compared to +rDNA controls

(**Figure 2A-B;** +rDNA mean=1.78μm, -rDNA mean=3.69μm, p<0.0001). This result suggests that

Fibrillarin and HP1a proteins do not interact directly and, thus, are unlikely to mediate the

canonical surrounded organization. Instead, we propose that when factors responsible for

nucleolus nucleation (rDNA/rRNA) are not present, Fibrillarin self-associations (Berry et al. 2015)

and/or secondary affinities with other structures or molecules are sufficient to cause the formation

of new enrichments not found in wild-type cells (hereafter termed 'neocondensates').

Most importantly, PCH condensate morphology is significantly perturbed in embryos lacking

rDNA. The "extended" HP1a conformation normally observed in cycles 14 and 15 is replaced by

a more compact, rounded structure at the apical end of -rDNA nuclei (**Figure 2A, C**). The aspect

Figure 2: Increased PCH compaction in embryos lacking rDNA. (**A**) Maximum intensity projections of live nuclei from wildtype embryos (+rDNA) and mutant embryos lacking rDNA (-rDNA) at Cycle 14, showing RFP-HP1a (magenta) and GFP-Fibrillarin (green). The dashed box highlights the nuclei enlarged in the panel below. (**B**) Distance between the centers of geometry of HP1a and Fibrillarin in +rDNA and -rDNA embryos. n>200 nuclei (from 5 embryos) for each genotype. (**C**) Maximum intensity projections from live imaging of nuclei from +rDNA and -rDNA embryos showing the reassembly of RFP-HP1a (magenta) and GFP-Fibrillarin (green) at the indicated minutes after the start of Cycle 15. (**D**) Quantification of the aspect ratio (major axis over minor axis) of HP1a segments 15 minutes after the start of Cycle 15 in +rDNA and -rDNA embryos. n>30 nuclei in each genotype. (**E**) Maximum intensity projections of FISH (Fluorescence In Situ Hybridization) of 1.686 (magenta) and AAGAG (green) in DAPI (blue)-stained nuclei in +rDNA and -rDNA Stage 10 embryos. The dashed boxes mark the nuclei enlarged on the right. (**F**) Quantification of the distance between AAGAG and its nearest 1.686 locus. n>40 pairs of loci in each genotype.

ratio (major axis/minor axis) of the HP1a domain is significantly reduced (**Figure 2D;** +rDNA

mean=1.96, -rDNA mean=1.40, p<0.0001), and the distance between pericentromeric repeats

1.686 (Chr 2 and 3) and AAGAG (all chromosomes) is decreased in -rDNA nuclei compared to

+rDNA controls (**Figure 2E-F;** +rDNA mean=0.74μm, -rDNA mean=0.32μm, p<0.0001)). We

conclude that nucleoli are required to avoid hyper-compaction of PCH condensates in early

Drosophila embryos (before the end of cycle 14).

Figure 3: A protein-filled core reshapes the PCH condensate to a sphere-like structure in -rDNA developing embryos. (**A**) Representative stills of live +rDNA and -rDNA nuclei at Cycle 14 and in Stage 16 (late stage) embryos showing RFP-HP1a (magenta) and GFP-Fibrillarin (green). Nuclei marked with the yellow arrowhead are enlarged on the right. (**B**) Line scans of the intensity along the dashed yellow line in the optical slices shown in (A)**.** (**C**) Time-lapse stills (single slices) of a -rDNA amnioserosa nucleus with RFP-HP1a (magenta) and GFP-Fibrillarin (green). (**D**) Orthogonal views (single slices) of RFP-HP1a in a -rDNA live amnioserosa nucleus. The yellow lines indicate the location of the orthogonal slices. (**E**) Late embryos (Stage 16) with the +rDNA or -rDNA genotype was co-stained with a pan-protein label, 488 NHS Ester (green) and DAPI (magenta). Nuclei marked with the black dashed box are enlarged on the right.

Unexpectedly, in late-stage -rDNA Drosophila embryos, the PCH morphed from a hyper-

compacted cluster to a hollow sphere-like structure, surrounding a core devoid of HP1a (hereafter

'PCH void') (**Figure 3A-B**). To visualize this transformation at increased spatial resolution, we

performed live imaging in the amnioserosa, whose large nuclei form a monolayer on the dorsal

surface of the embryo during gastrulation (Lacy & Hutson 2016). Live imaging of the amnioserosa

nuclei in -rDNA Drosophila embryos revealed the stepwise transformation of the compacted PCH

to the formation of a progressively larger PCH void (**Figure 3C-D and Supplementary Movie 6**).

In addition to the absence of HP1a, the PCH void lacked the major nucleolar proteins Fibrillarin

(**Figure 3A-B**) and Modulo (**Extended Figure 4A**), which formed neocondensates or dispersed

in the nuclear space, respectively. The PCH void did not contain DNA (DAPI staining), histones
marked with H3K9me2 (IF), or any significant accumulation of RNAs (Propidium Iodide staining)
(**Extended Figures 4B-D**). However, treatment with 488 NHS Ester, a pan-protein label (M'Saad
& Bewersdorf 2020), revealed that the PCH void was enriched for unknown proteins (**Figure 3E**).
We conclude that PCH initially displays an atypical, hyper-compacted morphology in embryos
lacking rDNA and nucleoli. Further, the PCH morphs into an abnormal sphere-like structure whose
central core (the PCH void) lacks HP1a, nucleolar proteins, chromatin, and RNA but is filled with
unknown protein(s) that may represent another neocondensate. Importantly, these results
suggest that nucleoli are required for the normal distribution and organization of PCH
condensates in nuclear space and that disrupting interactions within or between condensates can
create new nuclear structures.

Coarse-grained modeling recapitulates *in vivo* phenotypes and highlights a potential role
for amphiphilic proteins in mediating PCH-nucleolar organization
We used coarse-grained molecular dynamics simulations and physical modeling to identify the
molecular interactions and biophysical parameters that could mediate nucleolar-PCH
associations in wild-type embryos, as well as PCH disorganization in the absence of rDNA. Based
on our experimental observations and previous studies, the model was constrained by four main
features. First, some nucleolar proteins like Fibrillarin have an intrinsic tendency to form
condensates (Figure 2A) (Falahati et al. 2016; Feric et al. 2016). Second, PCH forms a
condensate that associates with the nucleolus (Figure 1) (Strom et al. 2017). Third, PCH and
Fibrillarin have no direct associations (Figure 2A). Fourth, unknown proteins accumulate in the
PCH void in -rDNA embryos (Figure 3E). We thus modeled four minimal components (see details
in Materials and Methods) (**Figure 4A**): (i) PCH (H) as a long, self-interacting polymer, (ii) rDNA
(rD) as a polymeric block within the chromatin fiber that experiences good solvent conditions and
is flanked by two PCH blocks, (iii) Fibrillarin (F), representative of a self-associating nucleolar
protein, and (iv) an 'amphiphilic' protein (X), defined here as having shared affinities for both PCH
and nucleolar components (etymology: Greek for 'likes both'). Similar 'amphiphilic' proteins have
previously been synthesized and demonstrated to form multiphasic condensates in vitro (Kelley
et al. 2021).

The theory of three-phase wetting of simple liquids demonstrates that a balance between the
interfacial tensions of the different liquids determines the spatial configurations of the phases
(Torza & Mason 1969). Thus, we propose that to form the layered PCH-nucleolar organization,
the self-associations of Fibrillarin must be stronger than those of the amphiphilic protein, which in

Figure 4: A hierarchy of interaction strengths between PCH, nucleoli and 'amphiphilic' protein(s) recapitulates in vivo +rDNA and -rDNA phenotypes. (**A**) Course-grained modelling of nucleolar-PCH assembly with four minimal components: PCH (H) as a self-interacting polymer (magenta) rDNA (rD) as polymeric block embedded within PCH (red), Fibrillarin (F) as a representative of a self-associating nucleolar protein (green), and an 'amphiphilic' protein (X) that has an affinity for both PCH and nucleolar components (yellow). (**B and C**) The matrix indicates the strength of interaction (blue gradient, units: k_BT) between Fibrillarin (F), rDNA (rD), PCH (H) and an unknown amphiphilic protein (X), with simulated outcomes for +rDNA and -rDNA conditions on the right. (**D**) Distance (d_{HF}) between the centers of mass of PCH (H) and Fibrillarin (F) clusters in +rDNA and -rDNA simulations.

turn must be stronger than the self-interactions of PCH, since interfacial tension is proportional to
the relative interaction strengths. These considerations led us to define a hierarchy of interaction
strengths (rD-F=F-F>X-X>F-X>X-H>H-H) as a foundation for simulations using the parameters
listed in the 4x4 interaction matrix (**Figure 4B**). Importantly, simulating this affinity hierarchy
recapitulated the canonical PCH-nucleolar organization observed in +rDNA animals *in vivo*
*(***Figure 4B and Supplementary Movie 7***)*. Various scenarios incompatible with the experimental
observations of the layered organization of the nucleolus and PCH arise by changing the
hierarchy of self-interaction strengths in the simulations (**Extended Figure 5**). The detailed
rationale for the choice of parameters is described in the Methods section.

Setting the rDNA-Fibrillarin interactions to zero to model the consequences of eliminating the
rDNA resulted in the formation of the protein X neocondensate within the PCH void and the
spatially separated Fibrillarin neocondensate (**Figure 4C-D**), as observed in vivo (Figures 2 and
3). These phenotypes are observed in the simulations whenever the protein X – PCH interaction
strength is greater than or equal to the X – Fibrillarin interactions (i.e., X-H≥F-X) (**Extended**

Figure 5E-F). In the model, the nucleation of the new phase of amphiphilic protein X occurs within
the PCH to reduce the cost of creating an interface of protein X facing the solvent. Instead, being
surrounded by the PCH can reduce this interfacial energy, as expected from a wetting picture of
simple liquids (Torza & Mason 1969). In sum, simply eliminating the rDNA from this minimal
model, and thus the ability to recruit nucleolar components such as Fibrillarin, recapitulates both
the PCH void filled with X protein and Fibrillarin neocondensate phenotypes observed in -rDNA
nuclei *in vivo*.

We conclude that incorporating affinity hierarchies and an amphiphilic protein X into a minimal
model recapitulates the observed *in vivo* organization of PCH and nucleolar condensates in both
+rDNA and -rDNA conditions (**Figure 4B, C**). Further, the experimental and simulation results
indicate that the same amphiphilic protein(s) could mediate the surrounded configuration in wild-
type and accumulation in the PCH void in the -rDNA condition. This led us to predict that protein
X should be a self-associating, outer nucleolar/granular compartment protein with an affinity for
PCH components, such as HP1a.

The DEAD-box RNA Helicase Pitchoune is a granular compartment protein that forms a
neocondensate within PCH in -rDNA embryos
The physical modeling results predicted that amphiphilic proteins interacting with both PCH and
the nucleolus would be enriched in the PCH void in -rDNA embryos. A candidate for such a protein
emerged in Falahati & Wieschaus (2017) who reported that the DEAD-Box RNA Helicase
nucleolar component Pitchoune (Pit) (Zaffran et al. 1998) mislocalized in cycle 14 -rDNA embryos
to the apical part of the nucleus. We reasoned that it could be co-localizing with the hyper-
compacted PCH observed in the absence of nucleoli, suggesting an affinity for PCH (**Figure 2A**).
Moreover, Pitchoune and its human ortholog DDX18 have an N-terminal disordered domain that
could mediate self-association and/or nucleolar interactions, a helicase core, and a consensus
PxVxL motif found in many HP1a binding partners (**Figure 5A**) (Thiru et al. 2004). To test the
hypothesis that Pitchoune acts as the predicted amphiphilic protein X *in vivo*, we performed live
imaging in Drosophila embryos expressing RFP-Fibrillarin and Pitchoune-GFP transgenes
(Falahati & Wieschaus 2017). In +rDNA, Pitchoune localizes to the nucleolus with enrichment at
the edge of Fibrillarin, suggesting that Pitchoune is enriched in the granular compartment, thus
juxtaposed with the surrounding PCH (**Figure 5B**). However, in -rDNA embryos, Pitchoune and
Fibrillarin do not associate and form separate, high-intensity spherical bodies (**Figure 5B**),
suggesting that Pitchoune does not interact with Fibrillarin in the absence of rDNA and rRNA.

Figure 5: The DEAD-Box RNA Helicase Pitchoune is enriched in the PCH void in -rDNA embryos. (**A**) Protein subdomains including a conserved PXVXL HP1-interaction motif in Pitchoune (Pit) and its human ortholog DDX18. (**B**) Representative images of epidermal nuclei from late-stage Drosophila embryos with +rDNA and -rDNA genetic backgrounds with RFP-Fibrillarin (magenta) and Pitchoune-GFP (green). Nuclei marked with the white dashed box are enlarged on the right. (**C**) Maximum intensity projections of Cycle 14, Cycle 17 and amnioserosa nuclei from +rDNA and -rDNA embryos showing RFP-HP1a (magenta) and Pitchoune-GFP (green). (**D**) Time-lapse stills (single slices) of a -rDNA amnioserosa nuclei with RFP-HP1a (magenta) and Pitchoune-GFP (green). (**E**) Mean intensity, (**F**) Circularity, and (**G**) Area of projections of Pitchoune in -rDNA embryos. Mean of 35-60 nuclei at each time point in n= 3 embryos. Error Bars: SEM.

To investigate whether Pitchoune is enriched in the PCH void in -rDNA embryos, we performed

live imaging in +rDNA and -rDNA embryos co-expressing Pit-GFP and RFP-HP1a transgenes. In

cycle 14 +rDNA nuclei, Pitchoune localizes to the nucleolus while HP1a is in the extended form

away from the nucleolus. In contrast, in -rDNA nuclei, a faint Pitchoune signal co-localizes with

HP1a (**Figure 5C**). As the embryo develops, HP1a appears in the canonical form surrounding

Pitchoune in +rDNA embryos. However, in -rDNA embryos, high-intensity Pitchoune puncta

appear within each compacted HP1a condensate in cycle 17 epidermal nuclei and amnioserosa

nuclei (**Figure 5C**). This revealed that the Pitchoune puncta filling the PCH void are first visible in

cycle 17 in -rDNA embryos, the same developmental stage where the stable surrounded
configuration is established in +rDNA embryos.

Time-lapse imaging of amnioserosa nuclei in -rDNA embryos revealed that the intensity of
Pitchoune within HP1a is initially low and increases ~2.5-fold over an hour (**Figure 5D-E and**
Supplementary Movie 8). Furthermore, the circularity of Pitchoune increases and approaches 1
(in projections) (**Figure 5F**), and the size of Pitchoune decreases over time (**Figure 5G**),
suggesting that Pitchoune demixes from PCH to form a sphere, thereby minimizing its surface
area due to its interfacial tension. Together, these data suggest that the nucleation and growth of
Pitchoune within PCH creates a neocondensate whose emergence coincides with the formation
of the PCH-void in -rDNA embryos. Notably, the predictions of the modeling results were validated
in vivo and led to the discovery of Pitchoune as an example of a new class of amphiphilic proteins
that interact with the nucleolus and PCH.

Knockdown of Pitchoune decreases PCH-nucleolar associations
While the *in vivo* experiments and simulations aligned to reveal that Pitchoune has affinities for
both PCH and the nucleolus, we next asked whether Pitchoune is required for PCH-nucleolar
associations. We first investigated this by decreasing the concentration of the amphiphilic protein
in our physical model while keeping all other parameters the same as in Figure 4B. The
simulations revealed a progressive clustering of PCH away from the nucleolus as Pitchoune levels
decreased (**Figure 6A-B**). However, unlike -rDNA (Figure 4C), PCH maintained a restricted
attachment to Fibrillarin in these conditions since the rDNA polymer is embedded between PCH
blocks. These results led us to hypothesize that Pitchoune could act as a linker protein for
stabilizing PCH-nucleolar associations in the 'surrounded' conformation.

To test this hypothesis *in vivo*, we depleted Pitchoune protein levels using RNAi in Drosophila
tissues. When we knocked down Pitchoune in the female germline using a maternal GAL4 driver,
no eggs were laid (**Extended Figure 6A**), and ovary development was halted (**Extended Figure**
6B), precluding further assessment of the effects of Pitchoune knockdown in early embryos.
Therefore, we used eyeless-GAL4 to knock down Pitchoune in eye-antennal discs, representing
a tissue where nuclei form the canonical surrounded PCH-nucleolar conformation. In third-instar
larvae, upon Pitchoune knockdown, the normal development of the eye-antennal discs is severely
perturbed (**Extended Figure 6C**), consistent with defects in cell proliferation (Zaffran et al. 1998).
Crucially, the PCH-nucleolar organization recapitulated the physical prediction for the depletion
of an amphiphilic molecule linking the PCH and nucleolus. Control samples showed H3K9me2
organized around the nucleolus. However, upon Pitchoune knockdown, both DAPI and H3K9me2

Figure 6: Knockdown of Pitchoune decreases PCH association with the nucleolus. (**A**) Simulation endpoint snapshots demonstrate how decreasing the concentration (c_P) of Pitchoune (yellow) reduces the association of PCH (H) (magenta) around the Fibrillarin condensate (F) (green). (**B**) The average distance (d_{HF}) between the PCH (H) and Fibrillarin (F) clusters from their center of mass is measured at different Pitchoune concentrations (c_P). Error Bars represent standard deviation (SD). (**C**) Immunofluorescence of H3K9me2 (magenta), Fibrillarin (green) and DAPI in nuclei from dissected eye-discs in third instar larvae from UAS-Pitchoune RNAi and UAS-mCherry RNAi (control) driven by the eyeless-GAL4 driver. (**D**) Quantification of the fraction of the nucleolar edge occupied by H3K9me2 in 3D. Each data point indicates the mean value from one animal. n=6 animals with over 500 nuclei in total for each experimental group. Error bars indicate SEM. (**E**) Schematic of the model showing the impact of -rDNA and Pitchoune knockdown on the organization of PCH and nucleoli.

patterns changed, becoming more compact and rounded while still attached to Fibrillarin (**Figure**

6C). We observed a 50% reduction in the fraction of the nucleolar edge in 3D occupied by

H3K9me3 after Pitchoune knockdown (control RNAi mean=0.296, pit RNAi mean=0.146, p-

value=0.0011) (**Figure 6D**). Together, these results reveal a new mechanism for the association

of two distinct biomolecular condensates and demonstrate a previously unappreciated role for the

RNA Helicase Pitchoune in promoting PCH organization around the nucleolus.

Discussion

This study presents four main findings about the organization of pericentromeric heterochromatin

(PCH) and nucleolar condensates in the developing Drosophila embryo. Firstly, we observed that

PCH-nucleolar associations are highly dynamic during initial establishment in early embryos,

characterized by cycle-specific spatio-temporal differences in transitions between the extended

(nuclear periphery) and surrounded (nucleolar periphery) configurations (Figure 1). Secondly, we

have established a structural relationship between these two condensates, discovering that while

PCH plays a role in restricting nucleolar size, the nucleolus facilitates the distribution of PCH

components and repeats within the nuclear space (Figure 2). Thirdly, we have shown that a
hierarchy of interaction strengths between nucleolar and PCH components could regulate their
layered organization. Notably, the model predicted previously unrecognized protein(s) with
'amphiphilic' properties, which exhibit binding affinities to both the nucleolus and PCH (Figure 3-
4). Lastly, we identified Pitchoune, a DEAD-box RNA-Helicase, as a potential amphiphilic protein
that stabilizes PCH-nucleolar associations and demonstrated that Pitchoune removal reduces
nucleolar-PCH interactions and disrupts the surrounded configuration (Figure 5-6). These findings
provide mechanistic insights into the co-assembly dynamics and organization of PCH and
nucleolar condensates, contributing to our understanding of a prominent conserved feature of
nuclear architecture (**Figure 6E**).

The multiphasic nature of the nucleolus has been elegantly demonstrated, where multiple,
immiscible layered phases form due to localized transcription and processing of rRNA, advective
flow or vectorial transport, and layer-specific protein-protein and protein-rRNA binding (Feric et
al. 2016; King et al. 2022; Lafontaine et al. 2020; Riback et al. 2023). Our study introduces a new
mechanism for layering PCH around the nucleolus in which some nucleolar components with
affinities for PCH can stabilize physical associations between the nucleolus and PCH. Further,
we identified the DEAD-Box RNA Helicase, Pitchoune, as such an amphiphilic protein that
localizes to the granular compartment of the nucleolus, contains a C-terminal PXVXL HP1a
binding motif, and importantly, is required for PCH-nucleolar 'surrounded' configurations.
Pitchoune's affinity for PCH is further supported by observing initial mixing with HP1a in cycle 14
and, later, the consistent formation of new spherical structures (neocondensates) within PCH in
nuclei lacking rDNA. It is important to emphasize that Pitchoune is not restricted only to the
nucleolar-PCH interface but is localized through the outer nucleolar layer, reminiscent of the
behavior of synthetic amphiphilic proteins at higher concentrations that form 3D multiphasic
protein condensates instead of monolayer films (Kelley et al. 2021). Together, we posit that PCH-
nucleolar condensate juxtaposition arises from the balance between Pitchoune nucleolar
associations and weaker interactions with HP1, and that Pitchoune's amphiphilic nature stabilizes
physical associations between immiscible condensates by simple minimization of the interfacial
energy. Further studies are required to determine if this phenomenon of amphiphilic protein-
mediated interactions between distinct condensates is generalizable.

The physical model accounted for the formation of the PCH void in -rDNA embryos through self-
association of the amphiphilic protein, combined with lower affinity binding to PCH. Although
Pitchoune self-association has not been demonstrated, oligomerization of orthologous RNA-

dependent DEAD-Box ATPases (DDXs) does promote the formation of RNA-rich condensates

(Overwijn & Hondele, 2023). Interestingly, Pitchoune neocondensates and the PCH void emerge

in –rDNA embryos at the same timepoint (cycle 17) when the canonical 'surrounded' PCH-

nucleolar conformation appears in +rDNA embryos. We thus hypothesize that Pitchoune's ability

to mediate PCH-nucleolar associations is altered at this developmental stage, perhaps via

increased self-association, abundance (beyond saturation), or increased affinity for HP1. Such

changes could elevate the energy cost of creating an interface with the nucleoplasm/solvent and

promote associations with PCH condensates by minimizing this interfacial energy. Finally, we

note the structural similarity between the PCH voids observed in -rDNA Drosophila embryos and

Nucleolus Precursor Bodies (NPBs) observed in mammalian oocytes and early embryos (Fulka

& Aoki 2016). Since they appear before the initiation of rRNA synthesis, we speculate that NPBs

are formed by DDX-rich neocondensates, as observed for Pitchoune in the absence of rDNA and

nucleoli.

Our research unveils the process of *de novo* co-assembly of the nucleolus and PCH at high spatial

and temporal resolution in the naïve chromatin landscape of a developing Drosophila embryo.

PCH and nucleolar condensates are nucleated independently and are initially spatially separated.

As they grow and coarsen (self-fuse), one end of PCH is tethered to the nucleolus via rDNA while

the rest spreads along the nuclear edge to form an 'extended' intermediate. Subsequently, PCH

from all chromosomes transitions to the canonical 'surrounded' configuration around the

nucleolus. Associations between uniquely mapped heterochromatin and the lamina (nuclear

periphery) have been well characterized (van Steensel & Belmont, 2017), and recent studies have

demonstrated overlaps between Nucleolus-associated domains (NADs) and Lamina-associated

domains (LADs) (Bersaglieri et al., 2022; Bizhanova et al., 2020; Vertii et al., 2019). Our work

uniquely captures the dynamics of the transitions of PCH between the nuclear and nucleolar

periphery during embryonic development. Although the precise mechanisms of these dynamics

are yet to be elucidated, we speculate that changes in chemical and biophysical properties, such

as varying nucleolar and PCH condensate volumes or material properties, increasing H3K9me2/3

levels, rRNA expression levels, or post-translational modifications of molecular factors like

Pitchoune, might alter binding affinities, influencing the balance of PCH's association with the

nucleolus versus the lamina.

The nucleolus has been recognized as a nuclear hub that attracts heterochromatin (Padeken &

Heun 2014; Quinodoz et al. 2018). Our study expands on this knowledge by revealing that the

nucleolus is required to distribute PCH condensates and repeats in the 3D nuclear space since

removing the nucleolus causes increased PCH compaction, potentially allowing PCH self-

attractions to dominate (Figure 3). The distribution of PCH repeats around the nucleolus is likely

functionally important, as the increased compaction could restrict critical nuclear processes like

PCH transcription, repair, or replication. Reciprocally, genetic depletion experiments in the early

Drosophila embryo revealed that the loss of H3K9me2/3 and HP1a causes rDNA decondensation

and nucleolar enlargement (Supplementary Figure 3). This result could indicate that PCH

condensates directly impact nucleolar structure and function. However, we favor the view that

nucleolar enlargement reflects the role of heterochromatin in silencing the transcription of subsets

of rDNA repeats and the partitioning of silent rDNA to nucleolus-associated heterochromatin

(Németh & Längst 2011; Pontvianne et al. 2013; Santoro et al. 2002; Van Koningsbruggen et al.

2010). This does not preclude the possibility that the surrounding PCH has independent roles in

nucleolar functions, such as impact on ribosome export. Together, we propose a model where

PCH and nucleoli connected via a chromatin polymer (rDNA) regulate each other's spatial

organization; the nucleolus distributes the PCH in the nucleus while the PCH constrains nucleolar

size by limiting rDNA transcription and may impact other nucleolar functions.

Finally, we predict that during aging and senescence, and perhaps other disease states, the

balance of interactions between PCH and nucleoli is perturbed to cause cellular phenotypes such

as enlarged nucleoli (Buchwalter & Hetzer 2017), decreased HP1a and heterochromatin (Larson

et al., 2012), and the dissociation of PCH repeats from the nucleolus (Dillinger et al. 2017). While

these cellular aging phenotypes are correlated, our model predicts a causative relationship

between PCH and nucleolar changes, which remains to be tested. Additionally, our findings

demonstrate how disrupting condensate nucleation sites or interaction hierarchies can form new,

abnormal nuclear structures or 'neocondensates' through inherent self-associations or secondary

interactions with other molecules. Removing rDNA causes Modulo to become broadly dispersed

in the nuclear volume, while Fibrillarin forms a spherical body separated from the PCH, and a

Pitchoune neocondensate transforms the compacted PCH into a shell. Since the cell contains

several multi-component condensates, our results may have broader relevance since perturbing

one set of interactions could give rise to new condensates/aggregates that cause defective

cellular phenotypes or behaviors. Understanding these outcomes will also be relevant in cellular

stress responses when new condensates are often formed or the composition of existing

condensates changes (Alberti & Hyman 2021).

MATERIALS AND METHODS

Reagent or Resource	**Source**	**Identifier**
446 **Antibodies**		
447 Rabbit anti-Fibrillarin	Abcam	ab5821
448 Mouse anti-H3K9me2	Abcam	ab8898
449 Mouse anti-Modulo	Gift from Mellone Lab (gift of Jacques Pradel, Chin-Chi 450 Chen et al., 2012)	
451 Goat-anti-Mouse, Alexa Fluor 488	Invitrogen	A-11001
452 Goat-anti-Mouse, Alexa Fluor 568	Invitrogen	A-11004
453 Goat-anti-Rabbit, Alexa Fluor 488	Invitrogen	A-11034
454 Donkey-anti-Rabbit, Alexa Fluor 568	Invitrogen	A-10042
455 **Bacterial Strains**		
456 DH5α Competent Cells	This study	N/A
457 **Chemicals**		
458 DAPI	Sigma	D9542-10MG
459 Triton-X	Sigma	T8787
460 Commercial Bleach	Clorox	N/A
461 Formaldehyde Solution	Sigma	F8775
462 Halocarbon Oil 27	Sigma	H8773
463 Gas-permeable Lumox Film	Sarstedt	Catalog # 94.6077.317
464 Heptane	Sigma	246654
465 Methanol	VWR (EMD)	EM-MX0485-9
466 Bovine Serum Albumin	VWR	0332
467 VECTASHIELD® Antifade Mounting 468 Medium	Vector Laboratories	H-1000-10
469 Atto 488 NHS Ester	Sigma	41698
470 Propidium Iodide Solution	Invitrogen	P3566
471 **Critical Commercial Assays**		
472 FISH Tag™ DNA Multicolor Kit, 473 Alexa Fluor™ dye combination	Thermo Fisher	F32951
474 pGEM T-Easy Cloning Kit	Promega	A1360
475 **Recombinant DNA**		
476 pGEM-rDNA	This study	N/A
477 **FISH Probes**		

478	359bp	This study	N/A
479	1.686	This study	N/A
480	AAGAG	This study	N/A
481	rDNA	This study	N/A

482 Experimental Models and Subject Details

483	**Fly Genotype**	**Source**
484	Oregon-R	BDSC # 5
485	RFP-HP1a (2nd Chr)	Lipsick Lab (Wen et al., 2008)
486	GFP-HP1a (3rd Chr)	Lipsick Lab (Wen et al., 2008)
487	GFP-Fibrillarin	Weischaus Lab (Falahati and Wieschaus, 2017)
488	RFP-Fibrillarin	Weischaus Lab (Falahati et al., 2016)
489	GFP-Modulo	Weischaus Lab (Falahati and Wieschaus, 2017)
490	Pitchoune-GFP	Weischaus Lab (Falahati and Wieschaus, 2017)
491	GFP-Rpl135	Weischaus Lab (Blythe et al., 2015)
492	Su(var)3-906/ TM3Ser	Reuter Lab (Tschiersch et al., 1994)
493	Su(var)3-917/ TM3Ser	Reuter Lab (Tschiersch et al., 1994)
494	Mat alpha GAL4	BDSC # 7063
495	HP1a RNAi	BDSC # 33400
496	FM6/C(1)DX, y[*] f[1]	BDSC # 784
497	C(1)RM/C(1;Y)6,y[1]w[1]f[1]/0	BDSC # 9460
498	Eyeless-GAL4	BDSC # 5534
499	Pitchoune RNAi VAL20	BDSC # 80368
500	Pitchoune RNAi VAL22	BDSC # 43984

501 Primers

502 rDNA FISH: 5'-ACGGTTGTTTCGCAAAAGTT-3' and 5'-TGTTGCGAAATGTCTTAGTTTCA-3'

503 Software and Algorithms

504	ImageJ (Fiji)	NIH	
505	ZEN Zeiss	Zeiss	https://www.zeiss.com/microscopy/int/products/microscope-software/zen.html
507	Arivis Vision 4D	Arivis	https://www.arivis.com/
508	GraphPad Prism	GraphPad	https://www.graphpad.com/

Method Details

Drosophila Stocks and Genetics

All crosses were maintained at 25° C. To visualize the dynamics of HP1a and nucleolar assembly, live embryos from stocks containing *RFP-HP1a; GFP-Fib*, *RFP-Fib; GFP-HP1a* and *RFP-HP1a; GFP-Mod* were imaged. Embryos lacking rDNA were obtained as described in Falahati () by crossing C(1)DX/Y; *RFP-HP1a; GFP-Fib* or C(1)DX/Y; *RFP-HP1a; Pit-GFP* or C(1)DX/Y; *RFP-Fib; Pit-GFP* virgins to C(1;Y)6,y[1]w[1]f[1]/0 males. 1/4th of the resulting embryos from this cross lack rDNA, and -rDNA embryos were selected based on the presence of Fibrillarin neocondensates in live and fixed embryos and DAPI morphology in fixed embryos.

To knockdown HP1a in Drosophila embryos, we crossed matα-GAL4 (BDSC #7063) females to HP1a RNAi males (BDSC #33400) to generate an F1 generation with HP1a depleted in the female germlines. F1 siblings were crossed to collect embryos depleted for HP1a, most of which exhibit severe developmental defects (Park et al. 2019). We selected embryos that reached nuclear cycle 14 for analyses of nucleolar morphology. The study of the effects of HP1a knockdown on the nucleolus was also limited to cycle 14 since most embryos were arrested by this stage. To analyze nucleolar phenotypes in Su(var)3-9^{null} embryos, we crossed Su(var)3-9^{06} /TM3Ser virgins to Su(var)3-9^{17} /TM3Ser males. After selecting Su(var)3-9$^{06/17}$ adults in the F1 generation, we intercrossed these siblings to collect Su(var)3-9$^{06/17}$ maternal null embryos. However, H3K9me2/3 levels partially recover by the gastrula stage (cycle 15) (through the action of other H3K9me2/3 methyltransferases like Eggless (Seller et al. 2019)), which limited this analysis to cycle 14. To knockdown Pitchoune in eye discs, eyeless-GAL4 virgin females were crossed with Pitchoune RNAi VAL20 males, and eye discs were dissected from F1 third instar larvae.

Live Imaging of Drosophila Embryos

To collect Drosophila embryos for live imaging, males and females of the desired genotype were added to a plastic cage covered with apple juice agar plates and left for at least 3 days at 25°C. On the day of imaging, a fresh plate was added to the cage, and embryos were collected for two hours. After removing the plate, embryos were coated with Halocarbon oil 27 (Sigma) for staging using a dissection scope with transillumination, then stage-selected embryos were placed on a 2 in x 2 in paper towel square and dechorionated in 50% bleach for 1 min. Bleach was wicked off with a Kimwipe after 1 min, the square was washed with a small amount of distilled water, and excess water was wicked off. Dechorionated embryos were secured onto a semipermeable membrane (Lumox film, Starstedt) on a membrane slide holder using Heptane Glue. These embryos were mounted in Halocarbon oil 27 (Sigma) between the membrane and a coverslip.

Live imaging was performed on a Zeiss LSM880 Airy Scan microscope (Airy Fast mode) with a
63X oil immersion objective at room temperature.

Immunostaining
Embryos were collected on apple juice-agar plates and aged till the appropriate stage,
dechorionated in 50% bleach, fixed in 1:1 heptane:4% formaldehyde in 1XPBS for 25 mins,
devitellinized in a 1:1 mixture of methanol:heptane, and stored at -20°C in methanol. Embryos
were rehydrated by washing in 1xPBS+0.2% Triton (PBT), blocked for ~1hr with 2% BSA in PBT,
then incubated with the primary antibody at 4°C overnight. Following washes with PBT, embryos
were incubated with the appropriate secondary antibody at room temperature for 2 hrs, stained in
DAPI, and mounted onto a slide using VectaShield (Vector Laboratories) mounting medium. All
primary antibodies were used at 1:250 dilution and secondaries at 1:1000. Imaging was
performed on a Zeiss LSM880 Airy Scan microscope (Airy Fast mode) with 63X oil immersion
objective at room temperature.

DNA Fluorescent in situ hybridization (FISH) and combined Immuno-FISH
<u>Probe Labelling</u>
A probe for ITS-1 rDNA was prepared by amplifying an ~800 bp fragment by PCR from genomic
DNA   using   the   primers   5'-ACGGTTGTTTCGCAAAAGTT-3'   and   5'-
TGTTGCGAAATGTCTTAGTTTCA-3' that was cloned into a pGEM-Teasy vector (Promega).
Using this plasmid as a template, ITS-1 rDNA probe was labeled with Alexa 488, Alexa 555, or
Alexa 648 using the FISH Tag™ DNA Multicolor Kit (ThermoFisher) following the manufacturer's
protocol. Locked nucleic acid (LNA) oligonucleotides (Integrated DNA technologies) conjugated
with Cy5 or FAM were used as probes for 359bp, 1.686, and AAGAG satellite DNA repeats.
<u>Hybridization</u>
Embryos were collected on apple juice-agar plates and aged till the appropriate stage,
dechorionated in 50% bleach, then fixed in 1:1 heptane: 4% formaldehyde in 1XPBS for 25 mins,
devitellinized in a 1:1 mixture of methanol: heptane, and stored in -20°C. Embryos were washed
in 2xSSC-T (2xSSC containing 0.1% Tween-20) with increasing formamide concentrations (20%,
40%, then 50%) for 15 min each. 100ng of DNA probes in 40 µl of hybridization solution (50%
formamide, 3× SSCT, 10% dextran sulfate) was added, denatured together with the embryos at
95°C for 5 min and incubated overnight at 37°C. Following hybridization, embryos were washed
twice in 2xSSCT for 30 mins at 37°C and thrice in PBT for 5 mins at room temperature. After
completing washes, embryos were stained in DAPI and mounted onto a slide using VectaShield
(Vector Laboratories) mounting medium.

<u>Combined Immuno-FISH</u>

For combined in situ detection of proteins and DNA sequences, immunofluorescence was

performed first, embryos were post-fixed in 4% formaldehyde for 25 mins, then processed for

FISH.

Pan-Protein Staining using 488 NHS Ester

Formaldehyde-fixed embryos were devitellinized in a 1:1 mixture of methanol: heptane and stored

in methanol. Embryos were rehydrated by washing in 1xPBS+0.2% Triton (PBT). After washing

off methanol, embryos were stained in the diluted 488 NHS ester fluorophore (1:50) from a

10mg/ml stock in 0.1% PBST for 6 h at 4 °C followed by washing in PBT three times for 30 mins

each at room temperature. Embryos were stained in DAPI for ten minutes and mounted onto a

slide using VectaShield (Vector Laboratories) mounting medium.

Propidium Iodide Staining

Formaldehyde-fixed and Heptane devitellinized embryos were rehydrated by washing in

1xPBS+0.2% Triton (PBT). Samples were equilibrated in 2X SSC. RNase-treated controls alone

were incubated in 100 µg/mL DNase-free RNase in 2X SSC for 20 minutes at 37°C. After washing

away the RNase with 2X SSC, embryos were incubated in 500nM of Propidium Iodide in 2X SSC

for 10 mins at room temperature. Samples were rinsed in 2X SSC, stained with DAPI, and

mounted on a slide with VectaShield (Vector Laboratories) mounting medium.

Modeling

To better understand the association of PCH with the nucleolus, we developed a physical model

that simulates the interactions between different types of molecules found in these biomolecular

condensates. In our physical model, we simulate four components of the nucleus: PCH (H) and

ribosomal DNA (rD) as long polymers, and Fibrillarin (F) and an amphiphilic protein (X) as

independent, single monomers, which we hereafter refer to as beads. Since the experiments

focused on the PCH domain of Drosophila, our physical model only simulates this specific region

of the genome (30% of the genome), not the entire genome. This allows us to study the dynamics

of this particular region of the genome more accurately and efficiently. PCH is modeled as a

semiflexible bead-spring polymer chain in which N beads are connected by N-1 harmonic springs.

Each bead of the chain represents a cluster of PCH containing approximately 5 kilo base pairs of

DNA, with a diameter of approximately 30 nm. The semi-flexibility of the chain is determined by

its persistence length, which is taken to be 60 nm (2 beads) in accordance with previous studies

that indicate chromatin has a persistence length between 50 to 100 nm (Wachsmuth et al. 2016).

We represent rDNA as a self-avoiding chain that occupies approximately 20% of the middle
domain of PCH. Fibrillarin and the amphiphilic protein X are modeled using single, diffusive beads
where each protein has distinct interactions with the polymer and other proteins.

To simplify the model, we assume that the size of each protein bead is equal to the size of the
heterochromatin bead, with both having a diameter σ. The non-bonding interactions between
polymer-polymer, protein-protein, and polymer-protein are modeled using a standard Lennard-
Jones (LJ) potential. The LJ potential is truncated at the distance 2.5 σ, meaning that the
interaction between the beads is only non-zero if they are within 2.5 σ distance. At very short
distances between the beads, the LJ potential is strongly repulsive (representing the excluded
volume of two molecules). At intermediate spacings, the LJ potential is attractive, with a strength
adjusted to model the different states of chromatin, such as its compaction or decondensation
and the phase-separating tendency of the proteins. In our model, the polymer and the proteins
are confined within a spherical boundary that represents the nucleus of the cell. This boundary
mimics the effect of the nuclear envelope, which constrains the movement of these beads and
affects their interactions with each other. In our study, we used the LAMMPS (Large-scale
Atomic/Molecular Massively Parallel Simulator) package to simulate the behavior of our
biomolecular system (Thompson et al. 2022). LAMMPS uses Brownian dynamics, which accounts
for the viscous forces acting on the beads, and a stochastic force (Langevin thermostat) to ensure
that the system of beads and solvent is maintained at a constant temperature (NVT ensemble).
This allows us to model the interactions between polymer-polymer, polymer-protein, and protein-
protein beads accurately and study the behavior of the system over time.

Rationale for the choice of parameters in the coarse-grained model
To analyze the experimental observations of phase separation in the nucleus, we study a minimal
model with four crucial components: PCH (H), rDNA (rD), Fibrillarin (F), and an amphiphilic protein
(X) that binds both nucleolar and PCH components. The parameters in the simulations include
the number of molecules of each component α (N_α), the strength of the attractions between two
components (denoted by indices β and γ, $\epsilon_{\beta\gamma}$), and the size of the confinement (R_c). The fraction
of the nucleus that is hydrated (does not contain PCH or the other proteins) is obtained from the
relative difference between the confinement volume and the volumes of PCH and the other
proteins.
<u>Bonding potential between monomers in the polymer made of PCH and rDNA:</u> Adjacent beads
on the polymer chain are interconnected by harmonic springs using the potential function:

$$639 \qquad V_s = \sum_{i=1}^{N-1} k_s (r_i - \sigma)^2$$

Here, r_i represents the distance between the i-th and $(i + 1)$-th beads. The spring constant and

equilibrium distance between neighboring beads are denoted as k_s and σ respectively. In our

simulations, the spring constant k_s is set to $100 k_B T / \sigma^2$ to ensure the presence of rigid bonds

between adjacent beads of the polymer chain.

<u>Attraction strength</u> ($\epsilon_{\beta\gamma}$): The Lennard-Jones potential is used to model the attraction between

any two non-bonded beads:

$$647 \qquad V_{\beta\gamma}(r) = \begin{cases} 4\epsilon_{\beta\gamma} \left[\left(\frac{\sigma}{r}\right)^{12} - \left(\frac{\sigma}{r}\right)^6 \right] & \text{for } r \le r_c \\ 0 & \text{for } r > r_c \end{cases}$$

Here, the symbol $\epsilon_{\beta\gamma}$ represents the attraction strength between beads of type β and γ, where

$\beta, \gamma \in \{H, rD, F, X\}$ represent PCH, rDNA, Fibrillarin, and amphiphilic protein, respectively. For

instance, ϵ_{FX} represents the attraction strength between Fibrillarin and amphiphilic protein beads.

When dealing with attractive interactions between chromatin-chromatin and chromatin-protein

beads, a distance cutoff of $r_c = 2.5\sigma$ is used for the Lennard-Jones potential, beyond which the

interaction is set to zero. To account for only excluded volume interactions (with no attractions)

using the same potential, a cutoff of $r_c = 2^{1/6}\sigma$ and $\epsilon = 1 k_B T$ are employed. This choice is made

because the potential energy is at its minimum at that point, and the resulting force on a bead is

zero. As there are four components in our model, there are a total of 10 combinations ((n(n+1)/2),

where n is number of components) of attraction strength parameters between the different

components.

<u>Confinement size:</u> The size of the confinement is determined by defining the volume fraction of

chromatin ϕ:

$$661 \qquad \phi = N_G \times \frac{\text{volume of 1 bead}}{\text{volume of confinement}} = N_G \times \frac{(\frac{4}{3}\pi\sigma^3)}{(\frac{4}{3}\pi R_c^3)}$$

Here, N_G represents the total number of beads in the Drosophila genome, where each bead

corresponds to 5 kilobase pairs (kbps) of DNA. The diameter of a spherical bead, denoted as σ,

is taken to be 30 nm (see Bajpai & Safran, 2023 for further explanation). The total length of the

diploid Drosophila genome is 360 Mbps, and N_G can be calculated by dividing the Drosophila

genome length by the amount of DNA represented by one bead (5 kbps). The volume fraction of

chromatin (ϕ) is commonly reported as ~ 0.1 in existing literature (Qi & Zhang, 2021; Tripathi &

Menon, 2019). Using the equation above, the calculated radius of the confinement (R_c) is found
to be 45σ.

<u>Number of molecules in PCH and rDNA:</u> We model the PCH domain as a polymer chain
composed of $N = 10,000$ beads, which represents approximately 30% of the beads in the entire
genome. Among these 10,000 beads, 20% are designated as rDNA, resulting in a total of 4,000
rDNA beads. We do not explicitly model the rest of the genome since the experiments show that
the nucleolar components are localized near PCH.

<u>Concentration of Fibrillarin and amphiphilic protein:</u> The concentration of Fibrillarin and
amphiphilic protein is calculated as follows:

677
$$\text{concentration } (c_\alpha) = \frac{N_\alpha}{\text{volume of confinement}}$$

Here, the symbol α represents the type of protein, where $c_\alpha = c_F$ for the fibrillarin concentration
and $c_\alpha = c_X$ for the amphiphilic protein concentration. After defining the parameters and obtaining
the value for the confinement radius parameter from the assumed volume fraction of PCH, we
proceed with an initial simulation, focusing on a single component, namely the Fibrillarin protein,
within the confinement. During this simulation, we vary the concentration (c_F) and the attraction
strength (ϵ_{FF}) of Fibrillarin. Our experimental results demonstrate that Fibrillarin undergoes phase
separation independently of the other components (rDNA or PCH) during cycle 14 (refer to Figure
2A). These initial-stage simulation results yield a phase diagram, which indicates that a minimum
attraction strength of $1.3 - 2.0k_\text{B}\text{T}$ is required to condense fibrillarin particles within the
concentration range of $c_F = 0.0013 - 0.013$. The reported concentration of nucleolar particles is
$c = 0.015$ (Qi & Zhang, 2021). Consequently, we maintain a fixed concentration of Fibrillarin at
$c_F = 0.013$ when simulating all the other protein components (**Extended Figure 5A**). Since the
concentration of amphiphilic protein is not determined from the experiments, we conducted
multiple simulations, systematically varying the amphiphilic protein concentration (c_X) within the
range of $0.0013 - 0.13$. The results of these simulations are discussed in **Figure 6A**.

<u>Parameter range of ϵ_{HH}:</u> To understand the behavior of each component separately (before
including interactions between different components) in the next stage, we conducted simulations
specifically focusing only on the PCH chain. The PCH chain represents a condensed chromatin
region within the nucleus which implies self-attractive interactions of the beads representing the
polymer. During these simulations, we varied the attraction strength between PCH beads (ϵ_{HH})
from 0 to $0.5k_\text{B}\text{T}$. Our results revealed that within the range of $\epsilon_{HH} = 0.35 - 0.5 \ k_\text{B}\text{T}$ (**Extended**
Figure 5B), the PCH chain underwent collapse, resulting in a condensed conformation that is
phase-separated from the aqueous component of the system (not simulated explicitly).

<u>Parameter value of</u> ϵ_{rD-F}: We next simulate the self-organization due to the interactions between

the three components (where each component so far was considered alone): PCH, rDNA, and

Fibrillarin. We set $\epsilon_{HH} = 0.35\,k_\mathrm{B}T$ and $\epsilon_{FF} = 2\,k_\mathrm{B}T$, incorporating only excluded-volume

interactions (hard-core, repulsive interactions) between H-F, i.e., there is no direct attraction

between PCH and Fibrillarin as implied by the experiments. By varying the attraction strength

between rDNA and Fibrillarin (ϵ_{rD-F}), we made the following observations based on our

simulation results: a weaker attraction (and $\epsilon_{rD-F} = 0.75\,k_\mathrm{B}T$) resulted in rDNA wrapping around

the condensed Fibrillarin phase, while a stronger attraction (and $\epsilon_{rD-F} = 2\,k_\mathrm{B}T$) led to the

condensation of rDNA within the Fibrillarin complex. This latter observation aligns with our

experimental findings. Therefore, we select $\epsilon_{rD-F} = 2\,k_\mathrm{B}T$ as the parameter value in the

subsequent simulations (**Extended Figure 5C**).

<u>Parameter ranges of</u> ϵ_{FF} and ϵ_{XX}: Finally, we introduce the fourth component, an amphiphilic

protein 'X', which we suggest may interact attractively with both PCH and Fibrillarin. Initially, we

investigated the relative attraction strengths between Fibrillarin and protein X when considering

the same concentration for both. We explore all possible combinations of ϵ_{FF} both greater than

and less than ϵ_{XX}. For $\epsilon_{XX} \geq \epsilon_{FF}$, we observe PCH surrounding the amphiphilic protein-rich phase,

but just a partial wetting between the amphiphilic protein-rich phase and the Fibrillarin-rich phase

(**Extended Figure 5D**). Only when $\epsilon_{XX} < \epsilon_{FF}$ do we observe PCH surrounding the amphiphilic

protein-rich phase, which in turn surrounds the Fibrillarin condensate, consistent with the

experimental results in wild type embryos.

<u>Parameter range of</u> ϵ_{FX}: We proceeded to vary the attraction strength between Fibrillarin and

amphiphilic protein X within the range of $\epsilon_{FX} = 1.5\,k_\mathrm{B}T$. When the attraction strength is relatively

low ($\epsilon_{FX} \leq 0.75\,k_\mathrm{B}T$), the fibrillarin-rich phase and the amphiphilic protein-rich phase do not

associate with each other. At moderate attraction strengths ($\epsilon_{FX} \geq 1\,k_\mathrm{B}T$ and ($\epsilon_{FX} \leq 1.25\,k_\mathrm{B}T$), the

Fibrillarin-rich and amphiphilic protein-rich phases partially wet each other. Finally, at higher

attraction strengths ($\epsilon_{FX} = 1.5\,k_\mathrm{B}T$), the amphiphilic protein completely wets Fibrillarin (**Extended**

Figure 5E-F).

<u>Parameter range of</u> c_X: Additionally, we explored variations in the concentration of amphiphilic

protein X (c_X). Notably, for higher concentrations of protein X ($c_X = 0.005 - 0.013$), we observed

that heterochromatin tends to completely engulf the amphiphilic protein-rich phase, which in turn

engulfs Fibrillarin (**Figure 6A**).

Quantitative Image Analysis

3D measurements were performed using Arivis Vision4D (Zeiss) while ImageJ was used for all 2D measurements. The details of each analysis pipeline used in this study are listed below:

Measuring the fraction of the nucleolar edge occupied by HP1a or H3K9me2

Figure 1B: A cluster of nuclei was manually chosen for analyses ~15-30 mins into the specified interphase and defined as "Early Cycle", while those observed from ~50-70 mins were defined as "Late Cycle". Preprocessing steps include background subtraction and denoising using Gaussian blur. HP1a and Fibrillarin were segmented in 3D using the "Intensity Threshold Segmenter" with the Auto segmentation method. To calculate the fraction of the nucleolar edge occupied by HP1a, the nucleolus was dilated by 1 pixel and subtracted from the dilated object to create a 1-pixel shell around the nucleolus. The nucleolus shell was intersected with HP1a segments to calculate the volume of the nucleolar shell that overlaps with HP1a.

Figure 6D: Same as above, except the nucleolus was dilated by 2 pixels to create a 2-pixel shell around the nucleolus. The shell was intersected with H3K9me2 segments in nuclei from eye-discs to calculate the fraction of the nucleolar shell that overlaps with H3K9me2.

Measuring Distances

Extended Figure 1F: Preprocessing steps include background subtraction and denoising using Gaussian blur. 1.686 or 359bp foci and Fibrillarin were manually segmented using the "Intensity Threshold Segmenter" with the Simple segmentation method. The distance between a Fibrillarin segment and its nearest 1.686 or 359bp locus was measured in 3D using the "Distances" feature in Arivis.

Figure 2B: HP1a and Fibrillarin were segmented using the "Intensity Threshold Segmenter" with the Auto segmentation method. The distance between HP1a and its nearest Fibrillarin segment was measured in 3D using the "Distances" feature in Arivis.

Figure 2F: AAGAG and 1.686 were segmented using the "Intensity Threshold Segmenter" with the Auto segmentation method. The distance between AAGAG and its nearest 1.686 segment was measured in 3D using the "Distances" feature in Arivis.

Measuring Aspect Ratio of HP1a

Figure 2D: Individual nuclei were manually selected 15 mins after the start of Cycle 15. Preprocessing steps include background subtraction and denoising using Gaussian blur. HP1a was segmented using Auto thresholding using Otsu. The aspect ratio of the segment was determined using the Analyze Particles feature in Fiji.

<u>Line Scans:</u>

<u>Figure 3B:</u> Line scans were generated using the "Plot profile" feature in Fiji. The highest intensity

values in the green and magenta channels were each set to 1.0, and the remaining intensities

were normalized to it.

<u>Pitchoune void formation measurements</u>

<u>Figure 5E-G:</u> To measure the dynamics of the formation of Pitchoune in -rDNA embryos,

maximum intensity projections of Amnioserosa were first preprocessed in Fiji using Subtract

background and a Gaussian Blur filter. Auto thresholding for each time point was performed using

the Yen method. Using Analyze Particles, the area, circularity, and mean intensity of each

segment of Pitchoune was extracted. The mean intensity over time was normalized to its value at

T=0.

<u>Measuring Volume</u>

<u>Supplementary Figure 3:</u> Preprocessing steps include background subtraction and denoising

using Gaussian blur. The volume of segments was extracted using Arivis after Fibrillarin was

segmented in 3D using the "Intensity Threshold Segmenter" with the Auto segmentation method

(Otsu feature).

Statistical Analysis

Data were plotted, and statistical analyses were performed using GraphPad Prism8. P-values

were calculated using unpaired two-tailed t-tests.

Data availability statement

All data supporting the results of this study are included in the manuscript. Reagents used in this

study are available upon request.

Code availability statement

All the simulations, analysis, and visualization in this study were conducted using publicly

available software packages or custom codes. We utilized LAMMPS (version: 23 June 2022) for

performing Langevin Dynamics simulations and Ovito (version 3.7.11) for visualization. Custom

codes were employed for all other analyses. All code and raw data will be deposited on Zenodo

(https://zenodo.org/).

ACKNOWLEDGEMENTS

We thank Eric Weischaus (Princeton University) and Shelby Blythe (Northwestern University) for generously sharing the fluorescently tagged nucleolar fly lines. We are grateful to Karpen Lab members for their helpful discussions and to colleagues Lucy Brennan, Coral Zhou, and Amy Strom for their valuable feedback on this manuscript. The Karpen lab acknowledges critical support from the National Institutes of Health (R35GM139653), and SS and GHK are grateful for the support of the Volkswagen Stiftung (98196).

AUTHOR CONTRIBUTIONS

SR: conceptualization, experimentation, data acquisition, formal analysis, validation, investigation, methodology, and writing

OA: conceptualization, investigation, formal analysis, and writing

GB: simulations, data acquisition, formal analysis, methodology, and writing

KL: experimentation and data acquisition

SC: reagent preparation and formal analysis

SS: writing and funding acquisition

GHK: conceptualization, writing, and funding acquisition

COMPETING INTERESTS

No competing interests to declare.

REFERENCES

Ahmad K, Henikoff S. 2022. The Drosophila embryo as a tabula rasa for the epigenome.
Fac Rev. 11(40):

Alberti S, Hyman AA. 2021. Biomolecular condensates at the nexus of cellular stress,
protein aggregation disease and ageing. *Nature Reviews Molecular Cell Biology 2021*
22:3. 22(3):196–213

Ballmer D, Tardat M, Ortiz R, Graff-Meyer A, Ozonov EA, et al. 2023. HP1 proteins
regulate nucleolar structure and function by secluding pericentromeric constitutive
heterochromatin. *Nucleic Acids Res*. 51(1):117–43

Berry J, Weber SC, Vaidya N, Haataja M, Brangwynne CP, Weitz DA. 2015. RNA
transcription modulates phase transition-driven nuclear body assembly. *Proc Natl*
Acad Sci U S A. 112(38):E5237–45

Bizhanova A, Kaufman PD. 2021. Close to the edge: Heterochromatin at the nucleolar and
nuclear peripheries. *Biochim Biophys Acta Gene Regul Mech*. 1864(1):194666

Brangwynne CP, Mitchison TJ, Hyman AA. 2011. Active liquid-like behavior of nucleoli
determines their size and shape in Xenopus laevis oocytes. *Proceedings of the*
National Academy of Sciences. 108(11):4334–39

Buchwalter A, Hetzer MW. 2017. Nucleolar expansion and elevated protein translation in
premature aging. *Nat Commun*

Dillinger S, Straub T, Németh A. 2017. Nucleolus association of chromosomal domains is
largely maintained in cellular senescence despite massive nuclear reorganisation.
PLoS One. 12(6):e0178821

Ditlev JA, Case LB, Rosen MK. 2018. Who's In and Who's Out—Compositional Control of
Biomolecular Condensates. *J Mol Biol*. 430(23):4666–84

Eissenberg JC, Elgin SC. 2000. The HP1 protein family: getting a grip on chromatin. *Curr*
Opin Genet Dev. 10(2):204–10

Falahati H, Pelham-Webb B, Blythe S, Wieschaus E. 2016. Nucleation by rRNA dictates
the precision of nucleolus assembly. *Current Biology*. 26(3):277–85

Falahati H, Wieschaus E. 2017. Independent active and thermodynamic processes govern
the nucleolus assembly in vivo. *Proceedings of the National Academy of Sciences*.
114(6):1335–40

Fare CM, Villani A, Drake LE, Shorter J. 2021. Higher-order organization of biomolecular
condensates. *Open Biol*. 11(6):210137

Farrell JA, O'Farrell PH. 2014. From Egg to Gastrula: How the Cell Cycle Is Remodeled
During the Drosophila Mid-Blastula Transition. *Annu Rev Genet*. 48(1):269–94

Fefelova EA, Pleshakova IM, Mikhaleva EA, Pirogov SA, Poltorachenko VA, et al. 2022.
Impaired function of rDNA transcription initiation machinery leads to derepression of

ribosomal genes with insertions of R2 retrotransposon. *Nucleic Acids Res*. 50(2):867–
84

Feric M, Vaidya N, Harmon TS, Mitrea DM, Zhu L, et al. 2016. Coexisting Liquid Phases
Underlie Nucleolar Subcompartments. *Cell*. 165(7):1686–97

Fulka H, Aoki F. 2016. Nucleolus precursor bodies and ribosome biogenesis in early
mammalian embryos: Old theories and new discoveries. *Biol Reprod*. 94(6):143–44

Holmberg Olausson K, Nistér M, Lindström MS. 2014. Loss of nucleolar histone chaperone
NPM1 triggers rearrangement of heterochromatin and synergizes with a deficiency in
DNA methyltransferase DNMT3A to drive ribosomal DNA transcription. *Journal of*
Biological Chemistry. 289(50):34601–19

Janssen A, Colmenares SU, Karpen GH. 2018. Heterochromatin: Guardian of the
Genome. *Annu Rev Cell Dev Biol*. 34(1):265–88

Keenen MM, Brown D, Brennan LD, Renger R, Khoo H, et al. 2020. HP1 proteins compact
DNA into mechanically and positionally stable phase separated domains. . 1–30

Kelley FM, Favetta B, Regy RM, Mittal J, Schuster BS. 2021. Amphiphilic proteins
coassemble into multiphasic condensates and act as biomolecular surfactants. *Proc*
Natl Acad Sci U S A. 118(51):e2109967118

King MR, Lin AZ, Ruff KM, Farag M, Ouyang W, et al. 2022. Uncovering molecular
grammars of intrinsically disordered regions that organize nucleolar fibrillar centers.
bioRxiv. 2022.11.05.515292

Lacy ME, Hutson MS. 2016. Amnioserosa development and function in Drosophila
embryogenesis: Critical mechanical roles for an extraembryonic tissue.
Developmental Dynamics. 245(5):558–68

Lafontaine DLJ, Riback JA, Bascetin R, Brangwynne CP. 2020. The nucleolus as a
multiphase liquid condensate. *Nat Rev Mol Cell Biol*. 22(3):165–82

Larson AG, Elnatan D, Keenen MM, Trnka MJ, Johnston JB, et al. 2017. Liquid droplet
formation by HP1α suggests a role for phase separation in heterochromatin. *Nature*.
547(7662):236–40

Larson K, Yan SJ, Tsurumi A, Liu J, Zhou J, et al. 2012. Heterochromatin formation
promotes longevity and represses ribosomal RNA synthesis. *PLoS Genet*

Lieberman-Aiden E, Van Berkum NL, Williams L, Imakaev M, Ragoczy T, et al. 2009.
Comprehensive mapping of long-range interactions reveals folding principles of the
human genome. *Science*. 326(5950):289–93

Lyon AS, Peeples WB, Rosen MK. 2021. A framework for understanding functions of
biomolecular condensates on molecular to cellular scales. *Nat Rev Mol Cell Biol*.
22(3):215

M'Saad O, Bewersdorf J. 2020. Light microscopy of proteins in their ultrastructural context.
Nature Communications 2020 11:1. 11(1):1–15

Németh A, Conesa A, Santoyo-Lopez J, Medina I, Montaner D, et al. 2010. Initial
Genomics of the Human Nucleolus. *PLoS Genet*. 6(3):e1000889

Németh A, Längst G. 2011. Genome organization in and around the nucleolus. *Trends in*
Genetics. 27(4):149–56

Padeken J, Heun P. 2014. Nucleolus and nuclear periphery: Velcro for heterochromatin.
Curr Opin Cell Biol. 28(1):54–60

Padeken J, Mendiburo MJ, Chlamydas S, Schwarz H-J, Kremmer E, Heun P. 2013. The
Nucleoplasmin Homolog NLP Mediates Centromere Clustering and Anchoring to the
Nucleolus. *Mol Cell*. 50(2):236–49

Park AR, Liu N, Neuenkirchen N, Guo Q, Lin H. 2019. The Role of Maternal HP1a in Early
Drosophila Embryogenesis via Regulation of Maternal Transcript Production.
Genetics. 211(1):201

Peng JC, Karpen GH. 2007. H3K9 methylation and RNA interference regulate nucleolar
organization and repeated DNA stability. *Nat Cell Biol*. 9(1):25–35

Politz JCR, Scalzo D, Groudine M. 2013. Something Silent This Way Forms: The
Functional Organization of the Repressive Nuclear Compartment. *Annu Rev Cell Dev*
Biol. 29(1):241–70

Pontvianne F, Blevins T, Chandrasekhara C, Mozgová I, Hassel C, et al. 2013. Subnuclear
partitioning of rRNA genes between the nucleolus and nucleoplasm reflects
alternative epiallelic states. *Genes Dev*. 27(14):1545–50

Potapova TA, Gerton JL. 2019. Ribosomal DNA and the nucleolus in the context of
genome organization. . 109–27

Quinodoz SA, Ollikainen N, Tabak B, Palla A, Schmidt JM, et al. 2018. Higher-Order Inter-
chromosomal Hubs Shape 3D Genome Organization in the Nucleus. *Cell*. 174(3):744-
757.e24

Riback JA, Eeftens JM, Lee DSW, Kleiner R, Kriwacki R, et al. 2023. Viscoelasticity and
advective flow of RNA underlies nucleolar form and function. *Mol Cell*. 83:3095–3107

Santoro R, Li J, Grummt I. 2002. The nucleolar remodeling complex NoRC mediates
heterochromatin formation and silencing of ribosomal gene transcription. *Nature*
Genetics 2002 32:3. 32(3):393–96

Schotta G, Ebert A, Krauss V, Fischer A, Hoffmann J, et al. 2002. Central role of
Drosophila SU(VAR)3-9 in histone H3-K9 methylation and heterochromatic gene
silencing. *EMBO J*. 21(5):1121–31

Seller CA, Cho CY, O'Farrell PH, Armstrong RL, Duronio RJ, et al. 2019. Rapid embryonic
cell cycles defer the establishment of heterochromatin by eggless/SetDB1 in
Drosophila. *Genes Dev*. 33(7–8):403–17

Shin Y, Brangwynne CP. 2017. Liquid phase condensation in cell physiology and disease.
Science (1979). 357(6357):eaaf4382

Srivastava R, Srivastava R, Ahn SH. 2016. The Epigenetic Pathways to Ribosomal DNA
Silencing. *Microbiol Mol Biol Rev*. 80(3):545

Strom AR, Emelyanov A V., Mir M, Fyodorov D V., Darzacq X, Karpen GH. 2017. Phase
separation drives heterochromatin domain formation. *Nature*. 547(7662):241–45

Thiru A, Nietlispach D, Mott HR, Okuwaki M, Lyon D, et al. 2004. Structural basis of
HP1/PXVXL motif peptide interactions and HP1 localisation to heterochromatin.
EMBO J. 23(3):489

Thiry M, Lafontaine DLJ. 2005. Birth of a nucleolus: The evolution of nucleolar
compartments. *Trends Cell Biol*. 15(4):194–99

Thompson AP, Aktulga HM, Berger R, Bolintineanu DS, Brown WM, et al. 2022. LAMMPS
- a flexible simulation tool for particle-based materials modeling at the atomic, meso,
and continuum scales. *Comput Phys Commun*. 271:108171

Tortora MMC, Brennan LD, Karpen G, Jost D. 2023. HP1-driven phase separation
recapitulates the thermodynamics and kinetics of heterochromatin condensate
formation. *bioRxiv*. 2022.07.11.499635

Torza S, Mason SG. 1969. Coalescence of Two Immiscible Liquid Drops. *Science (1979)*.
163(3869):813–14

Van Koningsbruggen S, Gierliński M, Schofield P, Martin D, Barton GJ, et al. 2010. High-
Resolution Whole-Genome Sequencing Reveals That Specific Chromatin Domains
from Most Human Chromosomes Associate with Nucleoli. *Mol Biol Cell*. 21(21):3735

Wachsmuth M, Knoch TA, Rippe K. 2016. Dynamic properties of independent chromatin
domains measured by correlation spectroscopy in living cells. *Epigenetics Chromatin*.
9(1):1–20

Wang L, Gao Y, Zheng X, Liu C, Dong S, et al. 2019. Histone Modifications Regulate
Chromatin Compartmentalization by Contributing to a Phase Separation Mechanism.
Mol Cell. 1–14

Zaffran S, Chartier A, Gallant P, Astier M, Arquier N, et al. 1998. A Drosophila RNA
helicase gene, pitchoune, is required for cell growth and proliferation and is a potential
target of d-Myc. *Development*. 125(18):3571–84

EXTENDED FIGURES

Extended Figure 1: PCH reorganizes from the nuclear to the nucleolar edge during early Drosophila development. (**A**) Schematic of the locations of pericentromeric satellite repeats (359bp and 1.686) and rDNA repeats in Drosophila melanogaster chromosomes. Schematic of ribosomal DNA (rDNA) arrays in *Drosophila melanogaster* showing the position of the ITS-1 rDNA probe. (**B**) Combined immuno-FISH of Drosophila nuclei in Cycle 14 stained for Fibrillarin (green), DAPI (gray), (Left) ITS-1 rDNA (magenta), (Right) 359bp satellite DNA (magenta) on Chr X. Dashed white lines represent the edge of the nucleus determined by DAPI staining. (**C**) Combined immuno-FISH of Drosophila nuclei in Cycle 14 stained for Fibrillarin (green) and 1.686 satellite DNA (magenta) on Chr 2 and 3. (**D**) Representative stills of nuclei showing the conformation of the Fibrillarin and HP1a in progressive cycles 13-17. Nuclei marked with a dashed yellow boundary in the xy view are shown in their xz view below. (**E**) Immunofluorescence of H3K9me2 (magenta), Fibrillarin (green) and DAPI (gray) in nuclei from a Stage 8 (~Cycle 15) and eye disc from a third instar larva. (**F**) Combined immuno-FISH of 1.686 (magenta arrows), Fibrillarin (green arrow) and DAPI (grey) in a nucleus from epidermal cells of Stage 8 (~Cycle 15) and Stage 16 (Late) Drosophila embryos. (**G**) Distance between the centers of geometry of 1.686 and Fibrillarin in Stage 8 (~Cycle 15) and Stage 16 (Late) in Drosophila nuclei. n>60 loci (from 3 embryos) at each developmental stage.

Extended Figure 2: The nucleolus is surrounded by PCH in Drosophila nuclei. (**A**) Representative stills of nuclei from first instar larval epidermal cells (left), third instar larval eye disc (center), and S2 cell culture (right). Magenta, RFP-HP1a. Green, GFP-Fibrillarin. (**B**) Representative images of nucleoli showing the distribution of GFP-Modulo (green) and RFP-Fib (magenta) (top), and GFP-Rpl135 (green) and RFP-Fib (magenta) (bottom) in a live late (Stage 16) embryo. (**C**) Representative maximum intensity projections of PCH (RFP-HP1a, magenta) and nucleoli (GFP-Modulo, green) in live epidermal nuclei from a late Drosophila embryo (Stage 16, ~14-16hr). The nucleus within the white dashed box is magnified on the right and presented in xy and xz view. White dashed lines indicate the nuclear boundary.

Extended Figure 3: The nucleolus is enlarged and rDNA is decondensed when H3K9 methylation or HP1a is reduced. (**A**) Maximum intensity z projections of WT and Su(var)3-9[06/17] nuclei immunostained with H3K9me2 (green) and co-stained with DAPI (blue) at nuclear cycle 14. (**B**) Maximum intensity z projections of nuclei immunostained with Fibrillarin (green) and costained with DAPI (blue) in WT and Su(var)3-9[06/17] embryos at Cycle 14. (**C**) Quantification of the volume of Fibrillarin in WT and Su(var)3-9[06/17] embryos at Cycle 14. (**D**) Combined immuno FISH (maximum intensity z projections) of Fibrillarin (green), rDNA (magenta) and DAPI (blue) in WT and Su(var)3-9[06/17] embryos at Cycle 14. (**E**) Maximum intensity z projections of nuclei from wildtype and HP1a RNAi embryos at Cycle 14. Fibrillarin (green), rDNA (red), DAPI (blue)**.** (**F**) Volume of Fibrillarin in WT and HP1a RNAi in embryos at Cycle 14. (**G**) Schematic of the observed phenotypes depicting decondensation of rDNA and nucleolar enlargement when H3K9me2/3 or HP1a is decreased.

Extended Figure 4: The HP1a-void in -rDNA embryos does not stain for DAPI, Fibrillarin, Modulo or Propidium Iodide (RNA). (**A**) Representative images of fixed nuclei from wildtype late embryos and mutant embryos lacking rDNA stained for Modulo (magenta), Fibrillarin (green) and DAPI (blue). (**B**) Left: Representative images of fixed nuclei from Stage 14-16 (late) wildtype embryos and mutant embryos lacking rDNA showing Fibrillarin (green) and DAPI (blue). Right: Nuclei marked with white dashed outlines on the left are shown in the z plane. (**C**) H3K9me2 immunofluorescence (magenta) and DAPI (blue) staining in -rDNA nuclei in Stage 14-16 (late) Drosophila embryos. (**D**) Representative images of fixed nuclei from Stage 14-16 (late) wildtype and mutant embryos lacking rDNA stained with Propidium Iodide (green) and DAPI (blue) without (left) and with (right) RNaseA. The yellow arrow points to the RNA staining in the nucleolus in +rDNA and lack of propodium iodide staining in -rDNA nuclei.

Extended Figure 5: Coarse-grained model for the assembly of the nucleolus and PCH. (**A**) The phase diagram illustrates the minimum attraction strength required to condense Fibrillarin for different concentrations. (**B**) Simulation endpoint snapshots depict the outcomes of varying the attraction strengths between beads of the PCH (H) polymer chain (ϵ_{HH}). (**C**) Simulation snapshots depict varying attraction strengths between rDNA-Fibrillarin (top to bottom) and beads of the PCH polymer chain (left to right). (**D**) Simulation endpoint snapshots depict the outcomes of varying X-X attraction strengths (ϵ_{XX}). (**E**) Simulation endpoint snapshots depict the outcomes of varying attraction strengths between Fibrillarin and protein X (ϵ_{FX}) in the -rDNA condition. (**F**) The average distance (d_{FX}) between Fibrillarin and protein X condensates from their center of mass is measured for different attraction strengths. Error Bars represent SD.

Extended Figure 6: Knockdown of Pitchoune in the female germline and eye-discs prevents their normal development. (**A**) Number of eggs laid on an apple juice plate after a 3hr collection in mataGAL4 (driver only) and mataGAL4 driving UAS-Pitchoune RNAi (VAL20 and VAL22 lines). (**B**) Representative images of dissected ovaries in control (mataGAL4, driver only) and after Pitchoune knockdown stained with DAPI. (**C**) Representative images of dissected eye antennal discs (yellow arrow) in control (ey-GAL4>mCherry RNAi) and after Pitchoune knockdown (ey-GAL4>pit RNAi, VAL20) stained with DAPI.

LEGENDS FOR SUPPLEMENTARY MOVIES

Supplementary Movie 1: Live imaging of RFP-Fib (green) and GFP-HP1a (magenta) in nuclear
cycle 13.

Supplementary Movie 2: Live imaging of RFP-Fib (green) and GFP-HP1a (magenta) in nuclear
cycle 14.

Supplementary Movie 3: Live imaging of RFP-Fib (green) and GFP-HP1a (magenta) in nuclear
cycle 15.

Supplementary Movie 4: Live imaging of RFP-Fib (green) and GFP-HP1a (magenta) in nuclear
cycle 16.

Supplementary Movie 5: Live imaging of RFP-Fib (green) and GFP-HP1a (magenta) in nuclear
cycle 17.

Supplementary Movie 6: Live imaging of GFP-Fib (green) and RFP-HP1a (magenta) in nuclear
amnioserosa nuclei in embryos lacking rDNA.

Supplementary Movie 7: Simulations of coarse-grained modeling of rDNA (red), Fibrillarin
(green), PCH (magenta), and an amphiphilic protein (yellow) with the parameters listed in the 4X4
matrix in Figure 5B.

Supplementary Movie 8: Live imaging of Pitchoune-GFP (green) and RFP-HP1a (magenta) in
nuclear amnioserosa nuclei in embryos lacking rDNA.