Filter-aided extracellular vesicle enrichment (FAEVEr) for proteomics

Jarne Pauwels[1,2], Tessa Van de Steene[1,2], Jana Van de Velde[1,2,3], Freya De Muyer[1,2], Danaë De Pauw[1,2], Femke Baeke [4,5], Sven Eyckerman[1,2] & Kris Gevaert[1,2,$]

[1]VIB-UGent Center for Medical Biotechnology, VIB, 9052 Ghent, Belgium

[2]Department of Biomolecular Medicine, Ghent University, 9052 Ghent, Belgium

[3]Current address: Eurofins Scientific, 9052 Gent, Belgium

[4]Ghent University Expertise Center for Transmission Electron Microscopy and VIB BioImaging Core, 9000 Ghent, Belgium.

[5]Department of Biomedical Molecular Biology, Ghent University, VIB Center for Inflammation Research, 9052 Ghent, Belgium.

[$]Corresponding author: Prof. Dr. Kris Gevaert, VIB-UGent Center for Medical Biotechnology and Department of Biomolecular Medicine, Ghent University, Technologiepark-Zwijnaarde 75, 9052 Ghent, Belgium; e-mail: kris.gevaert@vib-ugent.be

ORCID ID JP: 0000-0002-8525-7109

ORCID ID TVDS: 0000-0001-7915-2055

ORCID ID FDM: 0009-0008-2311-1965

ORCID ID DDP: 0009-0001-4093-6802

ORCID ID FB: 0000-0002-0761-9338

ORCID ID SE: 0000-0001-7525-0096

20   ORCID ID KG: 0000-0002-4237-0283

A. Abstract

Extracellular vesicles (EVs), membrane-delimited nanovesicles that are secreted by cells into the extracellular environment, are gaining substantial interest due to their involvement in cellular homeostasis and their contribution to disease pathology. The latter in particular has led to an exponential increase in interest in EVs as they are considered to be circulating packages containing potential biomarkers and are also a possible biological means to deliver drugs in a cell-specific manner. However, several challenges hamper straightforward proteome analysis of EVs as they are generally low abundant and reside in complex biological matrices. These matrices typically contain abundant protein concentrations that vastly exceed those of the EV proteome. Therefore, extensive EV isolation and purification protocols are imperative and many have been developed, including (density) ultracentrifugation, size-exclusion and precipitation methods. Here, we describe filter-aided extracellular vesicle enrichment (FAEVEr) as an approach based on 300 kDa MWCO filtration that allows the processing of multiple samples in parallel within a reasonable timeframe and at moderate cost. We demonstrate that FAEVEr is capable of quantitatively retaining EV particles on filters, whilst allowing extensive washing with the mild detergent TWEEN-20 to remove interfering non-EV proteins. The retained particles are directly lysed on the filter for a complete recovery of the EV protein cargo towards proteome analysis. Here, we validate and optimize FAEVEr on recombinant EV material and apply it on conditioned medium as well as on complex serum. Our results indicate that EVs isolated from MCF7 cells cultured with or without serum have a drastic different proteome because of nutrient deprivation.

B. Keywords

Extracellular vesicles, 300 kDa MWCO ultrafiltration, conditioned medium, starvation, EV-depleted serum, TWEEN-20, LC-MS/MS, proteomics

C. Background

Extracellular vesicles (EVs) make up a heterogeneous population of membrane-enclosed particles that are released by virtually all cells into their extracellular (EC) environment and in bodily fluids, such as plasma, urine and cerebrospinal fluid. In general, EVs are subdivided in three main categories according to their biogenesis and size (1). The smallest (30 – 150 nm), termed exosomes, are generated by invaginations at the endosomal membrane, in a transport (ESCRT) dependent (2,3) or independent manner (4). During this process, early endosomes mature into a multivesicular body (MVB) that holds multiple membrane delineated vesicles termed intraluminal vesicles (ILVs). Once the MVB fuses with the plasma membrane (PM), these ILVs are released into the EC environment as exosomes. Microvesicles (100 – 500 nm) are formed by an immediate outward budding of the PM, orchestrated by the ESCRT machinery (5). These protrusions are ultimately pinched off and released from the cell. Finally, apoptotic bodies (50 – 5,000 nm) are formed during apoptotic cell disassembly with large extrusions at the PM, fragmenting dying cells in large blobs that may even contain organelles (6). Ultimately, in the body, different EV categories are released from millions of individual cells, forming an immensely heterogeneous EV pool. This also results in the loss of information on their biogenesis and therefore, in practice, EVs are distinguished from one another by their size; small (sEV, < 200 nm) or large (lEV, > 200 nm), as described in the MISEV2018 guidelines (7). Note that since there is a certain level of size overlap between the different subtypes, none of these subpopulations is exclusive for either the small or the large EV group (8).

Due to their biogenesis, EVs have some important biophysical properties. First, EVs are

delineated by a phospholipid bilayer, effectively shielding their cargo (consisting of proteins,

oligonucleotides (RNA & DNA), lipids and metabolites) from the hostile extracellular

environment, thereby, maintaining the functionality and stability of biomolecules (9). Therefore,

the cargo can be considered to represent the molecular state of the parental cell at the moment

of EV formation. In addition, (trans)membrane proteins are incorporated into the lipid bilayer,

which, in turn, allows targeted signal transduction with distant EV receiving cells and/or allows

the selective delivery of their cargo in such cells (10,11). In physiological conditions, this cargo

takes part in cellular communication and homeostasis (12,13). In disease, however, the EV cargo

can contribute to the pathology, for example by initiating a pre-metastatic niche at a distant

location during cancer progression (14,15).

Both of these EV characteristics are appealing for clinical implementations, offering great

potential for both therapeutic applications (i.e. targeted drug delivery) as well as for biomarker

discovery (e.g. disease diagnosis, prognosis and therapy guidance) (16–18). Although the

therapeutic applications remain rather hypothetical or, at best, experimental to date (19,20), the

biomarker potential of circulating EVs has known a substantial increase in the past decade.

Indeed, EVs are present in different biofluids, such as plasma and urine, which can be isolated

in a low to non-invasive way at multiple time points.

In general, considering the biophysical properties of EVs, the identification and characterization

of protein biomarkers within or on circulating EVs from liquid biopsies might lead to (early)

disease diagnosis, monitoring of disease progression and therapeutic response, all critical

aspects towards the establishment of precision treatment (21). In this respect, EVs have shown

clinical potential as a source of circulating biomarkers for different cancers (e.g. breast (18),

urinary tract (22) and bone cancer (23)) and other diseases such as Alzheimer's disease (24,25),

chronic liver disease (26) and multiple sclerosis (27).

Unfortunately, multiple challenges complicate EV protein research, especially with regard to

proteome analysis by bottom-up mass spectrometry (MS), which is the method-of-choice for

unbiased protein biomarker discovery (28). Indeed, once EVs are released by the cell they form

a heterogeneous pool that resides in biological complex matrices that contain proteins at levels

that exceed those of the low abundant EV proteins by several orders of magnitude (29), such as

albumin in plasma and uromodulin in urine. In addition, they co-exist with non-EV particles that

overlap in size and density such as apolipoprotein particles. As a result, an adequate EV

enrichment and purification strategy remains essential for in-depth EV-associated biomarker

discovery (30). Of note, in MS-based proteomics, the recent advent of data-independent

acquisition (DIA) MS due to computational (31,32) and instrumental (33) improvements has

proven to cope more successfully with the above-mentioned challenges compared to traditional

data-dependent acquisition (DDA), as it is less biased to relative high protein levels.

Over the years, different EV enrichment strategies have been developed to enrich, purify and

concentrate EVs for in-depth proteome analysis and quantitative profiling, with the most

prominent strategies being based on size (ultracentrifugation and size-exclusion

chromatography) (34,35), density (density gradient ultracentrifugation) (36), precipitation (37,38)

or capture of molecular markers present on the EV surface (immuno-capture) (39). To further

enhance the purity of EV preparations, different orthogonal approaches can be combined (40,41).

However, a golden standard is yet to be determined, as the different strategies not only differ in

the consumption of resources and time, but also in terms of effectiveness and reproducibility.

Indeed, the different EV isolation techniques introduce variation between experiments in terms

of EV count and heterogeneity of molecular (protein) cargo. This has been emphasized in

different studies in which EV enrichment strategies were compared (42–46). Along this line,

initiatives such as MISEV and EV-TRACK were established within the EV research community to

encourage transparency on experimental procedures used to enrich EVs and on acquired

results, thereby enhancing inter-lab reproducibility (7,47). Importantly, purified, recombinant

EVs are now used as biological reference material to monitor technique development and

reproducibility, and improve data normalization (48,49). However, all enrichment strategies have

one major bottleneck in common; they lack the possibility for parallelization and high-

throughput set-ups. Indeed, enrichment strategies such as (density gradient) ultracentrifugation

require (multiple) lengthy centrifugation steps, are generally limited to six or eight samples and

require expensive instrumentation. Size-exclusion chromatography (SEC) often requires a

concentration step both prior and after EV enrichment and, in addition, also validation (often by

Western blotting) to identify those fractions containing EV material (50). Commercial EV

enrichment kits are again expensive and, more importantly, show high variability between

suppliers (51,52).

Ultrafiltration (UF) has thus far been mainly used as an intermediate step in EV enrichment

strategies to reduce the volume of conditioned medium or liquid biopsy either prior (e.g. density

gradient ultracentrifugation) or after EV enrichment (e.g. SEC) (53–55). The main challenges

associated with ultrafiltration for EV proteome analysis are the inefficient removal of non-EV

proteins, irreversible blocking of the filter membrane and low recovery of the retained particles

(56,57). However, it also offers inherent advantages, such as the limited cost, reduced sample

preparation time and the potential of analyzing multiple samples simultaneously using standard

lab equipment. In addition, UF has little bias towards EV subpopulations (58), in contrast to (for

instance) immuno-capture.

Here, we propose an ultrafiltration strategy for the isolation and purification of EVs from

conditioned medium and bovine serum, which we termed filter-aided extracellular vesicle

enrichment or FAEVEr. In short, pre-cleared conditioned medium or serum is loaded on 300 kDa

MWCO filters and centrifuged briefly at a moderate speed during which EVs are retained on the

filter, whereas the bulk of globular (non-EV) proteins do not. Remaining filter-retained non-EV

proteins are removed through subsequent purification steps that include TWEEN-20-containing

wash buffers. The retained (purified) EVs are then efficiently lysed on the filter in SDS-containing

buffer and the EV lysates are conveniently collected by centrifugation and processed for

proteomics applications, such as Western blotting and LC-MS/MS analysis. Recombinant EV

(rEV) material isolated from HEK293T conditioned medium was used for testing and optimizing

our strategy for comparison to ultracentrifugation, the most widely implemented EV enrichment

strategy. We demonstrate that the EV purity is considerably improved when including TWEEN-

20 in the wash steps without jeopardizing EV integrity, even at high concentrations of the

detergent. We then successfully applied FAEVEr to complex biofluids (bovine serum) and

investigated the effect on the EV protein cargo when using serum-depleted medium on MCF7

cell cultures.

D. Material and methods

1. Cell culture

All cell culture experiments were performed using standard protocol and lab equipment.

a. rEV production

A complete protocol for the transformation and production of recombinant extracellular vesicles (rEV) material is described in (49,59). In short, approximately 4 million HEK293T cells of low passage number (< 10) were seeded in a T75 falcon with Dulbecco's Modified Eagle Medium (DMEM, Gibco) containing 10% standard FBS and incubated for 48 h at 37 °C in 5% CO_2. The HEK293T cells were transfected by adding a mixture of polyethyleneimine (PEI) and the DNA bait construct (12x, pMET7- GAG-eGFP) in DMEM + 2% FBS for 6 h at 37 °C in 5% CO_2. The medium was then discarded and replaced with 8 mL fresh DMEM supplemented with 10% EV-depleted FBS (EDS, Thermo A2720801). After 48 h incubation at 37 °C in 5% CO_2, cell transfection was evaluated under UV light (excitation at 488 nm and emission at 507 nm) before the conditioned medium (CM) was isolated. The CM was immediately pre-cleared by centrifugation at 1,000 x g for 10 min and filtration of the supernatant using 0.22 µm filters (Millex). The pre-cleared CM was divided over different aliquots and frozen in -80°C until further use. Note that pure rEV extracts are commercially available (SAE0193, Sigma-Aldrich).

b. MCF7 cells

MCF7 cells of low passage number (< 10) were cultured using a standard protocol and equipment. The cells were seeded in T75 flasks with DMEM containing 10% standard FBS and incubated at 37 °C in 5% CO_2. When the cells reached 70-80% confluence, the medium was discarded and the

cells were washed three times with PBS. Experiments were performed in duplicate, adding 8

mL of DMEM supplemented with either 0%, 2%, 5% or 10% EDS (Thermo, A2720801) or OptiMEM

supplemented with 0% EDS. After 24 h, the CM containing 0% EDS were isolated, whereas the

2%, 5% and 10% EDS containing CM were isolated after 48 h. The CM was immediately pre-cleared

by two centrifugation rounds (500 x g for 5 min, and 10,000 x g for 10 min, room temperature)

and filtration using 0.22 µm filters (Millex).

2. EV enrichment

a. Ultrafiltration using 300 kDa MWCO filters

Pre-cleared CM was diluted 1:1 with PBS with or without supplemented TWEEN-20 and vortexed

for 5 – 10 s. The final volume was then transferred to 300 kDa MWCO polyethylene sulfone (PES)

filters (Vivaspin®6, Sartorius VS0652) and centrifuged. Centrifugation was done for 10 – 15 min

at room temperature either at 6,000 x g for fixed angle rotors (F14x50cy, Sorvall Lynx 4,000,

Thermo) or at 4,000 x g for swing-out rotors (e.g. A-4-44, 5804R, Eppendorf). The filter retentate

was washed several times with wash buffer (PBS with or without supplemented TWEEN-20) and

centrifuged for 5 – 7 min per wash step. After three rounds of washing, the retained particles

were either solubilized from the filter for nanoparticle tracking analysis or electron microscopy,

or immediately lysed on the filter for maximal recovery of the proteome. For recovery, 500 µL

50 mM HEPES was added on the filter and vortexed for 10 – 15 sec after which the solution was

transferred to a fresh tube and vacuum-concentrated (SpeedVac). To immediately collect the EV

lysate, 5% SDS in 50 mM TEAB is added and collected after 15 min incubation (37°C, 500 rpm) by

centrifugation (5 min at room temperature).

b.   Ultracentrifugation

Pre-cleared conditioned medium was transferred to polycarbonate thick-wall tubes (Beckman,

343778) and centrifuged for 90 min at 100,000 x g (4°C) using a Optima TLX ultracentrifuge

equipped with a TLA-120.2 rotor (Beckman). The supernatant was carefully removed and the

pellet washed with PBS with or without TWEEN-20 before a second round of ultracentrifugation

for 90 min at 100,000 x g (4°C). Again, the supernatant was carefully removed and the pellets

were either solubilized for NTA or EM, or immediately lysed for maximal recovery of the

proteome. The EVs were lysed by adding 5% SDS in 50 mM TEAB and the lysate collected after

15 min incubation (37 °C, 500 rpm).

3.   Proteomics

a.   Western blot

Samples for Western blot analysis were incubated for 5 min at 95 °C with XT sample buffer (4x)

and XT reducing agent (20x) (BioRad). The protein material was separated by SDS-PAGE (4%-

12%) for 80 min at 120 V prior to transfer onto a PVDF membrane (100 V, 30 min). The membrane

was incubated overnight at 4 °C with primary antibodies (rabbit monoclonal anti-Calnexin

(ab133615) and mouse monoclonal anti-HIV p24 39/5 4A (ab9071); 1:1,000). The next day, the

membrane was washed several times before incubation for 1 h at room temperature with

secondary antibodies (goat anti-rabbit IRDye 680 CW and goat anti-mouse IRDye 800 CW;

1:10,000). Visualization was done on an Odyssey infrared imager (v3.0.16, LI-COR Biosciences).

b. LC-MS/MS sample preparation

The proteome was prepared for LC-MS/MS analysis using S-Trap™ mini columns (ProtiFi, C02-mini-80) using the manufacturer's protocol. In brief, proteins in lysis buffer (5% SDS 50 mM TEAB, pH 7.4) were reduced and alkylated using 5 mM TCEP (10 min at 55 °C) and 20 mM iodoacetamide (IAA, 10 min at room temperature in the dark), respectively. The sample was acidified to 1.2% phosphoric acid and diluted seven-fold with binding/wash buffer (90% methanol in 100 mM TEAB, pH 7.4). The samples were transferred to the corresponding S-Trap™ columns and each time centrifuged at 4,000 x g for 30 s during sample loading and washing. For protein digestion, 0.5 µg of trypsin (Promega, V5111) was added to 125 µL 50 mM TEAB (pH 7.4) and samples were incubated overnight at 37 °C. The next day, the peptides were recovered by sequential elution by centrifugation (4,000 x g for 30 s at room temperature) using 80 µL 50 mM TEAB (pH 7.4), 80 µL 0.2% formic acid and 80 µL of 50% acetonitrile (ACN), 0.2% formic acid in dH$_2$O. The eluates (app. 365 µL) were transferred to MS vials, vacuum-dried in a SpeedVac and stored at -20 °C. We have submitted all relevant data of our experiments to the EV-TRACK knowledgebase (EV-TRACK ID: EV240045) [34].

c. LC-MS/MS analysis

Purified peptides were re-dissolved in loading solvent A (0.1% TFA in water/ACN (98:2, v/v)) and the peptide concentration was determined on a Lunatic instrument (Unchained Lab). For each sample the injection volume was adjusted to inject equal amounts of peptide material (500 ng) for LC-MS/MS analysis on an Ultimate 3000 RSLCnano system in-line connected to a QExactive Exploris (for optimization and validation) or an Orbitrap Fusion Lumos mass spectrometer (Thermo) (MCF7 cells and EVs). QCloud was used to control instrument longitudinal performance during the project [57, 58]. The mass spectrometry proteomics data have been deposited to the

237   ProteomeXchange Consortium via the PRIDE [59] partner repository with the dataset identifiers

238   PXD051938 (comparison FAEVEr with UC), PXD051955 (comparison MCF7 proteome of EVs and

239   cells under starving conditions) and PXD051956 (comparison FBS with EDS).

LC-MS/MS analysis using the Fusion Lumos instrument

LC-MS/MS data-independent (DIA) acquisition on the Fusion Lumos was initiated by trapping

the peptide material at 20 µl/min for 2 min in loading solvent A on a Pepmap column (300 µm

internal diameter (I.D.), 5 µm beads, C18, Thermo). The peptides were separated on a 110 cm µPAC

prototype column (Thermo), kept at a constant temperature of 50 °C. Peptides were eluted by a

non-linear gradient starting from 2% MS solvent B (0.1% formic acid (FA) in acetonitrile), reaching

26.4% MS solvent B in 82 min and 44% MS solvent B in 90 min and 56% MS solvent B in 100 min

starting at a flowrate of 600 nl/min for 5 min, and completing the run at a flow rate of 300 nl/min,

followed by a 5-minute wash at 56% MS solvent B and re-equilibration with MS solvent A (0.1%

FA in water). The mass spectrometer was operated in data-independent mode. Full-scan MS

spectra ranging from 400-900 m/z with a target value of 4E5, a maximum fill time of 50 ms and

a resolution of 60,000 were followed by 50 quadrupole isolations with a precursor isolation width

of 10 m/z for HCD fragmentation at an NCE of 34% after filling the trap at a target value of 4E5

for maximum injection time of 54 ms. MS2 spectra were acquired at a resolution of 30,000 at

200 m/z in the Orbitrap analyser without multiplexing. The isolation intervals ranged from 400 –

900 m/z, without overlap.

LC-MS/MS analysis using the Orbitrap Exploris 240 instrument

The peptide material was injected for LC-MS/MS analysis on a Vanquish™ Neo UHPLC System

in-line connected to an Orbitrap Exploris 240 mass spectrometer (Thermo). Injection was

performed in trap-and-Elute workflow in combined Control mode (maximum flow of 60 µl/min

and maximum pressure of 800 bar) in weak wash solvent (WW, 0.1% trifluoroacetic acid in

water/acetonitrile (ACN) (99.5:0.5, v/v)) on a 5 mm trapping column (Thermo scientific, 300 μm

internal diameter (I.D.), 5 μm beads). The peptides were separated on a 250 mm Aurora Ultimate,

1.7μm C18, 75 μm inner diameter (Ionopticks) kept at a constant temperature of 45 °C. Peptides

were eluted by a gradient starting at 0.5% MS strong wash solvent (SW) (0.1% FA in acetonitrile)

reaching 26% MS SW in 30 min, 44% MS SW in 38 min, 56% MS SW in 40 min followed by 5-minute

wash at 56% MS SW and column equilibration in Pressure Control mode (separation column:

fast equilibration, maximum pressure of 1500 bar, equilibration factor=2; Trap column: fast wash

and equilibration, wash factor=100) with MS WW . The flow rate was set to 300 nl/min. The mass

spectrometer was operated in data-independent mode, automatically switching between MS and

MS/MS acquisition. Full-scan MS spectra ranging from 400-900 m/z with a normalized target

value of 300%, a maximum fill time of 25 ms and a resolution at of 60,000 were followed by 30

quadrupole isolations with a precursor isolation width of 10 m/z for HCD fragmentation at an

NCE of 30% after filling the trap at a normalized target value of 2000% for maximum injection

time of 45 ms. MS2 spectra were acquired at a resolution of 15,000 with a scan range of 200-

1800 m/z in the Orbitrap analyser without multiplexing. The isolation intervals were set from 400

– 900 m/z with a width of 10 m/z using window placement optimization. EASY-IC^{TM} was used in

the start of the run as Internal Mass Calibration

d.   Database searching and data analysis

Data-independent acquisition (DIA) spectra were searched with the DIA-NN software (v1.8.2b)

[60] in library-free mode against the combination of two protein databases downloaded from

Swiss-Prot; the complete human protein sequence database (January 2021, 20,394 sequences)

supplemented with bovine serum protein sequences (March 2021, 618 sequences). For searches

that included rEV material, we included the Gag-eGFP protein sequence as well. The mass

accuracy was set to 10 ppm and 20 ppm for MS1 and MS2, respectively, with a precursor FDR of

0.01. Enzyme specificity was set to trypsin/P with a maximum of two missed cleavages. Variable

modifications were set to oxidation of methionine residues (to sulfoxides) and acetylation of

protein N-termini. Carbamidomethylation of cysteines was set as a fixed modification. The

peptide length range was set to 7-30 residues with a precursor charge state between 1 and 4

and an m/z range between 400-900 and 200-1,800 for the precursor and fragment ions,

respectively. Cross-run normalization was set to RT dependent with the quantification strategy

set to high accuracy and the neural network classifier to single-pass mode. The result file was

further processed in KNIME (v4.3.3) by removing non-proteotypic peptides and identifications

with a protein q-value above 0.01. Peptide quantifications were aggregated to protein group

quantifications using the median of the corresponding normalized LFQ values. Further data

analysis was performed with Perseus (version 1.6.14.0), GraphPad Prism (v9.4.1) and RStudio

(2023.12.1).

For the analysis of the EV and cellular proteomes across different experimental conditions,

proteins that were significantly differentially abundant were identified by multiple sample testing

(ANOVA) and selected for hierarchical clustering (Euclidean distance with averaged linkage)

after Z-scoring the individual proteins over the different conditions. Row clusters were defined

manually and used for gene ontology (GO) enrichment against the corresponding initial database

and included biological process (GOBP), molecular function (GOMF), cellular component (GOCC)

annotations and KEGG pathways. GO term enrichment was done using the online WebGestalt

tool for Gene ontology enrichment (GOA) (60).

4. EV characterization

a. NTA analysis

Recuperated EV particles were diluted to 1 mL using PBS (pH 7), injected in a calibrated Zetaview and analyzed with the corresponding software package (version 8.05.16 SP2). The temperature was maintained at 23°C and the analysis was done at a sensitivity of 70 and a shutter of 100 for 3 cycles at 60 fpm. The results were plotted and analyzed in GraphPad Prism (v10.2.2)

b. Scanning electron microscopy

Filters containing EVs were placed in 2 ml Eppendorf tubes and fixed with 4% paraformaldehyde and 2% glutaraldehyde in 0.1 M Sodium cacodylate buffer at room temperature. After one hour, the fixative was replaced and filters were washed 3 times for 5 min with dH_2O. After dehydration in 30%, 50%, 70%, 95% and 2x 100% ethanol for 30 min each, the filters were critical point dried (EM CPD300, Leica) and imaged on a Crossbeam 540 SEM (Zeiss) at 1kV.

c. Transmission electron microscopy

Aliquots (5 µl) of the EV solutions were blotted for 1 min on formvar- and carbon-coated Ni maze grids (EMS), which were glow discharged for 40 sec at 15 mA. The grids were then washed five times in droplets of dH_2O and stained in a droplet of 1/4 Uranyl Acetate Replacement Stain (EMS)/ dH_2O for 1 min. Excess stain was removed with filter paper and the grids were air dried for at least four hours before viewing with the TEM. Imaging was done at 80kV on a JEM1400plus (JEOL).

E. Result section

1. Use of 300 kDa MWCO filters to enrich EVs

Although ultrafiltration is a valid and well-implemented method to concentrate intact EVs, it holds many challenges as a stand-alone EV enrichment strategy. Nevertheless, ultrafiltration also has important advantages such as low cost and fast processing times, and it does not require specific instrumentation compared to, for instance, ultracentrifugation.

Here, we introduce Filter-Aided Extracellular Vesicle Enrichment (FAEVEr), a strategy that uses 300 kDa MWCO polyether sulfone (PES) filters to purify and concentrate intact EVs (Figure 1). In brief, pre-cleared conditioned media or biofluids are transferred onto the 300 kDa MWCO filter membrane and centrifuged at moderate speed (4,000 – 6,000 x g), retaining the EV particles on the filter (retentate) whereas globular non-EV proteins are removed (filtrate). By including three consecutive wash steps, non-EV proteins are increasingly removed, by which the EV retentate is further purified. Eventually, by using a lysis buffer with high percentages (5%) of SDS, EVs undergo efficient lysis and their cargo is conveniently recovered by centrifugation. The EV sample is then further processed for proteomics using S-Trap columns (61). Here, we opted for PES membranes, as filters based on such membranes are hydrophilic with low protein adsorption characteristics. In addition, PES material is resistant against high pH ranges and temperatures and can withstand a wide variety of solvents, buffers and detergents.

Recombinant EV (rEV) material was used for the initial optimization of the 300 kDa MWCO ultrafiltration strategy as it has similar biochemical and biophysical characteristics as sample EVs (48). The rEV material is derived from the conditioned medium of HEK293T cells, cultured in 10% EV-depleted serum, that overexpress the Gag-eGFP fusion protein (49). The Gag subunit

multimerizes at the plasma or endosomal membrane and initiates membrane curving, resulting

in the formation of microvesicles and intraluminal vesicles, respectively, commonly referred to

as virus-like particles (59,62) (**Error! Reference source not found.**). As a result, secreted rEV

material is overrepresented in the conditioned medium and contains high levels of luminal Gag-

eGFP, enabling the evaluation of EV integrity during the consecutive FAEVEr steps.

Figure 1: Schematic overview of the FAEVEr workflow. Conditioned media or biofluids are

precleared and transferred onto the 300 kDa MWCO PES filter (red). Driven by centrifugation,

the globular proteins pass through the filter (orange), whereas the EV particles are

quantitatively retained (grey). The retentate is subsequently washed three times using PBS

supplemented with TWEEN-20 to further increase the EV purity. Ultimately, intact particles are

recovered for qualitative analysis, or lysed (green) immediately on the filter. The lysate is

recovered by centrifugation and prepared for proteomic applications (e.g. LC-MS/MS analysis

or Western blotting). Figure created in Biorender.

We confirmed that rEV particles are retained on the 300 kDa MWCO mesh using scanning

electron microscopy (SEM) (Figure 2A and **Error! Reference source not found.**), indicating that

ultrafiltration is an efficient EV enrichment strategy. Indeed, after completing the FAEVEr

protocol, rEV particles were recovered from the filter for nanoparticle tracking analysis (NTA)

(Figure 2B) and transmission electron microscopy (TEM) characterization (Figure 2C). However,

in line with Vergauwen *et al.* (56), we noticed that the quantitative recuperation of particles from

the filter PES membrane is variable and that not all particles are recovered. When the remaining

rEV fraction on the filter was lysed and analyzed on Western blot, we observed that a substantial

share of the rEV material fails to be solubilized and remains on the filter (Figure 2D).

Figure 2: Qualitative analysis of FAEVEr retained particles. A) Scanning electron microscope of

an empty filter and after rEV enrichment. B) EV size distribution and count, and transmission

electron microscopy (C) of recovered rEV particles from the filter. D) Western blot reveals

incomplete recuperation of rEV particles from the filter. EDS: EV-depleted serum (negative

control), rEV material: commercial rEV particles (positive control), PBS: phosphate-buffered

saline washed rEV particles.

Concluding this part, we found that FAEVEr is an efficient approach for isolating EV particles

from conditioned medium for downstream proteome analysis and that we can successfully

recuperate intact particles from the filter (56). By immediately lysing the filter-retained EV

material using high percentages of SDS for proteomic applications, we ensure EV lysis,

comprehensive solubilization of (membrane) proteins (63) and maximal protein recovery. Note

that SDS is irreconcilable with LC-MS/MS analysis and we therefore introduced S-Trap columns

in our workflow (61,64). The S-Trap protocol is easy, fast and reproducible and has been

extensively compared to other methods in different studies (65,66).

2. Comparison of FAEVEr to ultracentrifugation

Using EV proteome analysis, we set out to compare our FAEVEr approach with

ultracentrifugation (UC), the most used strategy for EV enrichment. rEV material was isolated,

pre-cleared and divided over different aliquots before subjecting it to the corresponding EV

enrichment strategy in triplicate. In terms of sample preparation, the overall sample preparation

time differs noticeably. Ultracentrifugation required at least two rounds of centrifugation,

totaling to 3 h of centrifugation time, whereas FAEVEr was finished within the hour, including

centrifugation steps and hands-on time.

For an in-depth analysis of the proteome results, we examined the occupancy of the LC-MS/MS

spectral space, which is a measure of the recorded spectra and allows us to differentiate

between recorded spectra of bovine and human origin. Our results indicate that the relative

spectral space occupancy by bovine precursors was slightly reduced (from 35% to 30%) using

FAEVEr compared to UC (Figure 3A). In addition, approximately 15% more proteins were

identified proportionally distributed over the different cellular locations (Figure 3B). The six

commonly used EV markers are well represented and identified with similar relative intensities

between strategies (Figure 3C) and 85% of the identified proteins by the UC were found by

FAEVEr (Figure 3D). In addition, the same trend was observed for the number of unique peptides

(13%) and the number of the associated peptide-spectral matches (PSMs), the number of

recorded spectra that were translated in a confident peptide identification (14%) (Figure 3E).

Importantly, we observed a large decrease in the accompanying coefficient of variance (%CV)

for FAEVEr between the replicates as well as fewer missing values, which results in an

increased data completeness and therefore higher reproducibility with more accurate protein

quantifications (Figure 3F). From this experiment, we can thus conclude that ultrafiltration using

300 kDa MWCO filters allows for EV enrichment and is comparable to UC in terms of EV proteome

depth.

Figure 3: Overview of the comparison between FAEVEr and ultracentrifugation. A) Comparison

of the relative abundance of precursor material from bovine or human origin. B) Distribution of

the protein identifications across the cellular localizations of the parental cell. C) High relative

abundance of six molecular EV markers. D) Overlap of protein identifications between UC and

FAEVEr. E) Total number of identified human proteins, identified unique peptides and peptide-

spectral matches from these proteins between UC and FAEVEr. F) Decreased relative variance

(%CV) and fewer missing values are observed with FAEVEr compared to UC. UC:

ultracentrifugation (orange), FAEVEr: Filter-aided EV enrichment (teal), CC: cellular

compartment, LFQ: label-free quantification intensity, PSM: peptide-spectral match, %CV:

coefficient of variance.

3. TWEEN-20 improves the removal of non-EV proteins

One major pitfall of ultrafiltration is the inefficient removal of non-EV proteins, which has a

detrimental effect on the numbers of identified and quantified EV proteins following LC-MS/MS

analysis. Therefore, we explored the use of mild detergents to minimize the interaction between

proteins and the filter membrane. However, as EVs are delineated with a lipid bilayer and

therefore susceptible to detergent-mediated lysis, caution should be taken with the choice and

concentration of the detergent. A study by Osteikoetxea *et al.* explored the minimal concentration

of different detergents required for efficient lysis of EV particles in solution (67) and found that

this concentration greatly depends not only on the detergent used, but also on the EV

subpopulation (exosomes, microvesicles and apoptotic bodies). In particular, the smallest

population (exosomes) demonstrated increased resilience against detergents compared to

larger vesicles (microvesicles and apoptotic bodies) due to an increase in liquid order because

of an accumulation of cholesterol and sphingolipids during their formation (68–70). In their work,

Osteikoetxa *et al.* demonstrate that EVs are generally efficiently lysed even at low percentages

of SDS, Triton X-100 or sodium deoxycholate. On the contrary, an exceptional high tolerance is

observed towards TWEEN-20, a rather mild detergent that is commonly used in other

biochemical assays such as Western blotting to reduce unwanted adsorption of protein material

to membranes (71) and thereby decreases irreversible or total membrane fouling (72,73).

TWEEN-20 was explored here as an additive to improve the efficiency of the washing steps and

thus removal of non-EV proteins during the enrichment of EVs whilst keeping EVs intact. The

efficiency of washes with TWEEN-20-containing buffers was tested by Coomassie staining of the

different filtrate fractions of conditioned medium after completing the FAEVEr protocol using 0%, 0.1%, 0.5%, 1%, 2% and 5% TWEEN-20 in the wash buffer. We observed that PBS-only (the 0% TWEEN-20) condition is not capable of removing the bulk of bovine serum albumin (BSA; 65 kDa), whereas BSA-bands are more intense in the TWEEN-20 supplemented wash fractions. In addition, upon recovery of the retained material by lysis in 5% SDS, we observed an intense band for the PSB-only condition, which was absent in the TWEEN-20 supplemented ones (**Error! Reference source not found.**). This suggested that TWEEN-20 is indeed capable of removing the bulk of (contaminating) bovine serum material from the filter membrane and therefore should result in an EV proteome fraction that is more pure.

Next, we validated this beneficial effect of TWEEN-20 in wash buffers using rEV material, which also allows us to validate the integrity of the particles as they contain the Gag-eGFP protein on the luminal side. We thus argued that, when the rEV particles remain intact, no luminal Gag-eGFP should be present in the filtrates. In short, pre-cleared conditioned medium containing rEV material was filtered (300 kDa MWCO) and washed several times with up to 5% TWEEN-20 in PBS. After each centrifugation step, the filtrate was collected and the presence of luminal Gag-eGFP validated on Western blot, shown in Figure 4A. The almost complete absence of luminal Gag-eGFP in the flow-through and washes (1-3) does not only indicate that intact rEV particles are retained on the filter, but also that they maintain their integrity in concentrations up to 5% TWEEN-20, as validated by TEM, SEM and NTA (Figure 4B). In addition, it also indicates that the proteome of retained EVs is efficiently recovered from the filter for downstream proteomic applications using a lysis buffer containing 5% SDS.

Figure 4: The integrity of rEV particles remains even when washing with buffers supplemented with 5% TWEEN-20. A) Western blot of the different fractions during FAEVEr showing that no (or just marginal) luminal Gag-eGFP is observed in the flow-through and wash steps, implying that EVs remain intact. In addition, a strong signal is observed in the lysate fractions, indicating that

we can efficiently recover the rEV proteome. B) Qualitative analysis by TEM, NTA and SEM of

recovered rEV particles washed in PBS supplemented with 5% TWEEN-20.

We further extended our comparison between FAEVEr and UC to evaluate the effect of TWEEN-

20 on the purity of the final rEV proteome. We included different percentages of TWEEN-20 in

the washing buffer of both FAEVEr and UC and analyzed the corresponding proteomes by LC-

MS/MS. We initiated the analysis by comparing the spectral space distribution between human

and bovine precursors. Here, we did not only observe a remarkable decrease in bovine

precursor intensities using FAEVEr even at the lowest (0.1%) TWEEN-20 concentration, but also

that increasing the percentage further improved this (Figure 5A). The effect of supplementing

TWEEN-20 is summarized in a principle component analysis (PCA, Figure 5B) which illustrates

that supplementing TWEEN-20 has the most drastic change in combination with FAEVEr.

Indeed, by combining FAEVER with 5% TWEEN-20, we identified over 3,700 proteins with nearly

95% of all PSMs to be of human origin. For UC supplemented with 5% TWEEN-20 in the wash

buffer, we also observed an improvement in terms of decreased serum contamination, although

not to the same extent, where approximately 80% of the observed precursors were of human

origin. Interestingly, we identified over 4,400 proteins in the UC setup, or roughly 15% more

proteins in comparison to FAEVEr. We investigated this change in protein abundance further as

we expected these extra proteins not all to be serum proteins. We found, by initiating a multiple

sample test between the individual experimental setups, (as visualized in the PCA plot) three

major groups; UC with or without TWEEN-20, FAEVEr without TWEEN-20 and FAEVEr with

TWEEN-20. The setup in which FAEVEr was used without TWEEN-20 appeared to associate with

both groups to some extent (Figure 5C). By further examining gene ontology and KEGG terms

associated with the corresponding proteins, we found that in FAEVEr supplemented with

TWEEN-20, terms including cell-cell adhesion, ESCRT mechanism and (trans)membrane

proteins were more abundant (Figure 5D). On the other hand, we noticed that in UC, terms

including RNA binding, cell cycle, proteasome and metabolic activities were more prominent

(Figure 5E) as well as serum-related terms (e.g. blood microparticle, activation cascade).

In conclusion, our results indicate that FAEVEr in combination with TWEEN-20 in the washing

buffer improves the overall EV proteome purity, whereas this effect is less prominent in UC.

Figure 5: Overview of the effect of TWEEN-20 at different percentages on EV enrichment by UC

and FAEVEr. A) Relative share of bovine and human precursor abundance as a measure of

serum contamination in function of the enrichment strategy and the percentage of TWEEN-20 in the wash buffer. B) PCA plot of the shared protein intensities between the different experimental setups reveals three major groups. C) Heatmap of the significantly differential abundant proteins (multiple sample test, p-value < 0.05) per experimental setup with the accompanying GO analysis for FAEVEr enriched (D) and UC enriched (E) proteins. UC: ultracentrifugation (orange), FAEVEr: filter-aided EV enrichment (teal), PCA: principal component analysis, GO: Gene ontology, TM: transmembrane, CCA: cell-cell adhesion, ECM: extracellular matrix.

4. Effect of serum depletion (on the EV proteome)

Fetal bovine serum (FBS) is used as a standard supplement in cell culture media. Unfortunately, it is irreconcilable with EV analysis from conditioned cell media as it contains EVs from bovine origin that interfere with proteome analysis. Therefore, EV depleted serum (EDS) alternatives are used. Consequently, researchers often expand cells for several days in media containing EDS or even refrain from adding serum. To this day, it remains a controversial topic (74) with some researchers advocating against the use of serum because it remains a potential source of xeno-contamination. Others deem serum as an essential supplement for normal cell growth with serum depletion stressing cells due to starvation (75,76), which potentially biases the proteome of the cell culture-derived EVs. Benefiting from the EV-TRACE repository, we noticed that between 2018 and 2023 approximately 56% of the EV experiments that were conducted on conditioned medium of human cell lines were performed using EDS, whereas 30 – 35% of the EV experiments was performed using serum-free conditions.

Following the MISEV2018 (7) guidelines, we investigated whether the used EDS is indeed devoid of EV material and addressed this using FAEVEr comparing EDS with FBS. In addition, we

examined uses of FBS or EDS had an effect on the cellular and the EV proteome. For these

experiments, we made use of MCF7 breast cancer cells.

As the high protein concentrations and the accompanying viscosity of serum samples may result

in low filtration speed and irreversible blocking of the filter membrane, we diluted the sample

three times prior to filtration. After completing the FAEVEr protocol, the proteomes of the

FAEVEr filter-retained fraction from commercial FBS and EDS were compared by LC-MS/MS.

We validated that EV-specific molecular markers were absent in EDS as well as proteins

associated with EV biogenesis. In contrast, these proteins were found abundantly in the FBS

retained fraction (**Error! Reference source not found.**).

In conclusion, using FAEVEr, we here provide evidence at the LC-MS/MS level that EDS is indeed

devoid of EV material and that the FAEVEr strategy is capable of enriching EV particles from

complex biological fluids such as (bovine) serum.

The protein concentration of serum is reduced by a factor six because of the EV depletion

process (**Error! Reference source not found.**A). Indeed, FBS is diluted several times to prevent

unintended protein aggregation and pelleting by ultracentrifugation to remove EVs due to the

high viscosity of undiluted FBS (73). We decided to explore the effect on the MCF7 cellular

proteome once the medium was exchanged from 10% FBS to 10% EDS for 3 days. Besides no

change in cell morphology by visual inspection or a significant change in cell count or cell death

(**Error! Reference source not found.**B), the LC-MS/MS analysis indicated that only 10 out of more

than 7,000 quantified proteins appeared to be significantly differentially regulated between the

two conditions (**Error! Reference source not found.**C & 5D). We conclude that shifting from FBS

to EDS close to no influence on the viability, growth or proteome of MCF7 cells in culture.

We continued investigating the use of FAEVEr by exploring the effect of starvation on the EV

proteome by isolating EVs from the conditioned medium of MCF7 cells grown in DMEM

supplemented with 0%, 2%, 5% or 10% EDS (D0, D2, D5 and D10, respectively). The FAEVEr retained

EVs were purified with wash buffer containing 5% TWEEN-20 and either recovered for NTA

(Figure 6A) or lysed for LC-MS/MS analysis. The LC-MS/MS generated data immediately hinted

towards an important difference by protein content and quantity between serum-deprived and

EDS containing conditions as indicated in the number of identifications (Figure 6B) and the PCA

plot (Figure 6C). Over 5,191 proteins were identified in total, from which 4,887 were human and

304 were bovine. From the human proteins, 3,610 were found in duplicate in at least one

experimental set-up and were included for quantitative comparison of the EV proteomes.

Interestingly, we found that only 959 proteins were identified across all samples and that over

1,100 proteins were uniquely identified in D0. In addition, we found that the relative abundance of

transmembrane proteins gradually increased with increasing levels of EDS, whereas levels of

intracellular proteins decreased (Figure 6D). EV markers were abundantly picked-up across the

individual samples (Figure 6E). Proteins were then quantitatively compared (multiple sample

testing) and hierarchical clustered (Figure 6F) which lead to the conclusion that 333 proteins

were significantly differential abundant over two main clusters, between the serum depleted

and serum containing conditions. These two clusters were used for gene ontology analysis (GOA)

using the Webgestalt v2017 online tool (60). Note that the D0 unique protein list was joined with

the proteins that were significantly more abundant in the serum-depleted condition. This GOA

analysis revealed that EVs isolated from MCF7 cells cultured in serum-depleted conditions for

only 24 h are enriched in proteins involved in metabolism, RNA and DNA binding and autophagy,

including organelle fission. Indeed, serum deprived cells are known to rapidly adjust the cellular

proteome (**Error! Reference source not found.**), cells are known to upregulate autophagy related

(ATG) proteins to replenish essential building blocks and release non-ESCRT dependent

exosomes through the formation of autophagosomes (77), which can fuse with endosomes to

form amphisomes, similar to MVBs. In addition, the ATG5-ATG16L complex (78,79) influences the

EV secretion through dissociation of V-ATPases, thereby preventing MVB acidification and its

subsequent lysosomal degradation (80,81). Ultimately, this promotes MVB fusion with the PM

and secretion of the ILVs as exosomes. We found that multiple key ATG proteins and contributors

of the phagosome biogenesis are high abundant in the EVs isolated from the serum-starved

MCF7 cells yet nearly completely absent in the EVs isolated from the serum-containing

conditions (Figure 6G & 6H). These findings are in contrast to the serum-containing conditions,

where GO terms associated with (trans)membrane proteins (including transmembrane

transport, cell-cell adhesion, integrins, vesicle-mediated transport) are enriched as well as

proteins associated with the ESCRT complex.

Figure 6: Comparison between the EV proteomes isolated from MCF7 cells cultured in different

percentages of EV-depleted FBS (EDS) reveals drastic changes in protein heterogeneity and

abundances. A) NTA analysis of recovered EV particles. B) Number of protein identifications

between the conditions. C) PCA plot indicating large variation between the relative abundances

of proteins identified in serum-starved (D0) and serum-supplemented conditions (D2, D5 and

D10). D) Plot of the localization of the identified proteins. E) Profile plot of six commonly used EV

markers. F) Heatmap of the significantly differential abundant proteins between the different

conditions and the associated enriched GO terms for serum-starved (orange) and serum-

supplemented (teal). Detail of ATG related proteins in the serum-starved condition (G) and the

ESCRT associated proteins in the serum-supplemented conditions (H). EDS: EV-depleted FBS;

D0, D2, D5 & D10: DMEM supplemented with 0%, 2%, 5% and 10% EDS; NTA: nanoparticle tracking

analysis; PCA: principal component analysis; GO: Gene ontology.

F. Conclusion and discussion

Despite the potential of extracellular vesicles (EVs) for clinical applications such as being a

source of circulating biomarkers or therapeutic shuttles, their efficient isolation and purification

from complex biofluids in a cost and time efficient manner remains a major bottleneck. Here,

we propose filter-aided extracellular vesicle enrichment (FAEVEr) as an EV enrichment strategy

that uses 300 kDa MWCO filtration. By combining the filtration steps with wash buffers that

contain up to 5% TWEEN-20, we were able to improve the EV proteome purity towards LC-MS/MS

compatible levels, even in the presence of 10% EV-depleted FBS (EDS). Recombinant EVs (rEVs,

(48,49)) were used to initially establish and optimize our strategy. We made use of the

overexpression of a luminal Gag-eGFP construct to evaluate the rEV integrity during the

enrichment and purification and concluded that during FAEVEr the rEV particles indeed remain

intact. In addition, we concluded that the EVs can be recovered from the filter membrane for

further evaluation using common tools for EV characterization such as nanoparticle tracking

analysis (NTA) and scanning and transmission electron microscopy (SEM and TEM). In addition,

the rEV proteome was quantitatively recovered by immediately lysing the filter-retained EVs for

LC-MS/MS analysis and we identified over 5,000 human HEK293T proteins from 1 mL conditioned

medium. Amongst these proteins we found the different EV markers next to multiple

(trans)membrane proteins and proteins associated with EV biogenesis, which underlines the

potential of FAEVEr for EV proteomics.

We benchmarked our approach against ultracentrifugation (UC), a widespread implemented

method for EV enrichment, and showed that over 95% of the protein identifications overlap.

Furthermore, we found an increase in EV purity following FAEVEr compared to

ultracentrifugation as demonstrated by a reduced fraction of the LC-MS/MS spectral space

taken by bovine precursors, thereby limiting the competitive effect towards the identification of

low abundant (EV) proteins. In addition, we observed a drastic decrease in the relative variance

(%CV) between the number of identifications on both the protein, peptide and peptide-spectral

match level between UC and FAEVEr. This also indicates a higher reproducibility between

replicates, which in turn leads to improved data completeness (fewer missing values) and,

consequently, improved quantifications. We hypothesize that this improvement is due to the

absence of intermediate pipetting steps that remove the supernatant after ultracentrifugation,

which potentially leads to incomplete removal or unintended perturbations of the EV pellet. In

FAEVEr however, we create a distinct separation between the EV depleted filtrate and EV

enriched retentate, thereby omitting unnecessary pipetting steps, as the filtrate is simply

decanted.

We continued to improve the EV purity by implementing TWEEN-20 in wash buffers as this

detergent was previously shown not to lyse EVs, even in relatively high concentrations (67), and

it is an established supplement to reduce non-specific protein-membrane interactions, for

example in Western blotting. We provide insights in the gradual loss of bovine serum proteins

and secreted proteins, to which we refer here as non-EV proteins. This improved removal of

unwanted proteins is of high importance for LC-MS/MS based bottom-up proteomics, as

precursors from such high abundant proteins (e.g. albumin) dominate the spectral space,

outcompeting relevant EV precursors. Surprisingly, we observed that supplementing TWEEN-20

to the wash step of UC did not have the same profound effect compared to FAEVEr. Indeed, a

number of proteins associated with (amongst others) transcription and translation, proteasomal

degradation and metabolic activities were more prominent in all ultracentrifugation samples as

well as in the FAEVEr setup without TWEEN-20 supplemented to the wash buffers. In contrast,

the TWEEN-20 washed EV proteomes following the FAEVEr protocol were enriched in

(trans)membrane proteins and in proteins from the ESCRT machinery. We hypothesize that

during cell culture inevitable cell lysis occurs during which cytosolic proteins are released and

co-purified during ultracentrifugation. These insights raise the question whether or not such

proteins are effectively encapsulated in EVs or if they form non-specific protein interactions

with the EV membrane. However, additional studies are required to confirm this.

In EV research, many experiments exclude serum supplements in cell culture media to repulse

xeno-proteins and omit the suppressive competitive effect during LC-MS/MS analysis and the

potential biological influence on the EV proteomics outcome. We first investigated on the

difference between regular FBS and EV-depleted FBS (EDS) on the LC-MS/MS level, from which

we concluded that EV markers and biogenesis related proteins are virtually absent in EDS,

implying that we do not contaminate our EV preparation unintendedly with bovine EV particles.

We further elaborate that the depletion of serum during cell culture, even for a limited period of

time (e.g. 24 h), has a dramatic effect on the EV proteome of MCF7 cells. In line with Leidal *et al.*

(77) we observed an increase in protein heterogeneity associated with RNA binding,

biosynthesis, cell cycle and autophagy, pointing towards an alternative biogenesis of EV

particles through the formation of amphisomes. These findings are underlined by the unique

presence of several ATG proteins in the EV preparation of serum-starved MCF7 cells. In contrast

to this, the EV proteome from MCF7 cells supplemented with EV-depleted FBS, contained higher

levels of proteins associated with transmembrane transport, cell-cell adhesion and the ESCRT-

machinery were more abundant. We feel that the choice to add or exclude EDS during cell culture

for EV research is an important consideration when testing a preconceived hypothesis as it may

bias the interpretation of the EV (proteomics) data (82,83).

Our data shows that filter-aided EV enrichment, FAEVEr, using 300 kDa MWCO PES filters is an

efficient strategy to isolate EVs from complex medium and that, by including TWEEN-20, we

further improved the overall purity of the EV proteome without affecting the EV integrity, even

at relatively high TWEEN-20 concentrations. Furthermore, FAEVEr is conducted in a relatively

short timeframe and it offers the possibility of parallelization using standard lab equipment. In

comparison, ultracentrifugation is limited to six or ten samples and requires at least two rounds

of centrifugation.

G. Supplemental figures legends

Supplementary Figure 1: Schematic overview of the expression of GAG-eGFP in HEK293T cells.

Top: HEK293T cells are transfected with a Gag-eGFP construct, encoding for a Gag-eGFP fusion

protein that multimerizes at the endosomal and plasma membrane, inducing the formation of

virus-like particles packed with eGFP. Bottom: brightfield and eGFP pictures of HEK293T cells

after transfection, prior to the isolation of conditioned medium.

Supplementary Figure 2: Scanning electron microscopy of 300 kDa MWCO filters used in FAEVEr.

Blank: empty filter membrane with clear visualization of the 300 kDa MWCO filter membrane.

Synthetic particles: homogeneous synthetic particles used for NTA calibration were retained on

the filter after completing the FAEVEr protocol and serve as a positive control. rEV: retained rEV

particles after completing the FAEVEr protocol with a distinct delineation of the EV particles

against the 300 kDa MWCO membrane.

Supplementary Figure 3: SDS-PAGE with Coomassie staining indicates quantitative removal of

serum proteins with FAEVEr when TWEEN-20 is supplemented to the filter, indicating decreased

membrane fouling. In the PBS condition (no TWEEN-20) a faint band is observed at 65 kDa,

indicating that only a fraction of the bovine albumin is removed by the wash steps. This stands

in contrast to washing with buffers that are supplemented with of TWEEN-20, where more

intense bands are observed in the washes. Additionally, the lysate of PBS contains observable

more albumin material, further indicating that PBS alone fails to remove the bulk of

contaminants contrary to TWEEN-20 were the lysate does not contain such strong signal.

Supplementary Figure 4: Comparison of the identified proteins in standard fetal bovine serum

(FBS) and EV-depleted FBS (EDS). A) After completing FAEVEr and subjecting the lysate to LC-

MS/MS analysis, we observed a high number of identifications in FBS but not in EDS, indicating

a higher protein heterogeneity due to the presence of EV particles. B) EV markers or biogenesis

related proteins are virtually absent in EDS, yet highly abundant in FBS.

Supplementary Figure 5: Culturing MCF7 cells in 10% standard fetal bovine serum (FBS) or 10%

EV-depleted FBS (EDS) for three days, has no huge effect on the MCF7 cellular proteome. A)

Due to the EV depletion protocol, EDS contains approximately six times less protein material. B)

No significant changes in cell viability or cell count were observed. C) In both conditions, we

identified over 7,000 proteins from which only 10 were significantly differentially regulated (D).

Supplementary Figure 6: The cellular proteome of MCF7 cells cultured at different percentages

of EV-depleted FBS (EDS) changes between serum-starved (0% EDS for 24 h) and optimal

conditions (10% EDS for 48 h). A) Heatmap depicting the differentially regulated proteins between

the different conditions from which we further investigated two major clusters to evaluate the

contrast between 0% EDS and 10% EDS. Gene ontology (GO) analysis of the proteins up-regulated

in the serum-starved conditions (B) and optimal conditions (C).

H. List of abbreviations

DDA: Data dependent acquisition

DIA: Data independent acquisition

EC: extracellular

EDS: EV-depleted FBS

ESCRT: endosomal sorting complexes required for transport

EV: extracellular vesicle

FAEVEr: Filter-aided extracellular vesicle enrichment

FBS: fetal bovine serum

GO: Gene ontology

ILV: inter-luminal vesicles

LC-MS/MS: Liquid chromatography coupled tandem mass spectrometry

LFQ: label-free quantification

MVB: multivesicular body

MWCO: molecular weight cut-off

NTA: nanoparticle tracking analysis

PCA: principal component analysis

PES: Polyethersulfone

PM: plasma membrane

PSM: peptide-spectral match

rEV: recombinant EV

SDS: sodium dodecyl sulphate

SEC: size-exclusion chromatography

SEM: scanning electron microscopy

TEM: transmission electron microscopy

UC: ultracentrifugation

%CV: coefficient of variance

I. Ethical approval

Not applicable

J. Consent for publications

All co-authors have consented to publication

K. Data availability

L. Competing interest

The authors declare no conflict of interest.

M. Funding statement

K.G. and S.E. acknowledge support by a Strategic Basic Research project (EV-TRACE) from the Research Foundation -Flanders (FWO), project number S006319N.

N. Author contributions

JP planned, designed and performed the experiments and wrote the manuscript. JVDV, FDM and DDP assisted with the experiments. TVDS performed the HEK293T cell culture and production of

751 rEV material. FB performed the electron microscopy. All authors read and approved the

752 manuscript.

O. Acknowledgements

754 The authors thank all operators of the VIB Proteomics Core for their excellence and input. The

755 authors thank the lab of An Hendrix for kindly providing us with the required purified rEV

756 aliquots.

P. References

1. Dixson AC, Dawson TR, Di Vizio D, Weaver AM. Context-specific regulation of extracellular
vesicle biogenesis and cargo selection. Nat Rev Mol Cell Biol. 2023 Jul 1;24(7):454–76.

2. van Niel G, D'Angelo G, Raposo G. Shedding light on the cell biology of extracellular vesicles.
Nat Rev Mol Cell Biol. 2018 Apr 1;19(4):213–28.

3. Colombo M, Moita C, van Niel G, Kowal J, Vigneron J, Benaroch P, et al. Analysis of ESCRT
functions in exosome biogenesis, composition and secretion highlights the heterogeneity of
extracellular vesicles. J CELL Sci. 2013 Dec 15;126(24):5553–65.

4. Wei D, Zhan W, Gao Y, Huang L, Gong R, Wang W, et al. RAB31 marks and controls an ESCRT-
independent exosome pathway. Cell Res. 2021 Feb 1;31(2):157–77.

5. Mathieu M, Martin-Jaular L, Lavieu G, Théry C. Specificities of secretion and uptake of
exosomes and other extracellular vesicles for cell-to-cell communication. Nat Cell Biol. 2019
Jan 1;21(1):9–17.

6. Yu L, Zhu G, Zhang Z, Yu Y, Zeng L, Xu Z, et al. Apoptotic bodies: bioactive treasure left behind
by the dying cells with robust diagnostic and therapeutic application potentials. J
Nanobiotechnology. 2023 Jul 12;21(1):218.

7. Théry C, Witwer KW, Aikawa E, Alcaraz MJ, Anderson JD, Andriantsitohaina R, et al. Minimal
information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the
International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. J
Extracell Vesicles. 2018 Dec 1;7(1):1535750.

8. Cocucci E, Meldolesi J. Ectosomes and exosomes: shedding the confusion between
extracellular vesicles. Trends Cell Biol. 2015 Jun 1;25(6):364–72.

9. Hadisurya M, Lee ZC, Luo Z, Zhang G, Ding Y, Zhang H, et al. Data-Independent Acquisition
Phosphoproteomics of Urinary Extracellular Vesicles Enables Renal Cell Carcinoma Grade
Differentiation. Mol Cell Proteomics. 2023 May 1;22(5):100536.

10. Hoshino A, Costa-Silva B, Shen TL, Rodrigues G, Hashimoto A, Tesic Mark M, et al. Tumour
exosome integrins determine organotropic metastasis. Nature. 2015 Nov 1;527(7578):329–35.

11. Joshi BS, de Beer MA, Giepmans BNG, Zuhorn IS. Endocytosis of Extracellular Vesicles and
Release of Their Cargo from Endosomes. ACS Nano. 2020 Apr 28;14(4):4444–55.

12. Stahl PD, Raposo G. Extracellular Vesicles: Exosomes and Microvesicles, Integrators of
Homeostasis. Physiology. 2019 May 1;34(3):169–77.

13. Takahashi A, Okada R, Nagao K, Kawamata Y, Hanyu A, Yoshimoto S, et al. Exosomes maintain
cellular homeostasis by excreting harmful DNA from cells. Nat Commun. 2017 May
16;8(1):15287.

14. Amin S, Massoumi H, Tewari D, Roy A, Chaudhuri M, Jazayerli C, et al. Cell Type-Specific
Extracellular Vesicles and Their Impact on Health and Disease. Int J Mol Sci. 2024;25(5).

15. Hánělová K, Raudenská M, Masařík M, Balvan J. Protein cargo in extracellular vesicles as the
key mediator in the progression of cancer. Cell Commun Signal. 2024 Jan 10;22(1):25.

16. Hoffmann O, Wormland S, Bittner AK, Collenburg M, Horn PA, Kimmig R, et al. Programmed
death receptor ligand-2 (PD-L2) bearing extracellular vesicles as a new biomarker to identify
early triple-negative breast cancer patients at high risk for relapse. J Cancer Res Clin Oncol.
2023 Mar 1;149(3):1159–74.

17. Zhang XW, Qi GX, Liu MX, Yang YF, Wang JH, Yu YL, et al. Deep Learning Promotes Profiling of
Multiple miRNAs in Single Extracellular Vesicles for Cancer Diagnosis. ACS Sens [Internet].
2024 Mar 5; Available from: https://doi.org/10.1021/acssensors.3c02789

18. Dorado E, Doria ML, Nagelkerke A, McKenzie JS, Maneta-Stavrakaki S, Whittaker TE, et al.
Extracellular vesicles as a promising source of lipid biomarkers for breast cancer detection
in blood plasma. J Extracell Vesicles. 2024 Mar 1;13(3):e12419.

19. Ma D, Xie A, Lv J, Min X, Zhang X, Zhou Q, et al. Engineered extracellular vesicles enable high-
efficient delivery of intracellular therapeutic proteins. Protein Cell. 2024 Mar 22;pwae015.

20.   Attem J, Narayana RVL, Manukonda R, Kaliki S, Vemuganti GK. Small extracellular
vesicles loaded with carboplatin effectively enhance the cytotoxicity of drug-resistant cells
from Y79 cells-in vitro. Biomed Pharmacother. 2024 Apr 1;173:116403.

21. Lone SN, Nisar S, Masoodi T, Singh M, Rizwan A, Hashem S, et al. Liquid biopsy: a step closer
to transform diagnosis, prognosis and future of cancer treatments. Mol Cancer. 2022 Mar
18;21(1):79.

22.   K S, T D, M P. Small extracellular vesicles as a multicomponent biomarker platform in
urinary tract carcinomas. Front Mol Biosci [Internet]. 2022;9. Available from:
https://www.frontiersin.org/articles/10.3389/fmolb.2022.916666

23.   Gao X, Gao B, Li S. Extracellular vesicles: A new diagnostic biomarker and targeted drug
in osteosarcoma. Front Immunol [Internet]. 2022;13. Available from:
https://api.semanticscholar.org/CorpusID:252468251

24.   Hirschberg Y, Valle-Tamayo N, Dols-Icardo O, Engelborghs S, Buelens B, Vandenbroucke
RE, et al. Proteomic comparison between non-purified cerebrospinal fluid and cerebrospinal

fluid-derived extracellular vesicles from patients with Alzheimer's, Parkinson's and Lewy
body dementia. J Extracell Vesicles. 2023 Dec 1;12(12):12383.

25.   Vandendriessche C, Kapogiannis D, Vandenbroucke RE. Biomarker and therapeutic
potential of peripheral extracellular vesicles in Alzheimer's disease. Adv Drug Deliv Rev. 2022
Nov 1;190:114486.

26.   Newman LA, Muller K, Rowland A. Circulating cell-specific extracellular vesicles as
biomarkers for the diagnosis and monitoring of chronic liver diseases. Cell Mol Life Sci. 2022
Apr 10;79(5):232.

27.   Mazzucco M, Mannheim W, Shetty SV, Linden JR. CNS endothelial derived extracellular
vesicles are biomarkers of active disease in multiple sclerosis. Fluids Barriers CNS. 2022 Feb
8;19(1):13.

28.   Horvatovich PL, Bischoff R. Current Technological Challenges in Biomarker Discovery
and Validation. Eur J Mass Spectrom. 2010 Feb 1;16(1):101–21.

29.   Schiess R, Wollscheid B, Aebersold R. Targeted proteomic strategy for clinical biomarker
discovery. Mol Oncol. 2009 Feb 1;3(1):33–44.

30.   Burton JB, Carruthers NJ, Stemmer PM. Enriching extracellular vesicles for mass
spectrometry. Mass Spectrom Rev. 2023 Mar 1;42(2):779–95.

31. Demichev V, Messner CB, Vernardis SI, Lilley KS, Ralser M. DIA-NN: neural networks and
interference correction enable deep proteome coverage in high throughput. Nat Methods.
2020 Jan 1;17(1):41–4.

32.   Gessulat S, Schmidt T, Zolg DP, Samaras P, Schnatbaum K, Zerweck J, et al. Prosit:
proteome-wide prediction of peptide tandem mass spectra by deep learning. Nat Methods.
2019 Jun 1;16(6):509–18.

33.   Guzman UH, Martinez-Val A, Ye Z, Damoc E, Arrey TN, Pashkova A, et al. Ultra-fast label-
free quantification and comprehensive proteome coverage with narrow-window data-
independent acquisition. Nat Biotechnol [Internet]. 2024 Feb 1; Available from:
https://doi.org/10.1038/s41587-023-02099-7

34.   Yoshitake J, Azami M, Sei H, Onoshima D, Takahashi K, Hirayama A, et al. Rapid Isolation
of Extracellular Vesicles Using a Hydrophilic Porous Silica Gel-Based Size-Exclusion
Chromatography Column. Anal Chem. 2022 Oct 11;94(40):13676–81.

35.   Vanderboom PM, Dasari S, Ruegsegger GN, Pataky MW, Lucien F, Heppelmann CJ, et al.
A size-exclusion-based approach for purifying extracellular vesicles from human plasma.
Cell Rep Methods. 2021 Jul 26;1(3):100055.

36.   Onódi Z, Pelyhe C, Terézia Nagy C, Brenner GB, Almási L, Kittel Á, et al. Isolation of High-
Purity Extracellular Vesicles by the Combination of Iodixanol Density Gradient
Ultracentrifugation and Bind-Elute Chromatography From Blood Plasma. Front Physiol
[Internet]. 2018;9. Available from:
https://www.frontiersin.org/journals/physiology/articles/10.3389/fphys.2018.01479

37.   Gallart-Palau X, Serra A, Wong ASW, Sandin S, Lai MKP, Chen CP, et al. Extracellular
vesicles are rapidly purified from human plasma by PRotein Organic Solvent PRecipitation
(PROSPR). Sci Rep. 2015 Sep 30;5(1):14664.

38.   Rider MA, Hurwitz SN, Meckes DG. ExtraPEG: A Polyethylene Glycol-Based Method for
Enrichment of Extracellular Vesicles. Sci Rep. 2016 Apr 12;6(1):23978.

39.   Brambilla D, Sola L, Ferretti AM, Chiodi E, Zarovni N, Fortunato D, et al. EV Separation:
Release of Intact Extracellular Vesicles Immunocaptured on Magnetic Particles. Anal Chem.
2021 Apr 6;93(13):5476–83.

40.   Vergauwen G, Tulkens J, Pinheiro C, Avila Cobos F, Dedeyne S, De Scheerder MA, et al.
Robust sequential biophysical fractionation of blood plasma to study variations in the
biomolecular landscape of systemically circulating extracellular vesicles across clinical
conditions. J Extracell Vesicles. 2021 Aug 1;10(10):e12122.

41.Van Dorpe S, Lippens L, Boiy R, Pinheiro C, Vergauwen G, Rappu P, et al. Integrating automated
liquid handling in the separation workflow of extracellular vesicles enhances specificity and
reproducibility. J Nanobiotechnology. 2023 May 19;21(1):157.

42.   An M, Wu J, Zhu J, Lubman DM. Comparison of an Optimized Ultracentrifugation Method
versus Size-Exclusion Chromatography for Isolation of Exosomes from Human Serum. J
Proteome Res. 2018 Oct 5;17(10):3599–605.

43.   Turner NP, Abeysinghe P, Kwan Cheung KA, Vaswani K, Logan J, Sadowski P, et al. A
Comparison of Blood Plasma Small Extracellular Vesicle Enrichment Strategies for
Proteomic Analysis. Proteomes. 2022;10(2).

44.   Brennan K, Martin K, FitzGerald SP, O'Sullivan J, Wu Y, Blanco A, et al. A comparison of
methods for the isolation and separation of extracellular vesicles from protein and lipid
particles in human serum. Sci Rep. 2020 Jan 23;10(1):1039.

45.   Baranyai T, Herczeg K, Onódi Z, Voszka I, Módos K, Marton N, et al. Isolation of Exosomes
from Blood Plasma: Qualitative and Quantitative Comparison of Ultracentrifugation and Size
Exclusion Chromatography Methods. PLOS ONE. 2015 Dec 21;10(12):e0145686.

46.   Askeland A, Borup A, Østergaard O, Olsen JV, Lund SM, Christiansen G, et al. Mass-
Spectrometry Based Proteome Comparison of Extracellular Vesicle Isolation Methods:
Comparison of ME-kit, Size-Exclusion Chromatography, and High-Speed Centrifugation.
Biomedicines. 2020;8(8).

47.   Van Deun J, Mestdagh P, Agostinis P, Akay Ö, Anand S, Anckaert J, et al. EV-TRACK:
transparent reporting and centralizing knowledge in extracellular vesicle research. Nat
Methods. 2017 Mar 1;14(3):228–32.

48.   Geeurickx E, Lippens L, Rappu P, De Geest BG, De Wever O, Hendrix A. Recombinant
extracellular vesicles as biological reference material for method development, data
normalization and assessment of (pre-)analytical variables. Nat Protoc. 2021 Feb 1;16(2):603–
33.

49.   Geeurickx E, Tulkens J, Dhondt B, Van Deun J, Lippens L, Vergauwen G, et al. The
generation and use of recombinant extracellular vesicles as biological reference material.
Nat Commun. 2019 Jul 23;10(1):3288.

50.   Sidhom K, Obi PO, Saleem A. A Review of Exosomal Isolation Methods: Is Size Exclusion
Chromatography the Best Option? Int J Mol Sci. 2020;21(18).

51. Ding M, Wang C, Lu X, Zhang C, Zhou Z, Chen X, et al. Comparison of commercial exosome
isolation kits for circulating exosomal microRNA profiling. Anal Bioanal Chem. 2018 Jun
1;410(16):3805–14.

52. Royo F, Zuñiga-Garcia P, Sanchez-Mosquera P, Egia A, Perez A, Loizaga A, et al. Different
EV enrichment methods suitable for clinical settings yield different subpopulations of urinary
extracellular vesicles from human samples. J Extracell Vesicles. 2016 Jan 1;5(1):29497.

53. Lobb RJ, Becker M, Wen Wen S, Wong CSF, Wiegmans AP, Leimgruber A, et al. Optimized
exosome isolation protocol for cell culture supernatant and human plasma. J Extracell
Vesicles. 2015 Jan 1;4(1):27031.

54. Guan S, Yu H, Yan G, Gao M, Sun W, Zhang X. Characterization of Urinary Exosomes
Purified with Size Exclusion Chromatography and Ultracentrifugation. J Proteome Res. 2020
Jun 5;19(6):2217–25.

55. Gámez-Valero A, Monguió-Tortajada M, Carreras-Planella L, Franquesa M, Beyer K,
Borràs FE. Size-Exclusion Chromatography-based isolation minimally alters Extracellular
Vesicles' characteristics compared to precipitating agents. Sci Rep. 2016 Sep 19;6(1):33641.

56. Vergauwen G, Dhondt B, Van Deun J, De Smedt E, Berx G, Timmerman E, et al.
Confounding factors of ultrafiltration and protein analysis in extracellular vesicle research.
Sci Rep. 2017 Jun 2;7(1):2704.

57. Jordan KR, Hall JK, Schedin T, Borakove M, Xian JJ, Dzieciatkowska M, et al. Extracellular
vesicles from young women's breast cancer patients drive increased invasion of non-
malignant cells via the Focal Adhesion Kinase pathway: a proteomic approach. Breast Cancer
Res. 2020 Nov 23;22(1):128.

58. Allelein S, Medina-Perez P, Lopes ALH, Rau S, Hause G, Kölsch A, et al. Potential and
challenges of specifically isolating extracellular vesicles from heterogeneous populations.
Sci Rep. 2021 Jun 2;11(1):11585.

59. Eyckerman S, Titeca K, Van Quickelberghe E, Cloots E, Verhee A, Samyn N, et al. Trapping
mammalian protein complexes in viral particles. Nat Commun. 2016 Apr 28;7(1):11416.

60. Wang J, Vasaikar S, Shi Z, Greer M, Zhang B. WebGestalt 2017: a more comprehensive,
powerful, flexible and interactive gene set enrichment analysis toolkit. Nucleic Acids Res. 2017
Jul 3;45(W1):W130–7.

61. HaileMariam M, Eguez RV, Singh H, Bekele S, Ameni G, Pieper R, et al. S-Trap, an Ultrafast
Sample-Preparation Approach for Shotgun Proteomics. J Proteome Res. 2018 Sep
7;17(9):2917–24.

62. Ono A. HIV-1 assembly at the plasma membrane: Gag trafficking and localization. Future
Virol. 2009 May 1;4(3):241–57.

63. Chhuon C, Zhang SY, Jung V, Lewandowski D, Lipecka J, Pawlak A, et al. A sensitive S-
Trap-based approach to the analysis of T cell lipid raft proteome. J Lipid Res. 2020 Nov
1;61(11):1512–23.

64. Elinger D, Gabashvili A, Levin Y. Suspension Trapping (S-Trap) Is Compatible with Typical
Protein Extraction Buffers and Detergents for Bottom-Up Proteomics. J Proteome Res. 2019
Mar 1;18(3):1441–5.

65.   Ludwig KR, Schroll MM, Hummon AB. Comparison of In-Solution, FASP, and S-Trap
Based Digestion Methods for Bottom-Up Proteomic Studies. J Proteome Res. 2018 Jul
6;17(7):2480–90.

66.   Varnavides G, Madern M, Anrather D, Hartl N, Reiter W, Hartl M. In Search of a Universal
Method: A Comparative Survey of Bottom-Up Proteomics Sample Preparation Methods. J
Proteome Res. 2022 Oct 7;21(10):2397–411.

67.   Osteikoetxea X, Sódar B, Németh A, Szabó-Taylor K, Pálóczi K, Vukman KV, et al.
Differential detergent sensitivity of extracellular vesicle subpopulations. Org Biomol Chem.
2015;13(38):9775–82.

68.   Osteikoetxea X, Balogh A, Szabó-Taylor K, Németh A, Szabó TG, Pálóczi K, et al. Improved
Characterization of EV Preparations Based on Protein to Lipid Ratio and Lipid Properties.
PLOS ONE. 2015 Mar 23;10(3):e0121184.

69.   Choi DS, Kim DK, Kim YK, Gho YS. Proteomics, transcriptomics and lipidomics of
exosomes and ectosomes. PROTEOMICS. 2013 May 1;13(10–11):1554–71.

70.   Subra C, Laulagnier K, Perret B, Record M. Exosome lipidomics unravels lipid sorting at
the level of multivesicular bodies. Recent Dev Biosynth Storage Transp Lipids. 2007 Feb
1;89(2):205–12.

71. Minhua Feng, Berdugo Morales A, Poot A, Beugeling T, Bantjes A. Effects of Tween 20 on the
desorption of proteins from polymer surfaces. J Biomater Sci Polym Ed. 1996 Jan 1;7(5):415–
24.

72.   Amirilargani M, Saljoughi E, Mohammadi T. Improvement of permeation performance of
polyethersulfone (PES) ultrafiltration membranes via addition of Tween-20. J Appl Polym Sci.
2010 Jan 5;115(1):504–13.

73.   Lee KJ, Mower R, Hollenbeck T, Castelo J, Johnson N, Gordon P, et al. Modulation of
Nonspecific Binding in Ultrafiltration Protein Binding Studies. Pharm Res. 2003 Jul
1;20(7):1015–21.

74.   Urzì O, Olofsson Bagge R, Crescitelli R. The dark side of foetal bovine serum in
extracellular vesicle studies. J Extracell Vesicles. 2022 Oct 1;11(10):e12271.

75.   Drabovich AP, Pavlou MP, Dimitromanolakis A, Diamandis EP. Quantitative Analysis of
Energy Metabolic Pathways in MCF-7 Breast Cancer Cells by Selected Reaction Monitoring
Assay *. Mol Cell Proteomics. 2012 Aug 1;11(8):422–34.

76.   Nie S, McDermott SP, Deol Y, Tan Z, Wicha MS, Lubman DM. A quantitative proteomics
analysis of MCF7 breast cancer stem and progenitor cell populations. PROTEOMICS. 2015 Nov
1;15(22):3772–83.

77.   Leidal AM, Huang HH, Marsh T, Solvik T, Zhang D, Ye J, et al. The LC3-conjugation
machinery specifies the loading of RNA-binding proteins into extracellular vesicles. Nat Cell
Biol. 2020 Feb 1;22(2):187–99.

78.   Fujita N, Itoh T, Omori H, Fukuda M, Noda T, Yoshimori T. The Atg16L Complex Specifies
the Site of LC3 Lipidation for Membrane Biogenesis in Autophagy. Mol Biol Cell. 2008 May
1;19(5):2092–100.

79.   Matsushita M, Suzuki NN, Obara K, Fujioka Y, Ohsumi Y, Inagaki F. Structure of Atg5·Atg16,
a Complex Essential for Autophagy *. J Biol Chem. 2007 Mar 2;282(9):6763–72.

80.   Urbańska K, Orzechowski A. The Secrets of Alternative Autophagy. Cells. 2021;10(11).

81.Salimi L, Akbari A, Jabbari N, Mojarad B, Vahhabi A, Szafert S, et al. Synergies in exosomes
and autophagy pathways for cellular homeostasis and metastasis of tumor cells. Cell Biosci.
2020 May 13;10(1):64.

82.   Pirkmajer S, Chibalin AV. Serum starvation: caveat emptor. Am J Physiol–Cell Physiol.
2011 Aug 1;301(2):C272–9.

83.   Shin J, Rhim J, Kwon Y, Choi SY, Shin S, Ha CW, et al. Comparative analysis of differentially
secreted proteins in serum–free and serum–containing media by using BONCAT and pulsed
SILAC. Sci Rep. 2019 Feb 28;9(1):3096.

5% TWEEN-20
8.2e17 particles/mL