Proximal termination generates a transcriptional state that determines the

rate of establishment of Polycomb silencing

Govind Menon[1,†], Eduardo Mateo-Bonmati[2,5,†], Svenja Reeck[2], Robert Maple[2], Zhe Wu[3], Robert

Ietswaart[4], Caroline Dean[2,*], Martin Howard[1,*]

[1]Department of Computational and Systems Biology, John Innes Centre, Norwich Research Park NR4

7UH, UK

[2]Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park NR4

7UH, UK

[3]Key Laboratory of Molecular Design for Plant Cell Factory of Guangdong Higher Education

Institutes, Institute of Plant and Food Science, Department of Biology, School of Life Sciences,

Southern University of Science and Technology, Shenzhen, China

[4]Harvard Medical School, Department of Genetics, 77 Avenue Louis Pasteur, Boston, MA 02115,

USA

[5]Current address: Centro de Biotecnología y Genómica de Plantas, Universidad Politécnica de Madrid

(UPM) – Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA)/CSIC, 28223

Pozuelo de Alarcón, Madrid, Spain

[†]These authors contributed equally to this work

[*]Co-corresponding authors: caroline.dean@jic.ac.uk, martin.howard@jic.ac.uk

Abstract

Chromatin-mediated transcriptional silencing by Polycomb Repressive Complex 2 (PRC2) is critical

for gene regulation in development and environmental responses. However, the mechanism and

timescales controlling de novo establishment of PRC2 silencing are unclear. Here, we investigate

PRC2 silencing at Arabidopsis *FLOWERING LOCUS C* (*FLC*), known to involve co-transcriptional

RNA processing, histone demethylation activity, and PRC2 function; but so far not mechanistically

connected. We develop and then test a computational model that describes how proximal

polyadenylation/termination mediated by the RNA binding protein FCA induces H3K4me1 removal

by the histone demethylase FLD. H3K4me1 removal feeds back to reduce RNA Pol II processivity

and thus enhance early termination, thereby repressing productive transcription. The model predicts

that this transcription-coupled repression controls the level of transcriptional antagonism to Polycomb

action, Thus, the effectiveness of this repression dictates the timescale for establishment of Polycomb

H3K27me3 silencing. Experimental validation of these model predictions allowed us to

mechanistically connect co-transcriptional processing to setting the level of productive transcription at

the locus, which then determines the rate of the ON to OFF switch to PRC2 silencing.

Introduction

Transcriptional regulation is often mediated by chromatin states that are key to developmental

changes and environmental responses in many organisms. These chromatin states are characterised by

histone modifications, and trans-factors that recognise these modifications, which together influence

the level of productive transcription. Such chromatin states can be heritable across DNA replication

and mitosis, with controlled switching of key developmental loci between active and silenced states

underpinning the changes in gene expression programs that drive differentiation.

Polycomb Repressive Complex 2 (PRC2) mediated H3K27me3 is a system that generates such

heritable and switchable chromatin states. The mechanisms enabling inheritance in this case include

read-write reinforcement of existing histone modifications and antagonism between transcriptional

activity and silencing modifications (Berry et al., 2017; Dodd et al., 2007; Holoch et al., 2021). Such

mechanisms can generate bistability producing all or nothing (ON or OFF) transcriptional states

(Berry et al., 2017; Pease et al., 2021). Each gene copy can be stably maintained in either an ON or

OFF state even in dividing cells and copies can be independently switched between states. Such

switching has been achieved through experimental perturbations in various studies (Holoch et al.,

2021), but there are few examples where the mechanisms that enable the set-up and switching of these

states are studied in natural developmental contexts.

One such example is PRC2-mediated regulation at the Arabidopsis floral repressor gene

FLOWERING LOCUS C (*FLC*) (Menon et al., 2021). The winter cold-induced epigenetic silencing is

an ON/OFF cell-autonomous, stochastic switching of individual *FLC* copies into a Polycomb silenced

state (Fig. 1A) (Angel et al., 2011; Berry et al., 2015; Csorba et al., 2014; Lövkvist and Howard,

2021; Menon et al., 2021). In this process, the silencing histone modification H3K27me3 is first

nucleated (deposited at a specific intragenic site) during the cold, followed by spreading of

H3K27me3 across the locus through a cell-cycle dependent mechanism during growth after cold

(Yang et al., 2017). Establishment of Polycomb silencing requires specific Polycomb accessory

proteins and chromatin remodellers, some specific to the nucleation or spreading phases.

In rapid-cycling Arabidopsis accessions *FLC* silencing is constitutive, occurring early in

development. This silencing requires the RNA binding protein FCA, which interacts with Cleavage

and Polyadenylation Specificity Factor (CPF) machinery (Simpson et al., 2003). FCA promotes

proximal termination of *FLC* sense transcription in the developing embryo (Schon et al., 2021), and of

FLC antisense transcription in the seedling (Liu et al., 2010). *FLC* silencing also requires FLD (Liu et al., 2007), a plant homologue of the conserved mammalian histone K4 demethylase LSD1 (Martignago et al., 2019). The role of FCA and other co-transcriptional regulators in this mode of *FLC* silencing, and the observed FCA interactions with termination and 3' processing factors, suggests a transcription-coupled mechanism that links to Polycomb silencing. Indeed, the accompanying paper identifies a role for the CPF phosphatase module, a key regulator of RNA Pol II termination, in this mode of *FLC* silencing (Mateo-Bonmatí et al., 2023). This mechanism thus has the potential to provide a window into the link between transcription-coupled repression and Polycomb silencing.

Transcription-coupled mechanisms have been implicated in the establishment of heterochromatin in many organisms, through 3' processing and non-coding RNA. A recent study indicates a wide-ranging role for 3' processing in heterochromatin establishment in *S. pombe* (Shimada et al., 2021). For example, Clr4-mediated heterochromatin assembly by the RNAi-independent pathway involves non-canonical polyadenylation and 3' processing factors (Chalamcharla et al., 2015; Tucker et al., 2016; Vo et al., 2019). Transcription termination at non-canonical sites mediated by the conserved cleavage and polyadenylation factor (CPF) and the RNA-binding protein Mmi1 have been shown to be necessary for Clr4-driven assembly of facultative heterochromatin at meiotic and environmental response genes. Another RNA binding protein, Seb1, has been shown to promote heterochromatin formation, through a mechanism linked to termination of noncoding transcription (Marina et al., 2013; Wittmann et al., 2017). For PRC2-mediated silencing, early termination has been suggested to reduce nascent RNA inhibition of PRC2, by limiting the physical length of the transcript produced (Kaneko et al., 2014). Nevertheless, the exact mechanisms linking co-transcriptional processing to chromatin regulation are still poorly understood.

Exploiting an allelic series of *fca* mutations, we recently showed that the FCA-mediated mechanism can generate a graded regulation of *FLC* expression (Antoniou-Kourounioti et al., 2023). Thus, *FLC* can be regulated both in a graded manner (or *analogue* regulation) or by an ON/OFF mode of regulation (or *digital* regulation) as in the cold-induced silencing (Fig. 1A). The intermediate *fca-3* allele revealed that FCA-mediated analogue regulation precedes an ON/OFF developmental switch at *FLC* (Fig. 1B)(Antoniou-Kourounioti et al., 2023). Thus, the study highlighted the switch from analogue to digital regulation but did not mechanistically address HOW that switch occurred. In this study, we therefore asked: How can co-transcriptional processing mediated by FCA and the linked function of the chromatin modifier FLD achieve the observed *analogue* mode/graded manner of *FLC* regulation? Does the observed ON to OFF switch involve PRC2? If so, what is the mechanistic link between this analogue transcription-coupled mode of regulation and a Polycomb ON/OFF *digital* mode of regulation? Specifically, how is an FCA/FLD mediated *repressed* state (with enhanced early

termination of transcription enabling graded regulation of the amount of full length transcript

produced) linked to a PRC2 mediated *silenced* state (OFF state with a low frequency of transcription

initiation)?

Using mathematical modelling combined with molecular approaches, we dissect the interactions

between co-transcriptional processing and chromatin regulation that underpin how transcription-

coupled repression can lead to Polycomb-mediated silencing of *FLC*. Specifically, the model

describes sense and antisense transcription, alternative transcriptional termination, the dynamics of

activating (H3K4me1) and silencing (H3K27me) modifications, and the interactions between these

components (Fig. 1C). Importantly, the complexity of these feedbacks could not be interpreted

without the use of modelling approaches. We first showed that the model can successfully generate

intermediate transcriptional outputs at non-Polycomb silenced *FLC* copies (analogue/graded

regulation), while also allowing Polycomb nucleation to occur slowly over time (ON/OFF switch).

We then used the model to qualitatively predict changes in activating H3K4me1 and silencing

H3K27me3 modifications at *FLC* over time, and experimentally validated these predictions by

measuring changes in these modifications over a developmental time-course, showing further that the

model can be quantitatively fit to capture these changes. Further validation was provided by

measuring timecourse changes in *FLC* expression, where the model parameter fit obtained from

histone modification changes also achieves a satisfactory quantitative match with no additional

changes. The separate roles of the co-transcriptional repression and Polycomb silencing as predicted

by the model are further validated by examining a mutant of a Polycomb methyltransferase. The

model clearly elaborates how FCA/FLD-mediated transcription-coupled repression, induced by co-

transcriptional processing steps as described in the accompanying paper (Mateo-Bonmatí, 2023), can

influence the timing of the digital switch to Polycomb silencing. The combination of molecular

genetics and the theoretical modelling described here provide a quantitative description of how co-

transcriptional processing can quantitatively set transcriptional states that determine rates of

Polycomb silencing, a mechanism that is likely to be generally relevant in many organisms.

Results

Molecular framework governing the *FLC* transcription-coupled chromatin repression pathway

The RNA binding protein FCA promotes proximal polyadenylation/termination at many targets in the

Arabidopsis genome (Sonmez et al., 2011). FCA is known to directly interact *in vivo* with FY, a CPF

component needed for silencing homologous to yeast Pfs2 and mammalian WDR33 (Simpson et al.,

2003). FCA also indirectly interacts with other cleavage and polyadenylation specificity factors (Fang

et al., 2019), as well as the 3' processing factors CstF64 and CstF77 (Liu et al., 2010). FCA-mediated

silencing also requires the involvement of CDKC2, a component of the transcription elongation factor

P-TEFb (Wang et al., 2014), as well as the core spliceosome component PRP8 (Marquardt et al.,
2014), suggesting it is the process of transcription, co-transcriptional processing, and proximal
termination, rather than just the transcripts themselves, that drive silencing. FCA is proposed to
recognize any situation where RNA Pol II has stalled and transcription termination is required to clear
a 'tangle' in the chromatin (Baxter et al., 2021; Xu et al., 2021).

FLC silencing mediated by FCA requires the protein FLD, a homologue of the mammalian H3K4/K9
demethylase LSD1 (Liu et al., 2007). FLD is thought to demethylate H3K4me1 in Arabidopsis
(Inagaki et al., 2021; Martignago et al., 2019). FLD has been shown to associate *in vivo* with the SET
domain protein SDG26, which interacts with FY, indicating a physical link between processing of
antisense transcripts and chromatin remodelling (Fang et al., 2020). FLD also physically associates in
vivo with LD, a TFII-S domain containing protein and homolog of the CPF phosphatase subunit,
PNUTS in human and Ref2 in yeast. This strong association of chromatin modifiers such as FLD with
RNA Pol II regulators shown in the accompanying paper, identifies a mechanism where FCA-
mediated proximal cleavage and polyadenylation triggers FLD-mediated removal of H3K4me1
(Mateo-Bonmatí, 2023).The Arabidopsis H3K36 methyltransferase SDG8 is known to recognise
H3K4me1, so H3K4me1 removal might be expected to affect H3K36 methylation. *FLC* upregulation
in an *fca* mutant is rescued in an *sdg8 fca* double mutant (Fang et al., 2020), indicating that
H3K36me3 may be essential to set up an active transcriptional state at *FLC*. This is consistent with
our previous ChIP-qPCR data at *FLC,* showing high levels of H3K4me1 and H3K36me3 at the locus
in a high transcriptional state and low levels of both these modifications in a low transcriptional state
(Fang et al., 2020). Recent evidence from *Saccharomyces cerevisiae,* shows that H3K36me3 is
required for the recruitment of splicing factors and effective co-transcriptional splicing (Leung et al.,
2019). Based on this evidence, FLD-mediated removal of H3K4me1 may cause reduced H3K36
methylation, thus inhibiting co-transcriptional splicing, consequently reducing Pol II processivity and
promoting further early transcription termination. The FLD mechanism would thus form a feedback
loop where early termination enhances itself through H3K4me1 removal. The above molecular
mechanisms and possible feedback interaction are summarised in the schematic shown in Fig. 1C.

FCA promotes premature termination of both sense and antisense transcription at *FLC*
The *FLC* sense proximal transcripts are detectable in young embryos (Schon et al., 2021) but less so
in vegetative tissue, likely due to rapid nascent RNA turnover. In contrast, the *FLC* antisense RNAs
transcribed from the 3' end of *FLC* (collectively called *COOLAIR*), which are alternatively
polyadenylated (PA) at a proximal PA site within the *FLC* gene or a distal PA site in the vicinity of
the *FLC* promoter (Liu et al., 2010; Swiezewski et al., 2009), are both readily detectable. The
mechanism underlying the differential turnover of proximally polyadenylated transcripts is currently
unknown, but differential fates of coding and non-coding transcripts are not uncommon (Nojima and

Proudfoot, 2022). To assess the relative roles of FCA-mediated premature termination of

sense/antisense transcripts in silencing, we compared repression of a wild-type *FLC* transgene (*FLC-*

15), with that of an *FLC* transgene where the antisense promoter has been replaced (*TEX 1.0*) altering

the type and quantity of antisense transcript production (Csorba et al., 2014; Marquardt et al., 2014;

Wang et al., 2014). We performed the comparison in two genetic backgrounds: one with the

endogenous FCA carrying a null mutation (*fca-9* mutant in *Col-0*), thus having no functional FCA,

and the other with the same mutant of the endogenous FCA, but carrying a transgenic 35S::FCA

construct (Liu et al., 2007), giving FCA overexpression. We made two observations (Fig. 2A): (i) the

wild-type *FLC* transgene is significantly repressed by FCA overexpression (as previously reported

(Liu et al., 2007)); (ii) this FCA dependent repression of the wild-type *FLC* transgene is significantly

attenuated ~10 fold in *TEX 1.0,* but repression is not lost completely. Thus, the antisense transcription

mechanism only partially accounts for *FLC* repression; a fact supported by a CRISPR deletion of the

antisense promoter at the endogenous *FLC* locus causing a small *FLC* upregulation (Zhao et al.,

2021). To further explore the importance of FCA-mediated proximal termination we analysed

production of sense polyadenylated *FLC* transcripts by Quant-seq (Moll et al., 2014) in the *fca* allelic

series (Antoniou-Kourounioti et al., 2023; Koornneef et al., 1991).Three genotypes give different

levels of *FLC* expression at the whole plant level: the wild-type L*er*, with low *FLC* expression, the

fca-1 mutant, with high *FLC* expression, and the *fca-3* mutant, with intermediate *FLC* expression.

Proximally polyadenylated transcripts for the *FLC* transcript were detected for all genotypes analysed

at sites close to those reported in embryos (Fig. 2B) (Schon et al., 2021). Altogether these data

support a model whereby FCA-mediated proximal termination of both sense and antisense

transcription contributes to repression of *FLC* transcription.

Computational model of *FLC* chromatin silencing through alternative termination

The silenced state of *FLC* is associated with H3K27me3 coverage of the whole locus (Wu et al.,

2016). Genetic analysis also shows that the plant Polycomb methyltransferase CURLY LEAF (CLF)

is essential for FCA-mediated silencing of *FLC* (Fang et al., 2020). However, despite the detailed

molecular data on 3' processing-mediated silencing described above, and in the accompanying paper

(Mateo-Bonmatí, 2023), how the various elements linked to Polycomb silencing was unknown, due in

part to the complex feedbacks involved. To bridge this gap, we therefore constructed a computational

model based on the above features (Fig. 3A), using our prior knowledge of Polycomb dynamics

known to generate digital ON/OFF memory states. We then used this model to dissect the underlying

mechanism of the transition elucidating how digital silencing might be integrated into FCA-mediated

3' processing. Important features of the new model are described below:

The model includes both activating (H3K4me1) and repressive (H3K27me3) modifications at each

histone across the *FLC* locus, assumed mutually exclusive on the same H3 tail, consistent with the

reported mutual exclusivity of these modifications (Shema et al., 2016). The model also incorporates sense and antisense transcription at the locus. Importantly, the model assumes distinct roles for the two kinds of modifications in transcription regulation: H3K4me1 is assumed to affect RNA Pol II processivity rather than transcription initiation/transition to elongation. Higher overall coverage of H3K4me1 across the locus therefore promotes distal termination of transcription by reducing the probability of early termination. H3K4me1 modifications are added co-transcriptionally, allowing distal termination to feedback onto further H3K4me1 addition. Conversely, proximal termination of both sense and antisense transcription leads to removal of H3K4me1, in line with the potential function of FLD as a demethylase, allowing feedback onto further proximal termination. Higher H3K27me3 coverage, rather than affecting processivity, is assumed to decrease the frequency of both sense and antisense transcription events (representing either transcription initiation or transition to productive elongation). The model describes constitutive deposition of H3K27me by Polycomb in the *FLC* nucleation region. H3K27me spreads to the rest of the locus through interactions with the nucleation region, through read-write feedback interactions. An experimentally determined gene loop mediating an interaction between the 5' and 3' ends of *FLC* is also incorporated in the model. Finally, in line with experimental evidence from mammalian systems and previously developed models for Polycomb, the model assumes that productive transcription promotes removal of H3K27me through both histone exchange and co-transcriptional demethylation, thereby implementing antagonism between transcription and Polycomb silencing. These aspects are integrated into a fully stochastic spatiotemporal simulation of the *FLC* locus, where transcription events (sense and antisense) and histone modification levels (H3K4me1 and H3K27me0/1/2/3) are simulation outputs. Full details of the model can be found in the Methods and Supplementary Information.

FCA and FLD mediate a transcription-coupled repression mechanism that is sufficient to generate analogue regulation

Our previous analysis of the *fca* allelic series highlighted that differences in overall expression between genotypes appeared to have two components – a digital ON/OFF mode where we observed clear differences in the number of *FLC* OFF cells, and an analogue mode, with quantitatively different levels of *FLC* in the ON cells (Fig. 1B). Intermediate expression was associated with a slower switch off over time into the digital OFF state. Importantly, we also observed similar behaviour of *FLC* expression in single cells for a different intermediate mutant of *FCA*, called *fca-4* (Fig. S5C and (Antoniou-Kourounioti et al., 2023)), showing effects are not allele-specific, but related to the regulatory function of FCA at *FLC*. Overall, this demonstrates that FCA quantitatively modulates transcription and influences the timing of the switch into the digital OFF state.

We first tried to see if the model could capture the quantitative variation in *FLC* expression (analogue mode) in the absence of the Polycomb module (see model schematic in Fig. 3A). To represent the

difference in FCA function between the three genotypes, we used a single parameter controlling the
probability of FCA mediated proximal termination of transcription. We then simulated the model with
different values for this parameter representing different levels of functionality of the FCA protein.
We set the parameter to its highest value for L*er*, an intermediate value for *fca-3*, and lowest for *fca-1*
(Fig. 3B). With these parameter settings we found that the model, with just transcription-coupled
repression (Fig. 3A) was able to recapitulate the analogue mode of regulation, with the simulations
showing the lowest frequency of productive transcription events for L*er*, highest for *fca-1*, and
intermediate for *fca-3* (Fig. 3C). Consistent with the relationship between productive transcription and
H3K4me1 in the model, the levels of this modification were also predicted to be lowest in L*er,*
intermediate in *fca-3* and highest in *fca-1* (Fig. 3C). While the model simulation output shown in Fig.
3 is qualitatively consistent with our previous observations, note that the specific model parameter
values used here are chosen to obtain a quantitative fit for the full model to experimentally measured
histone modification levels as we explain below.

Model predicts differences in the rate of Polycomb silencing establishment in the different
genotypes
With the analogue module constrained in this way, we simulated the full model, including the
Polycomb digital switching module (Fig. 3A). Importantly, this model recapitulated the
experimentally observed reduction in *FLC* transcriptional output (Fig. 3D), as well as the digital
switch to stable silencing (one-way switch) over time (Fig. 3E) observed using *FLC*-Venus imaging
in root meristematic cells (Antoniou-Kourounioti et al., 2023). For L*er*, where productive
transcription is lowest, the model predicted a rapid switch to the digital OFF/Polycomb silenced state
(Fig. 3E, Fig. S4B(i)). For *fca-3*, the model predicted a slow switch to the digital OFF state, while
even slower switching is predicted for *fca-1* (Fig. 3E, Fig. S4B(i)). In all cases, the switching was
predicted to be one-way, with the Polycomb silenced state being highly stable.

Thus, the combined model of transcription-coupled repression and Polycomb silencing predicts that
the level of productive transcriptional activity set by FCA/co-transcriptional silencing/FLD (analogue
module) determines the level of transcriptional antagonism to H3K27me3. The level of transcriptional
antagonism in turn dictates the rate of establishment of the Polycomb silenced state.

Model predictions are consistent with observed chromatin changes over time
We next used the combined model to predict population levels of H3K4me1 and H3K27me3 at *FLC*
in the three genotypes, predicting low H3K4me1 and high H3K27me3 levels in L*er*, intermediate
levels of both modifications in *fca-3*, and high H3K4me1 and low H3K27me3 in *fca-1*. Starting from
an active chromatin state with full H3K4me1 coverage, the model also predicted a progressive
reduction in H3K4me1 over time in all genotypes*,* with a low steady state level quickly attained in

L*er,* and a slower reduction in *fca-3* and *fca-1,* with *fca-1* being the slowest (see red bars in Fig. 4D).

In all genotypes*,* this reduction is combined with an accumulation of H3K27me3 as more loci switch

to the Polycomb silenced state (see red bars in Fig. 4C).

To validate these predictions, we performed a time course measurement of silencing H3K27me3 and

the activating modifications H3K4me1 and H3K36me3 over the *FLC* locus by chromatin

immunoprecipitation followed by quantitative PCR (ChIP-qPCR), examining their levels in seedlings

at 7, 14 and 21 days after sowing. For each genotype and timepoint, we use the Polycomb silenced

STM locus as the positive control for H3K27me3 and the actively expressed *ACT7* locus as the

positive control for H3K4me1 (conversely *STM* and ACT7 function as negative controls for the

activating modifications and H3K27me3 respectively). See Fig. S4C(ii,iii,iv) for a comparison of the

levels measured at *FLC* and these control loci at the 7 day timepoint. Consistent with model

predictions, at all timepoints we see that *fca-3* shows intermediate levels of H3K4me1 and

H3K27me3, as compared to *fca-1* and Ler, with primers across the *FLC* locus showing a consistent

trend (Fig. 4A,B). This trend is reinforced by a statistical comparison between the three genotypes of

the averaged H3K27me3 and H3K4me1 levels across the locus (Fig. S4A (i,ii)).

For validating the model using the timecourse ChIP data we tried to fit the model (adjusting

parameters for the transcription-coupled repression mechanism) so that it captures the observed

trends. The fitted parameters are indicated in the supplementary tables (Tables 1-4). We note that a

direct quantitative comparison between the model predictions and the ChIP-qPCR data is difficult

because the simplest model would assume that the switching dynamics happens at all loci, while the

ChIP-qPCR analysis uses whole seedling tissue, including tissue where the switching may be

inhibited by other means. To account for this possibility in the model, when comparing the model

predictions to the ChIP-seq data, we include a subpopulation of 'active' *FLC* copies with full

H3K4me1 coverage and no H3K27me3 coverage, to compute a corrected model predicted level of

both modifications. The changes in histone modifications over time are then qualitatively consistent

with the model (Fig. 4C,D). Experimental H3K27me3 levels rise in *fca-3* and *fca-1* (the increase is

statistically significant in *fca-3* between 14 and 21 days, though narrowly not for *fca-1*: p=0.048, sig.

threshold=0.0253 (Holm-Sidak corrected), Fig. S4A(iii)). H3K4me1 levels fall in *fca-3* and *fca-1* over

time (statistically significant for both *fca-3* and *fca-1* between 14 and 21 days, Fig S4A(iv)) as more

loci switch to the digital OFF state. In L*er*, H3K4me1 shows essentially no change over the

experimental timepoints, while the H3K27me3 also does not show a statistically significant change on

average across the locus (Fig. S4A(iii, iv)). The lack of significant change in experimentally measured

H3K27me3 and H3K4me1 in L*er* are consistent with steady state levels having been reached by the 7-

day timepoint. The model predicts faster dynamics of H3K4me1 removal (through more frequent

proximal termination), and faster establishment of H3K27me3 in L*er,* so a rapid approach to steady

state is consistent. The activating H3K36me3 also shows a similar pattern, with *fca-3* exhibiting

intermediate levels, and the overall levels decreasing over time (Fig. S4B(iii)).

Overall, the combined model of transcription-coupled repression – mediated by proximal termination

of transcription, and Polycomb silencing, successfully predicts both the relative levels of the

modifications between genotypes and how these change over time.

Model predicts relative differences in proximal termination between genotypes

To validate the model predicted *FCA* dependent differences in proximal termination, we compared

these predictions to our Quant-seq data (data shown in Fig. 2B). We compared the predicted ratio of

proximal to distal termination events to the measured PA reads. To compute this ratio for the Quant-

seq data, we defined a Proximal termination cluster covering a range of termination sites close to the

5' end of the gene. Our definition of the proximal termination cluster is based on the proximal

termination of *FLC* sense transcripts we have previously reported (Schon et al., 2021). The genomic

coordinates are provided in a supplementary text file: FLC_termination_clusters.txt. A quantitative fit

of these ratios is not attempted here, as we do not know turnover rates of the different transcripts.

However, the predicted trends in the differences in the ratio of proximal to distal polyadenylation of

sense transcripts between the *FCA* mutants and the parental genotype qualitatively matched the trend

for the measured ratios of *FLC* sense transcription in our Quant-seq dataset (Fig. 4E,F; Fig. S4C(i),

see Fig. S4D for individual replicates).

Disrupting a Polycomb methyltransferase compromises the one-way switch to a Polycomb OFF

state but does not disrupt FCA mediated regulation

The model predicts that the transcription coupled repression mechanism is sufficient to generate the

analogue repression and that this repression controls the antagonism of transcription to Polycomb,

thus enabling the switch to Polycomb digital silencing. We tested these predictions using a Polycomb

mutant – a loss of function mutant of *CLF*. This methyltransferase is known to be required for

spreading of H3K27me3 from Polycomb nucleation peaks at many Arabidopsis targets (Shu et al.,

2019), and specifically at *FLC*, both in developmental (Shu et al., 2019) and cold induced silencing

contexts (Yang et al., 2017). This mutant is not expected to completely disrupt the ON/OFF silencing

mode, as H3K27me3 nucleation alone, which still occurs in *clf*, gives a metastable Polycomb silenced

state (Yang et al., 2017). However, only after the H3K27me3 spreads across the whole locus is the

Polycomb silenced state fully stable in rapidly dividing cells (Yang et al., 2017). Thus, the

maintenance of the OFF state is expected to be compromised in *clf* (Fig. 5A, B). Therefore, in a *clf*

mutant background, the model makes two predictions: (i) that the continued slow reduction in overall

FLC expression seen in L*er, fca-3*, and *fca-1*, i.e., an overall reduction in the number of actively

transcribed *FLC* copies – which is a consequence of a fully stable Polycomb digital switch – will be

missing in a *clf* mutant background, due to leakage out of the metastable Polycomb OFF state (Fig.

5A,B); (ii) that the analogue differences generated by *fca* mutants will persist, since the FCA

mediated repression can function independently of H3K27me3.

To validate this prediction, we measured *FLC* expression (measuring spliced and unspliced *FLC*

transcripts by qPCR) at the whole plant level, over the same developmental time course in five

different genotypes (Fig. 5C shows unspliced *FLC;* Fig. S5A shows spliced *FLC)*: in the parental L*er*,

clf-2 (Goodrich et al., 1997), *fca-3*, the *fca-3 clf-2* double mutant, and *fca-1*. We compared these data

to predicted *FLC* sense transcriptional activity from simulations of the full model for L*er, fca-3* and

fca-1, as well as for L*er* and *fca-3* with the PRC2 read-write feedback parameters changed to capture

a *clf* mutant. We did not attempt to fit to absolute levels of *FLC* transcripts here, since the measured

levels also depend on RNA stability, which we do not measure experimentally or include in our

model. Therefore, we compare changes relative to the measured levels at the first (10 day) timepoint.

We saw that, as predicted by the model, the slow reduction in expression is compromised in the *clf-2*

and *fca-3 clf-2* mutants (Fig. 5C, Fig. S5A) – neither of these mutants exhibits a statistically

significant *reduction* in *FLC* expression over the timecourse. We note that *clf-2* exhibits an increase

at the 14 day timepoint, (statistically significant only for unspliced *FLC*) that the model cannot

capture.

Also consistent with predictions for the *clf* mutants, we observe that the analogue differences

associated with FCA are maintained, i.e., the expression level in the *clf-2* mutant is clearly lower than

in *fca-3* at all timepoints (Fig. S5B), indicating that the FCA mediated transcription coupled

repression is the main factor underlying analogue differences. We also compared the model predicted

changes in *FLC* transcriptional output (i.e., frequency of distally terminated sense transcription

events) for *Ler, fca-3,* and *fca-1* to the above data (Fig. 5C, Fig. S5A). Consistent with previous data

(Antoniou-Kourounioti et al., 2023), and the observed chromatin changes (Fig. 4), all three genotypes

Ler, fca-3, and *fca-1* exhibit statistically significant changes in *FLC* expression over the time course

(both spliced and unspliced *FLC*). The relative changes predicted by the model are overall

quantitatively consistent with the measured changes in both unspliced (Fig. 5C) and spliced *FLC* (Fig.

S5A) (note that the model predictions are identical for spliced and unspliced – the model does not

distinguish between the two). We note that in some cases, between 10 days and 14 days, the model

predicted changes are larger than the measured changes, notably, for unspliced *FLC* in L*er* (Fig.5C)

and spliced *FLC* in *fca-3* and *fca-1* (Fig. S5A). However, since the model was only parametrised to fit

the ChIP time course data (Fig. 4), the overall satisfactory quantitative fit observed for *FLC*

expression with the same parameter values provides further support for the validity of the model.

In addition to *FLC* expression at the whole plant level, we also conducted time course imaging of

Venus tagged FLC in the roots in the *clf* mutants (Fig. 5D), following the same approach used to

analyse the *fca* alleles in (Antoniou-Kourounioti et al., 2023). Again, as predicted, *clf-2* showed clear

differences to the L*er* genotype, with ON cells being visible at all three timepoints. This is consistent

with *FLC* switching both ON to OFF and OFF to ON as described above, and the observed lack of

reduction in *FLC* expression at the whole plant level (Fig. 5C). We note that both *clf-2* and *fca-3 clf-2*

exhibited large variability between roots, with some roots showing essentially no *FLC* ON cells (Fig.

S5C). This is possibly due to the widespread misregulation of growth and developmental genes

resulting from the compromised Polycomb silencing in this mutant. Nevertheless, consistent with the

model predictions, we see that the slow Polycomb digital switch to OFF is disrupted in *fca-3 clf-2*:

when comparing roots exhibiting ON cells in the double mutant to *fca-3* roots, the double mutant

overall shows more ON cells at later timepoints.

The model also predicts a similar, high ratio of proximal to distal termination in the *clf* mutant and the

L*er* genotype (Fig. 4E), even though overall transcription is lower in L*er*. This is because the

FCA/FLD mediated transcription coupled mechanism is not disrupted in the *clf* mutant model, so that

the promotion of proximal termination by this mechanism, and the resulting feedback, remain intact.

We measured the ratio of proximal to distal termination of the *FLC* sense transcript in the *clf* mutant,

using our strand-specific Quant-seq data (Fig. S4C (i)). Consistent with the model prediction, it

exhibits a similar ratio to the L*er* genotype (Fig. 4F).

As further validation of the model, we then compared the predicted H3K27me3 levels at the locus in a

clf mutant to experimentally measured levels. The lack of stability of the nucleated state in the model

means that it predicts an H3K27me3 peak, but at significantly lower levels relative to a spread state.

The incorporation of the *FLC* gene loop in the model, as outlined above, causes the model to generate

a second peak in the *FLC* 3' region, in the region that interacts strongly with the nucleation region.

Consistent with these predictions, we find that the ChIP profile of H3K27me3 in *clf-2* exhibits two

peaks, at significantly lower levels relative to the spread state in L*er* (Fig. 5E). Our results also

aligned well with previously published genome-wide H3K27me3 data for a *clf* mutant in a Col-0

background (Shu et al., 2019). Overall, predictions from the model for *clf* mutants match well with

our experimental data.

The repressor FLD is co-transcriptionally targeted to *FLC* but requires FCA mediated

proximal termination for its repressive function

We next investigated the mechanism of FLD functionality in the co-transcriptional repression

mechanism. In apparent contradiction to its role as a repressor, in previously reported data, FLD

occupancy was found to be significantly higher in a high transcriptional state of *FLC*, relative to a

Polycomb silenced low transcriptional state. Specifically, high FLD levels were assayed by ChIP-seq of a transgenic *fld-4; 3xFLAG-FLD* in the Col*FRI* background, where the *FLC* activator FRIGIDA sets up a high transcriptional state (Inagaki et al., 2021). In contrast, in a Col-0 background, where FRI is not functional, *FLC* is Polycomb silenced and FLD shows low enrichment.

The model allows us to explain this counter-intuitive behaviour of FLD. Since FLD function is coupled to transcription and termination in the model, FLD targeting must be co-transcriptional, consistent with the molecular and genetic analysis in the accompanying paper (Mateo-Bonmatí, 2023). FLD levels at the locus must therefore be low in a Polycomb silenced state, where transcription events are infrequent. Such co-transcriptional FLD targeting is consistent with the genome-wide correlation between FLD enrichment and transcription as measured by Pol II ChIP and transcript amounts in (Inagaki et al., 2021). Furthermore, by capturing the digital switch off in L*er* and *fca-3*, the model predicts that FLD enrichment would reduce over time in these genotypes. In essence, FLD is only needed transiently to generate a lowly-transcribed state capable of nucleating Polycomb silencing, as transcription antagonises Polycomb. Following Polycomb nucleation and spreading of H3K27me3, FLD recruitment to the locus is reduced due to the lack of transcription.

The experimental validation of the model assumption of FLD targeting coupled to transcription termination, is presented in the accompanying paper (Mateo-Bonmatí, 2023), where we elucidate the links between the FLD complex (Fang et al., 2019) and CPF components. By ChIP-qPCR analysis of FLD enrichment at *FLC,* comparing high transcription states induced by two very different genetic backgrounds (*fca-9* and *FRI*) with a Polycomb silenced state (Col-0), we show that high FLD enrichment at *FLC* accompanies high transcription, irrespective of FCA function. Our model also supports the scenario proposed by Inagaki et al., (2021) in which FLD downregulates histone H3K4me1 around termination sites in regions with convergent overlapping transcription genome-wide in Arabidopsis.

We also tested whether FLD enrichment would change over time, as expected from the model predicted switch to the Polycomb silenced state. We measured 3xFLAG-FLD enrichment by ChIP-qPCR over a time course in a Col-0 background (as high-expressing *fca-9* and FRI are not expected to exhibit a digital switch OFF). Consistent with model predictions, we observed a reduction over time in FLD enrichment across *FLC* (Fig. 6B), mirroring the reduction in *FLC* transcription levels over time also observed in Col-0 by qPCR (Fig. 6A). Levels at the initial 7-day timepoint were high and declined to very low levels (almost background) by 21 days. Consistently, both 7-day and 14-day average FLD levels across the locus showed statistically significant differences to the control sample, while the 21-day levels did not (Fig. S6).

Together, these observations are consistent with co-transcriptional FLD targeting to the *FLC* locus,
with its repressive activity dependent on FCA-mediated proximal polyadenylation of sense and
antisense transcription. In genotypes such as *fca-9*, where FCA is non-functional, or in the presence of
the transcriptional activator FRIGIDA, which antagonises proximal PA by FCA (Schon et al., 2021),
FLD continues to be co-transcriptionally targeted, but fails to carry out its repressive function. This
highlights an important aspect relating genome-wide localisation of a regulator to its function: factors
that are found to localise at active transcription sites may not just be transcriptional activators. As is
the case for FLD, such factors could be essential components for shutting down transcription in
specific developmental contexts.

FCA mediated transcription-coupled repression potentially controls many Polycomb targets
We next analysed our genome-wide Quant-seq data to examine whether the FCA-mediated
transcription coupled mechanism could be applicable to a wider range of targets beyond *FLC,* and
whether this mechanism is also linked to Polycomb silencing at other targets. We first analysed
differential upregulation in *fca-1* (null) relative to L*er*, which showed that 130 targets were
significantly upregulated in *fca-1*. Of these 130 targets, 107 were upregulated to an intermediate level
in *fca-3* (without however showing statistically significant upregulation in *fca-3*, probably because of
the intermediate nature of this genotype*)*, hence behaving similar to *FLC* (Fig. 6C). We then
examined whether these 107 targets were also Polycomb targets. We used published ChIP-seq data
for the Arabidopsis PRC2 methyltransferases CLF and SWN (Shu et al., 2019) in the Col-0 genotype,
and generated a metagene plot for CLF and SWN enrichment across the 107 targets, comparing it to
the whole genome metaplot for CLF and SWN enrichment. This showed a clear higher enrichment of
these Polycomb components at these 107 putative FCA targets, particularly close to the Transcription
Start Site (TSS), compared to the genome-wide average (Fig. 6D).

In order to examine whether FCA and FLD function overlap in general, we used genome-wide Quant-
seq data in the null mutant genotypes *fca-9* and *fld-4* and the corresponding parental genotype Col-0,
to carry out a similar analysis. We found that 91 genes were significantly upregulated in *fca-9*
compared to the parental genotype, Col-0. We first examined whether these targets overlap with those
observed in *fca*-1. Our data shows extensive differential gene expression between the Col-0 and L*er*
parental genotypes (not shown), and these differences are reflected in the respective mutants in these
backgrounds. It is therefore not surprising that we find a relatively small overlap of 10 targets
between the *fca-1* upregulated set and the *fca-9* upregulated set, even though these are both null
mutants of *FCA*.

Analysis of *fld-4* relative to Col-0 showed a larger number of upregulated targets (193), consistent
with the proposed wider role for FLD in coordinating convergent transcription genome-wide (Inagaki

et al., 2021). We found that 48 targets were significantly upregulated in both *fca-9* and *fld-4*,
indicating that these factors may be working together at a number of genes. As above, we compared
metagene plots of CLF and SWN enrichment across these 48 targets, as well as for the *fca-9* and *fld-4*
singly upregulated subsets, in a whole genome metagene plot for CLF and SWN enrichment. For the
fca-9 upregulated genes, we saw slightly higher enrichment of SWN but not of CLF around the TSS
compared to the genome-wide average (Fig. 6E). Interestingly, for the *fld-4* upregulated subset, we
observe lower enrichment of both SWN and CLF compared to the genome-wide average (Fig. 6E).
This is consistent with the co-transcriptional targeting of FLD, but where the majority of the targets
do not couple to Polycomb. However, the metagene plot for the 48 overlapping targets compared to
the genome wide-average shows a further increase in SWN enrichment (and a smaller increase for
CLF) around the TSS than seen for the *fca-9* upregulated targets alone (Fig. 6E). This suggests that
FLD function may be linked to silencing by PRC2 in specific contexts, where FLD action is
facilitated by FCA mediated proximal polyadenylation.

Together, these analyses indicate that FCA/FLD mediated transcription-coupled repression may be
working at a broad range of targets, and that at these targets, as at *FLC,* this repression works in
concert with the Polycomb system. We also note that multiple genes of general interest are present in
the *fca-1* and *fca-9/fld-4* upregulated lists, including: a Trithorax homologue (ATX1) implicated in
development and multiple stress responses, genes such as GRP5 (Mangeon et al., 2010) and GASA9
(Yang et al., 2008) showing highly cell type specific expression, and several stress inducible genes
(see Supplementary Table S2 for gene names of selected targets), all features that are consistent with
Polycomb regulation.

Discussion

How co-transcriptional processing mechanistically influences transcriptional output, and how this is
linked to epigenetically stable Polycomb silencing is still not understood in any system. We have
exploited the Arabidopsis *FLC* system to investigate these questions. We had shown that a
transcription-coupled repression mechanism; specifically involving proximal transcription
polyadenylation/termination mediated by the RNA-binding protein FCA interacting with 3'
processing machinery, can generate graded (analogue) transcriptional output. We had also shown that
this mechanism was coupled to histone demethylase (FLD) activity. In the accompanying paper
(Mateo-Bonmatí, 2023), we showed that APRF1, a structural component of a CPF-like phosphatase
complex, directly links transcription termination with the FLD histone demethylase activity to alter
the local chromatin environment. But how FLD activity can affect proximal termination, how this
mechanism can generate analogue transcriptional output, and how this transcription-coupled
repression links to a digital ON to OFF switch to PRC2 silencing remained obscure.

We built on our previous using the computational model described here, and used it to reveal how the

level of productive transcriptional activity set by the transcription-coupled mechanism (FCA/FLD)

determines the level of transcriptional antagonism to Polycomb silencing. The different levels of

transcriptional activity set by the transcription coupled repression can dictate the timescale for

establishment of Polycomb silencing (Fig. 7). We validated the model by a variety of means,

including gene expression, histone modification and FLC-Venus imaging time courses in the wild-

type and in cases disrupted via a Polycomb mutation. Our model also resolves the apparent

contradiction between the observed FLD targeting to a transcriptionally active *FLC* locus, and its role

as a repressor: FLD's co-transcriptional recruitment, needed to lower transcriptional activity to pave

the way for Polycomb silencing, is eliminated once full Polycomb silencing is achieved. Hence,

factors localising at activated genes can in some instances be paradoxically essential for shutting

down transcription.

A key methodology to dissect the underlying mechanism was the construction of a computational

model. The model was used as a tool to predict the levels of dynamically changing transcriptional and

histone modification levels and to gain intuition about the underlying mechanisms. We chose to keep

the model simple and interpretable but the model was still able to reproduce all the statistically

significant trends in the data.

The transcriptional antagonism to Polycomb silencing, a fundamental aspect of the silencing

mechanism, is also important in the mammalian and Drosophila contexts (Klose et al., 2013), (Holoch

et al., 2021). In the plant context, this antagonism sets the timescale for the slow switch into the

digital Polycomb silenced state, in an interlinked three-way association between chromatin,

transcription and co-transcriptional processing (Graphical Abstract). Importantly, our model predicts

slow/inefficient establishment of Polycomb silencing in the *fca-1* mutant. As a result, if silencing can

be established in *fca-1* (independently of FCA), for example mediated by cold-specific PRC2

accessories in the case of cold-induced silencing, we predict that both silenced and active states can

each be stably maintained. Hence, our conceptual framework also explains why cold-induced

Polycomb silencing can be maintained in *fca* (or *fld*) mutants. The model predicts that these proteins

are needed only for Polycomb silencing establishment during development, but not for its

maintenance. In future work, it would be very interesting to specifically test this prediction by

removing FCA and/or FLD at a later developmental stage after silencing is established.

The putative H3K4 demethylase FLD is known to be a key component of the FCA mediated

transcription-coupled repression mechanism, indicating an important role for removal of H3K4me

associated with and downstream of proximal termination. A key mechanistic feature of the model is

the existence of a link also in the opposite direction: H3K4me1 levels influencing the processivity of

RNA Pol II – with low H3K4me1 levels reducing processivity. Thus, the model contains a positive

feedback loop (mutual inhibition) between proximal termination and H3K4me1. While chromatin

modifications have been mechanistically linked to co-transcriptional splicing (Leung et al., 2019),

links to Pol II processivity are only beginning to be explored, as seen from recent findings on Pol II

regulation by the Integrator complex in mammalian systems (Stein et al., 2022; Wang et al., 2023).

These links could also involve an effect on Pol II speed which can influence termination (Cortazar et

al., 2019).

Our model also explains how the *FLC* transcriptional activator FRIGIDA can set up an active

transcriptional state, preventing Polycomb silencing by functioning as an anti-terminator. Anti-

termination factors that prevent the use of early termination sites have been implicated in tuning

transcriptional output from eukaryotic polymerases (Gregersen et al., 2019). In the case of *FLC,* the

transcriptional activator FRIGIDA has recently been shown to function at least partly by antagonising

FCA mediated proximal cleavage and polyadenylation of *FLC* sense transcription at specific stages of

embryo development (Schon et al., 2021). The presence of active FRIGIDA promotes distal

polyadenylation and prevents *FLC* silencing, further supporting the view that FCA mediated proximal

termination is essential for repression. Interestingly, FRIGIDA expression outside of a specific

developmental stage fails to activate *FLC* expression (Schon et al., 2021): this reinforces the view that

once Polycomb epigenetic silencing has been established, reactivating transcription is difficult, with

simple upregulation of an activator being insufficient to overcome robust digital silencing. The model

predicts that the locus is bistable in the absence of proximal-termination mediated repression, so that

both active and Polycomb silenced states can be stably maintained (Fig. S4B(ii)). This is consistent

with the FCA mediated mechanism not having a role in maintaining the Polycomb silenced state, as

evidenced by the stable maintenance of cold-induced Polycomb silencing at *FLC* in both *fca* mutants

and genotypes with active FRIGIDA.

Overall, we have revealed the mechanism of co-transcriptional silencing at *FLC,* an essential

precursor for slow switching into the Polycomb silenced state. These features are included in an

animation to help explain these molecular feedbacks, and how this mechanism can lead to expression

of *FLC* in one of two stable expression states (supplementary file: Movie_1.mp4). It is particularly

intriguing that many elements of this pathway, such as FLD and FCA, are conserved

transcriptional/splicing regulators, that likely have functions at many more targets. Their core role in

an integrated sense/anti-sense transcriptional circuitry likely singles out *FLC* as being acutely

sensitive to their action. Indeed, our analysis of genome-wide Quant-seq data and overlap with ChIP-

seq data for Polycomb components indicates that this transcription-coupled mechanism targets many

genes and potentially works in concert with Polycomb at these targets. We therefore anticipate these

mechanisms will have even broader relevance, with additional examples of such mechanisms
emerging over time (Chen et al., 2023). Further investigating the links between co-transcriptional
mechanisms and Polycomb silencing will therefore be an important focus for future research.

Methods

624
Plant materials
All the plants were homozygous for the indicated genotype. Seeds were surface sterilized in 40 % v/v
commercial bleach for 10 min and rinsed 4 times with sterile distilled water. Seeds were then sown on
standard half-strength Murashige and Skoog (MS) medium (0.22% MS, 8% plant agar) media plates
and kept at 4°C in darkness for 3 days before being transferred to long day photoperiod conditions (16
h of light, 8 h dark).

631
632 **Plant growth (Imaging)**
633 Seeds were surface-sterilized with 5% (v/v) sodium hypochlorite for 5 min, rinsed with sterile water
634 4x for 1 min and stratified at 4°C for 2 days in water and in the dark. Seeds were plated on MES
635 buffered Murashige and Skoog (GM-MS) media, pH 5.8, containing respective antibiotics and grown
636 on vertically oriented petri plates for 7 days in long-day conditions (16 h light/8 h dark, 22/20°C).

637
Imaging
Time course imaging of FLC-Venus protein levels in the epidermis of L*er*, *fca* allelic mutants, the *clf-*
2 mutant as well as the *fca-3 clf-2* mutant, was performed using a Leica confocal Stellaris 8
microscope with a 20x multi-immersion objective (0.75 NA). Root tips (2 cm) were cut off and
immersed in 1.5 µg/mL Propidium Iodide (PI, Sigma–Aldrich, P4864) in 1x PBS before being imaged
immediately. The OPSL 514 laser was used at 5% power with 514 nm excitation for FLC-Venus and
PI. In photon counting mode, Venus was detected between 518-550 nm with the HyD SMD2 detector,
PI was detected between ~600-650 nm. Images were acquired with a laser speed of 400 Hz, line
accumulation of 6 (pixel dwell time of 2.43 µs), a Z-step size of 0.95 µm and a pinhole size of 1 AU.
The same settings were used at all imaging time points to allow direct comparison between genotypes
and time points. The representative images were projected using a single middle slice from the PI
channel to show the cell outline, and 10 slices of FLC-Venus channel were maximum intensity
projected. In standard image presentation, the dynamic range of the FLC-Venus channel was pushed
from 0-255 to 2-11, in enhanced image presentation, the dynamic ranged was pushed from 2-7. In
Supplementary Figure S5C, the dynamic range of FLC-Venus channel was enhanced from 0-255 to 2-
20 for all images.

654
RNA extraction and RT-qPCR

Total RNA was extracted as previously described (Box et al. 2011). Genomic DNA was digested with

TURBO DNA-free (Ambion Turbo Dnase kit AM1907) according to manufacturer´s guidelines,

before reverse transcription (RT) was performed. The RT reactions were performed with SuperScript

IV reverse transcriptase (ThermoFisher, 18090010) following the manufacturer´s instructions using

gene-specific primers. Primers are listed in Supplemental Table S1. For RNA sequencing

experiments, RNA was further purified with the Qiagen Rneasy miniprep kit (74106).

RNA sequencing

Library preparation and sequencing was performed by Lexogen GmbH. RNA integrity was assessed

on a Fragment Analyzer System using the DNF-471 RNA Kit (15 nt) (Agilent). Sequencing-ready

libraries were generated from 100ng of input RNA using a QuantSeq 3' mRNA-Seq Library Prep Kit

REV for Illumina (015UG009V0271) following standard procedures. Indexed libraries were assessed

for quality on a Fragment Analyzer device (Agilent), using the HS-DNA assay and quantified using a

Qubit dsDNA HS assay (Thermo Fisher). A sequencing-ready pool of indexed libraries were

sequenced on an Illumina NextSeq 2000 with a 100-cycle cartridge using the Custom Sequencing

Primer (CSP) at Lexogen GmbH.

Bioinformatics analysis

Data analysis of RNAseq data was performed by Lexogen GmbH using their standard

pipeline. Read Quality was assessed by FastQC and trimmed for quality and adapter content

with cutadapt version 1.18 (Martin, 2011). Clean reads were mapped to the Arabidopsis

reference genome (TAIR10) with STAR version 2.6.1a (Dobin et al., 2013). Raw reads have

been deposited on Short Read Archive (SRA) under the reference PRJNA980626, and

PRJNA1076161. To calculate proximal to distal count ratios, read ends were counted for

each identified polyA cluster ((Schon et al., 2021), FLC_polyA_cluster_coords.txt). Proximal

polyadenylation clusters are defined as termination sites within the first Exon and Intron of

FLC while distal clusters are in close proximity to the canonical polyadenylation site. For

each biological sample, ratios were calculated from the summed proximal and distal cluster

counts (FLC_termination_clusters.txt).

ChIP timecourse for histone modifications

2.5 g of 7-, 14-, or 21-day-old seedlings were crosslinked in 1% formaldehyde and grinded to fine

powder. Nuclei were extracted using Honda buffer as described previously in Sun et al. 2013. In all

histone ChIP reactions, sonication, immunoprecipitation, DNA recovery and purification were

performed as previously described in Wu et al. 2016. The antibodies used were anti-H3 (ab176842),

anti-H3K27me3 (Merck 07-449 and ab192985), anti-H3K36me3 (ab9050), anti-H3K4me1 (ab8895).

All ChIP data were quantified by qPCR with primers listed in Table S1. Values were normalized to
H3 and internal control to either STM or ACT7.
694

H3K27me3 ChIP in *clf* mutant
2.0 g of 14-day-old seedlings were crosslinked in 1% formaldehyde and grinded to fine powder.
Nuclei were extracted using Honda buffer as described previously in Sun et al. 2013. In all histone
ChIP reactions, sonication, immunoprecipitation, DNA recovery and purification were performed as
previously described in Wu et al. 2016. The antibodies used were anti-H3 (ab176842), and anti-
H3K27me3 (ab192985). All ChIP data were quantified by qPCR with primers listed in Table S1.
Values were normalized to H3 and internal control *STM*.
702

FLD ChIP
2.5 g seedlings were crosslinked in 0.685 mg/mL of EGS [ethylene glycol bis(succinimidyl
succinate)] and 1% formaldehyde and grinded to fine powder. Nuclei were extracted using Honda
buffer as described for histone ChIP. Nuclei pellets were resuspended in 600 µl of RIPA buffer (1%
NP-40, 0.5% Sodium Deoxycholate, 0.1 % SDS in PBS) and sonicated 5 times for 5 min (30s ON/
30s OFF) at high mode in a Bioruptor Pico (Diagenode). IP were performed overnight at 4°C using
1.5 mg of pre-coated epoxy Epoxy Dynabeads (M-270) with 1.5 µl of anti-FLAG (Sigma, F1804) per
reaction. Beads were then washed for 5 min twice with Low Salt Buffer (150 mM NaCl, 0.1% SDS,
1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl), twice with Hight Salt Buffer (500 mM NaCl,
0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl), and twice with TE buffer (1 mM
EDTA, 10 mM Tris-HCl). DNA recovery and purifications were performed as for Histone ChIP.
Results in Fig. 6B were normalized to IGN5.
717

718 **Computational model**
719 For a detailed description of the computational model, its assumptions, implementation, and analysis
720 of simulation output, see Supplementary Information. Additional experimental considerations for
721 modelling are also presented in the Supplementary Information.
722

Acknowledgements
The authors would like to thank members of the Dean and Howard labs for thoughtful inputs and
Shuqin Chen for excellent technical assistance. This work was funded by European Research Council
Advanced Grant (EPISWITCH, 833254), Wellcome Trust (210654/Z/18/Z), and a Royal Society
Professorship (RP\R1\180002) to CD, and BBSRC Institute Strategic Programmes (BB/J004588/1
and BB/P013511/1) and EPSRC/BBSRC Physics of Life grant (EP/T00214X/1) to MH and CD.

Eduardo Mateo-Bonmatí would like to thank grant RYC2021-030895-I, funded by MCIN/AEI and by

European Union NextGenerationEU/PRTR. Svenja Reeck was funded by the HORIZON EUROPE

Marie Sklodowska-Curie Actions PEP-NET (Predictive Epigenetics: Fusing Theory and Experiment)

grant 813282.

Figure Captions

Figure 1: **(A)** Schematic showing two possible modes of regulating population level expression –

Analogue and Digital. **(B)** Schematic of combined digital and analogue regulation observed

experimentally in *fca* mutants. **(C)** Schematic showing the transcription coupled repression

mechanism at *FLC*. Activating H3K4me1 can be added during productive transcription (green dashed

arrow). The repression mechanism consists of proximal polyadenylation and termination of sense and

antisense transcription mediated by the RNA binding protein FCA, which then enables removal of

transcriptionally activating H3K4me1 across the locus by the putative histone demethylase FLD. The

removal of H3K4me1 can inhibit co-transcriptional processing by preventing the addition of

H3K36me3, and consequently promotes further proximal termination. Thus, a feedback loop

(highlighted by the black dashed arrows) is formed, where the occurrence of proximal termination

promotes further proximal termination. Also shown is a schematic of the Polycomb silenced state at

FLC, but how this is linked to transcription-coupled repression has been unclear. Supplementary file:

Movie_1 contains an animation summarising the processes and interactions shown here.

Figure 2: **(A)** qPCR data for *FLC* expression, showing repression of a transgenic wild-type *FLC*

construct (*FLC-15* indicated by WT *FLC*) in an FCA overexpression genotype (*35S::FCA* indicated

by *FCA* OE), in 14 day old seedlings, and the attenuation of this repression for an *FLC* transgenic

construct where antisense transcription is disrupted (*TEX1*). All genotypes shown are in an *fca-9*

(*FCA* null mutant) background. Error bars represent mean +/- s.e.m. p-values are shown for a

statistical comparison of the mean values using the Student's t-test. **(B)** Quantseq data for 7-day old

seedlings (7 days after sowing), showing quantification of polyadenylated transcripts mapped to the

FLC locus in the parental Landsberg (L*er*) genotype and the *fca* mutants. Individual replicates are

shown in Fig. S4D.

Figure 3: **(A)** Schematic of the combined mathematical model incorporating the transcription-

coupled repression mechanism and PRC2 mediated silencing. The schematic highlights the key

processes and interactions captured by the mathematical model. For the transcription coupled

repression mechanism (as depicted in Fig. 1C) this includes sense and antisense transcription, co-

transcriptional addition of H3K4me1 (green dashed arrows), FCA mediated proximal termination of

transcription, FLD mediated removal of H3K4me1, and the promotion of proximal termination by

low H3K4me1 coverage. Feedback is again highlighted by black dashed lines. For PRC2 silencing the
model includes H3K27 methylation, PRC2 mediated read-write maintenance of these modifications,
and mutual antagonism between H3K27me3 and transcription. **(B)** FCA effectiveness parameter
values for the three genotypes. **(C)** Top panel: Model predicted frequency of distal transcription
events (steady state average over 1000 simulated loci) in an ON state, for the model of the
transcription-coupled repression mechanism shown in (A) but without the Polycomb silencing
module. Each simulated locus was started in an active state with full H3K4me1 coverage and allowed
to equilibrate over a duration of 5 cell cycles. Averaging was performed over the duration of the
following 10 cell cycles. Bottom panel: Model predicted fractional coverage of H3K4me1 in an ON
state from the above simulations. **(D)** Predicted transcriptional changes (quantified by frequency of
transcription events) over time obtained by simulating the combined model shown in (A). Frequencies
of the different transcription events are averaged over 1000 simulated copies of the *FLC* locus, all
started in an active state with full H3K4me1 coverage. **(E)** Predicted changes in the fraction of ON
copies in the *fca* mutants, computed from simulations of 1000 copies over 30 cell cycles, started in a
full H3K4me1 covered ON state. Fractions are shown at the starting timepoint (0), and at the end of 7,
14, and 21 days from the start. See Supplementary Information for the definition of ON and OFF
states used in this analysis.

Figure 4: (A,B) ChIP-qPCR time course measurements of H3K27me3 (A), and H3K4me1 (B) across
the *FLC* locus in the parental L*er* genotype and *fca* mutants. H3K27me3 levels are shown normalised
to H3 and to control locus *STM.* H3K4me1 levels are shown normalised to H3 and to the control locus
ACT7. Each dot represents an amplicon. Error bars in (A) represent mean +/- s.e.m. (n = 5 biological
replicates only for L*er* at 7 days and n=3 biological replicates in all other cases). Error bars in (B)
represent mean +/- s.e.m. (n = 5 biological replicates for L*er*, *fca-3*, and *fca-1* at 7 days and n=3
biological replicates in all other cases). **(C,D)** Experimental validation of model predictions. Vertical
red bars indicate model predicted level. All model predicted histone modification levels are averaged
over the locus and averaged over 1000 simulated copies of the locus started in an active state with full
H3K4me1 coverage. Horizontal black bars indicate median of the biological replicates. **(C)**
Comparison of model predicted H3K27me3 levels and changes over time in the different genotypes,
with the measured levels shown in (A), averaged across 15 primers covering the *FLC* locus. **(D)**
Similar comparison of model predicted H3K4me1 levels with the experimental data shown in (B).
(E,F) Model predicted (E) and measured (Quantseq) (F) ratios of proximal to distal termination for
FLC sense transcription, in the parental L*er* genotype, *fca* mutants and a *clf* mutant. Measured ratios
were computed using the Quantseq data for 7 day old seedlings shown in Fig. 2B and Fig. S4C(i).
Error bars in (F) represent mean +/- s.e.m. (n = 3 biological replicates). The genomic coordinates used
to define the proximal and distal termination clusters are shown in the supplementary file
FLC_polyA_cluster_coords.txt. The read counts within these clsuters for the different genotypes are

provided in the supplementary file FLC_termination_clusters.txt. Model predicted ratios are

normalised to the predicted ratio for L*er*. Model predicted ratios were computed at the 7 day

timepoint (averaged over the last 0.25 days leading up to this timepoint) for all four genotypes

(average over 1000 simulated copies started in an active state with full H3K4me1 coverage). The

model qualitatively captures the relative changes in the proximal to distal ratio between mutants. Note

that absolute predictions of the ratios would require estimates of lifetimes for the proximal and distal

polyadenylated forms in each mutant.

Figure 4 – Supplementary 4A: Statistical comparisons of average ChIP-qPCR measured H3K27me3

(i,iii) and H3K4me1 (ii,iv) levels across the locus, between genotypes (i, ii) and between timepoints

(iii,iv). In all cases the data points show the measured modification level averaged across 15 primers

covering the *FLC* locus (full modification profiles shown in Fig. 4A,B). Horizontal bars indicate

mean of replicates. Statistical comparisons for the experimental data were carried out using the

Student's t-test and corrected for multiple comparisons using the Holm-Sidak approach (*, **, ***

indicate significance thresholds of 0.05, 0.01, and 0.001 respectively).

Figure 4 – Supplementary 4B: (i) Histograms showing probability distribution of switch OFF times

for an individual locus, for L*er*, *fca-3,* and *fca-1*. Histograms are generated by simulating 1000 copies,

started in a full H3K4me1 covered ON state, and simulated over 60 cell cycles. **(ii)** Model predicts

that the locus is in a bistable regime in the absence of the proximal termination mechanism (proximal

termination probability set to residual value – see model description in Supplementary Information

and parameter definitions/values in Supplementary Table 3), with both ON state and OFF state stably

maintained across several cell cycles. Plots show fraction of ON copies (left column) and H3K27me3

fractional coverage (right column) for 100 simulated loci started in an ON state (full H3K4me1

coverage) (top row) or started in an OFF state (full H3K27me3 coverage) (bottom row). Fractions are

shown at the starting timepoint (0), and at the end of 5, 10, and 20 cell cycles from the start. During

steady state maintenance of the OFF state, the cell cycle-averaged H3K27me3 stabilises at an

intermediate level due to the periodic dilution by DNA replication and subsequent slow accumulation.

(iii) ChIP-qPCR time course measurements of H3K36me3 across the *FLC* locus in the parental L*er*

genotype and the *fca* mutants. Each dot represents an amplicon. Error bars represent mean +/- s.e.m.

(n = 3 biological replicates). H3K36me3 levels are shown normalised to H3 and to the control locus

ACT7.

Figure 4 – Supplementary 4C: (i) Quantseq data for 7-day old seedlings (7 days after sowing),
showing quantification of polyadenylated transcripts mapped to the *FLC* locus in the parental
Landsberg (L*er*) genotype, the *fca* mutants (L*er* and *fca* mutants data same as shown in Fig. 2B), and
the PRC2 methyltransferase mutant *clf-2*. **(ii)** ChIP-qPCR measured H3K27me3 levels (normalised to
H3) shown for the three *fca* genotypes at *FLC* as well as at the designated positive (+) and negative
control loci (-) for this modification. **(iii)** ChIP-qPCR measured H3K4me1 levels (normalised to H3)
shown for the three *fca* genotypes at *FLC* as well as at the designated positive (+) and negative control
loci (-) for this modification. **(iv)** ChIP-qPCR measured H3K36me3 levels (normalised to H3) shown
for the three *fca* genotypes at *FLC* as well as at the designated positive (+) and negative control loci (-
) for this modification.

Figure 4 – Supplementary 4D: Individual replicates for Quantseq data shown in Fig. S4C.

Figure 5: (A) Schematic showing the rationale for examining the *clf* mutant. Transition from a
Polycomb nucleated state to a spread state is prevented in a *clf* mutant. The lack of stability of a
nucleated state allows switching back to a digital ON state. Green circles: active modifications
(H3K4me1); red circles: repressive modifications (H3K27me3). **(B)** Consequently, the reduction over
time in the number of ON cells observed in the parental L*er* genotype is expected to be disrupted in
the *clf* mutant. **(C)** Comparison of model predicted changes in *FLC* transcription, with whole seedling
FLC unspliced transcript levels measured by qPCR in the *fca* genotypes (L*er*, *fca-3*, *fca-1*) and the *clf*
mutants *(clf-2* and *fca-3 clf-2* double mutant). Experimental data is normalised to measured levels for
UBQ10. Statistical comparisons for the experimental data were carried out using the Student's t-test
and corrected for multiple comparisons using the Holm-Sidak approach (*, **, *** indicate
significance thresholds of 0.05, 0.01, and 0.001 respectively). Model predictions correspond to the
predicted frequency of *FLC* sense distal transcription events at the corresponding timepoints,
averaged over 1000 simulated loci. Simulations for L*er*, *fca-3*, *fca-1* are identical to those used to
predict changes in histone modification levels (Fig. 4). Simulations for the *clf* mutants *(clf-2*, *fca-3 clf-*
2) use changed parameter values for reactions involving H3K27me3 addition by PRC2 – see
Supplementary Information and Supplementary Table 2. For comparison of relative changes, both
model predictions and data are normalised to the level at 10 days. **(D)** FLC-Venus time-course
imaging in the different genetic backgrounds. Representative confocal images of FLC-Venus signal at
three timepoints in the epidermis of the root meristem of the parental genotype Ler, the single mutants
clf-2, and *fca-3*, and the double mutant *fca-3 clf-2*. Grey shows the cell wall dye propidium iodide;
FLC-Venus intensity is indicated by colour maps for standard and enhanced images, respectively. The
same settings were used for imaging of all time points. The same image presentation is used for
standard and enhanced images. Brightness and contrast of FLC-Venus signal were increased to

generate the enhanced images (Note different scale on colour maps). Yellow boxes in *clf-2* highlight

ON cell files at DAS7/14. Scale bar, 50 μm. DAS stands for days after sowing. **(E)** The model can

capture the differences between L*er* and *clf-2,* in the spatial profile of H3K27me3 across the *FLC*

locus. Top panel: model predicted profiles of H3K27me3 coverage (see Supplementary Information

for detailed description of simulation). Model predicted profile is steady-state average over 100

simulated loci, started in an active chromatin state with full H3K4me1 and equilibrated over a time

duration of 10 cell cycles. Bottom panel: H3K27me3 profiles measured by ChIP-qPCR in 14-day old

seedlings. Error bars represent mean +/- s.e.m. (n = 3 biological replicates). Each dot represents an

amplicon.

Figure 5 – Supplementary 5A: Changes in Spliced *FLC* transcript level. Comparison of model

predicted changes in *FLC* transcription, with whole seedling *FLC* spliced transcript levels measured

by qPCR in the *fca* genotypes (L*er, fca-3, fca-1*) and the *clf* mutants (*clf-2* and *fca-3 clf-2* double

mutant). Experimental data is shown normalised to measured levels for *UBQ10.* Statistical

comparisons for the experimental data were carried out using the Student's t-test and corrected for

multiple comparisons using the Holm-Sidak approach (*, **, *** indicate significance thresholds of

0.05, 0.01, and 0.001 respectively). Model predictions correspond to the predicted frequency of *FLC*

sense distal transcription events at the corresponding timepoints, averaged over 1000 simulated loci.

Simulations for L*er, fca-3, fca-1* are identical to those used to predict changes in histone modification

levels (Fig. 4). Simulations for the *clf* mutants (*clf-2* and *fca-3 clf-2*) use changed parameter values for

reactions involving H3K27me3 addition by PRC2 – see Supplementary Information and

Supplementary Table 2. For comparison of reltive changes, both model predictions and data are

normalised to the level at 10 days.

Figure 5 – Supplementary 5B: (i) *FLC* unspliced transcript levels in whole seedlings measured by

qPCR. **(ii)** *FLC* spliced transcript levels in whole seedlings measured by qPCR. Data is shown

normalised to the measured levels for *UBQ10.*

Figure 5 - Supplementary 5C: FLC-Venus time course replicates with additional genotypes

Additional representative images of FLC-Venus signal in Ler, *clf-2*, *fca-3*, *fca-3 clf-2* as well as an

additional intermediate allele *fca-4* and the strong allele *fca-1* at three time points. FLC-Venus

intensity is indicated by colour map; grey shows the cell wall. Scale bar, 50 μm. DAS stands for

days after sowing.

Figure 6: (A) Changes over time in spliced *FLC* expression in Col-0 seedlings, measured by time-

course qPCR. Error bars represent mean +/- s.e.m. (n = 3 biological replicates**).** Data is shown

normalised to the measured levels for *UBQ10* and *PP2A.* **(B)** Changes over time in FLD targeting to

the *FLC* locus, measured by time-course ChIP-qPCR quantification of 3xFLAG-FLD in seedlings

(Col-0 background) (normalized to IGN5). Error bars represent mean +/- s.e.m. (n = 3 biological

replicates). **(C)** Heatmap generated from analysing genome wide Quant-seq data, showing changes in

transcriptional output at the 107 targets that are significantly upregulated in *fca-1* and also show

intermediate (but not necessarily statistically significant) upregulation in *fca-3*. **(D)** Metagene plots

showing enrichment of the PRC2 methyltransferases SWN and CLF at the 107 targets (Differentially

Expressed Genes, DEGs) shown in (C), compared to their enrichment genome-wide. **(E)** Metagene

plots comparing SWN and CLF enrichment at *fca-9* upregulated, *fld-4* upregulated, and overlapping

DEGs, with their genome-wide enrichment.

Figure S6: Statistical analysis of changes over time in 3xFLAG-FLD enrichment across the locus.

Data points indicate average measured level over 16 primers covering the *FLC* locus (full dataset

shown in Fig. 6B). Horizontal bar indicates mean over replicates. Statistical comparisons for the

experimental data were carried out using the Student's t-test and corrected for multiple comparisons

using the Holm-Sidak approach (*, indicates a significance threshold of 0.05).

Figure 7: Schematic showing how analogue and digital regulation can combine in the context of

Polycomb mediated silencing. The differences in thickness of the red arrows indicate the different

rates of switching to the Polycomb silenced OFF state.

Graphical Abstract: Schematic showing the general interplay between transcriptional activity,

chromatin state, and co-transcriptional processing that controls the transcriptional state, and the

specific aspects of this interplay that regulate transcription at the *FLC* locus.

References

Angel, A., Song, J., Dean, C., and Howard, M. (2011). A Polycomb-based switch underlying
quantitative epigenetic memory. Nature *476*, 105-108.

Antoniou-Kourounioti, R.L., Meschichi, A., Reeck, S., Berry, S., Menon, G., Zhao, Y.,
Fozard, J., Holmes, T., Zhao, L., Wang, H.*, et al.* (2023). Integrating analog and digital
modes of gene expression at Arabidopsis FLC. Elife *12*.

Baxter, C.L., Šviković, S., Sale, J.E., Dean, C., and Costa, S. (2021). The intersection of
DNA replication with antisense 3′ RNA processing in *Arabidopsis FLC* chromatin
silencing. Proceedings of the National Academy of Sciences *118*, e2107483118.

Berry, S., Dean, C., and Howard, M. (2017). Slow Chromatin Dynamics Allow Polycomb
Target Genes to Filter Fluctuations in Transcription Factor Activity. Cell Syst *4*, 445-
457 e448.

Berry, S., Hartley, M., Olsson, T.S., Dean, C., and Howard, M. (2015). Local chromatin
environment of a Polycomb target gene instructs its own epigenetic inheritance. eLife
4, e07205.

Chalamcharla, V.R., Folco, H.D., Dhakshnamoorthy, J., and Grewal, S.I. (2015). Conserved
factor Dhp1/Rat1/Xrn2 triggers premature transcription termination and nucleates

heterochromatin to promote gene silencing. Proceedings of the National Academy of
Sciences *112*, 15548-15555.
Chen, W., Zhu, T., Shi, Y., Chen, Y., Li, W.J., Chan, R.J., Chen, D., Zhang, W., Yuan, Y.A.,
Wang, X.*, et al.* (2023). An antisense intragenic lncRNA SEAIRa mediates
transcriptional and epigenetic repression of SERRATE in Arabidopsis. Proceedings of
the National Academy of Sciences *120*, e2216062120.
Cortazar, M.A., Sheridan, R.M., Erickson, B., Fong, N., Glover-Cutter, K., Brannan, K., and
Bentley, D.L. (2019). Control of RNA Pol II speed by PNUTS-PP1 and Spt5
dephosphorylation facilitates termination by a "sitting duck torpedo" mechanism.
Molecular cell *76*, 896-908. e894.
Csorba, T., Questa, J.I., Sun, Q., and Dean, C. (2014). Antisense *COOLAIR* mediates the
coordinated switching of chromatin states at *FLC* during vernalization. Proc Natl Acad
Sci U S A *111*, 16160-16165.
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15-21.
Dodd, I.B., Micheelsen, M.A., Sneppen, K., and Thon, G. (2007). Theoretical analysis of
epigenetic cell memory by nucleosome modification. Cell *129*, 813-822.
Fang, X., Wang, L., Ishikawa, R., Li, Y., Fiedler, M., Liu, F., Calder, G., Rowan, B., Weigel,
D., Li, P.*, et al.* (2019). Arabidopsis FLL2 promotes liquid-liquid phase separation of
polyadenylation complexes. Nature *569*, 265-269.
Fang, X., Wu, Z., Raitskin, O., Webb, K., Voigt, P., Lu, T., Howard, M., and Dean, C.
(2020). The 3' processing of antisense RNAs physically links to chromatin-based
transcriptional control. Proc Natl Acad Sci U S A *117*, 15316-15321.
Goodrich, J., Puangsomlee, P., Martin, M., Long, D., Meyerowitz, E.M., and Coupland, G.
(1997). A Polycomb-group gene regulates homeotic gene expression in Arabidopsis.
Nature *386*, 44-51.
Gregersen, L.H., Mitter, R., Ugalde, A.P., Nojima, T., Proudfoot, N.J., Agami, R., Stewart,
A., and Svejstrup, J.Q. (2019). SCAF4 and SCAF8, mRNA Anti-Terminator Proteins.
Cell *177*, 1797-1813.e1718.
Holoch, D., Wassef, M., Lövkvist, C., Zielinski, D., Aflaki, S., Lombard, B., Héry, T., Loew,
D., Howard, M., and Margueron, R. (2021). A cis-acting mechanism mediates
transcriptional memory at Polycomb target genes in mammals. Nature Genetics *53*,
1686-1697.
Inagaki, S., Takahashi, M., Takashima, K., Oya, S., and Kakutani, T. (2021). Chromatin-
based mechanisms to coordinate convergent overlapping transcription. Nat Plants *7*,
295-302.
Kaneko, S., Son, J., Bonasio, R., Shen, S.S., and Reinberg, D. (2014). Nascent RNA
interaction keeps PRC2 activity poised and in check. Genes Dev *28*, 1983-1988.
Klose, R.J., Cooper, S., Farcas, A.M., Blackledge, N.P., and Brockdorff, N. (2013).
Chromatin Sampling—An Emerging Perspective on Targeting Polycomb Repressor
Proteins. PLOS Genetics *9*, e1003717.
Koornneef, M., Hanhart, C.J., and van der Veen, J.H. (1991). A genetic and physiological
analysis of late flowering mutants in Arabidopsis thaliana. Mol Gen Genet *229*, 57-66.
Leung, C.S., Douglass, S.M., Morselli, M., Obusan, M.B., Pavlyukov, M.S., Pellegrini, M.,
and Johnson, T.L. (2019). H3K36 Methylation and the Chromodomain Protein Eaf3
Are Required for Proper Cotranscriptional Spliceosome Assembly. Cell Reports *27*,
3760-3769.e3764.

Liu, F., Marquardt, S., Lister, C., Swiezewski, S., and Dean, C. (2010). Targeted 3'
Processing of Antisense Transcripts Triggers *Arabidopsis FLC* Chromatin Silencing.
Science *327*, 94-97.
Liu, F., Quesada, V., Crevillén, P., Bäurle, I., Swiezewski, S., and Dean, C. (2007). The
Arabidopsis RNA-Binding Protein FCA Requires a Lysine-Specific Demethylase 1
Homolog to Downregulate FLC. Molecular Cell *28*, 398-407.
Lövkvist, C., and Howard, M. (2021). Using computational modelling to reveal mechanisms
of epigenetic Polycomb control. Biochemical Society Transactions *49*, 71-77.
Mangeon, A., Magioli, C., Tarre, E., Cardeal, V., Araujo, C., Falkenbach, E., Rocha, C.A.,
Rangel-Lima, C., and Sachetto-Martins, G. (2010). The tissue expression pattern of the
AtGRP5 regulatory region is controlled by a combination of positive and negative
elements. Plant Cell Rep *29*, 461-471.
Marina, D.B., Shankar, S., Natarajan, P., Finn, K.J., and Madhani, H.D. (2013). A conserved
ncRNA-binding protein recruits silencing factors to heterochromatin through an RNAi-
independent mechanism. Genes & Development *27*, 1851-1856.
Marquardt, S., Raitskin, O., Wu, Z., Liu, F., Sun, Q., and Dean, C. (2014). Functional
consequences of splicing of the antisense transcript COOLAIR on FLC transcription.
Mol Cell *54*, 156-165.
Martignago, D., Bernardini, B., Polticelli, F., Salvi, D., Cona, A., Angelini, R., and
Tavladoraki, P. (2019). The Four FAD-Dependent Histone Demethylases of
Arabidopsis Are Differently Involved in the Control of Flowering Time. Frontiers in
Plant Science *10*.
Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. 2011 *17*, 3.
Mateo-Bonmatí, E., Xiaofeng, F., Robert, M., Marc, F., Lori, A.P., and Caroline, A.D.
(2023). The CPSF phosphatase module links transcription termination to chromatin
silencing. bioRxiv, https://doi.org/10.1101/2023.07.06.547976.
Menon, G., Schulten, A., Dean, C., and Howard, M. (2021). Digital paradigm for Polycomb
epigenetic switching and memory. Current Opinion in Plant Biology *61*, 102012.
Moll, P., Ante, M., Seitz, A., and Reda, T. (2014). QuantSeq 3' mRNA sequencing for RNA
quantification. Nature Methods *11*, i-iii.
Nojima, T., and Proudfoot, N.J. (2022). Mechanisms of lncRNA biogenesis as revealed by
nascent transcriptomics. Nature Reviews Molecular Cell Biology *23*, 389-406.
Pease, N.A., Nguyen, P.H.B., Woodworth, M.A., Ng, K.K.H., Irwin, B., Vaughan, J.C., and
Kueh, H.Y. (2021). Tunable, division-independent control of gene activation timing by
a polycomb switch. Cell Rep *34*, 108888.
Schon, M., Baxter, C., Xu, C., Enugutti, B., Nodine, M.D., and Dean, C. (2021). Antagonistic
activities of cotranscriptional regulators within an early developmental window set FLC
expression level. Proc Natl Acad Sci U S A *118*.
Shema, E., Jones, D., Shoresh, N., Donohue, L., Ram, O., and Bernstein, B.E. (2016). Single-
molecule decoding of combinatorially modified nucleosomes. Science *352*, 717-721.
Shimada, Y., Carl, S.H., Skribbe, M., Flury, V., Kuzdere, T., Kempf, G., and Bühler, M.
(2021). An enhancer screen identifies new suppressors of small-RNA-mediated
epigenetic gene silencing. PLOS Genetics *17*, e1009645.
Shu, J., Chen, C., Thapa, R.K., Bian, S., Nguyen, V., Yu, K., Yuan, Z.-C., Liu, J., Kohalmi,
S.E., Li, C.*, et al.* (2019). Genome-wide occupancy of histone H3K27
methyltransferases CURLY LEAF and SWINGER in Arabidopsis seedlings. Plant
Direct *3*, e00100.

Simpson, G.G., Dijkwel, P.P., Quesada, V., Henderson, I., and Dean, C. (2003). FY Is an
RNA 3' End-Processing Factor that Interacts with FCA to Control the Arabidopsis
Floral Transition. Cell *113*, 777-787.
Sonmez, C., Bäurle, I., Magusin, A., Dreos, R., Laubinger, S., Weigel, D., and Dean, C.
(2011). RNA 3' processing functions of *Arabidopsis* FCA and FPA limit intergenic
transcription. Proceedings of the National Academy of Sciences *108*, 8508-8513.
Stein, C.B., Field, A.R., Mimoso, C.A., Zhao, C., Huang, K.L., Wagner, E.J., and Adelman,
K. (2022). Integrator endonuclease drives promoter-proximal termination at all RNA
polymerase II-transcribed loci. Mol Cell *82*, 4232-4245 e4211.
Swiezewski, S., Liu, F., Magusin, A., and Dean, C. (2009). Cold-induced silencing by long
antisense transcripts of an Arabidopsis Polycomb target. Nature *462*, 799-802.
Tucker, J.F., Ohle, C., Schermann, G., Bendrin, K., Zhang, W., Fischer, T., and Zhang, K.
(2016). A Novel Epigenetic Silencing Pathway Involving the Highly Conserved 5'-3'
Exoribonuclease Dhp1/Rat1/Xrn2 in Schizosaccharomyces pombe. PLOS Genetics *12*,
e1005873.
Vo, T.V., Dhakshnamoorthy, J., Larkin, M., Zofall, M., Thillainadesan, G., Balachandran, V.,
Holla, S., Wheeler, D., and Grewal, S.I.S. (2019). CPF Recruitment to Non-canonical
Transcription Termination Sites Triggers Heterochromatin Assembly and Gene
Silencing. Cell Reports *28*, 267-281.e265.
Wang, H., Fan, Z., Shliaha, P.V., Miele, M., Hendrickson, R.C., Jiang, X., and Helin, K.
(2023). H3K4me3 regulates RNA polymerase II promoter-proximal pause-release.
Nature *615*, 339-348.
Wang, Z.W., Wu, Z., Raitskin, O., Sun, Q., and Dean, C. (2014). Antisense-mediated FLC
transcriptional repression requires the P-TEFb transcription elongation factor. Proc Natl
Acad Sci U S A *111*, 7468-7473.
Wittmann, S., Renner, M., Watts, B.R., Adams, O., Huseyin, M., Baejen, C., El Omari, K.,
Kilchert, C., Heo, D.-H., Kecman, T.*, et al.* (2017). The conserved protein Seb1 drives
transcription termination by binding RNA polymerase II and nascent RNA. Nature
Communications *8*, 14861.
Wu, Z., Ietswaart, R., Liu, F., Yang, H., Howard, M., and Dean, C. (2016). Quantitative
regulation of *FLC* via coordinated transcriptional initiation and elongation. Proc Natl
Acad Sci U S A *113*, 218-223.
Xu, C., Wu, Z., Duan, H.C., Fang, X., Jia, G., and Dean, C. (2021). R-loop resolution
promotes co-transcriptional chromatin silencing. Nat Commun *12*, 1790.
Yang, H., Berry, S., Olsson, T.S.G., Hartley, M., Howard, M., and Dean, C. (2017). Distinct
phases of Polycomb silencing to hold epigenetic memory of cold in *Arabidopsis*.
Science *357*, 1142-1145.
Yang, Y., Costa, A., Leonhardt, N., Siegel, R.S., and Schroeder, J.I. (2008). Isolation of a
strong Arabidopsis guard cell promoter and its potential as a research tool. Plant
Methods *4*, 6.
Zhao, Y., Zhu, P., Hepworth, J., Bloomer, R., Antoniou-Kourounioti, R.L., Doughty, J.,
Heckmann, A., Xu, C., Yang, H., and Dean, C. (2021). Natural temperature fluctuations
promote *COOLAIR* regulation of *FLC*. Genes Dev *35*, 888-898.
1089

Supplementary Information

Proximal termination generates a transcriptional state that determines the rate of establishment of Polycomb silencing

Govind Menon[1], Eduardo Mateo-Bonmati[2,5], Svenja Reeck[2], Robert Maple[2], Zhe Wu[3], Robert Ietswaart[4], Caroline Dean[2], Martin Howard[1]

[1]Department of Computational and Systems Biology, John Innes Centre, Norwich Research Park NR4 7UH, UK

[2]Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park NR4 7UH, UK

[3]Key Laboratory of Molecular Design for Plant Cell Factory of Guangdong Higher Education Institutes, Institute of Plant and Food Science, Department of Biology, School of Life Sciences, Southern University of Science and Technology, Shenzhen, China

[4]Harvard Medical School, Department of Genetics, 77 Avenue Louis Pasteur, Boston, MA 02115, USA

[5]Current address: Centro de Biotecnología y Genómica de Plantas, Universidad Politécnica de Madrid (UPM) – Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA)/CSIC, 28223 Pozuelo de Alarcón, Madrid, Spain

Computational Model

We built a stochastic model describing the dynamics of the activating and repressive histone modifications of interest, across a single *FLC* locus. The model also describes transcription events at the locus – sense and antisense – as well as alternative polyadenylation during these transcription events.

Model features:

Activating and repressive histone modifications

The mathematical model describes the modification state of each H3 histone at the *FLC* locus, with the whole locus represented by an array of 30 nucleosomes, consistent with the approximately 6 kilobases extent of the locus (nucleosomal DNA + linker DNA spans approximately 200 bp). The modifications described are: (1) H3K4me1, an activating modification associated with gene bodies in Arabidopsis, and known to exhibit significant differences between active and silenced states of *FLC*; (2) the Polycomb-mediated repressive modification H3K27me3 as well as possible H3K27me2, H3K27me1 and me0 states. The model assumes mutual exclusivity of K4me and K27me on the same H3 tail in line with experimentally observed mutual exclusivity in mESCs (Shema et al., 2016; Voigt et al., 2012) (Kim et al., 2013; Schmitges et al., 2011). Evidence from mammalian and yeast studies indicate that H3K27me and H3K4me are added non-processively (McCabe et al., 2012; Soares et al., 2017). Therefore, the model assumes stepwise addition of these modifications.

Sense and Antisense transcription

The model describes sense and antisense transcription events at the locus. The frequency of these events (assumed to represent transcription initiation or transition to elongation) is determined by the coverage of repressive H3K27me3 across the locus. This is consistent with Polycomb (PRC1) binding to H3K27me3 and controlling accessibility to gene regulatory elements by mediating compaction (Blackledge and Klose, 2021), as well as other potential H3K27me3-mediated mechanisms, such as inhibition of RNA Pol II release from the Transcription Start Site (TSS) (Zhang et al., 2020), thus preventing transitions into an actively transcribing Pol II state (Brookes et al., 2012). Our data indicates that both sense and antisense transcription at *FLC* are silenced by Polycomb – both types of transcripts are present at higher levels in the absence of Polycomb silencing, as observed in genotypes containing the *FLC* activator FRIGIDA, and in loss of function mutants of FCA and FLD.

Co-transcriptional addition of activating modifications

Activating modification such as H3K4me and H3K36me in many contexts and across organisms are known to be added co-transcriptionally (Chan et al., 2022). The association of the H3K4 methyltransferase complex with elongating RNA Pol II and the addition of this modification during transcriptional elongation has been demonstrated in *S. cerevisiae* (Soares et al., 2017). In the case of Arabidopsis H3K4me1, a genome wide analysis of the localization patterns of H3K4me1, of RNA Pol II, and of the different H3K4 methyltransferases, indicates that the co-transcriptional addition of H3K4me1 over the gene body is mediated mainly by the methyltransferase ATXR7 (Oya et al., 2022). Consistent with this role, an *ATXR7* mutant has been shown to partially suppress *FLC* activation by the activator FRIGIDA (Tamada et al., 2009).

Alternative polyadenylation mediated by coverage of activating modifications

The model assumes that higher overall coverage of H3K4me1 across the locus promotes distal termination of transcription. This assumption is in line with a potential feedback mechanism where H3K4me1 enables binding of the H3K36me3 methyltransferase (SDG8) and consequent H3K36me3-mediated assembly of splicing factors (Leung et al., 2019) which facilitate co-transcriptional processing. Such a role for H3K4 methylation is also consistent with evidence in yeast (Soares et al., 2017) and mammalian cells (Wang et al., 2023), indicating a role in facilitating productive transcriptional elongation.

Proximal termination mediated removal of activating modifications

The model assumes proximal termination of both sense and antisense mediates removal of H3K4me1. This is in line with the potential function of FLD as a demethylase (Inagaki et al., 2021), and the fact that genetically, FLD relies on FCA-mediated proximal termination of

79   antisense (Liu et al., 2007). The removal of H3K4me1 resulting from proximal PA is assumed

80   to take place uniformly across the whole locus, as the data we currently have does not allow

81   us to observe the spatial details of this process.

Polycomb nucleation and spreading

The model describes nucleation of H3K27me by Polycomb in the *FLC* nucleation region,

where Polycomb silencing is known to be nucleated during cold-induced chromatin silencing

of *FLC*. Subsequent H3K27me3 spreading across the locus in this context is known to depend

on the plant PRC2 methyltransferase CLF and on an active cell cycle, while nucleation does

not depend on either. Experimental evidence from a genome-wide ChIP-seq study showing

H3K27me3 patterns in a *clf* mutant (Shu et al., 2019) indicates that a similar mechanism is

involved in the early developmental silencing of *FLC* – the *clf* mutant shows a clear

H3K27me3 peak in the *FLC* nucleation region, while spreading is compromised. Consistent

with this, in the wild-type, H3K27me3 is assumed to spread to the rest of the locus through

interactions with the nucleation region, through read-write feedback interactions. The

spreading is assumed to rely on an active cell cycle – in the model, this is achieved by having

the interactions with the nucleation region stronger in S/G2 phases.

Looping interactions mediating Polycomb activity

Experimental evidence indicates the existence of a gene loop mediating an interaction

between the 5' and 3' ends of *FLC* (Crevillen et al., 2013; Li et al., 2018), with the level of

interaction potentially depending the transcriptional state of the locus (Li et al., 2018).

Significantly, this looping interaction is present in the silenced *FLC* state in the reference

genotype Col-0, and consistently, a secondary peak of H3K27me3 is observed in published

ChIP-seq data in *clf* (Shu et al., 2019). Therefore, we capture this interaction in the model by

allowing for stronger interactions with the nucleation region for a designated "looping region"

within the gene body.

106
107 Productive transcription antagonises H3K27me

Experimental evidence from mammalian systems indicates that productive transcription can

promote removal of H3K27me through histone exchange. In addition, the H3K27me

demethylases have been shown to physically interact with and co-localise on chromatin with

elongating RNA Pol II, in *Drosophila* (Smith et al., 2008) and more specifically at the *FLC*

locus (Yang et al., 2016). Both features have been captured in previous models of Polycomb

silencing (Berry et al., 2017), and they are also incorporated in the current model. These

features allow the rate of H3K27me turnover across the locus to be determined by the

frequency of transcription events. Our model also incorporates alternative PA of both sense

and antisense transcription, allowing proximal PA and termination of RNA Pol II. The model
assumes that only distal PA and termination (i.e., productive transcription events) can remove
H3K27me through the above mechanisms.

Mutual exclusivity of sense and antisense at the same locus
We previously found that distal antisense transcription and sense *FLC* transcription are
mutually exclusive at individual *FLC* loci (Rosa et al., 2016). Consistent with this mutual
exclusivity, we have observed an anticorrelation between the levels of distal antisense
transcripts and sense transcripts in multiple contexts, including different chromatin states of
the *FLC* locus (Ietswaart et al., 2017; Zhao et al., 2021). While this evidence demonstrates a
mutual exclusivity between full length (distal) sense and antisense transcription events, we
currently do not have data to determine whether the mutual exclusivity applies to proximally
terminated transcription events. However, for simplicity, the model assumes mutual
exclusivity for both types of transcription events. This is realised by having a refractory
period after a sense (antisense) transcription event, during which antisense (sense) initiation is
not allowed to occur. The model assumes equal refractory periods for sense and antisense
transcription, as well as for distal and proximal termination.

Inheritance of modified histones during DNA replication
In line with recent evidence indicating that only H3-H4 tetramers carrying repressive
modifications are reliably inherited across DNA replication (Escobar et al., 2021), the model
assumes that H3 pairs corresponding to an individual nucleosome are inherited with
probability half to a daughter strand, only if at least one of the H3 carries K27 methylation.

139
Additional considerations for the model
Absence of a role for the L*er* *Mutator*-like transposable element at *FLC*
The *FLC* allele in the L*er* accession is known to contain a *Mutator*-like transposable element (TE) in
intron 1 that causes lower expression from this allele relative to *FLC* alleles in other accessions
(Michaels et al., 2003). This lower expression results from RNAi-mediated chromatin silencing of the
TE element (Liu et al., 2004). However, experimental evidence indicates that the intermediate *FLC*
expression levels in *fca* mutants are unlikely to be mediated by the TE. The TE has been shown to act
in cis (Michaels et al., 2003), as expression of an *FLC* allele from the Columbia-0 (Col-0) accession is
unaffected in F1 plants generated by a L*er* to Col-0 cross, while the L*er* *FLC* allele expression is
attenuated by the TE. In our case, the intermediate expression in *fca* mutants could be observed for
both the endogenous *FLC* (L*er* *FLC* containing TE) and the transgenic Venus tagged *FLC* (Col-0
FLC sequence), arguing against a role for the *cis*-acting TE mechanism in generating intermediate
transcription states.

Role for H3K36me3

The above model only considers the histone modifications H3K4me1 and H3K27me, and does not explicitly consider H3K36me3. As discussed in the text, the Arabidopsis H3K36 methyltransferase SDG8 is known to recognise H3K4me1, thus mechanistically linking the two modifications. Combined with the H3K36me3 requirement for effective co-transcriptional splicing and recruitment of splicing factors as reported in *S. cerevisiae* (Leung et al., 2019) this suggests a mechanism where FLD-mediated removal of H3K4me1 may cause reduced H3K36 methylation, which in turn promotes further early transcription termination by disrupting co-transcriptional splicing. For simplicity, the model omits the H3K36me3 step and assumes that H3K4me1 implicitly mediates this effect on termination.

Direct physical interaction between transcription-coupled repression and Polycomb

FCA has been reported to interact directly with the Arabidopsis PRC2 methyltransferase CLF *in vivo* (Tian et al., 2019). However, we have found no evidence of this interaction in our FCA immunoprecipitation followed by mass spectrometry (IP-MS) data or other published CLF IP-MS data (Fang et al., 2019; Liang et al., 2015), and hence this interaction is not further considered in our model or other analyses.

Model implementation

Locus description: The whole locus represented by an array of 30 nucleosomes (60 H3 histones), consistent with the approximately 6 kilobase extent of the locus (nucleosomal DNA + linker DNA spans approximately 200 bp).

Mutually exclusive H3 modifications: Each H3 histone can be in one of five modification mutually exclusive states: {-1,0,1,2,3} representing H3K4me1, H3 unmodified on K4 and K27, H3K27me1, me2, and me3 respectively. This means addition of K27me can take place only in the absence of H3K4me1 and vice versa. All transitions are assumed to be non-processive: only transitions involving addition/removal of a single methyl group are allowed. The modification state of the locus is representing by an array S representing the modification states of the 60 histones. Each entry S_i can take the values {-1,0,1,2,3}. The array S is used to store the current state of all 60 histones over the course of the simulation. Note that we do not describe H3K36me3 in the model, as the evidence indicates it is downstream of H3K4me1, as described above and in the main text.

Alternative termination of transcription: Each transcription event (sense or antisense) has a probability of being proximally terminated. This probability is determined by both the current H3K4me1 fractional coverage (whole locus) and the FCA effectiveness parameter, as follows:

$$P(\text{Proximal}) = \delta_{Proximal} + (p_0 + p_{FCA})(1 - P_{K4me1})(1 - \delta_{Proximal} - \mu_{Distal}).$$

190   The probability of distal termination is given by:

$$P(\text{Distal}) = 1 - P(\text{Proximal}).$$

192   Here p_{FCA} represents FCA effectiveness parameter and p_0 captures possible FCA independent
193   proximal termination. $(p_0 + p_{FCA})$ is constrained to be less than or equal to 1. P_{K4me1} represents
194   the fractional coverage of H3K4me1, $\delta_{Proximal}$ is a residual proximal termination probability (some
195   proximal termination occurs even when $P_{K4me1} = 1$), and μ_{Distal} is a residual probability for distal
196   termination (some distal termination occurs even when $p_{FCA} = 1$ or $P_{K4me1} = 0$).
197
Transcriptional antagonism of Polycomb: (1) Histone exchange: during each distally terminated
transcription event, iterating over the histones (entries of array S), for each histone, we generate a
pseudo-random number from a uniform distribution (0,1) and set both entries corresponding to the
current nucleosome to 0 if the generated number is less than $p_{ex}/2$ (see Supplementary Table). (2)
Co-transcriptional Demethylation: During each distally terminated transcription event, iterating over
the entries of array S, for each histone, we generate a pseudo-random number from a uniform
distribution (0,1) and reduce the methylation state by one if the current state is {1, 2, or 3} AND the
generated number is less than p_{dem} (see Supplementary Table).
206
207   **Transcription mediated effects on H3K4 methylation:** (1) Co-transcriptional addition of
208   H3K4me1: Each unmodified H3 histone has a fixed probability p_{K4me}^{Transc}, of being methylated during a
209   productive (distally terminated) transcription event (not in case of proximal termination). (2) H3K4
210   demethylation by FLD: Each H3 histone carrying H3K4me1 has a fixed probability p_{dem}^{FLD} of losing
211   this modification during a proximally terminated transcription event.
212
H3K4me1 dynamics: The background turnover rate (independent of proximal termination and FLD)
of H3K4me1 is set so that overall turnover rates of H3K27me3 and H3K4me1 are in the ratio
3.128/0.959, as measured in HeLa cells by a SILAC technique in (Zee et al., 2010). The H3K4
methylation propensities (background and transcription mediated) are chosen to be significantly
higher than the K27me3 propensities, consistent with faster H3K4 methylation seen in time course
evidence from mammalian systems.
219
H3K27me3 dynamics: Values for methylation rate constants are mostly chosen as in (Berry et al.,
2017). The overall rate β alone is adjusted from its value in (Berry et al., 2017) to allow for a changed
interaction paradigm – instead of just nearest neighbour interactions, the current model allows long
ranged interactions between nucleosomes across the whole locus, with any nucleosome being able to

interact with any other within the locus at some rate (see below). To account for nucleation, spreading

and the looping interaction, the nucleosome array representing the whole locus is divided into three

subsets: nucleation region (NR, nucleosomes 1 to 3, i.e., histones 1 to 6), looping region (LR,

nucleosomes 20 to 22, i.e., histones 39 to 44), and body region (Body, remaining nucleosomes). The

choice of nucleosomes 19 to 21 for the looping region is consistent with the observed secondary

nucleation peak observed around 4kb downstream of the *FLC* TSS in published data for the *clf* mutant

(Shu et al., 2019).

The methylation propensities at each histone *i* are computed as:

$$r_i^{\mathrm{me}} = \beta \left(\delta_{S_i,0} (\gamma_{\mathrm{me0\text{-}1}} + \mathrm{k}_{\mathrm{me0\text{-}1}} \mathrm{E_i}) + \delta_{S_i,1} (\gamma_{\mathrm{me1\text{-}2}} + \mathrm{k}_{\mathrm{me1\text{-}2}} \mathrm{E_i}) + \delta_{S_i,2} (\gamma_{\mathrm{me2\text{-}3}} + \mathrm{k}_{\mathrm{me2\text{-}3}} \mathrm{E_i}) \right)$$

233 where E_i represents the PRC2 read-write feedback contribution.

235 For nucleation region histones, the PRC2 read-write feedback mediated contribution to the H3K27

236 methylation propensity is computed as:

$$E_i = \xi_{NR}^{NR} \sum_{j \in NR} \left(\rho_{\mathrm{me2}} \delta_{S_j,2} + \delta_{S_j,3} \right) + \xi_{LR}^{NR} \sum_{j \in LR} \left(\rho_{\mathrm{me2}} \delta_{S_j,2} + \delta_{S_j,3} \right) + \xi_{Body}^{NR} \sum_{j \in Body} \left(\rho_{\mathrm{me2}} \delta_{S_j,2} + \delta_{S_j,3} \right),$$

238 where ρ_{me2} is the relative activation of PRC2 by H3K27me2 as compared to H3K27me3. For the

239 looping region histones, the PRC2 read-write feedback mediated contribution to the H3K27

240 methylation propensity is computed as:

$$E_i = \xi_{NR}^{LR} \sum_{j \in NR} \left(\rho_{\mathrm{me2}} \delta_{S_j,2} + \delta_{S_j,3} \right) + \xi_{LR}^{LR} \sum_{j \in LR} \left(\rho_{\mathrm{me2}} \delta_{S_j,2} + \delta_{S_j,3} \right) + \xi_{Body}^{LR} \sum_{j \in Body} \left(\rho_{\mathrm{me2}} \delta_{S_j,2} + \delta_{S_j,3} \right).$$

241 For the remaining gene body histones, the PRC2 read-write feedback mediated contribution to the

242 H3K27 methylation propensity is computed as:

$$E_i = \xi_{NR}^{Body} \sum_{j \in NR} \left(\rho_{\mathrm{me2}} \delta_{S_j,2} + \delta_{S_j,3} \right) + \xi_{LR}^{Body} \sum_{j \in LR} \left(\rho_{\mathrm{me2}} \delta_{S_j,2} + \delta_{S_j,3} \right) + \xi_{Body}^{Body} \sum_{j \in Body} \left(\rho_{\mathrm{me2}} \delta_{S_j,2} + \delta_{S_j,3} \right).$$

Here the parameters ξ are chosen so that interactions between nucleosomes within the nucleation and

looping region histones produce the highest contribution to feedback methylation, followed by the

contribution of NR and LR nucleosomes to methylation of other gene body nucleosomes. This is

consistent with the PRC2 enzymatic machinery localising mainly to the nucleation region. Also

consistent with this, we assume lower contributions from gene body histones to the methylation of NR

and LR nucleosomes, and the lowest contribution from interactions between gene body histones.

Hence, we set the above parameters as follows:

$$\xi_{NR}^{NR} = \xi_{LR}^{NR} = \xi_{NR}^{LR} = \xi_{LR}^{LR} > \xi_{NR}^{Body} = \xi_{LR}^{Body} > \xi_{Body}^{NR} = \xi_{Body}^{LR} > \xi_{Body}^{Body}.$$

Following (Berry et al., 2017), a background "noisy" removal of H3K27 methylation is also included

in the model, represented by the rate constant γ_{dem}.

Dependence on an active cell cycle: To implement the dependence of H3K27me3 spreading on an

active cell cycle (as observed experimentally in (Yang et al., 2017)), the total cell cycle duration is

divided into two parts: G1 and S/G2/M, with replication of the locus occurring at the midpoint of

S/G2/M. The duration of G1 is assumed to be approximately half the total duration of the cell-cycle.

The four parameters corresponding to interactions between the NR/LR and the body region (ξ_{NR}^{Body},

ξ_{LR}^{Body}, ξ_{Body}^{NR}, ξ_{Body}^{LR}) are reduced by a factor of 10 during the part of the cell cycle corresponding to

G1 phase.

H3K27me3 spreading mutant: For simulating the H3K27me3 spreading mutant (*clf-2*), we lower

the read-write feedback contributions from the NR and LR nucleosomes to methylation of gene body

nucleosomes (i.e., lower ξ_{NR}^{Body} and ξ_{LR}^{Body}), as well as the contribution from the gene body

nucleosomes to methylation of the NR and LR nucleosomes (i.e., lower ξ_{Body}^{NR} and ξ_{Body}^{LR}). Since the

loss of one of the PRC2 methyltransferases may also be expected to partially affect H3K27me3

addition within the nucleation region itself, we also slightly reduce the feedback contributions

between nucleosomes inside the NR and LR regions (i.e., lower ξ_{NR}^{NR}, ξ_{LR}^{NR}, ξ_{NR}^{LR}, ξ_{LR}^{LR}). These

parameter values are specified in supplementary Table 2.

De novo addition of H3K27me3: In addition to the differences in PRC2 feedback mediated

contributions, the nucleation region (first three nucleosomes) is assumed to have a significantly higher

de novo contribution to the H3K27 methylation propensity (γ_{me} parameters), consistent with evidence

for *de novo* PRC2 targeting to this region during early developmental silencing of *FLC* (Shu et al.,

2019).

H3K27me3 fractional coverage measure: Our previous studies of *FLC* silencing indicate that

nucleation region H3K27me3 alone can significantly repress transcription, with the body region

H3K27me3 making a relatively smaller additional contribution to silencing. This finding is different

to the assumptions of Berry et al., where both H3K27me2 and H3K27me3 could repress.

Furthermore, nucleation of H3K27me3 in the cold is sufficient to switch the locus to a digitally OFF

state, as seen by FLC Venus imaging in the roots (Lovkvist et al., 2021). Potentially different

contributions of the nucleation and looping region H3K27me3 versus the body region H3K27me3 to

the transcriptional silencing are captured by allowing a higher contribution from these NR and LR

nucleosomes when computing the overall fractional coverage of H3K27me3:

$$P_{K27me3} = \eta P_{K27me3}^{NR+LR} + (1 - \eta) P_{K27me3}^{Body}$$

Here P_{K27me3}^{NR+LR} and P_{K27me3}^{Body} represent fractional coverage of H3K27me3 in the nucleation

region/looping region and the rest of the locus, respectively, and $0 < \eta < 1$. The overall fractional

coverage, P_{K27me3} computed above, is then used to determine the transcription propensity (see

below).

Polycomb silencing of transcription: H3K27me3 coverage across the locus is assumed to determine

the "transcription initiation" frequency. This frequency is assumed to be a piecewise linear function of

H3K27me3 coverage with a saturation threshold (similar to the model in (Berry et al., 2017)). Note

that this frequency can also represent the transition of initiated Pol II into the elongating phase.

Nucleation region and looping region contributions to the fractional coverage are assumed to be

higher than the rest of the locus (see above). H3K27me3 coverage also has the same effect on sense

and antisense initiation frequency.

$$f = \alpha(f_{max} - \frac{P_{K27me3}}{P_T}(f_{max} - f_{min})) \text{ for } P_{K27me3} < P_T$$

and

$$f = \alpha f_{min} \text{ for } P_{K27me3} \geq P_T$$

We also allow for a slightly higher transcription frequency in the H3K27me3 nucleated state

compared to a fully H3K27me3 spread state. This is consistent with our previous datasets, where *FLC*

transcriptional output is observed to reduce further during the post-cold transition to a spread state

following nucleation. To enable this behaviour, we set $\eta < P_T$ so that the full silencing threshold is

not reached even with full coverage of H3K27me3 in the nucleation and looping regions (i.e., even

with $P_{K27me3}^{NR+LR} = 1$).

Published evidence based on sequencing of nascent RNA as well as our own qPCR data (Wu et al.,

2016) indicates lower frequency of antisense transcription compared to sense transcription. This

difference between sense and antisense transcription is captured by using different values of α for

sense and antisense initiation frequency. Note that H3K4me1 does not affect the initiation frequency.

Histone inheritance during DNA replication: For each individual simulation, the DNA replication

times are fixed deterministically, assuming a cell cycle duration of 22h (Rahni and Birnbaum, 2019).

At each DNA replication timepoint, the Gillespie Stochastic Simulation Algorithm (SSA) is

interrupted. We then iterate over the entries of the array S, where for each nucleosome at the locus

carrying at least one H3K27me, its modification state is set to 0, i.e., the pair of entries corresponding

to the nucleosome is set to 0, with a probability of 0.5. We assume here random inheritance of intact

319	H3-H4 nucleosomes, with equal probability to each daughter strand. For each nucleosome, this is
320	done by generating a pseudo-random number from a uniform distribution (0,1) and setting the
321	corresponding histone modification states to 0 if the generated number is less than 0.5. In contrast, for
322	each nucleosome carrying no H3K27me, the modification state is always set to zero, i.e., no
323	inheritance of nucleosomes carrying only H3K4me1.
324	

Table 1: Model parameter values: H3K27me

Parameter	Description	Value used in model	Details
N	Number of histones	60	Approximately 6kb size of the *FLC* locus
k_{me}	PRC2-mediated methylation rate (histone^{-1} s^{-1})	$8 \cdot 10^{-6}$	Parameter value fitted to genome-wide H3K27me3 dynamics in Berry et al., 2017
$k_{\mathrm{me0\text{-}1}}$	PRC2-feedback mediated methylation rate (me0 to me1) (histone^{-1} s^{-1})	$9k_{\mathrm{me}}$	Following the model in Berry et al., 2017
$k_{\mathrm{me1\text{-}2}}$	PRC2-feedback mediated methylation rate (me1 to me2) (histone^{-1} s^{-1})	$6k_{\mathrm{me}}$	Following the model in Berry et al., 2017
$k_{\mathrm{me2\text{-}3}}$	PRC2-feedback mediated methylation rate (me2 to me3) (histone^{-1} s^{-1})	k_{me}	Following the model in Berry et al., 2017
$\gamma_{\mathrm{me2\text{-}3}}$	Noisy methylation rate (me2 to me3) (histone^{-1} s^{-1})	$k_{\mathrm{me}}/20$	Following the model in Berry et al., 2017
$\gamma_{\mathrm{me1\text{-}2}}$	Noisy methylation rate (me1 to me2) (histone^{-1} s^{-1})	$k_{\mathrm{me1\text{-}2}}/20$	Following the model in Berry et al., 2017
$\gamma_{\mathrm{me0\text{-}1}}$	Noisy methylation rate (me0 to me1) (histone^{-1} s^{-1})	$k_{\mathrm{me0\text{-}1}}/20$	Following the model in Berry et al., 2017
γ_{me2-3}^{NR}	De novo methylation rate (me2 to me3) in Nucleation region (histone^{-1} s^{-1})	$0.35\, \gamma_{\mathrm{me2\text{-}3}}$	Fitted in this study
γ_{me1-2}^{NR}	De novo methylation rate (me1 to me2) in Nucleation region	$0.35\, \gamma_{\mathrm{me1\text{-}2}}$	Fitted in this study

	(histone^{-1} s^{-1})		
γ_{me0-1}^{NR}	De novo methylation rate (me0 to me1) in Nucleation region (histone^{-1} s^{-1})	0.35 $\gamma_{me0\text{-}1}$	Fitted in this study
β	Relative local PRC2-activity	0.7	Fitted in this study
ρ_{me2}	Relative activation of PRC2 by H3K27me2	0.1	Following the model in Berry et al., 2017
γ_{dem}	Noisy demethylation rate of H3K27 (histone^{-1} s^{-1})	$8.8 \cdot 10^{-7}$	Fitted in this study

Table 2: Model parameter values: Nucleosome interaction parameters for PRC2 read-write feedback

$\xi_{NR}^{NR}=\xi_{LR}^{NR}=\xi_{NR}^{LR}=\xi_{LR}^{LR}$	Interactions within nucleation and looping regions	1 (Ler, $fca\text{-}3$, $fca\text{-}1$), 0.8 ($clf\text{-}2$, $fca\text{-}3$ $clf\text{-}2$)	Fitted in this study
$\xi_{NR}^{Body}=\xi_{LR}^{Body}$	Nucleation and looping region contributions to methylation of gene body nucleosomes	0.5 (Ler, $fca\text{-}3$, $fca\text{-}1$) 0.05 ($clf\text{-}2$, $fca\text{-}3$ $clf\text{-}2$)	Fitted in this study
$\xi_{Body}^{NR}=\xi_{Body}^{LR}$	Gene body contributions to methylation of nucleation and looping region nucleosomes	0.2 (Ler, $fca\text{-}3$, $fca\text{-}1$) 0.12 ($clf\text{-}2$, $fca\text{-}3$ $clf\text{-}2$)	Fitted in this study
ξ_{Body}^{Body}	Interactions within gene body region	0.01	Fitted in this study

326
327

Table 3: Model parameter values: Transcription and PA site choice

α_{sense}	Trans-acting gene activation for sense transcription	1	Fixed to 1 for *FLC* sense transcription
$\alpha_{antisense}$	Trans-acting gene activation for antisense transcription	0.2	Set to a lower value relative to α_{sense}, consistent with lower frequency of antisense transcription compared to sense transcription under pre-cold warm conditions (Rosa et al., 2016)
f_{min}	Minimum transcription initiation	$f_{max}/60$	Fitted in this study

	rate (s^{-1})		
f_{max}	Maximum transcription initiation rate at *FLC* (s^{-1})	$0.58 \cdot 10^{-3}$	*FLC* sense transcription initiation frequency computed using the same approach as Ietswaart et al., 2017, but using single cell mRNA counts and *FLC* mRNA half-life measured for *fca-1* in Antoniou-Kourounioti et al., 2023
P_T	H3K27me3 coverage threshold for full silencing of transcription	1/3	Following the model in Berry et al., 2017
$\delta_{Proximal}$	Residual probability of proximal termination	0.05	Fixed in this study
μ_{Distal}	Residual probability of Distal termination	0.05	Fixed in this study
p_0	Parameter representing FCA independent proximal termination	0.8	Fitted in this study
p_{FCA}	Parameter representing FCA mediated proximal termination	0.2 (L*er, clf-2*), 0.15 (*fca-3*), 0 (*fca-1*)	Fitted in this study
η	Weight parameter for relative contribution of NR+LR H3K27me3 to transcriptional silencing	0.25	Fitted in this study

Table 4: Model parameter values: Transcription effects on modifications

p_{dem}	Demethylation probability of H3K27me (histone^{-1} distal transcription^{-1})	0.1	Fitted in this study
p_{ex}	Histone exchange probability (histone^{-1} distal transcription^{-1})	$1 \cdot 10^{-3}$	Following the model in Berry et al., 2017
p_{dem}^{FLD}	Demethylation probability of H3K4me1 (histone^{-1} proximal transcription^{-1})	0.5	Fitted in this study
p_{K4me}^{Transc}	Methylation probability for unmodified H3K4 during a distally terminated transcription event	$100\frac{k_{me}}{f_{max}}$	Computed as 100 times the PRC2-mediated methylation rate k_{me} (histone^{-1} s^{-1}), then normalised by transcription frequency (s^{-1}) to obtain

			probability per histone per transcription event.

Simulations

Model simulations were carried out using the Gillespie exact SSA (Gillespie, 1977). Simulations were

coded in C++ and compiled using Clang/LLVM (version 14.0.0). The Mersenne Twister 19937

algorithm (Matsumoto and Nishimura, 1998), implemented in the GNU Scientific Library (GSL

version 2.7), was used to generate pseudorandom numbers for the Gillespie SSA. The simulation code

can be accessed at:

https://github.com/gm1613/FLC_transcription_coupled_repression_model.git

Initial conditions

340 In all cases – both for the analog module in isolation and the full model, the initial state was set to full

341 coverage of H3K4me1.

Analysis of simulation output

Average levels of fractional coverage of histone modifications and average frequency of different

transcription events and termination site usage (sense distal, antisense distal, sense proximal,

antisense proximal) were computed by averaging over 500 individual trajectories. Individual

trajectories were started at random (uniformly distributed) times in the cell cycle to avoid having all

trajectories synchronised. To quantify switching to the digital OFF state (which in the model can

correspond either to a Polycomb nucleated state or a spread state), we have to define what qualifies as

an OFF state. In our analysis we classify an OFF state as any state for which:

$$P_{K27me3} > \chi P_T$$

351 where the parameter χ is chosen so that $\chi P_T < \eta$, and all the other parameters/quantities are as

352 defined above. This condition on χ ensures that an H3K27me3 build-up in the nucleation region (even

353 if this modification is absent across the rest of the locus), can be sufficient to qualify as a digitally

354 OFF state. In all simulation results shown , χ is set to 0.25.

Comparing simulation output to experimental data

Comparison to timecourse ChIP-qPCR data: In Fig. 4 we compare the model predicted levels of

H3K27me3 and H3K4me1 to the normalised levels measured by ChIP-qPCR (averaged over primers

across the locus). The model predicted levels consist of an average over multiple simulated copies

(1000 copies in all cases). While the model only describes a set of *FLC* copies in dividing cells, the

ChIP-qPCR measurement is performed on whole-seedling tissue. Therefore, we compare a model

predicted population average to the experimental data. Note that we expect there to be heterogeneity

in the modification levels between tissues, and differences in the dynamics between dividing and non-dividing cells. We have previously addressed these differences at *FLC* in detail using a population level Ordinary Differential Equation model of *FLC* copies switching between different states (Questa et al., 2020). However, capturing these aspects is beyond the scope of the current study. We therefore make a simpler assumption: that at all timepoints and across genotypes, there exist two subpopulations of *FLC* copies (at the whole plant level) in addition to the population of copies represented in the model. We assume that one of these subpopulations consists entirely of active *FLC* copies (with high H3K4me1 coverage) with no possibility of PRC2 silencing, and the other subpopulation consists of PRC2 silenced copies (with high H3K27me3 coverage), with no possibility of switching to an active state. We assume that these copies form a fixed fraction of the total *FLC* copies, fixed across timepoints and across genotypes (hence independent of FCA). Note that these subpopulations must also consist of dividing cells in order for their fractions to be maintained constant over time. We do not explicitly simulate these subpopulations. Instead, we account for their contribution to the averaged histone modification levels. The contribution from each copy in the active subpopulation to H3K4me1 is assumed to be the same as the average steady state level at non-Polycomb silenced copies in the *fca-1* (null mutant) model, and with no contribution to the H3K27me3 level. Similarly, the contribution from each copy in the silenced subpopulation to H3K27me3 is assumed to be the same as the average level at PRC2 silenced copies predicted by the L*er* model (these levels are similar in the L*er, fca-3, and fca-1* models). We assume that these silenced copies make no contribution to the H3K4me1 level. The sizes of these two subpopulations relative to the population of copies represented by the model form two model parameters that we change to fit the data. The fits presented in Fig. 4C,D assume that the active subpopulation is 0.2 times the size of the modelled population, while the silenced subpopulation is 0.1 times the size of the modelled population.

Comparison to ChIP profile across *FLC* : In Fig. 5E, we compare the model predicted H3K27me3 profiles at *FLC* in the L*er* parental genotype and the Polycomb spreading mutant *clf-2* to the ChIP-qPCR measured profiles in these genotypes. Note that the model predicted profile is an average across multiple simulated copies (1000 copies). To do this we used an approach developed in (Wu et al., 2016) for comparing model predicted Pol II profiles to those measured by ChIP. In our case, this method generates a predicted H3K27me3 ChIP profile by combining the fully spatially resolved model predicted profile with the resolution limiting effects arising from the sonication generated fragment size distribution inherent in ChIP based measurements. Mathematically, this method involves a convolution of the full model predicted profile with the probability distribution of fragment sizes generated during the ChIP experiment. Here we use the same fragment size distribution estimated in (Wu et al., 2016). The code used to carry out this analysis is also provided at:

https://github.com/gm1613/FLC_transcription_coupled_repression_model.git

Comparison of predicted *FLC* shutdown: In Fig. 5C and Fig. S5A, we compare the model
predicted *FLC* transcriptional shutdown over time to changes in *FLC* unspliced (Fig. 5C)and *FLC*
spliced (Fig. S5A) RNA levels, measured by qPCR at the whole plant level. The model predicted *FLC*
transcriptional output is computed as the average frequency of *FLC* sense distal transcription events
(transcription events per hour). This frequency is computed by averaging sense distal transcription
events over a 6hr time window (an approximation based on our estimate of ~6hr *FLC* mRNA half-
life) leading to the respective sampling timepoints (to match the experiments, we choose 10 day, 14
day, and 21 day timepoints). We note that the model is not designed to capture absolute RNA levels in
the different genotypes, since this may depend on differences in their stability, as well as
heterogeneity between tissue types which are not described in the model. What the model is intended
to capture, is the quantitative trends, including differences between genotypes and changes over time,
i.e. a reduction over time in some genotypes which may be disrupted in others. A whole-plant level
measurement of *FLC* RNA is an appropriate quantity for capturing these types of differences, and we
therefore use these measurements for comparison to the model predictions. Since the model does not
capture absolute RNA levels, for comparison of experimental timecourse data to model predictions,
we consider changes (i.e., fold changes) relative to the first timepoint (10 days).

Supplemental Table S1. Primer sets used in this work.

Purpose	Oligonucleotide name	Oligonucleotide sequences (5' → 3')	
		Forward primer (F)	Reverse primer (R)
Genotyping	clf-2	AGAGAAGCTCAAACAAGCCATCGA	TTAACCCGGACCCGCATTTGTTTCGG
	LB_clf-2	GTCGGCGTGCGGCTGGCGGCG	
RT	FLC spliced		TTTGTCCAGCAGGTGACATC
	FLC unspliced		CTTTGTAATCAAAGGTGGAGAGC
	UBC		TTGTGCCATTGAATTGAACCC
qPCR	FLC spliced	AGCCAAGAAGACCGAACTCA	TTTGTCCAGCAGGTGACATC
	FLC unspliced	CGCAATTTTCATAGCCCTTG	CTTTGTAATCAAAGGTGGAGAGC
	UBC	CTGCGACTCAGGGAATCTTCTAA	TTGTGCCATTGAATTGAACCC

ChIP	FLC_-2285	ATCCAGAAAAGGGCAAGGAG	CGAATCGATTGGGTGAATG
	FLC_-1788	GGATTGATGTGGGGCACTAT	AGTCATGGGTAGGGCATGTG
	FLC_-1555	TGGAGGGAACAACCTAATGC	TCATTGGACCAAACCAAACC
	FLC_-321	ACTATGTAGGCACGACTTTGGTAAC	TGCAGAAAGAACCTCCACTCTAC
	FLC_5	GCCCGACGAAGAAAAGTAG	TCCTCAGGTTTGGGTTCAAG
	FLC_246	CGACAAGTCACCTTCTCCAAA	AGGGGGAACAAATGAAAACC
	FLC_473	GGCGGATCTCTTGTTGTTTC	CTTCTTCACGACATTGTTCTTCC
	FLC_741	CGTGCTCGATGTTGTTGAGT	TCCCGTAAGTGCATTGCATA
	FLC_1212	CCTTTTGCTGTACATAAACTGGTC	CCAAACTTCTTGATCCTTTTTACC
	FLC_1613	TTGACAATCCACAACCTCAATC	TCAATTTCCTAGAGGCACCAA
	FLC_2094	AGCCTTTTAGAACGTGGAACC	TCTTCCATAGAAGGAAGCGACT
	FLC_2523	AGTTTGGCTTCCTCATACTTATGG	CAATGAACCTTGAGGACAAGG
	FLC_3276	GGGGCTGCGTTTACATTTTA	GTGATAGCGCTGGCTTTGAT
	FLC_3699	TGAAATGTTACGAATACTAGCGTGT	GGATCAAAACTACTAGCTAACCCTTG
	FLC_4406	AGAACAACCGTGCTGCTTTT	TGTGTGCAAGCTCGTTAAGC
	FLC_5090	CCGGTTGTTGGACATAACTAGG	CCAAACCCAGACTTAACCAGAC
ChIP (continuation)	FLC_5599	TGGTTGTTATTTGGTGGTGTG	ATCTCCATCTCAGCTTCTGCTC
	FLC_5715	CCTGCTGGACAAATCTCCGA	GGATTTTGATTTCAACCGCCGA
	FLC_6013	CGTGTGAGAATTGCATCGAG	AAAAACGCGCAGAGAGAGAG
	FLC_6189	TCCTAAACGCGTATGGTTGG	CCTTCATGGATGACGGAACT
	FLC_6480	TCCAGACGCCATTGTCATTA	AGAGTGCATTTTAACACTGACGA
	FLC_6810	TTGTAAAGTCCGATGGAGACG	ACTCGGCGAGAAAGTTTGTG

FLC_7091	CATCGCTGTGTTATGCGTTT	GGCAAGTGTCGCAGTAAACA
STM (for H3K27me3 normalization)	GCCCATCATGACATCACATC	GGGAACTACTTTGTTGGTGGTG
ACT_+122 (for H3K36me3 normalization)	CGTTTCGCTTTCCTTAGTGTTAGCT	AGCGAACGGATCTAGAGACTCACCTTG
ACT+939 (for H3K4me1 normalization)	TGCCCCGAGAGCAGTGTTCC	TGGACTGAGCTTCATCACCAACG
IGN5 (for FLD normalization)	CCCTTAAGCGGACATGGTT	AATGTCGGCCAATCTTCTTG

AT3G47470	LHCA4
AT3G54890	LHCA1
AT2G43970	LARP6B
AT1G22690	GASA9
AT3G25600	CML15
AT1G76100	PETE1
AT4G17970	ALMT2
AT2G33340	MAC3B
AT5G24290	MEB2
AT2G15880	LRX10
AT2G30980	GSK3 like kinase
AT4G15910	DI21
AT4G29190	OZF2
AT2G05520	GRP3
AT5G09400	KUP7
AT2G41090	CML10
AT1G64780	AMT12
AT1G66240	ATX1
AT5G22270	SEID1
AT1G80830	NRAMP1
AT4G39350	CESA2
AT2G33770	PHO2
AT1G1211	NRT1/CHL1
AT1G23020	FRO3
AT2G41240	BHLH100
AT1G68880	ATBZIP
AT5G53550	YSL3

AT5G09220	AAP2
AT3G16400	NSP1
AT3G20470	GRP5
AT3G28910	MYB30
AT2G21170	TIM
AT4G15093	LIGB
AT3G44735	PSK3
AT2G41420	WIH2
AT4G17870	PYR1
AT1G32450	NPF7
AT1G13280	AOC4
AT5G58670	PLC1
AT2G35880	WDL4
AT1G66180	ASP
AT3G20310	ERF7
AT1G49860	GSTF14
AT4G34710	ADC2
AT2G03090	EXP15
AT3G14310	PME3
AT5G18960	FRS12
AT1G56110	NOP56
AT3G17170	RFC3
AT3G14210	ESM1
AT4G26630	DEK3
AT4G13850	GRP2
AT1G11840	GLX1
AT2G19480	NAP1
AT3G09820	ADK1
AT4G13430	IIL1
AT1G80600	TUP5/WIN1
AT5G37780	CAM1
AT5G65670	IAA9

References

Berry, S., Dean, C., and Howard, M. (2017). Slow Chromatin Dynamics Allow Polycomb
Target Genes to Filter Fluctuations in Transcription Factor Activity. Cell Syst *4*, 445-
457 e448.
Blackledge, N.P., and Klose, R.J. (2021). The molecular principles of gene regulation by
Polycomb repressive complexes. Nature Reviews Molecular Cell Biology *22*, 815-833.
Brookes, E., de Santiago, I., Hebenstreit, D., Morris, K.J., Carroll, T., Xie, S.Q., Stock, J.K.,
Heidemann, M., Eick, D., Nozaki, N.*, et al.* (2012). Polycomb associates genome-wide
with a specific RNA polymerase II variant, and regulates metabolic genes in ESCs. Cell
Stem Cell *10*, 157-170.
Chan, J., Kumar, A., and Kono, H. (2022). RNAPII driven post-translational modifications of
nucleosomal histones. Trends in Genetics *38*, 1076-1095.
Crevillen, P., Sonmez, C., Wu, Z., and Dean, C. (2013). A gene loop containing the floral
repressor FLC is disrupted in the early phase of vernalization. The EMBO journal *32*,
140-148.
Escobar, T.M., Loyola, A., and Reinberg, D. (2021). Parental nucleosome segregation and
the inheritance of cellular identity. Nature Reviews Genetics *22*, 379-392.
Fang, X., Wang, L., Ishikawa, R., Li, Y., Fiedler, M., Liu, F., Calder, G., Rowan, B., Weigel,
D., Li, P.*, et al.* (2019). Arabidopsis FLL2 promotes liquid-liquid phase separation of
polyadenylation complexes. Nature *569*, 265-269.
Gillespie, D.T. (1977). Exact stochastic simulation of coupled chemical reactions. The
journal of physical chemistry *81*, 2340-2361.
Ietswaart, R., Rosa, S., Wu, Z., Dean, C., and Howard, M. (2017). Cell-Size-Dependent
Transcription of *FLC* and Its Antisense Long Non-coding RNA *COOLAIR* Explain
Cell-to-Cell Expression Variation. Cell Syst *4*, 622-635 e629.
Inagaki, S., Takahashi, M., Takashima, K., Oya, S., and Kakutani, T. (2021). Chromatin-
based mechanisms to coordinate convergent overlapping transcription. Nat Plants *7*,
295-302.
Kim, D.-H., Tang, Z., Shimada, M., Fierz, B., Houck-Loomis, B., Bar-Dagen, M., Lee, S.,
Lee, S.-K., Muir, T.W., Roeder, R.G.*, et al.* (2013). Histone H3K27 Trimethylation
Inhibits H3 Binding and Function of SET1-Like H3K4 Methyltransferase Complexes.
Molecular and Cellular Biology *33*, 4936-4946.
Leung, C.S., Douglass, S.M., Morselli, M., Obusan, M.B., Pavlyukov, M.S., Pellegrini, M.,
and Johnson, T.L. (2019). H3K36 Methylation and the Chromodomain Protein Eaf3
Are Required for Proper Cotranscriptional Spliceosome Assembly. Cell Reports *27*,
3760-3769.e3764.
Li, Z., Jiang, D., and He, Y. (2018). *FRIGIDA* establishes a local chromosomal environment
for *FLOWERING LOCUS C* mRNA production. Nature Plants *4*, 836-846.
Liang, S.C., Hartwig, B., Perera, P., Mora-García, S., de Leau, E., Thornton, H., de Alves,
F.L., Rapsilber, J., Yang, S., James, G.V.*, et al.* (2015). Kicking against the PRCs – A
Domesticated Transposase Antagonises Silencing Mediated by Polycomb Group
Proteins and Is an Accessory Component of Polycomb Repressive Complex 2. PLOS
Genetics *11*, e1005660.
Liu, F., Quesada, V., Crevillén, P., Bäurle, I., Swiezewski, S., and Dean, C. (2007). The
Arabidopsis RNA-Binding Protein FCA Requires a Lysine-Specific Demethylase 1
Homolog to Downregulate FLC. Molecular Cell *28*, 398-407.
Liu, J., He, Y., Amasino, R., and Chen, X. (2004). siRNAs targeting an intronic transposon in
the regulation of natural flowering behavior in Arabidopsis. Genes Dev *18*, 2873-2878.
Lovkvist, C., Mikulski, P., Reeck, S., Hartley, M., Dean, C., and Howard, M. (2021). Hybrid
protein assembly-histone modification mechanism for PRC2-based epigenetic
switching and memory. Elife *10*.

Matsumoto, M., and Nishimura, T. (1998). Mersenne twister: a 623-dimensionally
equidistributed uniform pseudo-random number generator. ACM Trans. Model.
Comput. Simul. *8*, 3–30.
McCabe, M.T., Graves, A.P., Ganji, G., Diaz, E., Halsey, W.S., Jiang, Y., Smitheman, K.N.,
Ott, H.M., Pappalardi, M.B., Allen, K.E.*, et al.* (2012). Mutation of A677 in histone
methyltransferase EZH2 in human B-cell lymphoma promotes hypertrimethylation of
histone H3 on lysine 27 (H3K27). Proc Natl Acad Sci U S A *109*, 2989-2994.
Michaels, S.D., He, Y., Scortecci, K.C., and Amasino, R.M. (2003). Attenuation of
FLOWERING LOCUS C activity as a mechanism for the evolution of summer-annual
flowering behavior in Arabidopsis. Proc Natl Acad Sci U S A *100*, 10102-10107.
Oya, S., Takahashi, M., Takashima, K., Kakutani, T., and Inagaki, S. (2022). Transcription-
coupled and epigenome-encoded mechanisms direct H3K4 methylation. Nature
Communications *13*, 4521.
Questa, J.I., Antoniou-Kourounioti, R.L., Rosa, S., Li, P., Duncan, S., Whittaker, C., Howard,
M., and Dean, C. (2020). Noncoding SNPs influence a distinct phase of Polycomb
silencing to destabilize long-term epigenetic memory at Arabidopsis *FLC*. Genes Dev
34, 446-461.
Rahni, R., and Birnbaum, K.D. (2019). Week-long imaging of cell divisions in the
Arabidopsis root meristem. Plant Methods *15*, 30.
Rosa, S., Duncan, S., and Dean, C. (2016). Mutually exclusive sense–antisense transcription
at *FLC* facilitates environmentally induced gene repression. Nature Communications *7*,
13031.
Schmitges, Frank W., Prusty, Archana B., Faty, M., Stützer, A., Lingaraju,
Gondichatnahalli M., Aiwazian, J., Sack, R., Hess, D., Li, L., Zhou, S.*, et al.* (2011).
Histone Methylation by PRC2 Is Inhibited by Active Chromatin Marks. Molecular Cell
42, 330-341.
Shema, E., Jones, D., Shoresh, N., Donohue, L., Ram, O., and Bernstein, B.E. (2016). Single-
molecule decoding of combinatorially modified nucleosomes. Science *352*, 717-721.
Shu, J., Chen, C., Thapa, R.K., Bian, S., Nguyen, V., Yu, K., Yuan, Z.-C., Liu, J., Kohalmi,
S.E., Li, C.*, et al.* (2019). Genome-wide occupancy of histone H3K27
methyltransferases CURLY LEAF and SWINGER in Arabidopsis seedlings. Plant
Direct *3*, e00100.
Smith, E.R., Lee, M.G., Winter, B., Droz, N.M., Eissenberg, J.C., Shiekhattar, R., and
Shilatifard, A. (2008). *Drosophila* UTX Is a Histone H3 Lys27 Demethylase That
Colocalizes with the Elongating Form of RNA Polymerase II. Molecular and Cellular
Biology *28*, 1041-1046.
Soares, L.M., He, P.C., Chun, Y., Suh, H., Kim, T., and Buratowski, S. (2017). Determinants
of Histone H3K4 Methylation Patterns. Molecular Cell *68*, 773-785.e776.
Tamada, Y., Yun, J.-Y., Woo, S.c., and Amasino, R.M. (2009). ARABIDOPSIS
TRITHORAX-RELATED7 Is Required for Methylation of Lysine 4 of Histone H3 and
for Transcriptional Activation of *FLOWERING LOCUS C* The Plant Cell *21*, 3257-
3269.
Tian, Y., Zheng, H., Zhang, F., Wang, S., Ji, X., Xu, C., He, Y., and Ding, Y. (2019). PRC2
recruitment and H3K27me3 deposition at *FLC* require FCA binding of *COOLAIR*.
Science Advances *5*, eaau7246.
Voigt, P., LeRoy, G., Drury, William J., III, Zee, Barry M., Son, J., Beck, David B., Young,
Nicolas L., Garcia, Benjamin A., and Reinberg, D. (2012). Asymmetrically Modified
Nucleosomes. Cell *151*, 181-193.

Wang, H., Fan, Z., Shliaha, P.V., Miele, M., Hendrickson, R.C., Jiang, X., and Helin, K.
(2023). H3K4me3 regulates RNA polymerase II promoter-proximal pause-release.
Nature *615*, 339-348.
Wu, Z., Ietswaart, R., Liu, F., Yang, H., Howard, M., and Dean, C. (2016). Quantitative
regulation of *FLC* via coordinated transcriptional initiation and elongation. Proc Natl
Acad Sci U S A *113*, 218-223.
Yang, H., Berry, S., Olsson, T.S.G., Hartley, M., Howard, M., and Dean, C. (2017). Distinct
phases of Polycomb silencing to hold epigenetic memory of cold in *Arabidopsis*.
Science *357*, 1142-1145.
Yang, H., Howard, M., and Dean, C. (2016). Physical coupling of activation and derepression
activities to maintain an active transcriptional state at *FLC*. Proc Natl Acad Sci U S A
113, 9369-9374.
Zee, B.M., Levin, R.S., Xu, B., LeRoy, G., Wingreen, N.S., and Garcia, B.A. (2010). In vivo
residue-specific histone methylation dynamics. J Biol Chem *285*, 3341-3350.
Zhang, Y.-Z., Yuan, J., Zhang, L., Chen, C., Wang, Y., Zhang, G., Peng, L., Xie, S.-S., Jiang,
J., Zhu, J.-K.*, et al.* (2020). Coupling of H3K27me3 recognition with transcriptional
repression through the BAH-PHD-CPL2 complex in Arabidopsis. Nature
Communications *11*, 6212.
Zhao, Y., Zhu, P., Hepworth, J., Bloomer, R., Antoniou-Kourounioti, R.L., Doughty, J.,
Heckmann, A., Xu, C., Yang, H., and Dean, C. (2021). Natural temperature fluctuations
promote *COOLAIR* regulation of *FLC*. Genes Dev *35*, 888-898.
547
548
549

Figure 1

Figure 3

A Transcription-coupled repression

Polycomb silencing

Antisense Transcription

Proximal Termination

H3K4 Methyltransferase

FCA

FLD

H3K4me1

FLD

FCA

H3K27me3 is self-reinforcing through PRC2 read-write feedback

H3K27me3 silences transcription

PRC2

H3K27me3

Proximal Termination

Proximal Termination

H3K4 Methyltransferase

Sense Transcription

Productive transcription antagonises H3K27me3

Analogue regulation

Digital regulation

B FCA effectiveness parameter

C Model predictions for transcription-coupled repression alone (without Polycomb silencing)

Can generate intermediate transcriptional output at a single locus

ON state distal transcription (h⁻¹)

Can generate intermediate levels of H3K4me1 at a single locus

ON state fractional coverage: H3K4me1

Predictions from full model (including Polycomb silencing)

D Predicted transcriptional changes over time in L*er*

Average transcription frequency (hr⁻¹)

Sense Proximal Termination

Sense Distal Termination

Antisense Proximal Termination

Antisense Distal Termination

Days

E Predicted transcriptional state changes over time in L*er*

Fraction of *FLC* ON copies

Days

L*er*
fca-3
fca-1

Figure 4

Figure S4A

Figure S4B

(i)

Model predicted differences in switch-OFF times

Ler

fca-3

fca-1

Probability density

Time taken to switch to Polycomb silenced
OFF state (days)

(ii) Bistability in the absence of the transcription-
coupled repression mechanism

Fraction of ON copies

**Fractional coverage of
H3K27me3**

Starting in ON state

Time (Cell Cycles)

Time (Cell Cycles)

Starting in OFF state

Time (Cell Cycles)

Time (Cell Cycles)

(iii)

7 days

14 days

21 days

Ler | *fca-3* | *fca-1*

H3K36me3/H3/*ACT7*

Distance from *FLC* TSS (kb)

Figure S4C

Figure S4D

Figure 5

Figure S5A

Comparing model predictions to *FLC* spliced transcript levels

Figure S5B

(i)

(ii)

Figure S5C

Figure 6

A *FLC* expression in Col-0 (*FLC* spliced)

B
- *3xFLAG-FLD* (fri 1W)
- *3xFLAG-FLD* (fri 2W)
- *3xFLAG-FLD* (fri 3W)
- Col-0 (non-transgenic)

C n=107 *fca-1* upregulated genes with intermediate upregulation in *fca-3*

D n=107 *fca-1* upregulated DEGs with intermediate upregulation in *fca-3*

CLF ChIP   SWN ChIP

— *fca-1* upregulated; *fca-3* intermediate upregulation
— All genes

E Upregulated DEGs in *fca-9, fld-4,* and the intersection of these subsets

CLF ChIP   SWN ChIP

— *fld-4* upregulated
— *fca-9* upregulated
— *fld-4* and *fca-9* upregulated
— All genes

Figure S6

Figure 7

ON state

Transcription-coupled repression sets level of productive transcription (FCA, FLD)

High transcription

Intermediate transcription

Low transcription

Polycomb mediated ON to OFF switch

High switching probability

Intermediate switching probability

Low switching probability

OFF state (Epigenetically silenced)

L*er*

fca-3

fca-1

Level of productive transcription at individual locus