1

NEURAL CORRELATES OF PERCEPTUAL SIMILARITY MASKING IN PRIMATE V1

Spencer C. Chen[1-5], Yuzhi Chen[1-4], Wilson S. Geisler[1-3], & Eyal Seidemann[1-4]

[1]Center for Perceptual Systems, [2]Center for Theoretical and Computational Neuroscience
[3]Department of Psychology, [4]Department of Neuroscience, University of Texas at Austin
[5]Department of Neurosurgery, Rutgers University-

Abstract

Visual detection is a fundamental natural task. Detection becomes more challenging as the similarity between the target and the background in which it is embedded increases, a phenomenon termed "similarity masking". To test the hypothesis that V1 contributes to similarity masking, we used voltage sensitive dye imaging (VSDI) to measure V1 population responses while macaque monkeys performed a detection task under varying levels of target-background similarity. Paradoxically, we find that during an initial transient phase, V1 responses to the target are enhanced, rather than suppressed, by target-background similarity. This effect reverses in the second phase of the response, so that in this phase V1 signals are positively correlated with the behavioral effect of similarity. Finally, we show that a simple model with delayed divisive normalization can qualitatively account for our findings. Overall, our results support the hypothesis that a nonlinear gain control mechanism in V1 contributes to perceptual similarity masking.

Introduction

Searching for, and detecting, visual targets in our environment is a ubiquitous natural task that our visual system performs exceptionally well. A key feature of behavioral detection performance is that the texture similarity between the target and the background in which it is embedded profoundly affects target detectability. The more similar are the target and the background, the harder it is to detect the target (Campbell & Kulikowski 1966, Foley 1994, Sebastian et al 2017, Stromeyer & Julesz 1972, Watson & Solomon 1997, Wilson et al 1983). This phenomenon, which is termed "similarity masking", is the foundation of camouflage.

An example of similarity masking is illustrated in Figure 1. Detecting a low contrast oriented visual target is easy on a uniform gray background (Fig. 1A). Detectability decreases when the target has a similar orientation as the background (Fig. 1B). The neural basis of similarity masking is not well understood. The main goal of the current study was to test the hypothesis that neural interactions between the representations of the target and background in the primary visual cortex (V1) contribute to the perceptual effect of similarity masking.

The responses of visual neurons to a target can be strongly modulated by the context in which the
stimulus is presented. Such contextual modulations have powerful, complex and diverse effects in the
visual cortex (Allman et al 1985, Angelucci et al 2017, Angelucci & Bressloff 2006, Bai et al 2021,
Cavanaugh et al 2002b, Henry et al 2020, Michel et al 2018, Sceniak et al 1999, Shushruth et al 2012).
Most of these effects reflect sublinear interactions between the target and the background, suggesting
that they could potentially contribute to behavioral masking effects. If nonlinear computations in V1
contribute to similarity masking, we would predict that the signals evoked by a target will be maximally
reduced by a background that is similar to the target.

We tested this hypothesis by measuring V1 population responses in macaque monkeys while they
performed a visual detection task under masking conditions (Fig. 1C-E). Because the nature of
contextual modulations in V1 is complex, a second goal of our study was to quantitatively characterize
the spatiotemporal dynamics of V1 population responses to different combinations of targets and
backgrounds.

As a first step, we characterized the behavioral effects of similarity masking in two macaque monkeys,
demonstrating clear effects of similarity on target detectability and reaction times. These results confirm
that macaque monkeys are a good animal model for human similarity masking.

Second, we used voltage-sensitive dye imaging (VSDI; (Grinvald & Hildesheim 2004, Seidemann et al
2002, Shoham et al 1999)) to measure V1 population responses at two scales: the scale of the
retinotopic map and the scale of orientation columns, while the monkeys performed the similarity
masking detection task. To study the effect of similarity masking on the neural detection sensitivity, we
constructed a task-specific decoder at each scale. Each decoder first pools the responses using a scale-
dependent spatial template and then combines these responses over time to form a decision variable.
The distributions of the decision variable in target-present vs. target-absent trials are used to compute
neural sensitivity that can be compared to behavioral sensitivity (Seidemann & Geisler 2018).

We found that V1 population responses to the target and background display two distinct phases. An
initial transient phase that starts at response onset, and a second phase that lasts until stimulus offset or
the animal's response. Surprisingly, the first phase displays a paradoxical effect; during this phase the
target evoked response is strongest when the target and background are similar and is therefore anti-
correlated with behavior. This effect reverses in the second phase so that in this phase the target-
evoked response is reduced with increased target-background similarity. V1 responses during this
second phase are therefore consistent with behavior.

We also observed complex spatiotemporal dynamics of the population response to the target and
background stimuli, including a repulsion of V1 columnar-scale representation of target orientation in
the direction away from the background orientation.

Finally, we show that a simple dynamic population gain control model can qualitatively account for our
physiological and behavioral results, and that the estimated properties of the gain-control mechanism
are consistent with a principled computational approach to feature encoding and decoding. Overall, our
results are consistent with the hypothesis that contextual interactions between the representations of
the target and background in V1 are likely to contribute to the perceptual phenomena of similarity
masking.

Results

Behavioral effect of target-background similarity masking

To study the neural basis of visual similarity masking, we trained two monkeys (macaca mulatta) to perform a visual detection task in which a small horizontal target appeared on a larger background at a known location in half of the trials (Fig. 1E). The monkey indicated target absence by maintaining fixation and target presence by making a saccadic eye movement to the target location as soon as it was detected. Within a block of trials, the contrast of the target and the background were fixed, while the orientation of the background varied randomly from trial to trial, allowing us to test for the effect of target-background orientation similarity on behavioral and neural detection sensitivities.

We tested the behavioral effect of similarity masking over five combinations of target and background contrasts (Fig. 2A). For each combination, we measured the behavioral sensitivity as a function of background orientation. Performance with no background (uniform gray screen) served as a baseline (Fig. 2A, dashed horizontal lines). Performance as a function of background orientation was fitted with an inverted Gaussian function. At all five target and background contrast combinations, detection sensitivity was lowest when background and target orientations matched, confirming the expected similarity masking effect from human subjects. These results demonstrate that macaque monkeys are a good animal model for studying the neural basis of human similarity masking.

The supplementary information includes a perceptual demonstration of similarity masking for a wide range of target amplitudes, orientations, and spatial-frequencies (Supplementary Material: "Orientation Masking Demo"). This demonstration can give the reader an intuitive sense of the masking effects studied here.

We also examined the effect of target-background orientation similarity on the monkeys' reaction times (Fig. 2C). We find two distinct effects of orientation similarity on reaction times. At higher target and background contrasts, reaction times are maximal when the background and target have the same orientation (when detectability is lowest and the task is hardest) and monotonically decrease as target-background similarity decreases (detectability increases and the task becomes easier). Surprisingly, at lower target and background contrasts, reaction times are low when the background matches target orientation, then increases as the background-target orientation difference increases, and then drops again when the background approaches the orthogonal orientation to the target. Thus, under these conditions, we see an interesting decoupling between difficulty and reaction time, so that reaction times can be shortest in the harder conditions. This surprising effect is present in both monkeys. Some of the complex neural dynamics described below could explain this interesting effect (see Discussion).

Our next goal was to test the hypothesis that contextual interactions between the representations of the target and background in V1 contribute to the observed behavioral similarity masking results.

Neural population responses to target and background stimuli in macaque V1

While the monkeys performed the similarity masking detection task, we used VSDI to measure V1 population responses to the target and the background. In each cranial window, we first used a fast and efficient VSDI protocol to obtain a detailed retinotopic map (Yang et al 2007). We then positioned the target so that its neural representation fell at the center of our imaging area.

The target elicits V1 population activity at two fundamental spatial scales. At the large retinotopic scale,
the target evokes an activity envelope that spreads over several mm^2 and is well fitted by a two-
dimensional (2D) Gaussian (Fig. 3B, top row; (Chen et al 2006, Chen et al 2012, Sit et al 2009)). Our 8x8
mm^2 imaging area allows us to capture this entire target-responsive region. Because the background is
much larger than the target and is centered at the same location in the visual field, it produces a
relatively uniform response within the imaging area (Fig. 3B, second row). Similarly, the target-plus-
background stimulus elicits activity within the entire imaged area, with a relatively elevated activity at
the retinotopic region corresponding to the target location (Fig. 3B, 3rd row). However, the target-
evoked response in the presence of the background (response to target plus background minus
response to the background alone) appears significantly weaker than the response to the target alone
(Fig. 3B, 1st vs. 4th row). This reduced target-evoked response in the presence of the background could
contribute to the perceptual masking effect of the background. Our goal here was to determine how this
sublinear interaction between the response to the target and background depends on target-
background similarity in orientation.

In addition to the retinotopic-scale response envelope, fine scale response modulations at a scale of
individual orientation columns (diameter of ~0.3 mm) reflect the orientation of the target and
background. These columnar scale modulations have a relatively small amplitude and therefore appear
as small ripples riding on top of the larger retinotopic response envelope. The relatively smaller VSDI
responses at the columnar scale is due to a mixture of robust non-orientation selective V1 population
responses in V1 as well as optical and biological blurring (Chen et al 2012). We can selectively access the
columnar scale signals by spatially filtering the responses at the scale of the orientation columns.
Despite their small relative amplitude, these columnar-scale signals provide high quality single-trial
orientation decoding (Benvenuti et al 2018).

138 Retinotopic-scale effect of target-background similarity masking

To study the effect of similarity masking on V1 responses at the retinotopic scale, we used an optimal
linear decoder of V1 population responses (Chen et al 2006, Chen et al 2008) that allows us to assess the
neural detection sensitivity of V1 population responses (i.e., how well one can detect the target from
single-trial V1 population responses) (Fig. 3C-D). The retinotopic decoder takes into account the location
and shape of the envelope of the target-evoked response (Fig. 3D), as well as the structure of the noise
covariance matrix (Fig. 3C; (Chen et al 2006)).

Figure 4 summarizes the dynamics of the retinotopic template output in response to the V1 signals
across all of our experiments for two combinations of target and background contrasts (see Table 1 and
Fig. 4 – figure supplement 1 for the full set of tested target/background combinations). When presented
on a uniform gray background, the target-related retinotopic signal begins to rise ~40 ms after target
onset, reaches its peak ~100 ms after stimulus onset, and remains high for the next 100 ms (Fig. 4B,H,
black curve). However, when the same target is added to the background, the target-related retinotopic
signals display a wide range of responses that depend on background orientation (Fig. 4B,H, colored
curves).

Our main interest here is in the target-evoked response in the presence of the background (Fig. 4C,I),
which can be extracted by subtracting the response to the background alone (Fig. 4A,G) from the
response to the target plus background (Fig. 4B,H). If V1 contextual interactions at the retinotopic scale
contribute to the behavioral effect of similarity masking, we would expect that target-evoked responses

would be weakest with high target-background similarity (similar target and background orientations)
and strongest with low target-background similarity (orthogonal target and background orientations).

Surprisingly, we find that the target-evoked response in V1 displays two distinct phases, with the early
phase showing a paradoxical neural dependence on target-background orientation similarity that is anti-
correlated with the behavioral masking effect, and with a later phase that is consistent with the
behavioral masking effect. Specifically, in the early phase which starts at response onset, the target-
evoked response is highest when the background matches the target orientation even though
behaviorally this is the condition in which detection performance is the worst. However, after this initial
phase, the high-similarity target-evoked response starts to drop, while the low-similarity target-evoked
response continues to build up, so that in the later stages the target-evoked response is strongest on the
dissimilar background and weakest on the similar background, consistent with the behavioral effect of
similarity.

To quantify the relation between the effects of target-background orientation similarity on V1
population responses and behavior, we computed the correlation between the effect of orientation
similarity on behavior (Fig. 4F,L) and its effect on the target-evoked neural responses in individual 10 ms
frames (Fig. 4E,K). This analysis reveals a robust paradoxical negative correlation between the early
neural V1 response and behavior, weak positive correlation between the late neural V1 response and
behavior, and no correlation between behavior and the integrated neural response. This result was
obtained from averaging the response across all trials irrespective of whether the monkey made the
correct decision.

To examined whether decision- and/or attention-related signals have a major contribution to the
observed biphasic dynamics, we repeated the analysis on only the hits and correct rejection trials (Fig. 4
– figure supplement 2B-C). Our results are qualitatively the same for the subset of correct trials,
indicating that decision- and/or attention-related signals are unlikely to play a major role in the
observed dynamics.

Because the target and background are defined by their orientation, the correspondence between the
neural signals in V1 and behavior may be better captured by V1 responses at the columnar scale. Our
next step was therefore to examine the dynamics of the columnar-scale target-evoked responses in V1.

Neural effects of target-background similarity masking at the scale of orientation columns

To study the effect of similarity masking on V1 responses at the columnar scale, we developed a linear
columnar decoder of the VSDI signals (Fig. 3E). The columnar decoder takes into account the location of
the orientation columns within the retinotopic envelope of the target-evoked response. Because the
target is horizontal, the output of the columnar template is expected to be positive for the horizontal
target and background stimuli and negative for the vertical background stimulus (since the horizontal
and vertical columnar maps are anti-correlated).

As with the output of the retinotopic-scale template, the output of the columnar-scale template displays
two distinct phases. Figure 5 shows the time course of the columnar template signals to background
alone (Fig. 5A,G), the target plus background (Fig. 5B,H), and target-evoked response in the presence of
the background (Fig. 5C,I). In the early phase, the target-evoked response is highest when the
background and target have similar orientations, producing a paradoxical neural response that is anti-

correlated with the behaviorial masking effect (Fig. 5E,K). In the second phase the trend reverses and
the target-evoked response is strongest on the dissimilar background and weakest on the similar
background, consistent with the behavioral effect of similarity (Fig. 5E,K). Similar results were obtained
with other target and background contrast combinations (Fig. 5 – figure supplement 1).

Again, to examined whether decision- and/or attention-related signals have a major contribution to the
observed biphasic dynamics at the columnar scale, we examined the behavioral correlations with hits
and correct rejection trials only. We found only minor differences in the target-evoked response and
behavioral correlations (Fig. 4 – figure supplement 2D-E), indicating that the observed biphasic dynamics
at the columnar scale are unlikely to have a major top-down contribution.

Because the first phase of the response is shorter than the second phase, when V1 response is
integrated over both phases, the overall response is positively correlated with the behavioral masking
effect (Fig. 5D,J, figure supplement 1C). Therefore, our results suggest that the neural masking effect at
the columnar scale in V1 could play a major role in the behavioral similarity masking effects.

Dynamics of columnar-scale orientation population trajectories

Our decoding analysis focuses on the columnar-scale orientation signals along the 0°-90° axis and
reveals complex columnar-scale dynamic interactions between the target-evoked response and the
response evoked by the background (Fig. 5). To examine these dynamics in more detail, we performed
two types of population-vector analyses (Fig. 6). We began by assigning each pixel within the retinotopic
footprint of the target-evoked response to one of 12 equally spaced preferred orientations, creating 12
orientation selective clusters of pixels (Fig. 6B-C). We then computed for each stimulus the response in
each orientation selective cluster in each frame and displayed, in the first analysis, the population
orientation tuning curve as a function of time (Fig. 6D), and in the second analysis the population vector
dynamic trajectory in the polar space spanned by the 12 orientations (Fig. 6E); i.e., the orientation θ
and magnitude R_θ of the peak of the population response over time.

The first population vector analysis reveals that V1 responses to the target or background alone are
consistent with the stimulus orientation. In background only trials, shortly after stimulus onset the peak
of the population tuning curve closely matches background orientation (Fig. 7A, F, top row, red arrow
and horizontal line). Similar results were obtained in the target only trials, where the peak of the
population response tuning curve matches target orientation (Fig. 7A, F, 4[th] row, green arrow and
horizontal line).

Target plus background responses display complex spatiotemporal dynamics. To examine the dynamics
of the target-evoked response in the presence of the background, we subtracted the background only
response from the target plus background response (Fig. 7A, F, 3[rd] row). The results reveal complex
target-background interactions which could lead to a population tuning peak (white curve) that
significantly deviates from the target orientation. For example, in some conditions, we observe an
orientation tuning peak that is repelled from target orientation in the direction away from background
orientation. An interesting goal of future studies would be to examine potential perceptual correlates of
these interactions.

In the second population vector analysis, we plotted the response trajectories for each stimulus using
the vector representation in polar coordinates (Fig. 7B-E, G-J). After stimulus onset, the population

vector for background only moves in the direction corresponding to the background orientation (Fig.
7B,G) and for target only moves in direction corresponding to the target (Fig. 7E,J).

The trajectories in the target plus background conditions are more complex. For example, when
background orientation is at +/- 45 deg to the target, the population response is initially dominated by
the background, but then in mid-flight, the population response changes direction and turns toward the
direction of the target orientation.

Such complex interactions can be used to constrain models of V1 population response.

244 Dynamic gain control model qualitatively captures similarity masking effects in V1

Our next goal was to determine whether the observed interactions between the background- and
target-evoked responses can be qualitatively captured by a gain control model. In this model,
orientation columnar response was tuned to one of 12 equally spaced orientations. The responses of
each orientation column were specified by the simple normalization model summarized in Figure 8A.
Specifically, the spatiotemporal input stimulus generates an excitation signal and a normalization signal
that are both linear with the input root mean square (rms) contrast. The normalization signal is then
combined with a normalization constant to obtain the normalization factor. The normalized response is
obtained by dividing the excitation signal by the normalization factor. The final response is then
obtained by applying a response exponent p, which is similar to applying a spiking nonlinearity.
Importantly, the excitation and normalization signals can differ in their spatial extent, orientation tuning
width, and temporal impulse response (see Methods for model parameters).

We find that this simple model can qualitatively captures our key results. First, in response to
background alone (Fig. 8B), the modeled population vector peaked at ~100 ms after stimulus onset and
then dropped to a lower amplitude, as in our data (Fig. 5A,G). This reduction in response amplitude was
due to normalization signal that was delayed relative to the excitation signal. Second, as in the real data,
response to the target plus background is less than the sum of the responses to each component
separately. Third, as in our physiological results, the target-evoked response in the presence of the
background is biphasic, having a brief early component in which the response is enhanced by target-
background similarity, and a longer-lasting late component in which the response is suppressed by
target-background similarity (Fig. 8D). This leads to an early phase in which the response is
anticorrelated with the behavioral effect of similarity masking, and a late phase and an integrated
response that are positively correlated with the behavioral effect of similarity masking (Fig. 8E,F).

Finally, this simple model can also display the curved trajectories of the population vector in response to
the target plus background (compare Fig. 8H to Fig. 7C,H).

Additional results show that the model is relatively insensitive to the spatial extent of the normalization
signal (Fig. 8 – figure supplement 1). The model predicts very similar temporal dynamics with the spatial
extent of the background mask as small as twice the size of the target (Fig. 8 – figure supplementary 2).
When the background and target are the same size, the model predicts that sufficiently high contrast
background will also drive the same biphasic temporal dynamics (fig. 8 – figure supplement 3, rows 7
and 8). To account for the orientation-dependent neural and behavioral masking effects, the model
requires an orientation tuned normalization (Fig. 8 – figure supplement 4).

Overall, our results suggest that a simple model with delayed and orientation tuned divisive gain control
can qualitatively capture the complex spatiotemporal dynamics of V1 population responses to localized
oriented targets added to oriented backgrounds.

Discussion

To test the hypothesis that nonlinear computations in V1 contribute to the perceptual effect of similarity
masking, we used voltage-sensitive dye imaging (VSDI) to measure neural population responses from V1
in two macaque monkeys while they performed a visual detection task in which a small oriented target
was detected in the presence of a larger background of varied orientations. Like human observers, the
monkeys were strongly affected by the orientation similarity of the target and the background (Fig. 2).
Their detection threshold increased with increased target-background orientation similarity, while their
reaction times showed complex, and in some cases non-monotonic, dependency on target-background
orientation similarity.

To quantify the neural effects of similarity masking, we measured neural sensitivity to the target at two
fundamental spatial scales of V1 topographic representations. The large scale of the retinotopic map
and the finer scale of the columnar orientation map. We discovered that at both scales, V1 population
responses to the target and background display two distinct phases (Fig. 4B, H, Fig. 5B, H). An initial
transient phase in which target-evoked V1 response is strongest when the target and background have
similar orientations. At this early phase, V1 responses are therefore paradoxically anti-correlated with
the behavioral effect of similarity masking (Fig. 4E, K, Fig. 5E, K). In the second phase, the masking effect
reverses, and the target-evoked response is maximally reduced when the target and background are
similar. In this second sustained phase V1 population responses are therefore consistent with the
behavioral similarity masking effect. To explore the possibility that these biphasic dynamics reflect
contributions from decision- and/or attention-related top-down signals rather than from low-level
nonlinear encoding mechanisms in V1, we re-examined our results while excluding error trials (Fig. 4 –
figure supplement 2). We found that the biphasic dynamics hold even for the subset of correct trials,
reducing the likelihood that decision/attention-related signals play a major role in explaining our results.

The positive correlation between the neural and behavioral masking effects occurred earlier (Fig. 5E, K
vs. Fig. 4E, K) and was more robust at the columnar scale than at the retinotopic scale (Fig. 5D, J vs. Fig.
4D, J; see also Fig. 4 – figure supplement 1C, and Fig. 5 – figure supplement 1C). In addition, while the
temporally integrated columnar response was positively correlated with behavior across all target and
background contrasts tested (Fig. 5E,K, figure supplement 1D), the integrated retinotopic responses
were uncorrelated, or in some cases anticorrelated, with behavior (Fig. 4E,K, figure supplement 1D).
These results suggest that behavioral performance in our task is dominated by columnar scale V1 signals
in the second phase of the response. To the best of our knowledge, this is the first demonstration of
such decoupling between V1 responses at the retinotopic and columnar scales, and the first
demonstration that columnar scale signals are a better predictor of behavioral performance in a
detection task.

Due to the challenges of setting up these experiments, we were unable to collect all target/background
contrast combinations from both monkeys. However, in the common conditions, the results appear
similar in the two animals, and the key results seem to be robust to the contrast combination in the

animal where a wider range of contrast combinations was tested (Fig. 4 – figure supplement 1, and Fig. 5
– figure supplement 1).

We find that when the target and background have similar orientations, columnar-scale information
about the target is restricted to the first phase of the response and then largely disappears during the
second phase of the response. These physiological results could be related to the surprising mismatch
between task difficulty and reaction times (Fig. 2C). Rather than having reaction times that
monotonically increase with task difficulty, in our masking detection task, reaction times can be shortest
when target and background orientations match, even though it is hardest to detect the target under
these conditions. The short reaction time to this stimulus may be the consequence of the target
information being best represented in the early phase of the response.

The nature of contextual modulations in V1 is quite complex (Angelucci et al 2017, Angelucci & Bressloff
2006, Bai et al 2021, Cavanaugh et al 2002b, Henry et al 2020, Michel et al 2018, Polat et al 1998,
Sceniak et al 1999, Shushruth et al 2012). A second goal of our study was to quantitatively characterize
the spatiotemporal dynamics of columnar-scale V1 population responses to targets and backgrounds of
different orientations and contrasts. Using a dynamic population vector analysis, we find that in the
presence of an oriented background, the peak of the population orientation tuning to the target can
deviate significantly from target orientation. For example, in some conditions, we observe a population
orientation tuning peak that is repelled away from target orientation in the direction opposite to
background orientation (Fig. 7E). These orientation-dependent interactions could contribute to non-
veridical perceptual representations of orientation such as in the well-known tilt illusion effect (Clifford
2014, Schwartz et al 2007, Wenderoth & Johnstone 1987). An important goal for future studies would
be to test for this possibility.

Using the population vector analysis, we find that columnar scale V1 representations are initially
dominated by the orientation of the background. The target orientation then appears in the second
phase of the response, which leads to curved population vector trajectories (Fig. 7C, H). Identifying
possible perceptual consequences of such dynamic and complex trajectories, and understanding the
neural circuit mechanisms that give rise to such responses, are two important goals for future work.

Nonlinear response properties in V1 are commonly modeled as a consequence of a divisive gain control
mechanism (Albrecht & Geisler 1991, Carandini & Heeger 1994, Heeger 1991, Heeger 1992, Sit et al
2009). As a first step toward understanding the mechanisms that could give rise to the observed V1
responses, we tested whether a simple dynamic gain control model could account for our findings (Fig.
8). We find that a simple gain control model can qualitatively account for our results, but that in order to
do so, the model has to display two important properties. First, to account for the biphasic nature of V1
response, the divisive normalization signals have to be delayed relative to the excitatory signal. Second,
in order to account for the reduced neural sensitivity with target-background similarity in the second
phase of the response, the divisive normalization signal has to be orientation selective (Fig. 8 – figure
supplement 4). Because in primates and carnivores, robust orientation selectivity first emerges in V1
(Hubel & Wiesel 1959, Hubel & Wiesel 1968), these results suggest that a significant portion of the
nonlinear interactions observed in the current study originate in V1 rather than being inherited from the
ascending inputs that V1 receives from the LGN. While our experimental and computational results
point to a delayed gain control signal that operates at the level of V1, they do not directly speak to the
circuit and biophysical mechanisms that contribute to the implementation of this gain control in V1.

Multiple candidate mechanisms for implementing gain control in V1 have been proposed (Angelucci et
al 2017, Angelucci & Bressloff 2006, Ozeki et al 2009, Rubin et al 2015, Tsodyks et al 1997). Our results
provide new and powerful constraints for such mechanistic models.

A key difference between our study and previous center-surround studies (e.g., (Cavanaugh et al 2002a,
Cavanaugh et al 2002b, Henry et al 2020, Shushruth et al 2012)) is the stimuli that we used. First, in our
experiments, the target and the mask were additive, while in most previous center-surround studies the
target occludes the background. Such studies therefore restrict the mask to the surround while our
study allows target-mask interactions at the center. Second, most previous center-surround studies
have a sharp-edged target/surround border, while in our experiments no sharp edges were present.
Unpublished results from our lab suggest that such sharp edges have a large impact on V1 population
responses. Third, our stimuli were flashed for a short interval of 250 ms corresponding to a typical
duration of a fixation in natural vision, while most previous center-surround studies used either longer-
duration drifting stimuli or very short-duration random-order stimuli for reverse-correlation analysis.

Because our targets are added to the background rather than occluding it, it is likely that a significant
portion of the behavioral and neural masking effects that we observe come from target-mask
interactions at the target location rather than from the effect of the mask in the surround. Several lines
of evidence support this possibility. First, in human subjects, perceptual similarity masking effects can be
almost entirely accounted for by target-mask interactions at the target location and are recapitulated
when the mask has the same size and location as the target (Sebastian et al 2017). There is a reduction
in masking when the background is windowed to the target envelope, but this effect is due to removing
background within the target envelope (Sebastian et al 2017). Second, in our computational model (Fig.
8), the effects of mask orientation on the dynamics of the response are qualitatively similar if the mask
is restricted to the size and location of the target and mask contrast is increased (Figs. 8 – figure
supplement 3). Third, in our model, the results are qualitatively the same when the spatial pooling
region for the normalization signal is the same as that for the excitation signal (Fig. 8 – figure
supplement 1). These considerations suggest that center-surround interactions may not be necessary
for neural and behavioral masking effects with additive targets.

Finally, we note that the tuned similarity normalization that explains the neural and behavioral
similarity-masking effects reported here is consistent with a principled encoding strategy for feature
detection under natural conditions. When viewing a static scene under natural conditions, human and
non-human primates make 3-4 saccadic eye movements per second, with fixations between saccades of
200-300 ms. Given the typical size of the saccades, most visual receptive fields are stimulated during
each fixation by a largely statistically independent random sample of natural image (Frazor & Geisler
2006). Analysis of the responses of linear receptive fields to random samples of natural image shows
that the standard deviation of the response increases in proportion to the product of the luminance, the
contrast and the similarity of the natural background to the receptive field (Sebastian et al 2017). Thus,
divisive normalization by the product of luminance, contrast and similarity causes the standard deviation
of the responses across natural images to be much more constant (i.e., nearly independent of the
luminance, contrast and similarity of the background within the receptive field). This more constant
standard deviation makes it possible, with relatively simple decoders, to reach near optimal feature-
detection performance, under the high levels of stimulus uncertainty that occur under natural
conditions (Schwartz & Simoncelli 2001, Sebastian et al 2017).

Acknowledgments

We thank We thank members of Seidemann and Geisler laboratories for their assistances with this project. This work was supported by NIH grants EY-016454 to E.S., EY-024662 to W.S.G. and E.S., BRAIN U01-NS099720 to E.S. and W.S.G., and DARPA-NESD0-N66001-17-C-4012 to E.S.

Figure 1 – Target-background similarity masking and behavioral task. (A) Low contrast orientated
targets can be easily detected on uniform background. (B) Similarity masking is induced by an orientated
background. The same additive target from (A) becomes hard to detect when target orientation
matches background orientation. (See also perceptual demonstration in the Supplementary
Information.) (C-E) Orientation masking was assessed in two awake behaving macaque monkeys
performing a target detection task. Monkey commence the task by fixating at the small bright square
(C). A few moments later, a 4° raise-cosine-masked background grating was flashed at ~3° eccentricity
for target detection (D). The horizontal white bar represents one degree of visual angle. In 50% of the
trials, a small additive horizontal Gabor target was also added to the background (E). The monkey
indicated the presence of the target by making a saccade to the target location, and indicated target
absent by maintaining gaze at the fixation point. The Gabor target was always the same – a cosine
centered, horizontal Gabor at 4cpd on 0.33° FWHM envelope. The background grating was also cosine-
centered at 4cpd such that the background completely aligned with the target when they were the same
orientation (as in B). Orientation of the grating ranged from 0° to 90° with respect to the Gabor target
and was randomized between trials. Bg – background; TBg – target plus background.

Figure 2 – Behavioral effect of target-background similarity masking. (A) Target detection performance in monkeys was affected by the orientation of the background grating over a variety of target contrast (T## = ##% contrast target) and background contrast levels (Bg## = ##% contrast background). Signal detection measure d-prime (d') of the target is plotted for uniform background (dotted lines), and for each background orientation (markers). In most cases there was a general performance reduction from uniform background to a grating background. Additionally, performance was further reduced when the background orientation was more aligned to the target (0°). A fitted Gaussian (solid line) illustrates the performance change due to orientation masking. d' was calculated from the hit rate (correctly reporting target present) and the false alarm rate (reporting target present when it was absent). The relationship between d' and optimum performance level in percent correct is plotted in (B). (C) Reaction time – calculated from stimulus onset to saccade initiation for Hit trials – is plotted for uniform background (dotted lines) and for each background orientation (solid lines). Error bars indicate the standard error of the mean. Data were pooled within each monkey across experiments. Each experiment contains a single combination of target and background contrast levels, with uniform background and orientated background trials assessed in separate blocks.

Figure 3 – VSD response and decoder schematic. (A) A cranial chamber and a transparent artificial dura
provided chronic imaging access to V1 of the monkey. V2 is completely hidden in the lunate sulcus
based on the retinotopic map taken in a separate imaging session (Fig. 3 – figure supplement 1). Imaging
ROI and target response at 2 SD of the fitted Gaussian from the example recording in B-E are illustrated.
(B) Example recording of the voltage-sensitive dye (VSD) response in an 8x8mm imaging region-of-
interest (ROI). Response to the Gabor target on uniform background could be easily identified in VSD
(top row). The visual span of the background grating extended beyond the coverage of the imaging
window, evoking an encompassing response over the entire ROI (2nd row). When the background was
presented with the additive target, the response to the target was diminished (3rd and 4th rows). Each
VSD response map were averaged across 50 trials, over 5 frames captured at 100Hz. T only – target only;
Bg – background only; TBg – target & background; ΔTBg – target & background minor background only.
(C) Target response was extracted at the retinotopic scale by estimating its response profile with a two-
dimensional Gaussian. The profile was estimated from response from a separate recording block on
each experiment day. To optimized signal-to-noise, in this recording block, the target was flashed
repeatedly at 5Hz while the monkey maintained fixation. The effect of spatially correlated VSD noise was
minimized by estimating a whitening kernel from trials without stimulus presentation (see Methods). (D)
Target response was extract at the columnar scale by estimating the orientation map within the imaging
area. This was constructed from full-field gratings flashed at 5Hz in a separate recording block on each
experiment day (see Methods). The columnar map in the 0°-90° axis was extracted and windowed down
to the retinotopic profile to identify the columnar scale response of the target.

Figure 4 – Retinotopic template dynamics and correlation to behavior. (A-F) Average response and dynamics from recordings with 12% target contrast (T12) and 7% background contrast (Bg7). (A) Response time course from stimulus onset (t=0ms) for background only trials. Background orientation are identified by color. Backgrounds with the same clockwise and anticlockwise orientation disparity from the target were pooled. (B) Response time course with the same additive target on different oriented background. Response of the target on uniform gray background is illustrated in black. (C) The target-evoked response time course was obtained by subtracting background only response (A) from response to target & background (B). Target evoked response was initially strongest for backgrounds close to 0° (red), then inverted around t=100ms such that response became the stronger for background closer to 90° (blue). (D) Response was averaged over 50-200ms and fitted with a Gaussian (gray) to illustrate the change in response magnitude with respect to background orientation. The neural-behavioral correlation (*r*) of the response against behavioral response (F) is printed with *p* significance value. Here, response to clockwise and anti-clockwise background orientations are plotted separately. Size of markers indicate the number of trials tested for each orientation. Black line indicates the response of the target only trials integrated over the same window. (E) The animals' behavior performance was anti-correlated with the initial phase of the retinotopic response, and was more

aligned in the latter phase. Correlation coefficient was calculated across background orientations
between each frame of the retinotopic response in (C) against the overall behavior performance in (F).
Red dots indicate frames reaching statistical significance ($p < 0.05$, t-test for correlation coefficient, see
Methods). Behavioral correlation crosses from negative to positive at t=130ms. (F) Behavior
performance in d' was calculated as described in Figure 2. Size of markers indicate the number of trials
tested for each orientation. Data was pooled across 8 experiments from both monkeys (see Table 1). (G-
L) Same as (A-F) for recordings with 24% target contrast (T24) and 12% background contrast (Bg12).
Similar trends were observed. Behavioral correlation crosses from negative to positive at t=96ms in (K).

Figure 5 – Columnar template dynamics and correlation to behavior. Same format as Figure 4 with the response examined at the columnar scale. The biphasic response time course observed in the retinotopic scale was more pronounced at the columnar scale. (A-F) Averaged response and dynamics from recordings with 12% target contrast (T12) and 7% background contrast (Bg7). (G-L) Averaged response and dynamics from recordings with 24% target contrast (T24) and 12% background contrast (Bg12). Here positive response represents relatively stronger activation of the neurons tuned to the target orientation (0°), and negative response represent stronger activation for neurons tuned to the orthogonal orientation (90°). Data pooling and counts are the same as reported in Figure 4. Behavioral correlation crosses from negative to positive at t=99ms in (E), and t=66ms in (K).

Figure 6 – Columnar orientation estimation by populations tuning. (A) The orientation map obtained for each experiment as described in Figure 3 was windowed to the retinotopic profile of the target. (B) Each pixel was assigned to one of 12 equally spaced orientation selective cluster maps by its preferred orientation. (C) The orientation selective decomposition of VSD response. To a grating stimulus oriented at 0°, the population tuning curve peaks at 0° (solid curve); likewise, the population peak would shift to 45° for a 45° grating (dotted curve). Note that this population response only represents the relative difference in preferred orientation (balanced positive and negative values); the overall neural response offset (retinotopic response) is not captured by this approach. (D) Example of a full population response time course from stimulus onset (t=0ms). (E) Population response can be summed to a complex vector representing the overall population tuning orientation and magnitude.

Figure 7 – Dynamics of orientation population response. (A) Population tuning time course for trials
with 12% contrast targets (T12) and 7% contrast backgrounds (Bg7). Averaged response time courses
are presented as a heatmap with the y-axis representing the preferred orientation (from Fig. 6A-C) and
the x-axis time (see key on bottom right) to illustrate the change in the tuning over time. *Row 1:*
Heatmaps for background only trials exhibit clear population tuning in the orientation of the background
grating (red horizontal line). *Row 2:* Heatmaps for background with additive target showing population
response dominated by the background orientation rather than the target orientation (0°). *Row 3:* The
target evoked response is obtained by subtracting the background only response Row 1 from the target
& background response in Row 2. Masking of the target evoked response was strong for backgrounds
oriented near the target orientation (0°). With the background orthogonal to the target, population
tuning in the target orientation can be identified. White line identifies the orientation of the population
vector (peak tuning) wherever the normalized amplitude of the vector average was great than 0.2 (see
Methods). Depending on background orientation, peak tuning appears to be offset from the orientation
of the target (e.g. at Bg -45°). *Row 4:* Heatmap for the target only trials demonstrated clear population
tuning in the target orientation (0°, green horizontal line). (B-E) Averaged response in (A) represented as
a population vector form and illustrated as a continuous trajectory for each background orientation
(color coded). (B) Population tuning trajectory for background only trials. The trajectories commenced in

the center of the circle (white dot) and adhered closely to the orientation of the background. Dot on
each trajectory indicates the position of the population tuning vector at 100 ms. (C) Population tuning
trajectory for background with additive target illustrating the biphasic response of this combined
stimulus. In the early phase, the heading of the trajectory was a mixture of the background and target
(0°) orientations, dominated more by the background. In the late phase, the trajectory made a sharp
turn (t≈100ms) such that trajectories appeared to head towards a convergent point on the positive x-
axis. (D) The trajectory for the target evoked response, calculated by subtracting the background only
response (B) from the corresponding background & target (C). The target evoked response was weak
and noisy, but was heading in the general direction of the target orientation (0°). (E) The population
tuning trajectory for the target only trials illustrating clear tuning in the target orientation (0°). (F-J)
Same as (A-E) for trials with 24% contrast targets (T24) and 12% contrast backgrounds (Bg12). Data was
pooled and averaged across both monkeys.

Figure 8 – Delayed normalization model qualitatively captures key orientation masking response features. (A) Schematic of the normalization model showing the visual input being processed by separate excitatory and normalization signal pathways. The normalization pathway in particular was modelled with a slightly delayed temporal kinetics and wider orientation tuning curves. The excitatory signal undergoes divisive normalization prior to neural output. (B-D) Modelled columnar response output with target contrast of 24% and background contrast of 12%. Modelled response normalized to the target only response averaged over 50-200ms as plotted in (E). (B) Modelled response of oriented backgrounds as in Figure 5A. (C) Modelled response of background with additive target as in Figure 5B, and the modelled response of the small Gabor target in black. (D) The target evoked response by subtracting the background only response (B) from (C), matching the biphasic observation in Figure 5C. (E) Model response integrated over 50-200ms in the same format as Fig. 5D. (F) Correlation of modelled behavioral performance (Gaussian fit in Fig. 5L) against each time frame of the modelled response, illustrating the early phase where the response was negatively correlated to behavioral choice, and the late phase with positive correlations. (G-J) Modelled time course of the population tuning vector. (G) Modelled populating tuning trajectory of the background only stimuli (color coded) as in Figure 7B. (H) Modelled populating tuning trajectory of the background with additive target illustrating the turn towards a convergent point on the x-axis as in Figure 7C. (I) Modelled target evoked response trajectory from subtracting (F) from (G). (J) Modelled populating tuning trajectory of the target only trials as in Figure 7E.

	Target Contrast	Background Contrast	# Experiments	# Total Trials	# Hits	# Misses	# Correct Rejects (CR)	# False Alarms (FA)
Monkey H	T12%	---	5	52	26	0	25	1
	T12%	Bg7%	3	593	214	80	253	46
	T12%	Bg12%	2	397	125	74	121	77
Monkey T	T12%	---	11	490	185	59	228	18
	T12%	Bg7%	5	1660	644	189	740	87
	T12%	Bg12%	6	1748	704	169	683	192
	T24%	---	16	546	272	0	267	7
	T24%	Bg7%	5	1012	501	5	450	56
	T24%	Bg12%	6	1320	654	12	599	55
	T24%	Bg24%	4	388	204	2	155	27

561   **Table 1 – Experiment summary.** Experiment counts and the total number of trials included in the
562   analysis presented in Figures 2, 4, 5, 7; Fig. 4 – figure supplement 1 and 2; and Fig. 5 – figure supplement
563   1.

Figure 3 – figure supplement 1 – Retinotopy of imaging chambers and target placement positions. A
separate VSD imaging experiment was conducted for each chamber with special retinotopic scanning
stimuli (see Methods). (A) Photograph of the imaging chamber overlaid with contour lines (green) of
polar angles relative to the horizontal meridian of the visual field. Angular values were anti-clockwise
from the right horizontal meridian. The V1-V2 border (dashed orange line) is at 270°. This border is hard
to estimate and is believed to be 0.7 mm from the lunate in Monkey T's left visual cortex chamber (col
2). For the other two chambers, the V1-V2 border is estimated to be inside the lunate sulcus (dashed red
line). (B) The same imaging areas as in A marked with both the angular (green) and eccentricity (red)
contour lines. Area with good VSD staining on the day of retinotopy is marked in white. Red dots in A
and B indicate where the target was placed. The position of the target in cols 2-3 was moved between
experiments to optimize for the imaging outcome as a response to the quality of VSD staining of the day.
Scalebars indicate 4 mm in length.

Figure 4 – figure supplement 1 – Retinotopic template dynamics and correlation to behavior for all combinations of background and target contrast levels. Retinotopic response time course using doubly-whitened retinotopic decoding template illustrated in Figure 3. (A) Response time course from stimulus onset (t=0ms), grouped by different combinations of target and background contrast levels. Background orientation is indicated by color. Backgrounds with the same clockwise and anticlockwise orientation disparity from the target were pooled. Response of the target on uniform gray background is illustrated in black. (B) The target-evoked response time course was obtained by subtracting the background only response (A, top row) from the background-plus-target response (rest of A) from the same experiments. (C) Response was averaged over 50-200ms and fitted with a Gaussian (gray) to illustrate the change in response magnitude with respect to background orientation. Neural-behavioral correlation (*r*) of the integrated response against the behavior performance (Figure 2A pooled across monkeys) are printed with p significance values. Here, response to clockwise and anti-clockwise background orientations are plotted separately. Size of markers indicate the number of trials tested for each orientation. Black line indicates the response of the target only trials integrated over the same window. (D) Correlation coefficient was calculated across background orientations between each frame of the retinotopic response in (B) against the overall behavioral performance for each combination of background and target contrast (Figure 2A pooled across monkeys). Red dots indicate frames reaching statistical

599   significance ($p < 0.05$, t-test for correlation coefficient, see Methods). Data was pooled across
600   experiments from both monkeys (see Table 1).

Figure 4 – figure supplement 2 – Retinotopic and columnar integrated response and correlation to behavior. Response dynamics calculated over all trials as shown in Fig. 4 and Fig. 5 are replotted in odd rows for comparison against the same stimulus conditions using only correct trials (hits and correct rejections only, even rows). (A) Behavioral performance replotted from Fig. 4F and L. Behavioral performance was calculated using all trials (correct and incorrect), and was used for the calculation of correlations when the template response included all trials or just the correct trials. (B,D) Integrated response over 50-200ms from stimulus onset in the same format as Fig. 4D. (C,E) Behavioral correlations in the same format as Fig. 4E. The vertical lines mark the time of the cross-over from negative to positive correlations (the vertical lines are labelled with the time in ms).

Figure 5 – figure supplement 1 – Columnar template dynamics and correlation to behavior for all
combinatory background and target contrast levels. (A-D) Same format as Fig. 4 – figure supplement 1
with the averaged response examined at the columnar scale. The columnar template extracts the
relative strength between neural activity aligned and orthogonal to the target orientation. Positive
response represents relatively stronger activation of the neurons tuned to the target orientation (0°),
and negative response represent stronger activation for neurons tuned to the orthogonal orientation
(90°). Data pooling and counts are the same as reported in Fig. 4 – figure supplement 1.

Figure 8 – figure supplement 1 – Divisive normalization model with different normalization spatial
extents. The spatial extent of the normalization kernel was varied to determine its effect on key model
outputs from Fig. 8. The size of s_n was varied while all other parameters were at the same values as in
Fig. 8. Target contrast was modelled at 24% and background contrast at 12% (as in Fig. 5G-L). (A)
Illustration of the size of the normalization spatial kernel. Fig. 8 was simulated with $s_n = 0.28°$ (row 1). (B-
D) Model response corresponding to Fig. 8B-D. (E) Model response integrated over 50-200ms in the
same format as Fig. 8E. (F) Correlation of modelled behavioral performance (Gaussian fit in Fig. 5L)
against each time frame of the modelled response. (G) Modelled populating tuning trajectory
corresponding to Fig. 8H.

Figure 8 – figure supplement 3 – Divisive normalization model with different background contrasts.
The contrast level of the background grating was explored for its effect on key model outputs from Fig. 8
(row 2). Rows 1-4: The contrast of the background was varied while all other parameters were at the
same values as in Fig. 8. Rows 5-8: Contrast of background was varied with the background set to the
same size as the Gabor target, while all other parameters were at the same values as in Fig. 8. Target
contrast was 24% for all rows (as in Fig 8), and the background contrast for each row is as labeled. (A)
Illustration of the additive Gabor target on the background grating. (B-D) Model response corresponding
to Fig. 8B-D. (E) Model response integrated over 50-200ms in the same format as Fig. 8E. (F) Modelled
populating tuning trajectory corresponding to Fig. 8H.

Figure 8 – figure supplement 4 – Divisive normalization model with different normalization signal

orientation tuning width. The orientation tuning width (σ_n) of the normalization signal was explored for

its effect on key model outputs from Fig. 8. The size of σ_n was varied while all other parameters were at

the same values as in Fig. 8. Target contrast was modelled at 24% and background contrast at 12% (as in

Fig. 5G-L). (A) Illustration of the normalization signal orientation tuning. Fig. 8 was simulated with σ_e =

15° and σ_n = 20° (row 1). (B-D) Model response corresponding to Fig. 8B-D. (E) Model response

integrated over 50-200ms in the same format as Fig. 8E. (F) Correlation of modelled behavioral

performance (Gaussian fit in Fig. 5L) against each time frame of the modelled response. (G) Modelled

populating tuning trajectory corresponding to Fig. 8H.

Methods

All procedures have been approved by the University of Texas Institutional Animal Care and Use
Committee and conform to NIH standards.

Widefield Voltage-Sensitive Dye Imaging

The experimental technique for widefield voltage-sensitive dye (VSD) imaging of neural response in
awake, behaving macaques was adapted from previous studies (Bai et al 2021, Chen et al 2006, Chen et
al 2008, Chen et al 2012). Briefly, two adult male macaque monkeys (Monkey H and Monkey T) were
implanted with a metal head post and metal recording chambers located over the dorsal portion of V1, a
region representing the lower contralateral visual field at eccentricities of 2–5°. Craniotomy and
durotomy were performed. A transparent artificial dura made of silicone was used to protect the brain
while allowing optical access for imaging (Arieli et al 2002)(Fig. 3A). Experiments were conducted in the
left hemisphere chamber of Monkey H, and in both left and right hemisphere chambers of Monkey T.

VSD imaging was used to record neural population activity at a high resolution in space and time
(Shoham et al 1999). Before each experiment, VSD (RH1691 or RH1838) was topically applied on the
cortex for ~2 hours to allow the VSD molecules to bind to cellular membranes. In Monkey H,
fluorescence from neural activity was recorded using Imager 3001 (Optical Imaging, Inc.) using a
tungsten-halogen light source (Zeiss). An infrared eye-tracker (Dr Bouis Inc.) was used to monitor eye
position. In Monkey T, florescence was recorded using custom Matlab software interfaced to PCO Edge
4.2 sCMOS camera (Excelitas PCO GmbH) using X-Cite 110LED light source (Excelitas Technologies Corp).
Eye position was monitored using an Eyelink 100 Plus video eye-tracker (SR Research Ltd).

Both imaging systems were interfaced to a double-SLR-lens-macro system with housing for dichroic
mirrors in between the two SLR lenses. The combination of a 50 mm fixed-focus objective lens (cortex
end, Nikkor 50mm f/1.2) and an 85 mm fixed-focused (Canon EF 85mm f/1.2L USM) camera lens
provided 1.7x magnification, corresponding to imaging approximately an 8 x 8 mm^2 area of the cortex.
Fluorescence signals were measured through a dichroic mirror (650-nm long-pass filter) and an emission
filter (RG 665). VSD molecules were excited by light at 630 nm. Imaging data were collected at 512 ×
512 resolution at 100 Hz. Data acquisition was time locked to the animal's heartbeat (EKG QR up-stroke,
HP Patient Monitor HP78352C). More details about optical imaging with VSD in behaving monkeys are
described elsewhere (Bai et al 2021, Chen et al 2006, Chen et al 2008, Chen et al 2012).

Prior to the main experiments, VSD imaging was used to obtain a precise retinotopic map of the entire
recording (Fig. 3 – figure supplement 1; (Yang et al 2007)). In two out of three chambers, retinotopic
maps indicate that V1 extended into the lunate sulcus. In the third chamber, V1 terminated ~0.75mm
from the lunate sulcus. The area used for decoding analysis was chosen to entirely lie within V1.

Behavioral Task with Optical Stimulation

Monkeys were trained to detect a small additive horizontal Gabor target (4cpd, with σ = 0.14°, 0.33°
FWHM envelope) centered on a sinusoidal grating background mask of the same spatial frequency (4°
raised-cosine windowed). The background grating was oriented at 0°, ±15°, ±30°, ±45°, ±60, and 90°
from the Gabor target orientation. Both the Gabor and background grating were bright-centered – that
is, the 0° orientation background was completely in phase with the target. The contrast of the target and
background were varied in combinations of levels, reported in Michelson contrast:

$$c = \frac{L_{max} - L_{min}}{L_{max} + L_{min}} = \frac{L_{max} - L_{background}}{L_{background}}$$

For each experiment, a Fixation recording block and a Detection recording block were made using the
same target and background conditions. In both blocks, the target and background were centered at a
fixed position for each experiment corresponding to the working cortical chamber. This position varied
between experiments from 1.6 to 3 deg of visual angle eccentricity from the fixation point and from 20
to 50 deg of polar angle from the vertical meridian in the corresponding hemifield (Fig. 3 – figure
supplement 1). At these coordinates, the spatial extent of the target was fully imaged through the
cortical window, and the larger oriented background uniformly activated the entire imaging area.

In the Fixation block, the monkeys were required to remain fixated for each imaging trial while either
the target or full-field sinusoidal gratings at 100% contrast were flashed at 5Hz (60ms ON, 140ms OFF)
for 1.0s. These recordings were processed to obtain retinotopic and columnar orientation response
maps that were used to decode the detection recording responses from the same experiment day (Fig.
3C-E).

In the Detection recordings blocks, a background with random orientation appeared on every trial, with
a 50% chance of an accompanying Gabor target (Fig. 1C-E). The monkeys were tasked to report the
presence of the target. Each trial began with fixation on a bright 0.1° square. An auditory tone and the
dimming of the fixation square cued the monkey to the start of the detection task trial. 250ms later, the
background with or without the target was presented. The monkeys were trained to maintain gaze at
the fixation cue on target absent trials or saccade to and hold gaze (for 150ms) at the target position to
indicate target detection (with a 75ms minimum allowed reaction time). When the target was present, it
remained on screen for a maximum of 250ms or was extinguished immediately upon the monkeys'
saccade initiation. The monkey was given 600ms to make the saccade or to hold fixation and was
subsequently rewarded on correct choices: stay (correct reject) on target absent trials, or saccade to
target (hit) on target present trials. The target and background contrast level were fixed for each
recording block. The probability of each orientated background and target presence were balanced for
each recording block.

A separate target only Detection block on uniform gray background was also taken on each experiment
day using the same routine as above. This block was later used as the reference data to normalize
response amplitudes across experiment days.

Experiments were conducted with custom code using TEMPO real-time control system (Reflective
Computing). The visual stimulus was presented on a Sony CRT (1024x768 @ 100 Hz), distanced 108 cm
from the animal (50 pixels-per-degree), with mean luminance 50 cd/m^2. The visual stimulus was
generated using in-house real-time graphics software (glib).

<u>Behavior Performance and Reaction Time</u>

Behavior performance was calculated for each background orientation and reported in units of
detection sensitivity index d' (d-prime). D-prime and criterion were estimated as:

$$d' = \Phi^{-1}\big(P(HIT)\big) - \Phi^{-1}\big(P(FA)\big), \qquad criterion = -\frac{\Phi^{-1}\big(P(HIT)\big) + \Phi^{-1}\big(P(FA)\big)}{2}$$

where $\Phi^{-1}(x)$ is the inverse transform of the cumulative normal distribution; and $P(HIT)$ and $P(FA)$
represent the proportion of hits and false alarms respectively. To avoid leaving out the data in some

conditions (e.g., when there are no false alarms), we scaled all the proportions to be between 0.005 and
0.995 ($\widehat{P}(x) = 0.005 + 0.99 \cdot P(x)$).

Mapping between the unbiased percentage correct response and d′ is as follows and is depicted in
Figure 2B:

$$d' = 2 \cdot \Phi^{-1}(PC_{max}); \qquad PC_{max} = \Phi\left(\frac{d'}{2}\right)$$

D-prime performance across orientations was fitted with an inverted, dc-shifted Gaussian:

$$y = Ae^{-\frac{\theta^2}{2\sigma^2}} + c$$

The reaction times of the animals were calculated from the stimulus onset to the onset of saccade.
Consequently, there was no measurement of reaction time on trials where the animals remain fixated at
the fixation point.

<u>VSD Imaging</u>

For each trial, an image sequence was captured for a total of 1.2 seconds including pre-stimulus and
post-stimulus frames. The image sequence was analyzed to extract the response using a variant of the
previous reported routines (Bai et al 2021, Chen et al 2006, Chen et al 2008, Chen et al 2012).

Image stabilization was introduced as the first stage of pre-processing to de-accentuate blood vessel
edges in the ΔF/F response map caused by micro movements of the camera and/or the cortex during
imaging. The image intensity across time at each individual pixel was modelled with separable motion-
free ($I_{\mathbf{x}0}(t)$) and motion-related ($\vec{\alpha}_{\mathbf{x}}. \vec{v}(t)$) components as follows:

$$I_{\mathbf{x}}(t) = I_{\mathbf{x}0}(t) + \vec{\alpha}_{\mathbf{x}}\vec{v}(t)$$

For each trial, a single global motion vector $\vec{v}(t)$ was obtained by estimating the translational motion of
the center portion of the images (1/4 of the imaging area). The motion coefficients $\vec{\alpha}_{\mathbf{x}}$ for each pixel was
then obtained using least squares fitting to the model. The motion-corrected image is $I_{\mathbf{x}0}(t)$. This
approach to image stabilization (compared to traditional image registration approach) has the
advantage of correcting for non-rigid movements (rotations, expansion/contractions, affine
transformation, local distortions, etc.) and sub-pixel motion.

<u>Retinotopic and Columnar Template Decoding</u>

Template decoding was used to summarize the retinotopic and columnar response for each image
frame. The retinotopic response map of the target and columnar orientation map of the imaging area
were estimated from the Fixation recording blocks, in which response were stimulated with visual
presentation at 5Hz. The preprocessing steps for the Fixation recording blocks were: image stabilization,
5Hz FFT response extraction, ΔF/F normalization, then down-sampling to 128x128 pixels (from
512x512). Baseline florescence (F_0) was estimated by the average florescence over frames -80 to 0ms
relative to stimulus onset. ΔF/F was calculated as:

$$\frac{\Delta F}{F}(t) = \frac{F(t) - F_0}{F_0}$$

The retinotopic response map of the Gabor target ($H_{ret}(\mathbf{x})$) was estimated by fitting a 2D Gaussian over
the 5Hz flashing Gabor amplitude response. The full-ROI (8x8mm^2 imaging area) columnar orientation

response maps ($H_{ori}(\mathbf{x})$) were estimated from the flashing full-field grating, where the 5Hz FFT grating
response amplitudes were bandpass filtered from 0.8 to 3.0 cycles/mm and the orientation tuning of
each pixel was estimated as described previously (Chen et al 2012). Subsequently, the full-ROI
orientation map was windowed by the retinotopic map to co-localize the retinotopic and the columnar
decoders. The columnar map comprised of pixelwise response magnitude ($A(\mathbf{x})$) and tuning angle
($\theta(\mathbf{x})$), represented in modified Euler's form:

$$H_{ori}(\mathbf{x}) = H_{ret}(\mathbf{x})A(\mathbf{x})e^{2i\theta(\mathbf{x})}$$

These maps served as templates for decoding the retinotopic and columnar response from the
Detection blocks. To further reduce the effect of motion artefacts, a pixel-wise reliability weighted
approach was adopted. The Detection blocks was preprocessed with image stabilization, down-sampling
to 128x128, and ΔF/F normalization as above. From the pre-processed images, the retinotopic scale
variance $\sigma_{ret}^2(\mathbf{x})$ and columnar scale variance $\sigma_{col}^2(\mathbf{x})$ were calculated to be used as reliability weights.
The variance was obtained from condition-mean subtracted residuals taken across all frames across
trials, with the ΔF/F response ($R_{ret}(\mathbf{x}, t)$) bandpass filtered between 0.8 to 3.0 cycles/mm ($R_{col}(\mathbf{x}, t)$)
for the columnar scale variance. Reliability weighting was implemented by normalizing each template
pixel by the corresponding pixel-wise variance.

795
$$\hat{H}(\mathbf{x}) = \frac{H(\mathbf{x})}{\sigma^2(\mathbf{x})}$$

Two different columnar decoding methods were employed. The first examined the overall response
aligned to the orientation of the Gabor (0° orientation tuning axis). A second columnar decoding scheme
was employed to examine the full orientation population response. This second scheme was comprised
of 12 decoders that evenly partitioned the orientation space, such that each decoder contained the
column response magnitude in a subset of pixels tuned with ±7.5° centered at -75° to 90° in 15° steps.
Each decoder therefore represents the population response of similarly tuned neural ensembles
spanning the orientation space every 15°. Each decoder was normalized by the summed response
magnitude of its pixel subset; in this way, each sub-population response was equally represented in the
population tuning curve.

The formulation for the reliability-weighted templates and the decoding is summarized below for the
retinotopic response time course ($r_{ret}(t)$), columnar response time course ($r_{090}(t)$), columnar
population tuning time course ($r_\theta(t)$).

$$r_{ret}(t) = R_{ret}(\mathbf{x}, t)\frac{\hat{H}_{ret}(\mathbf{x})}{\|\hat{H}_{ret}\|}, \qquad \hat{H}_{ret}(\mathbf{x}) = \frac{H_{ret}(\mathbf{x})}{\sigma_{ret}^2(\mathbf{x})}$$

$$r_{090}(t) = R_{col}(\mathbf{x}, t)\frac{\hat{H}_{090}(\mathbf{x})}{\|\hat{H}_{090}\|}, \qquad \hat{H}_{090}(\mathbf{x}) = \frac{real(H_{ori}(\mathbf{x}))}{\sigma_{col}^2(\mathbf{x})}$$

$$r_\theta(t) = R_{col}(\mathbf{x}, t)\frac{\hat{H}_\theta(\mathbf{x})}{\|\hat{H}_\theta\|}, \qquad \hat{H}_\theta(\mathbf{x}) = \frac{H_\theta(\mathbf{x})}{\sigma_{col}^2(\mathbf{x})}, \qquad H_\theta(\mathbf{x}) = \frac{|H_{ori}(\mathbf{x}|\theta)|}{\sum_\mathbf{x}|H_{ori}(\mathbf{x}|\theta)|}$$

<u>Response Pooling across Experiments</u>

Across experiments, the varying effectiveness of VSD staining led to large variations in the noise level
and amplitude of the ΔF/F response. For pooling data across experiments, response amplitudes were
normalized based on singular value decomposition (SVD). Here, the decoded response of the reference
target only Detection block was used. These were trials of varying target contrast levels on the uniform,
mean luminance background instead of the oriented grating background, collected on the same day.

The assumption is that the neural response amplitude and dynamics to the target should be the same
irrespective of the experiment day. Retinotopic and columnar decoder responses for the target only
block were calculated as with the background detection block as described above. Target contrast
response for each experiment were interpolated so that all assessed target contrast levels across
experiments were represented, then SVD was performed over the frames -100 to 200ms about the
stimulus onset across all experiments. In this way, the first component of the SVD (SVD1) represented
the average neural dynamics of the target contrast response, and the SVD1 coefficients represent the
magnitude of this target contrast response each experiment day. Response from each experiment was
then scaled to match the experiment with the largest SVD1 coefficient. This was performed for each
decoded response separately. When pooling data for the mean and standard deviation, the inverse of
the scaling factor squared was used as the reliability weighting.

$$\bar{r} = \frac{1}{n}\sum_{exp,trial} w_{exp} r_{exp,trial}, \qquad \sigma_r = \sqrt{\frac{1}{n}\sum_{exp,trial}\left(w_{exp} r_{exp,trial} - \bar{r}\right)^2}$$

$$w_{exp} = \frac{1}{scaler_{exp}^2} = \frac{1}{\left(SVD1_{max}/SVD1_{exp}\right)^2}$$

Lastly, responses are expressed in a modified z-score, obtained by normalizing the response by its
standard deviation. To calculate this standard deviation, responses were grouped by presented stimuli,
and the means of each group was subtracted. The residuals from the mean-subtraction from frames 50
to 250ms post stimulus onset was pooled according to the aforementioned experiment reliability
weights w_{exp} to obtain the response standard deviation.

Response beyond the saccade may contain unwanted signals. Frames beyond the reaction time for each
trial were therefore omitted from summary statistics. For the integrating response within trials,
responses were averaged up to the frame of saccade. For frame-by-frame averaging across trials for
response time course, trials were dropped out from the averaging beyond their reaction time frame.

<u>Descriptive Trends across Background orientation</u>

Trends were fitted to the normalized VSD response across background orientations (e.g., gray curve in
Fig. 4D). VSD responses were first averaged over a specific range of frames, and for illustration only the
trends of the averaged response across orientations were fitted with either a flat line or a Gaussian.

$$\tilde{r}_{flat} = m\theta + c$$

$$\tilde{r}_{gau} = Ae^{-\frac{\theta}{2\sigma^2}} + c$$

The best fitting trend by the F-test were chosen for display in the figures.

$$\frac{\frac{SS_2 - SS_1}{df_2 - df_1}}{\frac{SS_1}{df_1}} \sim F(df_1, df_2 - df_1)$$

<u>Behavior Correlation</u>

The retinotopic and the columnar response time courses were correlated against the monkey's behavior
judgement. Pearson's correlation coefficient was estimated between the instantaneous response in
time and overall behavior sensitivity index d'. To account for the different trial counts between
background orientations, a trial count weighted version of the correlation coefficient was adopted:

$$\rho = \frac{cov(x,y;w)}{\sqrt{cov(x,x;w)cov(y,y;w)}},$$

$$cov(x,y;w) = \frac{\sum\left(w_i\left(x_i - m(x;w)\right)\right)\left(w_i\left(y_i - m(y;w)\right)\right)}{\sum w_i}, \qquad m(x;w) = \frac{\sum w_i x_i}{\sum w_i}$$

The p-value for the weighted coefficients were estimated using the standard t-score replacing the
degree of freedom with an entropy based effective estimate from the weights.

$$t_\rho(\rho;w) = \frac{\rho\sqrt{n_{eff}-2}}{\sqrt{1-\rho^2}} \qquad \sim tcdf(n_{eff}-2)$$

$$n_{eff} = exp(H) = exp(\sum n_{wi}ln(n_{wi})), \qquad n_{wi} = w_i/\sum w_i$$

<u>Normalization model of orientation masking dynamics</u>

A simple model of the neuronal population response with divisive normalization described our results
qualitatively. In this model, orientation columnar response was tuned to one of 12 different
orientations: -75° to 90° in 15° increments. The responses of each orientation column were specified by
the simple normalization model summarized in Figure 8. The input stimulus is specified by the contrast
and orientation of the target, the contrast and orientation of the background, and the duration of the
stimulus: $\mathbf{s} = (C_T, \theta_T, C_B, \theta_B, D)$. In the model, the input stimulus generates and excitation signal
$r_e\left(t|\mathbf{s},\theta_{max}\right)$ that is linear with stimulus contrast and a normalization signal $r_n\left(t|\mathbf{s},\theta_{max}\right)$ that is also linear
with stimulus contrast. Without loss of generality, all signals were scaled by the 24% contrast target only
response averaged over 50-200ms. These excitation and normalization signals are controlled by the
excitation and normalization parameters, $\boldsymbol{\Omega}_e$ and $\boldsymbol{\Omega}_n$, described below. The normalization signal is then
combined with a normalization constant r_0 to obtain the normalization factor. The normalization
constant limits how small the normalization factor can become. The normalized response is obtained by
dividing the excitation signal by the normalization factor. The final response is then obtained by applying
a response exponent p, which is similar to applying a spiking nonlinearity:

$$r\left(t|\mathbf{s},\theta_{max}\right) = \left[\frac{r_e\left(t|\mathbf{s},\theta_{max}\right)}{normalization\ factor}\right]^p = \left[\frac{r_e\left(t|\mathbf{s},\theta_{max}\right)}{\sqrt[p]{r_n^p\left(t|\mathbf{s},\theta_{max}\right)+r_0^p}}\right]^p = \frac{r_e^p\left(t|\mathbf{s},\theta_{max}\right)}{r_n^p\left(t|\mathbf{s},\theta_{max}\right)+r_0^p}$$

In Figure 8, we show the final responses corresponding to the center of the stimulus $\mathbf{x}_0$. The excitatory
response at that location for background, and target plus background, is obtained by convolving the
effective input contrast signals with the spatial-temporal impulse-response function and evaluating at $\mathbf{x}_0$

$$r_e\left(t|\mathbf{s},\theta_{max}\right) = \begin{cases} \left(c_{Be} * h_e\right)(\mathbf{x}_0,t) \\ \left(c_{TBe} * h_e\right)(\mathbf{x}_0,t) \end{cases}$$

where $c_{Be}\left(\mathbf{x},t\right)$ and $c_{TBe}\left(\mathbf{x},t\right)$ are the effective input contrast signals, and $h_e(\mathbf{x},t)$ is the spatiotemporal
impulse response function. The effective excitatory contrast of the background for the orientation
channel with preferred orientation θ_{max} is given by

881
$$c_{Be}(\mathbf{x},t) = C_B(\mathbf{x})\exp\left(-\frac{(\theta_B - \theta_{\max})^2}{2\sigma_e^2}\right)w(t;D)$$

882 where $C_B(\mathbf{x})$ is the background contrast, σ_e is the falloff parameter of column's orientation turning
883 function, and $w(t;D)$ is a temporal pulse function of width D. Similarly, the effective target contrast is
884 given by

885
$$c_{Te}(\mathbf{x},t) = C_T(\mathbf{x})\exp\left(-\frac{(\theta_T - \theta_{\max})^2}{2\sigma_e^2}\right)w(t;D)$$

886 The effective excitatory contrast for target plus background is slightly more complicated because there
887 can be some contrast summation or cancellation depending on the phase and orientation of the target
888 relative to the background

889
$$c_{TBe}(\mathbf{x},t) = \lambda(\mathbf{x})\sqrt{c_{Be}^2(\mathbf{x},t) + c_{Te}^2(\mathbf{x},t)}$$

890 where $\lambda(\mathbf{x})$ is the contrast correction factor,

891
$$\lambda(\mathbf{x}) = \frac{C_{TB}(\mathbf{x})}{\sqrt{C_B^2(\mathbf{x}) + C_T^2(\mathbf{x})}}$$

892 The spatiotemporal impulse response function is the separable product of a Gaussian distribution and a
893 gamma distribution

894
$$h_e(\mathbf{x},t) = gauss(\mathbf{x};s_e)\,gamma(t;a_e,b_e)$$

where s_e is the standard deviation of the 2D Gaussian and a_e, b_e are the two parameters of a gamma
distribution: $gamma(t;a_e,b_e) = b_e^{-a_e}t^{a_e-1}\exp(-t/b_e)/\Gamma(a_e)$.

The formulas for the normalization signal are the same as for the excitatory signal, except the four
parameters are allowed to differ: $\mathbf{\Omega}_e = (\sigma_e, s_e, a_e, b_e)$, $\mathbf{\Omega}_n = (\sigma_n, s_n, a_n, b_n)$.

Parameter values from known properties of single neurons in primary visual cortex were adopted. The
response exponent p was constrained to 2.0, consistent with single neuron contrast-response functions
(Albrecht & Hamilton 1982, Geisler & Albrecht 1997, Sclar et al 1990). It was assumed that the peak
orientation of the excitatory signal and suppressive normalization signal θ_{max} were the same
(Cavanaugh et al 2002a), but that the orientation bandwidth of the normalization signal was greater
than that of the excitation signal, $\sigma_n > \sigma_e$ (Cavanaugh et al 2002b), and that the spatial pooling region
for the normalization signal was larger than that for the excitation signal, $s_n > s_e$ (Cavanaugh et al
2002a, Cavanaugh et al 2002b, Levitt & Lund 2002, Sceniak et al 2001). As shown in Fig. 8 – figure
supplement 1, we found that setting $s_n > s_e$ had little effect on the modeling of our empirical results
(Figs. 5); for simplicity, our final model assumed $s_n = s_e$. Finally, it was assumed that the temporal
dynamics of the normalization signal, determined by parameters a_n and b_n, are slower than those for
the excitation signal, determined by parameters a_e and b_e (Groen et al 2022, Zhou et al 2019).

The following were the parameter values used in Figure 8: $r_0 = 0.03125$,

$$\mathbf{\Omega}_e = (\sigma_e = 15°,\ a_e = 9,\ b_e = 8ms,\ s_e = 0.28°)$$

$$\boldsymbol{\Omega}_n = (\sigma_n = 20°, \; a_n = 9, \; b_n = 10.64ms, \; s_n = 0.28°)$$

All analyses were done using Matlab R2018a.

<u>Statistics</u>

Two animals were examined to verify the consistency of experimental approach and results. Multiple
recordings were made from the same animals. The number of recordings were based on previous
experience; no statistical method was used to predetermine sample size.

Statistical analyses were conducted in Matlab (R2018a).

<u>Data and Code Availability</u>

The data and custom code from this study will be made available upon manuscript acceptance.

References

References

Albrecht DG, Geisler WS. 1991. Motion Selectivity and the Contrast-Response Function of Simple Cells in
the Visual-Cortex. *Visual Neuroscience* 7: 531-46

Albrecht DG, Hamilton DB. 1982. Striate Cortex of Monkey and Cat - Contrast Response Function.
Journal of Neurophysiology 48: 217-37

Allman J, Miezin F, McGuinness E. 1985. Stimulus specific responses from beyond the classical receptive
field: neurophysiological mechanisms for local-global comparisons in visual neurons. *Annu Rev*
Neurosci 8: 407-30

Angelucci A, Bijanzadeh M, Nurminen L, Federer F, Merlin S, Bressloff PC. 2017. Circuits and Mechanisms
for Surround Modulation in Visual Cortex. *Annu Rev Neurosci* 40: 425-51

Angelucci A, Bressloff PC. 2006. Contribution of feedforward, lateral and feedback connections to the
classical receptive field center and extra-classical receptive field surround of primate V1. *Visual*
Perception, Part 1, Fundamentals of Vision: Low and Mid-Level Processes in Perception 154: 93-
120

Arieli A, Grinvald A, Slovin H. 2002. Dural substitute for long-term imaging of cortical activity in behaving
monkeys and its clinical implications. *Journal of Neuroscience Methods* 114: 119-33

Bai Y, Chen S, Chen Y, Geisler WS, Seidemann E. 2021. Similar masking effects of natural backgrounds on
detection performances in humans, macaques, and macaque-V1 population responses. *J*
Neurophysiol 125: 2125-34

Benvenuti G, Chen Y, Ramakrishnan C, Deisseroth K, Geisler W, Seidemann E. 2018. Scale-Invariant
Visual Capabilities Explained by Topographic Representations of Luminance and Texture in
Primate V1. *Neuron* 100: 1-9

Campbell FW, Kulikowski JJ. 1966. Orientational selectivity of the human visual system. *J Physiol* 187:
437-45

Carandini M, Heeger DJ. 1994. Summation and Division by Neurons in Primate Visual-Cortex. *Science*
264: 1333-36

Cavanaugh JR, Bair W, Movshon JA. 2002a. Nature and interaction of signals from the receptive field
center and surround in macaque V1 neurons. *Journal of Neurophysiology* 88: 2530-46

Cavanaugh JR, Bair W, Movshon JA. 2002b. Selectivity and spatial distribution of signals from the
receptive field surround in macaque V1 neurons. *Journal of Neurophysiology* 88: 2547-56

Chen Y, Geisler WS, Seidemann E. 2006. Optimal decoding of correlated neural population responses in
the primate visual cortex. *Nat Neurosci* 9: 1412-20

Chen Y, Geisler WS, Seidemann E. 2008. Optimal Temporal Decoding of V1 Population Responses in a
Reaction-Time Detection Task. *J Neurophysiol* 99: 1366-79

Chen Y, Palmer CR, Seidemann E. 2012. The relationship between voltage-sensitive dye imaging signals
and spiking activity of neural populations in primate V1. *Journal of Neurophysiology*: 3281-95

Clifford CW. 2014. The tilt illusion: phenomenology and functional implications. *Vision Res* 104: 3-11

Foley JM. 1994. Human luminance pattern-vision mechanisms: masking experiments require a new
model. *Journal of the Optical Society of America. A, Optics, image science, and vision* 11: 1710-9

Frazor RA, Geisler WS. 2006. Local luminance and contrast in natural images. *Vision Res* 46: 1585-98

Geisler WS, Albrecht DG. 1997. Visual cortex neurons in monkeys and cats: Detection, discrimination,
and identification. *Visual Neuroscience* 14: 897-919

Grinvald A, Hildesheim R. 2004. VSDI: A new era in functional imaging of cortical dynamics. *Nature*
Reviews Neuroscience 5: 874-85

Groen IIA, Piantoni G, Montenegro S, Flinker A, Devore S, et al. 2022. Temporal dynamics of neural
responses in human visual cortex. *J Neurosci* 42: 7562-80

Heeger DJ. 1991. Nonlinear model of neural responses in cat visual cortex. *Computational models of*
visual processing: 119-33
Heeger DJ. 1992. Normalization of Cell Responses in Cat Striate Cortex. *Visual Neuroscience* 9: 181-97
Henry CA, Jazayeri M, Shapley RM, Hawken MJ. 2020. Distinct spatiotemporal mechanisms underlie
extra-classical receptive field modulation in macaque V1 microcircuits. *eLife* 9
Hubel DH, Wiesel TN. 1959. Receptive fields of single neurones in the cat's striate cortex. *The Journal of*
Physiology 148: 574-91
Hubel DH, Wiesel TN. 1968. Receptive fields and functional architecture of monkey striate cortex.
Journal of Physiology 195: 215-43
Levitt JB, Lund JS. 2002. The spatial extent over which neurons in macaque striate cortex pool visual
signals. *Visual Neuroscience* 19: 439-52
Michel MM, Chen Y, Seidemann E, Geisler WS. 2018. Nonlinear Lateral Interactions in V1 Population
Responses Explained by a Contrast Gain Control Model. *J Neurosci* 38: 10069-79
Ozeki H, Finn IM, Schaffer ES, Miller KD, Ferster D. 2009. Inhibitory Stabilization of the Cortical Network
Underlies Visual Surround Suppression. *Neuron* 62: 578-92
Polat U, Mizobe K, Pettet MW, Kasamatsu T, Norcia AM. 1998. Collinear stimuli regulate visual
responses depending on cell's contrast threshold. *Nature* 391: 580-84
Rubin DB, Van Hooser SD, Miller KD. 2015. The Stabilized Supralinear Network: A Unifying Circuit Motif
Underlying Multi-Input Integration in Sensory Cortex. *Neuron* 85: 402-17
Sceniak MP, Hawken MJ, Shapley R. 2001. Visual spatial characterization of macaque V1 neurons. *J*
Neurophysiol 85: 1873-87
Sceniak MP, Ringach DL, Hawken MJ, Shapley R. 1999. Contrast's effect on spatial summation by
macaque V1 neurons. *Nat Neurosci* 2: 733-39
Schwartz O, Hsu A, Dayan P. 2007. Space and time in visual context. *Nat Rev Neurosci* 8: 522-35
Schwartz O, Simoncelli EP. 2001. Natural signal statistics and sensory gain control. *Nature Neuroscience*
4: 819-25
Sclar G, Maunsell JH, Lennie P. 1990. Coding of image contrast in central visual pathways of the macaque
monkey. *Vision research* 30: 1-10
Sebastian S, Abrams J, Geisler WS. 2017. Constrained sampling experiments reveal principles of
detection in natural scenes. *Proc Natl Acad Sci U S A* 114: E5731-e40
Seidemann E, Arieli A, Grinvald A, Slovin H. 2002. Dynamics of depolarization and hyperpolarization in
the frontal cortex and saccade goal. *Science* 295: 862-65
Seidemann E, Geisler WS. 2018. Linking V1 Activity to Behavior. *Annual review of vision science* 4: 287-
310
Shoham D, Glaser DE, Arieli A, Kenet T, Wijnbergen C, et al. 1999. Imaging cortical dynamics at high
spatial and temporal resolution with novel blue voltage-sensitive dyes. *Neuron* 24: 791-802
Shushruth S, Mangapathy P, Ichida JM, Bressloff PC, Schwabe L, Angelucci A. 2012. Strong recurrent
networks compute the orientation tuning of surround modulation in the primate primary visual
cortex. *J Neurosci* 32: 308-21
Sit Y-F, Chen Y, Geisler WS, Miikkulainen R, Seidemann E. 2009. Complex dynamics of V1 population
responses explained by a simple gain-control model. *Neuron* 24: 943-56
Stromeyer CF, 3rd, Julesz B. 1972. Spatial-frequency masking in vision: critical bands and spread of
masking. *Journal of the Optical Society of America* 62: 1221-32
Tsodyks MV, Skaggs WE, Sejnowski TJ, McNaughton BL. 1997. Paradoxical Effects of External Modulation
of Inhibitory Interneurons. *The Journal of Neuroscience* 17: 4382-88
Watson AB, Solomon JA. 1997. Model of visual contrast gain control and pattern masking. *Journal of the*
Optical Society of America. A, Optics, image science, and vision 14: 2379-91

Wenderoth P, Johnstone S. 1987. Possible neural substrates for orientation analysis and perception.
1018   *Perception* 16: 693-709
1019   Wilson HR, McFarlane DK, Phillips GC. 1983. Spatial-Frequency Tuning of Orientation Selective Units
1020   Estimated by Oblique Masking. *Vision Research* 23: 873-82
1021   Yang Z, Heeger DJ, Seidemann E. 2007. Rapid and precise retinotopic mapping of the visual cortex
1022   obtained by voltage sensitive dye imaging in the behaving monkey. *J Neurophysiol* 98: 1002-14
1023   Zhou J, Benson NC, Kay K, Winawer J. 2019. Predicting neuronal dynamics with a delayed gain control
1024   model. *PLoS Comput Biol* 15: e1007484

1025