NEURAL CORRELATES OF PERCEPTUAL SIMILARITY MASKING IN PRIMATE V1

Spencer C. Chen[1-5], Yuzhi Chen[1-4], Wilson S. Geisler[1-3], & Eyal Seidemann[1-4]

[1]Center for Perceptual Systems, [2]Center for Theoretical and Computational Neuroscience
[3]Department of Psychology, [4]Department of Neuroscience, University of Texas at Austin
[5]Department of Neurosurgery, Rutgers University-

Abstract

Visual detection is a fundamental natural task. Detection becomes more challenging as the similarity between the target and the background in which it is embedded increases, a phenomenon termed "similarity masking". To test the hypothesis that V1 contributes to similarity masking, we used voltage sensitive dye imaging (VSDI) to measure V1 population responses while macaque monkeys performed a detection task under varying levels of target-background similarity. Paradoxically, we find that during an initial transient phase, V1 responses to the target are enhanced, rather than suppressed, by target-background similarity. This effect reverses in the second phase of the response, so that in this phase V1 signals are positively correlated with the behavioral effect of similarity. Finally, we show that a simple model with delayed divisive normalization can qualitatively account for our findings. Overall, our results support the hypothesis that a nonlinear gain control mechanism in V1 contributes to perceptual similarity masking.

Introduction

Searching for, and detecting, visual targets in our environment is a ubiquitous natural task that our visual system performs exceptionally well. A key feature of behavioral detection performance is that the texture similarity between the target and the background in which it is embedded profoundly affects target detectability. The more similar are the target and the background, the harder it is to detect the target (Campbell & Kulikowski 1966, Foley 1994, Sebastian et al 2017, Stromeyer & Julesz 1972, Watson & Solomon 1997, Wilson et al 1983). This phenomenon, which is termed "similarity masking", is the foundation of camouflage.

An example of similarity masking is illustrated in figure 1. Detecting a low contrast oriented visual target is easy on a uniform gray background (Fig. 1A). Detectability decreases when the target has a similar orientation as the background (Fig. 1B). The neural basis of similarity masking is not well understood. The main goal of the current study was to test the hypothesis that neural interactions between the representations of the target and background in the primary visual cortex (V1) contribute to the perceptual effect of similarity masking.

The responses of visual neurons to a target can be strongly modulated by the context in which the stimulus is presented. Such contextual modulations have powerful, complex and diverse effects in the visual cortex (Allman et al 1985, Angelucci et al 2017, Angelucci & Bressloff 2006, Bai et al 2021,

Cavanaugh et al 2002b, Michel et al 2018, Sceniak et al 1999). Most of these effects reflect sublinear
interactions between the target and the background, suggesting that they could potentially contribute
to behavioral masking effects. If nonlinear computations in V1 contribute to similarity masking, we
would predict that the signals evoked by a target will be maximally reduced by a background that is
similar to the target.

We tested this hypothesis by measuring V1 population responses in macaque monkeys while they
performed a visual detection task under masking conditions (Fig. 1C-E). Because the nature of
contextual modulations in V1 is complex, a second goal of our study was to quantitatively characterize
the spatiotemporal dynamics of V1 population responses to different combinations of targets and
backgrounds.

As a first step, we characterized the behavioral effects of similarity masking in two macaque monkeys,
demonstrating clear effects of similarity on target detectability and reaction times. These results confirm
that macaque monkeys are a good animal model for human similarity masking.

Second, we used voltage-sensitive dye imaging (VSDI; (Grinvald & Hildesheim 2004, Seidemann et al
2002, Shoham et al 1999)) to measure V1 population responses at two scales: the scale of the
retinotopic map and the scale of orientation columns, while the monkeys performed the similarity
masking detection task. To study the effect of similarity masking on the neural detection sensitivity, we
constructed a task-specific decoder at each scale. Each decoder first pools the responses using a scale-
dependent spatial template and then combines these responses over time to form a decision variable.
The distributions of the decision variable in target-present vs. target-absent trials are used to compute
neural sensitivity that can be compared to behavioral sensitivity (Seidemann & Geisler 2018).

We found that V1 population responses to the target and background display two distinct phases. An
initial transient phase that starts at response onset, and a second phase that lasts until stimulus offset or
the animal's response. Surprisingly, the first phase displays a paradoxical effect; during this phase the
target evoked response is strongest when the target and background are similar and is therefore anti-
correlated with behavior. This effect reverses in the second phase so that in this phase the target-
evoked response is reduced with increased target-background similarity. V1 responses during this
second phase are therefore consistent with behavior.

We also observed complex spatiotemporal dynamics of the population response to the target and
background stimuli, including a repulsion of V1 columnar-scale representation of target orientation in
the direction away from the background orientation.

Finally, we show that a simple dynamic population gain control model can qualitatively account for our
physiological and behavioral results, and that the estimated properties of the gain-control mechanism
are consistent with a principled computational approach to feature encoding and decoding. Overall, our
results are consistent with the hypothesis that contextual interactions between the representations of
the target and background in V1 are likely to contribute to the perceptual phenomena of similarity
masking.

Results

Behavioral effect of target-background similarity masking

To study the neural basis of visual similarity masking, we trained two monkeys (macaque mulatta) to perform a visual detection task in which a small horizontal target appeared on a larger background at a known location in half of the trials (Fig. 1E). The monkey indicated target absence by maintaining fixation and target presence by making a saccadic eye movement to the target location as soon as it was detected. Within a block of trials, the contrast of the target and the background were fixed, while the orientation of the background varied randomly from trial to trial, allowing us to test for the effect of target-background orientation similarity on behavioral and neural detection sensitivities.

We tested the behavioral effect of similarity masking over five combinations of target and background contrasts (Fig. 2A). For each combination, we measured the behavioral sensitivity as a function of background orientation. Performance with no background (uniform gray screen) served as a baseline (Fig. 2A, dashed horizontal lines). Performance as a function of background orientation was fitted with an inverted Gaussian function. At all five target and background contrast combinations, detection sensitivity was lowest when background and target orientations matched, confirming the expected similarity masking effect from human subjects. These results demonstrate that macaque monkeys are a good animal model for studying the neural basis of human similarity masking.

The supplementary information includes a perceptual demonstration of similarity masking for a wide range of target amplitudes, orientations, and spatial-frequencies (Supplementary Material: "Orientation Masking Demo"). This demonstration can give the reader an intuitive sense of the masking effects studied here.

We also examined the effect of target-background orientation similarity on the monkeys' reaction times (Fig. 2C). We find two distinct effects of orientation similarity on reaction times. At higher target and background contrasts, reaction times are maximal when the background and target have the same orientation (when detectability is lowest and the task is hardest) and monotonically decrease as target-background similarity decreases (detectability increases and the task becomes easier). Surprisingly, at lower target and background contrasts, reaction times are low when the background matches target orientation, then increases as the background-target orientation difference increases, and then drops again when the background approaches the orthogonal orientation to the target. Thus, under these conditions, we see an interesting decoupling between difficulty and reaction time, so that reactions times can be shortest in the harder conditions. This surprising effect is present in both monkeys.

Our next goal was to test the hypothesis that contextual interactions between the representations of the target and background in V1 contribute to the observed behavioral similarity masking results.

Neural population responses to target and background stimuli in macaque V1

While the monkeys performed the similarity masking detection task, we used VSDI to measure V1 population responses to the target and the background. In each cranial window, we first used a fast and efficient VSDI protocol to obtain a detailed retinotopic map (Yang et al 2007). We then positioned the target so that its neural representation fell at the center of our imaging area.

The target elicits V1 population activity at two fundamental spatial scales. At the large retinotopic scale,
the target evokes an activity envelope that spreads over several mm^2 and is well fitted by a two-
dimensional (2D) Gaussian (Fig. 3B, top row; (Chen et al 2006, Chen et al 2012, Sit et al 2009)). Our 8x8
mm^2 imaging area allows us to capture this entire target-responsive region. Because the background is
much larger than the target and is centered at the same location in the visual field, it produces a
relatively uniform response within the imaging area (Fig. 3B, second row). Similarly, the target-plus-
background stimulus elicits activity within the entire imaged area, with a relatively elevated activity at
the retinotopic region corresponding to the target location (Fig. 3B, 3rd row). However, the target-
evoked response in the presence of the background (response to target plus background minus
response to the background alone) appears significantly weaker than the response to the target alone
(Fig. 3B, 1st vs. 4th row). This reduced target-evoked response in the presence of the background could
contribute to the perceptual masking effect of the background. Our goal here was to determine how this
sublinear interaction between the response to the target and background depends on target-
background similarity in orientation.

In addition to the retinotopic-scale response envelope, fine scale response modulations at a scale of
individual orientation columns (diameter of ~0.3 mm) reflect the orientation of the target and
background. These columnar scale modulations have a relatively small amplitude and therefore appear
as small ripples riding on top of the larger retinotopic response envelope. The relatively smaller VSDI
responses at the columnar scale is due to a mixture of robust non-orientation selective V1 population
responses in V1 as well as optical and biological blurring (Chen et al 2012). We can selectively access the
columnar scale signals by spatially filtering the responses at the scale of the orientation columns.
Despite their small relative amplitude, these columnar-scale signals provide high quality single-trial
orientation decoding (Benvenuti et al 2018).

Retinotopic-scale effect of target-background similarity masking

To study the effect of similarity masking on V1 responses at the retinotopic scale, we used an optimal
linear decoder of V1 population responses (Chen et al 2006, Chen et al 2008) that allows us to assess the
neural detection sensitivity of V1 population responses (i.e., how well one can detect the target from
single-trial V1 population responses) (Fig. 3C-D). The retinotopic decoder takes into account the location
and shape of the envelope of the target-evoked response (Fig. 3D), as well as the structure of the noise
covariance matrix (Fig. 3C; (Chen et al 2006)).

Figure 4 summarizes the dynamics of the retinotopic template output in response to the V1 signals
across all of our experiments for two combinations of target and background contrasts (see Table 1 and
Fig. S1 for the full set of tested target/background combinations). When presented on a uniform gray
background, the target-related retinotopic signal begins to rise ~40 ms after target onset, reaches its
peak ~100 ms after stimulus onset, and remains high for the next 100 ms (Fig. 4B,H, black curve).
However, when the same target is added to the background, the target-related retinotopic signals
display a wide range of responses that depend on background orientation (Fig. 4B,H, colored curves).

Our main interest here is in the target-evoked response in the presence of the background (Fig. 4C,I),
which can be extracted by subtracting the response to the background alone (Fig. 4A,G) from the
response to the target plus background (Fig. 4B,H). If V1 contextual interactions at the retinotopic scale
contribute to the behavioral effect of similarity masking, we would expect that target-evoked responses

would be weakest with high target-background similarity (similar target and background orientations)
and strongest with low target-background similarity (orthogonal target and background orientations).

Surprisingly, we find that the target-evoked response in V1 displays two distinct phases, with the early
phase showing a paradoxical neural dependence on target-background orientation similarity that is anti-
correlated with the behavioral masking effect, and with a later phase that is consistent with the
behavioral masking effect. Specifically, in the early phase which starts at response onset, the target-
evoked response is highest when the background matches the target orientation even though
behaviorally this is the condition in which detection performance is the worst. However, after this initial
phase, the high-similarity target-evoked response starts to drop, while the low-similarity target-evoked
response continues to build up, so that in the later stages the target-evoked response is strongest on the
dissimilar background and weakest on the similar background, consistent with the behavioral effect of
similarity.

To quantify the relation between the neural and behavioral effects of similarity masking, we computed
the correlation between the effect of orientation similarity on behavior and its effect on the integrated
neural responses (Fig. 4 D,J) in individual 10 ms frames (Fig. 4 E,K). This analysis reveals a robust
paradoxical negative correlation between the early neural V1 response and behavior, weak positive
correlation between the late neural V1 response and behavior, and no correlation between behavior
and the integrated neural response.

Because the target and background are defined by their orientation, the correspondence between the
neural signals in V1 and behavior may be better captured by V1 responses at the columnar scale. Our
next step was therefore to examine the dynamics of the columnar-scale target-evoked responses in V1.

Neural effects of target-background similarity masking at the scale of orientation columns

To study the effect of similarity masking on V1 responses at the columnar scale, we developed a linear
columnar decoder of the VSDI signals (Fig 3E). The columnar decoder takes into account the location of
the orientation columns within the retinotopic envelope of the target-evoked response. Because the
target is horizontal, the output of the columnar template is expected to be positive for the horizontal
target and background stimuli and negative for the vertical background stimulus (since the horizontal
and vertical columnar maps are anti-correlated).

As with the output of the retinotopic-scale template, the output of the columnar-scale template displays
two distinct phases. Figure 5 shows the time course of the columnar template signals to background
alone (Fig. 5A,G), the target plus background (Fig. 5B,H), and target in the presence of the background
(Fig. 5C,I). In the early phase, the target-evoked response is highest when the background and target
have similar orientations, producing a paradoxical neural response that is anti-correlated with the
behavioral masking effect (Fig. 5D,E,J,K). In the second phase the trend reverses and the target-evoked
response is strongest on the dissimilar background and weakest on the similar background, consistent
with the behavioral effect of similarity (Fig. 5D,E,J,K). Similar results were obtained with other target and
background contrast combinations (Figure S2).

Because the first phase of the response is shorter than the second phase, when V1 response is
integrated over both phases, the overall response is positively correlated with the behavioral masking

effect. Therefore, our results suggest that the neural masking effect at the columnar scale in V1 could
play a major role in the behavioral similarity masking effects.

Dynamics of columnar-scale orientation population trajectories

Our decoding analysis focuses on the columnar-scale orientation signals along the 0°-90° axis and
reveals complex columnar-scale dynamic interactions between the target-evoked response and the
response evoked by the background (Fig. 5). To examine these dynamics in more details, we performed
two types of population-vector analyses (Fig. 6). First, we assigned each pixel within the retinotopic
footprint of the target-evoked response to one of 12 equally spaced preferred orientations, creating 12
orientation selective clusters of pixels. We then computed for each stimulus the response in each
orientation selective cluster in each frame and displayed the population orientation tuning curve as a
function of time (Fig. 6C) and as population vector dynamic trajectory in the polar space spanned by the
12 orientations (Fig. 6D); i.e., the orientation θ and magnitude R_θ of the peak of the population
response over time.

The population vector analysis reveals that V1 responses to the target or background alone are
consistent with the stimulus orientation. In background only trials, shortly after stimulus onset the peak
of the population tuning curve closely matches background orientation (Fig. 7A, F, top row, red arrow
and horizontal line). Similar results were obtained in the target only trials, where the peak of the
population response tuning curve matches target orientation (Fig. 7A, F, 4th row, green arrow and
horizontal line).

Target plus background responses display complex spatiotemporal dynamics. To examine the dynamics
of the target-evoked response in the presence of the background, we subtracted the background only
response from the target plus background response (Fig. 7A, F, 3rd row). The results reveal complex
target-background interactions which could lead to a population tuning peak (white curve) that
significantly deviates from the target orientation. For example, in some conditions, we observe an
orientation tuning peak that is repelled from target orientation in the direction away from background
orientation. An interesting goal of future studies would be to examine potential perceptual correlates of
these interactions.

To further examine the dynamics of the population vector, we plotted the response trajectories for each
stimulus using the vector representation in polar coordinates (Fig. 7B-E, G-J). After stimulus onset, the
population vector for background only moves in the direction corresponding to the background
orientation (Fig. 7B,G) and for target only moves in direction corresponding to the target (Fig. 7E,J).

The trajectories in the target plus background conditions are more complex. For example, when
background orientation is at +/- 45 deg to the target, the population response is initially dominated by
the background, but then in mid-flight, the population response changes direction and turns toward the
direction of the target orientation.

Such complex interactions can be used to constrain models of V1 population response.

Dynamic gain control model qualitatively captures similarity masking effects in V1

Our next goal was to determine whether the observed interactions between the background- and
target-evoked responses can be qualitatively captured by a gain control model. In this model,

orientation columnar response was tuned to one of 12 equally spaced orientations. The responses of
each orientation column were specified by the simple normalization model summarized in Figure 8.
Specifically, the spatiotemporal input stimulus generates an excitation signal and a normalization signal
that are both linear with the input root mean square (rms) contrast. The normalization signal is then
combined with a normalization constant to obtain the normalization factor. The normalized response is
obtained by dividing the excitation signal by the normalization factor. The final response is then
obtained by applying a response exponent p, which is similar to applying a spiking nonlinearity.
Importantly, the excitation and normalization signals can differ in their spatial extent, orientation tuning
width, and temporal impulse response (see Methods for model parameters).

We find that this simple model qualitatively captures our key results. First, in response to background
alone (Fig. 8B), the modeled population vector peaked at ~100 ms after stimulus onset and then
dropped to a lower amplitude, as in our data (Fig. 5A,G). This reduction in response amplitude was due
to normalization signal that was delayed relative to the excitation signal. Second, as in the real data,
response to the target plus background is less than the sum of the responses to each component
separately. Third, as in our results, the target-evoked response in the presence of the background is
biphasic, having a brief early component in which the response is enhanced by target-background
similarity, and a longer-lasting late component in which the response is suppressed by target-
background similarity (Fig. 8D). This leads to an early phase in which the response is anticorrelated with
the behavioral effect of similarity masking, and a late phase that is positively correlated with the
behavioral effect of similarity masking (Fig. 8E).

Finally, this simple model can also display the curved trajectories of the population vector in response to
the target plus background (compare Fig. 8G to Fig. 7C,H). Overall, our results suggest that a simple gain
control model with different spatial extent, orientation tuning width, and temporal dynamics of
excitation and divisive gain control, can qualitatively capture the complex spatiotemporal dynamics of
V1 population responses to localized oriented targets added to a larger oriented background.

Discussion

To test the hypothesis that nonlinear computations in V1 contribute to the perceptual effect of similarity
masking, we used voltage-sensitive dye imaging (VSDI) to measure neural population responses from V1
in two macaque monkeys while they performed a visual detection task in which a small oriented target
was detected in the presence of a larger background of varied orientations. Like human observers, the
monkeys were strongly affected by the orientation similarity of the target and the background. Their
detection threshold increased with increased target-background orientation similarity, while their
reaction times showed complex, and in some cases non-monotonic, dependency on target-background
orientation similarity.

To quantify the neural effects of similarity masking, we measured neural sensitivity to the target at two
fundamental spatial scales of V1 topographic representations. The large scale of the retinotopic map
and the finer scale of the columnar orientation map. We discovered that at both scales, V1 population
responses to the target and background display two distinct phases. An initial transient phase in which
target-evoked V1 response is strongest when the target and background have similar orientations. At
this early phase, V1 responses are therefore paradoxically anti-correlated with the behavioral effect of

similarity masking. In the second phase, the masking effect reverses, and the target-evoked response is
maximally reduced when the target and background are similar. In this second sustained phase V1
population responses are therefore consistent with the behavioral similarity masking effect.

The positive correlation between the neural and behavioral masking effects occurred earlier and was
more robust at the columnar scale than at the retinotopic scale, suggesting that behavioral performance
in our task is dominated by columnar scale signals in the second phase of the response. To the best of
our knowledge, this is the first demonstration of such decoupling between V1 responses at the
retinotopic and columnar scales, and the first demonstration that columnar scale signals are a better
predictor of behavioral performance in a detection task.

We find that when the target and background have similar orientations, columnar-scale information
about the target is restricted to the first phase of the response and then largely disappears during the
second phase of the response. These physiological results could be related to the surprising mismatch
between task difficulty and reaction times. Rather than having reaction times that monotonically
increase with task difficulty, in our masking detection task, reaction times can be shortest when target
and background orientations match, even though it is hardest to detect the target under these
conditions. The short reaction time to this stimulus may be the consequence of the target information
being best represented in the early phase of the response.

The nature of contextual modulations in V1 is quite complex (Angelucci et al 2017, Angelucci & Bressloff
2006, Bai et al 2021, Cavanaugh et al 2002b, Michel et al 2018, Polat et al 1998, Sceniak et al 1999). A
second goal of our study was to quantitatively characterize the spatiotemporal dynamics of columnar-
scale V1 population responses to targets and backgrounds of different orientations and contrasts. Using
a dynamic population vector analysis, we find that in the presence of an oriented background, the peak
of the population orientation tuning to the target can deviate significantly from target orientation. For
example, in some conditions, we observe a population orientation tuning peak that is repelled away
from target orientation in the direction opposite to background orientation (Fig. 7E). These orientation-
dependent interactions could contribute to non-veridical perceptual representations of orientation such
as in the well-known tilt illusion effect (Clifford 2014, Schwartz et al 2007, Wenderoth & Johnstone
1987). An important goal for future studies would be to test for this possibility.

Using the population vector analysis, we find that columnar scale V1 representations are initially
dominated by the orientation of the background. The target orientation then appears in the second
phase of the response, which leads to curved population vector trajectories (Fig. 7C, H). Identifying
possible perceptual consequences of such dynamic and complex trajectories and understanding the
neural circuit mechanisms that give rise to such responses are two important goals for future work.

Nonlinear response properties in V1 are commonly modeled as a consequence of a divisive gain control
mechanism (Albrecht & Geisler 1991, Carandini & Heeger 1994, Heeger 1991, Heeger 1992, Sit et al
2009). As a first step toward understanding the mechanisms that could give rise to the observed V1
responses, we tested whether a simple dynamic gain control model could account for our findings (Fig.
8). We find that a simple gain control model can qualitatively account for our results, but that in order to
do so, the model has to display two important properties. First, to account for the biphasic nature of V1
response, the divisive normalization signals have to be delayed relative to the excitatory signal. Second,
in order to account for the reduced neural sensitivity with target-background similarity in the second
phase of the response, the divisive normalization signal has to be orientation selective. Because in

primates and carnivores, robust orientation selectivity first emerges in V1 (Hubel & Wiesel 1959, Hubel & Wiesel 1968), these results suggest that a significant portion of the nonlinear interactions observed in the current study originate in V1 rather than being inherited from the ascending inputs that V1 receives from the LGN. While our experimental and computational results point to a delayed gain control signal that operates at the level of V1, they do not directly speak to the circuit and biophysical mechanisms that contribute to the implementation of this gain control in V1. Multiple candidate mechanisms for implementing gain control in V1 have been proposed (Angelucci et al 2017, Angelucci & Bressloff 2006, Ozeki et al 2009, Rubin et al 2015, Tsodyks et al 1997). Our results provide new and powerful constraints for such mechanistic models.

Finally, we note that the tuned similarity normalization that explains the neural and behavioral similarity-masking effects reported here is consistent with a principled encoding strategy for feature detection under natural conditions. When viewing a static scene under natural conditions, human and non-human primates make 3-4 saccadic eye movements per second, with fixations between saccades of 200-300 ms. Given the typical size of the saccades, most visual receptive fields are stimulated during each fixation by a largely statistically independent random sample of natural image (Frazor & Geisler 2006). Analysis of the responses of linear receptive fields to random samples of natural image shows that the standard deviation of the response increases in proportion to the product of the luminance, the contrast and the similarity of the natural background to the receptive field (Sebastian et al 2017). Thus, divisive normalization by the product of luminance, contrast and similarity causes the standard deviation of the responses across natural images to be much more constant (i.e., nearly independent of the luminance, contrast and similarity of the background within the receptive field). This more constant standard deviation makes it possible, with relatively simple decoders, to reach near optimal feature-detection performance, under the high levels of stimulus uncertainty that occur under natural conditions (Schwartz & Simoncelli 2001, Sebastian et al 2017).

Acknowledgments

We thank We thank members of Seidemann and Geisler laboratories for their assistances with this project. This work was supported by NIH grants EY-016454 to E.S., EY-024662 to W.S.G. and E.S., BRAIN U01-NS099720 to E.S. and W.S.G., and DARPA-NESD0-N66001-17-C-4012 to E.S.

Figure 1 – Target-background similarity masking and behavioral task. (A) Low contrast orientated
targets can be easily detected on uniform background. (B) Similarity masking is induced by an orientated
background. The same additive target from (A) becomes hard to detect when target orientation
matches background orientation. (See also perceptual demonstration in the Supplementary
Information.) (C-E) Orientation masking was assessed in two awake behaving macaque monkeys
performing a target detection task. Monkey commence the task by fixating at the small bright square
(C). A few moments later, a 4° raise-cosine-masked background grating was flashed at ~3° eccentricity
for target detection (D). In 50% of the trials, a small additive horizontal Gabor target was also added to
the background (E). The monkey indicated the presence of the target by making a saccade to the target
location, and indicated target absent by maintaining gaze at the fixation point. The Gabor target was
always the same – a cosine centered, horizontal Gabor at 4cpd on 0.33° FWHM envelope. The
background grating was also cosine-centered at 4cpd such that the background completely aligned with
the target when they were the same orientation (as in B). Orientation of the grating ranged from 0° to
90° with respect to the Gabor target and was randomized between trials. Bg – background; TBg – target
plus background.

Figure 2 – Behavioral effect of target-background similarity masking. (A) Target detection performance in monkeys was affected by the orientation of the background grating over a variety of target contrast (T## = ##% contrast target) and background contrast levels (Bg## = ##% contrast background). Signal detection measure d-prime (d') of the target is plotted for uniform background (dotted lines), and for each background orientation (markers). In most cases there was a general performance reduction from uniform background to a grating background. Additionally, performance was further reduced when the background orientation was more aligned to the target (0°). A fitted Gaussian (solid line) illustrates the performance change due to orientation masking. d' was calculated from the hit rate (correctly identify target present) and the false alarm rate (incorrectly identify the target when it was absent). The relationship between d' and optimum performance level in percent correct is plotted in (B). (C) Reaction time – calculated from stimulus onset to saccade initiation for Hit trials – is plotted for uniform background (dotted lines) and for each background orientation (solid lines). Error bars indicate the standard error of the mean. Data were pooled within each monkey across experiments. Each experiment contains a single combination of target and background contrast levels, with uniform background and orientated background trials assessed in separate blocks.

Figure 3 – VSD response and decoder schematic. (A) A cranial chamber and a transparent artificial dura provided chronic imaging access to V1 of the monkey. (B) Example recording of the voltage-sensitive dye (VSD) response in an 8x8mm imaging region-of-interest (ROI). Response to the Garbor target on uniform background could be easily identified in VSD (top row). The visual span of the background grating extended beyond the coverage of the imaging window, evoking an encompassing response over the entire ROI (2nd row). When the background was presented with the additive target, the response to the target was diminished (3rd and 4th rows). Each VSD response map were averaged across 50 trials, over 5 frames captured at 100Hz. T only – target only; Bg – background only; TBg – target & background; ΔTBg – target & background minor background only. (C) Target response was extracted at the retinotopic scale by estimating its response profile with a two-dimensional Gaussian. The profile was estimated from response from a separate recording block on each experiment day. To optimized signal-to-noise, in this recording block, the target was flashed repeatedly at 5Hz while the monkey maintained fixation. The effect of spatially correlated VSD noise was minimized by estimating a whitening kernel from trials without stimulus presentation (see Methods). (D) Target response was extract at the columnar scale by estimating the orientation map within the imaging area. This was constructed from full-field gratings flashed at 5Hz in a separate recording block on each experiment day (see Methods). The columnar map in the 0°-90° axis was extracted and windowed down to the retinotopic profile to identify the columnar scale response of the target.

Figure 4 – Retinotopic template dynamics and correlation to behavior. (A-F) Average response and
dynamics from recordings with 12% target contrast (T12) and 7% background contrast (Bg7). (A)
Response time course from stimulus onset (t=0ms) for background only trials. Background orientation
are identified by color. Backgrounds with the same clockwise and anticlockwise orientation disparity
from the target were pooled. (B) Response time course with the same additive target on different
oriented background. Response of the target on uniform gray background is illustrated in black. (C) The
target-evoked response time course was obtained by subtracting background only response (A) from
response to target & background (B). Target evoked response was initially strongest for backgrounds
close to 0° (red), then inverted around t=100ms such that response became the stronger for background
closer to 90° (blue). (D) Response was averaged over 50-200ms and fitted with a Gaussian to illustrate
the change in response magnitude with respect to background orientation. Here, response to clockwise
and anti-clockwise background orientations are plotted separately. Size of markers indicate the number
of trials tested for each orientation. (E) The animals' behavior performance was anti-correlated with the
initial phase of the retinotopic response, and was more aligned in the latter phase. Correlation
coefficient was calculated across background orientations between each frame of the retinotopic

response in (C) against the overall behavior performance in (F). Red dots indicate frames reaching

statistical significance ($p < 0.05$, t-test for correlation coefficient, see Methods). The vertical offset of the

gray horizontal line indicates the correlation between the integrated response (over the time interval

shown in (C)) and the behavior. (F) Behavior performance in d' was calculated as described in Figure 2.

Size of markers indicate the number of trials tested for each orientation. Data was pooled across 8

experiments from both monkeys (see Table 1). (G-L) Same as (A-F) for recordings with 24% target

contrast (T24) and 12% background contrast (Bg12). Similar trends were observed.

Figure 5 – Columnar template dynamics and correlation to behavior. Same format as Figure 4 with the
response examined at the columnar scale. The biphasic response time course observed in the
retinotopic scale was more pronounced at the columnar scale. (A-F) Averaged response and dynamics
from recordings with 12% target contrast (T12) and 7% background contrast (Bg7). (G-L) Averaged
response and dynamics from recordings with 24% target contrast (T24) and 12% background contrast
(Bg12). Here positive response represents relatively stronger activation of the neurons tuned to the
target orientation (0°), and negative response represent stronger activation for neurons tuned to the
orthogonal orientation (90°). Data pooling and counts are the same as reported in Figure 4.

Figure 6 – Columnar orientation estimation by populations tuning. (A) The orientation map obtained
for each experiment as described in Figure 3 was windowed to the retinotopic profile of the target. (B)
Each pixel was assigned to one of 12 equally spaced orientation selective cluster maps by its preferred
orientation. (C) The orientation selective decomposition of VSD response. To a grating stimulus oriented
at 0°, the population tuning curve peaks at 0° (solid curve); likewise, the population peak would shift to
45° for a 45° grating (dotted curve). Note that this population response only represents the relative
difference in preferred orientation (balanced positive and negative values); the overall neural response
offset (retinotopic response) is not captured by this approach. (D) Example of a full population response
time course from stimulus onset (t=0ms). (E) Population response can be summed to a complex vector
representing the overall population tuning orientation and magnitude.

Figure 7 – Dynamics of orientation population response. (A) Population tuning time course for trials
with 12% contrast targets (T12) and 7% contrast backgrounds (Bg7). Averaged response time courses
are presented as a heatmap with the y-axis representing the preferred orientation and the x-axis time
(see key on bottom right) to illustrate the change in the tuning over time. *Row 1:* Heatmaps for
background only trials exhibit clear population tuning in the orientation of the background grating (red
horizontal line). *Row 2:* Heatmaps for background with additive target showing population response
dominated by the background orientation rather than the target orientation (0°). *Row 3:* The target
evoked response is obtained by subtracting the background only response Row 1 from the target &
background response in Row 2. Masking of the target evoked response was strong for backgrounds
oriented near the target orientation (0°). With the background orthogonal to the target, population
tuning in the target orientation can be identified. White line identifies the orientation of the population
vector (peak tuning) wherever the normalized amplitude of the vector average was great than 0.2 (see
Methods). Depending on background orientation, peak tuning appears to be offset from the orientation
of the target (e.g. at Bg -45°). *Row 4:* Heatmap for the target only trials demonstrated clear population
tuning in the target orientation (0°, green horizontal line). (B-E) Averaged response in (A) represented as
a population vector form and illustrated as a continuous trajectory for each background orientation
(color coded). (B) Population tuning trajectory for background only trials. The trajectories commenced in

the center of the circle (white dot) and adhered closely to the orientation of the background. Dot on
each trajectory indicates the position of the population tuning vector at 100 ms. (C) Population tuning
trajectory for background with additive target illustrating the biphasic response of this combined
stimulus. In the early phase, the heading of the trajectory was a mixture of the background and target
(0°) orientations, dominated more by the background. In the late phase, the trajectory made a sharp
turn (t≈100ms) such that trajectories appeared to head towards a convergent point on the positive x-
axis. (D) The trajectory for the target evoked response, calculated by subtracting the background only
response (B) from the corresponding background & target (C). The target evoked response was weak
and noisy, but was heading in the general direction of the target orientation (0°). (E) The population
tuning trajectory for the target only trials illustrating clear tuning in the target orientation (0°). (F-J)
Same as (A-E) for trials with 24% contrast targets (T24) and 12% contrast backgrounds (Bg12). Data was
pooled and averaged across both monkeys.

Figure 8 – Delayed normalization model qualitatively captures key orientation masking response features. (A) Schematic of the normalization model showing the visual input being processed by separate excitatory and normalization signal pathways. The normalization pathway in particular was modelled with a slightly delayed temporal kinetics and wider orientation tuning curves. The excitatory signal undergoes divisive normalization prior to neural output. (B-D) Modelled columnar response output. (B) Modelled response of oriented backgrounds as in Figure 5A. (C) Modelled response of background with additive target as in Figure 5B, and the modelled response of the small Gabor target in black. (D) The target evoked response by subtracting the background only response (B) from (C), matching the biphasic observation in Figure 5C. (E) Correlation of modelled behavioral performance against each time frame of the modelled response, illustrating the early phase where the response was negatively correlated to behavioral choice, and the late phase with positive correlations. (F-I) Modelled time course of the population tuning vector. (F) Modelled populating tuning trajectory of the background only stimuli (color coded) as in Figure 7B. (G) Modelled populating tuning trajectory of the background with additive target illustrating the turn towards a convergent point on the x-axis as in Figure 7C. (H) Modelled target evoked response trajectory from subtracting (F) from (G). (I) Modelled populating tuning trajectory of the target only trials as in Figure 7E.

	Target Contrast	Background Contrast	# Experiments	# Total Trials
Monkey H	T12%	---	5	52
	T12%	Bg7%	3	593
	T12%	Bg12%	2	397
Monkey T	T12%	---	11	490
	T12%	Bg7%	5	1660
	T12%	Bg12%	6	1748
	T24%	---	15	546
	T24%	Bg7%	5	1012
	T24%	Bg12%	6	1320
	T24%	Bg24%	4	388

Table 1 – Experiment summary. Experiment counts and the total number of trials included in the analysis presented in Figures 2, 4, 5, 7 and Supplementary Figures S1 and S2.

Figure S1 – Retinotopic template dynamics and correlation to behavior for all combinatory
background and target contrast levels. Retinotopic response time course using doubly-whitened
retinotopic decoding template illustrated in Figure 3. (A) Response time course from stimulus onset
(t=0ms), grouped by different combinations of target and background contrast levels. Background
orientation are identified by color. Backgrounds with the same clockwise and anticlockwise orientation
disparity from the target were pooled. Response of the target on uniform gray background is illustrated
in black. (B) The target-evoked response time course was obtained by subtracting the background only
response (A, top row) from the background with target response (rest of A) from the same experiments.
(C) Response was averaged over 50-200ms and fitted with a Gaussian to illustrate the change in
response magnitude with respect to background orientation. Here, response to clockwise and anti-
clockwise background orientations are plotted separately. Size of markers indicate the number of trials
tested for each orientation. (D) Correlation coefficient was calculated across background orientations
between each frame of the retinotopic response in (B) against the overall behavior performance for
each combinatory background and target contrasts (Figure 2A pooled across monkeys). Red dots
indicate frames reaching statistical significance ($p < 0.05$, t-test for correlation coefficient, see Methods).
The vertical offset of the gray horizontal line indicates the correlation between the integrated response
(50-200ms) and the behavior. Data was pooled across experiments from both monkeys (see Table 1).

Figure S2 – Columnar template dynamics and correlation to behavior for all combinatory background and target contrast levels. (A-D) Same format as Figure S1 with the averaged response examined at the columnar scale. The columnar template extracts the relative strength between neural activity aligned and orthogonal to the target orientation. Positive response represents relatively stronger activation of the neurons tuned to the target orientation (0°), and negative response represent stronger activation for neurons tuned to the orthogonal orientation (90°). Data pooling and counts are the same as reported in Figure S1.

Methods

All procedures have been approved by the University of Texas Institutional Animal Care and Use Committee and conform to NIH standards.

Widefield Voltage-Sensitive Dye Imaging

The experimental technique for widefield voltage-sensitive dye (VSD) imaging of neural response in awake, behaving macaques was adapted from previous studies (Bai et al 2021, Chen et al 2006, Chen et al 2008, Chen et al 2012). Briefly, two adult male macaque monkeys (Monkey H and Monkey T) were implanted with a metal head post and metal recording chambers located over the dorsal portion of V1, a region representing the lower contralateral visual field at eccentricities of 2–5°. Craniotomy and durotomy were performed. A transparent artificial dura made of silicone was used to protect the brain while allowing optical access for imaging (Arieli et al 2002)(Fig. 3A). Experiments were conducted in the left hemisphere chamber of Monkey H, and in both left and right hemisphere chambers of Monkey T.

VSD imaging was used to record neural population activity at a high resolution in space and time (Shoham et al 1999). Before each experiment, VSD (RH1691 or RH1838) was topically applied on the cortex for ~2 hours to allow the VSD molecules to bind to cellular membranes. In Monkey H, fluorescence from neural activity was recorded using Imager 3001 (Optical Imaging, Inc.) using a tungsten-halogen light source (Zeiss). An infrared eye-tracker (Dr Bouis Inc.) was used to monitor eye position. In Monkey T, florescence was recorded using custom Matlab software interfaced to PCO Edge 4.2 sCMOS camera (Excelitas PCO GmbH) using X-Cite 110LED light source (Excelitas Technologies Corp). Eye position was monitored using an Eyelink 100 Plus video eye-tracker (SR Research Ltd).

Both imaging systems were interfaced to a double-SLR-lens-macro system with housing for dichroic mirrors in between the two SLR lenses. The combination of a 50 mm fixed-focus objective lens (cortex end, Nikkor 50mm f/1.2) and an 85 mm fixed-focused (Canon EF 85mm f/1.2L USM) camera lens provided 1.7x magnification, corresponding to imaging approximately an 8 x 8 mm^2 area of the cortex. Fluorescence signals were measured through a dichroic mirror (650-nm long-pass filter) and an emission filter (RG 665). VSD molecules were excited by light at 630 nm. Imaging data were collected at 512 × 512 resolution at 100 Hz. Data acquisition was time locked to the animal's heartbeat (EKG QR up-stroke, HP Patient Monitor HP78352C). More details about optical imaging with VSD in behaving monkeys are described elsewhere (Bai et al 2021, Chen et al 2006, Chen et al 2008, Chen et al 2012).

Behavioral Task with Optical Stimulation

Monkeys were trained to detect a small additive horizontal Gabor target (4cpd, with σ = 0.14°, 0.33° FWHM envelope) centered on a sinusoidal grating background mask of the same spatial frequency (4° raised-cosine windowed). The background grating was oriented at 0°, ±15°, ±30°, ±45°, ±60, and 90° from the Gabor target orientation. Both the Gabor and background grating were bright-centered – that is, the 0° orientation background was completely in phase with the target. The contrast of the target and background were varied in combinations of levels, reported in Michelson contrast:

$$c = \frac{L_{max} - L_{min}}{L_{max} + L_{min}} = \frac{L_{max} - L_{background}}{L_{background}}$$

For each experiment, a Fixation recording block and a Detection recording block were made using the same target and background conditions. In both blocks, the target and background were centered at 1.6°

to 3° eccentricity from the fixation point and -40° to -65° (from the right horizontal meridian) in the visual
hemifield corresponding to the working cortical chamber. At these coordinates, the spatial extent of the
target was fully imaged through the cortical window, and the larger oriented background uniformly
activated the entire imaging area.

In the Fixation block, the monkeys were required to remain fixated for each imaging trial while either the
target or full-field sinusoidal gratings at 100% contrast were flashed at 5Hz (60ms ON, 140ms OFF) for 1.0s.
These recordings were processed to obtain retinotopic and columnar orientation response maps that
were used to decode the detection recording responses from the same experiment day (Fig. 3C-D).

In the Detection recordings blocks, a background with random orientation appeared on every trial, with a
50% chance of an accompanying Gabor target (Fig. 1C-E). The monkeys were tasked to report the presence
of the target. Each trial began with fixation on a bright 0.1° square. An auditory tone and the dimming of
the fixation square cued the monkey to the start of the detection task trial. 250ms later, the background
with or without the target was presented. The monkeys were trained to maintain gaze at the fixation cue
on target absent trials or saccade to and hold gaze (for 150ms) at the target position to indicate target
detection (with a 75ms minimum allowed reaction time). When the target was present, it remained on
screen for a maximum of 250ms or was extinguished immediately upon the monkeys' saccade initiation.
The monkey was given 600ms to make the saccade or to hold fixation and was subsequently rewarded on
correct choices: stay (correct reject) on target absent trials, or saccade to target (hit) on target present
trials. The target and background contrast level were fixed for each recording block. The probability of
each orientated background and target presence were balanced for each recording block.

A separate target only Detection block on uniform gray background was also taken on each experiment
day using the same routine as above. This block was later used as the reference data to normalize
response amplitudes across experiment days.

Experiments were conducted with custom code using TEMPO real-time control system (Reflective
Computing). The visual stimulus was presented on a Sony CRT (1024x768 @ 100 Hz), distanced 108 cm
from the animal (50 pixels-per-degree), with mean luminance 50 cd/m². The visual stimulus was generated
using in-house real-time graphics software (glib).

<u>Behavior Performance and Reaction Time</u>

Behavior performance was calculated for each background orientation and reported in detection
sensitivity index d' (d-prime). D-prime and criterion were estimated as:

594
$$d' = \Phi^{-1}\big(P(HIT)\big) - \Phi^{-1}\big(P(FA)\big), \qquad criterion = -\frac{\Phi^{-1}(P(HIT)) + \Phi^{-1}(P(FA))}{2}$$

where $\Phi^{-1}(x)$ is the inverse transform of the cumulative normal distribution; and $P(HIT)$ and $P(FA)$
represent the proportion of hits and false alarms respectively. To avoid leaving out the data in some
conditions (e.g., when there are no false alarms), we scaled all the proportions to be between 0.005 and
0.995 $(\widehat{P}(x) = 0.005 + 0.99 \cdot P(x))$.

Mapping between the unbiased percentage correct response and d' is as follows and is depicted in Figure
2B:

601
$$d' = 2 \cdot \Phi^{-1}(PC_{max}); \qquad PC_{max} = \Phi\left(\frac{d'}{2}\right)$$

602   D-prime performance across orientations was fitted with an inverted, dc-shifted Gaussian:

603
$$y = Ae^{-\frac{\theta^2}{2\sigma^2}} + c$$

The reaction times of the animals were calculated from the stimulus onset to the onset of saccade.
Consequently, there was no measurement of reaction on trials where the animals remain fixated at the
fixation point.

<u>VSD Imaging</u>

For each trial, an image sequence was captured for a total of 1.2 seconds including pre-stimulus and post-
stimulus frames. The image sequence was analyzed to extract the response using a variant of the previous
reported routines (Bai et al 2021, Chen et al 2006, Chen et al 2008, Chen et al 2012).

Image stabilization was introduced as the first stage of pre-processing to de-accentuate blood vessel
edges in the ΔF/F response map caused by micro movements of the camera and/or the cortex during
imaging. The image intensity across time at each individual pixel was modelled with separable motion-
free ($I_{\mathbf{x}0}(t)$) and motion-related ($\vec{\alpha}_{\mathbf{x}}. \vec{v}(t)$) components as follows:

$$I_{\mathbf{x}}(t) = I_{\mathbf{x}0}(t) + \vec{\alpha}_{\mathbf{x}}\vec{v}(t)$$

For each trial, a single global motion vector $\vec{v}(t)$ was obtained by estimating the translational motion of
the center portion of the images (1/4 of the imaging area). The motion coefficients $\vec{\alpha}_{\mathbf{x}}$ for each pixel was
then obtained using least squares fitting to the model. The motion-corrected image is $I_{\mathbf{x}0}(t)$. This
approach to image stabilization (compared to traditional image registration approach) has the advantage
of correcting for non-rigid movements (rotations, expansion/contractions, affine transformation, local
distortions, etc.) and sub-pixel motion.

<u>Retinotopic and Columnar Template Decoding</u>

Template decoding was used to summarize the retinotopic and columnar response for each image frame.
The retinotopic response map of the target and columnar orientation map of the imaging area were
estimated from the Fixation recording blocks, in which response were stimulated with visual presentation
at 5Hz. The preprocessing steps for the Fixation recording blocks were: image stabilization, 5Hz FFT
response extraction, ΔF/F normalization, then down-sampling to 128x128 pixels (from 512x512). Baseline
florescence (F_0) was estimated by the average florescence over frames -80 to 0ms relative to stimulus
onset. ΔF/F was calculated as:

$$\frac{\Delta F}{F}(t) = \frac{F(t) - F_0}{F_0}$$

The retinotopic response map of the Gabor target ($H_{ret}(\mathbf{x})$) was estimated by fitting a 2D Gaussian over
the 5Hz flashing Gabor amplitude response. The full-ROI (8x8mm^2 imaging area) columnar orientation
response maps ($H_{ori}(\mathbf{x})$) were estimated from the flashing full-field grating, where the 5Hz FFT grating
response amplitudes were bandpass filtered from 0.8 to 3.0 cycles/mm and the orientation tuning of each
pixel was estimated as described previously (Chen et al 2012). Subsequently, the full-ROI orientation map
was windowed by the retinotopic map to co-localize the retinotopic and the columnar decoders. The
columnar map comprised of pixelwise response magnitude ($A(\mathbf{x})$) and tuning angle ($\theta(\mathbf{x})$), represented
in modified Euler's form:

$$H_{ori}(\mathbf{x}) = H_{ret}(\mathbf{x})A(\mathbf{x})e^{2i.\theta(\mathbf{x})}$$

These maps served as templates for decoding the retinotopic and columnar response from the Detection
blocks. To further reduce effect of motion artefacts, a pixel-wise reliability weighted approach was
adopted. The Detection blocks was preprocessed with image stabilization, down-sampling to 128x128,
and ΔF/F normalization as above. From the pre-processed images, the retinotopic scale variance $\sigma_{ret}^2(\mathbf{x})$
and columnar scale variance $\sigma_{col}^2(\mathbf{x})$ were calculated to be used as reliability weights. The variance was
obtained from condition-mean subtracted residuals taken across all frames across trials, with the ΔF/F
response ($R_{ret}(\mathbf{x},t)$) bandpass filtered between 0.8 to 3.0 cycles/mm ($R_{col}(\mathbf{x},t)$) for the columnar scale
variance. Reliability weighting was implemented by normalizing each template pixel by the corresponding
pixel-wise variance.

649
$$\widehat{H}(\mathbf{x}) = \frac{H(\mathbf{x})}{\sigma^2(\mathbf{x})}$$

Two different columnar decoding methods were employed. The first examined the overall response
aligned to the orientation of the Gabor (0° orientation tuning axis). A second columnar decoding scheme
was employed to examine the full orientation population response. This second scheme comprised of 12
decoders that evenly partitioned the orientation space, such that each decoder contained the column
response magnitude in a subset of pixels tuned with ±7.5° centered at -75° to 90° in 15° steps. Each
decoder therefore represents the population response of similarly tuned neural ensembles spanning the
orientation space every 15°. Each decoder was normalized by the summed response magnitude of its pixel
subset; in this way, each sub-population response is equally represented in the population tuning curve.

The formulation for the reliability weighted templates and the decoding is summarized below for the
retinotopic response time course ($r_{ret}(t)$), columnar response time course ($r_{090}(t)$), columnar population
tuning time course ($r_\theta(t)$).

661
$$r_{ret}(t) = R_{ret}(\mathbf{x},t)\frac{\widehat{H}_{ret}(\mathbf{x})}{\|\widehat{H}_{ret}\|}, \qquad \widehat{H}_{ret}(\mathbf{x}) = \frac{H_{ret}(\mathbf{x})}{\sigma_{ret}^2(\mathbf{x})}$$

662
$$r_{090}(t) = R_{col}(\mathbf{x},t)\frac{\widehat{H}_{090}(\mathbf{x})}{\|\widehat{H}_{090}\|}, \qquad \widehat{H}_{090}(\mathbf{x}) = \frac{real(H_{ori}(\mathbf{x}))}{\sigma_{col}^2(\mathbf{x})}$$

663
$$r_\theta(t) = R_{col}(\mathbf{x},t)\frac{\widehat{H}_\theta(\mathbf{x})}{\|\widehat{H}_\theta\|}, \qquad \widehat{H}_\theta(\mathbf{x}) = \frac{H_\theta(\mathbf{x})}{\sigma_{col}^2(\mathbf{x})}, \qquad H_\theta(\mathbf{x}) = \frac{|H_{ori}(\mathbf{x}|\theta)|}{\sum_\mathbf{x}|H_{ori}(\mathbf{x}|\theta)|}$$

Response Pooling across Experiments

Across experiments, the varying effectiveness of VSD staining led to large variations in the noise level and
amplitude of the ΔF/F response. For pooling data across experiments, response amplitudes were
normalized based on singular value decomposition (SVD). Here, the decoded response of the reference
target only Detection block was used. These were trials of varying target contrast levels on the uniform,
mean luminance background instead of the oriented grating background, collected on the same day. The
assumption is that the neural response amplitude and dynamics to the target should be the same
irrespective of the experiment day. Retinotopic and columnar decoder responses for the target only block
were calculated as with the background detection block as described above. Target contrast response for
each experiment were interpolated so that all assessed target contrast levels across experiments were
represented, then SVD was performed over the frames -100 to 200ms about the stimulus onset across all
experiments. In this way, the first component of the SVD (SVD1) represented the average neural dynamics
of the target contrast response, and the SVD1 coefficients represent the magnitude of this target contrast
response each experiment day. Responses from each experiment was then scaled to match the

experiment with the largest SVD1 coefficient. This was performed for each decoded response separately.
When pooling data for the mean and standard deviation, the inverse of the scaling factor squared was
used as the reliability weighting.

$$\bar{r} = \frac{1}{n}\sum_{exp,trial} w_{exp} r_{exp,trial}, \qquad \sigma_r = \sqrt{\frac{1}{n}\sum_{exp,trial}\left(w_{exp} r_{exp,trial} - \bar{r}\right)^2}$$

$$w_{exp} = \frac{1}{scaler_{exp}^2} = \frac{1}{\left(SVD1_{max}/SVD1_{exp}\right)^2}$$

Lastly, responses are expressed in a modified z-score, obtained by normalizing the response by its
standard deviation. To calculate this standard deviation, responses were grouped by presented stimuli,
and the means of each group was subtracted. The residuals from the mean-subtraction from frames 50
to 250ms post stimulus onset was pooled according to the aforementioned experiment reliability weights
w_{exp} to obtain the response standard deviation.

Response beyond the saccade may contain unwanted signals. Frames beyond the reaction time for each
trial were therefore omitted from summary statistics. For the integrating response within trials, responses
were averaged up to the frame of saccade. For frame-by-frame averaging across trials for response time
course, trials were dropped out from the averaging beyond their reaction time frame.

<u>Descriptive Trends across Background orientation</u>

Trends were fitted to the normalized VSD response across background orientations (e.g., gray curve in Fig.
4D). VSD responses were first averaged over a specific range of frames, and for illustration only the trends
of the averaged response across orientations were fitted with either a flat line, a Gaussian or a difference
of Gaussians curve.

$$\tilde{r}_{flat} = m\theta + c$$

$$\tilde{r}_{gau} = Ae^{-\frac{\theta}{2\sigma^2}} + c$$

$$\tilde{r}_{dog} = Ae^{-\frac{\theta}{2\sigma_A^2}} - Be^{-\frac{\theta}{2\sigma_B^2}} + c$$

The best fitting trend by the F-test were chosen for display in the figures.

$$\frac{\frac{SS_2 - SS_1}{df_2 - df_1}}{\frac{SS_1}{df_1}} \sim F(df_1, df_2 - df_1)$$

<u>Behavior Correlation</u>

The retinotopic and the columnar response dynamics time courses were correlated against the monkey's
behavior judgement. Pearson's correlation coefficient was estimated between the instantaneous
response in time and overall behavior sensitivity index d'. To account for the different trial counts between
background orientations, a trial count weighted version of the correlation coefficient was adopted:

$$\rho = \frac{cov(x,y;w)}{\sqrt{cov(x,x;w)cov(y,y;w)}},$$

$$cov(x,y;w) = \frac{\sum\left(w_i(x_i - m(x;w))\right)\left(w_i(y_i - m(y;w))\right)}{\sum w_i}, \qquad m(x;w) = \frac{\sum w_i x_i}{\sum w_i}$$

The p-value for the weighted coefficients were estimated using the standard t-score replacing the degree
of freedom with an entropy based effective estimate from the weights.

$$t_\rho(\rho; w) = \frac{\rho\sqrt{n_{eff}-2}}{\sqrt{1-\rho^2}} \quad \sim tcdf(n_{eff} - 2)$$

$$n_{eff} = exp(H) = exp(\sum n_{wi} ln(n_{wi})), \qquad n_{wi} = w_i / \sum w_i$$

<u>Normalization model of orientation masking dynamics</u>

A simple model of the neuronal population response with divisive normalization described our results
qualitatively. In this model, orientation columnar response was tuned to one of 12 different orientations:
-75° to 90° in 15° increments. The responses of each orientation column were specified by the simple
normalization model summarized in Figure 8. The input stimulus is specified by the contrast and
orientation of the target, the contrast and orientation of the background, and the duration of the stimulus:
$\mathbf{s} = (C_T, \theta_T, C_B, \theta_B, D)$. In the model, the input stimulus generates and excitation signal $r_e(t|\mathbf{s}, \theta_{max})$ that
is linear with stimulus contrast and a normalization signal $r_n(t|\mathbf{s}, \theta_{max})$ that is also linear with stimulus
contrast. These excitation and normalization signals are controlled by the excitation and normalization
parameters, $\mathbf{\Omega}_e$ and $\mathbf{\Omega}_n$, described below. The normalization signal is then combined with a
normalization constant r_0 to obtain the normalization factor. The normalization constant limits how small
the normalization factor can become. The normalized response is obtained by dividing the excitation
signal by the normalization factor. The final response is then obtained by applying a response exponent
p, which is similar to applying a spiking nonlinearity:

$$r(t|\mathbf{s}, \theta_{max}) = \left[\frac{r_e(t|\mathbf{s}, \theta_{max})}{normalization\ factor}\right]^p = \left[\frac{r_e(t|\mathbf{s}, \theta_{max})}{\sqrt[p]{r_n^p(t|\mathbf{s}, \theta_{max}) + r_0^p}}\right]^p = \frac{r_e^p(t|\mathbf{s}, \theta_{max})}{r_n^p(t|\mathbf{s}, \theta_{max}) + r_0^p}$$

In Figure 8, we show the final responses corresponding to the center of the stimulus $\mathbf{x}_0$. The excitatory
response at that location for background, and target plus background, is obtained by convolving the
effective input contrast signals with the spatial-temporal impulse-response function and evaluating at $\mathbf{x}_0$

$$r_e(t|\mathbf{s}, \theta_{max}) = \begin{cases} (c_{Be} * h_e)(\mathbf{x}_0, t) \\ (c_{TBe} * h_e)(\mathbf{x}_0, t) \end{cases}$$

where $c_{Be}(\mathbf{x}, t)$ and $c_{TBe}(\mathbf{x}, t)$ are the effective input contrast signals, and $h_e(\mathbf{x}, t)$ is the spatiotemporal
impulse response function. The effective excitatory contrast of the background for the orientation channel
with preferred orientation θ_{max} is given by

$$c_{Be}(\mathbf{x}, t) = C_B(\mathbf{x}) exp\left(-\frac{(\theta_B - \theta_{max})^2}{2\sigma_e^2}\right) w(t; D)$$

where $C_B(\mathbf{x})$ is the background contrast, σ_e is the falloff parameter of column's orientation turning
function, and $w(t; D)$ is a temporal pulse function of width D. Similarly, the effective target contrast is
given by

740
$$c_{Te}(\mathbf{x},t) = C_T(\mathbf{x}) \exp\left(-\frac{(\theta_T - \theta_{\max})^2}{2\sigma_e^2}\right) w(t;D)$$

741   The effective excitatory contrast for target plus background is slightly more complicated because there
742   can be some contrast summation or cancellation depending on the phase and orientation of the target
743   relative to the background

744
$$c_{TBe}(\mathbf{x},t) = \lambda(\mathbf{x})\sqrt{c_{Be}^2(\mathbf{x},t) + c_{Te}^2(\mathbf{x},t)}$$

745   where $\lambda(\mathbf{x})$ is the contrast correction factor,

746
$$\lambda(\mathbf{x}) = \frac{C_{TB}(\mathbf{x})}{\sqrt{C_B^2(\mathbf{x}) + C_T^2(\mathbf{x})}}$$

747   The spatiotemporal impulse response function is the separable product of a Gaussian distribution and a
748   gamma distribution

749
$$h_e(\mathbf{x},t) = gauss(\mathbf{x};s_e)\, gamma(t;a_e,b_e)$$

where s_e is the standard deviation of the 2D Gaussian and a_e, b_e are the two parameters of a gamma
distribution: $gamma(t;a_e,b_e) = t^{a_e-1} \exp(-t/b_e)$.

The formulas for the normalization signal are the same as for the excitatory signal, except the four
parameters are allowed to differ: $\mathbf{\Omega}_e = (\sigma_e, s_e, a_e, b_e)$, $\mathbf{\Omega}_n = (\sigma_n, s_n, a_n, b_n)$.

Parameter values from known properties of single neurons in primary visual cortex were adopted. The
response exponent p was constrained to 2.0, consistent with single neuron contrast-response functions
(Albrecht & Hamilton 1982, Geisler & Albrecht 1997, Sclar et al 1990). It was assumed that the peak
orientation of the excitatory signal and suppressive normalization signal θ_{max} were the same (Cavanaugh
et al 2002a), but that the orientation bandwidth of the normalization signal was greater than that of the
excitation signal, $\sigma_n > \sigma_e$ (Cavanaugh et al 2002b), and that the spatial pooling region for the
normalization signal was larger than that for the excitation signal, $s_n > s_e$ (Cavanaugh et al 2002a,
Cavanaugh et al 2002b, Levitt & Lund 2002, Sceniak et al 2001). Finally, it was assumed that the temporal
dynamics of the normalization signal, determined by parameters a_n and b_n, are slower than those for the
excitation signal, determined by parameters a_e and b_e (Groen et al 2022, Zhou et al 2019).

The following were the parameter values used in Figure 8:

$\mathbf{\Omega}_e = (\sigma_e = 15°,\, a_e = 9,\, b_e = 8ms,\, s_e = 0.28°)$

$\mathbf{\Omega}_n = (\sigma_n = 20°,\, a_n = 9,\, b_n = 10.64ms,\, s_n = 0.28°)$

All analyses were done using Matlab R2018a.

<u>Statistics</u>

Two animals were examined to verify the consistency of experimental approach and results. Multiple
recordings were made from the same animals. The number of recordings were based on previous
experience; no statistical method was used to predetermine sample size.

Statistical analyses were conducted in Matlab (R2018a).

Data and Code Availability

The data and custom code from this study will be made available upon manuscript acceptance.

References

Albrecht DG, Geisler WS. 1991. Motion Selectivity and the Contrast-Response Function of Simple Cells in
the Visual-Cortex. *Visual Neuroscience* 7: 531-46
Albrecht DG, Hamilton DB. 1982. Striate Cortex of Monkey and Cat - Contrast Response Function.
Journal of Neurophysiology 48: 217-37
Allman J, Miezin F, McGuinness E. 1985. Stimulus specific responses from beyond the classical receptive
field: neurophysiological mechanisms for local-global comparisons in visual neurons. *Annu Rev*
Neurosci 8: 407-30
Angelucci A, Bijanzadeh M, Nurminen L, Federer F, Merlin S, Bressloff PC. 2017. Circuits and Mechanisms
for Surround Modulation in Visual Cortex. *Annu Rev Neurosci* 40: 425-51
Angelucci A, Bressloff PC. 2006. Contribution of feedforward, lateral and feedback connections to the
classical receptive field center and extra-classical receptive field surround of primate V1. *Visual*
Perception, Part 1, Fundamentals of Vision: Low and Mid-Level Processes in Perception 154: 93-
120
Arieli A, Grinvald A, Slovin H. 2002. Dural substitute for long-term imaging of cortical activity in behaving
monkeys and its clinical implications. *Journal of Neuroscience Methods* 114: 119-33
Bai Y, Chen S, Chen Y, Geisler WS, Seidemann E. 2021. Similar masking effects of natural backgrounds on
detection performances in humans, macaques, and macaque-V1 population responses. *J*
Neurophysiol 125: 2125-34
Benvenuti G, Chen Y, Ramakrishnan C, Deisseroth K, Geisler W, Seidemann E. 2018. Scale-Invariant
Visual Capabilities Explained by Topographic Representations of Luminance and Texture in
Primate V1. *Neuron* 100: 1-9
Campbell FW, Kulikowski JJ. 1966. Orientational selectivity of the human visual system. *J Physiol* 187:
437-45
Carandini M, Heeger DJ. 1994. Summation and Division by Neurons in Primate Visual-Cortex. *Science*
264: 1333-36
Cavanaugh JR, Bair W, Movshon JA. 2002a. Nature and interaction of signals from the receptive field
center and surround in macaque V1 neurons. *Journal of Neurophysiology* 88: 2530-46
Cavanaugh JR, Bair W, Movshon JA. 2002b. Selectivity and spatial distribution of signals from the
receptive field surround in macaque V1 neurons. *Journal of Neurophysiology* 88: 2547-56
Chen Y, Geisler WS, Seidemann E. 2006. Optimal decoding of correlated neural population responses in
the primate visual cortex. *Nat Neurosci* 9: 1412-20
Chen Y, Geisler WS, Seidemann E. 2008. Optimal Temporal Decoding of V1 Population Responses in a
Reaction-Time Detection Task. *J Neurophysiol* 99: 1366-79
Chen Y, Palmer CR, Seidemann E. 2012. The relationship between voltage-sensitive dye imaging signals
and spiking activity of neural populations in primate V1. *Journal of Neurophysiology*: 3281-95
Clifford CW. 2014. The tilt illusion: phenomenology and functional implications. *Vision Res* 104: 3-11
Foley JM. 1994. Human luminance pattern-vision mechanisms: masking experiments require a new
model. *Journal of the Optical Society of America. A, Optics, image science, and vision* 11: 1710-9
Frazor RA, Geisler WS. 2006. Local luminance and contrast in natural images. *Vision Res* 46: 1585-98
Geisler WS, Albrecht DG. 1997. Visual cortex neurons in monkeys and cats: Detection, discrimination,
and identification. *Visual Neuroscience* 14: 897-919
Grinvald A, Hildesheim R. 2004. VSDI: A new era in functional imaging of cortical dynamics. *Nature*
Reviews Neuroscience 5: 874-85
Groen IIA, Piantoni G, Montenegro S, Flinker A, Devore S, et al. 2022. Temporal dynamics of neural
responses in human visual cortex. *J Neurosci* 42: 7562-80

Heeger DJ. 1991. Nonlinear model of neural responses in cat visual cortex. *Computational models of*
visual processing: 119-33
Heeger DJ. 1992. Normalization of Cell Responses in Cat Striate Cortex. *Visual Neuroscience* 9: 181-97
Hubel DH, Wiesel TN. 1959. Receptive fields of single neurones in the cat's striate cortex. *The Journal of*
Physiology 148: 574-91
Hubel DH, Wiesel TN. 1968. Receptive fields and functional architecture of monkey striate cortex.
Journal of Physiology 195: 215-43
Levitt JB, Lund JS. 2002. The spatial extent over which neurons in macaque striate cortex pool visual
signals. *Visual Neuroscience* 19: 439-52
Michel MM, Chen Y, Seidemann E, Geisler WS. 2018. Nonlinear Lateral Interactions in V1 Population
Responses Explained by a Contrast Gain Control Model. *J Neurosci* 38: 10069-79
Ozeki H, Finn IM, Schaffer ES, Miller KD, Ferster D. 2009. Inhibitory Stabilization of the Cortical Network
Underlies Visual Surround Suppression. *Neuron* 62: 578-92
Polat U, Mizobe K, Pettet MW, Kasamatsu T, Norcia AM. 1998. Collinear stimuli regulate visual
responses depending on cell's contrast threshold. *Nature* 391: 580-84
Rubin DB, Van Hooser SD, Miller KD. 2015. The Stabilized Supralinear Network: A Unifying Circuit Motif
Underlying Multi-Input Integration in Sensory Cortex. *Neuron* 85: 402-17
Sceniak MP, Hawken MJ, Shapley R. 2001. Visual spatial characterization of macaque V1 neurons. *J*
Neurophysiol 85: 1873-87
Sceniak MP, Ringach DL, Hawken MJ, Shapley R. 1999. Contrast's effect on spatial summation by
macaque V1 neurons. *Nat Neurosci* 2: 733-39
Schwartz O, Hsu A, Dayan P. 2007. Space and time in visual context. *Nat Rev Neurosci* 8: 522-35
Schwartz O, Simoncelli EP. 2001. Natural signal statistics and sensory gain control. *Nature Neuroscience*
4: 819-25
Sclar G, Maunsell JH, Lennie P. 1990. Coding of image contrast in central visual pathways of the macaque
monkey. *Vision research* 30: 1-10
Sebastian S, Abrams J, Geisler WS. 2017. Constrained sampling experiments reveal principles of
detection in natural scenes. *Proc Natl Acad Sci U S A* 114: E5731-e40
Seidemann E, Arieli A, Grinvald A, Slovin H. 2002. Dynamics of depolarization and hyperpolarization in
the frontal cortex and saccade goal. *Science* 295: 862-65
Seidemann E, Geisler WS. 2018. Linking V1 Activity to Behavior. *Annual review of vision science* 4: 287-
310
Shoham D, Glaser DE, Arieli A, Kenet T, Wijnbergen C, et al. 1999. Imaging cortical dynamics at high
spatial and temporal resolution with novel blue voltage-sensitive dyes. *Neuron* 24: 791-802
Sit Y-F, Chen Y, Geisler WS, Miikkulainen R, Seidemann E. 2009. Complex dynamics of V1 population
responses explained by a simple gain-control model. *Neuron* 24: 943-56
Stromeyer CF, 3rd, Julesz B. 1972. Spatial-frequency masking in vision: critical bands and spread of
masking. *Journal of the Optical Society of America* 62: 1221-32
Tsodyks MV, Skaggs WE, Sejnowski TJ, McNaughton BL. 1997. Paradoxical Effects of External Modulation
of Inhibitory Interneurons. *The Journal of Neuroscience* 17: 4382-88
Watson AB, Solomon JA. 1997. Model of visual contrast gain control and pattern masking. *Journal of the*
Optical Society of America. A, Optics, image science, and vision 14: 2379-91
Wenderoth P, Johnstone S. 1987. Possible neural substrates for orientation analysis and perception.
Perception 16: 693-709
Wilson HR, McFarlane DK, Phillips GC. 1983. Spatial-Frequency Tuning of Orientation Selective Units
Estimated by Oblique Masking. *Vision Research* 23: 873-82
Yang Z, Heeger DJ, Seidemann E. 2007. Rapid and precise retinotopic mapping of the visual cortex
obtained by voltage sensitive dye imaging in the behaving monkey. *J Neurophysiol* 98: 1002-14

Zhou J, Benson NC, Kay K, Winawer J. 2019. Predicting neuronal dynamics with a delayed gain control
model. *PLoS Comput Biol* 15: e1007484

875