Experimental procedures for flow cytometry of wild type mouse brain:

A systematic review

Robert C. Sharp[1], Dylan T. Guenther[1]and Matthew J. Farrer[1*]

[1]McKnight Brain Institute, Department of Neurology, University of Florida, Gainesville, FL., USA

[*]**Correspondence:**

Matthew J. Farrer

m.farrer@ufl.edu

Keywords: Neuroimmune, Flow Cytometry, Mouse Studies, Immunity, Systematic Review,

Inflammation, Neurological Disorders, Methodology

Main Text Word Count: 5463

Figures: 7 and Suppl. Figures: 3

Tables: 1 and Suppl. Tables: 3

Abstract

Objective: To systematically review the neuroimmunology literature to determine the average

immune cell counts reported by flow cytometry in wild-type (WT) homogenized mouse brain.

Background: Mouse models of gene dysfunction are widely used to study age-associated

neurodegenerative disorders, including Alzheimer's disease and Parkinson's disease. The importance

of the neuroimmune system in these multifactorial disorders has become increasingly evident, and

methods to quantify resident and infiltrating immune cells in brain, including flow cytometry, are

necessary. However, there appears to be no consensus on the best approach to perform flow

cytometry or quantify/report immune cell counts. The development of more standardized methods

would accelerate neuroimmune discovery/validation by meta-analysis.

Methods: Examination of the PROSPERO registry confirmed a systematic review of

'neuroimmunology' by 'flow cytometry' has yet to be reported. A protocol for a systematic review

was subsequently based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses

(PRISMA) using Studies, Data, Methods and Outcomes (SDMO) criteria. Literature searches were

conducted in Google Scholar and PubMed databases. From that search, 900 candidate studies were

identified and 437 studies were assessed for eligibility based on formal exclusion criteria.

Results: Out of the 437 studies reviewed, 58 were eligible for inclusion and comparative analysis.

Each study assessed immune cell subsets within homogenized mouse brain and used flow cytometry.

Nonetheless, there was considerable variability in methods, data analysis, reporting and results.

Descriptive statistics are presented on study designs and results, including medians with interquartile

ranges (IQRs) and overall means with standard deviations (SD) for specific immune cell counts and

their relative proportions, within and between studies. Results derived from 58 studies reported the

most abundant immune cells within the brains were TMEM119$^+$ microglia, bulk CD4$^+$ T cells, and

bulk CD8$^+$ T cells.

Conclusion: Experiments to conduct and report flow cytometry data, derived from WT homogenized

mouse brains, would benefit from a more standardized approach. While within study comparisons are

valid, variability in methods counts of immune cell populations are too broad for meta-analysis.

Inclusion of a minimal protocol with more detailed methods, controls and standards could enable this

nascent field to compare results across studies.

1   Introduction

The mouse has been used to model neurological disorders for many decades, whether lesion models

(i.e. inducing a stroke in a mouse specimen), using transgenic or more physiologic gene knock-out or

mutant knock-in approaches (1–3). Some illustrations of mouse modeling for neurologic and

neurodegenerative disorders include work in Alzheimer's disease (AD), Parkinson's disease (PD),

Huntington's disease (HD), multiple sclerosis (MS), psychological/intellectual disabilities (i.e.

depression or Down's syndrome), traumatic brain injuries (TBI) and prion diseases (1–3). Most of

these disorders are multifactorial, with genetic and environmental components, for which the immune

system may provide some integration. Consequently, there has been growing interest in

neuroimmunology, and not just focused on resident immune cells within the brain, but also on

infiltrating peripheral immune cells (4–7). Recent studies of gut microbiota have highlighted nervous

and immune communication between the gut and central nervous system (CNS) (8–10). With this

resurgence, researchers have employed traditional (i.e. immunofluorescence/histology slide staining

and Western blotting) and contemporary methods (i.e. single-cell genomics and single-cell

sorting/staining via flow cytometry) to better characterize specific immune cell subsets within the

body and brain.

Specifically, brain resident immune cells, including microglia and astrocytes, have been

comprehensively examined in many mouse models of neurologic disorders (4–10). However, the

characterization of other immune cell subsets in peripheral blood, the CNS, and within the brain

(both residential and infiltrated) has yet to be fully described (4–10). These immune cell subsets

include, but are not limited to, natural killer cells (NK cells), T cells, and B cells (4–10). However,

validation and comparison through meta-analysis might be enabled if researchers utilize more

standardized methods and reporting.

Technologic developments for single-cell isolation and analysis, including single-cell RNAseq, mass

cytometry (CYTOF), fluorescence-activated cell sorting (FACS) and multi-color flow cytometry

derived immunophenotyping, have illuminated a wide variety of scientific fields (11,12). Of these

techniques, FACS and flow cytometry immunophenotyping are most frequently used. Reasons

include the ease of use in setting up a flow cytometer or sorter for a variety of applications, the

sensitivity of the assay, the specificity of the antibodies used, the potential of those antibodies to be

used for both flow cytometry and Western blotting for the same target, and the ability to produce

qualitative and quantitative data (12,13).

Nevertheless, flow cytometry has its pitfalls as analytic interpretation of the data depends on the

user's preference for gating and target choices (13). Experimentally, flow cytometry is also

dependent on the fluorophores and cytometers used, and variation between these instruments may

result in false positives and negatives. Nevertheless, such issues can be circumvented by providing

multiple controls, such as unstained, isotype antibody-stained and fluorescence minus one (FMO)

controls, more rigorously described methods and standardization of flow cytometry protocols.

FACs and flow cytometry immunophenotyping have been insightful and widely used in basic

immunology that other scientific fields, including neurology, have begun to adopt these methods

(14–16). However, the use of flow cytometry for detecting cell types within the brain and CNS

warrants more standardized protocols and reporting, now that their utility has been demonstrated.

Although mouse immune profiles within the brain have been identified by high-dimensional single-

cell mapping using techniques like mass cytometry (CYTOF)(17), at the time of writing and despite

numerous publications, the expected cell counts and proportions of each immune subset within the

brain of a wild type (WT) mouse have not yet been clearly defined by standard FACS and flow

cytometry. Consequently, we have performed a systematic review focused on neuroimmunology and

the use of flow cytometry to detect immune cells derived from WT homogenized mouse brain. In

our results, we summarize the number of immune cells overall and estimate the immune subsets that

can be detected via flow cytometry immunophenotyping. Our findings demonstrate a critical need

for more standardized methods and reporting, and led to best practice recommendations for future

publications.

2   Methods

2.1   Study design

Study design was informed by prior literature (18–20) and based upon the Preferred Reporting Items

for Systematic Reviews and Meta-Analyses (PRISMA) criteria (21) and the Cochrane Handbook of

reporting methodology reviews, employing SDMO (Types of Studies, Types of Data, Types of

Methods, and Types of Outcome Measures) criteria (22,23). Bias assessment for each individual

study selected for systematic review inclusion was also conducted using Systematic Review Centre

for Laboratory animal Experimentation (SYRCLE) (24) methodology, and subsequent reporting uses

the *robvis* R package and Shiny web app (25).

2.2   Search strategy

Google Scholar and PubMed databases were used to identify all studies for this systematic review,

published between Jan 2013 and July 2023. The search protocol and study design were also assessed

within the National Institute of Health (NIH) PROSPERO registry database, which confirmed a

review of this topic has not been previously conducted. For database searches, the following

combination of keywords were used to identify eligible studies: "flow cytometry" AND "immune

subset name examined in study" AND "mouse brain." For example, a keyword search containing

"flow cytometry CD4 T cells mouse brain" was used. "Immune subset name examined in study" is

defined as one of the following immune cell subsets:," "CD4 T cells," "CD8 T cells," "double

negative DN T cells," "regulatory T cells TREG," "follicular helper T cells TFH," "T helper 1 T cells

TH1," "T helper 2 T cells TH2," "T helper 17 T cells TH17," "naïve T cells," "naïve CD4 T cells,"

"naïve CD8 T cells," "naïve-like T cells," "central memory T cells TCM," "central memory CD4 T

cells TCM," "central memory CD8 T cells TCM," "effector memory T cells TEM," "effector

memory CD4 T cells TEM," " effector memory CD8 T cells TEM," "effector memory T cells re-

expressing CD45RA TEMRA," "effector memory CD4 T cells re-expressing CD45RA TEMRA,"

"effector memory CD8 T cells re-expressing CD45RA TEMRA," "TEMRA-like T cells," "natural

killer cells NK cells," "dendritic cells DC," "B cells," "monocytes," "macrophages," "M1

macrophages," "M2 macrophages," "TMEM119 microglia," or "neutrophils."

2.3   Selection and exclusion criteria

Studies were selected from search results employing the following inclusion criteria: 1) any study

performed between 2013-2023; 2) any study that contained flow cytometry data identifying immune

cell subsets and counts in mouse brain; 3) any study that reported total cell numbers or total live cell

percentages for one or more immune cell subsets, and; 4) any study that had WT mice or a non-

treated control (when reporting on transgenic mouse models). Studies were excluded from search

results based upon the following criteria: 1) studies performed in 2012 and prior; 2) any study not

focused on flow cytometry of homogenized mouse brains; 3) rat studies; 4) human studies, or; 5)

studies not reporting controls or the background mouse strain.

2.4   Data extraction

For each study, information on the mouse strain used, age, and sex were recorded. In addition,

methodological information on mouse perfusion, brain extraction and homogenization were recorded.

Flow cytometry methods were recorded when the following was reported: 1) cytometer make and

model; 2) software for data collection/analysis; 3) full gating strategy; 4) total cells collected per

sample; 5) total live cell counts; 6) total immune cell subset percentages calculated directly from live

cell counts, and; 7) methods for determining cell counts and/or mean fluorescence intensity (MFI)

readings.

Studies were subsequently examined for immune cell subset counts from WT/control mouse brains

directly from the main text, supplementary materials, and/or extrapolated from the figures and graphs

presented. In this systematic review, "raw" total immune cell subset counts were reported for each

immune cell subset as a median with an interquartile range (IQR; defined as: 75[th] percentile upper

quartile [Q3] – 25[th] percentile lower quartile [Q1]) and as the combined mean with standard deviation

(SD) of multiple studies (n). The "raw" total overall cell count collected per sample by flow

cytometry was also reported from these studies.

2.5   Data analysis

After the median with IQR and combined means with SD from the "raw" total immune cell subset

counts and from the "raw" total overall cell count collected per sample were recorded from each

study, we standardized an approach to estimate how many immune cells of each subset would be

counted if the total homogenized mouse brain cells collected by flow cytometry equaled a total of 1 x

10^5 collected cells. The value of cells required, per sample, for an accurate flow cytometry reading is

stated to be between 1 x 10^4 total cells (minimum) and 1 x 10^6 total cells (maximum) hence we used

the median (26).

Standardized total cell counts for each immune cell subset reported in this systematic review, were

calculated using the following equation:

153
$$\frac{a}{b} = \frac{X}{1 \ x \ 10^5 \ Total \ Cells \ Collected}$$

154   Where "a" is the [Raw Total Immune Cell Subset Count], "b" is the [Raw Total Cell Count

155   Collected] per sample, where "X" is the [Standardized Total Immune Cell Subset Count].

156   Rearranging to solve for X gives the following equation:

157
$$\frac{(a)(1 \; x \; 10^5 \; Total \; Cells \; Counted)}{b} = X$$

Once all total immune cell subset counts were standardized, we determined the theoretical

standardized percentages of each immune cell subset within the entire WT mouse brain, and report

those results as means with SD.

3   Results

3.1   Literature search and study selection

A PRISMA based flow diagram illustrates our screening methodology for study identification

(**Figure 1**) (21). The two databases (Google Scholar and PubMed) were searched using keywords, as

defined in Methods, and highlighted 900 articles for systematic review. Of those 900 reports, 223

were removed as they supplied only an abstract or were duplicated between databases, and 677

studies remained. Of these an additional 240 studies were removed as they were published in 2012 or

prior. This cut off is arbitrary, but was used to identify more contemporary flow cytometry

immunophenotyping publications and resulted in 437 eligible studies. Further inclusion and

exclusion criteria removed: 133 studies in which flow cytometry of homogenized mouse brains was

not a main focus; 129 studies with insufficient information and/or results from controls or in which

the background mice strain was not specified; 94 human studies, and; 23 rat studies. Consequently,

58 studies that passed our inclusion and exclusion criteria were incorporated in this systematic review

(27–84).

3.2   Reporting of mouse information and perfusion/tissue processing is inconsistent between studies.

Of the 58 studies selected, we reviewed the basic characteristics of the mice strains used (**Figure 2A**, **Suppl. Table 1**). The inbred congenic C57BL/6 mouse line was used most (26/58 studies [44.8%]) as a control and as the background for genetically modified mice (27,28,30,31,36,41–47,49,51,54,58,66–69,72,74–76,79,82). However, other C57BL/6 mice sub-strains were used throughout the 58 studies that include: C57BL/6J (19/58 studies [32.8%]) (29,32,33,39,52,56,60,62–65,70,71,73,78,80,81,83,84); C57/BL6 (2/58 studies [3.44%]) (34,40); C57BL/6 (H-2b) (2/58 studies [3.44%]) (35,53); and C57BL/6J (B6) (2/58 studies [3.44%]) (38,61). The majority of studies that reported mouse sex (**Figure 2B, Suppl. Table 1**) used only male mice (33/58 studies [56.9%]) (29,30,33,34,38,40,41,44,45,47,49,50,54,58–63,65,67–71,74,77–80,82–84), although 10/58 studies [17.2%] reported mixed results of male and female animals together (27,31,32,36,37,39,43,51,72,75). The age of the mice (**Figure 2C, Suppl. Table 1**) within studies varied but the majority reported findings at 8-12 week (2-3 months) (9/58 studies [15.5%]) (29,33,56,57,65,72,75,80,83). However, studies characterized animals over a wide range of age, from 1-2-weeks (32,39,81) and between 3-26 months (37).

Perfusion (**Suppl. Figure 1, Suppl. Table 1**) and brain tissue processing methods (**Figure 3A-C, Suppl. Table 1**) also varied across the 58 studies. A majority of studies (36/58 studies [62.1%], **Suppl. Figure 1, Suppl. Table 1**) used cold PBS for perfusion (27,29,31–33,35,39,43–48,50–58,61,63–67,70,73–76,79,81,84). Of the 58 studies reviewed, 6/58 [10.3%] (**Figure 3A, Suppl. Table 1**) used a commercial kit, such as the Neural Tissue Dissociation Kit P (Miltenyi Biotec), to isolate immune cells from mouse brain (34,40,41,61,67,71). Nevertheless, in neurology it remains unclear how to best process mouse brain to dissociate the tissue and leave cell types intact (85–87).

Researchers may use mechanical homogenization, enzymatic homogenization, or both

homogenization techniques (85–87). For the studies reviewed, 18/58 [31.0%] (**Suppl. Table 1**) used

some type of mechanical homogenization (i.e. Glass-Teflon homogenizer, an 18-gauge needle, etc.)

before filtering through a cell strainer, prior to enzyme treatment and the use of a myelin

removal/immune isolation gradient (i.e. Percoll gradient) (31,33,37,44,48,50,53,55–

57,59,60,63,64,66,69,81,84). The enzymatic solutions used in the reports also vary widely (**Figure**

3B, Suppl. Table 1). Most studies used collagenase (I, II, IV, D, I-S, or liberase) alone or combined

with additional enzymes (30/58 studies [51.7%]) (27–31,33,46–48,53,54,60,62–64,66–68,70,73–

80,82–84). The enzyme most used in combination with collagenase (or another tissue digestion

enzyme) was DNase I (27/58 studies [46.6%]) (27–31,47,48,51,53,54,60,62,63,66,68,70,73–80,82–

84). After homogenization of mouse brain, cell strainers are generally used to remove dead cells and

myelin (**Figure 3A, Suppl. Table 1**), and a 70 μm filter was used in the majority (23/58 studies

[39.7%]) of studies (15,30,32,33,38,39,43,45,46,48,52,53,56,58,59,62–64,67,72,74,82,84). To further

remove myelin from mouse brain homogenate along with isolating immune cells, researchers employ

a myelin removal kit or Percoll gradient solutions. Specific cell types can be isolated while the

myelin layer rises to the top of the sample tube with centrifuge, depending upon the percentage of

Percoll. Again, in the studies reviewed, the percentages of Percoll (**Figure 3C, Suppl. Table 1**)

varied with most using a 30%/70% Percoll gradient solution (17/58 studies [29.3%])

(15,27,29,33,43,45,53,57,62,63,71,72,74,77,79,81,84). Overall, the age of mice and methodology for

isolating immune cell counts from brain for flow cytometry varied greatly between studies.

3.3   Flow cytometry methodology used and cell counts are inconsistently reported between

studies

After considering mice strain and brain homogenization methods, we examined the flow cytometry

instruments used and data reporting (**Figure 4A-C, Suppl. Figures 2A-B and 3A-B, Suppl. Table**

2). The make and model of flow cytometer/FACS sorter (**Suppl. Figure 2A, Suppl. Table 2**) and

analysis software (**Suppl. Figure 2B, Suppl. Table 2**) used in each of studies also varied widely. The

flow cytometer most reported was the BD LSRII Flow Cytometer (12/58 studies [20.7%])

(30,32,33,35,36,39,43,51,60,61,63,65) whereas the FACS sorter was the BD FACS Aria III (8/58

studies [13.8%])(31,38,47,49,54,55,64,83). The analysis software most generally used was a version

of FlowJo (Tree Star) for flow cytometry immunophenotyping (38/58 studies [65.5%]) (27,29–

31,33–36,38,41,43–45,47,49–54,56,58,62,63,65,68–74,76,77,79,81,83,84) and BD FACSDiva

specifically for FACS analysis (11/58 studies [19.0%]) (32,35,37,39,42,45,60–62,76,77).

After reviewing methodology in all 58 studies, we assessed the quality of flow cytometry reporting

(**Figure 4A-C, Suppl. Figure 3A-B, Suppl. Table 2**). For each study, we scored the following

parameters: 1) whether a full gating strategy was reported; 2) the total cells collected per sample for

flow cytometry; 3) the total live cell counts during flow cytometry sample collection; 4) if the total

immune cell subset percentage was calculated directly from the live cell count reported or was it

derived from another gate (i.e. was the immune cell subset percentage reported derived from the

CD45$^+$ gate or from the live/dead gate); and 5) whether the methods used to determine cell counts

and/or MFI readings were documented. Of the 58 studies reviewed, 35/58 [60.3%] included a full

gating strategy (**Figure 4A, Suppl. Table 2**) (27,30,32–34,36,37,41,43–45,47,50,51,54,55,58,60–

64,67,71–79,81,82,84) , while 20/58 [34.5%] reported a partial gating strategy

(28,29,35,38,40,42,46,48,49,52,53,56,57,59,66,68–70,80,83) and 3/58 [5.17%] of studies did not

include this information (31,39,65). When documented, the flow antibody clone and gating strategy

was reported (44/58 studies; 75.9%) (**Suppl. Table 2**) (27,29–33,35–40,42,43,45–55,57,58,60–

62,66–69,74–80,82–84). Most studies used similar clones to identify specific immune cell subsets

(**Supple. Table 2**).

On reporting total cells collected per sample during flow cytometry collection (**Figure 4B, Suppl.**

Table 2), only 16/58 [27.6%] of studies provided this data

(30,36,37,41,44,47,52,57,58,61,63,67,71,73,74,82). Total live cell counts collected during flow

cytometry (**Suppl. Figure 3A, Suppl. Table 2**) was only reported in 9/58 [15.5%] of the studies

reviewed (30,37,57,58,61,67,71,73,74). Out of the 58 selected studies, only 12/58 [20.7%] expressed

their results in terms of total immune cell subset percentages derived from live cells only (**Figure 4C,**

Suppl. Table 2) and not from another gate (such as deriving from the CD45$^+$ gate)

(35,37,43,47,52,58,67,68,71,72,74,76). Lastly, 35/58 [60.3%] of studies (**Suppl. Figure 3B, Suppl.**

Table 2) reported methods on how cell counts and/or MFI readings were recorded

(27,28,30,32,33,35,37,39,41–44,49–52,54,55,57–59,61–64,67,70–77,82). Overall, the

cytometer/FACS sorter used, and the reporting of total cells collected, total live cells and immune

cell subset percentages was not standardized in the 58 studies examined.

3.4   Reported immune cell counts from WT/control homogenized mouse brain is highly

variable between studies.

Across all studies we then examined the total immune cell counts reported in WT/control mouse

models, as detected by flow cytometry immunophenotyping and/or by FACS (**Suppl. Table 3**). As

described in Methods and **Suppl. Table 3**, studies were selected based on their reporting of a wide-

variety of immune cell subsets. These included, T cell subsets/memory T cells (naïve-like, central

memory [T_{CM}], effector memory [T_{EM}] and effector memory re-expressing CD45RA [T_{EMRA}]),

natural killer cells (NK cells), dendritic cells (DCs), B cells, monocytes, macrophages (M1

[predominately CD86$^+$] and M2 [predominately CD163$^+$] phenotypes), TMEM119$^+$ microglia, and

neutrophils. Of note, the M1/M2 nomenclature for macrophages is being updated currently in the

immunology field(88)

Many immune subsets were examined but relatively few were reported in sufficient studies to be able

to calculate representative medians with IQR and means with SD (**Figure 5 and Suppl. Table 3**).

Immune cell subsets with at least two or more references to derive a median and mean cell count for

WT/control mice include: bulk $CD4^+$ T cells (27–33), bulk $CD8^+$ T cells (27,28,30,34–37), double

negative (DN) T cells (33,38), regulatory T cells (T_{REG}) (29,39–42), T helper 1 cells (T_{H1}) (27,40), T

helper 17 cells (T_{H17}) (44,45), NK cells (33,41,46,47,49–51), DCs (33,52–55), B cells (33,56–60),

monocytes (61–67), bulk macrophages (64,65,68–72), $TMEM119^+$ microglia (73–76) and

neutrophils (46,77–84). Immune cell subset counts that were derived from only one study, such as

follicular T cells (T_{FH}) (43) and T_{EM} bulk $CD4^+$ and $CD8^+$ T cells (30), were still included in this

review as a representation of the possible median/average cell count for these subsets.

We first calculated the overall medians for each examined immune subset by collecting all of the

"raw" total immune subset cell counts from each of the 58 studies (**Figure 5**). By doing this, we

discovered that out of these studies, there were some that were outliers (outside of the IQR) that

appear to heavily altered the overall total immune cell counts determined by flow cytometry

(27,33,36,41,47,53,58,61,69,72,76,79). Out of all the immune cell subsets from the 58 studies, the

highest median was $TMEM119^+$ microglia (72,300 [IQR=194,038]; n = 4 studies) (73–76). The

lowest median that was calculated was T_{FH} T cells (8 [IQR=0]; n = 1 study) (43). Memory T cell

subsets found within both $CD4^+$ and $CD8^+$ T cells were not reported as an immune cell subset in any

study. Similarly, T helper 2 cells (T_{H2}) were not reported in any studies.

The 58 studies had highly variable ranges for each immune cell subset found within mouse brain, for

which most of the SD calculated were greater than the means (**Suppl. Table 3**). As with the

calculated medians, the highest mean cell count reported from the 58 studies was TMEM119$^+$

microglia (90,323 ± 104,555; n = 4 studies) (73–76). The lowest mean cell counts reported were for

T$_{FH}$ T cells (8 ± 0; n = 1 study) (43). Subsequently, we also calculated the means of total overall cell

counts, collected per sample, as reported in the 58 studies reviewed (**Suppl. Table 3**). Although not

as variable as immune cell subset counts, the total overall cell counts collected by flow cytometry

ranged from 1 x 10^4 to ~3 x 10^6 cells (**Suppl. Table 3**). In some studies, the actual number of cells

collected by flow cytometry was not specified, but the total cell counts could be extrapolated from

data given in the main text, figures, and/or supplementary material. Overall, we conclude the immune

cell subset counts collected by flow cytometry immunophenotyping are highly variable between

studies, and there is currently insufficient data on specific T cell subsets/memory subsets and specific

macrophage phenotype counts within the WT mouse brain.

3.5   Standardizing immune cell counts and percentages within WT/control mouse brain.

We devised a method to standardize counts of immune cells and subset percentages within

WT/control mouse brain in each study (**Figure 6, Suppl. Table 3**). In flow cytometry, cell type

percentages should be reported as values from the total number of cells collected (on average 1 x 10^5

cells) rather than from a sub-partition within the gating strategy (26,89). Thus, based on the

assumption the total cells collected per sample was 1 x 10^5 cells, we were able to estimate a

standardized total immune cell subset count for the data provided in each study.

From these standardized total immune cell subset counts, we are able to determine the proportion of

immune cell subsets within the brain by simply dividing the standardized count by 1 x 10^5 total cells

collected to obtain percentages (**Figure 6**, **Suppl. Table 3**). The immune subset with the highest total

cell percentages that were found within WT/control mouse brain were TMEM119$^+$ microglia (28.5%

± 33.0). Of non-neural/glial specific immune cells, bulk CD4$^+$ T cells (6.41% ± 6.15) and bulk CD8$^+$

T cells (4.00% ± 5.34), were most often counted within mouse brain compared to other

adaptive/innate immune cells. Overall, we were able to calculate the average percentage of immune

cells found within WT/control mouse brain from the 58 selected studies. Hence, we are able to report

a more reliable estimate of the immune cell composition within mouse brain despite their wide SD.

3.6   Evaluating risk of bias of all included studies

As per PRISMA and Cochrane criteria for systematic reviews, it is important to evaluate the risk of

bias for all studies cited (20,21,24,90). Here we utilize the SYRCLE's risk of bias tools for animal

studies (24) to create a summary graph and "stop-light" figure (**Figure 7**) to highlight the overall bias

of each study assessed within the following domains: D1: Sequence Generation (randomization

methods used to choose animals for comparable groups); D2: Baseline Characteristics (full

description of animal characteristics from all comparable groups); D3: Allocation Concealment

(methods used to conceal how animals are distributed to researchers, i.e. using a coding method for

each animal) ; D4: Random Housing (housing all animal groups randomly within the animal room);

D5: Blinding (blinding methods used on researchers, such as blinding the knowledge of intervention

or transgenic model used and blinding the outcome assessors); D6: Random Outcome Assessment

(methods on how the animals were selected at random for outcome assessment); D7: Incomplete

Outcome Data (description of the completeness of the data outcome, i.e., stating if data was excluded

or if animals were removed from study at any point); D8: Selective Outcome Reporting (the

completeness of the study protocols); D9: Other Sources of Bias (examples include: confounders,

contamination problems, analysis errors, design-specific risk of bias, etc.).

The overall bias of all selected studies was deemed predominantly "unclear" (~over 50%) due to lack

of reporting on specific data/methodology required to pass the "high" or "low" bias questionnaire in

each study (**Figure 7**). The most biased domains (~over 50% high bias scoring) from the selected

studies were Baseline Characteristics (i.e. it was largely unclear how sex, age, weight or other

baseline characteristics or confounders were adjusted for in each analysis) and Incomplete Outcome

Data (i.e. it was generally unclear whether all animals were included in each analysis, and if not,

whether there any report on why they were missing outcome data or how that missing data influenced

the study). The lowest biased domain (~over 75% low bias scoring) was Selective Outcome

Reporting (i.e. did the results reported reflect on the methods described in the selective studies). For

further clarification of methods used in order to clear-up bias reporting, we contacted all 58

corresponding authors to be able to provide us with more information (all names and affiliations of

the authors that responded are included in the Acknowledgements section). Overall, the reported bias

from all included studies were largely considered to be "unclear" due to the lack of reporting and

transparency in the methods and results described.

4   Discussion and Recommendations

A prominent role for the innate and acquired immune system in brain health, neurologic and age-

associated neurodegenerative disorders has become increasing apparent. In part, this has been driven

by the immunologic role of several variant gene discoveries including triggering receptor expressed

on myeloid cells 2 (*TREM2*) in Alzheimer's disease, granulin (*GRN*) in frontotemporal dementia and

leucine rich repeat kinase 2 (*LRRK2*) in Parkinson's disease (91–94). Despite this burgeoning interest

in neuroimmunology, and many published studies, results from flow cytometry immunophenotyping

from homogenized mouse brain are highly variable. Although this does not invalidate 'within study'

comparisons of specific immune subsets, such variability is a challenge for reproducibility, meta-

analysis across studies and interpretation (27–84). Reliable data on residential and infiltrating

immune cells within WT/control mouse brain would be of benefit (1–10). One step towards that goal

is standardized reporting of flow cytometry methods and results as well as requiring that to become a

prerequisite for peer-reviewed publications. In this systematic review, we demonstrate most studies

that apply flow cytometry methods to neurology and neuroimmunology, specifically to homogenized

mouse brain, share little to no consensus on methods, analysis or results. Here we summarize our

findings and make a series of recommendations for future studies (**Table 1**).

We retrieved 58 neurological/neuroimmunology studies that utilized flow cytometry to identify or

sort multiple immune cell subsets from WT/control mouse brain, which generally compared within

study results to an experimental mouse model (27–84). We compared mouse strains, perfusion and

tissue processing methods, and note the age of mice and methods for tissue homogenization are

variable (96–99). Vivarium conditions, such as group housing within ventilated racks in a pathogen

free barrier facility versus more conventional non-barrier non-ventilated caging, were seldom

documented. Corresponding authors from the 10/58 studies (17.2%) indicated the majority of studies

utilized a barrier facility with HEPA-filtered air, where each cage was individually ventilated and had

sterilized bedding and chow (27,39,45,51,52,58,62,73,75,76). Housing conditions, age and sex

influences the immune cell subsets that can be identified by flow cytometry (95–98), and should be

carefully considered, documented in experimental protocols and adjusted for as a covariate in

subsequent analyses. When deciding on immune cell isolation methods for mouse brains, both

mechanical homogenization or/and enzymatic tissue digestion are appropriate. Nevertheless, each

approach has pros and cons on immune cell retrieval and phenotypic expression, and depending on

the specific research question, must be carefully considered (99–101).

Once the mouse lines and homogenization/isolation methods were analyzed, we compared flow

cytometry techniques and data reporting across the 58 studies. Cytometers/FACS sorters come in a

variety of makes and models, but essentially do the same function, and should be calibrated using

universal standards (89,102,103). Many useful guidelines exist for reporting and include the

Minimum Information about a Flow Cytometry Experiment (MIFlowCyt) criteria (26,89,102,104).

These recommend researchers present flow cytometry data and methods by reporting: 1)

sample/specimen used for experiments; 2) how the samples were treated (storage, processing,

stained, etc.); 3) what reagents were used and which antibody clones; 4) what controls were used

(unstained controls, FMOs, single color controls, etc.) with a demonstration of the full gating strategy

for each panel; 5) what instrument was used and details about it; 6) how many total cells and/or live

cells were collected in each sample (either exact cell counts or overall estimated counts); 7) what

analysis software was used and how compensation was calculated. Although these guidelines do

recommend reporting the total cells and/or live cells collected in each sample, this can be misleading

for brain homogenate studies that use different tissue dissociation and cell isolation techniques. For

example, a 30%/70% Percoll gradient solution will preferentially isolate immune cells, whereas a

30% solution will isolate immune cells and other residential brain cells (neurons, astrocytes, etc.).

Hence, it would be beneficial to report all CD45$^+$ (infiltrating/residential immune cells within the

brain) and CD45$^{-/lo}$ (microglia and other glial/brain cells). Better documentation would enable

replication, more reliable and accurate results, and enable subsequent meta-analysis.

In the 58 selected studies, we next examined the median and average count and percentages of

immune cell subsets reported from WT/control mouse brain. The immune cell subsets selected

(**Figure 5**, **Figure 6**, **Suppl. Table 3**) consist of brain/CNS-only residential immune cells (microglia)

and immune cells considered to be both residential and peripheral (i.e. T cells, B cells, macrophages,

NK cells) (1–10). As expected, of all the immune subsets examined, microglia are the most populous

immune cells within WT/control mouse brain (15,74–76). For microglia markers, we searched for

publications that showed expression of TMEM119, which is expressed in more stable, non-reactive

microglia, (105–108). Traditional methods of detecting microglia by flow cytometry use CD11b$^+$

CD45$^{lo/-}$ phenotyping. However, we favor microglia-specific markers, such as TMEM119 as these

differentiate microglia from other phagocytic cell types.

For other immune cell subsets, T cells make up the second most abundant immune cell within

WT/control mouse brains, with more bulk CD4$^+$ T cells (27–37). In the brain, it is likely both

CD4$^+$/CD8$^+$ T cells are comprised of resident memory T cells (T$_{RM}$) and might be classified as an

even more specialized subset (i.e. T$_{CM}$, T$_{EM}$, and T$_{EMRA}$) (109–111). For the innate immune cell

populations (excluding microglia), NK cells were observed more frequently in WT/control mouse

brain than other innate immune cells (33,46–51). Many studies report NK cells as most abundant

within the brain's parenchyma, and more often than other innate immune populations (excluding

microglia) or adaptive immune cells (T cell and B cells) (112–114). NK cells were the most reported

innate immune cell subset within WT/control mouse brains in the 58 studies reviewed, but there were

still more bulk CD4$^+$/CD8$^+$ T cell counts reported. We cautiously include neutrophils in our

systematic review (46,77–84). These polynucleated cells are challenging to detect with flow

cytometry as neutrophil extracellular traps (NETs) cause them to be extremely "sticky", to bind onto

each other and to bind non-specifically to flow antibodies (causing false positives) (115–118). The

variability in nomenclature/targets to identify neutrophils, their short life span and sensitivity towards

purification methods are additional limitations (115–118).

The presence of less abundant immune cell subsets found within the brain, or infiltrating the brain,

including specific CD4$^+$ T cell subsets, T cell memory subsets, and DCs, were also assessed.

Unpredictably, it appears that T$_{H1}$ T cells are more abundant in WT/control mouse brains than T$_{H2}$ T

cells (not reported in the 58 selected studies) (27,40). Conventionally, the T$_{H1}$/T$_{H2}$ ratio is used to

determine whether an individual has a bacterial/viral infection (higher ratios are indicative or greater

infection) although higher ratios of these subsets are also found in aged subjects, once again

highlighting the importance of defining age in studies of immunity (119). None of the 58 studies

reported specific T cell memory subsets besides bulk T_{EM} CD4$^+$/CD8$^+$ T cells (30) within WT/control

mice. Given the importance of specific T cell memory subsets to overall immunity, future flow

cytometry analysis of the brain may benefit from their inclusion. Surprisingly, a few reports listed

migratory/residential DCs within WT/control mouse brains, albeit at extremely low levels (33,52–

55), as microglia are thought to mediate brain immune surveillance (120–122).

Our systematic review has some limitations. Our analysis and database search were not automated to

update figures from more recent research (123,124) and potentially, this may be considered a

selection bias. The key word searches we conducted for each immune cell subset has been reported in

our methods but different variations of these names, or use of other abbreviations when searching,

could alter what literature is identified in each database, thus could also be considered as some level

of bias. As several studies selected failed to report sufficient details within their main text, figures, or

supplementary materials, whenever possible, data was extrapolated. We have attempted to avoid any

author bias or incorrect statements as two independent reviewers assessed all manuscripts. Original

authors were also contacted when further clarifications were required.

We recommend supplementary data includes all raw cell counts and documents which data are used

in the main figures and text, to provide more transparency and enable reproducibility. Overall, there

was tremendous variability in immune cell subset counts in the 58 studies reviewed, such that the SD

often exceeded mean estimates. Hence medians with IQRs are provided throughout this review.

Although heterogeneity in instrumentation is unavoidable and necessary in experimental design, and

within study comparisons are still valid, it would be helpful to standardize some aspects of how

immunophenotyping and cell counts are reported. Minimally, publications must include details on

vivarium housing, age, sex, and strain. In our analysis, we have included a series of recommendations

to aid interpretation of results, reproducibility and meta-analysis (**Table 1**). It would be insightful to

compare flow cytometry results across studies, not only within them. Adherence to reporting

guidelines will ultimately improve our understanding of the dynamic role of immunity in mouse

brain.

 5   **Figure Legends**

Figure 1. Prisma Flow Diagram. Guidelines provided by PRISMA (21).

Figure 2. Mouse information reported between studies. Reported baseline mouse information

described as percentages out of the 58 studies: **A.** mouse strains; **B.** mouse sex; and **C.** mouse age.

Figure 3. Brain tissue dissociation and cell isolation methods reported between studies. Tissue

processing techniques were reported as percentages from the 58 studies: **A.** brain dissociation filters

utilized; **B.** digestion enzymes used for brain homogenization; and **C.** Percoll gradients used to

remove myelin layer and isolate immune cells.

Figure 4. Reporting of flow cytometry results and cell counts between the studies. Differences in

flow cytometry results and cell counts reported as within study percentages (n=58 studies): **A.** gating

strategy reported; **B.** total cells per sample collected from flow cytometer/sorter reported; and **C.** total

immune cell subset counts derived from the live cell gate reported.

Figure 5. Calculated medians with interquartile ranges (IQRs) of immune cells quantified by

flow cytometry within wild-type/control mouse brains. Medians with IQRs (defined as: 75th

percentile upper quartile [Q3] – 25th percentile lower quartile [Q1]) of immune cell subset counts

found with wild-type (WT)/control homogenized mouse brain were calculated from data extrapolated

from the 58 studies selected for inclusion in this systematic review. Immune cell subsets were

organized as: **A.** bulk adaptive immune cells; **B.** specialized T cells; **C.** innate immune cells; and **D.**

microglia. Total n-values (number of studies for each identified subset) are reported along with the

median and IQR for each immune subset above each bar graph.

Figure 6. Standardized total cell percentages of immune cells quantified by flow cytometry

within wild-type/control mouse brains. Estimated percentages of immune cell subset counts found

within wild-type (WT)/control homogenized mouse brain were calculated from data extrapolated

from the 58 studies selected for inclusion in this systematic review. The equations used to determine

the standardized cell counts can be found in the Methods section. Briefly, "raw" total immune cell

subset count and "raw" total cell count collected via flow cytometry were standardized assuming 1 x

10^5 total cells were collected. Results were reported as combined means of percentages with standard

deviations (±SD) from the standardized totals from each study assuming 1 x 10^5 total cells were

collected.

Figure 7. SYRCLE's risk of bias: Tools for animal studies summarized for each study. Both the

summary graph and "stop-light" figure was created via the *robvis* R package and Shiny web app (25).

Criteria are based upon the Systematic Review Centre for Laboratory animal Experimentation

(SYRCLE) methodology to detect bias (24). All 58 studies selected were reviewed and analyzed

using SYRCLE criteria domains: D1: Sequence Generation; D2: Baseline Characteristics; D3:

Allocation Concealment; D4: Random Housing; D5: Blinding; D6: Random Outcome Assessment;

D7: Incomplete Outcome Data; D8: Selective Outcome Reporting; and D9: Other Sources of Bias.

Red: High Bias; Yellow: "Unclear" Bias; Green: Low Bias.

Tables

Table 1. Minimum recommendations and sources for standards for future mouse brain flow

cytometry reports

Table 1. Minimum recommendations and sources for standards for future mouse brain flow cytometry reports(4,5,13,15,26,89,95–98,102,125,126)	
Criteria	**Recommendations and Notes**
Mouse Strain	• Report background mouse strain being used, as different mouse strains have different immune backgrounds and responses. • Ensure a WT or a non-treated mouse is included, and housed in the same vivarium as transgenic/treated mice, in order to enable cross-study comparison. • Report specific details of the vivarium and the caging conditions used. • DO NOT switch mouse strains in the middle of an experiment, as immune profiles will vary drastically. • Ensure genetic, phenotypic, and supplier information is provided for each mouse strain used.
Mouse Age	• Report the ages of the mice used for the study, as immune profiles are altered by age. • For a more naïve, younger immune cell population, it is best to utilize mice aged ~3 months. • For a more mature, older immune cell population, it is best to utilize mice aged ~18 months. • Age can affect neurological phenotype and behavior; thus, needs to be included as a confounder.
Mouse Sex	• Report the sex of the mice used, as male and female mice have different immune responses. • Ensure that the same sex of mice is used throughout all experiments within the study, unless the study is focused on sex differences. • Use power analysis based on sex-specific means and their variance to estimate the of male and female required.
Perfusion Techniques	• Perfusion is recommended before brain homogenization, to avoid circulating blood/immune cell contamination. • Use cold Hank's balanced salt solution (HBSS) with 10% serum (i.e. heat-inactive fetal bovine serum [FBS], mouse serum, etc.) via cardiac perfusion to ensure residential immune cells and neurological cells remain intact and viable for flow cytometry. • 4% paraformaldehyde (PFA) in any perfusion mixture will affect flow cytometry staining, and should only be included AFTER staining for surface antigens.
Brain Dissociation and Cell Isolation Techniques	• Mechanical homogenization (i.e. Dounce tissue grinder, bead beating, sonication, etc.), enzymatic homogenization (i.e. collagenase, DNase I, trypsin-EDTA, etc.), or the combination of both can alter the phenotypes of immune cells, thus the it is important to describe the technique was used. • Use one or more cell strainers (preferably using a 100 μm filter first, then a 70 μm filter) in order to remove as much excess dead cells, fats, myelin, etc. as possible, as these components will alter flow cytometry collection and readings.

	- Use an isotonic Percoll gradient in order to isolate immune cells and other cells of interest. At minimum a 37% isotonic Percoll gradient is required to remove all myelin (top layer) and isolate all brain cells (bottom layer/cell pellet). - A 30%/37%/70% isotonic Percoll solution is also recommended if microglia isolation is the only objective.
Flow Cytometer/Sorter Equipment and Analysis Software Reporting	- Report the exact specifications of the flow cytometer/sorter used in the study. - DO NOT switch between cytometers/sorters throughout experimentation, as different models can produce different results. Ideally, calibration and optimization of the machine should be performed daily to ensure accurate results. - Report flow cytometry antibody information, such as clone ID, fluorophore conjugated, targets, and source, etc. - Analysis software and version number should be reported.
Flow Cytometry Result Reporting	- Include a representation of the entire gating strategy (from forward scatter/side scatter, to single cells, to live dead, to beginning of gating strategy and beyond) for each type of experiment conduct. These figures should come from a WT/non-treated control mouse. - State that the results provided are standardized and comparable to multiple flow cytometry controls, such as unstained, FMO, and single-color controls. At a minimum, either unstained and/or FMO control gating should be included in the report. - Provide cell counts and cell percentages for each gate reported. - Histograms should include unstained and FMO results along with the subjects' data. Indicate MFI readings for each histogram included. - Report exact or estimated total cells per sample collected by flow cytometry either as a unified cell count (i.e. all samples had 1×10^5 cells collected during flow cytometry) or for each individual sample. Report the total number of $CD45^+$ cells (all infiltrating/residential immune cells) and $CD45^{-/lo}$ cells (all microglia and other glial cells) when counting immune cells within mouse brain. - Use a live/dead dye to separate out living cells for counts. Include the total amount of live cells counted on average and/or from each sample. - Report cell subsets as either total cell counts or by percentages derived from either live cells or from total cells collected. - Methods used to describe how cell counts were reported (i.e. by use of flow cytometry cell counting beads, counts provided by the machine, counts calculated, etc.) or how MFI's were calculated is recommended. - Deposit raw data, such as FCS files, in a database such as the FlowRepository (http://flowrepository.org/) after full analysis is completed.

Conflict of Interest

The authors declare no potential conflict of interest.

Author Contributions

RCS performed the initial search and analysis of this systematic review, where DTG reviewed all

literature to confirm findings and reach a consensus. RCS and MJF contributed equally on the

conception, design, organization. All authors contributed to manuscript revision, read, and approved

the submitted version.

Funding

MJF is supported by the Lee and Lauren Fixel Chair for Parkinson's disease research.

Acknowledgments

The authors would like to thank John P.A. Ioannidis from the Meta-Research Innovation Center at

Stanford (METRICS) and the Departments of Medicine, Epidemiology and Population Health,

Biomedical Data Science, and Statistics at Stanford University, Stanford, CA., U.S.A., for his review

and helpful comments. The authors would like to also thank Etty (Tika) Benveniste from the

University of Alabama at Birmingham (UAB), Birmingham, AL., U.S.A. and Malu Tansey from the

University of Florida, Gainesville, FL., U.S.A., for recommendations on flow cytometry and

neuroimmunology. The following corresponding authors from the 58 studies reviewed are thanked

for providing additional information about their publications: Ashley S. Harms (University of

Alabama at Birmingham, Birmingham, AL., U.S.A.); Josephine Herz (University Hospital Essen,

University Duisburg-Essen, Germany); Eleonore Beurel (University of Miami, Miami, FL., U.S.A.);

Jae-Kyung Lee (University of Georgia, Athens, GA., U.S.A.); Paul G. McMenamin (Monash

University, Clayton, Victoria, Australia); Asya Rolls (Technion-Israel Institute of Technology, Haifa,

Israel); Anna M. Planas (Institut d'Investigacions Biomediques August Pi I Sunyer [IDIBAPS] and

Institute for Biomedical Research of Barcelona [IIBB]-Spanish Research Council [CSIC], 08036

Barcelona, Spain); Cecile Delarasse (University of Paris, Paris France); Yao Yao (University of

South Florida, Tampa, FL., U.S.A.); and Alanna G. Spiteri (University of Sydney, Sydney,

Australia). The authors would also like to thank the Farrer lab at the University of Florida for their

encouragement and support to perform this systematic review.

Data Availability Statement

All relevant data is contained within the article: The original contributions presented in the study are

included in the article/supplementary material, further inquiries can be directed to the corresponding

author/s.

References

1. Dawson TM, Golde TE, Lagier-Tourenne C. Animal models of neurodegenerative diseases.
Nat Neurosci (2018) 21:1370–1379. doi: 10.1038/s41593-018-0236-8

2. Chesselet M-F, Carmichael ST. Animal Models of Neurological Disorders. *Neurotherapeutics*
(2012) 9:241–244. doi: 10.1007/s13311-012-0118-9

3. Hafezparast M, Ahmad-Annuar A, Wood NW, Tabrizi SJ, Fisher EM. Mouse models for
neurological disease. *Lancet Neurol* (2002) 1:215–224. doi: 10.1016/S1474-4422(02)00100-X

4. Croxford JL, Miyake S. "Animal Models for the Study of Neuroimmunological Disease.,"
Neuroimmunological Diseases. Tokyo: Springer Japan (2016). p. 33–54 doi: 10.1007/978-4-
431-55594-0_3

5. da Silva APB, Silva RBM, Goi LDS, Molina RD, Machado DC, Sato DK. Experimental
Models of Neuroimmunological Disorders: A Review. *Front Neurol* (2020) 11: doi:
10.3389/fneur.2020.00389

6. Grotemeyer A, McFleder RL, Wu J, Wischhusen J, Ip CW. Neuroinflammation in Parkinson's
Disease – Putative Pathomechanisms and Targets for Disease-Modification. *Front Immunol*
(2022) 13: doi: 10.3389/fimmu.2022.878771

7. Saito T, Saido TC. Neuroinflammation in mouse models of Alzheimer's disease. *Clin Exp*
Neuroimmunol (2018) 9:211–218. doi: 10.1111/cen3.12475

8. Passaro AP, Lebos AL, Yao Y, Stice SL. Immune Response in Neurological Pathology:
Emerging Role of Central and Peripheral Immune Crosstalk. *Front Immunol* (2021) 12: doi:
10.3389/fimmu.2021.676621

9. Tan EK, Chao YX, West A, Chan LL, Poewe W, Jankovic J. Parkinson disease and the
immune system — associations, mechanisms and therapeutics. *Nat Rev Neurol* (2020) 16:303–
318. doi: 10.1038/s41582-020-0344-4

10. Ma Q, Xing C, Long W, Wang HY, Liu Q, Wang R-F. Impact of microbiota on central
nervous system and neurological diseases: the gut-brain axis. *J Neuroinflammation* (2019)
16:53. doi: 10.1186/s12974-019-1434-3

11. Hu P, Zhang W, Xin H, Deng G. Single Cell Isolation and Analysis. *Front Cell Dev Biol*
(2016) 4: doi: 10.3389/fcell.2016.00116

12. Lanz T V., Pröbstel A-K, Mildenberger I, Platten M, Schirmer L. Single-Cell High-
Throughput Technologies in Cerebrospinal Fluid Research and Diagnostics. *Front Immunol*
(2019) 10: doi: 10.3389/fimmu.2019.01302

13. Drescher H, Weiskirchen S, Weiskirchen R. Flow Cytometry: A Blessing and a Curse.
Biomedicines (2021) 9:1613. doi: 10.3390/biomedicines9111613

14.   de Graaf MT, de Jongste AHC, Kraan J, Boonstra JG, Smitt PAES, Gratama JW. Flow
cytometric characterization of cerebrospinal fluid cells. *Cytometry B Clin Cytom* (2011)
80B:271–281. doi: 10.1002/cyto.b.20603

15.   Martin D, Xu J, Porretta C, Nichols CD. Neurocytometry: Flow Cytometric Sorting of
Specific Neuronal Populations from Human and Rodent Brain. *ACS Chem Neurosci* (2017)
8:356–367. doi: 10.1021/acschemneuro.6b00374

16.   Brioschi S, d'Errico P, Amann LS, Janova H, Wojcik SM, Meyer-Luehmann M, Rajendran L,
Wieghofer P, Paolicelli RC, Biber K. Detection of Synaptic Proteins in Microglia by Flow
Cytometry. *Front Mol Neurosci* (2020) 13: doi: 10.3389/fnmol.2020.00149

17.   Mrdjen D, Pavlovic A, Hartmann FJ, Schreiner B, Utz SG, Leung BP, Lelios I, Heppner FL,
Kipnis J, Merkler D, et al. High-Dimensional Single-Cell Mapping of Central Nervous System
Immune Cells Reveals Distinct Myeloid Subsets in Health, Aging, and Disease. *Immunity*
(2018) 48:380-395.e6. doi: 10.1016/j.immuni.2018.01.011

18.   Kim H, Sefcik JS, Bradway C. Characteristics of Qualitative Descriptive Studies: A
Systematic Review. *Res Nurs Health* (2017) 40:23–42. doi: 10.1002/nur.21768

19.   Hooijmans CR, IntHout J, Ritskes-Hoitinga M, Rovers MM. Meta-Analyses of Animal
Studies: An Introduction of a Valuable Instrument to Further Improve Healthcare. *ILAR J*
(2014) 55:418–426. doi: 10.1093/ilar/ilu042

20.   Vesterinen HM, Sena ES, Egan KJ, Hirst TC, Churolov L, Currie GL, Antonic A, Howells
DW, Macleod MR. Meta-analysis of data from animal studies: A practical guide. *J Neurosci*
Methods (2014) 221:92–102. doi: 10.1016/j.jneumeth.2013.09.010

21.   Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic
Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* (2009) 6:e1000097. doi:
10.1371/journal.pmed.1000097

22.   Clarke M, Oxman A, Paulsen E, Higgins J, Green S. Guide to the contents of a Cochrane
Methodology protocol and review. *The Cochrane Collaboration* (2020) 5.1.0:

23.   Munn Z, Stern C, Aromataris E, Lockwood C, Jordan Z. What kind of systematic review
should I conduct? A proposed typology and guidance for systematic reviewers in the medical
and health sciences. *BMC Med Res Methodol* (2018) 18:5. doi: 10.1186/s12874-017-0468-4

24.   Hooijmans CR, Rovers MM, de Vries RB, Leenaars M, Ritskes-Hoitinga M, Langendam MW.
SYRCLE's risk of bias tool for animal studies. *BMC Med Res Methodol* (2014) 14:43. doi:
10.1186/1471-2288-14-43

25.   McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): An R package and Shiny
web app for visualizing risk-of-bias assessments. *Res Synth Methods* (2021) 12:55–61. doi:
10.1002/jrsm.1411

26.   Cossarizza A, Chang H, Radbruch A, Abrignani S, Addo R, Akdis M, Andrä I, Andreata F,
Annunziato F, Arranz E, et al. Guidelines for the use of flow cytometry and cell sorting in

immunological studies (third edition). *Eur J Immunol* (2021) 51:2708–3145. doi:
10.1002/eji.202170126

27. Williams GP, Schonhoff AM, Jurkuvenaite A, Gallups NJ, Standaert DG, Harms AS. CD4 T
cells mediate brain inflammation and neurodegeneration in a mouse model of
Parkinson's disease. *Brain* (2021) 144:2047–2059. doi: 10.1093/brain/awab103

28. Ferretti MT, Merlini M, Späni C, Gericke C, Schweizer N, Enzmann G, Engelhardt B, Kulic
L, Suter T, Nitsch RM. T-cell brain infiltration and immature antigen-presenting cells in
transgenic models of Alzheimer's disease-like cerebral amyloidosis. *Brain Behav Immun*
(2016) 54:211–225. doi: 10.1016/j.bbi.2016.02.009

29. Ito M, Komai K, Mise-Omata S, Iizuka-Koga M, Noguchi Y, Kondo T, Sakai R, Matsuo K,
Nakayama T, Yoshie O, et al. Brain regulatory T cells suppress astrogliosis and potentiate
neurological recovery. *Nature* (2019) 565:246–250. doi: 10.1038/s41586-018-0824-5

30. Daglas M, Draxler DF, Ho H, McCutcheon F, Galle A, Au AE, Larsson P, Gregory J,
Alderuccio F, Sashindranath M, et al. Activated CD8+ T Cells Cause Long-Term Neurological
Impairment after Traumatic Brain Injury in Mice. *Cell Rep* (2019) 29:1178-1191.e6. doi:
10.1016/j.celrep.2019.09.046

31. Pasciuto E, Burton OT, Roca CP, Lagou V, Rajan WD, Theys T, Mancuso R, Tito RY, Kouser
L, Callaerts-Vegh Z, et al. Microglia Require CD4 T Cells to Complete the Fetal-to-Adult
Transition. *Cell* (2020) 182:625-640.e24. doi: 10.1016/j.cell.2020.06.026

32. Herz J, Köster C, Crasmöller M, Abberger H, Hansen W, Felderhoff-Müser U, Bendix I.
Peripheral T Cell Depletion by FTY720 Exacerbates Hypoxic-Ischemic Brain Injury in
Neonatal Mice. *Front Immunol* (2018) 9: doi: 10.3389/fimmu.2018.01696

33. Chu HX, Kim HA, Lee S, Moore JP, Chan CT, Vinh A, Gelderblom M, Arumugam T V,
Broughton BR, Drummond GR, et al. Immune Cell Infiltration in Malignant Middle Cerebral
Artery Infarction: Comparison with Transient Cerebral Ischemia. *Journal of Cerebral Blood*
Flow & Metabolism (2014) 34:450–459. doi: 10.1038/jcbfm.2013.217

34. Zhou Y, Wang X, Tang D, Li Y, Jiao Y, Gan Y, Hu X, Yang L, Yu W, Stetler RA, et al. IL-
2mAb reduces demyelination after focal cerebral ischemia by suppressing CD8 $^+$ T cells. *CNS*
Neurosci Ther (2019) 25:532–543. doi: 10.1111/cns.13084

35. Shaw TN, Stewart-Hutchinson PJ, Strangward P, Dandamudi DB, Coles JA, Villegas-Mendez
A, Gallego-Delgado J, van Rooijen N, Zindy E, Rodriguez A, et al. Perivascular Arrest of
CD8+ T Cells Is a Signature of Experimental Cerebral Malaria. *PLoS Pathog* (2015)
11:e1005210. doi: 10.1371/journal.ppat.1005210

36. Jurado KA, Yockey LJ, Wong PW, Lee S, Huttner AJ, Iwasaki A. Antiviral CD8 T cells
induce Zika-virus-associated paralysis in mice. *Nat Microbiol* (2017) 3:141–147. doi:
10.1038/s41564-017-0060-z

37. Unger MS, Li E, Scharnagl L, Poupardin R, Altendorfer B, Mrowetz H, Hutter-Paier B,
Weiger TM, Heneka MT, Attems J, et al. CD8+ T-cells infiltrate Alzheimer's disease brains

and regulate neuronal- and synapse-related gene expression in APP-PS1 transgenic mice.
Brain Behav Immun (2020) 89:67–86. doi: 10.1016/j.bbi.2020.05.070

38. Meng H, Zhao H, Cao X, Hao J, Zhang H, Liu Y, Zhu M, Fan L, Weng L, Qian L, et al.
Double-negative T cells remarkably promote neuroinflammation after ischemic stroke.
Proceedings of the National Academy of Sciences (2019) 116:5558–5563. doi:
10.1073/pnas.1814394116

39. Beckmann L, Obst S, Labusek N, Abberger H, Köster C, Klein-Hitpass L, Schumann S,
Kleinschnitz C, Hermann DM, Felderhoff-Müser U, et al. Regulatory T Cells Contribute to
Sexual Dimorphism in Neonatal Hypoxic-Ischemic Brain Injury. *Stroke* (2022) 53:381–390.
doi: 10.1161/STROKEAHA.121.037537

40. Li W, Luo Y, Xu H, Ma Q, Yao Q. Imbalance between T helper 1 and regulatory T cells plays
a detrimental role in experimental Parkinson's disease in mice. *Journal of International*
Medical Research (2021) 49:030006052199847. doi: 10.1177/0300060521998471

41. Chen C, Chencheng Z, Cuiying L, Xiaokun G. Plasmacytoid Dendritic Cells Protect Against
Middle Cerebral Artery Occlusion Induced Brain Injury by Priming Regulatory T Cells. *Front*
Cell Neurosci (2020) 14: doi: 10.3389/fncel.2020.00008

42. Maes W, Verschuere T, Van Hoylandt A, Boon L, Van Gool S. Depletion of Regulatory T
Cells in a Mouse Experimental Glioma Model through Anti-CD25 Treatment Results in the
Infiltration of Non-Immunosuppressive Myeloid Cells in the Brain. *Clin Dev Immunol* (2013)
2013:1–6. doi: 10.1155/2013/952469

43. Rayasam A, Kijak JA, Kissel L, Choi YH, Kim T, Hsu M, Joshi D, Laaker CJ, Cismaru P,
Lindstedt A, et al. CXCL13 expressed on inflamed cerebral blood vessels recruit IL-21
producing TFH cells to damage neurons following stroke. *J Neuroinflammation* (2022)
19:125. doi: 10.1186/s12974-022-02490-2

44. Guo Y, Chen X, Li D, Liu H, Ding Y, Han R, Shi Y, Ma X. PR-957 mediates neuroprotection
by inhibiting Th17 differentiation and modulating cytokine production in a mouse model of
ischaemic stroke. *Clin Exp Immunol* (2018) 193:194–206. doi: 10.1111/cei.13132

45. Beurel E, Lowell JA, Jope RS. Distinct characteristics of hippocampal pathogenic TH17 cells
in a mouse model of depression. *Brain Behav Immun* (2018) 73:180–191. doi:
10.1016/j.bbi.2018.04.012

46. He H, Geng T, Chen P, Wang M, Hu J, Kang L, Song W, Tang H. NK cells promote
neutrophil recruitment in the brain during sepsis-induced neuroinflammation. *Sci Rep* (2016)
6:27711. doi: 10.1038/srep27711

47. Jin W-N, Shi K, He W, Sun J-H, Van Kaer L, Shi F-D, Liu Q. Neuroblast senescence in the
aged brain augments natural killer cell cytotoxicity leading to impaired neurogenesis and
cognition. *Nat Neurosci* (2021) 24:61–73. doi: 10.1038/s41593-020-00745-w

48. Zhang Y, Fung ITH, Sankar P, Chen X, Robison LS, Ye L, D'Souza SS, Salinero AE,
Kuentzel ML, Chittur S V., et al. Depletion of NK Cells Improves Cognitive Function in the

Alzheimer Disease Mouse Model. *The Journal of Immunology* (2020) 205:502–510. doi:
10.4049/jimmunol.2000037

49. Li Z, Li M, Shi SX, Yao N, Cheng X, Guo A, Zhu Z, Zhang X, Liu Q. Brain transforms
natural killer cells that exacerbate brain edema after intracerebral hemorrhage. *Journal of*
Experimental Medicine (2020) 217: doi: 10.1084/jem.20200213

50. D'Alessandro G, Antonangeli F, Marrocco F, Porzia A, Lauro C, Santoni A, Limatola C. Gut
microbiota alterations affect glioma growth and innate immune cells involved in tumor
immunosurveillance in mice. *Eur J Immunol* (2020) 50:705–711. doi: 10.1002/eji.201948354

51. Earls RH, Menees KB, Chung J, Gutekunst C-A, Lee HJ, Hazim MG, Rada B, Wood LB, Lee
J-K. NK cells clear α-synuclein and the depletion of NK cells exacerbates synuclein pathology
in a mouse model of α-synucleinopathy. *Proceedings of the National Academy of Sciences*
(2020) 117:1762–1771. doi: 10.1073/pnas.1909110117

52. Dando SJ, Naranjo Golborne C, Chinnery HR, Ruitenberg MJ, McMenamin PG. A case of
mistaken identity: CD11c-eYFP [+] cells in the normal mouse brain parenchyma and neural
retina display the phenotype of microglia, not dendritic cells. *Glia* (2016) 64:1331–1349. doi:
10.1002/glia.23005

53. Clarkson BD, Walker A, Harris MG, Rayasam A, Hsu M, Sandor M, Fabry Z. CCR7 deficient
inflammatory Dendritic Cells are retained in the Central Nervous System. *Sci Rep* (2017)
7:42856. doi: 10.1038/srep42856

54. Gelderblom M, Gallizioli M, Ludewig P, Thom V, Arunachalam P, Rissiek B, Bernreuther C,
Glatzel M, Korn T, Arumugam TV, et al. IL-23 (Interleukin-23)–Producing Conventional
Dendritic Cells Control the Detrimental IL-17 (Interleukin-17) Response in Stroke. *Stroke*
(2018) 49:155–164. doi: 10.1161/STROKEAHA.117.019101

55. Göbel K, Pankratz S, Asaridou C-M, Herrmann AM, Bittner S, Merker M, Ruck T, Glumm S,
Langhauser F, Kraft P, et al. Blood coagulation factor XII drives adaptive immunity during
neuroinflammation via CD87-mediated modulation of dendritic cells. *Nat Commun* (2016)
7:11626. doi: 10.1038/ncomms11626

56. Brioschi S, Wang W-L, Peng V, Wang M, Shchukina I, Greenberg ZJ, Bando JK, Jaeger N,
Czepielewski RS, Swain A, et al. Heterogeneity of meningeal B cells reveals a lymphopoietic
niche at the CNS borders. *Science (1979)* (2021) 373: doi: 10.1126/science.abf9277

57. Pierson ER, Stromnes IM, Goverman JM. B Cells Promote Induction of Experimental
Autoimmune Encephalomyelitis by Facilitating Reactivation of T Cells in the Central Nervous
System. *The Journal of Immunology* (2014) 192:929–939. doi: 10.4049/jimmunol.1302171

58. Korin B, Ben-Shaanan TL, Schiller M, Dubovik T, Azulay-Debby H, Boshnak NT, Koren T,
Rolls A. High-dimensional, single-cell characterization of the brain's immune compartment.
Nat Neurosci (2017) 20:1300–1309. doi: 10.1038/nn.4610

59. Liu C, Yang J, Zhang C, Geng X, Zhao H. Remote ischemic conditioning reduced cerebral
ischemic injury by modulating inflammatory responses and ERK activity in type 2 diabetic
mice. *Neurochem Int* (2020) 135:104690. doi: 10.1016/j.neuint.2020.104690

60. Mracsko E, Javidi E, Na S-Y, Kahn A, Liesz A, Veltkamp R. Leukocyte Invasion of the Brain
After Experimental Intracerebral Hemorrhage in Mice. *Stroke* (2014) 45:2107–2114. doi:
10.1161/STROKEAHA.114.005801

61. Trahanas DM, Cuda CM, Perlman H, Schwulst SJ. Differential Activation of Infiltrating
Monocyte-Derived Cells After Mild and Severe Traumatic Brain Injury. *Shock* (2015) 43:255–
260. doi: 10.1097/SHK.0000000000000291

62. Miró-Mur F, Pérez-de-Puig I, Ferrer-Ferrer M, Urra X, Justicia C, Chamorro A, Planas AM.
Immature monocytes recruited to the ischemic mouse brain differentiate into macrophages
with features of alternative activation. *Brain Behav Immun* (2016) 53:18–33. doi:
10.1016/j.bbi.2015.08.010

63. Ritzel RM, Patel AR, Grenier JM, Crapser J, Verma R, Jellison ER, McCullough LD.
Functional differences between microglia and monocytes after ischemic stroke. *J
Neuroinflammation* (2015) 12:106. doi: 10.1186/s12974-015-0329-1

64. Cazareth J, Guyon A, Heurteaux C, Chabry J, Petit-Paitel A. Molecular and cellular
neuroinflammatory status of mouse brain after systemic lipopolysaccharide challenge:
importance of CCR2/CCL2 signaling. *J Neuroinflammation* (2014) 11:132. doi:
10.1186/1742-2094-11-132

65. Chu HX, Broughton BRS, Ah Kim H, Lee S, Drummond GR, Sobey CG. Evidence That Ly6C
[hi] Monocytes Are Protective in Acute Ischemic Stroke by Promoting M2 Macrophage
Polarization. *Stroke* (2015) 46:1929–1937. doi: 10.1161/STROKEAHA.115.009426

66. Peralta Ramos JM, Iribarren P, Bousset L, Melki R, Baekelandt V, Van der Perren A.
Peripheral Inflammation Regulates CNS Immune Surveillance Through the Recruitment of
Inflammatory Monocytes Upon Systemic α-Synuclein Administration. *Front Immunol* (2019)
10: doi: 10.3389/fimmu.2019.00080

67. Li Q, Lan X, Han X, Wang J. Expression of Tmem119/Sall1 and Ccr2/CD69 in FACS-Sorted
Microglia- and Monocyte/Macrophage-Enriched Cell Populations After Intracerebral
Hemorrhage. *Front Cell Neurosci* (2019) 12: doi: 10.3389/fncel.2018.00520

68. Hsieh CL, Kim CC, Ryba BE, Niemi EC, Bando JK, Locksley RM, Liu J, Nakamura MC,
Seaman WE. Traumatic brain injury induces macrophage subsets in the brain. *Eur J Immunol*
(2013) 43:2010–2022. doi: 10.1002/eji.201243084

69. Hsieh CL, Niemi EC, Wang SH, Lee CC, Bingham D, Zhang J, Cozen ML, Charo I, Huang
EJ, Liu J, et al. CCR2 Deficiency Impairs Macrophage Infiltration and Improves Cognitive
Function after Traumatic Brain Injury. *J Neurotrauma* (2014) 31:1677–1688. doi:
10.1089/neu.2013.3252

70. DePaula-Silva AB, Gorbea C, Doty DJ, Libbey JE, Sanchez JMS, Hanak TJ, Cazalla D,
Fujinami RS. Differential transcriptional profiles identify microglial- and macrophage-specific
gene markers expressed during virus-induced neuroinflammation. *J Neuroinflammation* (2019)
16:152. doi: 10.1186/s12974-019-1545-x

71. Lehmann ML, Cooper HA, Maric D, Herkenham M. Social defeat induces depressive-like
states and microglial activation without involvement of peripheral macrophages. *J
Neuroinflammation* (2016) 13:224. doi: 10.1186/s12974-016-0672-x

72. Cai W, Liu S, Hu M, Sun X, Qiu W, Zheng S, Hu X, Lu Z. Post-stroke DHA Treatment
Protects Against Acute Ischemic Brain Injury by Skewing Macrophage Polarity Toward the
M2 Phenotype. *Transl Stroke Res* (2018) 9:669–680. doi: 10.1007/s12975-018-0662-7

73. Martin E, El-Behi M, Fontaine B, Delarasse C. Analysis of Microglia and Monocyte-derived
Macrophages from the Central Nervous System by Flow Cytometry. *Journal of Visualized
Experiments* (2017) doi: 10.3791/55781

74. Honarpisheh P, Lee J, Banerjee A, Blasco-Conesa MP, Honarpisheh P, d'Aigle J, Mamun AA,
Ritzel RM, Chauhan A, Ganesh BP, et al. Potential caveats of putative microglia-specific
markers for assessment of age-related cerebrovascular neuroinflammation. *J
Neuroinflammation* (2020) 17:366. doi: 10.1186/s12974-020-02019-5

75. Nirwane A, Yao Y. SMAlow/undetectable pericytes differentiate into microglia- and
macrophage-like cells in ischemic brain. *Cellular and Molecular Life Sciences* (2022) 79:264.
doi: 10.1007/s00018-022-04322-1

76. Spiteri AG, Terry RL, Wishart CL, Ashhurst TM, Campbell IL, Hofer MJ, King NJC. High-
parameter cytometry unmasks microglial cell spatio-temporal response kinetics in severe
neuroinflammatory disease. *J Neuroinflammation* (2021) 18:166. doi: 10.1186/s12974-021-
02214-y

77. Perez-de-Puig I, Miró-Mur F, Ferrer-Ferrer M, Gelpi E, Pedragosa J, Justicia C, Urra X,
Chamorro A, Planas AM. Neutrophil recruitment to the brain in mouse and human ischemic
stroke. *Acta Neuropathol* (2015) 129:239–257. doi: 10.1007/s00401-014-1381-0

78. Garcia-Bonilla L, Moore JM, Racchumi G, Zhou P, Butler JM, Iadecola C, Anrather J.
Inducible Nitric Oxide Synthase in Neutrophils and Endothelium Contributes to Ischemic
Brain Injury in Mice. *The Journal of Immunology* (2014) 193:2531–2537. doi:
10.4049/jimmunol.1400918

79. Makinde HM, Cuda CM, Just TB, Perlman HR, Schwulst SJ. Nonclassical Monocytes
Mediate Secondary Injury, Neurocognitive Outcome, and Neutrophil Infiltration after
Traumatic Brain Injury. *The Journal of Immunology* (2017) 199:3583–3591. doi:
10.4049/jimmunol.1700896

80. Garcia-Bonilla L, Racchumi G, Murphy M, Anrather J, Iadecola C. Endothelial CD36
Contributes to Postischemic Brain Injury by Promoting Neutrophil Activation via CSF3. *The
Journal of Neuroscience* (2015) 35:14783–14793. doi: 10.1523/JNEUROSCI.2980-15.2015

81. Yao H-W, Kuan C-Y. Early neutrophil infiltration is critical for inflammation-sensitized
hypoxic-ischemic brain injury in newborns. *Journal of Cerebral Blood Flow & Metabolism*
(2020) 40:2188–2200. doi: 10.1177/0271678X19891839

82. Pösel C, Möller K, Boltze J, Wagner D-C, Weise G. Isolation and Flow Cytometric Analysis
of Immune Cells from the Ischemic Mouse Brain. *Journal of Visualized Experiments* (2016)
doi: 10.3791/53658

83. García-Culebras A, Durán-Laforet V, Peña-Martínez C, Moraga A, Ballesteros I, Cuartero MI,
de la Parra J, Palma-Tortosa S, Hidalgo A, Corbí AL, et al. Role of TLR4 (Toll-Like Receptor
4) in N1/N2 Neutrophil Programming After Stroke. *Stroke* (2019) 50:2922–2932. doi:
10.1161/STROKEAHA.119.025085

84. Roy-O'Reilly MA, Ahnstedt H, Spychala MS, Munshi Y, Aronowski J, Sansing LH,
McCullough LD. Aging exacerbates neutrophil pathogenicity in ischemic stroke. *Aging* (2020)
12:436–461. doi: 10.18632/aging.102632

85. Mattei D, Ivanov A, van Oostrum M, Pantelyushin S, Richetto J, Mueller F, Beffinger M,
Schellhammer L, vom Berg J, Wollscheid B, et al. Enzymatic Dissociation Induces
Transcriptional and Proteotype Bias in Brain Cell Populations. *Int J Mol Sci* (2020) 21:7944.
doi: 10.3390/ijms21217944

86. Herron S, Delpech J-C, Madore C, Ikezu T. Using mechanical homogenization to isolate
microglia from mouse brain tissue to preserve transcriptomic integrity. *STAR Protoc* (2022)
3:101670. doi: 10.1016/j.xpro.2022.101670

87. Calvo B, Rubio F, Fernández M, Tranque P. Dissociation of neonatal and adult mice brain for
simultaneous analysis of microglia, astrocytes and infiltrating lymphocytes by flow cytometry.
IBRO Rep (2020) 8:36–47. doi: 10.1016/j.ibror.2019.12.004

88. Nahrendorf M, Swirski FK. Abandoning M1/M2 for a Network Model of Macrophage
Function. *Circ Res* (2016) 119:414–417. doi: 10.1161/CIRCRESAHA.116.309194

89. Herzenberg LA, Tung J, Moore WA, Herzenberg LA, Parks DR. Interpreting flow cytometry
data: a guide for the perplexed. *Nat Immunol* (2006) 7:681–685. doi: 10.1038/ni0706-681

90. Drucker AM, Fleming P, Chan A-W. Research Techniques Made Simple: Assessing Risk of
Bias in Systematic Reviews. *Journal of Investigative Dermatology* (2016) 136:e109–e114. doi:
10.1016/j.jid.2016.08.021

91. Colonna M. TREMs in the immune system and beyond. *Nat Rev Immunol* (2003) 3:445–453.
doi: 10.1038/nri1106

92. Keating CE, Houser MC, MacPherson KP, Herrick MK, Coomes A, Joers VL, Oliver D,
Johnson AG, Chang J, McEachin Z, et al. Loss of progranulin leads to dysregulation of innate
and adaptive immune cell populations, increased susceptibility to experimental colitis, and
brain infiltration of peripheral immune cells. *Alzheimer's & Dementia* (2020) 16: doi:
10.1002/alz.042177

93.   Rhinn H, Tatton N, McCaughey S, Kurnellas M, Rosenthal A. Progranulin as a therapeutic
823   target in neurodegenerative diseases. *Trends Pharmacol Sci* (2022) 43:641–652. doi:
824   10.1016/j.tips.2021.11.015

825   94.   Wallings RL, Tansey MG. LRRK2 regulation of immune-pathways and inflammatory disease.
826   *Biochem Soc Trans* (2019) 47:1581–1595. doi: 10.1042/BST20180463

827   95.   Golomb SM, Guldner IH, Zhao A, Wang Q, Palakurthi B, Aleksandrovic EA, Lopez JA, Lee
828   SW, Yang K, Zhang S. Multi-modal Single-Cell Analysis Reveals Brain Immune Landscape
829   Plasticity during Aging and Gut Microbiota Dysbiosis. *Cell Rep* (2020) 33:108438. doi:
830   10.1016/j.celrep.2020.108438

831   96.   Kovacs EJ, Palmer JL, Fortin CF, Fülöp T, Goldstein DR, Linton P-J. Aging and innate
832   immunity in the mouse: impact of intrinsic and extrinsic factors. *Trends Immunol* (2009)
833   30:319–324. doi: 10.1016/j.it.2009.03.012

834   97.   Zhang X, Pearsall VM, Carver CM, Atkinson EJ, Clarkson BDS, Grund EM, Baez-Faria M,
835   Pavelko KD, Kachergus JM, White TA, et al. Rejuvenation of the aged brain immune cell
836   landscape in mice through p16-positive senescent cell clearance. *Nat Commun* (2022)
837   13:5671. doi: 10.1038/s41467-022-33226-8

838   98.   Erickson M, Banks W. Age-Associated Changes in the Immune System and Blood–Brain
839   Barrier Functions. *Int J Mol Sci* (2019) 20:1632. doi: 10.3390/ijms20071632

840   99.   Cumba Garcia LM, Huseby Kelcher AM, Malo CS, Johnson AJ. Superior isolation of antigen-
841   specific brain infiltrating T cells using manual homogenization technique. *J Immunol Methods*
842   (2016) 439:23–28. doi: 10.1016/j.jim.2016.09.002

843   100.   Srakočić S, Josić P, Trifunović S, Gajović S, Grčević D, Glasnović A. Proposed practical
844   protocol for flow cytometry analysis of microglia from the healthy adult mouse brain:
845   Systematic review and isolation methods' evaluation. *Front Cell Neurosci* (2022) 16: doi:
846   10.3389/fncel.2022.1017976

847   101.   Molina Estevez FJ, Mathews TD, Biffi A, Peviani M. Simultaneous Flow Cytometric
848   Characterization of Multiple Cell Types Retrieved from Mouse Brain/Spinal Cord Through
849   Different Homogenization Methods. *Journal of Visualized Experiments* (2019) doi:
850   10.3791/60335

851   102.   Kalina T. Reproducibility of Flow Cytometry Through Standardization: Opportunities and
852   Challenges. *Cytometry Part A* (2020) 97:137–147. doi: 10.1002/cyto.a.23901

853   103.   Parks DR, Moore WA, Brinkman RR, Chen Y, Condello D, El Khettabi F, Nolan JP, Perfetto
854   SP, Redelman D, Spidlen J, et al. Methodology for evaluating and comparing flow cytometers:
855   A multisite study of 23 instruments. *Cytometry Part A* (2018) 93:1087–1091. doi:
856   10.1002/cyto.a.23605

857   104.   Lee JA, Spidlen J, Boyce K, Cai J, Crosbie N, Dalphin M, Furlong J, Gasparetto M, Goldberg
858   M, Goralczyk EM, et al. MIFlowCyt: The minimum information about a flow cytometry
859   experiment. *Cytometry Part A* (2008) 73A:926–930. doi: 10.1002/cyto.20623

860 105. Ruan C, Elyaman W. A New Understanding of TMEM119 as a Marker of Microglia. *Front*
861   *Cell Neurosci* (2022) 16: doi: 10.3389/fncel.2022.902372

862 106. Young KF, Gardner R, Sariana V, Whitman SA, Bartlett MJ, Falk T, Morrison HW. Can
863   quantifying morphology and TMEM119 expression distinguish between microglia and
864   infiltrating macrophages after ischemic stroke and reperfusion in male and female mice? *J*
865   *Neuroinflammation* (2021) 18:58. doi: 10.1186/s12974-021-02105-2

866 107. Jurga AM, Paleczna M, Kuter KZ. Overview of General and Discriminating Markers of
867   Differential Microglia Phenotypes. *Front Cell Neurosci* (2020) 14: doi:
868   10.3389/fncel.2020.00198

869 108. Hopperton KE, Mohammad D, Trépanier MO, Giuliano V, Bazinet RP. Markers of microglia
870   in post-mortem brain samples from patients with Alzheimer's disease: a systematic review.
871   *Mol Psychiatry* (2018) 23:177–198. doi: 10.1038/mp.2017.246

872 109. Fransen NL, Hsiao C-C, van der Poel M, Engelenburg HJ, Verdaasdonk K, Vincenten MCJ,
873   Remmerswaal EBM, Kuhlmann T, Mason MRJ, Hamann J, et al. Tissue-resident memory T
874   cells invade the brain parenchyma in multiple sclerosis white matter lesions. *Brain* (2020)
875   143:1714–1730. doi: 10.1093/brain/awaa117

876 110. Steinbach K, Vincenti I, Kreutzfeldt M, Page N, Muschaweckh A, Wagner I, Drexler I,
877   Pinschewer D, Korn T, Merkler D. Brain-resident memory T cells represent an autonomous
878   cytotoxic barrier to viral infection. *Journal of Experimental Medicine* (2016) 213:1571–1587.
879   doi: 10.1084/jem.20151916

880 111. Prasad S, Lokensgard JR. Brain-Resident T Cells Following Viral Infection. *Viral Immunol*
881   (2019) 32:48–54. doi: 10.1089/vim.2018.0084

882 112. Sedgwick AJ, Ghazanfari N, Constantinescu P, Mantamadiotis T, Barrow AD. The Role of
883   NK Cells and Innate Lymphoid Cells in Brain Cancer. *Front Immunol* (2020) 11: doi:
884   10.3389/fimmu.2020.01549

885 113. Cuapio A, Ljunggren H-G. Activated Natural Killer Cells Hit Neurogenesis in the Aging
886   Brain. *Neurosci Bull* (2021) 37:1072–1074. doi: 10.1007/s12264-021-00654-3

887 114. Earls RH, Lee JK. The role of natural killer cells in Parkinson's disease. *Exp Mol Med* (2020)
888   52:1517–1525. doi: 10.1038/S12276-020-00505-7

889 115. McKenna E, Mhaonaigh AU, Wubben R, Dwivedi A, Hurley T, Kelly LA, Stevenson NJ,
890   Little MA, Molloy EJ. Neutrophils: Need for Standardized Nomenclature. *Front Immunol*
891   (2021) 12: doi: 10.3389/fimmu.2021.602963

892 116. Fine N, Barzilay O, Sun C, Wellappuli N, Tanwir F, Chadwick JW, Oveisi M, Tasevski N,
893   Prescott D, Gargan M, et al. Primed PMNs in healthy mouse and human circulation are first
894   responders during acute inflammation. *Blood Adv* (2019) 3:1622–1637. doi:
895   10.1182/bloodadvances.2018030585

896 117. McGill CJ, Lu RJ, Benayoun BA. Protocol for analysis of mouse neutrophil NETosis by flow
897   cytometry. *STAR Protoc* (2021) 2:100948. doi: 10.1016/j.xpro.2021.100948

898   118.   Manda-Handzlik A, Ostafin M, Bystrzycka W, Sieczkowska S, Moskalik A, Demkow U,
899   Ciepiela O. Flow cytometric quantification of neutrophil extracellular traps: Limitations of the
900   methodological approach. *Am J Hematol* (2016) 91:E9–E10. doi: 10.1002/ajh.24257

901   119.   Sakata-Kaneko S, Wakatsuki Y, Matsunaga Y, Usui T, Kita T. Altered Th1/Th2 commitment
902   in human CD4+ T cells with ageing. *Clin Exp Immunol* (2001) 120:267–273. doi:
903   10.1046/j.1365-2249.2000.01224.x

904   120.   D'Agostino PM, Gottfried-Blackmore A, Anandasabapathy N, Bulloch K. Brain dendritic
905   cells: biology and pathology. *Acta Neuropathol* (2012) 124:599–614. doi: 10.1007/s00401-
906   012-1018-0

907   121.   Constant O, Maarifi G, Blanchet FP, Van de Perre P, Simonin Y, Salinas S. Role of Dendritic
908   Cells in Viral Brain Infections. *Front Immunol* (2022) 13: doi: 10.3389/fimmu.2022.862053

909   122.   Ludewig P, Gallizioli M, Urra X, Behr S, Brait VH, Gelderblom M, Magnus T, Planas AM.
910   Dendritic cells in brain diseases. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of*
911   *Disease* (2016) 1862:352–367. doi: 10.1016/j.bbadis.2015.11.003

912   123.   Clark J, McFarlane C, Cleo G, Ishikawa Ramos C, Marshall S. The Impact of Systematic
913   Review Automation Tools on Methodological Quality and Time Taken to Complete
914   Systematic Review Tasks: Case Study. *JMIR Med Educ* (2021) 7:e24418. doi: 10.2196/24418

915   124.   Marshall IJ, Wallace BC. Toward systematic review automation: a practical guide to using
916   machine learning tools in research synthesis. *Syst Rev* (2019) 8:163. doi: 10.1186/s13643-019-
917   1074-9

918   125.   Hammond TR, Dufort C, Dissing-Olesen L, Giera S, Young A, Wysoker A, Walker AJ,
919   Gergits F, Segel M, Nemesh J, et al. Single-Cell RNA Sequencing of Microglia throughout the
920   Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes. *Immunity*
921   (2019) 50:253-271.e6. doi: 10.1016/j.immuni.2018.11.004

922   126.   Ritzel RM, Lai Y-J, Crapser JD, Patel AR, Schrecengost A, Grenier JM, Mancini NS, Patrizz
923   A, Jellison ER, Morales-Scheihing D, et al. Aging alters the immunological response to
924   ischemic stroke. *Acta Neuropathol* (2018) 136:89–110. doi: 10.1007/s00401-018-1859-2

A. Mouse Strain Used %

B. Mouse Gender Used %

1.72% B6129SF2/J
1.72% C3H/Fej
1.72% BKS.Cg-Dock7m+/+Leprdb/Nju,db/db
1.72% BALB/C
1.72% Not Reported

91.4%

C57BL/6
44.8% C57BL/6
3.49% C57BL/6N
32.8% C57BL/6J
3.45% C57/BL6
3.45% C57BL/6 (H-2b)
3.45% C57BL/6J (B6)

Not Reported

Female 12.07%

13.79%

Male and Female 17.24%

56.90%

Male

C. Mouse Age

A.

Brain Dissociation Filter Used %

- 5.17% MACS Strainer System
- 40 μm Filter
- 10.34%
- 70 μm Filter
- 39.66%
- 100 μm Filter
- 13.79%
- Not Reported or Used Dissocation Kit Only
- 31.03%

B.

Digestion Enzymes Used %

- Collagenase Alone
  - 1.72% Collagenase D
  - 1.72% Collagenase II
  - 1.72% Collagenase IV
  - 5.17%
- Collagenase + Other Enzymes
  - 6.90% Collagenase/DNase I
  - 15.51% Collagenase/Dispace/DNase I
  - 1.72% Collagenase DNase I
  - 8.62% Collagenase IV/DNase I
  - 1.72% Collagenase IX/Hyalurondise/Collagenase I-S
  - 12.07% Liberase/DNase I
  - 46.55%
- Not Reported
  - 45.00%
- Other Enzymes
  - 1.72% Tryspsin-EDTA
  - 1.72% DNase I/Dispace II/Papain
  - 3.45%

C.

Percoll Gradients Used

Percentage (%)

Standardized Total Cell% in Mouse Brain

Legend:
- 1.82% ± 1.57 DN T cells
- 0.86% ± 1.33 T_{REG} T cells
- 0.008% ± 0 T_{FH} T cells
- 2.39% ± 2.70 T_{H1} T cells
- 0.32% ± 0.06 T_{H17} T cells
- 1.01% ± 0 T_{EM} Bulk CD4$^+$ T cells
- 0.75% ± 0 T_{EM} Bulk CD8$^+$ T cells
- 1.28% ± 2.64 NK cells
- 0.04% ± 0.05 DCs
- 1.28% ± 2.54 B cells
- 0.89% ± 2.11 Monocytes
- 0.33% ± 0.41 Bulk Macrophages
- 0.19% ± 0.25 Neutrophils

6.41% ± 6.15 Bulk CD4$^+$ T cells

4.00% ± 5.34 Bulk CD8$^+$ T cells

Other Neural/Glial Cell Types Not Reported — 49.83%

28.5% ± 33.0

TMEM119$^+$Microglia

Risk of bias

- Sequence Generation
- Baseline Characterisitcs
- Allocation Concealment
- Random Housing
- Blinding
- Random Outcome Assessment
- Incomplete Outcome Data
- Selective Outcome Reporting
- Other Sources of Bias
- **Overall**

0%   25%   50%   75%   100%

High   Unclear   Low