The CPSF phosphatase module links transcription termination to chromatin silencing

Eduardo Mateo-Bonmatí[1,3], Xiaofeng Fang[1,4], Robert Maple[1], Marc Fiedler[2], Lori A Passmore[2], and

Caroline Dean[1]*

[1]John Innes Centre, Norwich Research Park, Norwich NR4 7UH, United Kingdom;

[2]MRC Laboratory of Molecular Biology, Cambridge CB2 0QH, United Kingdom;

[3]Present address: Centro de Biotecnología y Genómica de Plantas, Universidad Politécnica de Madrid
(UPM) – Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA)/CSIC, 28223
Pozuelo de Alarcón, Madrid, Spain;

[4]Present address: Centre for Plant Biology, School of Life Sciences, Tsinghua University, Beijing
100084, China.

*Correspondence to: Caroline Dean (caroline.dean@jic.ac.uk).

Keywords: RNA processing, chromatin, transcription, *FLC*, *COOLAIR*, noncoding RNA, CPSF

14
15
ORCID (E.M.-B.): 0000-0002-2364-5173   @emateobonmati
ORCID (X.F.): 0000-0003-1161-9186   @XiaofengFangLab
ORCIDC (R.M.): 0000-0002-9175-5024   @Bobby_Mapes
ORCID (M.F.): 0000-0003-4269-7873
ORCID (L.P.): 0000-0003-1815-3710   @lapassmore
ORCID (C.D.): 0000-0002-6555-3525   @CarolineDeanLab
22

Word count (total): 9214

ABSTRACT

RNA-mediated chromatin silencing plays important roles in development and environmental response

in many organisms. However, it is still unclear how the RNA, and/or the transcriptional processes

producing them, trigger chromatin silencing. Through the study of developmental timing in plants we

have found co-transcriptional RNA processing modulates a chromatin regulatory mechanism

controlling quantitative expression of Arabidopsis *FLOWERING LOCUS C* (*FLC*). Here, we show

that the phosphatase module of the mRNA 3' processing CPSF complex is central to this mechanism.

The phosphatase module components APRF1 (yeast Swd2/human WDR82), TOPP4 (Glc7/PP1) and

LD (which we show is functionally equivalent to Ref2/PNUTS) all function genetically downstream

of the CPSF-mediated cleavage and polyadenylation and are necessary for chromatin modification

that results in *FLC* transcriptional silencing. LD has been previously shown to stoichiometrically co-

associate *in vivo* with the histone demethylase FLD. We show both proteins result in removal of

H3K4 monomethylation in the central section of the *FLC* locus. This links transcription termination to

delivery of chromatin modifications that then influence subsequent transcription. Thus, physical

association of the CPSF phosphatase module with chromatin modifiers generates a transcription-

coupled silencing mechanism. The conservation of the factors involved suggest this mechanism may

be generally relevant to transcription-mediated chromatin silencing.

INTRODUCTION

How chromatin and transcription interact to regulate gene expression is a major area of interest.

Extensive feedback mechanisms make this interaction difficult to dissect. Chromatin affects co-

transcriptional processes that influence RNA Polymerase II (RNA Pol II) elongation rate, which can

lead to alternative splicing and early termination/processivity changes i.e., the likelihood of

transcription reaching the end of the gene. In turn, these co-transcriptional processes feed-back to

influence the chromatin state (Berry *et al.* 2015, Muniz *et al.* 2021, Zumer *et al.* 2021). One locus

where this complexity has been studied in detail is Arabidopsis *FLC*. *FLC* encodes a floral repressor

and quantitative variation in *FLC* transcription has been an important determinant in adaptation of

Arabidopsis accessions to a wide range of climates. For example high *FLC* expression necessitates an

overwintering reproductive strategy through winter-induced epigenetic silencing of *FLC* (Menon *et*

al. 2021). In contrast, low *FLC* expression leads to a rapid-cycling reproductive strategy, enabling

multiple generations a year in some climates.

There has been extensive analysis of the components regulating *FLC* transcription. Forward genetic

screens identified repressors that include RNA binding proteins, RNA 3' processing factors and

chromatin modifiers (Wu *et al.* 2020). These factors have been found to promote proximal

termination of both the antisense transcript *COOLAIR*, but also the sense *FLC* transcript in the

embryo (Schon *et al.* 2021). Suppressor genetics and chromatin analysis showed that these co-

transcriptional activities function through the chromatin modifiers FLD (a H3K4 demethylase)-LD

(TFIIS domain protein)-SDG26 (a SET domain protein) (Liu *et al.* 2007, Fang *et al.* 2020) to suppress

FLC. Proximal termination induces histone H3K4 demethylation across the locus, delivered by FLD,

LD and SDG26. These three factors tightly associate with each other *in vivo* based on proteomic

interactions, physically associate with Cleavage and Polyadenylation Specificity Factor (CPSF; CPF

in yeast) factors after cross-linking (Fang *et al.* 2020), and together reduce H3K4me1 and H3K36me3

accumulation at *FLC*. This antagonizes transcription enabling a switch to PRC2-mediated H3K27me3

accumulation, so reducing transcriptional initiation and elongation rates (Wu et al 2016). The *FLC*

silencing mechanism thus involves co-transcriptional processes that mediate chromatin modifications,

which in turn feedback to reinforce the co-transcriptional processing. The factors involved are all

evolutionarily conserved, raising the likelihood that this mechanism will be broadly relevant for gene

regulation in many organisms; for example, there are parallels with mammalian aging processes

(Gyenis *et al.* 2023).

A key question that remains is how proximal termination delivers a changed chromatin environment.
Here, we used proteomic immunoprecipitation experiments to identify additional partners of the FLD-
LD-SDG26 complex. We find robust interaction with Arabidopsis APRF1 and TOPP4. These proteins
are homologous to different members of the CPF phosphatase module (Swd2 and Glc7) that
associates with RNA Pol II and mediates transcription termination and RNA polyadenylation in yeast
(Carminati *et al.* 2022). We use molecular and genetic tools to show that these components function
in the same co-transcriptional pathway with the FLD-LD-SDG26 complex to alter *FLC* chromatin
environment, and thus transcriptional output. Upon close examination of all the phosphatase module
components we also show that the LD component of the complex is structurally related to Ref2 in
yeast and PNUTS in mammals; they all contain intrinsically disordered regions and a TFIIS domain.
Ref2/PNUTS act in yeast/mammalian complexes as the regulatory subunit of the phosphatase module
that influences phosphorylation of the RNA Pol II and its partners via the Glc7/PP1 phosphatase. This
activity promotes transcription termination through phosphorylation-dependent effects on RNA Pol II
elongation/processivity (Rodríguez-Molina *et al.* 2023). We show that LD-APRF1-TOPP4 have
similar functional consequences promoting proximal termination of the antisense *COOLAIR*
transcripts. Thus, our proteomic and genetic dissection identifies many factors of this floral repression
pathway as conserved members of the 3' CPSF phosphatase module. This clarifies the overall
mechanism of how RNA processing is linked to chromatin silencing at Arabidopsis *FLC*. The CPSF
phosphatase module directly links RNA processing with histone demethylase activity to alter the local
chromatin environment as the RNA Pol II transcribes downstream of the polyadenylation site. This
chromatin environment reinforces proximal termination choice so providing the molecular feedback
necessary to stably maintain a low transcription state. Our accompanying paper describes a
mathematical model of this co-transcriptional regulatory mechanism revealing how co-transcriptional
3' processing sets the level of productive transcription via H3K4 monomethylation, thus dictating the
degree of antagonism to Polycomb silencing, which then determines the rate at which Polycomb
repression is established (Menon *et al.* 2023). Overall, our studies shed light on the tight
interconnections/feedbacks between co-transcriptional RNA processing, chromatin environment and
transcriptional mode. These have proven difficult to dissect at the molecular level but are likely to be
at the basis of most epigenetic switching mechanisms.
105

RESULTS

The FLD complex robustly associates with the homolog of yeast Swd2.
We previously reported that the histone demethylase homolog FLD (FLOWERING LOCUS D)
associates with LD (LUMINIDEPENDENS) and SDG26 (SET DOMAIN GROUP 26) *in vivo*; each
tagged version of these three proteins stoichiometrically enrich the other two partners in co-
immunoprecipitation experiments (Fang *et al.* 2020). Each of these proteins also co-

immunoprecipitated with ANTHESIS PROMOTING FACTOR1 (APRF1; Figure 1A). This was also found in another analysis of proteins that co-immunoprecipitate with the FLD complex [Figure 1B; Qi et al. (2022)].

APRF1 is a WD40-repeat protein encoded by At5g14530. We obtained a T-DNA insertion line [WiscDsLox_489; *aprf1-9*; Kapolas et al. (2016)] and analysed the flowering time and *FLC* expression. Both *FLC* spliced and unspliced transcripts were significantly upregulated in the mutant line, and accordingly, *aprf1-9* plants were late flowering (Figure 1C-E). Given that the insertion is predicted to interrupt the 9^{th} of 10 exons, the plants might retain some APRF1 function. To address this, we designed a CRISPR-Cas9 transgene to generate a mutant null for *APRF1*. Using a sgRNA targeting the second exon and screening for edited plants, we found transgene-free T_2 plants carrying a 5-nt deletion which creates an in frame early stop codon (Figure S1). This new allele was named *aprf1-10* and was as late flowering as *aprf1-9* (Figure 1D). We crossed *aprf1-9* to *aprf1-10* and analysed *FLC* expression levels of the single and heterozygous mutants. All of them had similarly upregulated *FLC,* confirming their allelism and the role of APRF1 on *FLC* repression (Figure 1E).

APRF1 is one of the two Arabidopsis orthologs of the *Saccharomyces cerevisiae* Swd2 [(Figure S2A)(Fiorucci *et al.* 2019)]. Swd2 has been found to play a role in two very different complexes; it is one of the WD40 components of the COMPlex of Proteins Associated with Set1 (COMPASS). This complex catalyses the co-transcriptional deposition of H3K4me3 and Swd2 has been shown to play a key role in the interaction between Set1 and the RNA Pol II Carboxy Terminal Domain (CTD) (Bae *et al.* 2020). Swd2 has also been found as a structural component of the phosphatase module of the Cleavage and Polyadenylation Factor (CPF; CPSF in higher eukaryotes), which signals transcriptional termination (Casanal *et al.* 2017). This apparently opposite role on transcription in yeast made us test if the Arabidopsis orthologs *APRF1* (also known as *Swd2-like A/S2LA*) and *S2LB* have functionally diverged. *FLC* spliced and unspliced levels were assayed in an insertional allele of *S2LB* (Figure S2B) and the double mutant *aprf1-9 s2lb*. Opposite to *aprf1-9*, *s2lb* showed a significant reduction in *FLC* expression, while the double mutant shows up-regulated *FLC* indistinguishable from the *aprf1-9* single mutant (Figure S2C). These contrasting phenotypes suggest that in Arabidopsis, the ancestral Swd2 may have been sub-functionalized, with S2LB working through the COMPASS complex and therefore activating *FLC,* and APRF1 working with the 3' processing machinery to silence the locus (Figure S2D).

FLD-mediated repression of *FLC* requires H3K4me1 removal (Fang *et al.* 2020, Inagaki *et al.* 2021). Considering the tight link between the FLD complex and APRF1, we performed chromatin immunoprecipitation coupled with quantitative PCR (ChIP-qPCR) to quantify H3K4me1 at *FLC*. We found a H3K4me1 increase across the locus in *aprf1-10* compared to Col-0 (Figure 1G). H3K4me1 is thought to be read by SDG8 (Liu and Huang 2018), a histone methyltransferase that deposits

H3K36me3, an active histone modification, which is mutually exclusive to the repressive PRC2-
deposited H3K27me3 (Yang *et al.* 2014). Consistent with the *FLC* upregulation and the increased
H3K4me1 levels in *aprf1-10*, ChIP-qPCR analyses also showed that H3K36me3 and H3K27me3
were upregulated and downregulated, respectively. Thus, APRF1 has a role in establishing a silent
chromatin status at *FLC* (Figure 1H, I).

Working upstream of the FLD complex, FCA promotes proximal termination on both strands of *FLC*
(Simpson *et al.* 2003, Schon *et al.* 2021). To obtain genetic evidence of the relationship between
APRF1 and *FLD, LD* or *FCA*, we crossed *aprf1-9* with mutants in subunits of the FLD complex such
as *fld-4* or *ld-1*, and the core component of the pathway *fca-9*. *FLC* levels in the double mutants
revealed an epistatic relationship between *FLD, LD* or *FCA* and *APRF1* (Figure 1J, K), further
demonstrating the genetic connection between *APRF1* and the *FLC* silencing machinery. To study the
effects of loss of *APRF1* in a high transcriptional environment, but without perturbing the FCA
pathway, we introgressed a functional *FRIGIDA* (*FRI*) into *aprf1-9*. FRI functions as an *FLC*
transcriptional activator, and like other anti-terminators (Gregersen *et al.* 2019), promotes distal
polyadenylation of both *FLC* and *COOLAIR*, thus antagonizing the co-transcriptional repression
mechanism (Schon *et al.* 2021). Levels of *FLC* in *aprf1-9 FRI* were the same as Col*FRI,* likely due to
an overwriting effect of *FRI* compared to the relatively modest *FLC* upregulation of *aprf1-9* (Figure
1L, M). Finally, we crossed *aprf1-10* into the C2 background which harbours a *35S_{pro}:FCA* transgene
in the presence of a functional *FRI*. *aprf1-10* significantly released the *35S_{pro}:FCA* transgene-
mediated *FLC* silencing convincingly demonstrating a role for APRF1 downstream of FCA (Figure
1N, O).

APRF1, LD, and TOPP4 form a plant equivalent of a CPSF phosphatase sub-module.
To further characterize the role of APRF1 on *FLC* silencing we generated a translational fusion of
APRF1 (*APRF1_{pro}:APRF1-VENUS*) and performed CrossLinked Nuclear ImmunoPrecipitation and
Mass Spectrometry (CLNIP-MS) (Fang *et al.* 2020), detecting LD in the pulldown, thus confirming
their reciprocal interaction (Figure 2A). We also detected a significant number of peptides for TYPE
ONE SERINE/THREONINE PROTEIN PHOSPHATASE4 (TOPP4) (Figure 2A). These interactions
were recently validated by independent experiments of differentially tagged lines [(Qi et al. 2022;
Figure 2B)].

180
TOPP4 shares high sequence and structural homology with Glc7, another member of the CPF
phosphatase module in yeast (Figure S3A). The similarity was even higher than that showed by
APRF1 and Swd2 (Figure S4). Identifying homologues of two members of the CPF phosphatase
module prompted us explore further parallels. Swd2^{APRF1} and Glc7^{TOPP4} are tightly associated with
each other in yeast, and to Ref2 (Casanal *et al.* 2017). Ref2 is key for the interaction between CPF and

the RNA Pol II and behaves as a phosphatase regulatory subunit, potentially providing substrate

specificity to the phosphatase Glc7 (Nedea *et al.* 2008, Carminati *et al.* 2022). Ref2 and its putative

metazoan ortholog PNUTS, are largely disordered proteins apart from a TFIIS motif at the N terminal

region of the protein which consists of a compact four-helix bundle. Intriguingly, despite its apparent

key role in an otherwise highly conserved 3' processing machinery, Ref2 shows no obvious homology

with an Arabidopsis protein. We performed BLAST searches to find potential orthologs in

Arabidopsis for PNUTS, the counterpart of Ref2 in mammals. BLAST algorithms showed the best hit

for PNUTS in Arabidopsis is LD (Figure S5). We then aligned the sequence of Ref2, PNUTS and LD.

The degree of conservation is low (Figure S6A), although slightly improved when aligning only LD

and PNUTS (Figure S7). However, all of them share similar features, a TFIIS domain in the N-

terminal region of the protein and a highly disordered region (Figure 2D, S6B-D). PNUTS is known

to provide substrate specificity to PP1 phosphatases, like Glc7 and TOPP4 (Figure S3), and to interact

with WDR82, a close ortholog of Swd2/APRF1 (Figure S4). Further exploring these parallels, we

carried out an *in-silico* prediction using AlphaFold2 of the interaction between APRF1-LD and their

yeast and human counterparts (Figure 2E). Despite the low sequence homology, predictions support

conservation of structural features and contact points between APRF1/Swd2/WDR82 and

LD/Ref2/PNUTS supporting their functional equivalence (Figure 2F). Taken together, the robust

immunoprecipitation of LD and TOPP4 by APRF1, the structural parallels between Ref2-PNUTS-LD

and the similar roles in co-transcriptional processing lead us to propose that LD, APRF1, and TOPP4

may form a functional CPSF phosphatase module in plants.

FLD complex and RNA Pol II co-occupy *FLC* chromatin independently of FCA function.

Available genome-wide data shows FLD binding correlates with actively transcribed genes and

elongating RNA Pol II CTD phosphorylated at Ser2 or Ser5 (Inagaki *et al.* 2021). RNA Pol II

occupancy in the Arabidopsis genome often shows peaks near transcription termination sites (TTS),

potentially linked to slow co-transcriptional termination events (Wu *et al.* 2016). We found this was

the case for *FLC* in Col*FRI* vs Col-0 (Figure 3A) and has been shown to be the case in *fca-9* or *fld-4*

(Wu *et al.* 2016). To address co-occupancy with FLD, we performed ChIP-qPCR using a transgenic

FLAG-tagged FLD with and without the transcriptional activator FRI, and non-transgenic control

plants. Agreeing with the reported data (Inagaki *et al.* 2021), FLAG-FLD showed high enrichment at

the 3' region of *FLC* in a *FRI* genotype compared to Col-0 (*fri*), with the latter close to the

background signal (Figure 3B). To rule out the possibility that this enrichment was an effect of the

transcriptional activator FRI and not the high transcription itself, we introgressed the FLAG-FLD

transgene into the *fca-9* background. FLD enrichment at the 3´end of *FLC* in *fca-9 FLAG-FLD* was

indistinguishable from *FRI FLAG-FLD* confirming FLD enrichment primarily associated to

transcriptional activity (Figure 3B). *FLD* functions genetically downstream of *FCA* (Liu *et al.* 2007)

so it was interesting that FLD-Pol II co-occupancy association was not affected by *fca-9*. Thus, even

when FLD is located at *FLC*, it cannot function properly without FCA-mediated 3' processing of the
nascent transcript.

We introgressed a GFP-tagged LD (Fang *et al.* 2020) into Col*FRI* to create a line with high *FLC*
transcription and performed ChIP-qPCR experiments. LD enrichment was only detected in the line
where *FLC* transcription is high, peaking at the *FLC* TTS (Figure 3C). The shared enrichment of
elongating Pol II, FLD- and LD is consistent with an *in vivo* association of the transcriptional
machinery and the FLD complex as suggested by genome-wide data (Inagaki *et al.* 2021). Thus, we
propose that FLD and LD associate with RNA Pol II, and therefore work co-transcriptionally, fitting
with LD working as a CPSF equivalent to Ref2 and PNUTS.

Inefficient termination in *aprf1* mutants leads to transcriptional read-through.
Previous work on the FCA pathway found that most of the factors involved were directly affected on
FLC antisense transcription (Wu *et al.* 2020). To pursue a potential role of LD-APRF1-TOPP4 as a
CPSF phosphatase module we analysed the transcription and processing of *COOLAIR*. In contrast to
FLC levels which did not show any significant difference (Figure 1J, K), total *COOLAIR* levels were
upregulated in double mutants containing *aprf1-9* (Figure 4A, B). This was particularly striking for
the *aprf1-9 FRI* combination. *COOLAIR* transcripts are generally polyadenylated at either a cluster of
proximal sites (Class I) or a cluster of distal sites coincident with the *FLC* promoter (Class II) (Liu *et*
al. 2010). A moderate increase in proximal *COOLAIR* (Figure 4C) and no change in distal *COOLAIR*
were found in *aprf1-9 FRI* (Figure 4D), with the proximal / distal ratio increased in *aprf1-9* double
mutant combinations compared to the single mutant (Figure 4E), unlike other mutations affecting the
FCA pathway (Xu *et al.* 2021a). However, like *PEP1* in *Arabis alpina* (Castaings *et al.* 2014), we had
also detected low abundance spliced *COOLAIR* transcripts polyadenylated at a medial site (Liu and
Dean, unpublished), which we had termed *COOLAIR* Class III (Figure 4A). This isoform was
significantly up-regulated specifically in *aprf1-9 FRI* (Figure 4F), with one specific spliced variant
COOLAIR Class III.3 becoming the most abundant isoform, enriched in high transcription situations
such as *fca-9* or Col*FRI* (Figure 4G, S8). We interpret the high levels of the *COOLAIR* class III
isoform as reflecting inefficient polyadenylation/transcript termination at the proximal site. To
analyse the polyadenylation site choice in an unbiased and strand-specific manner in these two
genotypes, we carried out a Quant-seq analysis, which detects polyadenylation events, of Col*FRI* and
aprf1-9 FRI. For the *FLC* (sense) transcripts there were a few reads potentially indicating
transcriptional readthrough events (Figure S9). However, the differences for *COOLAIR* were more
dramatic with reads compatible with the use of the medial poly A site, others pointing to distal
COOLAIR readthrough, and even alternative *COOLAIR* transcriptional start sites, potential
consequence of *FLC* sense readthrough (Figure 4H, S9).

To understand which were likely to be co-transcriptional changes we analysed chromatin-bound RNA

(chRNA) of Col-0, *aprf1-9*, Col*FRI* and *aprf1-9 FRI*. Not unexpectedly, the *FLC* (sense) chRNA

levels of *aprf1-9* were higher than Col-0, as *aprf1-9* compromises *FLC* silencing. PCR amplicons for

sense *FLC* levels in Col*FRI* and *aprf1-9 FRI* gave similar values at the 5' and 3' ends of the locus, but

there were differences in the centre, pointing to complex effects of APRF1 on *FLC* Pol II processivity

(Figure 5A). However, the most striking differences were on the *COOLAIR* strand (antisense) (Figure

5A). RNA was detected in regions corresponding to the 5' half of *COOLAIR* in Col*FRI*, but in *aprf1-9*

genotypes (*fri* and *FRI*) this signal extended to regions that cover both Class I and Class III *COOLAIR*

transcripts. This likely represents antisense transcriptional readthrough, supporting a role for APRF1

as part of the Pol II termination machinery (Figure 5A). Overall, the loss of *APRF1* appears to affect

the ability of CPSF to facilitate transcription termination leading to transcriptional readthrough.

LD and APRF1 affect RNA Pol II CTD Ser5 hyperphosphorylation

Different mechanistic models, not mutually exclusive, have been proposed for transcription

termination (Mo *et al.* 2021, Rodríguez-Molina *et al.* 2023). Current thinking based on studies in

different organisms is that CPSF-RNA Pol II recognition of polyadenylation site (PAS) triggers the

nascent RNA cleavage. This generates a free 5' end on the cleaved RNA which is a substrate for 5'-3'

ribonucleases such as XRNs that degrade the nascent transcript and eventually dislodge RNA Pol II

from the chromatin (Cortazar *et al.* 2019, Rodríguez-Molina *et al.* 2023). Transcription of the PAS

also triggers a conformational change in RNA Pol II, potentially driven by dephosphorylation of the

CTD and or co-factors, which slows-down RNA Pol II. To ascertain if *aprf1* mutants affect RNA Pol

II CTD modifications, we carried out ChIP-qPCR experiments using antibodies targeting the RNA

Pol II CTD phosphorylated residues Ser2, Ser5 and Tyr1 comparing *aprf1-10* with Col-0. Not

surprisingly, all the modified RNA Pol II were detected at higher levels in *aprf1-10,* since the locus is

more actively transcribed (Figure 5B, S10A). We repeated our analysis comparing Col*FRI* and *aprf1-*

9 FRI, given that both genotypes have similar *FLC* RNA levels. In this case, RNA Pol II Ser2P and

Tyr1P levels were identical (Figure 5C, S10B), while there was a slight increase in RNA Pol II Ser5P

towards the 5' end of the locus (Figure 5C). FRI forms a super-molecular complex recruiting

transcriptional activators and repressing proximal termination so may mask APRF1 direct effects on

RNA Pol II phosphorylation status (Li *et al.* 2018, Zhu *et al.* 2021). Thus, to investigate direct effects

of APRF1-LD (a potential CPSF phosphatase module) on RNA Pol II CTD phosphorylation we

compared the FRI mutant Col-0 [which is a *fri* null (Johanson *et al.* 2000)], *ld-1* and *aprf1-9 ld-1*.

Both, *ld-1* and *aprf1-9 ld-1* showed similarly high *FLC* expression levels (Figure 1J, K), similar levels

of total Pol II, Ser2P and Tyr1P (Figure 5D, S10C), but *aprf1-9 ld-1* showed higher RNA Pol II Ser5P

(Figure 5D). This suggests a role for the Arabidopsis CPSF phosphatase module in removal of RNA

Pol II Ser5P at *FLC,* however, we cannot rule out the possibility that the CTD Ser5 phosphorylation

effects are indirect, through phosphorylation of elongation factors such as SPT5 (Cortazar *et al.*
2019).
298

DISCUSSION

The study of developmental timing in plants has led to mechanistic dissection of chromatin silencing
mechanisms at the gene encoding the floral repressor Arabidopsis FLC. Since quantitative variation of
FLC expression affects the reproductive strategy of Arabidopsis any molecular variation can be
subject to strong evolutionary selection. *FLC* is thus an excellent system to dissect RNA processing,
chromatin regulation, and their interconnections and molecular feedbacks that generate low and high
transcriptional states that underpin adaptively important variation in transcriptional output.
306

The work here identifies the importance of the CPSF phosphatase module, a regulator of RNA Pol II,
in *FLC* regulation. We identify LD, a protein characterized as a flowering regulator nearly 20 years
ago as part of the CPSF phosphatase module, functionally equivalent to yeast Ref2 and metazoan
PNUTS proteins. Given its interaction partners, LD thus acts as a bridge between chromatin modifiers
and the Pol II-machinery (Figure 6). We propose that actively transcribed *FLC* chromatin is poised
with H3K4me1, which promotes RNA Pol II processivity, i.e. the likelihood of transcription reaching
the end of the gene. Any pause in RNA Pol II functioning, for example coincident with the formation
of the R-loop during *COOLAIR* transcription, would stimulate the 3' processing machinery (carried
along in the RNA Pol II supercomplex) to polyadenylate the transcript at the proximal site. PAS
recognition would be signalled by the CPSF phosphatase module to Pol II via dephosphorylation of
either elongating factors or the CTD or both and trigger a conformational change on RNA Pol II (as
proposed by Carminati *et al.* 2022) to activate FLD function. RNA Pol II downstream of the PAS
would proceed slowly due to the conformational change, causing FLD to remove H3K4me1 co-
transcriptionally until the RNA Pol II is terminated by 5'-3' XRNs. This creates a less processive
chromatin environment for subsequent rounds of transcription. Each round of proximal
polyadenylation-termination-H3K4me1 removal will reinforce the next round of transcription,
creating an intrinsic feedback loop in the mechanism (Menon *et al.* 2023). Intriguingly, our chRNA
data detected very clear readthrough effects on the antisense (*COOLAIR*) strand, but also in certain
parts of the sense (*FLC*) transcription unit when *APRF1* was disrupted. Proximal termination of *FLC*
sense transcription during early embryo development is associated with establishment of the silenced
state (Schon *et al.* 2021). Proximally polyadenylated sense transcripts do not accumulate in seedling
tissue, but it is possible that these are particularly sensitive to RNA degradation pathways. We thus
envisage the same mechanism functioning on both *FLC* and *COOLAIR* transcription to both establish
and then maintain the transcriptionally silenced state, with the conserved CPSF phosphatase factors
central to that chromatin silencing mechanism.
332

Arabidopsis genome-wide data indicates that FLD shows a clear enrichment at sites of convergent

transcription (Inagaki *et al.* 2021) and CPSF factors have been shown to resolve DNA

transcription/replication conflicts in the Arabidopsis genome (Baxter *et al.* 2021). Thus, FCA and the

CPSF may constitute a plant equivalent to the metazoan Integrator-PP2C (INTAC) (Hu *et al.* 2023,

Wagner *et al.* 2023). Indeed, FCA and FPA, have been previously shown to play a role in defining the

transcriptional boundaries, with transcriptional readthrough in hundreds of genes detected in the

double mutant *fca fpa* (Sonmez *et al.* 2011).

The evolutionary variation in the co-transcriptional complexes provides regulatory flexibility. *APRF1*

and *S2LB* constitute an unusually clear example of gene sub-functionalization. In yeast, Swd2 is a

single copy gene whose product was found in two protein complexes with antagonistic functions:

COMPASS (Bae *et al.* 2020) and the mRNA 3'-end processing machinery (Casanal *et al.* 2017).

Likely due to this dual role, Swd2 is the only essential COMPASS component in yeast (Soares and

Buratowski 2012). S2LB was found to interact *in vivo* with the methyltransferase SDG2 and the

structural component of COMPASS WDR5, controlling H3K4me3 levels genome-wide at thousands

of gene promoters (Fiorucci *et al.* 2019). In contrast, APRF1 interacts with LD as well as the other

components of the FLD complex, and with TOPP4, functionally equivalent to the CPF phosphatase,

Glc7 (Fang *et al.* 2020, Qi *et al.* 2022). The functional divergence suggested by the differential *in vivo*

association was confirmed through analysis of mutants affecting *FLC* expression. Like other mutants

in the Arabidopsis COMPASS machinery (Jiang *et al.* 2011), *s2lb* mutants reduce *FLC* expression

due to their impaired ability to deposit H3K4me3 at the locus and promote transcriptional initiation.

The accompanying paper (Menon *et al.* 2023) describes a theoretical analysis producing a dynamic

mathematical model of this co-transcriptional mechanism. The model quantifies the molecular

feedbacks linking co-transcriptional processing, chromatin modification and transcriptional output in

an interlinked three-way association that sets the timescale for the slow switch into the digital

Polycomb silenced state. The model explains how the low transcriptional state influenced by proximal

termination and H3K4 demethylation reduces the antagonism to Polycomb silencing and leads to a

stable PRC2 epigenetic switch, with sufficient feedbacks to maintain the silenced state through DNA

replication and cell division. We have generated an animation (weblink) to help explain these

molecular feedbacks, and how this mechanism leads to the stable expression of *FLC* in one of two

stable expression states.

This transcription-coupled silencing mechanism is likely to be generally relevant to heterochromatin

establishment in genomes of many eukaryotic organisms. A clear genome-wide correlation has been

found between human FLD and APRF1 homologs (LSD1 and WDR82) and Pol II (Kim *et al.* 2022,

Estell *et al.* 2023), and an unbiased search for LSD1 regulators in human cells found a direct

relationship between the histone demethylase, and an RNA helicase involved in R-loop resolution and
PRC2-mediated silencing (Pinter *et al.* 2021). Indeed, 3' processing has been extensively linked to
chromatin silencing in *S. pombe* (Kowalik *et al.* 2015, Grewal 2023). Continued mechanistic
dissection is therefore likely to elaborate concepts important in chromatin silencing generally.
374

MATERIALS AND METHODS

Plant material and growth conditions

All the plants were homozygous for the indicated genotype. The *aprf1-9* mutant seeds
(WiscDsLox489-492K11, N858279) were obtained from NASC (Nottingham Arabidopsis Stock
Centre). The *s2lb* mutant and the *aprf1-9 s2lb* double mutant, previously described in (Fiorucci *et al.*
2019), were kindly provided by Fredy Barneche. Transgenic *fld-4*; *FLD_{pro}:3xFLAG-FLD* were
previously described in (Inagaki *et al.* 2021) and shared by Soichi Inagaki.

The *aprf1-10* mutant harbours a 5-nt deletion in *APRF1*. The *aprf1-10* deletion creates a new target
for the MaeII (target ACGT) restriction enzyme. To generate the *aprf1-10* we employed the
CRISPR/Cas9 plasmid pKI1.1R following the protocol described (Tsutsui and Higashiyama 2017).
Briefly, pKI1.1R plasmid (Addgene #85808) was linearized by incubating 1.5 µg of the plasmid with
AarI restriction enzyme for 16 h, and then dephosphorylated using the alkaline phosphatase rAPid
(Roche). A target-specific gRNA was designed using CRIPR-P 2.0
(http://crispr.hzau.edu.cn/CRISPR2). Oligonucleotides harbouring the gRNA target
(sgRNA_APRF1_F and sgRNA_APRF1_R; Table S1) were hybridised by slow cooling down from
95-25°C and then phosphorylated using the T4 Polynucleotide Kinase (NEB). The digested plasmid
and the hybridised oligonucleotides were ligated using the T4 ligase (NEB) and then transformed in
Escherichia coli HST08 competent cells (Takara). The sequence integrity of inserts carried by
transformants were verified by Sanger sequencing. The plasmid was then transfer to *Agrobacterium*
tumefaciens C58C1 strain by electroporation. T$_1$ plants carrying the construct were selected on MS
media supplemented with 15 µg/ml of Hygromycin. Next generation plants were counter-selected to
find transgene-free individuals carrying the homozygous mutation. The *aprf1-9 APRF1_{pro}-APRF1-*
mVENUS transgenic was generated by cloning the *APRF1pro:APRF1* genomic sequence without the
stop codon upstream of the coding sequence of *mVENUS* at pCAMBIA1300. Vector was mobilized to
Agrobacterium C58C1 and transform to *aprf1-9* by floral dipping. T$_1$ transformant plants were
selected on Hygromycin.

Seeds were surface sterilized in 40 % v/v commercial bleach for 10 min and rinsed 4 times with
sterile distilled water. Seeds were then sown on standard half-strength Murashige and Skoog (MS)
medium (0.22% MS, 8% plant agar) media plates and kept at 4°C in darkness for 3 days before being
transferred to long day photoperiod conditions (16 h of light, 8 h dark). All RNA and protein
experiments were done using 14-days old seedlings unless otherwise specified.
406

Gene expression analyses

Seedlings were harvested, and RNA was extracted with the hot phenol method as previously described (Zhu *et al.* 2021). TURBO DNase (Ambion) was used to remove genomic DNA contamination before reverse transcription. cDNA was synthesized using SuperScript IV (Invitrogen) and gene-specific primers (Table S1). qPCR analyses were performed, and data was normalized to the indicated housekeeping gene or genes.

ChIP

2.5 gr of seedlings were crosslinked with 1% formaldehyde in 1X PBS for 12 min by vacuum infiltration, followed by addition of glycine (final concentration 125 mM) with another 7 min of vacuum infiltration. Tissue was then ground to fine powder with liquid nitrogen. Ground tissue was resuspended in 35 mL of Honda Buffer (20 mM Hepes, 0.44 M sucrose, 1.25 % Ficoll, 2.5% Dextran, 10 mM $MgCl_2$, 0.5% Triton X-100, 5 mM DTT, 1x Roche protease inhibitor mixture), filtered through two layers of Miracloth, and centrifuged at 2500 xg for 15 min. Nuclei pellet was then washed once more with 1.6 mL of Honda Buffer.

For histone ChIP, nuclear pellets were resuspended in Nuclei Lysis Buffer (50 mM Tris-HCl pH 8, 10 mM EDTA, 1 % SDS), and sonicated 4 x 5 min (30 sec ON/ 30 sec OFF) using a Diagenode Bioruptor on Medium setting. IP was performed by incubating 140 µl of sonicated chromatin diluted ten times with ChIP dilution buffer (16.7 mM Tris-HCl pH 8, 1.2 mM EDTA, 1.1 % Triton X-100, 167 mM NaCl) with 15 µl of Protein A-coated Dynabeads (Invitrogen) previously incubated for 2 h with either 2.5 µg of anti-H3 (ab176842), anti-H3K27me3 (ab192985), anti-H3K36me3 (ab9050), or H3K4me1 (ab8895) and incubated overnight at 4°C on a rotator wheel.

For FLAG-FLD and GFP-LD ChIP, nuclei were obtained as described above, but for FLAG-FLD the crosslinking buffer was supplemented with 1.5 mM of EGS [ethylene glycol bis(succinimidyl succinate); ThermoFisher]. Nuclear pellets were suspended in RIPA buffer (50 mM Tris-HCl, 150 mM NaCl, 1% Nonidet P-40, 0.5% NaDeoxycholate, 0.1% SDS, 1x Roche protease inhibitor mixture) and sonicated 5 times x 5 min (30 s ON/ 30 s OFF) with the Bioruptor in high setting. Undiluted chromatin was incubated overnight at 4°C with either 1.5 mg Dynabeads M-270 Epoxy preincubated with 1.5 µl anti-FLAG (Anti-FLAG® M2 / F1804, Merck) for FLAG-FLD or 15 µl Protein-A coated Dynabeads with 2.5 µl anti-GFP (ab290, Abcam), for GFP-LD.

Beads were then washed twice with Low Salt Wash Buffer (150 mM NaCl, 0.1 % SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8, 1x Roche protease inhibitor mixture), twice with High Salt Wash Buffer (500 mM NaCl, 0.1 % SDS, 1 % Triton X-100, 2 mM EDTA, 20 mM Tris-HCl 8, 1x Roche protease inhibitor mixture), and twice with TE wash buffer (10 mM Tris-HCl pH 8, 1 mM EDTA, 1x Roche protease inhibitor mixture).

For RNA Pol II ChIP, nuclei were obtained as described for histone ChIP, complementing the Honda Buffer with 1x of PhosSTOP (Merck). Nuclear pellet was suspended in 1 mL of TAP buffer (100 mM

NaCl, 20 mM Tris-HCl pH 8, 2.5 mM EDTA, 10 % Glycerol, 1 % Triton, 1x of PhosSTOP, 1x Roche

protease inhibitor mixture) and given 20 strokes with the Dounce Homogenizer. The resulting

solution was sonicated 4 x 10 min (15 s ON/ 45 s OFF) with the Bioruptor in low setting. 250 µl of

undiluted chromatin was incubated overnight at 4°C with either preincubated Protein-G coated

Dynabeads with 2 µl anti-Total Pol II (C15200004, Diagenode) or Dynabeads M-270 Epoxy

preincubated with anti-Tyr1P (MABE350, Merck), anti-Ser2P (C15200005-50, Diagenode), or anti-

Ser5P (C15200007-50, Diagenode). Then beads were washed twice for 15 min with Low Salt and

High Salt buffers like for histones but including 1x of PhosSTOP. Then beads were washed for 15

min with the LiCl buffer (250 mM LiCl, 0.5 % NP40, 2.5 mM EDTA, 0.05 % NaDeoxycholate, 20

mM Tris-HCl pH 8, 1x of PhosSTOP, 1x Roche protease inhibitor mixture), and the TE buffer (same

as for histones plus 1x PhosSTOP). In all cases, after IP, DNA was then eluted and reverse-

crosslinked by incubating the beads at 95°C for 10 min in presence of 100 µl of 10 % Chelex resin

(BioRad), treated with Proteinase K (Roche) for 1 h at 45°C, and incubated again at 95°C for 10 min

to inactivate the Proteinase K. Finally, DNA was purified using the ChIP DNA Clean & Concentrator

kit (Zymo Research).

CLINIP-MS

CLNIP-MS was performed as previously described in (Fang *et al.* 2019). Briefly, 12-days-old *aprf1-9*

APRF1$_{pro}$:APRF1-VENUS seedlings were crosslinked with 1% formaldehyde. Tissue was then ground

to a fine powder and nuclei isolated in lysis buffer (20 mM Tris HCl pH 7.5, 20 mM KCl, 2 mM

EDTA pH 8, 2.5 mM MgCl$_2$, 25 % glycerol, 250 mM sucrose, 1x Roche protease inhibitor mixture),

filtered through miracloth, washed three times with NRBT (20 mM Tris HCl pH 7.5, 2.5 mM MgCl2,

25 % glycerol, 0.2 % Triton X-100, 1x Roche protease inhibitor mixture), and resuspended in RIPA

(50 mM Tris-HCl, 150 mM NaCl, 1 % Nonidet P-40, 0.5 % NaDeoxycholate, 0.1 % SDS, 1x Roche

protease inhibitor mixture) before sonication. Nuclear extract was incubated with GFP-Trap Magnetic

Agarose beads (ChromoTek, gtma-10) for 2 h and washed sequentially with Low Salt, High Salt, and

TE buffers as for ChIP. The immunoprecipitates were then boiled at 95°C for 15 min to reverse

crosslinking. The protein samples were gel-purified and subjected to mass spectrometry analysis by

nanoLC-MS/MS on a Orbitrap Fusion Tribrid Mass Spectrometer coupled to an UltiMate 3000

RSLCnano LC system Thermo Scientific, Hemel Hampstead, UK. Data were searched using Mascot

server (Matrixscience, London, UK) and analysed using MaxQuant software.

Preparation of Chromatin-bound RNA

Chromatin-bound RNA was isolated as previously described (Wu *et al.* 2016). Nuclei from 2-2.5 gr of

non-crosslinked seedlings were obtained with Honda Buffer, supplemented with 20 U/mL RNase

inhibitor RNase Out (Invitrogen), 1 mM PMSF, and 50 ng/µL of yeast tRNA. Nuclear pellet was

rinsed with 500 µL of resuspension buffer (20 mM Tris pH 8, 75 mM NaCl, 0.5 mM EDTA, 1 mM

481 DTT, 0.125 mM PMSF, 50 % glycerol, 1x Roche complete, 20 U/mL RNase Out) and centrifuged at

482 4000 xg at 4°C for 3 min. Nuclei pellet was weighed and resuspended in an equal volume of

483 Resuspension Buffer. The suspension was then washed with two volumes of Urea Wash Buffer (20

484 mM Tris pH8, 300 mM NaCl, 7.5 mM MgCl2, 0.25 mM EDTA, 1 mM DTT, 1 M Urea, 1 % NP-40,

485 1x Roche complete, 20 U/mL RNase Out), pipetting up and down 30 times, and spun at 8000 xg for 1

486 min at 4°C. Supernatant was kept as the Free RNA samples. Nuclei pellet was resuspended again with

487 1 volume of Resuspension buffer and washed with 1 volume of Urea Wash Buffer, pipetting 30 times

488 up and down, and spun for 1 min at 8000 xg and 4°C. Supernatant was combined with the previous

489 one to constitute the Free RNA sample. Finally, nuclear pellet was dissolved in 1 mL of TRIzol

490 (Invitrogen), adding 0.2 mL of chloroform and shaking vigorously by hand for 15 s. Then the

491 suspension was incubated at RT for 2 min and centrifuged at 12 000 xg for 15 min at 4°C. The

492 aqueous phase was taken and mixed with an equivalent volume of Phenol:Chloroform:Isoamyl

493 alcohol (25:24:1, Sigma), shaken for 10 min at room temperature, and centrifuge at RT and 12 000 xg

494 for 10 min. The aqueous phase was then transferred to another tube and followed two precipitations,

495 first with isopropanol, sodium acetate, and RNA GlycoBlue and another one with LiCl. Finally, RNA

496 was dissolved and treated with DNase Turbo (Ambion) and used as template for reverse transcription

497 with gene-specific primers.

499 **Quant-seq**

500 For Quant-seq experiments, total RNA was isolated as for qPCR and further cleaned up using the

501 Qiagen RNeasy miniprep kit (74106). Library preparation, sequencing, and data analysis were carried

502 out by Lexogen GmbH (Austria). Sequencing-ready libraries were generated from 100 ng of input

503 RNA using a QuantSeq 3' mRNA-Seq Library Prep Kit REV for Illumina (015UG009V0271)

504 following standard procedures. RNA integrity, and Indexed libraries quality were assessed on a

505 Fragment Analyzer device (Agilent Technologies) using a DNF-471 RNA Kit and HS- DNA assay,

506 respectively. Libraries were quantified using a Qubit dsDNA HS assay (Thermo Fisher). A

507 sequencing-ready pool of indexed libraries were sequenced on an Illumina NextSeq 2000 with a 100-

508 cycle cartridge using the Custom Sequencing Primer (CSP). FastQC version v0.11.7 was used to

509 verify the read quality and cutadapt version 1.18 (Martin 2011) for read adapter trimming. Clean reads

510 were mapped to the latest version of the Arabidopsis genome (TAIR10) with a splice-aware aligner

511 STAR version 2.6.1a (Dobin *et al.* 2013). Raw reads have been deposited on Short Read Archive

512 (SRA) under the reference PRJNA978558.

514 **AlphaFold2 protein interaction prediction**

515 AlphaFold models were predicted using the Colab notebook running a slightly simplified version of

516 AlphaFold v2.3.2. (Jumper *et al.* 2021). For the Arabidopsis LD–APRF1 complex LD aa566-661 and

517 APRF1 aa1-330 were used. For the human PNUTS–WDR82 complex PNUTS aa380-530 and

WDR82 aa1-313, and for the yeast Ref2–Swd2 complex Ref2 aa406-533 and Swd2 aa1-329 were used. The number of recycles were set to 3 and models included a final relaxation stage. PAE plots indicating the quality of the prediction and LDDT plots are shown in Figure S11. Final models were imported into and colour figures were prepared with PyMOL (v2.1, Schrödinger).

Bioinformatic analyses

To design the sgRNA to edit *APRF1* via CRISPR/Cas9 we used the CRISPR-P 2.0 (http://crispr.hzau.edu.cn/cgi-bin/CRISPR2/CRISPR), selecting the canonical NGG PAM motif, and the Arabidopsis TAIR10 as a target genome. Protein alignments were performed with MEGA X (Kumar *et al.* 2018) using the MUSCLE (Edgar 2004) algorithm with default parameters, and shaded with the "Colour Align Conservation" tool from the Sequence Manipulation Suite (Stothard 2000). To find putative PNUTS homologs in Arabidopsis, BLASTP (Altschul *et al.* 1997), PSI-BLAST, and DELTA-BLAST (Boratyn *et al.* 2012) searches were performed at NCBI site using an alignment score threshold of 80 for DELTA-BLAST, and default parameters for the rest. Individual protein structure predictions were retrieved from the Alphafold website (https://alphafold.ebi.ac.uk/).

ACKNOWLEDGEMENTS

For genetic materials, we are indebted to Fredy Barneche for providing seeds of *s2lb* and *aprf1-9 s2lb* (Fiorucci *et al.* 2019) and Soichi Inagaki for sharing their *3xFLAG-FLD* [in both (*fld-4; FRI*) and (*fld-4; fri*) backgrounds] transgenic lines (Inagaki *et al.* 2021). We are indebted to Yusheng Zhao for initiating the introgression of the GFP-LD line into Col*FRI* and to Minglei Yang for writing the code used to analyse Quant-seq data. Authors would also thank Shuqin Chen, Aida Sánchez, and Tina Zhang for their excellent technical assistance. This work was funded by the European Research Council Advanced Grant (EPISWITCH, 833254), Welcome Trust (210654/Z/18/Z), BBSRC Institute Strategic Programmes (BB/J004588/1 and BB/P013511/1), and the Royal Society Professorship (RP\R1\180002) to Caroline Dean; and the Medical Research Council (MRC) as part of United Kingdom Research and Innovation (MRC file reference number MC_U105192715) and the Wellcome Trust (225217/Z/22/Z) to Lori A Passmore. Eduardo Mateo-Bonmatí would like to thank grant RYC2021-030895-I, funded by MCIN/AEI and by European Union NextGenerationEU/PRTR.

FIGURE LEGENDS

Figure 1. APRF1, a homolog of the yeast Swd2 is a robust interactor of the FLD complex, and genetically functions in the FCA pathway. Tables in (A) and (B) summarize LC-MS/MS proteomic experiments using tagged versions of the FLD complex components FLD, LD, SDG26. Experiments from (A) were performed on formaldehyde-treated seedlings (Fang et al. 2020), and from (B) were carried out using non-cross-linked tissue (Qi et al. 2022). Different columns represent different replicates. (C) Architecture of the *APRF1* gene and illustration of the nature and location of the mutations studied in this work. Boxes indicate exons and lines introns. White boxes represent

untranslated regions. The triangle represents a T-DNA insertion (*aprf1-9*); the red arrow points to the location of the CRISPR/Cas9-derived deletion (*aprf1-10*). (D) Boxplot representing the leaf number at bolting of the wild-type Col-0 and both *aprf1* mutants. Each dot represents the score for one plant. (E) Relative values of *FLC* spliced (left) and unspliced (right) in the wild-type Col-0, both *aprf1* mutants and F1 *aprf1-9/aprf1-10* hybrid plants. Values were normalized to the housekeeping *UBC* gene and to Col-0. (F) Schematic diagram showing *FLC* gene structure following the guidelines described for (C). +1 indicates the transcriptional start site (TSS). (G-I) ChIP analysis of H3K4me1 (G), H3K36me3 (H), and H3K27me3 (I) levels at *FLC* in Col-0 and *aprf1-10*. Numbers in x axis represent the distance in kilobases to the *FLC* TSS and numbers in the y axis correspond to relative enrichment of the corresponding histone mark. Each dot represents an amplicon. Values were normalized to H3 and to *ACT* (for H3K4me1 and H3K36me3) or *STM* (for H3K27me3). (J-O) Relative values of *FLC* spliced (J, L, N), and unspliced (K, M, O) in various genetic backgrounds. Values were normalized to the housekeeping gene *UBC* (J-O) and Col-0 (J, K) or *PP2A* (L-O). Asterisks indicate statistically significant differences to Col-0 (D, E), and to C2 (*35S_{pro}:FCA; FRI*) (K), in a two-way Student's t test (** $p < 0.01$, *** $p < 0.001$, and **** $p < 0.0001$). The n.s. stands for not statistically different. Scale bars indicate 500 bp (C, F). Experiments were performed in 2-week-old seedlings grown in long days conditions with n ≥ 3 (E, G-O).

Figure 2. LD-APRF1-TOPP4 form a plant equivalent of the CPSF phosphatase module. (A) Summary of the CLNIP results using *aprf1-9 APRF1_{pro}:APRF1-VENUS* transgenic line. (B) Summary of the different proteomic experiments obtained in Qi et al (2022) using an APRF1-FLAG transgenic line in different backgrounds. (C) Components of the CP(S)F phosphatase domain in yeast, humans and Arabidopsis. (D) Schematic representation of protein size, annotated domains and disorder probability score for LD, PNUTS (*Homo sapiens*) and Ref2 (*Saccharomyces cerevisiae*). Individual amino acid score for potential disorder were obtained with the online Protein DisOrder prediction System (PrDOS) and plotted using Graphpad. All the proteins harbour a TFIIS-I domain (IPR003617), followed by highly disordered region. Additionally, LD carries an upstream homeodomain (IPR001356), PNUTS a Zinc-finger domain in the C-terminal region, and it was reported that the C-terminal region (348-523) of Ref2 is key to interact with Glc7 (Nedea et al., 2008). (E) AlphaFold2 predictions of complexes between APRF1-LD, WDR82-PNUTS, and Swd2-Ref2. WD40 orthologs are shown in grey surface representation, whereas TFIIS orthologs are shown in cartoon; the N-terminal amino acid of the predicted TFIIS proteins are denoted by a black sphere. (F) Overlay of the three predictions shown in (E).

Figure 3. ChIP-qPCR co-occupancy profiles indicate that FLD and LD work co-transcriptionally to control *FLC* transcription. (A) Elongating RNAPol II (Ser2P) ChIP profiles over *FLC* in a high (Col*FRI*) and low (*fri* or Col-0) transcriptional background. (B) ChIP binding profile over *FLC* of the *fld-4 FLD_{pro}:3xFLAG-FLD* transgenic plants in different genetic backgrounds. (C) ChIP binding profile over *FLC* of the *ld-1 LD_{pro}:GFP-LD* transgenic plants in different genetic backgrounds. *FLC* gene structure following the indications for Figure 1F. All the experiments were done with 2-week-old seedlings. Dots and error bars represent average ± s.e.m. of three replicates. Results in (A) are normalized as in Mikulski *et al*. 2022.

Figure 4. Mutations in *APRF1* trigger *COOLAIR* upregulation and an increase in a new medially polyadenylated *COOLAIR* isoform. (A) *FLC* architecture following the representation of Figure 1J with indication of different *COOLAIR* isoforms in grey. For simplicity, Classes I and II (proximal and distal) are each represented by one isoform. The new three *COOLAIR* class III isoforms are highlighted in pale blue. Triangles represent primers pairs (not drawn to scale) used to measure the relative abundance by qRT-PCR. Dashed lines indicate primers spanning two *COOLAIR* exons. Green triangles indicate the primer used for *COOLAIR* class III retro-transcription. Red vertical bars represent polyadenylation sites. Scale bar indicates 500 bp. (B-D) Relative expression analyses of (B) total *COOLAIR,* (C) proximal *COOLAIR*, and (D) distal *COOLAIR* in different genotypes. (E) *COOLAIR* proximal-to-distal ratio in different genotypes. (F, G) Relative expression analyses of *COOLAIR* class (F) III-1 and III-2, and (G) III-3. Each dot represents a biological replicate analysed in triplicate. Expression values were normalized to the *UBC* housekeeping gene and to Col-0.

Asterisks indicate statistically significant differences to the indicated genotypes in a Student's t test (*
$p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, and **** $p < 0.0001$). N.s. stands for not statistically different.
(c, f). All the experiments were performed in 2-week-old seedlings grown in long days conditions. (H)
Quant-seq experiment results from the *COOLAIR* strand carried out on Col*FRI* and *aprf1-9 FRI*
seedlings. *FLC* locus is represented as in panel A. The blue spikes indicate reads supporting a
polyadenylation site. Vertical pale orange bars highlight polyA sites found in *aprf1-9 FRI* absent in
the wild type Col*FRI*.
618
Figure 5. *COOLAIR* transcriptional readthrough correlates with changed phosphorylation of the RNA
Pol II carboxy terminal domain (CTD). (A) Chromatin-bound RNA levels of *FLC* and *COOLAIR* in
aprf1-9 mutants with and without functional *FRI*. Data was normalized to *UBC* and *PP2A* and is
shown as fold-change to wildtype Col-0. Dots correspond to amplicons of the *FLC* (upper chart) and
COOLAIR (bottom chart) transcripts, represented as average ± s.e.m. of three biological replicates. (B,
C) RNA Pol II ChIP of Col-0 vs *aprf1-10* (B), and Col*FRI* vs *aprf1-9 FRI* (C) using antibodies raised
against different phosphoisoforms (Ser2P and Ser5P) of the CTD. (D) RNA Pol II ChIP of Col-0, *ld-*
1, and *ld-1 aprf1-9* using antibodies recognizing total CTD (left), CTD Ser2P (centre), and CTD
Ser5P (right). Pol II ChIP data was normalized as in Fig 3A.
628
Figure 6. Proposed model for transcription-mediated chromatin silencing. (A) Open *FLC* chromatin
represented by white nucleosomes and marked with H3K4me1 (green circles) is actively transcribed
by RNA Pol II machinery (solid maroon), which carries a non-active FLD complex (blue circle) as
well as the 3'-end processing machinery (CPSF), including the phosphatase module formed by LD-
APRF1-TOPP4, and elongation factors (green circle). Pol II CTD and elongation factors harbour
some posttranslational modifications including phosphorylation (yellow circles). Nascent *COOLAIR*
forms an R-loop in the 3' end of the locus (Sun *et al.* 2013, Xu *et al.* 2021b). (B) Formation of the R-
loop stimulates the 3'-end processing machinery to terminate transcription at the proximal PAS. This
termination is also signalled by the phosphatase module to Pol II via dephosphorylation of either
elongating factors or the CTD or both (dashed line). PAS recognition triggers a conformational
change on RNA Pol II, illustrated by a solid to pale maroon colour change, which also activates FLD
(now solid blue circle). (C) After *COOLAIR* is released, RNA Pol II continues transcribing an
uncapped transcript that is the substrate of 5'-3' exonucleases (XRNs). During this non-productive
transcription, the FLD complex co-transcriptionally removes H3K4me1 marks from nucleosomes,
creating a less processive chromatin environment for subsequent rounds of transcription.
644

REFERENCES

Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J.
(1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search
programs Nucleic Acids Res *25*, 3389-3402.
Bae, H.J., Dubarry, M., Jeon, J., Soares, L.M., Dargemont, C., Kim, J., Geli, V., and Buratowski, S.
(2020). The Set1 N-terminal domain and Swd2 interact with RNA polymerase II CTD to
recruit COMPASS. Nat Commun *11*, 2181.
Berry, S., Hartley, M., Olsson, T.S., Dean, C., and Howard, M. (2015). Local chromatin environment
of a Polycomb target gene instructs its own epigenetic inheritance. eLife *4*, e07205.
Boratyn, G.M., Schäffer, A.A., Agarwala, R., Altschul, S.F., Lipman, D.J., and Madden, T.L. (2012).
Domain enhanced lookup time accelerated BLAST. Biology Direct *7*, 12.
Carminati, M., Manav, M.C., Bellini, D., and Passmore, L.A. (2022). A direct interaction between
CPF and Pol II links RNA 3□-end processing to transcription.
https://doi.org/10.1101/2022.07.28.501803
Casanal, A., Kumar, A., Hill, C.H., Easter, A.D., Emsley, P., Degliesposti, G., Gordiyenko, Y.,
Santhanam, B., Wolf, J., Wiederhold, K., et al. (2017). Architecture of eukaryotic mRNA 3'-
end processing machinery. Science *358*, 1056-1059.
Castaings, L., Bergonzi, S., Albani, M.C., Kemi, U., Savolainen, O., and Coupland, G. (2014).
Evolutionary conservation of cold-induced antisense RNAs of *FLOWERING LOCUS C* in
Arabidopsis thaliana perennial relatives. Nat Commun *5*, 4457.
Cortazar, M.A., Sheridan, R.M., Erickson, B., Fong, N., Glover-Cutter, K., Brannan, K., and Bentley,
D.L. (2019). Control of RNA Pol II speed by PNUTS-PP1 and Spt5 dephosphorylation
facilitates termination by a "Sitting Duck Torpedo" mechanism. Mol Cell *76*, 896-908.
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and
Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.
Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res *32*, 1792-1797.
Estell, C., Davidson, L., Eaton, J.D., Kimura, H., Gold, V.A.M., and West, S. (2023). A restrictor
complex of ZC3H4, WDR82, and ARS2 integrates with PNUTS to control unproductive
transcription. Mol Cell.
Fang, X., Wang, L., Ishikawa, R., Li, Y., Fiedler, M., Liu, F., Calder, G., Rowan, B., Weigel, D., Li,
P., and Dean, C. (2019). Arabidopsis FLL2 promotes liquid-liquid phase separation of
polyadenylation complexes. Nature *569*, 265-269.
Fang, X., Wu, Z., Raitskin, O., Webb, K., Voigt, P., Lu, T., Howard, M., and Dean, C. (2020). The 3'
processing of antisense RNAs physically links to chromatin-based transcriptional control.
Proc Natl Acad Sci USA *117*, 15316-15321.
Fiorucci, A.S., Bourbousse, C., Concia, L., Rougee, M., Deton-Cabanillas, A.F., Zabulon, G., Layat,
E., Latrasse, D., Kim, S.K., Chaumont, N., et al. (2019). Arabidopsis S2Lb links
AtCOMPASS-like and SDG2 activity in H3K4me3 independently from histone H2B
monoubiquitination. Genome Biol *20*, 100.
Gregersen, L.H., Mitter, R., Ugalde, A.P., Nojima, T., Proudfoot, N.J., Agami, R., Stewart, A., and
Svejstrup, J.Q. (2019). SCAF4 and SCAF8, mRNA anti-terminator proteins. Cell *177*, 1797-
1813.
Grewal, S.I.S. (2023). The molecular basis of heterochromatin assembly and epigenetic inheritance.
Mol Cell *83*, 1767-1785.
Gyenis, A., Chang, J., Demmers, J., Bruens, S.T., Barnhoorn, S., Brandt, R.M.C., Baar, M.P., Raseta,
M., Derks, K.W.J., Hoeijmakers, J.H.J., and Pothof, J. (2023). Genome-wide RNA
polymerase stalling shapes the transcriptome during aging. Nat Genet *55*, 268-279.
Hu, S., Peng, L., Song, A., Ji, Y.X., Cheng, J., Wang, M., and Chen, F.X. (2023). INTAC
endonuclease and phosphatase modules differentially regulate transcription by RNA
polymerase II. Mol Cell *83*, 1588-1604.
Inagaki, S., Takahashi, M., Takashima, K., Oya, S., and Kakutani, T. (2021). Chromatin-based
mechanisms to coordinate convergent overlapping transcription. Nat Plants *7*, 295-302.

Jiang, D., Kong, N.C., Gu, X., Li, Z., and He, Y. (2011). Arabidopsis COMPASS-like complexes
mediate histone H3 lysine-4 trimethylation to control floral transition and plant development.
PLoS Genet *7*, e1001330.
Johanson, U., West, J., Lister, C., Michaels, S., Amasino, R., and Dean, C. (2000). Molecular analysis
of *FRIGIDA*, a major determinant of natural variation in *Arabidopsis* flowering time. Science
290, 344-347.
Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction
with AlphaFold. Nature *596*, 583-589.
Kapolas, G., Beris, D., Katsareli, E., Livanos, P., Zografidis, A., Roussis, A., Milioni, D., and
Haralampidis, K. (2016). *APRF1* promotes flowering under long days in *Arabidopsis*
thaliana. Plant Sci *253*, 141-153.
Kim, H.J., Li, P., Kim, T., Oldfield, A.J., Zheng, X., and Yang, P. (2022). Integrative analysis reveals
histone demethylase LSD1 promotes RNA polymerase II pausing. iScience *25*, 105049.
Kowalik, K.M., Shimada, Y., Flury, V., Stadler, M.B., Batki, J., and Bühler, M. (2015). The Paf1
complex represses small-RNA-mediated epigenetic gene silencing. Nature *520*, 248-252.
Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: molecular evolutionary
genetics analysis across computing platforms. Molecular Biology and Evolution *35*, 1547-
1549.
Li, Z., Jiang, D., and He, Y. (2018). FRIGIDA establishes a local chromosomal environment for
FLOWERING LOCUS C mRNA production. Nat Plants *4*, 836-846.
Liu, F., Marquardt, S., Lister, C., Swiezewski, S., and Dean, C. (2010). Targeted 3' processing of
antisense transcripts triggers *Arabidopsis FLC* chromatin silencing. Science *327*, 94-97.
Liu, F., Quesada, V., Crevillén, P., Bäurle, I., Swiezewski, S., and Dean, C. (2007). The Arabidopsis
RNA-binding protein FCA requires a lysine-specific demethylase 1 homolog to downregulate
FLC. Mol Cell *28*, 398-407.
Liu, Y., and Huang, Y. (2018). Uncovering the mechanistic basis for specific recognition of
monomethylated H3K4 by the CW domain of *Arabidopsis* histone methyltransferase SDG8. J
Biol Chem *293*, 6470-6481.
Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal *17*, 10-12.
Menon, G., Mateo-Bonmatí, E., Reeck, S., Maple, R., Wu, Z., Ietswaart, R., Dean, C., and Howard,
M. (2023). Alternative polyadenylation controls a transcriptional state that determines the rate
of Polycomb silencing *Co-submitted with this manuscript*.
Menon, G., Schulten, A., Dean, C., and Howard, M. (2021). Digital paradigm for Polycomb
epigenetic switching and memory. Curr Opin Plant Biol *61*, 102012.
Mikulski, P., Wolff, P., Lu, T., Nielsen, M., Echevarria, E.F., Zhu, D., Qüesta, J.I., Saalbach, G.,
Martins, C., and Dean, C. (2022). VAL1 acts as an assembly platform co-ordinating co-
transcriptional repression and chromatin regulation at Arabidopsis *FLC*. Nat Commun *13*,
5542.
Mo, W., Liu, B., Zhang, H., Jin, X., Lu, D., Yu, Y., Liu, Y., Jia, J., Long, Y., Deng, X., et al. (2021).
Landscape of transcription termination in Arabidopsis revealed by single-molecule nascent
RNA sequencing. Genome Biol *22*, 322.
Muniz, L., Nicolas, E., and Trouche, D. (2021). RNA polymerase II speed: a key player in controlling
and adapting transcriptome composition. EMBO J *40*, e105740.
Nedea, E., Nalbant, D., Xia, D., Theoharis, N.T., Suter, B., Richardson, C.J., Tatchell, K., Kislinger,
T., Greenblatt, J.F., and Nagy, P.L. (2008). The Glc7 phosphatase subunit of the cleavage and
polyadenylation factor is essential for transcription termination on snoRNA genes. Mol Cell
29, 577-587.
Pinter, S., Knodel, F., Choudalakis, M., Schnee, P., Kroll, C., Fuchs, M., Broehm, A., Weirich, S.,
Roth, M., Eisler, S.A., et al. (2021). A functional LSD1 coregulator screen reveals a novel
transcriptional regulatory cascade connecting R-loop homeostasis with epigenetic regulation.
Nucleic Acids Res *49*, 4350-4370.

Qi, P.L., Zhou, H.R., Zhao, Q.Q., Feng, C., Ning, Y.Q., Su, Y.N., Cai, X.W., Yuan, D.Y., Zhang,
Z.C., Su, X.M., et al. (2022). Characterization of an autonomous pathway complex that
promotes flowering in *Arabidopsis*. Nucleic Acids Res *50*, 7380-7395.
Rodríguez-Molina, J.B., West, S., and Passmore, L.A. (2023). Knowing when to stop: Transcription
termination on protein-coding genes by eukaryotic RNAPII. Mol Cell *83*, 404-415.
Schon, M., Baxter, C., Xu, C., Enugutti, B., Nodine, M.D., and Dean, C. (2021). Antagonistic
activities of cotranscriptional regulators within an early developmental window set *FLC*
expression level. Proc Natl Acad Sci USA *118*, e2102753118.
Simpson, G.G., Dijkwel, P.P., Quesada, V., Henderson, I., and Dean, C. (2003). FY is an RNA 3' end-
processing factor that interacts with FCA to control the *Arabidopsis* floral transition. Cell *113*,
777-787.
Soares, L.M., and Buratowski, S. (2012). Yeast Swd2 is essential because of antagonism between
Set1 histone methyltransferase complex and APT (associated with Pta1) termination factor. J
Biol Chem *287*, 15219-15231.
Sonmez, C., Bäurle, I., Magusin, A., Dreos, R., Laubinger, S., Weigel, D., and Dean, C. (2011). RNA
3' processing functions of *Arabidopsis* FCA and FPA limit intergenic transcription. Proc Natl
Acad Sci USA *108*, 8508-8513.
Stothard, P. (2000). The Sequence Manipulation Suite: JavaScript programs for analyzing and
formatting protein and DNA sequences. Biotechniques *28*, 1102-1104.
Sun, Q., Csorba, T., Skourti-Stathaki, K., Proudfoot, N.J., and Dean, C. (2013). R-loop stabilization
represses antisense transcription at the Arabidopsis *FLC* locus. Science *340*, 619-621.
Tsutsui, H., and Higashiyama, T. (2017). pKAMA-ITACHI Vectors for Highly Efficient
CRISPR/Cas9-Mediated Gene Knockout in *Arabidopsis thaliana*. Plant Cell Physiol *58*, 46-
56.
Wagner, E.J., Tong, L., and Adelman, K. (2023). Integrator is a global promoter-proximal termination
complex. Mol Cell *83*, 416-427.
Wu, Z., Fang, X., Zhu, D., and Dean, C. (2020). Autonomous pathway: *FLOWERING LOCUS C*
repression through an antisense-mediated chromatin-silencing mechanism. Plant Physiol *182*,
27-37.
Wu, Z., Ietswaart, R., Liu, F., Yang, H., Howard, M., and Dean, C. (2016). Quantitative regulation of
FLC via coordinated transcriptional initiation and elongation. Proc Natl Acad Sci USA *113*,
218-223.
Xu, C., Fang, X., Lu, T., and Dean, C. (2021a). Antagonistic cotranscriptional regulation through
ARGONAUTE1 and the THO/TREX complex orchestrates *FLC* transcriptional output. Proc
Natl Acad Sci USA *118*.
Xu, C., Wu, Z., Duan, H.C., Fang, X., Jia, G., and Dean, C. (2021b). R-loop resolution promotes co-
transcriptional chromatin silencing. Nat Commun *12*, 1790.
Yang, H., Howard, M., and Dean, C. (2014). Antagonistic roles for H3K36me3 and H3K27me3 in the
cold-induced epigenetic switch at *Arabidopsis FLC*. Curr Biol *24*, 1793-1797.
Zhu, P., Lister, C., and Dean, C. (2021). Cold-induced *Arabidopsis* FRIGIDA nuclear condensates for
FLC repression. Nature *599*, 657-661.
Zumer, K., Maier, K.C., Farnung, L., Jaeger, M.G., Rus, P., Winter, G., and Cramer, P. (2021). Two
distinct mechanisms of RNA polymerase II elongation stimulation *in vivo*. Mol Cell *81*, 3096-
3109.
795

A

Number of peptides

	APRF1-VENUS
	aprf1-9
FLD	0
LD	24
SDG26	0
APRF1	86
TOPP4	18

B

Number of peptides

	APRF1-FLAG		
	APRF1	*fld*	*ld*
FLD	15	0	0
LD	27	6	0
SDG26	8	4	0
APRF1	16	28	26
TOPP4	14	11	14

C

Glc7 / Swd2 / Ref2 — Yeast

PP1 / WDR82 / PNUTS — Human

TOPP4 / APRF1 / LD — Arabidopsis

D

LD

PNUTS

Ref2

E

APRF1-LD

180°

WDR82-PNUTS

180°

Swd2-Ref2

180°

F