Prospects for a sequence-based taxonomy of influenza A virus subtypes

Art F. Y. Poon[1,2,3]

[1]Department of Pathology & Laboratory Medicine, Western University, London, Canada

[2]Department of Microbiology & Immunology, Western University, London, Canada

[3]Department of Computer Science, Western University, London, Canada

Abstract

The hemagglutinin (HA) and neuraminidase (NA) proteins are the primary antigenic targets of influenza A virus (IAV) infections. IAV infections are generally classified into subtypes of HA and NA proteins, *e.g.*, H3N2. Most of the known subtypes were originally defined by a lack of antibody cross-reactivity. However, genetic sequencing has played an increasingly important role in characterizing the evolving diversity of IAV. Novel subtypes have recently been described solely by their genetic sequences, and IAV infections are routinely subtyped by molecular assays, *e.g.*, real-time PCR, or the comparison of sequences to references. In this study, I carry out a phylogenetic analysis of all available IAV protein sequences in the Genbank database (over 1.1 million records) to determine whether the serologically-defined subtypes can be reproduced with sequence-based criteria. I show that a robust genetic taxonomy of HA and NA subtypes can be obtained with a simple clustering method, namely by progressively partitioning the phylogeny on its longest internal branches. However, this taxonomy also requires some amendments to the current nomenclature. For example, two IAV isolates from bats previously characterized as a divergent lineage of H9N2 should be separated into their own subtype. With the exception of these small and highly divergent lineages, the phylogenies relating each of the other six genomic segments do not support partitions into major subtypes.

Introduction

Influenza A virus (IAV) genomes are comprised of eight single-stranded, negative sense RNA segments, which are numbered in decreasing order with respect to their lengths. The diversity of

IAV is usually characterized by the two segments (numbered 4 and 6) carrying the genes encoding the hemagglutinin (HA) and neuraminidase (NA) proteins, respectively. HA and NA are surface-exposed proteins responsible for binding and release of virus particles from host cells, respectively. As significant antigenic targets of the adaptive immune system, they are the most rapidly evolving proteins of the IAV genome [3]. The global diversities of both HA and NA sequences are clustered into serotypes or subtypes, terms which are used interchangeably in the literature; herein I will use the term 'subtypes' for consistency. These subtypes are identified by H and N prefixes for the respective proteins, followed by an integer suffix, producing the widely familiar HnNn nomenclature, *e.g.*, H3N2. Certain combinations of HA and NA subtypes are observed more frequently than others due to pandemic growth. Although these combinations are also referred to as subtypes, each genome segment has its own subtype nomenclature.

There are presently 18 described HA subtypes that are numbered H1 to H18 (Table 1). In addition, a potential new subtype H19 has only recently been described [7] from a partial HA sequence isolated in 2008 from waterfowl in Kazakhstan. In addition, there are 11 described subtypes of NA (N1 to N11). These subtypes were originally defined solely from their antigenic characteristics, predominantly from hemagglutination inhibition (HI, specifically for HA), complement fixation (CF), or double immunodefusion (DID [5]) testing. For instance, H2, initially labelled the 'Far East' or 'Asian' strain of IAV [6], was distinguished from H1 by the lack of HI activity by sera from hosts infected by the other subtype [14]. The nomenclature for HA and NA subtypes also formerly incorporated the host organism [34]. For instance, H1, Hsw1, Heq1 and Hav1 were originally different HA subtypes isolated from human, swine, equine and avian hosts, respectively [34]. These were subsequently merged by Schild et al. [26] into the current host-agnostic Hn and Nn nomenclature on the basis of the cross-reactivity of isolates from different host species.

Since the description of subtype H13 by Hinshaw et al. [13] in 1983, genetic sequence analysis has been increasingly incorporated into the characterization of novel HA and NA subtypes (Table 1). Indeed, the last three HA subtypes (H17-H19) have been described solely by their genetic

Subtype	Previous designation	Prototype strain	Methods	Publication
H1	H0	A/Puerto Rico/8/1934	HI, DID	[26]
	H1	A/Fort Monmouth/1/1947		
	Hsv1	A/Swine/Wisconsin/15/30		
H2	H2	A/Singapore/1/57	HI	[14]
H3	H3	A/Hong Kong/1/68	HI, DID	[26]
	Heq2	A/eq/Miami/1/63		
	Hav7	A/Duck/Ukraine/1/63		
H4	Hav4	A/Duck/Czech/56	HI, CF, DID	[21]
H5	Hav5	A/Tern/South Africa/61	HI, CF	[21]
H6	Hav6	A/Turkey/Mass/3740/65	HI, CF	[21]
H7	Heq1	A/Equine/Prague/1/56	HI, DID	[26]
	Hav1	A/FPV/Dutch/27		
H8	Hav8	A/Turkey/Ontario/6118/68	HI, CF, DID*	[18]
H9	Hav9	A/Turkey/Wisconsin/66	HI, DID	[31]
H10	Hav2	A/Chick/Germ/N/49	HI, CF	[21]
H11	Hav3	A/Duck/England/56	HI, CF	[21]
H12	Hav10	A/Duck/Alberta/60/76	HI, DID	[12]
H13		A/Gull/Md/704/77	HI, DID, AAI	[13]
H14		A/Mallard/Gurjev/263/82	HI, DID, P, AAI	[17]
H15		A/duck/Australia/341/83	HI, DID, P, AAI	[24]
H16		A/black-headed gull/Sweden/5/99	HI, DID, P, AAI	[8]
H17		A/little yellow-shouldered bat/Guatemala/153/2009	P, AAI	[29]
H18		A/flat-faced bat/Peru/033/2010	P, AAI	[30]
H19†		A/Common Pochard/Kazakhstan/KZ-52/2008	P, AAI	[7]

Table 1: Summary of the currently known HA subtypes and the methods used to describe them. Previous subtype and prototype strain designations for subtypes H1 to H12 were obtained from Schild et al. [26]. Methods: HI = hemagglutination inhibition; CF = complement-fixation; DID = double immunodiffusion; P = phylogeny; AAI = amino acid sequence identity. *The authors use an immunodiffusion assay that resembles DID (described in Beard [2]). †H19 has only recently been proposed from a single partial HA sequence.

sequences [7, 29, 30]. High-throughput genetic sequencing has become a ubiquitous feature of clinical and public health laboratories, boosted recently by the expansion of resources to confront the global SARS-CoV-2 pandemic. For instance, the number of IAV sequences deposited in the GISAID (Global Initiative for Sharing All Influenza Data, https://gisaid.org) database has grown exponentially from 943 sequences in 1992 to 307,198 in 2022 (Supplementary Figure S1).

Here, I investigate whether it is feasible to distinguish HA and NA subtypes based only on genetic diversity. This is a different problem than the supervised classification of genetic sequences to subtypes based on their similarity to a predefined set of reference sequences, commonly performed by BLAST search [27]. Instead, I ask whether an unsupervised method can robustly cluster HA and NA sequences into the antigenically-defined subtypes. Furthermore, I assess whether it is feasible to apply a similar classification scheme to the other six genomic segments. There is surpisingly limited work in this area. When describing serotype H14, for instance, Kawaoka et al. [17] proposed that HA sequences that differed at 30% of amino acids or more should be separated into different subtypes. However, this proportional (p) distance criterion is not attained for all pairwise comparisons of subtypes [24, 29]. This method also makes no adjustment for multiple substitutions at the same sites over long evolutionary time scales. Furthermore, the shortest p-distance between groups can depend by chance on which sequences have been incorporated into the analysis. Ideally the phylogenetic analysis should incorporate as many genetic sequences as possible.

Methods

I downloaded all available protein sequences associated with all eight IAV segments from the NCBI Genbank database (accessed June 6, 2023 except for HA, which were retrieved April 28, 2023), using a minimum length filter to exclude records with incomplete nucleotide sequences (Supplementary Table S1). For each segment, I used regular expressions to filter sequences from other segments based on gene and protein annotations in the sequence names. In addition, I excluded sequences derived from alternate open reading frames (PB1-F2 and PA-X), and concate-

nated sequences involved in alternative splicing with overlapping regions excluded (M1/M2 and NS1/NS2). Next, I discarded sequences with more than 10% ambiguous residues ('X') and compressed the remainder into unqiue sequences, recording the labels of duplicate sequences into a separate file. The remaining sequences were aligned using MAFFT (version 7.453) [16]. I manually evaluated and amended the resulting alignments with AliView (v.2018) [19]. Using FastTree, I reconstructed a preliminary tree from the alignment without maximum-likelihood or minimum-evolution steps (*i.e.*, neighbor-joining), and then visually assessed the tree using Taxonium [25]. After excluding any problematic sequences identified in the preliminary tree (*e.g.*, recombinant sequences produced by molecular cloning [11]), I re-ran FastTree with the default optimization steps to generate a maximum likelihood tree.

I investigated two different methods to partition each tree into monophyletic clades (*i.e.*, subtrees) as putative subtypes. First, I calculated the following quantities for each internal node in the tree:

1. y_i, the mean tip-to-tip (patristic) distance between every tip descending from node i;

2. d_i, the mean distance from node i to every descendant tip; and

3. v_i, the total distance from node i to its sibling node j.

The quantities d_i and y_i were calculated by postorder traversal of the tree for increased efficiency. For instance, y_i was calculated by this recurrence relation, where a and b are child nodes of i:

$$y_i = \frac{\binom{n_a}{2}y_a + \binom{n_b}{2}y_b + n_a n_b (d_a + d_b + l_a + l_b)}{\binom{n_a}{2} + \binom{n_b}{2} + n_a n_b}$$

where n_a is the number of tips descending from node a, and l_a is the length of the branch associated with (below) node a. Tip nodes were initialized with $n_i = 1$, $d_i = 0$, and $y_i = 0$. Hence, the y_i for the internal node of a cherry reduces to $l_a + l_b$. The quantity y_i is analogous to the mean pairwise distance within a subtype, while v_i is analogous to the shortest distance between two subtypes. Given minimum v and maximum y cutoffs, subtrees meeting these criteria were located

by evaluating nodes by preorder traversal of the tree. This method will be referred to as 'nodewise' clustering.

Second, I progressively cut the tree into subtrees on internal branches (edges) with a length exceeding a given cutoff, using the following algorithm:

1. initialize subtree list S with input tree

2. for each subtree s in S

   - locate longest internal branch b_s in s

   - if length of b_s exceeds cutoff

     - remove b_s to yield subtrees s_1 and s_2

     - for each new subtree, remove the node associated with b_s and join the branches

     - append s_1 and s_2 to new list S'

   - otherwise append s to new list S'

3. replace S with S'

4. repeat from step 2 until no internal branches exceed cutoff

This method will be referred to as 'edgewise' clustering. Both algorithms were implemented with the Phylo toolkit in Biopython [28].

To measure the correlation between the subtree clusters and the subtype annotations of sequences, I calculated the normalized mutual information (nMI) of these two partitions:

$$\text{nMI} = \frac{2\sum_i \sum_j p_{ij} \left(\log p_{ij} - \log p_i - \log p_j \right)}{-\sum_i p_i \log p_i - \sum_j p_j \log p_j}$$

where p_i is the marginal proportion of labeled tips in subtree i, p_j is the marginal proportion of tips labeled with subtype j, and p_{ij} is the number of tips in subtree i with label j divided by the total number of labels, *i.e.*, the joint proportion (see also Equation 14.4.17 in Press et al. [22]). The nMI adjusts for different numbers of subsets between these two partitions, and it ranges from 0 (no correlation) to 1 (for identical partitions).

Results

Maximum likelihood trees for $n = 66,864$ HA and $n = 52,146$ NA amino acid sequences are displayed in Figure 1 with branches coloured by subtype annotations, *i.e.*, NCBI Genbank source qualifiers. As anticipated, the subtrees labelled by subtype annotations are distinctly separated by long internal branches in both trees, implying that these subtypes should be readily distingushed by basic sequence-based criteria. The overall scale of the NA tree is substantially greater than the HA, driven by the enormous divergence of N10 and N11 from the other subtypes (4.23 expected substitutions per site). Based on their discordant locations in the tree, I identified 15 HA and 30 NA sequences with incorrect subtype annotations (Supplementary Tables S2 and S3). In addition, I classified 2,353 HA and 2,563 NA sequences with missing subtype annotations (*e.g.*, 'mixed', 'HX' or 'unknown') by locating their nearest neighbours in the respective phylogenies. These sequences were substantially less likely to assigned to the most common subtypes (Supplementary Figure S2). These inferred subtype labels were excluded from further analyses to mitigate potential biases.

I evaluated two different methods to partition a phylogeny into subtrees that are consistent with the antigenically-defined IAV hemagglutinin (HA) and neuraminidase (NA) subtypes. First, I assessed a nodewise method in which I calculated the mean tip-to-tip distance y and total branch length to the sibling node v for subtrees descending from every internal node. These quantites are analogous to the within- and between-group measures of genetic divergence that have been used to define clusters within IAV subtypes, *i.e.*, clades [1, 32]. Both v and y were required to partition the tree *de novo* into subtrees. For instance, using only a minimum cutoff on v tends to prematurely terminate a search by preorder traversal of nodes, *i.e.*, yielding two large subtrees that descend directly from the root node.

I evaluated the discordance between subtypes extracted from the sequence metadata (herein referred to as 'labels') and the phylogenetic subtrees produced at varying cutoffs of y and v (Supplementary Figure S3A-C). Overall, the discordance in the number of subtrees was minimized at a

Figure 1: Maximum likelihood trees reconstructed from unique amino acid sequences of HA ($n = 66,864$) and NA ($n = 52,146$). The trees are midpoint-rooted and ladderized. Branches are coloured by respective 'serotype' annotations from the source qualifiers in the NCBI Genbank database. Scale bars indicate the branch length corresponding to 0.1 expected amino acid substituitons per site. These plots were generated using the R package *ggfree* (https://github.com/ArtPoon/ggfree).

tip-to-tip distance cutoff of about $y = 1.2$. The results were less sensitive to varying cutoffs for v. However, the correspondence between subtrees and labels at the best settings was generally poor. Sequences labeled as subtype H9 were consistently partitioned into multiple subtrees. In addition, several groups of subtype labels become merged into single subtrees, namely H2/H5, H3/H4/H14, H7/H10, H8/H12, and H11/H13/H16 (Supplementary Figure S3D). These discrepancies could not resolved by varying the clustering settings for this method.

Since the results of nodewise clustering on the HA phylogeny were so poor, I abandoned this method and implemented a simpler edgewise method. This method partitions the phylogeny by progressively cutting internal branches with lengths exceeding some cutoff. The number of subtrees that result from applying this method to the HA phylogeny is summarized in Figure 2A. The target number of 18 subtrees is attained only within a narrow range of cutoffs around 0.18. Overall, the distribution of labels among the subtrees at this cutoff were generally concordant (Figure 2B),

Figure 2: (A) A line plot represents the number of subtrees produced by partitioning the phylogeny of HA sequences at varying branch length cutoffs (edgewise clustering). A dashed horizontal line is drawn at $n = 18$ to indicate the target number. The shaded region summarizes the normalized mutual information indicating the concordance between subtrees and HA subtype labels. (B) Distribution of HA subtype labels among subtrees produced at an internal branch length cutoff of 0.18. Boxes are shaded in proportion to relative counts of labels. Total counts are listed along the right side. Individual counts are printed in the respective boxes for low-frequency labels, most of which correspond to misclassified sequences (see Table S2).

with the exception of two major discrepancies. First, one subtree (indexed 8) was comprised of only two sequences corresponding to IAV isolated from bats resembling H9 [15, 23], which predominantly circulates in avian host species. This subtree does not merge with the adjacent subtree carrying the other 8,197 H9 labels until the cutoff is relaxed to 0.27, at which there is a sudden drop in the number of subtrees from 17 to 14 (Figure 2A). Second, the nine HA sequences labeled as H15 are grouped with H7 at this cutoff. They are split when the cutoff is decreased from 0.18 to 0.1775. If the cutoff is increased from 0.18 to 0.1825, then subtrees labeled H13 and H16 become merged into a single subtree.

Next, I applied the edgewise method to a maximum-likelihood phylogeny of $52,146$ NA sequences. This method yielded 12 subtrees for a broad range of cutoffs (from 0.222 to 0.4 expected substitutions, Figure 3A). Similarly to the case with the HA phylogeny, the target number of 11 subtrees could only be obtained within a narrow range of cutoffs (from 0.405 to 0.425). At a cutoff of 0.41, I observed the same two major discrepancies among the resulting 11 subtrees (Figure 3B). Two sequences labeled as N2 were separated from the subtree carrying the bulk of the reamining the N2 labels ($n = 23,803$). Not surprisingly, these corresponded to the same divergent N9N2-like viruses isolated from bats as noted above [15, 23]. In addition, 394 sequences labeled as N5 are placed in the same subtree as N8 unless the cutoff is reduced below 0.405.

Finally, I repeated this same process for the remaining IAV genome segments. Unlike the HA and NA phylogenies, there was limited support for partitioning these other phylogenies into distinct subtrees. Using the results in Figures 2A and 3A as a guide, I selected cutoffs at which the number of subtrees was relatively stable for each of these other segments (Supplementary Figure S4). These cutoffs tended to be around 0.05, yielding four to six subtrees (Supplementary Table S4), with the exception of the NS1/NS2 phylogeny where five subtrees were obtained at cutoffs ranging from 0.12 to 0.175. This difference in scale is consistent with previous findings that diversifying selection at the amino acid level for NS1 is comparable to the surface-exposed HA and NA proteins [4]. In every case, nearly all sequences were assigned to one of the five

Figure 3: (A) A line plot represents the number of subtrees produced by partitioning the phylogeny of NA sequences at varying branch length cutoffs (edgewise clustering). A dashed horizontal line is drawn at $n = 11$ to indicate the target number. The shaded region summarizes the normalized mutual information indicating the concordance between subtrees and NA subtype labels. (B) Distribution of NA subtype labels among subtrees produced at an internal branch length cutoff of 0.41. Boxes are shaded in proportion to relative counts of labels. Total counts are listed along the right side. Individual counts are printed in the respective boxes for minority serotypes. The smaller counts generally correspond to misclassified sequences (see Table S3).

subtrees. Two of the remaining small subtrees corresponded to the recently described H17N10 [29] and H18N11 [30] strains, respectively. Additionally, the divergent lineage of H9N2 that was consistently separated into its own HA and NA subtrees was also separated for all other segments except PB1. For all phylogenies except NP, one of the small subtrees corresponded to isolates of equine influenza (H7N7) from 1956. Finally, the phylogeny relating concatenated M1 and M2 protein sequences contained an additional subtree comprising three reassortant sequences isolated from swine in Australia [33].

Discussion

These results indicate that it is feasible to define subtypes on the basis of the variation among HA and NA protein sequences. Subtypes derived from phylogenies are largely consistent with the current definitions of 18 HA and 11 NA subtypes, which are mostly derived from serological data. However, there are some important discrepancies. The analysis presented here supports 17 genetically-defined HA subtypes, not 18. Two subtypes are removed by merging H15 with H7, and combining H13 and H16. This is consistent with previous work reporting average amino acid identities of 79.7% and 81.4% between these respective groups, where the typical mean identity was about 49% [29]. In addition, it supports the creation of a new subtype from a divergent lineage of H9N2-like isolates from bats. This lineage not only also creates an additional twelfth subtype for NA, but also forms distinct subtrees for nearly every other segment. The two sequences comprising this lineage (Genbank accessions MH376902 and OQ216561) were isolated from bats in Egypt in 2017 [15] and South Africa in 2018 [23], respectively. Kandeil et al. [15] reported the HA sequence was the most similar (73% nucleotide identity) to H9 sequences that circulate predominantly in avian host species. Even though this divergence was comparable to that between other HA subtypes, neither study made a case for defining a new subtype.

This is not the first attempt to cluster the genomic segments of IAV into subtypes. For instance, Lu et al. [20] performed a similar clustering analysis of roughly all segments for 2,300 complete

IAV genomes. For each segment, they used neighbor-joining to reconstruct trees from the HKY85 distance matrix for the multiple sequence alignment, and then extracted clusters based on a bootstrap support threshold of >90% and a nucleotide p-distance threshold of about 10%. Their analysis partitioned the HA sequence phylogeny into 78 subtrees, for example. Unfortunately, the web interface for classifying IAV sequences under this system (https://www.flugenome.org) was discontinued in 2016, and the expired domain name is presently being used by a laboratory reagents supplier under the guise of the original website (last accessed June 30, 2023). In addition, the classifier developed by Lu et al. [20] was not made publicly available, and it was not described with sufficient detail to reproduce.

Similarly, Zhuang et al. [35] used neighbor joining to reconstruct phylogenies relating between 4,194 to 33,066 sequences for each of the eight IAV segments. They partitioned each tree into clusters based on the presence of a "relatively separated branch", as well as the distribution of host species, sampling location and collection dates among tips. For example, they described three major clusters in the PB2 phylogeny. Unfortunately, their clustering method was not described with sufficient quantitative detail to reproduce, and the study data (*e.g.*, annotated trees) are not available online. Finally, there are multiple studies that have clustered sequences for all IAV genome segments for the purpose of detecting reassortment events. For example, Gong et al. [9] partitioned the maximum likelihood trees for 40,296 full IAV genomes into clusters based on the mean tip-to-tip distances associated with internal nodes, *i.e.*, the y statistic in this study. Using the overall mean of y as a cutoff for each segment, they obtained 493 to 663 clusters per phylogeny. Although this fine-grained clustering may confer greater sensitivity for detecting reassortment, it does not provide insight into the genetic characteristics of established IAV subtypes.

The process of updating the nomenclature of IAV subtypes has been complicated by the transition from serological testing to genetic sequencing to characterize novel variants (Table 1). Given the serological origins of most IAV subtype definitions, it is remarkable that these groupings can be readily recovered from genetic variation alone, although not all methods are equally success-

ful. There are significant advantages of developing a genetic taxonomy for IAV subtypes. The number of published IAV sequences is growing exponentially (Supplementary Figure S1). IAV infections are increasingly subtyped by genetic sequencing methods that are more scaleable than serological testing. At the current pace, we can expect nearly half a million new IAV sequences to have been deposited in GISAID by the end of 2023. We can also expect an increasing number of these sequences to be derived from environmental and metagenomic samples [10], where intact virus particles would not be available for serological characterization. Expanded sampling of non-human, non-domesticated host species with these technologies will increase our chances of encountering more novel and highly divergent lineages [7, 29, 30]. Finally, a major advantage of a genetic taxonomy is that sequence-based criteria for defining subtypes can be transparent and reproducible if the clustering methods are described sufficiently well, or better yet, made freely available online.

Data availability

The multiple sequence alignments and maximum likelihood phylogenies for all eight IAV segments in this study are available under a Creative Commons license at https://doi.org/10.5281/zenodo.8119571. The Python and R scripts used to produce and analyze these data are released for unrestricted use and modification under the MIT license at https://github.com/PoonLab/fluclades.

Acknowledgements

This study was motivated and informed by discussions with Richard Neher, who read an earlier version of this manuscript and suggested using mutual information to measure the correlation between phylogenetic subtrees and subtype labels.

References

[1] Tavis K Anderson, Catherine A Macken, Nicola S Lewis, Richard H Scheuermann, Kristien Van Reeth, Ian H Brown, Sabrina L Swenson, Gaëlle Simon, Takehiko Saito, Yohannes Berhane, et al. A phylogeny-based global nomenclature system and automated annotation

tool for H1 hemagglutinin genes from swine influenza A viruses. *MSphere*, 1(6):e00275–16, 2016.

[2] CW Beard. Demonstration of type-specific influenza antibody in mammalian and avian sera by immunodiffusion. *Bulletin of the World Health Organization*, 42(5):779, 1970.

[3] Samir Bhatt, Edward C Holmes, and Oliver G Pybus. The genomic rate of molecular adaptation of the human influenza A virus. *Molecular biology and evolution*, 28(9):2443–2451, 2011.

[4] Rubing Chen and Edward C Holmes. Avian influenza virus exhibits rapid evolutionary dynamics. *Molecular biology and evolution*, 23(12):2336–2341, 2006.

[5] WR Dowdle, JC Galphin, MT Coleman, and GC Schild. A simple double immunodiffusion test for typing influenza viruses. *Bulletin of the World Health Organization*, 51(3):213, 1974.

[6] Frederick L Dunn. Pandemic influenza in 1957: review of international spread of new asian strain. *Journal of the American Medical Association*, 166(10):1140–1148, 1958.

[7] Sasan Fereidouni, Elke Starick, Kobey Karamendin, Cecilia Di Genova, Simon D. Scott, Yelizaveta Khan, Timm Harder, and Aidyn Kydyrmanov. Genetic characterization of a new candidate hemagglutinin subtype of influenza A viruses. *Emerging Microbes & Infections*, page 2225645, 2023. doi: 10.1080/22221751.2023.2225645. URL https://doi.org/10.1080/22221751.2023.2225645. PMID: 37335000.

[8] Ron AM Fouchier, Vincent Munster, Anders Wallensten, Theo M Bestebroer, Sander Herfst, Derek Smith, Guus F Rimmelzwaan, Björn Olsen, and Albert DME Osterhaus. Characterization of a novel influenza A virus hemagglutinin subtype (H16) obtained from black-headed gulls. *Journal of Virology*, 79(5):2814–2822, 2005.

[9] Xingfei Gong, Mingda Hu, Wei Chen, Haoyi Yang, Boqian Wang, Junjie Yue, Yuan Jin, Long Liang, and Hongguang Ren. Reassortment network of influenza A virus. *Frontiers in Microbiology*, page 3759, 2021.

[10] Alexander L Greninger. A decade of RNA virus metagenomics is (not) enough. *Virus Research*, 244:218–229, 2018.

[11] Rong Hai, Florian Krammer, Gene S Tan, Natalie Pica, Dirk Eggink, Jad Maamary, Irina Margine, Randy A Albrecht, and Peter Palese. Influenza viruses expressing chimeric hemagglutinins: globular head and stalk domains derived from different subtypes. *Journal of virology*, 86(10):5774–5781, 2012.

[12] Virginia S Hinshaw and Robert G Webster. Characterization of a new avian influenza virus subtype and proposed designation of this haemagglutinin as Hav10. *Journal of General Virology*, 45(3):751–754, 1979.

[13] VS Hinshaw, GM Air, GC Schild, and RW Newman. Characterization of a novel haemagglutinin subtype (H13) of influenza A viruses from gulls. *Bulletin of the World Health Organization*, 61(4):677, 1983.

[14] Keith E Jensen. New set of type A influenza viruses. *Journal of the American Medical Association*, 164(18):2025–2029, 1957.

[15] Ahmed Kandeil, Mokhtar R Gomaa, Mahmoud M Shehata, Ahmed N El Taweel, Sara H Mahmoud, Ola Bagato, Yassmin Moatasim, Omnia Kutkat, Ahmed S Kayed, Patrick Dawson, et al. Isolation and characterization of a distinct influenza A virus from Egyptian bats. *Journal of Virology*, 93(2):e01059–18, 2019.

[16] Kazutaka Katoh and Daron M Standley. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, 30(4): 772–780, 2013.

[17] Yoshihiro Kawaoka, Svetlana Yamnikova, Thomas M Chambers, Dmitri K Lvov, and Robert G Webster. Molecular characterization of a new hemagglutinin, subtype H14, of influenza A virus. *Virology*, 179(2):759–767, 1990.

[18] G Lang, Bela Tumová, and GC Schild. A new subtype of type A influenzavirus isolated from turkeys. *Bulletin of the World Health Organization*, 47(4):515, 1972.

[19] Anders Larsson. AliView: a fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics*, 30(22):3276–3278, 2014.

[20] Guoqing Lu, Thaine Rowley, Rebecca Garten, and Ruben O Donis. FluGenome: a web tool for genotyping influenza A virus. *Nucleic Acids Research*, 35(suppl_2):W275–W279, 2007.

[21] HG Pereira, G Lang, OM Olesiuk, GH Snoeyenbos, DH Roberts, and BC Easterday. New antigenic variants of avian influenza A viruses. *Bulletin of the World Health Organization*, 35(5):799, 1966.

[22] William H Press, Saul A Teukolsky, W T Vetterling, and Brian P Flannery. Numerical Recipes in C, 1988.

[23] Rochelle Rademan, Marike Geldenhuys, and Wanda Markotter. Detection and Characterization of an H9N2 Influenza A Virus in the Egyptian Rousette Bat in Limpopo, South Africa. *Viruses*, 15(2):498, 2023.

[24] Carolin Röhm, Nannan Zhou, Jochen Süss, John Mackenzie, and Robert G Webster. Characterization of a novel influenza hemagglutinin, H15: criteria for determination of influenza A subtypes. *Virology*, 217(2):508–516, 1996.

[25] Theo Sanderson. Taxonium, a web-based tool for exploring large phylogenetic trees. *Elife*, 11, 2022.

[26] GC Schild, RW Newman, RG Webster, Diane Major, and Virginia S Hinshaw. Antigenic analysis of influenza A virus surface antigens: considerations for the nomenclature of influenza virus. *Comparative Immunology, Microbiology and Infectious Diseases*, 3(1-2):5–18, 1980.

[27] Erica Spackman. Influenza subtype identification with molecular methods. *Animal Influenza Virus*, pages 119–123, 2014.

[28] Eric Talevich, Brandon M Invergo, Peter JA Cock, and Brad A Chapman. Bio.Phylo: a unified toolkit for processing, analyzing and visualizing phylogenetic trees in Biopython. *BMC Bioinformatics*, 13:1–9, 2012.

[29] Suxiang Tong, Yan Li, Pierre Rivailler, Christina Conrardy, Danilo A Alvarez Castillo, Li-Mei Chen, Sergio Recuenco, James A Ellison, Charles T Davis, Ian A York, et al. A distinct lineage of influenza A virus from bats. *Proceedings of the National Academy of Sciences*, 109(11):4269–4274, 2012.

[30] Suxiang Tong, Xueyong Zhu, Yan Li, Mang Shi, Jing Zhang, Melissa Bourgeois, Hua Yang, Xianfeng Chen, Sergio Recuenco, Jorge Gomez, et al. New world bats harbor diverse influenza A viruses. *PLoS Pathogens*, 9(10):e1003657, 2013.

[31] RG Webster, B Tumova, VS Hinshaw, and G Lang. Characterization of avian influenza viruses. designation of a newly recognized haemagglutinin. *Bulletin of the World Health Organization*, 54(5):555, 1976.

[32] WHO/OIE/FAO H5N1 evolution working group. Continued evolution of highly pathogenic avian influenza A (H5N1): updated nomenclature. *Influenza and Other Respiratory Viruses*, 6(1):1–5, 2012.

[33] Frank YK Wong, Celeste Donato, Yi-Mo Deng, Don Teng, Naomi Komadina, Chantal Baas, Joyanta Modak, Mark O'Dea, David W Smith, Paul V Effler, et al. Divergent human-origin

influenza viruses detected in Australian swine populations. *Journal of Virology*, 92(16): e00316–18, 2018.

[34] World Health Organization. A revised system of nomenclature for influenza viruses. *Bulletin of the World Health Organization (WHO)*, pages 119–24, 1971.

[35] Qingye Zhuang, Suchun Wang, Shuo Liu, Guangyu Hou, Jinping Li, Wenming Jiang, Kaicheng Wang, Cheng Peng, Dong Liu, Aizhen Guo, et al. Diversity and distribution of type A influenza viruses: an updated panorama analysis based on protein sequences. *Virology Journal*, 16(1):1–38, 2019.

 Supplementary Figures

Figure S1: Summary of the number of influenza A virus sequences in the GISAID database by year of sample collection. Note that some sequences were collected prior to the establishment of this database in 2006, and submitted retrospectively at a later date. The number of sequences per isolate varies not only because of varying numbers of segments sequenced, but also because some isolates were sequenced more than once. This trend features a conspicuous surge in the number of sequences associated with the 2009 H1N1 pandemic, as well a drop in numbers in 2019-2020 in association with the SARS-CoV-2 pandemic. Overall, the number of sequences is exponentially growing over time (adjusted $R^2 = 0.926$).

Figure S2: Inferred subtypes for HA (left) and NA(right) sequences with missing subtype annotations (*e.g.*, 'unknown', 'HX' or 'mixed') compared to their respective known distributions. Subtypes were inferred by retrieving each sequence's nearest neighbour in the phylogeny. Dashed lines represent linear regressions on counts excluding the most common subtypes (H1, H3, H5 and H9 for hemagglutinin; N1 and N2 for neuraminidase).

Figure S3: Evaluation of concordance between HA subtype annotations (labels) and subtrees defined by node-level clustering, under varying thresholds. (A) The size of each point is scaled to the discordance between the number of subtrees and the target number ($n = 18$), *i.e.*, smaller is better. Pink indicates too many subtrees, and blue indicates too few. (B) The number of different labels increases as a function of the mean patristic distance cutoff (y). The ideal number is one label per subtree. The minimum divergence cutoff V had a limited effect on this outcome. (C) The proportion of labels that are not associated with any subtree at given cutoffs are represented by filled squares. For reference, an outline is drawn for each square to represent 100% loss. (D) The distribution of labels among subtrees defined by cutoffs at which the ideal number of subtrees is obtained ($y = 1.2$ and $v = 0.08$). Regions are shaded in proportion to the fraction of each label in the subtree, and the number of labels is displayed if the fraction is ≤ 0.5. The total number of labels per subtree is displayed along the right margin.

Figure S4: Number of subtrees as a function of internal branch length cutoffs applied to the phylogenies of other proteins. The plots are ordered with respect to genome segment numbering, *e.g.*, protein PB2 is encoded by segment 1. Solid triangles mark the cutoffs where the composition of the resulting subtrees are summarized in Supplementary Table S4.

 Supplementary Tables

Segment	Gene products	Min. length	Total	Unique
1	PB2	2000	93,415	32,718
2	PB1, *PB1-F2*	2000	160,506	29,790
3	PA, *PA-X*	2000	169,484	32,349
4	HA	1600	137,172	66,874
5	NP	1400	103,033	18,978
6	NA	1000	126,233	52,146
7	M1, M2	900	213,717	14,389
8	NS1, NS2	750	191,676	25,172

Table S1: Summary of amino acid sequences obtained for all eight IAV segments from the NCBI Genbank nucleotide database (date accessed, June 1, 2023 except for HA sequences, which were retrieved April 28, 2023). Amino acid sequences representing PB1-F2 and PA-X (italics) were excluded from subsequent analyses. The M1/M2 and NS1/NS2 sequences were concatenated with amino acid residues within overlapping regions excluded. Min. length = minimum sequence length in nucleotides. Total = number of sequence records returned by query. Unique = number of unique sequences after filtering.

Accession	Strain	Original	Corrected
ON427060	A/swine/Kansas/A02711700/2022	H1	H3
MF098827	A/swine/Chile/VN1401-587/2014	H1	H3
MT377993	A/swine/Thailand/CU22337/2018	H1	H3
MH596892	A/ruddy turnstone/Delaware/650622/2002	H11	H7
MN569599	A/Newcastle/177/2017	H3	H1
MN574095	A/Victoria/79/2017	H3	H1
KM110063	A/swine/Hanoi/568/2013	H3	H1
KM110062	A/swine/Hanoi/424/2013	H3	H1
KM110061	A/swine/Hanoi/411/2013	H3	H1
CY251407	A/Victoria/1003A/2012	H3	H1
OP888980	A/yellow-billed	H6	H1
LC500374	A/duck/Vietnam/HU8-1918/2017	H3	H5
MH598081	A/ruddy turnstone/New Jersey/650677/2002	H11	H6
MH598105	A/ruddy turnstone/New Jersey/828212/2001	H5	H12
MH130194	A/mallard/Korea/M198/2013	H4	H9

Table S2: Reclassification of IAV HA sequences with serotype (subtype) annotations that are not consistent with their location in the phylogeny. Discordant sequences were detected as outliers in box-and-whisker plots of their vertical positions in the rectangular layout of the tree, grouped by subtype annotations. The predicted subtypes were estimated from the nearest labeled tips in the tree. These predictions were manually verified using nucleotide BLAST.

Accession	Strain	Original	Corrected
MK960206	A/Texas/9437/2019	N1	N2
OP940105	A/swine/Iowa/G92995/2022	N1	N2
OP645900	A/swine/Iowa/C61728/2022	N1	N2
OP940090	A/swine/Illinois/E80174/2022	N1	N2
MW839927	A/swine/Iowa/CD81015/2015	N1	N2
MW839656	A/swine/North Carolina/LJ8316/2016	N1	N2
MT377995	A/swine/Thailand/CU22337/2018	N1	N2
MF099208	A/swine/Chile/VN1401-587/2014	N1	N2
MT378776	A/swine/France/22-180459/2018	N1	N2
MW362711	A/swine/Belgium/Gent-36/2016	N1	N2
MT818840	A/Mallard/North Carolina/AH0114261I.6.A/2017	N8	N2
MW513533	A/green-winged teal/Alaska/UGAI16-4602/2016	N8	N2
MF613906	A/ruddy turnstone/Delaware Bay/303/2016	N4	N2
MH597119	A/ruddy turnstone/Delaware/1016391/2003	N5	N2
MH597724	A/ruddy turnstone/New Jersey/1321395/2005	N8	N6
MH597785	A/ruddy turnstone/New Jersey/1321408B/2005	N8	N6
MH598073	A/ruddy turnstone/New Jersey/650675/2002	N9	N5
MH597447	A/ruddy turnstone/Delaware/650647/2002	N9	N5
L06581	A/equine/New Market/1979	N3	N8
MH597692	A/ruddy turnstone/New Jersey/1148671/2004	N7	N8
MH596859	A/ruddy turnstone/Delaware/1016396/2003	N9	N8
LC431470	A/pintail/Alberta/121/1979	N8	N4
CY251409	A/Victoria/1003A/2012	N2	N1
MH598004	A/ruddy turnstone/New Jersey/650584/2002	N9	N1
MH596894	A/ruddy turnstone/Delaware/650622/2002	N4	N1
MH579375	A/wild waterfowl/Korea/M54-F-1/2010	N2	N1
MK636253	A/New York/7745/2018	N2	N1
ON848942	A/Homosapiens/India/149/2017	N2	N1
MN568417	A/Brisbane/144/2017	N2	N1
MN574097	A/Victoria/79/2017	N2	N1

Table S3: Reclassification of IAV NA sequences with serotype (subtype) annotations using the same methods described in Table S2.

Protein	# tips	Example strain	HnNn
PB2	32,700		
	4	A/equine/Prague/1/1956	H7N7
	4	A/dark fruit-eating bat/Bolivia/PBV780-781/2011	H18N11
	2	A/Bat/Egypt/381OP/2017	H9N2
	2	A/little yellow-shouldered bat/Guatemala/164/2009	H17N10
PB1	29,685		
	5	A/equine/Prague/1/1956	H7N7
	4	A/flat-faced bat/Peru/033/2010	H18N11
	2	A/little yellow-shouldered bat/Guatemala/164/2009	H17N10
PA	32,325		
	3	A/equine/Prague/1/1956	H7N7
	3	A/dark fruit-eating bat/Bolivia/PBV780-781/2011	H18N11
	2	A/Bat/Egypt/381OP/2017	H9N2
	2	A/little yellow-shouldered bat/Guatemala/164/2009	H17N10
NP	18,970		
	4	A/dark fruit-eating bat/Bolivia/PBV780-781/2011	H18N11
	2	A/Bat/Egypt/381OP/2017	H9N2
	2	A/little yellow-shouldered bat/Guatemala/164/2009	H17N10
M1/M2	14,367		
	12	A/equine/Lexington/1/1966	H7N7
	3	A/swine/Western Australia/AS12-2111-02/2012	H3N2
	3	A/Artibeus lituratus/Brazil/2344/2012	H18N11
	2	A/Bat/Egypt/381OP/2017	H9N2
	2	A/little yellow-shouldered bat/Guatemala/164/2009	H17N10
NS1/NS2	23,380		
	4	A/equine/Prague/1956	H7N7
	2	A/Bat/Egypt/381OP/2017	H9N2
	2	A/little yellow-shouldered bat/Guatemala/164/2009	H17N10
	2	A/flat-faced bat/Peru/033/2010	H18N11

Table S4: Composition of subtrees produced at selected branch length cutoffs for phylogenies relating other protein sequences. Cutoffs were chosen based on the trends displayed in Supplementary Figure S4. Example strain and HA/NA subtype labels (HnNn) are omitted for the largest subtrees.