The Intricate Relationship of G-Quadruplexes and

Pathogenicity Islands: A Window into Bacterial Pathogenicity

Bo Lyu*

Division of Plant Science and Technology, University of Missouri, Columbia, MO

65211, USA

*Corresponding author: Bo Lyu; E-mail: bl3pt@missouri.edu; Phone number: +1 573

289 3566;

Abstract

The dynamic interplay between guanine-quadruplex (G4) structures and pathogenicity islands (PAIs) represents a captivating area of research with implications for understanding the molecular mechanisms underlying pathogenicity. This study conducted a comprehensive analysis of a large-scale dataset from reported 89 pathogenic strains to investigate the potential interactions between G4 structures and PAIs. G4 structures exhibited an uneven and non-random distribution within the PAIs and were consistently conserved within the same pathogenic strains. Additionally, this investigation identified positive correlations between the number and frequency of G4s and the GC content across different genomic features, including the genome, promoters, genes, tRNA, and rRNA regions, indicating a potential relationship between G4 structures and the GC-associated regions of the genome. The observed differences in GC content between PAIs and the core genome further highlight the unique nature of PAIs and underlying factors, such as DNA topology. High-confidence G4 structures within regulatory regions of *Escherichia coli* were identified, modulating the efficiency or specificity of DNA integration events within PAIs. Collectively, these findings pave the way for future research to unravel the intricate molecular mechanisms and functional implications of G4-PAI interactions, thereby advancing our understanding of bacterial pathogenicity and the role of G4 structures in infectious diseases.

Key Words: Guanine quadruplex; Pathogenicity island; Genome; *Escherichia coli*

Introduction

The discovery of the DNA double helix by Watson and Crick in 1953 revolutionized our understanding of genetics and laid the foundation for the modern field of molecular biology (*1*). Nonetheless, the intricate nature of DNA continues to surprise us even today. One such captivating feature is the DNA guanine (G)-quadruplex (G4) structure, a unique arrangement that defies the conventional double helix (*2, 3*). A G4 consists of four guanine bases and stabilized by Hoogsteen hydrogen bonding. These stacked tetrads are interconnected by loop regions, which can vary in length and sequence, adding further complexity to the structure (Fig. 1). It is important to consider the inherent directionality of nucleic acids, with all four strands having the possibility to run in the same 5' to 3' direction, referred to as "parallel", or alternatively, they can run in different directions, known as "antiparallel". G4 regions can be very stable *in vitro*, particularly in the presence of K^+ (*4*).

Pathogenicity islands (PAIs) are genomic regions that contribute to the virulence and pathogenic potential of various microorganisms (*5, 6*). PAIs often exhibit close proximity to tRNA genes, suggesting a putative mechanism where tRNA genes act as anchor points for the integration of foreign DNA acquired through horizontal gene transfer (Fig. 1E). One notable feature is their variable GC content, which tends to deviate from the average GC content of the genome in various organisms, such as *Streptomyces* (*7*), *Salmonella* (*8*), and *Yersinia* (*9*). One question raised in PAI is that PAIs often exhibit distinct base composition (G+C contents) compared to the core genome. The underlying reasons for this variation remain unknown, but the

preservation of a genus- or species-specific base composition represents a noteworthy

characteristic of bacteria (*5*). Herbert Schmidt et al. (2004) proposed a hypothetical

mechanism to explain the observed variation, suggesting that factors such as DNA

topology and codon message in the virulence regions present could contribute to the

preservation of the distinct base composition (*5*). Hopefully, the availability of

genome sequences from pathogenic bacteria and their non-pathogenic counterparts

currently presents an exceptional opportunity to explore the intricate structure

variance and underlying mechanisms within PAIs.

The possibility of interactions between G4 structures and pathogens has been

suggested, although this field of study is still in its nascent phase. Some studies

observed that bacterial genomes possess G4-forming sequences within their genome

regions (*10, 11*). G4 structures are formed by G-rich DNA sequences, and their

stability is influenced by the G+C content and arrangement of G tetrads. Interestingly,

PAIs often exhibit an altered GC content, putatively contributing to the propensity of

G4 structure formation within these regions. The G4 structures in PAIs might

modulate the accessibility of transcription factors, DNA-binding proteins, or RNA

polymerase in pathogens, as documented in eukaryotes (*2, 12*), thereby influencing

the expression of virulence-associated genes. The formation of G4 structures within

PAIs may serve as an additional layer of regulation that fine-tunes the expression of

genes critical for pathogenesis. Hence, the investigation of G4 structures within PAIs

may open new avenues for the development of therapeutic strategies aimed at

disrupting the regulatory mechanisms of pathogenicity-associated genes.

Results and Discussion

Genomic information, PAI patterns, and the presence of G4s in 89 reported

pathogenic strains

A dataset of PAIs was compiled from 89 reported pathogenic strains, encompassing

222 distinct types of PAIs. Those pathogens that share closely related PAIs on

phylogenetic branches, such as LEE in *E. coli* strains (Fig. 2A), may have acquired

these elements through recent common ancestors or horizontal gene transfer events (*5,*

13, 14). Additional information, including the genome length (bp), G+C content (%),

rRNA density, tRNA density, and PAI length (bp), was present and showed conserved

patterns in the same species (Fig. 2A & Table S1). PAIs commonly exhibit mosaic-

like patterns, exemplified by the presence of distinct PAIs like FPI in *Francisella*

tularensis, SaPIbov in *Staphylococcus aureus*, and Hrp PAI in *Xanthomonas*

campestris (Fig. 2B). Many PAIs were present associated with tRNAs, such as the

insertions of tRNA^{Thr}, tRNA^{Phe}, and tRNA^{Gly} in *E. coli* strains (Fig. 2B & Table S2).

The presence of PAIs distributes in similar genomic regions across different

pathogens or strains, showing non-random patterns and functionally clustered.

Employing the G4Hunter search algorithm, the study identified a total of 225,376

putative G4 sequences in these 89 pathogenic genomes (Table S1). The heatpmap

showed that the number of G4 structures was diverse in the pathogen genomes (Fig.

2C), similar to previous reports that showed that the G4 structures display uneven

distribution patterns in eukaryotic and prokaryotic genomes and are conserved in

evolutionary groups (*15-17*).

Interaction between PAIs and G4s in different genomic features

The analysis of G4 structures across all pathogen species demonstrated a positive

correlation between the number of G4s and the GC content in various genomic

features, including the whole genome, gene, promoter, rRNA, and tRNA regions (Fig.

2D). Furthermore, the density of G4s, measured as the frequency of predicted G4-

forming sequences per 1000 base pairs (bp), also showed a positive correlation with

the GC content across the analyzed genomic elements (Fig. 2E). The presence of G4

structures within the promoter, rRNA, and tRNA regions may have functional

implications for the regulation of DNA replication, ribosome biogenesis, protein

synthesis, and other RNA-related processes (*18-20*). Throughout evolution, there

seems to be a greater enrichment of G4 structures in regulatory genes compared to

other genes, enabling intricate control of gene expression in cascade signal

transduction pathways (*21*).

This study observed that the GC contents in the genome region were significantly

higher compared to the corresponding PAIs region that was classified into five parts

according to the genome datasets (Fig. 3A-E & Table S1). This suggests that the

genomic regions surrounding the PAIs (i.e., core genome) tend to have a higher GC

content than PAI regions, which was consistent with the fact that PAIs often exhibit

distinct base composition compared with the core genome (*5*). The variation was

explained by the presence of G4 sequences within the PAIs, whereas the results were

surprising. This study observed a distinct pattern in the frequency of G4 structures

within different regions of the PAIs. This differential distribution of G4 structures

suggests that (i) specific segments (e.g., <30% and >60% GC contents) within the

PAIs may be more prone to induce G4 formation discrepancy, as these segments

showed unstable gene structure; (ii) the variation of base composition between core

genome and PAIs is partially correlated with the presence of G4s; (iii) the frequency

of G4s in PAIs present stable as the core genome in the most situation; (iv) an

alternative hypothesis, other factors, such as i-motif (i.e., the anti-G4 structure) and

CpG island, may work synergistically with G4 and potentially contribute the base

composition variation (*22, 23*).

Putative origin, transfer mechanisms, and functions of G4s in PAIs

To understand the origin of G4 structures within PAIs, the study hypothesized that

these G4 sequences could be acquired through three types of horizontal gene transfer

mechanisms: conjugation, transformation, and transduction (Fig. 3F). These

mechanisms serve as means for genetic material exchange between different

organisms. Considering the presence of G4 sequences within the PAIs, it is plausible

that these sequences are transferred along with the PAIs through these horizontal gene

transfer mechanisms. Later, the study used *E. coli as an example* to investigate the

potential regulatory role and function of genes covered by G4 structures in PAIs. *E.*

coli contains at least ten types of PAI in different strains, and one of the well-known

PAI is LEE (Fig. 3G), harboring genes responsible for causing attaching and effacing

lesions (*14, 24*). One stable G4 structure with a G4hunter score of 1.6 was identified

at position 37,085 in the LEE PAI of *E. coli* str. O103:H2 12009 (Fig. 3H), located

between an IS element and a tRNA insertion site. The tRNA region generally contains

a higher G4 frequency compared with transfer-messenger RNA (tmRNA) and rRNA

regions in the bacterial genome (*16*). Interestingly, this G4 structure found in *E. coli*

str. O103:H2 12009 was present in close proximity to a tRNA region, suggesting a

potential regulatory role of G4s in the tRNA gene, or upstream- and downstream-

genes that are responsible for LEE virulence. Additionally, another stable G4

sequence with a score of 1.381 was discovered at position 12,457 in the *E. coli str.*

CFT073 PAI II to provide one more evidence of G4 in PAI regions (Fig. 3I).

Functional enrichment analysis was conducted to explore the putative functions

of G4-covered genes in two *E. coli* strains (Table S3&S4). The results revealed that

the genes covered by G4 structures were predominantly involved in genetic

information processes, including DNA binding, DNA integration, and nucleic acid

metabolism processes (Fig. 3J & K). This suggests that G4 structures may play a

regulatory role in these essential cellular processes, especially gene expression and

DNA-related functions. For instance, G4 structures in the promoter regions of certain

transcription factors may influence their binding affinity to DNA and subsequently

affect downstream gene expression patterns (*25, 26*). These elements frequently

utilize DNA integration mechanisms mediated by integrases, recombinases, or

transposases to transfer or incorporate genetic material into the bacterial genome (*27,*

28).

Overall, the conserved evolutionary relatedness of PAIs, the detection of stable

G4 structures in specific genomic positions, and the enrichment of G4-covered genes

in genetic information processes collectively support the hypothesis that G4 structures

may have regulatory functions in key biological processes in pathogens.

Methods

Selection and extraction of DNA Sequences

A total of 89 genomes corresponding to the identified pathogens from the

Pathogenicity Island Database (PAIDB) were included in the study. The complete

bacterial genomic DNA sequences and their corresponding annotation files in .gff

and .fna formats were obtained from the Genome database of the National Center for

Biotechnology Information (NCBI). To ensure the reliability and completeness of the

dataset, only completely assembled genomes were included in the analysis. To avoid

redundancy and incomplete sequences, one representative genome was selected for

each species or strain. The selection of representative genomes was based on a careful

examination of the supplementary material (Table S1) accompanying the study. The

study employed TBtools (https://cj-chen.github.io/tbtools), a versatile bioinformatics

tool, to investigate the presence of G-quadruplex (G4) sequences in specific genomic

features. The gene regions, promoters (2 kb upstream of the genes), tRNA regions,

and rRNA regions were extracted from the selected genomes using TBtools. PAI

regions were downloaded following previously documented information in PAIDB

(Table S1&S2). Default thresholds and parameters were applied during extraction to

maintain consistency across all genomes.

Data process and detection of G4s in genomic features

The G4Hunter algorithm, a widely used tool for G4 prediction, was employed for the

identification of G4 motifs in the genomic sequences. The G4Hunter parameters were

set to a window size of "25" and a G4 score threshold of 1.2, which ensured the

identification of potential G4 sequences. The analysis was performed for each species

group, where the genomic DNA size and the number of putative G4 sequences found

were determined. The study quantified the predicted number of putative G4-forming

sequences within different genomic features, including the whole genome, gene,

promoter, tRNA, rRNA, and PAI regions. The density of G4 motifs was determined

by dividing the number of G4 sequences by the total length of the genome, while the

length ratio of G4 motifs was calculated by dividing the total length of the G4

sequences by the total length of the genome.

Relationship between G4s and PAIs

The heatmap was used to show the distribution of G4 motifs in the genome divided by

ten parts as PAI regions using R package "pheatmap". The correlation between the

number of G4 structures and the GC content was analyzed across various genomic

elements, including the whole genome, gene, promoter, rRNA, and tRNA regions.

The analysis utilized the R-squared value (R^2) to determine the strength of the

correlation. The significance of the correlation was assessed using p-values. The GC

content in the genome regions and corresponding PAI regions was compared and

classified into different ranges to explore the variation in base composition. GraphPad

Prism (V.5.02, GraphPad Software, Inc) was employed to conduct Normality and

Lognormality Tests. The K-S test and F-test were used to assess normal distribution

and variances, and Student's t-test was used to identify significant difference.

Phylogenetic Tree Construction

The exact Taxonomy ID (taxid) for each analyzed group was obtained from the NCBI

Taxonomy Database using the Taxonomy Browser. The Neighbor-Joining (NJ)

method was employed to construct the phylogenetic trees for the analyzed groups.

The phylogenetic trees were generated using MEGA11 software

(www.megasoftware.net), which offers robust algorithms and comprehensive tools for

phylogenetic analysis. To assess the reliability and statistical support of the

phylogenetic tree branches, bootstrap analysis was performed. One thousand

bootstrap replicates were used to estimate the confidence levels of the branching

patterns in the phylogenetic trees. The phylogenetic trees, along with the bootstrap

support values, were displayed and visualized using the Interactive Tree Of Life

(ITOL) platform (https://itol.embl.de/).

Gene functional annotation

The gene sequences covered by G4 structures within PAIs were subjected to gene

ontology annotation. The gene sequences were translated into protein sequences using

the Expasy online toolkit (https://web.expasy.org/translate/). This tool performs the

translation based on the standard genetic code, converting the DNA nucleotide

sequence into its corresponding amino acid sequence. The GO annotation database

assigned GO terms to the protein sequences based on their predicted functions and

known biological process (BP), molecular function (MF), and cellular component

(CC). Fisher's exact test was employed to determine the statistical significance of the

enrichment results. The obtained p-values indicated the overrepresentation of specific

GO terms, with lower p-values suggesting higher significance.

Statistics and reproducibility

All genomic data utilized in this study, including the species-specific datasets, were

obtained from publicly available sources. Statistical analyses, such as the Kruskal-

Wallis test, F test, Student's t-test, Wilcoxon test, and chi-square test, were performed

using GraphPad Prism software. The samples used in the statistical analyses

corresponded to the genomic data, PAIs, or specific genes under investigation.

Acknowledgment

The sincere appreciation extends to Dr. Sung Ho Yoon and his colleagues for their

dedicated efforts in identifying PAIs and establishing the Pathogenicity Island

Database for public analysis. Their commitment to advancing the field of pathogen

genomics has greatly facilitated this research. This study would like to thank Prof.

Qisheng Song and Dr. Jingjing Li for their insightful suggestions and constructive

comments regarding the exploration of G4 structures in genomes. Their expertise and

guidance have significantly enriched the understanding of the potential roles and

implications of G4 structures in the context of PAIs. The study is grateful to Dr.

Minyu Zhou for her valuable discussions and exchange of ideas regarding the

interplay between G4 structures and pathogenicity islands.

Competing Interests

The author declares no competing interests.

Funding information

This research received no external funding.

Credit author statement

Bo Lyu: writing original draft, data analysis, and visualization.

Ethical approval

Ethical review and approval were not required for this study.

Availability of data and materials

The original reported PAIs datasets analyzed in this study are available from the

publication with the DOI: 10.1093/nar/gku985. Additionally, supplementary materials

accompanying this article provide further PAIs data analyzed in the study. These

supplementary materials contain detailed information and datasets that support the

findings and conclusions presented in this research. Researchers and interested

individuals can access supplementary materials to explore and validate the results

obtained in this study.

Consent for publication

Not applicable.

1.   J. D. Watson, F. Crick, A structure for deoxyribose nucleic acid.
(1953).

2.   D. Rhodes, H. J. Lipps, G-quadruplexes and their regulatory roles in
biology. *Nucleic acids research* **43**, 8627-8637 (2015).

3.   J. Spiegel, S. Adhikari, S. Balasubramanian, The structure and
function of DNA G-quadruplexes. *Trends in Chemistry* **2**, 123-136 (2020).

4.   O. Stegle, L. Payet, J.-L. Mergny, D. J. MacKay, J. L. Huppert,
Predicting and understanding the stability of G-quadruplexes. *Bioinformatics*
25, i374-i1382 (2009).

5.   H. Schmidt, M. Hensel, Pathogenicity islands in bacterial
pathogenesis. *Clinical microbiology reviews* **17**, 14-56 (2004).

6.   E. A. Groisman, H. Ochman, Pathogenicity islands: bacterial
evolution in quantum leaps. *Cell* **87**, 791-794 (1996).

7.   J. A. Kers *et al.*, A large, mobile pathogenicity island confers plant
pathogenicity on Streptomyces species. *Molecular Microbiology* **55**, 1025-
1033 (2005).

8.   S. Kombade, N. Kaur, in *Salmonella spp.-A Global Challenge*.
(IntechOpen, 2021).

9.   E. Carniel, The Yersinia high-pathogenicity island. *Int Microbiol* **2**,
161-167 (1999).

10. P. Yadav *et al.*, G-quadruplex structures in bacteria: Biological
relevance and potential as an antimicrobial target. *Journal of Bacteriology* **203**,
10.1128/jb. 00577-00520 (2021).

11. L. M. Harris, C. J. Merrick, G-quadruplexes in pathogens: a
common route to virulence control? *PLoS pathogens* **11**, e1004562 (2015).

12. D. Varshney, J. Spiegel, K. Zyner, D. Tannahill, S. Balasubramanian,
The regulation and functions of DNA and RNA G-quadruplexes. *Nature*
reviews Molecular cell biology **21**, 459-474 (2020).

13. E. L. Hartland, J. M. Leong. (Frontiers Media SA, 2013), vol. 3, pp.
15.

14. J. Jores, L. Rumer, L. H. Wieler, Impact of the locus of enterocyte
effacement pathogenicity island on the evolution of pathogenic Escherichia

coli. *International Journal of Medical Microbiology* **294**, 103-113 (2004).

15. Z. Du, Y. Zhao, N. Li, Genome-wide colonization of gene regulatory
elements by G4 DNA motifs. *Nucleic acids research* **37**, 6784-6798 (2009).

16. M. Bartas *et al.*, The presence and localization of G-quadruplex
forming sequences in the domain of bacteria. *Molecules* **24**, 1711 (2019).

17. E. P. Lombardi, A. Londoño-Vallejo, A. Nicolas, Relationship
between G-quadruplex sequence composition in viruses and their hosts.
Molecules **24**, 1942 (2019).

18. P. Ivanov *et al.*, G-quadruplex structures contribute to the
neuroprotective effects of angiogenin-induced tRNA fragments. *Proceedings*
of the National Academy of Sciences **111**, 18201-18206 (2014).

19. S. Mestre-Fos *et al.*, G-quadruplexes in human ribosomal RNA.
Journal of molecular biology **431**, 1940-1955 (2019).

20. B. Zybailov, M. Sherpa, G. Glazko, K. Raney, V. Glazko, G4-
quadruplexes and genome instability. *Molecular Biology* **47**, 197-204 (2013).

21. F. Wu *et al.*, Genome-wide analysis of DNA G-quadruplex motifs
across 37 species provides insights into G4 evolution. *Communications*
biology **4**, 98 (2021).

22. A. M. Deaton, A. Bird, CpG islands and the regulation of
transcription. *Genes Dev* **25**, 1010-1022 (2011).

23. N. Sushmita *et al.*, I-motif DNA: significance and future prospective.
Exploratory Animal and Medical Research **10**, 18-23 (2020).

24. F. M. Franzin, M. P. Sircili, Locus of enterocyte effacement: a
pathogenicity island involved in the virulence of enteropathogenic and
enterohemorragic Escherichia coli subjected to a complex network of gene
regulation. *Biomed Res Int* **2015**, 534738 (2015).

25. K. Niu *et al.*, BmILF and i-motif structure are involved in
transcriptional regulation of BmPOUM2 in Bombyx mori. *Nucleic Acids*
Research **46**, 1710-1723 (2018).

26. L. Xiang *et al.*, DNA G-quadruplex structure participates in
regulation of lipid metabolism through acyl-CoA binding protein. *Nucleic*
Acids Research **50**, 6953-6967 (2022).

323 27. I. R. Arkhipova, P. A. Rice, Mobile genetic elements: in silico, in
324 vitro, in vivo. *Mol Ecol* **25**, 1027-1031 (2016).

325 28. R. A. Wozniak, M. K. Waldor, Integrative and conjugative elements:
326 mosaic mobile genetic elements enabling dynamic lateral gene flow. *Nature*
327 *Reviews Microbiology* **8**, 552-563 (2010).

Figure 1. Structural and Functional Aspects of G-quadruplex (G4) Structures

and Pathogenicity Islands (PAIs). (A) Schematic representation of a guanine tetrad

stabilized by Hoogsten base pairing and a positively charged central ion, illustrating

the key elements of G4 structures. (B) Structural heterogeneity of G4s. G-quadruplex

structures exhibit polymorphism and can be categorized into different families, such

as parallel or antiparallel, based on the orientation of the DNA strands. They can fold

either intramolecularly or intermolecularly, leading to diverse structural

configurations. (C) General sequence formula for G4s, highlighting the repeated

occurrence of guanine-rich sequences that form G-quartets. (D) Regulatory roles of

G-quadruplexes in transcription. G4 structures can regulate transcription by blocking

RNA polymerase from binding to promoter sequences or aiding in single-stranded

DNA (ssDNA) formation, thereby enhancing transcription. (E) General structure of

PAI. PAIs are characteristic regions of DNA found within the genomes of pathogenic

bacteria, distinguishing them from nonpathogenic strains of the same or related

species. Repeat sequences are DNA segments duplicated within the PAI and can serve

as recognition sites for various enzymes involved in the integration and excision of

the PAI from the bacterial chromosome. tRNA genes act as anchor points for the

insertion of foreign DNA acquired through horizontal gene transfer. Virulence genes

encode proteins or factors that play crucial roles in the virulence and pathogenicity of

the bacterium, contributing to adhesion, invasion, immune evasion, toxin production,

or other pathogenic mechanisms. Insertion elements include transposons,

bacteriophages, or plasmids, enabling the PAI to be transferred between bacterial cells

and potentially disseminated to different strains or species.

Figure 2. Analysis of Pathogenicity Islands (PAIs) and G-quadruplexes (G4s) in Pathogen Genomes. (A) Phylogenetic analysis of pathogen genomes based on 89 bacterial strains, showing the evolutionary relationships among species. Additional genomic information, including genome size, GC content, rRNA density, tRNA density, and PAI length, is provided. The same color indicates the same species. (B) Genomic location of specific PAIs in bacterial genomes, divided into ten regions. PAIs are represented by green triangles, and their names are indicated. The tRNA insertion sites are also marked. (c) Heatmap illustrating the relative abundance of G4s in bacterial genomes, divided into ten regions. Red indicates a higher relative abundance, while blue indicates a lower relative abundance. (D & E) Correlation analysis between the number of G4s, the frequency of G4s and GC content in various genomic features, including the whole genome, genes, promoters, rRNA, and tRNA.

Figure 3. Putative Origin and Functional Annotation of G-quadruplexes (G4s)

within Pathogenicity Islands (PAIs) in *Escherichia coli*. (A-E) Comparison of GC

content (left panel) and GC frequency (right panel) between the genome and

pathogenicity islands (PAIs), categorized into five regions (20%-30%, 30%-40%,

40%-50%, 50%-60%, and 60%-70%). */**/***/**** indicates significant difference

($p < 0.05/0.01/0.001/0.0001$). (F) Hypotheses on the origin of G4 structures within

PAIs, involving gene horizontal transfer mechanisms (conjugation, transduction, and

transformation). (G) Evolutionary relatedness of 10 types of PAIs (categorized into

six main categories) in E. coli strains. (H & I) Examples of G4 structures within PAIs

in E. coli strains. The grey bar represents the virulence region, the red box indicates a

50   virulence gene, the blue box represents an insertion site region or repeat, the green

51   box denotes an integrase, the purple triangle indicates a tRNA insertion site, and the

52   yellow triangle indicates an effector. (J &K) Functional annotation analysis of G4-

53   covered genes within PAIs in two E. coli strains, including biological process (BP),

54   cellular component (CC), and molecular function (MF) categories.

Flagellum

Capsule Nucleoid

Pili

Ribosome

G4

K⁺

phe | Integrase | tRNA | Vir I | IS | | G4 | Effector | Vir II | Vir III

GGGGTGGGCGAGAGCCGGGACAGGG

Transcription

Translation