CLIPB4 is a central node in the protease network that regulates humoral immunity in *Anopheles gambiae* mosquitoes

Xiufeng Zhang[a], Shasha Zhang[a,b], Junyao Kuang[c], Kathleen A. Sellens[d], Bianca Morejon[a], Sally A. Saab[e], Miao Li[a], Eve C. Metto[d], Chunju An[b], Christopher T. Culbertson[d], Mike A. Osta[e], Caterina Scoglio[c], Kristin Michel[a]

[a] Division of Biology, Kansas State University, Manhattan, KS, USA
[b] Department of Entomology, College of Plant Protection, China Agricultural University, Beijing, China
[c] Department of Electrical and Computer Engineering, Kansas State University, Manhattan, KS 66506, USA
[d] Department of Chemistry, Kansas State University, Manhattan, KS, USA
[e] Department of Biology, American University of Beirut, Beirut, Lebanon

Author contact information: Xiufeng Zhang, xiufengz@ksu.edu; Shasha Zhang, zhangshasha@bjfc.cc; Junyao Kuang, junyao.kuang@crownbio.com; Kathleen A. Sellens, sellens2@yahoo.com; Bianca Morejon, bmorejon744@ksu.edu; Sally A. Saab, ssaab4@jh.edu; Eve C. Metto, emetto@d.umn.edu; Miao Li, limiao@ksu.edu; Mike A. Osta, mo07@aub.edu.lb; Chunju An, 201118@cau.edu.cn; Christopher T. Culbertson, culbert@ksu.edu; Caterina Scoglio, caterina@ksu.edu; Kristin Michel, kmichel@ksu.edu.

Short Title: CLIPB4's central role in mosquito melanization

Corresponding Author:
Kristin Michel
Division of Biology
Kansas State University
Manhattan, KS 66506, USA
Telephone: +1-785-532-0161
Email: kmichel@ksu.edu

Number of Tables: 1 main and 4 online supplementary tables
Number of Figures: 6 main and 6 online supplementary figures

Word count: 7,788

Key words: *Anopheles gambiae*, clip domain serine protease, insect immunity, melanization, phenoloxidase, serpin

Abstract

Insect humoral immune responses are regulated in part by protease cascades, whose components circulate as zymogens in the hemolymph. In mosquitoes, these cascades consist of <u>c</u>lip domain <u>s</u>erine <u>p</u>roteases (cSPs) and/or their non-catalytic <u>h</u>omologs (cSPHs), which form a complex network, whose molecular make-up is not fully understood. Using a systems biology approach, based on a co-expression network of gene family members that function in melanization and co-immunoprecipitation using the serine protease inhibitor (SRPN)2, a key negative regulator of the melanization response in mosquitoes, we identify the cSP CLIPB4 from the African malaria mosquito *Anopheles gambiae* as a central node in this protease network. *CLIPB4* is tightly co-expressed with *SRPN2* and forms protein complexes with SRPN2 in the hemolymph of immune-challenged female mosquitoes. Genetic and biochemical approaches validate our network analysis and show that CLIPB4 is required for melanization and antibacterial immunity, acting as a prophenoloxidase (proPO)-activating protease, which is inhibited by SRPN2. In addition, we provide novel insight into the structural organization of the cSP network in *An. gambiae*, by demonstrating that CLIPB4 is able to activate proCLIPB8, a cSP upstream of the proPO-activating protease CLIPB9. These data provide the first evidence that, in mosquitoes, cSPs provide branching points in immune protease networks and deliver positive reinforcement in proPO activation cascades.

Introduction

The African malaria mosquito, *Anopheles gambiae*, is one of the efficient vectors of life-threatening human malaria parasites. In 2021, approximately 247 million cases and 619,000 deaths were reported of malaria worldwide (1). Despite the availability of an effective, albeit non-sterilizing, vaccine, vector population control remains the most common and effective malaria prevention tool, which is threatened by insecticide resistance in target mosquito populations. Novel vector population control schemes, utilizing gene drive, male sterility, and *Wolbachia*-based methods (2–4) in part rely on detailed molecular understanding of mosquito physiology. For this reason, mosquito innate immunity has received much attention, as it is a major determinant of vector competence and survival of mosquitoes within and across their life stages (5–7).

Protease cascades play a central role in innate immunity, regulating blood clotting and complement activation in mammals, hemolymph clotting in horseshoe crabs, and melanization and Toll signaling pathway activation in insects (8–10). Their activation leads to proteolytic cleavage of a series of inactive protease zymogens between a prodomain and the protease domain, which then cleave downstream key immune factors, ensuring a fast, often within minutes, activation of antimicrobial immunity. In insects, these cascades consist of so-called clip-containing serine proteases (cSPs), a large protein family (consisting of 5 subfamilies, A-E), which is characterized by one or more clip pro-domains and one or more protease domains with trypsin-like specificity, and their homologs (cSPHs), which have lost one or more residues of the catalytic triad required for proteolytic activity. Our knowledge of the molecular makeup of these cascades is mainly derived from insect model species that are either large in size, enabling biochemical investigations of their hemolymph (e.g. *Manduca sexta* (11) and *Tenebrio molitor* (12)), or hold an excellent genetic and molecular tool box (in case of *Drosophila melanogaster* (13,14)).

At their core, canonical insect immune protease cascades consist of a modular serine protease (ModSP), which upon recognition of foreign molecular patterns, self-activates and then cleaves and activates a cSP in the subfamily C that in turn cleaves and activates a terminal cSP in the subfamily B (cSP-B) (15–17). Activation of melanization, an insect specific immune response proceeds through the terminal cSP-B pro-phenoloxidase activating protease (PAP) that cleaves the zymogen pro-phenoloxidase (proPO) to active phenoloxidase, a key enzyme in the conversion of aromatic amino acids to quinones that ultimately polymerize to eumelanin on foreign surfaces. Clip serine protease homologs support the proPO cascade, as they are required for effective

activation of PO on foreign surfaces (18–20). To prevent uncontrolled spread and over-activation
of responses, serine protease inhibitors in the serpin family tightly control mammalian and
arthropod immune protease cascades.

In *An. gambiae*, melanization is a readily selectable parasite resistance mechanism that
garnered much attention in hopes of exploiting the molecular underpinnings of this resistance
mechanism for malaria control purposes (21,22). In addition, the gene families encoding cSPs and
cSPHs in *An. gambiae*, known as CLIPs, and proPOs have undergone large expansions, raising
the question whether regulation of melanization is more complex than in other insects (23–26). To
date, the melanization response in *An. gambiae* has been analyzed genetically using RNAi (27)
combined with at least five experimental readouts, often with the implicit or explicit assumption
that identified CLIPs contribute biochemically to proPO activation. Thus far, two *An. gambiae*
cSPs, CLIPB9 and B10 have been identified as PAPs, based on their ability to revert increased
systemic melanization induced genetically by knockdown (kd) of *An. gambiae serpin-2* (*SRPN2*)
and to cleave purified *M. sexta* proPO *in vitro* (28,29). These data indicate that, based on the
canonical model of proPO activation cascades, melanization in *An. gambiae* depends on at least
two such cascades, characterized by CLIPB9 and B10, respectively. Based on genetic and
biochemical data, CLIPB8 is located further upstream of CLIPB9 (30). All three CLIPB proteases
required for proteolytic activation of proPO are inhibited by serpins (SRPNs). Of the 18 SRPNs
annotated in the *An. gambiae* genome, only SRPN2 has been shown to control systemic
melanization (7,31). It inhibits effectively both PAPs, CLIPB9 and B10, and also CLIPB8, albeit
to a lesser degree (28–30). Several additional CLIPBs, including CLIPB3, 4, 14 and 17 also
contribute to melanization; however the molecular function and placement of these additional
CLIPBs in proPO activation cascades is unknown (32,33). The only cSP in the CLIPC family to
contribute to melanization is CLIPC9, presumably through the action of cleavage of either PAP,
which however remains to be demonstrated (34).

In addition to cSPs, melanization in *An. gambiae* is regulated positively by a core module of
cSPHs, consisting, in hierarchical order, of SPCLIP1, CLIPA8 and CLIPA28. This module is
located genetically upstream of CLIPC9 and downstream of the thioester-containing protein TEP1
(6,32,34–36). Melanization is also regulated negatively by cSPHs, including CLIPA2 and A14
(32,37). While the precise mechanism(s) by which *An. gambiae* cSPHs regulate melanization
(positively or negatively) are unknown, observations in *An. gambiae* suggest that cSPHs contribute

to the opsonization of microbial surfaces by TEP1 and the leucine-rich repeat proteins LRIM1 and

APLC1 (35,38), and, based on data from other insect species, to PO activity by mediating proPO

cleavage by PAPs (19,39,40). In addition, melanization of rodent and human malaria parasites in

An. gambiae is inhibited by two c-type lectins, CTL4 and CTLMA2 (41,42), which presumably

bind to sugar moieties on the parasite surface and prevent melanization locally. The rodent malaria

parasite melanization phenotype induced by *CTL4* kd has been used successfully to identify some

of the cSPs and cSPHs mentioned above (e.g. (32)).

However, despite the significant number of studies summarized above, our understanding of

cSP cascades that control melanization in *An. gambiae* remains incomplete, and more than 80% of

the 110 annotated CLIPs await experimental exploration. To guide the experimental analysis of

An. gambiae cSPs and to identify putative key regulators of the melanization response, we describe

here a novel systems biology approach to study melanization in insects, combining transcriptomic

network and proteomic analyses. Using this approach, we identify the cSP CLIPB4 as a central

node in the protease network that regulates melanization in *An. gambiae* and characterize its

placement in the proPO activation cascade using a combination of genetic and biochemical

approaches.

Materials and Methods

Mosquito cultures

The *An. gambiae* G3 strain was obtained from the Malaria Research and Reference Reagent

Resource Center (MR4) at CDC and reared as described previously (29). For egg production,

females were provided a blood meal of commercial heparinized horse blood (Plasvacc USA) using

parafilm as a membrane stretched across glass feeders connected to a 37°C circulating water bath.

Anopheles gambiae *Melanization Gene Co-expression Network (AgMelGCN)*

The AgMelGCN was constructed according to (43). Genes of interest (GOIs), represented as nodes

were selected based on published evidence either confirming experimentally their role in

melanization or confirming phylogenetically their membership in protein families known to

contribute to melanization in mosquitoes. The GOI list contained 191 genes, including 9

prophenoloxidases (PPO), 14 thioester-containing proteins (TEP), 23 C-type lectins (CTL), 16

leucine-rich repeat immune proteins (LRIM), 18 serpins (SRPN), 4 putative modular serine

proteases (ModSP), 20 clip domain serine protease A family proteins (CLIPA), 29 CLIPB, 12

CLIPC, 13 CLIPD, 32 CLIPE, and SP24D, a putative ModSP. Their raw expression values were

collated from 257 unique physiological conditions assessed in 30 separate transcriptome studies,

and z-score conversion was performed for normalization under each condition. Two genes were

considered co-expressed and connected by edges if the Pearson Correlation coefficient (PCC) of

their expression vectors was above the 3% fitted sliding threshold. PCCs were rescaled based on

the reversed curve of the fitted sliding thresholds to assign edge weights. The network was

visualized using Gephi 0.9.2 (44) using the Fruchterman-Reingold force-directed algorithm (45).

Network communities were assigned with the Girvan-Newman algorithm (46). Four measures

were applied for node centrality, evaluating the edge abundance connected to a single node: node

strength (sum of connected edge weights), eigenvector centrality, betweenness centrality and

harmonic closeness centrality (47). In addition, s-core decomposition analysis was performed. In

the case of a weighted network, the s-core subgraph consists of all nodes i with node strengths $s(i) >$

s, where s is a threshold value. We define the threshold value of the s_n-core as $s_{(n-1)} = \min s(i)$,

among all nodes i belonging to the $s_{(n-1)}$-core network. The s_n-core is found by the iterative removal

of all nodes with strengths $s(i) \leq s_{(n-1)}$ (48).

SRPN2 Co-immunoprecipitation (Co-IP) and Mass Spectrometric Analysis

Female mosquitoes were challenged by microinjection of 69 nl lyophilized *Micrococcus luteus* in

water ($OD_{600} = 0.55$, Sigma-Adrich) per mosquito, and hemolymph was collected from 420 female

mosquitoes into 1x Roche Protease Inhibitor in PBS by proboscis clipping 19 h post infection. The

SRPN2 antiserum, produced previously (29), was affinity purified using the pH 7.2 coupling buffer

with AminoLink[TM] Plus Immobilization kit (ThermoFisher Scientific), and stored in PBS. Co-IP

was performed with Pierce™ Protein A Magnetic Beads (ThermoFisher Scientific), and beads

were rinsed with the buffer PBSOg (PBS with 0.7% w/v n-Octyl-BD- Glucopyranoside). Mosquito

hemolymph was equally split for two co-IP reactions: one with 40 µl 1:1 slurry of PBSOg-

containing magnetic beads cross-linked with 30 µg of purified SRPN2 antibody; and the other with

40 µl 1:1 slurry of PBSOg-containing control magnetic beads. Hemolymph was incubated with

the beads for 1.5 h at room temperature with constant rotation. The unbound fractions were

removed and magnetic beads were washed twice with PBSOg followed by four washes with

UltraPure[TM] water (Invitrogen). Bound proteins were eluted into 20 µl 0.1% TFA (w/v,

trifluoroacetic acid in water) in each co-IP. Samples were shipped to the Nevada Proteomics Center

(University of Nevada, Reno) for identification via LC-Orbitrap-MS. Contaminants were removed

with 2-D Clean-up kit (GE Healthcare), and in-tube trypsin digestion was performed. Resultant

peptides were run through a Michrom Paradigm Microdialysis Liquid Chromatography and

Michrom CaptiveSpray coupled to a Thermo LTQ Orbitrap XL with Electron-transfer dissociation

(ETD). MS spectra were analyzed using Sequest (IseNode in Proteome Discoverer version

2.2.0.388, Thermo Fisher Scientific). Sequest searches of ANOGA_uniprot_20210804.fasta

(25565 entries) were performed with the following settings: digestion enzyme trypsin, maximal

missed cleavage = 2, fragment ion mass tolerance = 0.60 Da, and a parent ion tolerance = 50 PPM.

Carbamidomethyl of cysteine was specified as a fixed modification, while oxidation of methionine,

acetylation of the N-terminus and phosphorylation of serine were specified as variable

modifications. Scaffold (version 5.1.0, Proteome Software Inc.) was used to validate peptide and

protein identifications. Peptide probability was calculated by the Peptide Prophet algorithm (49)

with Scaffold delta-mass correction, and a ≥95.0% identification cut-off. Protein probabilities were

assigned by the Protein Prophet algorithm (50). Proteins that contained similar peptides and could

not be differentiated based on MS analysis alone were grouped to satisfy the principles of

parsimony. In the SRPN2 Co-IP sample, a protein was considered present if protein probability

was greater than 99.0%, and 2 minimal unique peptides were identified in the sequence. A protein

was removed from the SRPN2 Co-IP list if at least one minimal peptide was detected in the control

sample. Proteome Discoverer SequestHT scores were listed at the protein level to rank the final

identification list.

Expression and purification of recombinant proteins proCLIPB4 and proCLIPB4$_{Xa}$

To produce recombinant proCLIPB4, its full coding region, including the signal peptide, was

amplified by PCR using gene specific primers (Table S1) and *An. gambiae* adult cDNA as template.

The forward primer contained a NotI site at 5'-end, and the reverse primer included three glycine

codons, six histidine codons, a stop codon, and a HindIII site at the 3'-end. The PCR product was

digested by NotI and HindIII, and inserted into the same restriction sites in the transfer vector

pFastBac1 (Invitrogen). The resulting proCLIPB4-6His-pFastBac1 plasmid was used as a template

to produce the mutant proCLIPB4$_{Xa}$-6His-pFastBac1.To utilize commercial Factor Xa (New

England Biolabs), the predicted activation site LTDR^{107} of proCLIPB4 was replaced with IEGR^{107}.

Site-directed mutagenesis was introduced to proCLIPB4-6His-pFastBac1 by Vent DNA

polymerase (New England Biolabs) with mutagenic primers (Table S1), DpnI (New England

Biolabs) digestion and NEB10β transformation. The resulting proCLIPB4$_{Xa}$-6His-pFastBac1 was

confirmed by Sanger Sequencing. The DNA segments containing proCLIPB4-6His and

proCLIPB4$_{Xa}$-6His, respectively, were transferred from pFastBac1 to pOET3 (Oxford Expression

Technologies) by BamHI and HindIII sites, creating proCLIPB4-6His-pOET3 and proCLIPB4$_{Xa}$-

6His-pOET3, respectively. The pFastBac1 and pOET3 transfer vectors were used to produce

corresponding CLIPB4 baculoviruses using the Sf9 cell line by means of Bac-to-Bac system

(Invitrogen) or flashBAC-Baculovirus Expression system (Genway Biotech). Large-scale

expression was performed using Sf9 suspension cultures at 2×10^6 cells/mL cell density in Sf-900

II serum-free medium (Invitrogen). Baculoviruses were inoculated at multiplicity of infection

(MOI) of 1, and expression media were harvested 48 h post infection. Recombinant proteins were

purified as described previously (28–30). In brief, cells were cleared by centrifugation, and

expression medium was dialyzed and subjected to column chromatography with Ni-NTA agarose

(Qiagen) and Q Sepharose Fast Flow (Cytiva). Eluted fractions were screened by immunoblot with

THE™ His Tag Antibody. Combined fractions were concentrated with buffer exchange (20 mM

Tris, 50 mM NaCl, pH 8.0) in Amicon™ Ultra-10K centrifugal units. Recombinant proteins were

stored at -80°C until future use. Recombinant proCLIPB8$_{Xa}$, proCLIPB8, proCLIPB9$_{Xa}$,

proCLIPB9 and SRPN2 were expressed and purified as described previously (29,30).

SDS-PAGE and Immunoblot

Protein samples were mixed with SDS sample buffer containing β-mercaptoethanol, and denatured

at 95°C for 5 min. Proteins were separated on SDS-PAGE gels (Bio-Rad) with constant voltage at

120V followed by 180V. SDS-PAGE gels were stained with EZ-Run Protein Gel Staining Solution

(Fisher) and de-stained with Milli-Q water. For immunoblot, proteins were transferred onto

nitrocellulose membranes (GE Healthcare) at 10V constant voltage using Owl™ Semidry

Electroblotting Systems (ThermoFisher Scientific). Membranes were incubated with 5% dry milk

in 1xTBST (0.05% Tween-20) at room temperature (RT) for 1 h, followed by primary antibody

incubation diluted in 0.5% dry milk in 1xTBST (4°C overnight or RT for 2 h). Blots were washed

three times with 1xTBST for 20 min each at RT. Membranes were further incubated with diluted

secondary antibodies in 0.5% dry milk 1xTBST at RT for 1 h. Protein signals were developed by

Alkaline Phosphate (AP) Conjugate Substrate Kit (Bio-Rad) and Amersham™ ECL™ Western

Blotting Detection Reagents (GE Healthcare) for AP-conjugated and HRP-conjugated secondary

antibodies, respectively. Images were captured by CCD camera of Azure Biosystems 300 for ECL

development.

Factor Xa activation of IEGR-mutated recombinant proteins

Factor Xa activation of recombinant proCLIPB4$_{Xa}$ is described in the legend of online suppl. Fig.

S3. Activation of proCLIPB8$_{Xa}$ and proCLIPB9$_{Xa}$ was performed as described previously (29,30).

SRPN2 inhibition of CLIPB4

To produce active CLIPB4, 630 ng of recombinant proCLIPB4$_{Xa}$ was incubated with 1 µg of

Factor Xa in a total volume of 11 µl with Factor Xa activation buffer at 37℃ overnight. Two

negative controls were set up in parallel, where proCLIPB4$_{Xa}$ or Factor Xa was replaced with the

same volume of Factor Xa activation buffer. Activated CLIPB4$_{Xa}$ (154 ng/per reaction) was mixed

with recombinant (r)SRPN2 at molar ratios of 1:1 or 1:10 (proCLIPB4$_{Xa}$:SRPN2). After incubation

at room temperature for 15 min, the reaction mixtures were subjected to immunoblot analysis with

mouse anti-His, rabbit anti-SRPN2 or rabbit anti-CLIPB4 as primary antibodies. For Mass

spectrometry identification, reactions were set up as described as above, and subjected to SDS-

PAGE. After Coomassie brilliant blue staining, the band of interest (~72 kDa) was excised. In-gel

trypsin digestion and Electrospray ionization mass spectrometry (ESI-MS, Bruker Daltonics HCT

Ultra) were performed at the Biotechnology/Proteomics Core Facility (Kansas State University).

MS results were analyzed in Scaffold 4.0. To determine the stoichiometry of inhibition, 90 ng of

Factor Xa-activated rCLIPB4$_{Xa}$ was mixed with rSRPN2 at molar ratios of 0:1, 0.5:1, 1:1, 2:1, 4:1,

and 6:1 (rSRPN2:rCLIPB4$_{Xa}$) in reaction buffer 20 mM Tris, 150 mM NaCl, pH 8.0 supplemented

with 3 µg BSA. Reactions were incubated at room temperature for 15 min. The residual amidase

activity was determined with IEARpNA substrate at OD$_{405}$ as described above. IEARase activity

of Factor Xa was measured in parallel and subtracted from the above reactions of

rSRPN2:rCLIPB4$_{Xa}$. Amidase activity at molar ratio of 0:1 (rSRPN2:rCLIPB4$_{Xa}$) was defined as

100%. All assays were performed in triplicate.

Purification of PPO1&2 from M. sexta larval hemolymph

Prophenoloxidase 1&2 (*Ms*PPO1&2) was purified from *M. sexta* hemolymph similarly to

published methodology (51). Briefly, hemolymph was collected into 100% ammonium sulfate (AS)

at 0℃ from 5th instar Day 2 *M. sexta* larvae. Fraction of 38-48% AS precipitation was dissolved in

buffer A (10 mM KPi, pH 6.8, 500 mM NaCl, 0.5 mM reduced L-Glutathione), and dialyzed

against buffer A. Bio-Gel HT Hydroxyapatite column (Bio-Rad) was utilized to separate proteins

via a linear gradient of 10-100 mM KPi in buffer A. Two microliters per fraction was assayed with

200 µl 2 mM Dopamine hydrochloride in 50 mM NaPi, pH 6.5 with and without 0.1%

cetylpyridinium chloride (CPC, w/v) in 96-well plates, and absorbance change was monitored at

470 nm in EPOCH 2 microplate reader (BioTek). Fractions containing CPC-activated PO activity

were combined, and concentrations of $MgCl_2$ and $CaCl_2$ were adjusted to 1 mM each. The protein

solution was applied twice through a Concanavalin A Sepharose 4B column (GE Healthcare),

equilibrated in 20 mM Tris, pH 7.4, 0.5 M NaCl, 1 mM $MgCl_2$, 1 mM $CaCl_2$. Column flow-through

fraction was adjusted to 1 M AS, and loaded onto a Phenyl Sepharose 6 Fast Flow low sub column

(Cytiva), equilibrated in 0.1 M KPi, pH 7.1, 1 M AS. Elution was performed with a descending

linear gradient of 0.1 M KPi, pH 7.1, 1 M AS to 0.01 M KPi, pH 7.1. Fractions with CPC-activated

PO activity were pooled and concentrated to 1.1 ml using Amicon^{TM} Ultra-15 30K centrifugal

units (Merck Millipore). Proteins were separated by Sephacryl S-300 column (Cytiva),

equilibrated in 10 mM MOPS, pH 7.2, 0.5 M NaCl. Fractions containing CPC-activated PO

activity were pooled and concentrated to 4.6 mg/ml by Amicon^{TM} Ultra-15 30K units. Equal

volume of glycerol was mixed with purified *Ms*PPO1&2, and the purified zymogen was stored at

-80°C at 2.3 mg/ml in 5 mM MOPS, pH 7.2, 0.25 M NaCl, 50% glycerol.

ProPO activation by CLIPB4$_{Xa}$ using M. sexta plasma and purified MsPPO1&2

To evaluate the contribution of CLIPB4 on proPO activity, *M. sexta* plasma and purified

*Ms*PPO1&2 were utilized as described previously (29). Naïve hemolymph was collected from a

cut proleg of 5th instar Day 3 *M. sexta* larvae, and centrifuged at 5,000 g, 4°C for 5 min to remove

hemocytes and cell debris. Factor Xa was used to activate proCLIPB4$_{Xa}$ as described in online

suppl. Fig. S3. For PO enzymatic activity plate assay, in each well, 1 µl plasma was mixed with 3

µl Factor Xa-proCLIPB4$_{Xa}$ activation mixture per well, supplemented to 10 µl with buffer B (20

mM Tris, 20 mM NaCl, 6.7 mM $CaCl_2$, pH 7.5). Three negative controls were set up, replacing

Factor Xa, proCLIPB4$_{Xa}$, and both with buffer B, respectively for a final reaction volume of 10 µl.

Three technical replicates were set up per reaction, and incubated at RT for 15 min. Per well, 200

µl 2 mM Dopamine hydrochloride was added in the buffer 50 mM NaPi, pH 6.5. Absorbance was

monitored at 470 nm at RT for 1 h. For PO activity assay with purified MsPPO1&2, per well, 1

µg purified *Ms*PPO1&2 (4 µl at 250 ng/µl) was mixed with 5 µl of activation mixture and 3 µl of

buffer B for a total reaction volume of 12 µl. One additional negative control was included by

mixing 1 µg purified *Ms*PPO1&2 with 8 µl buffer B. Three technical replicates were set up for

each reaction and incubated at RT for 1 h. Dopamine hydrochloride was added as described above

and absorbance was monitored at 470 nm at RT for 1 h. One unit of PO activity was defined as

$\Delta A470/min = 0.001$.

ProPO cleavage by CLIPB4$_{Xa}$ using M. sexta plasma and purified MsPPO1&2

To detect cleavage of *Ms*PPO1&2, we performed immunoblots with rabbit anti-*Ms*PPO1&2

primary antibody (51). Naïve *M. sexta* plasma was first diluted to 1:20 in 20 mM Tris, 150 mM

NaCl, pH 8.0. Reactions were set up by mixing 1 µl diluted plasma, 3 µl Factor Xa activation

buffer or 3 µl each from Factor Xa proCLIPB4$_{Xa}$ RT activation mixtures (including negative

controls), and 6 µl buffer B for a 10 µl total reaction volume. Three additional samples were

prepared by mixing 3 µl each from Factor Xa proCLIPB4$_{Xa}$ RT activation mixtures and 7 µl buffer

B. Reactions were incubated at RT for 20 min. For purified *Ms*PPO1&2, reactions were set up by

mixing 2 µl purified *Ms*PPO1&2 (250 ng/µl, 500 ng) with 6 µl Factor Xa activation buffer or 6 µl

each from Factor Xa proCLIPB4$_{Xa}$ activation mixtures (see online suppl. Fig. 3). Reactions were

incubated at RT for 20 min. For immunoblot, 1.2 µl of each reaction (containing 75 ng of

*Ms*PPO1&2) was mixed with 8.8 µl water, and boiled at 95°C for 5 min with SDS sample buffer.

Samples were separated on 7.5% SDS-PAGE gels, and probed with *Ms*PPO1&2 antibody. In

addition, CLIPB9$_{Xa}$-*Ms*PPO1&2 reactions were included as a positive control. Briefly, to obtain

activated CLIPB9$_{Xa}$, 2.5 µg of purified recombinant proCLIPB9$_{Xa}$ was mixed with 1 µg Factor Xa,

and Factor Xa activation buffer in a total reaction volume of 25 µl, and incubated at 37°C overnight.

Two negative control reactions were set up and incubated in parallel, in which recombinant

proCLIPB9$_{Xa}$ and Factor Xa were replaced with the same volume of Factor Xa activation buffer,

respectively.

CLIPB4 activation of proCLIPB8 and proCLIPB9

To test whether CLIPB4 cleaves recombinant proCLIPB8 and/or proCLIPB9, proCLIPB4$_{Xa}$ was

activated by Factor Xa at 37°C. 154 ng of Factor Xa-activated CLIPB4$_{Xa}$ was incubated with 100

ng proCLIPB8 or 100 ng proCLIPB9 at RT for 15 min. Recombinant proCLIPB8$_{Xa}$ or

proCLIPB9$_{Xa}$ was activated by Factor Xa similarly as described previously (29,30), and 100 ng of

activated CLIPB8$_{Xa}$ or CLIPB9$_{Xa}$ was used as a positive control. Samples were subjected to SDS-

PAGE and immunoblot using mouse anti-His, rabbit anti-CLIPB8, or rabbit anti-CLIPB9

antibodies. To measure the enzymatic activity of cleaved CLIPB8, 610 ng of recombinant

proCLIPB4$_{Xa}$ (102 ng/µl) was incubated at 37 °C for 10 h with 1 µg (1 µg/µl) Factor Xa,

supplemented to 21 µl with Factor Xa activation buffer. 72 ng of activated CLIPB4$_{Xa}$ was

incubated with 450 ng proCLIPB8 (225 ng/µl) at room temperature for 1 h. Reactions were

supplemented with 200 µl of 100 µM N-benzoyl-Phe-Val-Arg-p-nitroanilide (FVR-pNA, EMD

Chemicals) in 0.1 M Tris-HCl, pH 8.0, 0.1 M NaCl, 5 mM CaCl$_2$ and amidase activity was

measured continuously at OD$_{405}$ for 30 min. 450 ng of activated CLIPB8$_{Xa}$ was used as a positive

control. One unit of CLIPB8 activity was defined as $\Delta A405/min = 0.001$. All activity assays were

performed in triplicate.

Double-stranded RNA (dsRNA) synthesis and RNAi experiments

DsRNA synthesis and adult female mosquito injections were performed as described previously

(29). Primers are listed in Table S1 for T7-tagged gene-specific amplification of *CLIPB4*. Primers

were adapted from previous studies for dsRNA synthesis of *SRPN2*, *GFP*, *CLIPB8*, and *LacZ*

(29,30,52). In single gene knockdown (kd) experiments, 3 to 4-day old adult female mosquitoes

were injected each with 207 ng of dsRNA in a total volume of 69 nl. For kd control, mosquitoes

were injected with the same quantity of the non-related dsRNA of either *Green Fluorescent*

Protein (ds*GFP*) or *β-galactosidase* (ds*LacZ*). For double knockdowns (dkd), mosquitoes were

injected with 138 nL of a 1:1 dsRNA mixture containing 1.5 µg/µl of each dsRNA. For dkd

controls, ds*GFP* was added to compensate the total dsRNA dose to 414 ng/mosquito.

Melanization phenotype analysis

To assess CLIPB4's impact on melanization, two assays were utilized, Melanization-associated

Spot Assay (MelASA) and SRPN2 depletion-induced melanotic pseudotumor formation. To

quantify melanotic excreta, MelASA was performed in the standard format (50

mosquitoes/experimental cup) as described previously (34), with or without bacterial challenge.

For MelASAs after bacterial challenge, dsRNA-injected mosquitoes were allowed to recover for

60 h and injected each with 50.6 nl of 1x PBS, OD$_{600}$=1.5 *Escherichia coli* (resuspended in 1x

PBS, same below), OD$_{600}$=1.5 *Staphylococcus aureus*, and OD$_{600}$= 0.4 *M. luteus* (*Micrococcus*

lysodeikticus lyophilized cells ATCC, No. 4698, Sigma - Aldrich), respectively. Clean P8-grade

white filter papers (Fisher) were inserted into the bottom of the experimental cups after bacterial

challenge and removed 12 h later. For MelASAs without bacterial challenge, clean filter papers

were inserted immediately after dsRNA injection, and removed 60 h later. All MelASAs were

performed with five independent biological replicates using different mosquito generations.

Collected filters papers were imaged under white epi-illumination without filters in an Alpha

Imager System, and analyzed by Fiji is Just Image J (FIJI, Version 2.1.0) (53) using custom FIJI

macros. To quantify melanotic pseudotumor formation, mosquitoes were dissected when mortality

reached 70% in the ds*SRPN2*/ds*GFP* group. Abdominal walls were prepared as described

previously (7) and examined under 40× magnification with Axio Imager A1 microscope (Zeiss)

equipped with AxioCam MR5 (Zeiss). Image processing, and abdomen melanized area calculation

were performed in ImageJ software as described previously (29,54). Five independent biological

replicates were performed using 20 mosquitoes per treatment. All statistical analyses were

performed using GraphPad Prism (ver. 9.5.1, GraphPad Software).

Results

Co-expression analysis of putative melanization genes identifies CLIPB4 as a central node

In the absence of a genome-wide protein-protein interaction network for *An. gambiae*, we used

gene co-expression as a proxy to identify proteins that are likely to inhabit the same space at the

same time. To facilitate the identification of possible protein-protein interactions, and specifically

SRPN2-cSP interactions controlling melanization, we therefore constructed an undirected gene

co-expression network of all *An. gambiae* genes belonging to gene families experimentally shown

to contribute to melanization, which we named the *Anopheles gambiae* <u>Mel</u>anization <u>G</u>ene <u>C</u>o-

expression <u>N</u>etwork (AgMelGCN) (Fig. 1a). In total, the AgMelGCN contains 178 nodes (genes)

and 1229 edges (co-expression links between gene pairs) (online suppl. Tables S2 and S3). To

reveal the central nodes of AgMelGCN, we performed an *s*-core analysis (*s*=11.14). The core

contained 25 genes, of which nine play a role in melanization (APL1C, CLIPA2, A8, B4, B10,

CLIPC9, CTL4, CTLMA2, LRIM1; online suppl. Table S2). To identify key cSPs in the network,

we calculated two centrality measures, node strength (weight sum of all node's edges) and

harmonic closeness centrality (sum of shortest paths to all other nodes within the full network)

(47). Of all annotated cSPs, CLIPB4 was the highest ranked node in either centrality measure

(online suppl. Table S2, Fig. 1b). In AgMelGCN, CLIPB4 was connected by a weighted edge to

44 other nodes, including seven SRPNs; notably, the SRPN2-CLIPB4 edge had the highest weight

among all SRPN nodes connected to CLIPB4 (Fig. 1b; online suppl. Table S3). SRPN2 was

connected to 18 nodes, including five cSPs, all in the CLIPB sub-family (Fig.1c; online suppl.

Table S3), of which the SRPN2-CLIPB4 edge possessed the highest weight. Overall, these results

suggested that CLIPB4 is a central node in the AgMelGCN whose expression most closely

resembled that of SRPN2.

SRPN2 and CLIPB4 interact in adult mosquito hemolymph

Since SRPN2 is a critical factor affecting the melanization process, we aimed to identify its closely

associated proteins in the hemolymph of adult mosquitoes. Using co-immunoprecipitation, we

identified 21 proteins bound directly or indirectly to SRPN2 in the hemolymph of adult female

mosquitoes 24 h post challenge with lyophilized *M. luteus* bacteria (Table 1). Among these, six

genes were co-expressed with SRPN2 in AgMelGCN, including TEP15, CLIPB4, CLIPA14,

SRPN11, CLIPA9 and CLIPA6 (Fig. 1c, Table 1 and online suppl. Table S3). Among annotated

cSPs and cSPHs, CLIPB4 was ranked the highest in SRPN2 Co-IP MS protein scores, warranting

further examination of the interaction between SRPN2 and CLIPB4.

CLIPB4 is a serine protease with trypsin-like specificity

To investigate its biochemical function(s), we expressed recombinant *An. gambiae* proCLIPB4

zymogen *in vitro*. The determinants of enzyme specificity were predicted to be D^{305}, G^{332} and G^{343}

(online suppl. Fig. S1a), suggesting that CLIPB4 is a trypsin-like protease that cleaves its substrate

after arginine or lysine residues (55). As the endogenous activating enzyme of CLIPB4 is unknown,

a mutant (proCLIPB4$_{Xa}$) was produced by replacing its putative activation site ^{104}LTDR^{107} (56)

with ^{104}IEGR^{107} (online suppl. Fig. S1b), enabling recognition and cleavage by commercially

available bovine Factor Xa. We produced and purified recombinant proCLIPB4$_{Xa}$ zymogen

(rproCLIPB4$_{Xa}$) tagged with 6His at its C-terminus (Fig. S2a). Coomassie stain, His antibody and

anti-CLIPB4 antibody detected rproCLIPB4$_{Xa}$ (37.9 kDa) at the expected molecular weight.

Recombinant proCLIPB4$_{Xa}$ was cleaved by Factor Xa after incubation at 37°C and RT, generating

a fragment corresponding to the expected size of the CLIPB4 catalytic domain (rCLIPB4$_{Xa}$, Fig.

S3a and c). Recombinant CLIPB4$_{Xa}$ exhibited amidase activity, using IEAR*p*NA substrate, while

rproCLIPB4$_{Xa}$ did not (Fig. S3b and S3d). As noted in previous works, Factor Xa also utilized

IEAR*p*NA as a substrate (28–30). These results together support that, consistent with our construct

design, Factor Xa is able to cleave and activate rCLIPB4$_{Xa}$-6His.

SRPN2 irreversibly inhibits CLIPB4 in vitro

Based on the AgMelGCN and co-IP results, we hypothesized that SRPN2 is a direct inhibitor of

CLIPB4. To test this hypothesis, we determined (i) if SRPN2 and CLIPB4 form inhibitory protein

complexes and (ii) if SRPN2 decreases CLIPB4 amidase activity in a concentration-dependent

manner. Purified rSRPN2, tagged N-terminally with 6-His tag, was incubated with rCLIPB4$_{Xa}$ at
room temperature for 15 min at two different molar ratios, 1:1 or 10:1 (rSRPN2: rCLIPB4$_{Xa}$). In
SDS-PAGE and immunoblot analyses of these reactions a higher molecular weight band appeared
at 72 kDa, which was detected with anti-His, anti-SRPN2 and anti-CLIPB4 antibodies (Fig. 2A,
2B and 2C). ESI-MS analysis on the 72 kDa band (online suppl. Table S4) detected 15 and 2
peptides for SRPN2 and CLIPB4, respectively, corresponding to 11 and 2 unique peptides for each,
confirming that SRPN2 and CLIPB4 form a SDS-stable protein complex. CLIPB4$_{Xa}$ amidase
activity against IEARpNA decreased with increasing amount of rSRPN2, with a stoichiometry of
inhibition (SI) of 1.4 (Fig. 2d), demonstrating that SRPN2 functions as an efficient inhibitor of
CLIPB4.

CLIPB4kd *is required for the phenotypes induced by SRPN2 depletion*
Previously, *SRPN2*kd was shown to reduce the longevity of female mosquitoes and to induce
melanotic pseudotumor formation (7). Our biochemical data support that SRPN2 directly inhibits
the activity of CLIPB4, and we hypothesized that CLIPB4 is required for the phenotypes induced
by SRPN2 depletion. To test this hypothesis, we performed genetic epistasis experiments, inducing
gene knockdown by long dsRNA injection. Adult female *An. gambiae* were injected with dsRNAs
targeting *CLIPB4*, *SRPN2*, and both *CLIPB4* and *SRPN2*, respectively. Injection of *dsGFP* was
used as a control. *CLIPB4* knockdown was significant and specific as determined by immunoblot
(online suppl. Fig. S4). Injection of ds*SRPN2* did not affect CLIPB4 protein levels.

We first evaluated the effect of *CLIPB4*kd on survival comparing the survival curves between
the four different treatment groups, and we also monitored the survival of uninjected mosquitoes.
*SRPN2*kd significantly shortened longevity as compared to the uninjected controls and ds*GFP*-
control treatment (Fig. 3a and b). In contrast, *CLIPB4*kd had no effect on lifespan as compared to
the controls. Importantly, ds*CLIPB4/SRPN2*-treatment substantially and significantly increased
mosquito longevity as compared to *SRPN2*kd mosquitoes, returning longevity to those of
uninjected and ds*GFP*-control treated mosquitoes.

We then evaluated the effect of *CLIPB4*kd on melanization, using the appearance of melanotic
tumors as a read-out (Fig. 3c). Injection of ds*GFP* or ds*CLIPB4* alone did not cause melanization
(data not shown). In contrast, *CLIPB4*kd in a *SRPN2*-depleted background, significantly affected
the *SRPN2*kd melanization phenotype, reducing the average total melanotic area for each dissected
mosquito abdomen by 54 %, however, not completely reverting the SRPN2 depletion phenotype.

We also used the recently established melanization-associated spot assay (MelASA) to evaluate

the contribution of CLIPB4 on the SRPN2 depletion phenotype (34). The standard format (50

mosquitoes/cup) was adopted and melanotic excreta were recorded for the four treatment groups.

*CLIPB4*kd alone exhibited similar melanotic excreta area compared to that of the negative control,

ds*GFP*. SRPN2 depletion, however, significantly increased melanotic excreta of female

mosquitoes. Double knockdown of *SRPN2* and *CLIPB4* significantly reduced the excreta

compared to *SRPN2* single knockdown, but did not completely abolish the SRPN2 depletion

phenotype (Fig. 3d). Taken together, the silencing experiments strongly suggest that *CLIPB4* is

epistatic to *SRPN2* and therefore downstream of this serpin in the proPO activation cascade.

CLIPB4 is required for melanization after microbial challenge

Since CLIPB4 is required for melanization induced by SRPN2 depletion, we hypothesized that

CLIPB4 is broadly required for melanization in antimicrobial immunity. To test the hypothesis,

we first conducted MelASAs after challenge with bacteria (*S. aureus*, *M. luteus*, and *E. coli*)

(online suppl. Fig. S5). Without bacterial challenge, only small amounts of melanotic excreta were

observed in the ds*GFP*-injected and *CLIPB4*kd treatment groups, consistent with our previous

observation in Fig. 3d. Each bacterial challenge increased significantly melanotic excreta in the

ds*GFP*-injected group as compared to ds*GFP*-injected mosquitoes treated with PBS, supporting

that bacterial treatment indeed induces the melanization immune response. The induction of

melanotic excreta by bacterial challenge was dampened in the *CLIPB4*kd treatment groups.

Independent of bacterial species, melanotic excreta were reduced by about 50% in *CLIPB4*kd

mosquitoes as compared to ds*GFP*-injected mosquitoes, demonstrating that melanization in

response to bacterial infection is broadly dependent on CLIPB4.

CLIPB4 functions as a proPO activating enzyme

The shortage of hemolymph collectable from *An. gambiae* mosquitoes can be overcome by using

the *M. sexta* model system as a source of proPO, as *M. sexta* PPO1&2 share the conserved cleavage

site with eight of the nine proPOs encoded in the *An. gambiae* genome (7). Activated CLIPB4$_{Xa}$

was incubated with *M. sexta* plasma, followed by immunoblot analysis using anti-*M. sexta* proPO

antibody. A doublet band around 80 kDa in *M. sexta* plasma represents heterodimeric proPO

consisting of 79-kDa proPO1 and 80-kDa proPO2 (51). Addition of activated CLIPB4$_{Xa}$ to *M.*

sexta plasma resulted in the appearance of a 70-kDa doublet band corresponding to *M. sexta* active

PO (Fig. 4a). The same doublet band was observed in the plasma supplemented with activated

CLIPB9$_{Xa}$, which we identified previously as a functional PAP in *An. gambiae* (29). PO activity

of plasma increased in the presence of active CLIPB4$_{Xa}$ (Fig. 4c). These results confirm that the

proteolytic activity of CLIPB4 promotes proPO cleavage and PO activity. To determine the

placement of CLIPB4 in the proPO activation cascade in *An. gambiae*, we next tested whether

CLIPB4 can directly cleave and activate proPO. ProPO1&2 was purified from 5th instar *M. sexta*

larvae (Fig. S2c). By Coomassie Blue staining and immunoblot analysis with MsPPO antibody, a

dimer was observed around 80 kDa, consistent with previous observations of purified full length

M. sexta proPO1 and 2 (51). We then incubated Factor Xa-activated rCLIPB4$_{Xa}$ with purified *M.*

sexta proPO, using again rCLIPB9$_{Xa}$ as a positive control. Neither rproCLIPB9$_{Xa}$ zymogen nor

Factor Xa alone were able to cleave MsPPO1&2, while active rCLIPB9$_{Xa}$ cleaved purified

MsPPO1&2 to produce the PO1&2 band at 70 kDa (Fig. 4b). Similarly, neither rproCLIPB4$_{Xa}$

zymogen nor Factor Xa alone cleaved PPO1&2, but active rCLIPB4$_{Xa}$ cleaved proPO1&2 (Fig.

4b), leading to an increase in PO activity (Fig. 4d), demonstrating that CLIPB4 functions as a PAP.

CLIPB4 cleaves and activates CLIPB8

We next explored whether CLIPB4 also promotes PO activity indirectly by activating the PAP

zymogen proCLIPB9 (29) and/or proCLIPB8, a protease upstream of CLIPB9 in the melanization

cascade in *An. gambiae* (30). To test whether CLIPB4 can cleave proCLIPB8 and/or B9 *in vitro*,

active CLIPB4$_{Xa}$ was incubated with wild-type recombinant proCLIPB8 and proCLIPB9,

respectively, followed by immunoblot analysis using either anti-His antibody or anti-CLIPB8 and

anti-CLIPB9 antibodies, respectively. In the reaction of active CLIPB4$_{Xa}$ with proCLIPB8

zymogen, anti-His and anti-CLIPB8 antibodies detected a 36 kDa band, corresponding to the

CLIPB8 protease domain, which was also detected in the control reaction of proCLIPB8$_{Xa}$

incubated with Factor Xa (Fig. 5a and b). In contrast, addition of active CLIPB4$_{Xa}$ (or Factor Xa)

did not result in the cleavage of proCLIPB9 (Fig. 5c and d), suggesting that CLIPB4 is not directly

upstream of CLIPB9. To evaluate whether cleavage of recombinant proCLIPB8 by CLIPB4$_{Xa}$

results in CLIPB8 activation, we measured the amidase activity of the reaction mixtures using

FVR as a substrate (30). Each of the five control reactions, containing proCLIPB8, proCLIPB8$_{Xa}$,

Factor Xa, proCLIPB4$_{Xa}$, and proCLIPB4$_{Xa}$/proCLIPB8, respectively did not exhibit amidase

activity. Active CLIPB4$_{Xa}$ showed limited amidase activity, which was increased threefold when

active CLIPB4$_{Xa}$ was incubated with proCLIPB8 prior to the addition of substrate (Fig. 5e).

Together with the immunoblot results (Fig. 5a and 5b), FVRase activity assay supports that
CLIPB4 specifically cleaves and activates CLIPB8.
To examine this phenomenon *in vivo*, ds*LacZ*, ds*CLIPB4* and ds*CLIPB8* were injected in adult
female mosquitoes that were challenged by *S. aureus* post the dsRNA injection recovery.
Hemolymph was collected 10 h after challenge and subjected to SDS-PAGE and immunoblot
detection (Fig. 6). The Apolipophorin-II (APOII) was utilized as a loading control. CLIPB8 was
depleted with ds*CLIPB8* treatment, serving as a positive control for CLIPB8 antibody detection.
Ten hours after bacterial challenge, a clear reduction of CLIPB8 was observed in ds*LacZ* compared
to the naïve control in two independent experiments, suggesting that CLIPB8 in the hemolymph
was rapidly utilized in the acute immune response (57). The CLIPB8 reduction, mediated by *S.*
aureus challenge, was rescued with ds*CLIPB4* treatment in two independent experiments,
providing additional support that CLIPB4 acts upstream of CLIPB8 in the hierarchical cSP cascade
regulating melanization in *An. gambiae*.
539

Discussion
Mosquitoes interact with a broad range of organisms across the symbiosis continuum and deploy
melanization as a broad-spectrum immune defense against many organisms they encounter (58).
The melanization of microbial surfaces in *An. gambiae* requires opsonization through TEP1 and
the activity of PO to produce eumelanin, which depends on the action of cascades made up of cSPs
and cSPHs. This study identifies CLIPB4 as an important factor required for melanization in
response to microbial infection. CLIPB4 was previously shown to contribute to melanization of
glass beads and malaria parasites in *CTL4* kd mosquitoes (32,33). Here we show that CLIPB4 is
also required for melanotic tumor formation exerted by the depletion of SRPN2 in adult female
mosquitoes, demonstrating that its contribution to melanization is not limited to specific surfaces,
and suggesting that CLIPB4 is fundamental to eumelanin production. Indeed, our data demonstrate
that CLIPB4 functions as a PAP, mediating proPO activation *ex vivo* and *in vitro*. In addition to
CLIPB9 and B10, CLIPB4 is the third PAP identified in *An. gambiae* (28,29). In all three cases,
their ability to cleave and activate proPO *in vitro*, was demonstrated using purified proPO1 and 2
from the hemolymph of *M. sexta*. Given that *An. gambiae* PPO5 and 6, the most abundant proPOs
in the hemolymph of adult female *An. gambiae*, share the same activation cleavage site with *M.*
sexta proPO1&2, it is highly likely that all three CLIPBs are bona fide PAPs in *An. gambiae* (7,59).

The existence of three PAPs, each constituting a terminal protease in a cSP cascade, suggests that melanization in adult female *An. gambiae* is regulated by at least three proPO activation cleavage cascades. Similarly, *M. sexta* proPO activation is regulated by at least two proPO activation cascades, utilizing either PAP3 or PAP1/PAP2 as the terminal proteases to cleave proPO (60,61). Likewise, two distinct proPO activation cascades regulate melanization in the lepidopteran species, *Helicoverpa amigera* (15,62). In *An. gambiae*, possibly not all three PAPs contribute equally to activate proPO. Expression of CLIPB4 in adult female *An. gambiae*, as measured by RNA-Seq is much higher than that of CLIPB9, B10, and any other CLIPB whose kd causes a melanization phenotype (Buchon and Michel, unpublished), suggesting that CLIPB4 protein, based on its abundance, contributes more to proPO activation than other PAPs. This observation may also explain why neither CLIPB8, B9 nor B10 were identified in the SRPN2 co-IP analyses, despite previous data demonstrating that CLIPB9 and SRPN2 form inhibitory complexes in *An. gambiae* hemolymph (29). Whether, in addition to protein concentration, the activation of the three *An. gambiae* PAPs depends on the type of microbial challenge, as observed in *H. amigera* (62), is currently under investigation.

In addition to its PAP function, CLIPB4's central role in melanization in female *An. gambiae* is further manifested by its ability to cleave and activate proCLIPB8. This function was demonstrated *in vitro* by biochemical studies and *in vivo*, as *CLIPB4* kd reverts the consumption of CLIPB8 in the hemolymph of adult female *An. gambiae* upon microbial infection. This is of not only the first observation of multiple substrates for a single *An. gambiae* CLIP protease, but also the first demonstration of a CLIPB protease activating another proCLIPB zymogen in mosquitoes, breaking the mold of the classical hierarchy of CLIPC-CLIPB in insect immune protease cascades (63). The observation of CLIPB4 cleavage of CLIPB8 is paralleled in *M. sexta*, where PAP3 cleaves proHP5, a cSP-B family member (64). This cleavage of HP5, provides positive reinforcement of proPO activation, as HP5 in turn cleaves HP6, a cSP in the C family, upstream of PAP-1, linking the two proPO activation cascades in *M. sexta* (64). Based on these similar patterns in two evolutionarily distant insect species, it is tempting to speculate that positive reinforcement of proPO activation by cSP-B mediated activation of other cSP-Bs is a common theme in the melanization immune response.

Given that the regulation of melanization through proPO activation requires cSPs and cSPHs circulating in mosquito hemolymph prior to immune challenge, we posited that identification of

important members in the cSP and cSPH cascades could be identified based on their co-expression.

Indeed, the topology of the AgMelGCN, the co-expression network of members of gene families

known to contribute to melanization clustered genes with known immune phenotypes into a single

community. Using network science approaches, we then identified highly-connected nodes,

referred to as hubs, which are integral to scale-free network stability (65). In gene co-expression

networks, hub genes and are often relevant for network functionality, either as regulators or

because they encode proteins with essential functions (66,67). The nodes with highest connectivity

in the AgMelGCN encoded two cSPHs (CLIPA9 and 8), APLC1, and CLIPB4. All four genes

encode proteins with known functions in innate immunity. *CLIPA9* kd in *An. gambiae* reduces *P.*

falciparum oocysts, a phenotype that seems dependent on the presence of an unperturbed

microbiota through an unknown mechanism (68). CLIPA8 is a key positive regulator of

melanization and part of the cSPH cascade upstream of CLIPC9 (34,36). CLIPA8 is required for

melanization of malaria parasites after *CTL4* kd, filamentous fungi, and bacteria (32,69,70). The

leucine-rich repeat protein APLC1 contributes to antimicrobial immunity in mosquitoes by

stabilizing TEP1 in the hemolymph, preventing its premature activation, and facilitating its

delivery to microbial surfaces for opsonization (38,71,72). Likewise, CLIPB4 plays an important

role in melanization through its integral roles in the cSP cascade that activates proPO described in

this current study. Combined, these data support that gene co-expression networks can indeed be

used to identify genes with important functions in mosquito innate immunity, and provide a

blueprint for analyses of innate immunity in other insect species and likely other complex

physiologies in mosquitoes.

The topology of the AgMelGCN also demonstrates the integration of the cSPH and cSP

cascades through co-expression of members of each cascade. Our co-IP data indicate that these

cascades also at least partially form protein complexes in the hemolymph of *An. gambiae*. We

recently showed that CLIPB4 also cleaves CLIPA8 *in vitro*, which is required for CLIPA8 function

(57), suggesting that the activation of cSPHs required for melanization is mediated by cSPs and

potentially integrated with proPO activation through the formation of higher molecular weight

protein complexes. Indeed, higher molecular weight complexes between cSPs and cSPHs and

proPO have been described in other insect species, including *M. sexta* (73) and *Bombyx mori* (74),

indicating that the formation of such immune complexes is likely an evolutionarily conserved

process that is used to optimize proPO activity to microbial surfaces.

Based on our results presented herein and published data, we propose the following model of

proPO activation in the melanization response of *An. gambiae* (Fig. S6). In adult female

mosquitoes, circulating proPO zymogen is activated through the proteolytic activity of three

CLIPB proteases, CLIPB4, B9 and B10 (28,29). CLIPB4 also activates proCLIPB8, which

activates indirectly proCLIPB9 (30), providing positive reinforcement of proPO activation. Each

of these CLIPBs provides an intervention point, as their activity is inhibited directly by SRPN2,

reinforcing its role as the main negative regulator of melanization and survival in *An. gambiae*

(7,31). This observation is paralleled in *M. sexta*, where serpin-3, the ortholog of SRPN2 inhibits

all three identified PAPs (75,76). CLIPC9, currently the only known CLIPC protease required for

melanization, is located likely upstream of one or more of the three PAPs and responsible for their

proteolytic activation. In turn, based on the canonical proPO activation cascades in other insects,

we hypothesize that proCLIPC9 is activated proteolytically through the action of an unknown

ModSP. Additionally, melanization of microbial surfaces is further regulated through the activity

of a cSPH cascade upstream of CLIPC9 (34,36). Whether, cSPHs also support melanization

downstream of PAPs, as observed in other insect species (20,40,62) is currently under

investigation. However, despite the existing complexity of this immune protease network, the

presence of 110 CLIP genes in the *An. gambiae* genome suggests that additional CLIPB and

CLIPC proteases may be involved (23,25,56). We are currently performing a reverse genetic

screen to identify the contribution of all annotated *An. gambiae* cSPs and cSPHs to melanization

and antimicrobial activity, leveraging the AgMelGCN and recently established *in vivo* bioassays

and (34,77). Completion of this screen will describe the entire molecular make-up of the *An.*

gambiae immune protease network and enable us to identify the molecular interactions within that

are critical for vector competence and survival in this important vector species.

Acknowledgements

We would like to thank all members of the Michel laboratory for mosquito rearing. We thank the

former Michel laboratory members, Ms. Erin Peel for help with the reverse genetic experiments,

and Dr. Xin Zhang for initial plasmid construction and expression of recombinant proteins. We

are grateful to Dr. John Tomich and the members of the Biotechnology/Proteomics Core Facility,

Kansas State University for the MS analysis of the SRPN2/CLIPB4 complexes. We thank Dr.

Susan Paskewitz, University of Wisconsin for the kind gift of the CLIPB4 antibody. We

gratefully acknowledge Ms. Rebekah J. Woolsey and Dr. David R. Quilici at the Mick

Hitchcock, Ph.D. Nevada Proteomics Center Nevada Proteomics Center, University of Nevada,

Reno, for providing the MS analysis of the co-IP experiments. Finally, sincere thanks got to Dr.

Michael R. Kanost, Kansas State University, for his continued support and mentorship.

Statement of Ethics

Not applicable to the enclosed manuscript.

Conflict of Interest Statement

Authors disclosed all potential conflict of interest, and none existed for the presented work.

Funding Sources

This work was supported by National Institutes of Health R01AI095842 and

R01AI140760, and the USDA National Institute of Food and Agriculture Hatch project 1021223

(all to KM). S.Z. was supported by a scholarship from the China Scholarship Council. This is

contribution No. 22-296-J from the Kansas Agricultural Experiment Station. The contents of this

article are solely the responsibility of the authors and do not necessarily represent the official views

of the funding agencies.

Author Contributions

K.M. conceived the study and C.A., C.T.C., M.A.O., C.S. and K.M. designed the experiments and

supervised their execution. X.Z., S.Z., J.K., K.A.S., B.M. V., S.A.S., M.L., E.C.M., and K.M.

performed the experiments and analyzed the data. X.Z., S.Z., and K.M. prepared the manuscript,

and X.Z., S.Z., M.A.O., C.S., and K.M. revised the manuscript. All authors approved the submitted

version.

Data Availability

All generated or analyzed data are included within this article and its supplementary material files.

Further enquiries should be directed to the corresponding author.

References

1. World malaria report 2022 [Internet]. [cited 2023 Jan 16]. Available from:
https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2022

2. Nolan T. Control of malaria-transmitting mosquitoes using gene drives. Philos Trans R Soc
Lond B Biol Sci. 2021 Feb 15;376(1818):20190803.

3. Ickowicz A, Foster SD, Hosack GR, Hayes KR. Predicting the spread and persistence of
genetically modified dominant sterile male mosquitoes. Parasit Vectors. 2021 Sep
16;14(1):480.

4. Sicard M, Bonneau M, Weill M. Wolbachia prevalence, diversity, and ability to induce
cytoplasmic incompatibility in mosquitoes. Curr Opin Insect Sci. 2019 Aug;34:12–20.

5. Huff CGG. Studies on the infectivity of plasmodia of birds for mosquitoes, with special
reference to the problem of immunity in the mosquito. The American Journal of Hygiene.
1927;7(6):706–34.

6. Blandin S, Shiao SH, Moita LF, Janse CJ, Waters AP, Kafatos FC, et al. Complement-Like
Protein TEP1 Is a Determinant of Vectorial Capacity in the Malaria Vector Anopheles gambiae.
Cell. 2004 Mar 5;116(5):661–70.

7. Michel K, Budd A, Pinto S, Gibson TJ, Kafatos FC. Anopheles gambiae SRPN2 facilitates
midgut invasion by the malaria parasite Plasmodium berghei. EMBO Rep. 2005 Sep;6(9):891–
7.

8. Pryzdial ELG, Leatherdale A, Conway EM. Coagulation and complement: Key innate defense
participants in a seamless web. Front Immunol. 2022;13:918775.

9. Cerenius L, Kawabata S ichiro, Lee BL, Nonaka M, Söderhäll K. Proteolytic cascades and
their involvement in invertebrate immunity. Trends in Biochemical Sciences.
2010;35(10):575–83.

10. Kawabata S ichiro, Muta T. Sadaaki Iwanaga: Discovery of the lipopolysaccharide- and beta-
1,3-D-glucan-mediated proteolytic cascade and unique proteins in invertebrate immunity. J
Biochem. 2010 May;147(5):611–8.

11. Kanost MR, Jiang H, Yu XQ. Innate immune responses of a lepidopteran insect, Manduca
sexta. Immunol Rev. 2004;198:97–105.

12. Park JW, Kim CH, Rui J, Park KH, Ryu KH, Chai JH, et al. Beetle immunity. Adv Exp Med
Biol. 2010/01/01 ed. 2010;708:163–80.

13. Buchon N, Silverman N, Cherry S. Immunity in Drosophila melanogaster - from microbial
recognition to whole-organism physiology. Nature Reviews Immunology. 2014;14(12):796–
810.

14. Lemaitre B, Hoffmann J. The host defense of Drosophila melanogaster. Annu Rev Immunol.
2007/01/05 ed. 2007 Jan;25:697–743.

15. Wang Q, Yin M, Yuan C, Liu X, Hu Z, Zou Z, et al. Identification of a Conserved
Prophenoloxidase Activation Pathway in Cotton Bollworm Helicoverpa armigera. Front
Immunol. 2020;11:785.

16. Kanost MR, Jiang H. Clip-domain serine proteases as immune factors in insect hemolymph.
Current Opinion in Insect Science. 2015 Oct 1;11:47–55.

17. Buchon N, Poidevin M, Kwon HM, Guillou A, Sottas V, Lee BL, et al. A single modular
serine protease integrates signals from pattern-recognition receptors upstream of the
Drosophila Toll pathway. Proceedings of the National Academy of Sciences of the United
States of America. 2009 Jul 28;106(30):12442–7.

18. Kwon TH, Kim MS, Choi HW, Joo CH, Cho MY, Lee BL. A masquerade-like serine
proteinase homologue is necessary for phenoloxidase activity in the coleopteran insect,
Holotrichia diomphalia larvae. Eur J Biochem. 2000;267(20):6188–96.

19. Yu XQ, Jiang H, Wang Y, Kanost MR. Nonproteolytic serine proteinase homologs are
involved in prophenoloxidase activation in the tobacco hornworm, Manduca sexta. Insect
Biochem Mol Biol. 2003;33(2):197–208.

20. Jin Q, Wang Y, Hartson SD, Jiang H. Cleavage activation and functional comparison of
Manduca sexta serine protease homologs SPH1a, SPH1b, SPH4, and SPH101 in conjunction
with SPH2. Insect Biochem Mol Biol. 2022 May;144:103762.

21. Collins FH, Sakai RK, Vernick KD, Paskewitz S, Seeley DC, Miller LH, et al. Genetic
selection of a Plasmodium-refractory strain of the malaria vector Anopheles gambiae. Science.
1986 Oct 31;234(4776):607–10.

22. Hurd H, Taylor PJ, Adams D, Underhill A, Eggleston P. Evaluating the costs of mosquito
resistance to malaria parasites. Evolution; international journal of organic evolution. 2005
Dec;59(12):2560–72.

23. Christophides GK, Zdobnov E, Barillas-Mury C, Birney E, Blandin S, Blass C, et al.
Immunity-related genes and gene families in Anopheles gambiae. Science. 2002/10/05 ed.
2002;298(5591):159–65.

24. Neafsey DE, Waterhouse RM, Abai MR, Aganezov SS, Alekseyev MA, Allen JE, et al. Highly
evolvable malaria vectors: The genomes of 16 Anopheles mosquitoes. Science. 2015 Jan
2;347(6217):1258522–1258522.

25. Waterhouse RM, Kriventseva EV, Meister S, Xi ZY, Alvarez KS, Bartholomay LC, et al.
Evolutionary dynamics of immune-related genes and pathways in disease-vector mosquitoes.
Science. 2007/06/26 ed. 2007;316(5832):1738–43.

26. Bartholomay LC, Waterhouse RM, Mayhew GF, Campbell CL, Michel K, Zou Z, et al.
Pathogenomics of Culex quinquefasciatus and Meta-Analysis of Infection Responses to
Diverse Pathogens. Science. 2010/10/12 ed. 2010;330(6000):88–90.

27. Blandin SA, Levashina EA. Reverse genetics analysis of antiparasitic responses in the malaria
vector, Anopheles gambiae. Methods Mol Biol. 2008/03/29 ed. 2008;415:365–77.

28. Zhang X, Li M, El Moussawi L, Saab S, Zhang S, Osta MA, et al. CLIPB10 is a Terminal
Protease in the Regulatory Network That Controls Melanization in the African Malaria
Mosquito Anopheles gambiae. Front Cell Infect Microbiol. 2020;10:585986.

29. An C, Budd A, Kanost MR, Michel K. Characterization of a regulatory unit that controls
melanization and affects longevity of mosquitoes. Cell Mol Life Sci. 2011 Jun;68(11):1929–
39.

30. Zhang X, An C, Sprigg K, Michel K. CLIPB8 is part of the prophenoloxidase activation system
in Anopheles gambiae mosquitoes. Insect Biochem Mol Biol. 2016 Apr;71:106–15.

31. Michel K, Suwanchaichinda C, Morlais I, Lambrechts L, Cohuet A, Awono-Ambene PH, et
al. Increased melanizing activity in Anopheles gambiae does not affect development of
Plasmodium falciparum. Proc Natl Acad Sci U S A. 2006/10/27 ed. 2006;103(45):16858–63.

32. Volz J, Müller HM, Zdanowicz A, Kafatos FC, Osta MA. A genetic module regulates the
melanization response of Anopheles to Plasmodium. Cellular Microbiology. 2006;8(9):1392–
405.

33. Paskewitz SM, Andreev O, Shi L. Gene silencing of serine proteases affects melanization of
Sephadex beads in Anopheles gambiae. Insect Biochemistry and Molecular Biology.
2006;36(9):701–11.

34. Sousa GL, Bishnoi R, Baxter RHG, Povelones M. The CLIP-domain serine protease CLIPC9
regulates melanization downstream of SPCLIP1, CLIPA8, and CLIPA28 in the malaria vector
Anopheles gambiae. PLOS Pathogens. 2020 Oct 12;16(10):e1008985.

35. Povelones M, Bhagavatula L, Yassine H, Tan LA, Upton LM, Osta MA, et al. The CLIP-
domain serine protease homolog SPCLIP1 regulates complement recruitment to microbial
surfaces in the malaria mosquito Anopheles gambiae. PLoS Pathog. 2013;9(9):e1003623.

36. Moussawi LE, Nakhleh J, Kamareddine L, Osta MA. The mosquito melanization response
requires hierarchical activation of non-catalytic clip domain serine protease homologs. PLOS
Pathogens. 2019 Nov 25;15(11):e1008194.

37. Nakhleh J, Christophides GK, Osta MA. The serine protease homolog CLIPA14 modulates
the intensity of the immune response in the mosquito Anopheles gambiae. The Journal of
biological chemistry. 2017 Nov 3;292(44):18217–26.

38. Povelones M, Upton LM, Sala KA, Christophides GK. Structure-Function Analysis of the
Anopheles gambiae LRIM1/APL1C Complex and its Interaction with Complement C3-Like
Protein TEP1. PLOS Pathogens. 2011 Apr 14;7(4):e1002023.

39. Kim MS, Baek MJ, Lee MH, Park JW, Lee SY, Soderhall K, et al. A new easter-type serine
protease cleaves a masquerade-like protein during prophenoloxidase activation in Holotrichia
diomphalia larvae. J Biol Chem. 2002;277(42):39999–40004.

40. Ji Y, Lu T, Zou Z, Wang Y. Aedes aegypti CLIPB9 activates prophenoloxidase-3 in the
presence of CLIPA14 after fungal infection. Front Immunol. 2022;13:927322.

41. Simões ML, Dong Y, Mlambo G, Dimopoulos G. C-type lectin 4 regulates broad-spectrum
melanization-based refractoriness to malaria parasites. PLoS Biol. 2022 Jan;20(1):e3001515.

42. Osta MA, Christophides GK, Kafatos FC. Effects of mosquito genes on Plasmodium
development. Science. 2004 Mar 26;303(5666):2030–2.

43. Kuang J, Buchon N, Michel K, Scoglio C. A global gene co-expression network constructed
from hundreds of experimental conditions with missing values. BMC Bioinformatics. 2022
May 9;23(1):170.

44. Bastian M, Heymann S, Jacomy M. Gephi : An Open Source Software for Exploring and
Manipulating Networks. :2.

45. Fruchterman TMJ, Reingold EM. Graph drawing by force-directed placement. Software:
Practice and Experience. 1991;21(11):1129–64.

46. Girvan M, Newman MEJ. Community structure in social and biological networks. Proceedings
of the National Academy of Sciences. 2002 Jun 11;99(12):7821–6.

47. Rodrigues FA. Network centrality: an introduction. arXiv:190107901 [physics] [Internet].
2019 Jan 22 [cited 2022 Mar 11]; Available from: http://arxiv.org/abs/1901.07901

48. Eidsaa M, Almaas E. s-core network decomposition: A generalization of k-core analysis
to weighted networks. Phys Rev E. 2013 Dec 30;88(6):062819.

49. Keller A, Nesvizhskii AI, Kolker E, Aebersold R. Empirical statistical model to estimate the
accuracy of peptide identifications made by MS/MS and database search. Anal Chem. 2002
Oct 15;74(20):5383–92.

50. Nesvizhskii AI, Keller A, Kolker E, Aebersold R. A statistical model for identifying proteins
by tandem mass spectrometry. Anal Chem. 2003 Sep 1;75(17):4646–58.

51. Jiang H, Wang Y, Ma C, Kanost MR. Subunit Composition of Pro-phenol Oxidase from
Manduca sexta: Molecular Cloning of Subunit ProPO-p1. Insect Biochemistry and Molecular
Biology. 1997 Oct 1;27(10):835–50.

52. Kamareddine L, Nakhleh J, Osta MA. Functional Interaction between Apolipophorins and
Complement Regulate the Mosquito Immune Response to Systemic Infections. JIN.
2016;8(3):314–26.

53. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012 Jul;9(7):676–82.

54. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.
Nat Methods. 2012 Jul;9(7):671–5.

55. Perona JJ, Craik CS. Structural basis of substrate specificity in the serine proteases. Protein
Sci. 1995 Mar;4(3):337–60.

56. Cao X, Gulati M, Jiang H. Serine protease-related proteins in the malaria mosquito, Anopheles
gambiae. Insect Biochemistry and Molecular Biology. 2017 Sep 1;88:48–62.

57. Saab SA, Zhang X, Morejon B, Michel K, Osta MA. Insight into the structural hierarchy of
the protease cascade that regulates the mosquito melanization response. bioRxiv. 2023;

58. Bartholomay LC, Michel K. Mosquito Immunobiology: The Intersection of Vector Health and
Vector Competence. Annu Rev Entomol. 2018;63:145–67.

59. Muller HM, Dimopoulos G, Blass C, Kafatos FC. A hemocyte-like cell line established from
the malaria vector Anopheles gambiae expresses six prophenoloxidase genes. J Biol Chem.
1999;274(17):11727–35.

60. Jiang H, Wang Y, Yu XQ, Zhu Y, Kanost M. Prophenoloxidase-activating proteinase-3 (PAP-
3) from Manduca sexta hemolymph: a clip-domain serine proteinase regulated by serpin-1J
and serine proteinase homologs. Insect Biochemistry and Molecular Biology. 2003 Oct
1;33(10):1049–60.

61. Jiang H, Wang Y, Kanost MR. Pro-phenol oxidase activating proteinase from an insect,
Manduca sexta: a bacteria-inducible protein similar to Drosophila easter. Proc Natl Acad Sci
U S A. 1998;95(21):12220–5.

62. Wang Q, Yin M, Yuan C, Liu X, Jiang H, Wang M, et al. The Micrococcus luteus infection
activates a novel melanization pathway of cSP10, cSP4, and cSP8 in Helicoverpa armigera.
Insect Biochemistry and Molecular Biology. 2022 Aug 1;147:103775.

63. Kanost MR, Jiang H. Clip-domain serine proteases as immune factors in insect hemolymph.
Curr Opin Insect Sci. 2015 Oct 1;11:47–55.

64. Wang Y, Yang F, Cao X, Zou Z, Lu Z, Kanost MR, et al. Hemolymph protease-5 links the
melanization and Toll immune pathways in the tobacco hornworm, Manduca sexta. Proc Natl
Acad Sci U S A. 2020 Sep 22;117(38):23581–7.

65. Barabási AL. Network science. Philosophical Transactions of the Royal Society A:
Mathematical, Physical and Engineering Sciences. 2013 Mar 28;371(1987):20120375.

66. Hahn MW, Kern AD. Comparative Genomics of Centrality and Essentiality in Three
Eukaryotic Protein-Interaction Networks. Molecular Biology and Evolution. 2005 Apr
1;22(4):803–6.

67. Pierson E, Consortium   the Gte, Koller D, Battle A, Mostafavi S. Sharing and Specificity of
Co-expression Networks across 35 Human Tissues. PLOS Computational Biology. 2015 May
13;11(5):e1004220.

68. Dong Y, Manfredini F, Dimopoulos G. Implication of the Mosquito Midgut Microbiota in the
Defense against Malaria Parasites. PLOS Pathogens. 2009 May 8;5(5):e1000423.

69. Schnitger   a. KD, Kafatos FC, Osta M a. The melanization reaction is not required for
survival of Anopheles gambiae mosquitoes after bacterial infections. J Biol Chem.
2007;282:21884–8.

70. Yassine H, Kamareddine L, Osta MA. The mosquito melanization response is implicated in
defense against the entomopathogenic fungus Beauveria bassiana. PLoS Pathog. 2012/11/21
ed. 2012;8(11):e1003029.

71. Fraiture M, Baxter RHG, Steinert S, Chelliah Y, Frolet CC, Quispe-Tintaya W, et al. Two
Mosquito LRR Proteins Function as Complement Control Factors in the TEP1-Mediated
Killing of Plasmodium. Cell Host and Microbe. 2009/03/17 ed. 2009;5(3):273–84.

72. Ruiz VMR, Sousa GL, Sneed SD, Farrant KV, Christophides GK, Povelones M. Stimulation
of a protease targeting the LRIM1/APL1C complex reveals specificity in complement-like
pathway activation in Anopheles gambiae. PLOS ONE. 2019 Apr 8;14(4):e0214753.

73. Gupta S, Wang Y, Jiang H. Manduca sexta prophenoloxidase (proPO) activation requires
proPO-activating proteinase (PAP) and serine proteinase homologs (SPHs) simultaneously.
Insect Biochemistry and Molecular Biology. 2005 Mar;35(3):241–8.

74. Clark KD, Strand MR. Hemolymph melanization in the silkmoth Bombyx mori involves
formation of a high molecular mass complex that metabolizes tyrosine. J Biol Chem. 2013
May 17;288(20):14476–87.

75. Zhu Y, Wang Y, Gorman MJ, Jiang H, Kanost MR. Manduca sexta serpin-3 regulates
prophenoloxidase activation in response to infection by inhibiting prophenoloxidase-
activating proteinases. J Biol Chem. 2003 Nov 21;278(47):46556–64.

76. Christen JM, Hiromasa Y, An C, Kanost MR. Identification of plasma proteinase complexes
with serpin-3 in Manduca sexta. Insect Biochem Mol Biol. 2012 Dec;42(12):946–55.

77. Morejon B, Michel K. A zone-of-inhibition assay to screen for humoral antimicrobial activity
in mosquito hemolymph. Front Cell Infect Microbiol. 2023;13:891577.

Tables

Table 1. List of proteins that co-immunoprecipitate with SRPN2

AGAP number*	NCBI Acc. #	Gene Name	Molecular Weight (kDa)	Scaffold			Proteome Discoverer			
				Percent Coverage (%)	Unique Peptide Count	Total Spectrum Count	Score	Percent Coverage (%)	Total Peptide Count	# PSMs
AGAP006911	Q7QIJ8	SRPN2	47	52.8	13	26	99.29	53	13	28
AGAP008364	A0A453YTR1	TEP15	164	19.4	21	27	97.21	21	24	31
AGAP000376	Q7QF98	Transferrin	69	26.9	14	19	73.08	34	19	24
AGAP003250	C0L2J9	CLIPB4	39	33.1	8	11	45.76	36	9	14
AGAP006258	O44250	PPO1**	78	16.4	7	12	43.8	16	7	12
AGAP009217	Q7PVQ0	CLIPB12	38	22.3	5	6	32.9	34	8	11
AGAP004977	O96753	PPO6	79	14.1	6	9	30.87	16	7	10
AGAP011788	A0A1S4H7K2	CLIPA14	46	25.8	7	8	29.89	26	7	9
AGAP028138	Q5TPW1	Gelsolin	83	8.19	5	7	26.84	11	7	9
AGAP008060	Q6JEK4	Imaginal disc growth factor 2	48	8.66	3	3	16.12	16	6	6
AGAP003095	Q8MZM5	Dopachrome conversion enzyme	53	9.09	3	3	14.4	16	5	5
AGAP001377	Q005M5	SRPN11	57	14	5	5	13.85	14	5	5
AGAP003139	A0A1S4GHQ9	SRPN9	50	10.3	3	3	13.67	10	3	4
AGAP002564	F5HKV6	Fructose biphosphate aldolase	39	7.71	2	3	10.56	10	3	4
AGAP010968	A0A1S4H5E1	CLIPA9	40	7.18	2	3	9.22	7	2	3
AGAP011789	Q7PV63	CLIPA6	46	7.03	2	2	9.2	7	2	3
AGAP007406	Q7PT29	Elongation factor 1-alpha	50	7.56	2	2	8.92	11	3	3
AGAP010815	A0A2P1E488	TEP1	152	1.87	2	2	8.4	3	3	3
AGAP007037	F2Y0G4	APL1C	82	5.7	3	3	7.95	6	3	3
AGAP003689	A0A1S4GIW4	CLIPC7	67	4.19	2	2	5.82	4	2	2
AGAP010724	Q7PCX0	Histon H4	11	23.3	2	2	5.19	23	2	2
AGAP003691	A0A1S4GJE4	CLIPE12	98	3.13	2	2	4.88	3	2	2
*AGAP number based on AgamP4.13										
**named PPO2 in VectorBase										

887
888

Figures

Fig. 1. Gene co-expression network of genes with putative roles in melanization (AgMelGCN). (A) The network topography of AgMelGCN containing 178 nodes and 1229 edges. Of all annotated putative CLIP serine proteases, CLIPB4 has the highest node degree in the network and is co-expressed with SRPN2. (B) Genes in the network that are co-expressed with CLIPB4, which includes SRPN2. (C) Genes in the network that are co-expressed with SRPN2, which includes CLIPB4. The position of every node in the network visualization is identical in all three panels.

Fig. 2. SRPN2 forms SDS-stable complex with CLIPB4 and inhibits its activity. (**a**, **b**, **c**) Detection of CLIPB4-SRPN2 complex by immunoblot analysis. Recombinant proCLIPB4$_{Xa}$ was activated by Factor Xa, and then incubated with recombinant SRPN2 with a molar ratio of 1:1 or 10:1 (rSRPN2: proCLIPB4$_{Xa}$) at room temperature for 15 min. The mixtures were subjected to 12% SDS-PAGE followed by immunoblot analysis with mouse anti-His (**a**) or rabbit anti-SRPN2 (**b**) or rabbit anti-CLIPB4 (**c**) antibodies. Circles, the proCLIPB4$_{Xa}$ zymogen; asterisks, catalytic domain of proCLIPB4$_{Xa}$; solid triangles, recombinant SRPN2; hollow triangles, truncated SRPN2; *arrows*, CLIPB4/SRPN2 complex. (D) Stoichiometry of SRPN2 inhibition of CLIPB4. Factor Xa-activated CLIPB4 was incubated with recombinant SRPN2 at various molar ratios at room temperature for 15 min. Residual activities of CLIPB4 were measured with IEAR*p*NA as substrate and plotted as mean±1 SD (n=3) against the corresponding molar ratios of rSRPN2 and proCLIPB4$_{Xa}$.

Fig. 3. ds*CLIPB4* partially recovers the phenotype of SRPN2 depletion. (**a**) *CLIPB4*kd alone

did not affect survival rates, but reverted the increased mortality induced by SRPN2 depletion.

Survival curves were plotted as means with 95% confidence intervals (n=3). (**b**) Median survivals

in days were plotted (mean ± 1 SEM) based on results shown in (**a**); statistically significantly

differences are indicated by different letters (one-way ANOVA followed by Newman-Keuls test,

P<0.01). (**c**) Total melanotic area per abdomen decreased due to ds*CLIPB4* in SRPN2-depleted

genetic background. Medians and interquartile ranges are marked (n=104, 5 biological replicates),

and statistical analysis was performed with the Mann-Whitney U test. (**d**) SRPN2 depletion

increased melanotic excreta, which was tempered partially by ds*CLIPB4* treatment. Data are

shown as dots for individual replicates, bars represent means ± 1 SD (n = 5); Data were distributed

normally (Shapiro-Wilk test, *P*<0.0001), and statistically significantly differences are indicated by

different letters (one-way ANOVA followed by Newman-Keuls test, P<0.01).

Fig. 4. CLIPB4 cleaves and activates *M. sexta* PPO1&2. Factor Xa-activated proCLIPB4$_{Xa}$

cleaves MsPPO1&2 in plasma (**a**) and purified protein (**b**), detected by immunoblot using rabbit

anti-MsPPO1&2 antibody. The light and dark gray arrows indicate full and cleaved MsPPO1&2,

respectively. CLIPB9 cleavage of *M. sexta* PPO was used as a positive control. The cleavage by

CLIPB4 leads to increased PO activity in plasma (**c**) and with purified PPO protein (**d**). Bars

represent mean ± 1 SD (n = 3); statistically significantly differences are indicated by different

letters (one-way ANOVA followed by Newman-Keuls test, P<0.05).

Fig. 5. CLIPB4 activates B8, not B9. Recombinant proCLIPB4$_{Xa}$ was activated by Factor Xa,

and subsequently incubated with 100 ng of recombinant proCLIPB8 or proCLIPB9 at room

temperature for 15 min. Samples were subjected to 12% SDS-PAGE followed by immunoblot

with mouse anti-His (**a**, **c**), rabbit anti-CLIPB8 (**b**), and rabbit anti-CLIPB9 (**d**) antibodies,

respectively. In parallel, 100 ng of Factor Xa-activated proCLIPB8$_{Xa}$ or proCLIPB9$_{Xa}$ were used

as positive controls. Circles, proCLIPB4$_{Xa}$ zymogen; asterisks, catalytic domain of CLIPB4;

hollow squares, recombinant proCLIPB8 or proCLIPB9 zymogen; solid squares, catalytic domain

of proCLIPB8; hollow triangles, proCLIPB8$_{Xa}$ or proCLIPB9$_{Xa}$ zymogen; solid triangles, catalytic

domain of proCLIPB8$_{Xa}$ or proCLIPB9$_{Xa}$. (**e**) 72 ng of Factor Xa-activated proCLIPB4$_{Xa}$ was

incubated with 450 ng of recombinant proCLIPB8 at room temperature for 1 h, and the catalytic

activity of CLIPB8 was determined using FVR as substrate; 450 ng of Factor Xa-activated

proCLIPB8$_{Xa}$ was used as a positive control. ars represent mean $\pm$ 1 SD (n = 3); statistically

significantly differences are indicated by different letters (one-way ANOVA followed by

Newman-Keuls test, P<0.05).

Fig. 6. CLIPB4 is required for infection-induced depletion of CLIPB8 from the hemolymph

of *An. gambiae*. Female mosquitoes were treated by ds*LacZ*, ds*CLIPB4* and ds*CLIPB8*. Posted

recovery, mosquitoes were challenged by *S. aureus* (OD_{600}=0.8). Ten hours post infection,

hemolymph was collected by proboscis clipping, separated by SDS-PAGE, and subjected to

immunoblot using rabbit anti-CLIPB8 and rabbit anti-APOII antibodies. Results were shown from

two independent experiments.