Title: A growth chart of brain function from infancy to adolescence based on electroencephalography

Authors: Kartik K. Iyer[1, 2]*; James A. Roberts[1]; Michaela Waak[2,3]; Simon J. Vogrin[4]; Ajay Kevat[2,3]; Jasneek Chawla[2,3]; Leena M. Haataja[5]; Leena Lauronen[5] ; Sampsa Vanhatalo[5] ; Nathan J. Stevenson[1]*

Affiliations:

[1] Brain Modeling Group, QIMR Berghofer Medical Research Institute; Brisbane, Australia.

[2] Faculty of Medicine, The University of Queensland; Brisbane, Australia.

[3] Queensland Children's Hospital; Brisbane, Australia.

[4] St Vincent's Hospital; Melbourne, Australia.

[5] BABA Center, Pediatric Research Center, Departments of Physiology and Clinical Neurophysiology, Children's Hospital, Helsinki University Hospital and University of Helsinki; Helsinki, Finland.

*Corresponding authors. Email: Kartik.Iyer@qimrberghofer.edu.au ;
Nathan.Stevenson@qimrberghofer.edu.au

One Sentence Summary: Brain age from electroencephalography provides an accurate functional measure of cortical maturation from infancy to adolescence

Abstract:

In children, objective, quantitative tools that determine functional neurodevelopment are scarce and rarely scalable for clinical use. Direct recordings of cortical activity using routinely acquired electroencephalography (EEG) offer physiologically reliable measures of brain function. Here, we develop a novel measure of functional brain age (FBA) using a residual neural network based interpretation of the pediatric EEG. We show that the FBA from a 10 to 15 minute segment of 18-channel EEG during light sleep (stages 1 and 2) in typically developing children and adolescents was strongly associated with chronological age (R^2 = 0.96, 95%CI: 0.94 - 0.96, n = 1062, age range: 1 month to 18 years). The mean absolute error (MAE) between FBA and age was 0.6 years (n = 1062), with an MAE of 2.1 years following validation on an independent set of EEG recordings (n = 723). The FBA detected group level maturational delays in a small cohort of children with abnormal neurodevelopment (p = 0.00053, n = 40). Our work offers a practical, scalable and powerful automated tool for tracking maturation of brain function throughout childhood with an accuracy comparable to that of widely used physical growth charts.

INTRODUCTION

Deriving brain age in relation to a person's chronological age has become a burgeoning focus for scientists, clinicians, and education providers (*1*). The physiological maturation of the brain is fundamentally shaped by an interaction between an individual's genetic traits and acquired/environmental effects (*2, 3*). The chronological age of an individual is a particularly important benchmark in the assessment of pediatric populations, given that measures of brain age, whether functional or structural, are always judged as being 'correct, or delayed for age'.

The best known example of this practice is the use of anatomical growth charts, such as height, weight and head circumference, which provide an important, widely-used tool for early childhood assessment (*4*). In recent times, there has been a surge of alternative physiological and phenotypic measures aimed at capturing developmental trajectories across the lifespan (*5-9*). Biological age surrogates have been advanced by the development of measures across system-wide ageing processes occurring in cardiovascular, metabolic, and brain systems, demonstrating that tracking biological predictors of ageing may provide an important proxy for quantifying health status (*9, 10*).

Assessment of brain age has gained the most attention in recent studies of biological age predictors. Brain age models are predominantly built upon morphometric measurements of brain structure such as cortical thickness, grey matter, white matter, and intracranial volumes (*2, 3, 11-13*) derived from large bio-banking collections of magnetic resonance imaging (MRI) scans. Across these studies, the key biomarker assessed is referred to as 'brain age delta', "brain age gap" or 'brain-predicted age difference' ('brain-PAD'), and defines the difference between predicted brain age and chronological age as a biomarker (*14*). Prior work has also focused on adult populations to show that individual brain age trajectories are associated with

neurodegenerative illness and other conditions associated with compromised neurological health

(*15, 16*) and can be used to monitor the response to intervention (*13, 17, 18*) .

The applicability of measures of functional brain maturation is arguably more promising in

pediatrics than later parts of the lifespan, due to rapid functional and structural changes occurring

in the brain during childhood and adolescence (*19-22*) and given that over 10% of children

worldwide (*23*) are affected by neurodevelopmental problems. Anatomical measures of MRI

modalities of brain age have emerged in detecting deviations in brain structure (*24-26*), but brain

structure is not perfectly correlated with function; the pre-dominant concern for clinicians. The

desire to assess function has promoted the use of behavioral assessments such as

neuropsychological testing and gross motor tests. While these tests are well-aligned with primary

clinical tools that assess core developmental functions, they often succumb to subjective

interpretations that affect inter-rater reliability and diverge due to the evolving nature of the

toolsets that practitioners employ (*23, 27*). Objective tools are necessary to overcome potentially

confounding effects (e.g. anatomical variations unrelated to development) and account for effects

of plasticity in the developing brain that accurately reflects the true neurodevelopment stage at

the time of the test (*28*). Furthermore, recent studies have also revealed that altered gene

expression and genetic mutations, such as those present in trisomy 21, autism spectrum disorder

or attention-deficit/hyperactivity disorder, effect brain function at the neuronal level (*29-31*),

suggesting that more direct measurement of brain function through techniques such as the

electroencephalogram (EEG) provide an opportunity for neurodevelopmental assessment. The

EEG's high temporal resolution enables the capture of subtle changes in brain function across

dynamic brain states such as evoked responses, resting states and sleep providing valuable

insights into typical and atypical maturation during neurodevelopment (*32-35*). Developing

neurodevelopmental biomarkers that are associated with these cellular-level changes to brain

functions could thus provide key actionable indicators that guide early intervention and

personalization of clinical care to ultimately impact long term outcomes.

Our study aimed to design a 'functional brain age growth chart' (FBA) for children and

adolescents based on the cortical activity measured by the electroencephalogram (EEG). The

EEG is well-suited for scalable and non-invasive measures of brain function in children, with

wide availability in clinical settings. We used EEG activity during sleep to harmonize vigilance

states across participants at varying stages of childhood. We collected EEG from three cohorts of

children: two large cohorts of typical neurological development to develop and validate the FBA

and a small cohort of atypical neurodevelopment to test the diagnostic potential of the FBA. Our

modeling approach sought to develop a normative brain age measure with modern machine

learning (ML) methods, and rigorous statistical analysis (*36, 37*), based on (i) a deep learning,

graph neural network architecture with the raw EEG signals as input and, (ii) a combination of

EEG summary metrics linked to spontaneous cortical activity features combined via

multivariable regression models. These approaches yielded a 'functional brain age growth chart'

of typical development.

RESULTS

107
The framework for predicting functional brain age (FBA) from the sleep EEG is presented in

Figure 1. A FBA was developed using a supervised residual neural network architecture (Res-

NN). A 'bag of features', Gaussian process regression (GPR) predictor of age was also assessed

for further benchmarking of age estimation methods. Our input for the FBA model consisted of

60 second epochs of EEG, where an FBA per recording was calculated as an average across

multiple epochs extracted from a 10 or 15 minute segment. We developed the FBA on a primary

training dataset (D1) which comprised 15 minutes of 18-channel EEGs from 1062 subjects

recorded at the Helsinki University Children's Hospital, Finland. We then validated our trained

model on a dataset (D2) which comprised of 2-channel EEGs recorded from 723 subjects at

Queensland Children's Hospital, Brisbane, Australia.

118
Figure 1 – Study design. **a.** EEG acquisition from routine sleep studies from 2 sites (D1 – Helsinki and D2 –

Brisbane). The development of a functional brain age (FBA) growth chart within a supervised learning framework.

Single, 60 s EEG epochs per recording were used to form a preliminary estimate of FBA that was then averaged

across all available epochs within a 10-15 minute segment of EEG. Model evaluation (FBA) involved cross-

validation procedures from the primary developmental dataset (D1) with external validation on an independently

collected dataset (D2). Growth charts for D1, D2 and a combined D1/D2 dataset were computed. **b.** Distributions for

training (D1) and validation (D2) datasets across age and the EEG channel montages of acquisition. Data consisted

of 5 minute epochs of N1 sleep followed by 10 minute epochs of N2 sleep from 18 channels for D1, and 10 minute

epochs of N2 sleep from 2 channels for D2. **c.** A trained Residual Neural Network (Res-NN) was our primary

method of feature extraction for FBA estimation. Features were combined to form an estimate of FBA using linear

regression for Res-NN outputs, with performance benchmarked against a Gaussian process regression model on *a*

priori engineered EEG summary measures (*38*); (see Methods for further details). **d**. Performance of the FBA was

assessed via measures including the mean and weighted absolute error (MAE, wMAE) and predicted age differences

(PAD = FBA minus actual age). Effects of age, sex, and recording site were examined using statistical tests. The

behavior of the final trained Res-NN was explained using links between network activation, age, and engineered

EEG features across sites.

EEGs from D1 contained recordings that captured the transition between N1 and N2 sleep and

were used to optimize the FBA. Each 15 minute segment consisted of 5 minutes of N1 followed

by 10 minutes of N2. Recordings in D2 contained 10 minutes of EEG from 2-channels during the

N2 sleep state. The performance accuracies of all models tested were evaluated via mean

absolute error (MAE), an age-adjusted MAE or weighted (wMAE; the average MAE across

discrete age bins), and root mean square error (RMSE).

Functional brain age.

The FBA generated by the Res-NN was the most accurate predictor of chronological age (**Figure**

2a-b; MAE = 0.57 years, weighted MAE = 0.88 years, RMSE = 0.83 years; 10 fold cross-

validation on D1, *n = 1062*), providing relatively uniform age predictions across the age range,

with a median deviation of 13.9% from chronological age (see **Figure 3a**). Our comparative

benchmarking of the Res-NN against a GPR approach and physical growth measures indicated

that the neural network architecture exceeded (i) the age estimation accuracy of a GPR model

(Cohen's $d = 0.14$, $p = 1.2 \times 10^{-3}$, $n = 1062$, df $= 1060$) which had a MAE of 0.78 years

(**Supplementary Figure 1c, d**; wMAE $= 1.1$ years, RMSE $= 1.08$ years); (ii) the highest-

performing individual EEG feature predictor (5[th] percentile of EEG amplitude; MAE $= 1.3$ years,

wMAE $= 1.6$ years, RMSE $= 1.7$ years; **Supplementary Table 1**); iii) a prediction based on head

circumference (estimated MAE $= 1.72$ years, wMAE $= 2.54$ years – see Methods for details);

and iv) a prediction based on the mean age (MAE $= 3.52$ years, wMAE $= 4.47$ years; n.b. this

was performed to determine an upper bound for MAE based on the age distribution of the

cohort).

160
161 ***Figure 2** – Functional brain age estimated in children and performance in training dataset (D1).* **a.** Res-NN model

162 ($R^2=0.96$, MAE$=0.57$ years). The dashed line indicates an error bound of ±1.5 years. Individual EEG recordings

163 (blue filled circles) are plotted across ages. **b.** The residual error (predicted age difference; PAD) represents the

164 difference between the FBA measure and chronological age of an individual, in years.

165
FBA model interpretability.

To ascertain the presence of any bias in the data, we first examined the effects of age and sex on

the FBA. The residual error between FBA and chronological age (referred to here as the

predicted age difference or PAD) indicated a small bias in FBA estimates as age increased ($\beta = -$

0.031, $p = 4$ x 10^{-9}, $n = 1062$, df $= 1060$). There were no significant differences in PAD between

males and females ($p = 0.1$, t-statistic $= 1.7$, unpaired t-test, $n = 1062$, df $= 1060$). Interactions

between age and sex did not confound the relationship between PAD and age but a significant

interaction with age alone was observed; PAD increased with age (PAD ~ sex: $\beta = 0.06$, $p =$

0.37; PAD ~ age*sex: $\beta = 0.011$, $p = 0.32$, PAD ~ age^2: $\beta = -0.012$, $p = 3.2 \times 10^{-17}$, $n = 1062$, df $=$

1060; **Figure 3a**).

We next examined the behavior of the trained Res-NN using a dimensionality reduction method

(UMAP (*39*)) (**Figure 1c**; see also **Supplementary Figure 2** for general architectures). We

observed that early layers of network activations showed distinct age related clustering, a pattern

that becomes increasingly resolved with network depth (**Figure 3b**). We also observed that

network activations were related to a range of EEG amplitude, frequency and entropy-based

features. We found that EEG features, used in our GPR model were highly correlated with

activations present across early, middle and late stages of the NN architecture (median $R^2 = 0.48$,

IQR $= 0.31$, across all 32 features), suggesting that the Res-NN architecture captures several

fundamental time and frequency domain characteristics of the EEG signal that are measureable

with independent summary measures within its training phases (see **Supplementary Figure 3**

and **Supplementary Figure 4**). These relationships suggest that EEG features can be viewed as

data surrogates that track the behavior of neural network activations.

In addition to these tests, we also examined whether EEG electrode location on the scalp and the

timing of an epoch during an EEG segment (i.e., with respect to transitions between N1 and N2

sleep) influenced the FBA. FBA accuracy was higher for posterior channels (**Figure 3c**), e.g.

central, occipital (average MAE $= 0.63$ years), whereas anterior channels (e.g. frontal) had lower

accuracies (average MAE = 0.79 years). Furthermore, these spatial differences contributed to

prediction accuracy (p = 3 x 10^{-24}, Kruskal-Wallis test). Applying the predictor to data with

swapped anterior and posterior channel positions resulted in a reduced performance accuracy

(MAE = 1.10 years) whereas data with left hemisphere channels swapped with right hemisphere

channels did not alter overall performance accuracy (MAE = 0.55 years). Temporally, the

accuracy of the FBA was highest during the transition between N1 and N2 sleep states (**Figure**

3d; MAE = 0.61 years). Taken together, averaging FBA estimates across time and space

improved overall accuracy.

Figure 3 – *Interpreting the EEG-based FBA*. **a.** The mean absolute error (MAE) of the Res-NN (black circles, left axis) across 2 yearly age bins and the relative accuracy of PAD (in blue, right axis) across 2 yearly age bins. For both plots, the mean and standard error of the mean (SEM) are plotted to reflect the sample distribution within age bins. **b.** A lower-dimensional representation of the Res-NN network generated by UMAP on 18-channel D1 data. The network is composed of 63 layers represented by early (layer 6), middle (layer 29) and late stages (layer 61). Here, the Res-NN of the EEG clusters into younger age groups (blue) and older age groups (red) throughout the

training phase, MAE was calculated using UMAP values as predictors (no cross-validation). **c.** The performance accuracy of the Res-NN model (MAE, years) following individual cross-validation per EEG channels. Colors are ordered by right/left channel derivations, where frontal channels (Fp and F in yellow) collectively had higher MAEs than temporal (T in teal), central (C in light blue) and occipital channels (in dark blue) which had comparatively lower MAEs. The average MAE across channel regions is shown with the standard deviation shown as error bars. **d.** The temporal change in MAE of epoch sequences obtained sequentially from N1 (light pink) and N2 (light purple) indicated that the lowest MAEs were observed during a transition between sleep N1 and N2 stages. The MAE is shown with error bars indicating the standard deviation.

Validation.

We then validated the FBA using a secondary dataset (D2) composed of 723 subjects recorded at the Queensland Children's Hospital. To homogenize datasets across the different recording configurations in D1 and D2, we re-trained D1 on a 2-channel bipolar montage (F4-C4, C4-O2) and selected only periods of N2 sleep. In applying this retrained model to the validation set (trained D1 2-channel, tested D2 2-channel), the MAE of the FBA was significantly higher (**Supplementary Figure 5a;** MAE = 2.15 years; wMAE = 2.28 years) when compared to 2-channel models trained and tested on D1 and D2 individually (**Table 1**). Notably though, the validation performance still outperformed the estimated accuracy of simulated head circumference based models (wMAE = 2.54 years; see *Growth Charts* in Methods).

The decrease in FBA performance was attributed to three key site differences: age distribution, number of EEG channels, and site-specific differences of the EEG across age. In comparing MAE to wMAE, distributions of age contribute approximately 0.2 years to the increase in MAE (**Table 1**). Similarly, a reduction in the number of EEG channels from 18 to 2 resulted in an increase of 0.2 years in the wMAE (**Table 1**). Despite the effectiveness of the Res-NN in

capturing fundamental characteristics of the EEG signal (**Supplementary Figure 5b**), several

age-specific differences in individual summary EEG features were observed across sites

(**Supplementary Figure 5c** and **d.** Additionally, we found noteworthy distinctions in spectral

estimates of the EEG recording noise floor between sites, whereby EEG recordings in D1

exhibited a higher frequency noise floor compared to D2 (70 to 80 Hz band power; $p = 3 \times 10^{-9}$,

unpaired t-test). Training a FBA model constructed from a combined dataset D1 and D2

considerably improved overall prediction accuracy with a MAE of 1.04 years for D2 ($n = 723$)

suggesting site specific differences were incorporated into the model.

Functional growth charts.

We next combined both datasets (D1 + D2) to generate a FBA 'growth chart', wherein

generalized additive models were applied to construct age-appropriate centiles (*4, 40*). The

resultant FBA had an MAE of 1.04 years with a wMAE of 1.32 years (**Figure 4a**; see also **Table**

1). Further, FBA growth charts based on an age-stratification of infants (0 to 2 years; MAE =

0.55 years) and children (2 to 18 years; MAE = 1.23 years) indicated a high degree of accuracy

relative to their respective age group (**Supplementary Figure 6**).

The practical utility of an FBA is that it enables stratification of children, by quantifying brain

functions associated with a child's diagnostic status and underlying neurodevelopmental issues.

To demonstrate this, we compared typically developing children from D2 with an additional

small cohort of children from the same site whom were diagnosed with Trisomy 21 ($n = 40$;

29/40 were recorded at less than 7 years of age). The FBAs (trained on D2 only and tested on

children diagnosed with Trisomy 21 children) relative to age, despite having normal sleep

studies, were found to be significantly lower than typically developing children (**Figure 4b**; $p = $

0.00053, t-statistic = 3.5, unpaired t-test, n = 763, df = 761). Similar comparisons using the FBA

on (i) trained on D1 and tested on D2 + Trisomy 21; and, (ii) trained on D1 + D2 and tested on

D2 + Trisomy 21 also indicated significant differences (i): p = 0.012 (t-statistic = 2.5, unpaired t-

test, df = 761) and (ii): p = 0.03 (t-statistic = 2.2, unpaired t-test, df = 761). The observations

suggest that at the group level, deviant neurodevelopmental trajectories in Trisomy 21 may

translate to delayed maturation of their cortical function.

Finally, we benchmarked our FBA model against conventional growth chart trajectories of head

circumference and height in children (*41, 42*). Here, the maximal variation of age for the FBA

(difference between 97[th] and 3[rd] centiles) falls between head circumference and height for age for

a simulated cohort with similar age and sex demographics to D1 and D2 combined (**Figure 4c**).

Our FBA growth chart thus exhibits comparable age variability to that of widely-used physical

growth charts. Code for converting EEG into FBA and centiles is available at

https://github.com/brain-modelling-group/functional-brain-age.

267
268
Figure 4 – The functional brain age (FBA) growth chart derived from cross-validation of a combination of D1 and

D2 datasets. **a.** The chart is based on the FBA derived from the Res-NN model. Translucent colored triangles

represent individual EEG recordings from D1 (blue) and D2 (red) datasets. The 3rd (red), 15th (yellow), 50th (green),

85th (purple) and 97th (blue) centiles are indicated. **b.** Differences in relative PAD between subjects with typically

developing neurodevelopment and Trisomy 21. Significance values ($p < 0.05$) were determined by conducting an

unpaired t-test between groups, where all data was checked for normality. Typically developing groups ($n = 723$;

blue) and Trisomy groups ($n = 40$; pink) from D2 are plotted as violin plots, with the median (black line) and

interquartile ranges (rectangles) shown, **c.** Variation in pediatric predictors of age. The variation (in years) represents

the difference between the 97th and 3rd centiles across age for the FBA (in blue), head circumference (HC, in red)

and height (in green). HC and Height predictors were simulated with the same age distribution as the combined D1

and D2 dataset.

284   **Table 1** – *Overall performance of FBA across datasets.* The performances of the FBA, using a Res-NN
285   model, across all training, test, and combined datasets. MAE, wMAE and RMSE are shown. CI is
286   confidence interval, [a]only N2 from D1 used, [b]10-fold cross-validation.

Train	# channels, # recordings, # epochs	Test	# channels, # recordings, # epochs	MAE (in years; 95% CI)	wMAE (in years; 95% CI)	RMSE (in years)
D1	19, 1062, 30798	D1[b]	19, 1062, 31001	0.57 (0.53 - 0.61)	0.88 (0.73 – 1.03)	0.83
D1	2, 1062, 20311	D1[b]	2, 1062, 20311	0.65 (0.60 – 0.69)	1.10 (0.88 – 1.29)	0.88
D1[a]	2, 1062, 20311	D2	2, 723, 13737	2.15 (2.03 – 2.28)	2.28 (1.87 – 2.70)	2.06
D2	2, 723, 13737	D2[b]	2, 723, 13737	1.46 (1.38 – 1.55)	1.73 (1.45 – 2.04)	1.49
D1+D2	2, 1792, 34390	D1+D2[b]	2, 1792, 34048	1.04 (0.99 – 1.09)	1.32 (1.14 – 1.51)	1.35

DISCUSSION

In infants, children and adolescents, it is well known that EEG activity exhibits clear, consistent, and rapid changes with age (*33, 43-45*). We formalize this knowledge using a targeted deployment of deep learning algorithms to form a prediction of EEG age (FBA) underpinned by key human operator expertise and decision-making in specific stages of the process. We achieved an accurate FBA prediction when applying deep neural networks directly to the EEG signal, relying on a summary of only brief epochs (60 seconds) within a 10 - 15 minute segment of routinely recorded sleep EEG during N1/N2. The proposed FBA demonstrated state-of-the-art age prediction accuracy, was validated in an independent cohort and detected group level maturational delays in a small cohort of young children with a defined neurodevelopmental disorder. Tracking maturational delays in young children is of particular interest to clinicians, with the FBA demonstrating its potential for providing an objective, early correlate for neurological development. Our study presents a conceptual advance on prior EEG knowledge by showing that an FBA prediction with modern machine learning and artificial intelligence algorithms facilitates a highly accurate prediction of developmental age, providing similar accuracies with widely used anatomical growth charts (*4, 41, 42*).

We show that the application of deep learning directly to the EEG signal improves the accuracy of the FBA over traditional 'bag of features' approaches. Improved accuracy via these methods suggest that deep neural networks are better able to cope with the non-monotonic relationships between key EEG characteristics such as EEG amplitude, sample entropy, fractal dimension and relative spectral power and age that pose challenges for feature based age predictions (**Supplementary Figure 4**). Our FBA estimates had MAEs comparable to the highest performing MRI-based (*2, 3, 12, 13, 16, 46*) and EEG-based studies (*47-51*) with reported MAEs

in the literature ranging from 1.0 to 4.6 years compared to our best wMAE of 0.88 years. The

accuracy of the FBA may be directly attributable to the use of residual neural network

architectures over conventional multivariable age regression approaches that are typically used in

brain age studies (e.g. GPR). We show that deep learning networks capture well-established EEG

characteristics (such as amplitude, frequency, bursting behavior, and entropy) and that the

representation of these latent patterns improve FBA estimates through higher-dimensional

abstractions of the EEG signal. It is also important to recognize that the fundamental physical

and physiological differences in measures of brain function (EEG) versus structure (MRI): the

EEG captures neuronal level signatures that reflect rapidly maturing cortical function (*52*), which

correlate well with multiple domains of neurocognitive performance and associated

neurocognitive conditions (*53, 54*). The underlying gross brain structure measured by MRI is at

best only sensitive to larger structural alterations such as malformations of cortical development

(*55*).

The present FBA measures were developed solely using large EEG datasets that are routinely

collected and widely available in hospitals worldwide. Although the performance error of the

FBA increases with age, the relative accuracy of the FBA is comparatively uniform across the

age range. Here, additional variations to FBA accuracy were linked to the spatial and temporal

organization of the EEG. The effect of spatial organization for instance was primarily a frontal-

occipital gradient which is a well-established phenomenon in the maturing EEG within this age

group (*56, 57*). Temporally, the accuracy of the FBA was maximum at the onset of N2 sleep

characterized by the presence of sleep spindles which are key cortical signatures that emerge in

the first few months of life and remain present in the EEG through adulthood (*58*). Waking EEG

could also be used in older cooperative children, but collecting several minutes of good quality

EEG signal from alert infants and toddlers is difficult. Thus, the only manual selection done prior

to our computational analyses was the identification of the first sleep spindle as a sign of N2

sleep, which was necessary to harmonize vigilance state across a cohort with a wide age range.

The N2 sleep state offers a well standardized vigilance state that can be considered much more

homogeneous across individuals than the wake or other sleep states (*35*). A brief period of N2

sleep is also often recorded in routine EEG studies as it is rich in EEG signatures, relatively

homogeneous across individuals, known to be sensitive for observing pathological phenomena,

(such as epileptiform events (*59, 60*)), and is also minimally contaminated by the common

artefacts due to subjects' movements. Our additional sleep stage-specific tests (**Supplementary**

Table 4) show that deeper sleep stages such as N3 and REM are potentially valuable for further

disentangling the relationship between finer grained cortical maturation processes related to

growth and homeostatic regulation of sleep during childhood (*61*). Our findings support N2 sleep

as an ideal target for the measurement of cortical activity, suggesting that the FBA can be further

optimized, providing potential avenues for finer-grained studies of brain maturation.

It is likely that a multitude of age-related changes in the synaptic circuitry underpinning the EEG

signal contribute to the high performance of our FBA measure. The development of the EEG

signal across childhood is well-established and includes, for instance, notable increases in the

posterior dominant frequency during wakefulness and sleep, as well as a decrease in the posterior

amplitudes during early childhood (*62, 63*). These developmental changes coincide with

maturing brain cytoarchitecture, including changes in neuronal cell density, dendritic growth,

myelination, and synaptic remodeling (*52*). However, despite concerted efforts to quantify

visually observable changes to the pediatric EEG (*43, 45, 64, 65*), establishing normative values

to directly assess brain maturity have largely been challenging. In addition, biophysical

constraints on the underlying scalp EEG in childhood also change due to anatomical changes in

non-neuronal structures implicated in EEG propagation, such as skull ossification, cortical

thickness, and head circumference (*41, 52*). Our comparison to head circumference (**Figure 5c**)

shows, expectedly, that head size was associated with age, but provides higher variation in age

prediction in comparison to the EEG-based FBA.

The present study has some potential limitations. Firstly, the performance of external validation

was markedly lower than the overall performance of cross-validated results in each site

independently. The drop-off in accuracy due to several factors, namely: site specific differences

and differences across age, a lower electrode density, and inherent differences in acquisition of

the EEG recordings. We show that site differences in EEG features effect network activations

and, therefore, underlying age prediction. There were also differences in the neurodevelopmental

profiling at each site. Cohort selection of D2 was based on normal studies of children with no

overt neurological illnesses nor pathological EEG activity identified in their overnight PSG, and

were not subject to the four-year neurodevelopmental follow up performed in the D1.

Nevertheless, despite the trade-off in performance accuracy, externally validated results still

outperformed head circumference growth chart measures simulated over the same age range. The

combination of datasets largely corrects for these site differences, which suggests that functional

growth charts such as the 'normative' FBA proposed, are likely to benefit from assimilating

more diversity in datasets across multiple sites (*66*).

Across multiple sites, all children included in this study were identified as typically developing

and within clinical neurological norms (except for the trisomy 21 group). However, it is essential

to note that these children are not representative of the larger, healthy pediatric population but

rather a subgroup of children clinically referred from the primary care level to a tertiary care

center for diagnostic assessment. Estimates of FBA in such groups are clinically interesting;

however, it is essential to benchmark the FBA in healthy neurotypical cohorts to ensure further

clarity and confidence in applying FBA to a broader pediatric population. Conducting these

characterizations across wake and sleep states to establish the dynamic range within which EEG-

based brain age estimates can be effectively utilized and adequately interpreted. The FBA

measures in this study are also built upon well-resourced diagnostic centers placed within high-

income populations, but have equal applicability to providing rapid objective tools (e.g. EEG

wearables (*53*)) in lower middle income countries where there is a significant lack of diagnostic

resources (*67, 68*). Further studies are necessary to calibrate the FBA-derived growth charts for

populations with potentially different traits, including other aspects of population demographics

such as race and ethnicity. Continued development of tools such as the FBA in heterogeneous

clinical landscapes are likely to improve overall performance, resulting in better delineation of

diverse normative population profiles and atypical neurodevelopment.

Secondly, we did not find any significantly confounding effect of sex on our brain age estimates,

and have, therefore, generated a single FBA growth chart for both sexes. This does not discount

the fact that sex related differences in cortical activity exist across age, rather, that sex specific

effects in the EEG were accounted for and adjusted out by the model (*44*). Future applications of

the FBA could be used to study genuine sex-related differences in cortical maturation (*44*) by

separating data according to biological sex at the training stage. The FBA also appears to

compensate for, or is indifferent to, growth spurts, hormonal and pubertal changes in both sexes,

and changes to brain structure that occur at differing time points between the sexes, such as

increased rates of cortical thinning in males in adolescence (*69, 70*). This ability of trained

models to inherently adjust for potential confounders is a key aspect of artificial intelligence (AI)

methods in medicine and means that what we know about EEG and age should be reevaluated

singularly in the context of the AI outputs, requiring detailed new studies across a range of

exquisitely designed cohorts. Growth charts can also be calibrated for diagnosis, prognosis, and

stratification; here the optimal tradeoff between cohort heterogeneity, EEG acquisition, training

data size and MAE is not entirely resolved, with evidence from MRI-based studies suggesting

that the clinical utility is not necessarily inversely proportional to MAE (*12, 66*).

The FBA provides an accurate and objective maturational measure of cortical function that is

linked to a child's age. Future iterations of the FBA are needed to rigorously validate its

performance using independent clinical cohorts. While recognizing FBA's clinical potential, a

series of targeted evaluations of the FBA within clinical populations are necessary to determine

its efficacy prior to endorsing its widespread use. It is also essential to evaluate the links and

dependencies between FBA and the host of complementary modalities of neurodevelopmental

assessments, such as standardized neurological and behavioral assessments methods. Rather than

replacing or challenging the existing techniques, the EEG-derived FBA is perhaps best seen as a

valuable complement to support current neurodevelopmental assessment, offering a tool towards

personalization that both benefits the patient and healthcare practitioner alike. Further, the

intricacies of AI necessitate a new set of studies that are informed by, but unbiased to previous

understanding of EEG maturation in the developing brain. These extra studies are required not

only determine the clinical utility of the algorithm but to also enable other researchers and

institutions to identify appropriate safeguards that can be built to ensure decision safety and

efficacy. We, therefore, publicly release the FBA estimation algorithm to enable continued

validation and development via federated learning processes that utilize additional data collected

by independent researchers (*71*). Our approach ensures the preservation of earlier feature

extraction layers while retraining later layers using smaller data samples designed to address

clinical concerns related to demographics, diseases, or health outcomes. These methods-based

outcomes thus generate an 'online' resource that facilitates the continual refinement of targeted

algorithms for tracking childhood brain function and neurodevelopment.

MATERIALS AND METHODS

Study Design

Demographics. Our functional brain age chart aimed to cover the pediatric age range following

the neonatal period (4 weeks) to adulthood (18 years). Our primary training dataset comprised of

electroencephalography (EEG) recordings that were collected as routine sleep EEGs and judged

to be typically developing by visual review. These children were later followed up for at least

four years after the EEG to confirm typical neurodevelopment, which was established as the

absence of major neurological, developmental, or psychiatric diagnoses. The external validation

dataset comprised of EEG recordings obtained as a part of routine polysomnography (PSG;

overnight sleep study) in hospital-based sleep laboratories. Normal outcomes for the external

validation dataset were defined as the absence of sleep disorders based on clinical assessment of

the PSG, and absence of neurological diagnoses. Outcomes were defined by clinical experts

responsible for the EEG/PSG studies. Our functional brain age chart, hence, only included

studies that were defined as having: (i) EEG activity within normal ranges of background activity

for age; and (ii) children identified as being typically developing based on neurodevelopmental

and/or physical outcomes for their age. Children with any of the following outcomes were

excluded: (i) any seizure-related (epileptiform) or aberrant paroxysmal discharges present in the

EEG; (ii) underlying genetic or neurodevelopmental abnormalities disclosed prior to the

recording; and (iii) incomplete data due to technical difficulties (e.g. PSG was incompletely

recorded as the child did not tolerate the PSG monitoring leads). Site-specific characteristics for

each of the EEG datasets are described further below.

Datasets. Our primary training dataset (referred hereto as *D1*) was collected from a convenience

sample of EEGs collected at the Helsinki University Children's Hospital in Helsinki, Finland.

The original cohort included over 6800 recordings, out of which 3200 were reported to be

normal. Following the four-year follow-up period, children who presented with any diagnoses of

neurological diseases (including epilepsy), psychiatric disorders, brain-acting medications,

presence of tumors/cancer, congenital and/or perinatal disorders and malformations were

excluded (2131 recordings). A total of 1062 children with typical neurodevelopment were

available for analysis over the 18 year period. For D1, the Institutional Research Review Board

at Helsinki and Uusimaa Hospital district approved the study (HUS/244/2021) including waiver

of consent due to the retrospective collection of data acquired as part of standard of care. These

datasets were acquired and reviewed between 2011 and 2016. The ages of children for D1 ranged

between 4 weeks and 17 years of age (median=2.6 years, IQR=6.1 years), with 549 males

(min=1.5 months, max=15.8 years; median=2.9 years) and 520 females (min=1 month,

max=16.8 years; median=2.5 years). EEG was recorded using 10-20 electrode positions with an

electrode on the vertex as reference (either Fz or Cz) using a Nicolet One EEG (Natus Medical

Inc. Middleton, WI, USA). All EEG recordings were sampled at 250 Hz and the referential

montage was saved in a pseudonymized EDF file format. A total of 19 channels (Fp1, Fp2, F7,

F3, Fz, F4, F8, T3, C3, Cz, C4, T4, T5, P3, Pz, P4, T6, O1, O2) were collected as part of the

EEG. These recordings were clinically reviewed to define normality of the record, where the

most common montage was a standard longitudinal bipolar montage (double banana). Then, the

EEG record was re-reviewed by EEG technicians trained for the purpose (and approved by L.L),

and the first occurrence of sleep spindle or K-complex was annotated as the onset of an N2 sleep

state (see the guidelines of the American Academy of Sleep Medicine, AASM (*72*)). A 15

minute segment was then extracted and saved in EDF format, consisting of 5 minutes before the

sleep spindle and regarded as N1 sleep, as well as 10 minutes after the sleep spindle regarded as

N2 sleep. It should be noted that these 15 minute segments may include brief arousals or short

awake periods that arise during sleep state transitions. As the recordings were done in the routine

clinical EEG setting, the only preparatory measure given to patients' families was a request to

wake up the child several hours (2-4 hours) earlier than their normal wake-up time to ensure

sufficient sleep pressure for falling asleep in the laboratory; however, no external measures of

possible individual variations in the sleep pressure could be obtained. This period around N1/N2

transition was selected for FBA training because it allows the best comparability with respect to

behavioral states across the whole age range of interest, and because it is often the only state that

contains sufficient amount of artefact-free EEG signal in the younger children.

Our secondary dataset, *D2*, was used for external validation of the FBA algorithm, and included

723 children with normal outcomes following a PSG. Ethics approval for data in D2 was granted

by Children's Health Queensland (LNR/2021/QCHQ/73595) and the QIMR Berghofer Medical

Research Institute (No. P3736, P3727). PSG for these children were used to diagnose the

presence of a pediatric sleep disorder (e.g. central sleep apnea, nocturnal hypoventilation,

obstructive sleep apnea). For this dataset, we retrospectively screened a convenience sample of

3800 sleep studies from the Queensland Children's Hospital (Respiratory and Sleep Medicine

Department), reviewed between 2014 and 2021. Of those, 3376 were exclusively diagnostic PSG

studies, with the remaining group comprising of treatment studies (e.g. oxygen supplementation,

ventilator support). From 3800 studies, 1206 individual studies were identified as having a

'normal sleep study' or 'no evidence of sleep disordered breathing'. Out of these 1206 subjects,

465 children were identified as having neurological sequelae (e.g. ASD, Duchenne Muscular

dystrophy, spina bifida, spinal muscular atrophy, Trisomy 21, seizure activity, central sleep

apnea events higher than 5 events per hour (73)). Upon visual inspection of data, we excluded a

further 18 subjects for having any direct compromise to EEG activity during sleep segments

captured (i.e. EEG channel leads that lost connection). Following exclusion of these subjects, the

final cohort included 723 children who were identified as having a normal sleep study outcome

for age, without any underlying neurological and/or neurodevelopmental conditions.

The age range for D2 was between 2 months and 18 years of age (median age of 8.1 years,

IQR=5.6 years), with 291 females (min= 4 months; max 17.8 years; median=7.9 years) and 432

males (min= 3 months, max=17.9 years, median=8.3 years). As per D1, all sleep stages were

seen and scored by a clinician according to the AASM guidelines. PSG was acquired via the

EMBLA N7000 (Natus Neuro, Middleton, WI, USA). EEG was recorded using 10-20 electrode

positions with reference electrodes attached to the mastoid (A1/M1 and A2/M2). EEG recordings

were sampled at 200 Hz or 500 Hz. For D2, a total of 3 channels were recorded overnight (F4,

C4 and O2) as part of the PSG. Based on epoch and sleep stage availability, the first 10 minute

period of N2 classified sleep was selected from EEG recordings across all ages based on study

hypnograms.

The age of subjects in both D1 and D2 was resolved in months.

Comparison of datasets. While the overall age range, sample size, and the essential technical

descriptors of the EEG signals are comparable between the datasets, there are also some apparent

differences. First, the data from D1 consisted of a full routine EEG setup with 18 bipolar channel

derivations, while D2 only included three electrodes with two bipolar derivations (F4-C4, C4-

O2). Second, the normality of the EEG signal could be more thoroughly characterized in D1 due

to a higher channel count and a longer neurodevelopmental follow-up time, whereas normality in

D2 was based on normal-appearing signals, with additional physiological monitoring used to

assess sleep-disordered breathing and associated clinical outcomes, as well as the absence of

significant neurological illness by the time of PSG study.

530
Data processing. All EEG data were zero-phase filtered in both forward and reverse directions

with an infinite impulse response, bandpass, 4^{th} order Butterworth filter with a lower cutoff of

0.5 Hz and an upper cutoff of 30 Hz (Gaussian process regression) or 15 Hz (residual neural

network regression). EEG data were resampled to 64 Hz and 32 Hz as inputs to the feature

extraction components of the Gaussian process regression and residual network regression,

respectively. Residual network regression approaches employed EEG data inputs at a lower

sampling rate to maximize computational efficiency, with several qualitative tests performed to

ensure that important aspects of sleep EEG were retained (e.g. delta and alpha rhythms in sleep).

Our feature-based methods were examined at a higher sampling frequency (e.g. GPR;

Supplementary Table 1) and ensured that higher frequency components of the EEG were also

captured (e.g. beta activity). For D1, a bipolar montage was computed from the

monopolar/referential EEG data resulting in 18 EEG derivations/channels: Fp2-F4, F4-C4, C4-

P4, P4-O2, Fp1-F3, F3-C3, C3-P3, P3-O1, Fp2-F8, F8-T4, T4-T6, T6-O2, Fp1-F7, F7-T3, T3-

T5, T5-O1, Fz-Cz, and Cz-Pz.

Further, for D1, each 15 minute segment of EEG was divided into 60 s epochs for analysis.

Epochs were extracted with a 30 s overlap (29 epochs per recording). We assumed that 60 s was

sufficiently long to capture important EEG signal characteristics and sufficiently short to reduce

any effects of non-stationarity in the EEG while generating a sufficiently large and diverse set

for model training.

Due to the limited availability of channels for D2, we limited our analysis in this dataset to a

simplified bipolar derivation of the EEG; i.e., F4–C4, C4–O2. Each 10 minute EEG recording in

D2 was segmented into 60 s epochs with a 30 s overlap (19 epochs per recording) and used for

training and testing as described in *Cross-validation, optimization and training.*

At the end of these data curation steps, we then developed a residual neural network regression

(Res-NN) for age prediction (*74*). We also used Gaussian process regression (GPR) model as a

benchmark (GPR) (*75*).

Residual neural network regression. EEG epochs were first resampled to 32 Hz (resampling

included anti-aliasing filtering).

We added variability to the residual neural network by changing the temporal filter width (FW),

filter channel depth (FD) and filter number (FN) within the convolutional layers as well as

increasing the network depth (ND). We used the file *generate_networks_v2.m* to generate

networks with different configurations and architectures (see the GitHub page for more details).

The general structure of the residual network is shown in **Supplementary Figure 2**.

Several parameters specific to the definition of these neural network architectures were selected

during each training iteration. In general, parameters defined the filter size (temporal width and

channel depth), filter number and network depth. Training options such as solver type and mini-

batch size were selected based on preliminary analysis (see Supplementary Material for more

details; **Supplementary Figures 7 to 14**), and an alternate architecture based on inception layers

was also evaluated (*76*).

GPR. This approach combined several summary measures of the EEG features to form a

prediction of age. We chose GPR due to its consistent performance across various imaging

modalities in estimating brain age following feature extraction due to its capability to model the

underlying latent distribution (via Gaussian process prior distribution) and quantifying any

associated uncertainty to provide probabilistic predictions from data (*2, 3, 25, 26, 37*). All EEG

epochs were resampled to 64 Hz (resampling included anti-aliasing filtering). We used a total of

32 features from the EEG. Of these, 7 features were general measures of EEG amplitude activity,

12 features were frequency dependent representations present within the EEG, and the remaining

13 features were further computations of information content drawn from amplitude, frequency,

and entropy based transformations of the EEG signal. A descriptor of the EEG features used is

listed in **Supplementary Table 1** and detailed via our GitHub codes for their implementation in

MATLAB, see *Code Availability* section).

For both D1 and D2, the set of EEG features was estimated for each available channel and then

averaged across channels (with a median operation used). For training, each EEG recording was,

therefore, summarized by an M-by-32 feature matrix (where M was 29 epochs per recording for

D1 and 19 epochs per recording for D2, and there were 32 features per epoch). A combinatorial

function (shown in **Figure 1**) was used to determine the final brain age estimate. This was

achieved by first assuming a Gaussian prior (GP) where $P(f \,|X) = \mathrm{GP}(f\,|\mu, K)$ denotes the

dependence of f (i.e. FBA) on X (i.e. age) across all observed data points and μ and K represent

the mean function and kernel parameters, respectively.

Performance assessment, cross-validation, optimization, and independent validation.

Performance assessment: We utilized two commonly used measures to compare the predicted age to actual age and, therefore, define the accuracy of prediction: mean absolute error and root mean square error. The root mean square error (RMSE, Eq.1) between predicted age per recording and actual age was defined as

$$RMSE = \sqrt{\frac{1}{N}\sum_{\forall n}(a_n - b_n)^2} \qquad (Eq.1)$$

where a_n is the age of the n^{th} EEG recording in years, b_n is the median predicted age across all extracted EEG epochs per recording, and *N* is the number of EEG recordings (1062 for D1 and 723 for D2). The mean absolute error (MAE, Eq.2) was defined as:

$$MAE = \frac{1}{N}\sum_{\forall n}|a_n - b_n| \qquad (Eq.2)$$

We supported these performance measures with weighted MAE. The wMAE is a variation of the MAE that normalizes across the distribution of the data with respect to age; i.e., an approximation of the MAE for a cohort with uniformly distributed age. We defined it as the average MAE across age binned averages of the MAE (bin width = 2 years). We also computed the relative error with respect to age, by calculating the percentage change between predicted age (FBA) and chronological age.

Cross-validation, optimization and training:

We used D1 as our primary training set for developing a functional brain age. To evaluate the

accuracy of prediction, we used 10-fold cross-validation. This entailed a train-test split where

90% of the cohort of EEG recordings were used for training and 10% for testing. This process

was repeated 10 times so that all EEG recordings were excluded from training for at least one

iteration (fold).

For residual neural network regression, due to computational limits, initial experiments of

randomly selected combinations of network architectures were used to select several model

parameters based on maximum regression accuracy (minimum RMSE) on 10-fold cross-

validation outputs. These parameters included training solver and mini-batch size. It was

assumed that the selection of these parameters was less prone to overfitting than internal network

weights from variability in architecture. Architecture parameters were selected using an internal

10-fold cross-validation (9:1 training data split). We used a small fixed grid approach and

selected the parameter combination (filter width, filter depth, filter number, network depth) that

minimized the RMSE. A single optimal architecture was defined by averaging internal validation

RMSE across all 10 training iterations (a point of data leakage assumed to result in negligible

overfitting). Remaining training hyperparameters were not optimized: initial learning rate was

0.0001 for training for initial selection and nested architecture selection, training drop factor was

0.1 every 8 epochs, maximum number of training epochs was 50, 10% of recordings within the

training set were used for internal validation, and no changes in RMSE on the validation set

within 6 training iterations (epochs) stopped training. The squared gradient decay factor was 0.99

for the ADAM solver. Data extraction and network implementation and training was performed

in MATLAB (The MathWorks Inc, Natick, Massachusetts, USA: Deep Learning Toolbox;

R2020a or R2021a with GTX 1080 or 3070 graphics cards, for dataset D1 and datasets D1 and

D1+D2, respectively).

Additional parameters of the Res-NN model were also benchmarked to gauge performance

limits, including epoch length (30 s, 60 s, 120 s and 300 s in length; **Supplementary Figure 7**),

the type of solver of min-batch size (**Supplementary Figure 8**), the effect of uniform versus

non-uniform age distribution of ages during training (**Supplementary Note 1** and

Supplementary Figure 9) and various network parameters and architectures tested

(**Supplementary Figure 10 to Supplementary Figure 14**). Based on these extensive tests, we

use did not use data balancing, but ensured the distribution of age in each training fold was

similar with filter width = 9 samples, filter depth = 6 channels, filter number = 16, network depth

= 3, with a mini-batch size of 128 samples for our training dataset (D1; 18 channel EEG). For

training and testing on 2-channel EEG derivations, further optimization of network parameters

yielded a filter depth = 2 channels, network depth = 2, filter number = 2, with the remaining

parameters consistent with our training set. These additional robustness tests confirmed that the

bipolar montage with 60 s epochs offer optimal FBA estimates.

To explore how the EEG signal was represented by the neural network, we extracted activations

of network layers at several stages of the deep learning network when trained and tested on all

available data (for this exploratory analysis no cross-validation was used). Here, the optimal

network was composed of 63 layers, where an early layer output was taken at layer 6, the middle

layer output was taken at layer 29 and the late layer output was extracted at layer 61. A Uniform

Manifold Approximation and Project method (UMAP (*39*)) was used to reduce the high-

dimensional network activation space into a lower dimensional 2D space (UMAP1, UMAP2),

wherein the output space was coded according to age. The use of UMAP here enables network

activations to be condensed into low-dimensional manifolds (UMAP1, UMAP2), while retaining

the structural similarity of higher dimensional activations. Visualizations via UMAP allowed us

to qualitatively assess how the activations of a network layer cluster with respect to age (**Figure**

3b). To quantify the association between UMAP1 and UMAP2 dimensions and age, we used

GPR with UMAP1 and UMAP2 as inputs to predict age (via 10-fold cross-validation). The

accuracy of age prediction for each network layer (**Figure 3b**) was then derived. We used the

same approach to track how UMAP representations of network activations were linked to

individual features of the EEG (**Supplementary Figure 4 and Supplementary Figure 5b**).

For GPR, hyperparameters were selected using Bayesian optimization within a nested cross-

validation and included kernel function (e.g. rational quadratic) and sigma values. Shapley

values (*77*) were also computed to quantify each feature in terms of its contribution to the overall

prediction of FBA (**Supplementary Figure 5c**), extending on the notion of linear model

predictions.

Statistical analyses

We first evaluated the prediction accuracy of FBA between residual neural network and

Gaussian process regressions. The key variable used in this analysis was the residual error —

also referred to as the predicted age difference, PAD (the difference in years between the FBA

and chronological age). Comparisons were made using the non-parametric Mann-Whitney rank

sum test.

We then analyzed potential confounding effects of age and sex on PAD using multiple linear

regression. Here, our multiple regression models for D1 and D2 first tested the effect of sex and

age (Eq.3).

$$PAD \sim \beta_0 + \beta_1 age + \beta_2 sex + \beta_1 age * sex + \beta_1 age^2 \qquad \text{(Eq.3)}$$

We also assessed the performance of the FBA across spatial and temporal combinations of the

EEG. This included examining changes in prediction accuracy for limited 2-channel FBAs based

on bilateral channels of the bipolar montage. A FBA was trained and tested for each bilateral

channel and then examined with respect to its designation across the scalp to see if particular

regions of the brain were associated with higher accuracy. As a further adversarial test, we also

tested variations of channel laterality and channel location in 18 channel D1 data, to gauge

changes to performance accuracy with respect to the spatial location of EEG channels. Potential

differences in channel laterality were assessed by swapping left hemisphere EEG channels

swapped with right hemisphere EEG channels prior to training and testing. The same procedure

was used to assess differences in anteroposterior directions, where anteriorly positioned (e.g.

frontal) electrodes were swapped with posteriorly positioned electrodes (i.e. parietal, occipital)

prior to training and testing. We also examined the temporal variation in FBA across all available

epochs, i.e., from the start to the end of the recording period to observe optimal times to evaluate

FBAs during N1 to N2 transitions.

We also performed additional sleep stage-specific tests to ascertain the performance accuracy of

FBAs across wake, N2, N3 and REM sleep. Using Res-NN and GPR models, we estimated

FBAs for all available EEG epochs for subjects in D2 (based on availability of sleep states). Age

prediction within N2 sleep states were found to be the most accurate within our data

(**Supplementary Table 3**).

Validation: We re-trained models using all available data in D1 and performed an out-of-sample

validation on EEG data from D2. As D2 contained a limited number of recording electrodes, we

used the same electrodes when training on D1 in a bipolar configuration (F4-C4 and C4-O2).

Furthermore, only periods of N2 of D1 (10 minute segments) were included in the training

dataset. We compared the mean absolute error between predicted and chronological age from the

10-fold cross-validation results from D1 to D2 to determine if the accuracy of the FBA trained

on D1 was preserved when applied to D2. Mann-Whitney rank sum tests were used to compare

development versus validation errors. All validation analyses performed for the Res-NN model

were repeated for the GPR model (**Supplementary Figure 15**). **Supplementary Table 4**

summarizes model performances across test folds for all datasets, respectively.

To test site differences in EEG recordings (**Supplementary Figure 5c**), we compared the

distribution of EEG feature values using Kolmogorov-Smirnov tests corrected for multiple

comparisons (Bonferroni's method). We used a small random sampling of EEG features within

several age defined bins (bin width was 1 year) to minimize the effect of age on feature

distribution. Here, differences between EEG features per site and across age were denoted by the

number of times the null hypothesis (EEG features between sites were from the same

distribution) was rejected across 1000 samplings of each bin ($n = 30$ samples without

replacement per feature per bin). The comparisons were also encoded to delineate if the

distributional difference in EEG feature resulted in values greater than expected (older

appearing) or less than expected (younger appearing) when comparing D1 to D2. At an

acquisition level, we also estimated the total (summed) band power of the raw EEG signal,

between 70 and 80 Hz, using Welch's power density spectral estimate, to determine whether

significant differences in the high frequency noise floor were present across sites (unpaired t-

test).

Growth Charts: We generated growth charts for the FBA: a chart based on the full 10-20

International System (D1) and a limited 2-channel EEG growth chart for sleep recordings based

on the AASM criteria, in a combined D1 and D2 dataset (2-channel EEG). These charts are

available in our GitHub repository (see *Data Availability* section). Centiles were estimated using

generalized additive models with a protocol similar to the WHO (*4*) child growth charts for

height and weight for age. Here, we used the GAMLSS package (*78*) in RStudio (version

1.4.1717). These centiles were optimized by a Box-Cox-power exponential distribution with a

cubic spline smoothing function, with distribution GAMLSS parameters *sigma* set as a cubic

spline fit over age (df = 3) with *nu* = 1 and *tau* = 1. Centiles were estimated at the 3^{rd}, 15^{th}, 50^{th},

85^{th} and 97^{th} centiles as per WHO guidelines.

We also compared the performance of our FBA chart with pediatric head circumference and

height measures. Using respective reference centiles across the pediatric ranges (*41, 42*), we

generated 1000 simulated training and testing cohorts from similar distributions of age, sex and

cohort size to the combined D1 and D2 dataset. Using 10-fold cross-validation, we then trained

and tested a GPR model with head circumference or height as input, and evaluated the results

using MAE and wMAE.

To evaluate the FBA as a tool to determine group-wide differences amongst clinical cohorts, we

included an additional cohort of children from D2 who were diagnosed with Trisomy 21 (*n* =

40). This group of children were chosen as they are known to be predisposed to

neurodevelopmental delays. To test this 'proof-of-concept', we examined differences between

groups using unpaired t-tests, where variables were checked to ensure normality (*79*) (confirmed

via a Lilliefors test (*80*)). The null hypothesis tested was that children with Trisomy 21 will have

a PAD that is not different from children with typical neurodevelopment.

All statistical tests employed for analyses were two-sided and the level of significance was 0.05.

References

1. J. Jylhävä, N. L. Pedersen, S. Hägg, Biological age predictors. *EBioMedicine* **21**, 29-36 (2017).
2. J. H. Cole, R. E. Marioni, S. E. Harris, I. J. Deary, Brain age and other bodily 'ages': implications for neuropsychiatry. *Molecular psychiatry* **24**, 266-281 (2019).
3. J. H. Cole *et al.*, Brain age predicts mortality. *Molecular psychiatry* **23**, 1385-1392 (2018).
4. W. H. Organization, W. H. O. N. f. Health, *WHO child growth standards: growth velocity based on weight, length and head circumference: methods and development*. (World Health Organization, 2009).
5. T. Kaufmann *et al.*, Common brain disorders are associated with heritable patterns of apparent aging of the brain. *Nature neuroscience* **22**, 1617-1623 (2019).
6. D. W. Belsky *et al.*, Quantification of the pace of biological aging in humans through a blood test, the DunedinPoAm DNA methylation algorithm. *Elife* **9**, e54870 (2020).
7. D. Karasik, S. Demissie, L. A. Cupples, D. P. Kiel, Disentangling the genetic determinants of human aging: biological age as an alternative to the use of survival measures. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences* **60**, 574-587 (2005).
8. J. Van Dongen *et al.*, Genetic and environmental influences interact with age and sex in shaping the human methylome. *Nature communications* **7**, 1-13 (2016).
9. J. Rutledge, H. Oh, T. Wyss-Coray, Measuring biological age using omics data. *Nature Reviews Genetics*, 1-13 (2022).
10. Y. E. Tian *et al.*, Biological aging of human body and brain systems. *medRxiv*, (2022).
11. R. A. Bethlehem *et al.*, Brain charts for the human lifespan. *Nature* **604**, 525-533 (2022).
12. V. M. Bashyam *et al.*, MRI signatures of brain age and disease over the lifespan based on a deep brain network and 14 468 individuals worldwide. *Brain* **143**, 2312-2324 (2020).
13. K. Franke, C. Gaser, Ten years of BrainAGE as a neuroimaging biomarker of brain aging: What insights have we gained? *Frontiers in neurology*, 789 (2019).
14. S. M. Smith, D. Vidaurre, F. Alfaro-Almagro, T. E. Nichols, K. L. Miller, Estimation of brain age delta from brain imaging. *Neuroimage* **200**, 528-539 (2019).
15. S. Shahab *et al.*, Brain structure, cognition, and brain age in schizophrenia, bipolar disorder, and healthy controls. *Neuropsychopharmacology* **44**, 898-906 (2019).
16. J. Lee *et al.*, Deep learning-based brain age prediction in normal aging and dementia. *Nature Aging* **2**, 412-424 (2022).
17. D. Beck *et al.*, Cardiometabolic risk factors associated with brain age and accelerate brain ageing. *Human brain mapping* **43**, 700-720 (2022).
18. A.-M. G. De Lange *et al.*, Multimodal brain-age prediction and cardiovascular risk: The Whitehall II MRI sub-study. *NeuroImage* **222**, 117292 (2020).
19. Y. Chen, T. Z. Baram, Toward understanding how early-life stress reprograms cognitive and emotional brain networks. *Neuropsychopharmacology* **41**, 197-206 (2016).
20. M. d. Onis *et al.*, Development of a WHO growth reference for school-aged children and adolescents. *Bulletin of the World health Organization* **85**, 660-667 (2007).
21. A. B. Archibald, J. A. Graber, J. Brooks-Gunn, Pubertal processes and physiological growth in adolescence. *Blackwell handbook of adolescence*, 24-47 (2003).
22. R. M. Malina, C. Bouchard, O. Bar-Or, *Growth, maturation, and physical activity*. (Human kinetics, 2004).
23. D. E. Astle, J. Holmes, R. Kievit, S. E. Gathercole, Annual Research Review: The transdiagnostic revolution in neurodevelopmental disorders. *Journal of Child Psychology and Psychiatry* **63**, 397-417 (2022).
24. B. Tunç *et al.*, Deviation from normative brain development is associated with symptom severity in autism spectrum disorder. *Molecular autism* **10**, 1-14 (2019).
25. G. Ball, C. E. Kelly, R. Beare, M. L. Seal, Individual variation underlying brain age estimates in typical development. *Neuroimage* **235**, 118036 (2021).
26. C. Kelly *et al.*, Investigating brain structural maturation in children and adolescents born very preterm using the brain age framework. *NeuroImage* **247**, 118828 (2022).
27. P. J. Anderson, A. Burnett, Assessing developmental delay in early childhood—concerns with the Bayley-III scales. *The Clinical Neuropsychologist* **31**, 371-381 (2017).

809   28.   B. Kolb, A. Harker, R. Gibb, Principles of plasticity in the developing brain. *Developmental Medicine &*
810   *Child Neurology* **59**, 1218-1223 (2017).
811   29.   S. W. MacDonald, L. Nyberg, L. Bäckman, Intra-individual variability in behavior: links to brain structure,
812   neurotransmission and neuronal activity. *Trends in neurosciences* **29**, 474-480 (2006).
813   30.   S. K. Yadav *et al.*, Genetic variations influence brain changes in patients with attention-deficit
814   hyperactivity disorder. *Translational Psychiatry* **11**, 349 (2021).
815   31.   J. O. Edgin, Cognition in Down syndrome: a developmental cognitive neuroscience perspective. *Wiley*
816   *Interdisciplinary Reviews: Cognitive Science* **4**, 307-317 (2013).
817   32.   C. A. Nelson, J. P. McCleery, Use of event-related potentials in the study of typical and atypical
818   development. *Journal of the American Academy of Child & Adolescent Psychiatry* **47**, 1252-1261 (2008).
819   33.   A. J. Anderson, S. Perone, Developmental change in the resting state electroencephalogram: Insights into
820   cognition and the brain. *Brain and cognition* **126**, 40-52 (2018).
821   34.   M. Angriman, B. Caravale, L. Novelli, R. Ferri, O. Bruni, Sleep in children with neurodevelopmental
822   disabilities. *Neuropediatrics* **46**, 199-210 (2015).
823   35.   C. J. Chu, J. Leahy, J. Pathmanathan, M. Kramer, S. S. Cash, The maturation of cortical sleep rhythms and
824   networks over early development. *Clinical Neurophysiology* **125**, 1360-1370 (2014).
825   36.   A. F. Marquand *et al.*, Conceptualizing mental disorders as deviations from normative functioning.
826   *Molecular psychiatry* **24**, 1415-1424 (2019).
827   37.   S. Rutherford *et al.*, The normative modeling framework for computational psychiatry. *Nature Protocols*,
828   1-24 (2022).
829   38.   N. J. Stevenson *et al.*, Automated cot□side tracking of functional brain age in preterm infants. *Annals of*
830   *clinical and translational neurology* **7**, 891-902 (2020).
831   39.   L. McInnes, J. Healy, J. Melville, Umap: Uniform manifold approximation and projection for dimension
832   reduction. *arXiv preprint arXiv:1802.03426*, (2018).
833   40.   M. De Onis, A. W. Onyango, WHO child growth standards. *The Lancet* **371**, 204 (2008).
834   41.   A. F. Roche, D. Mukherjee, S. Guo, W. M. Moore, Head circumference reference data: birth to 18 years.
835   *Pediatrics* **79**, 706-712 (1987).
836   42.   R. J. Kuczmarski, *2000 CDC Growth Charts for the United States: methods and development*.
837   (Department of Health and Human Services, Centers for Disease Control and …, 2002).
838   43.   M. Eisermann, A. Kaminska, M.-L. Moutard, C. Soufflet, P. Plouin, Normal EEG in childhood: from
839   neonates to adolescents. *Neurophysiologie Clinique/Clinical Neurophysiology* **43**, 35-65 (2013).
840   44.   A. R. Clarke, R. J. Barry, R. McCarthy, M. Selikowitz, Age and sex effects in the EEG: development of the
841   normal child. *Clinical neurophysiology* **112**, 806-814 (2001).
842   45.   A. Kaminska, M. Eisermann, P. Plouin, Child EEG (and maturation). *Handbook of clinical neurology* **160**,
843   125-142 (2019).
844   46.   M. J. Lund *et al.*, Brain age prediction using fMRI network coupling in youths and associations with
845   psychiatric symptoms. *NeuroImage: Clinical* **33**, 102921 (2022).
846   47.   M. M. Vandenbosch, D. van't Ent, D. I. Boomsma, A. P. Anokhin, D. J. Smit, EEG□based age□prediction
847   models as stable and heritable indicators of brain maturational level in children and adolescents. *Human*
848   *brain mapping* **40**, 1919-1926 (2019).
849   48.   S. I. Dimitriadis, C. I. Salis, Mining time-resolved functional brain graphs to an EEG-based chronnectomic
850   brain aged index (CBAI). *Frontiers in human neuroscience* **11**, 423 (2017).
851   49.   O. Al Zoubi *et al.*, Predicting age from brain EEG signals—A machine learning approach. *Frontiers in*
852   *aging neuroscience* **10**, 184 (2018).
853   50.   D. A. Engemann *et al.*, A reusable benchmark of brain-age prediction from M/EEG resting-state signals.
854   *NeuroImage*, 119521 (2022).
855   51.   H. Sun *et al.*, Brain age from the electroencephalogram of sleep. *Neurobiology of aging* **74**, 112-120
856   (2019).
857   52.   G. Z. Tau, B. S. Peterson, Normal development of brain circuits. *Neuropsychopharmacology* **35**, 147-168
858   (2010).
859   53.   A. Lau-Zhu, M. P. Lau, G. McLoughlin, Mobile EEG in research on neurodevelopmental disorders:
860   Opportunities and challenges. *Developmental cognitive neuroscience* **36**, 100635 (2019).
861   54.   N. Hayashi-Kurahashi *et al.*, EEG for predicting early neurodevelopment in preterm infants: an
862   observational cohort study. *Pediatrics* **130**, e891-e897 (2012).
863   55.   T. J. Silk, A. G. Wood, Lessons about neurodevelopment from anatomical magnetic resonance imaging.
864   *Journal of Developmental & Behavioral Pediatrics* **32**, 158-168 (2011).

56. L. Novelli *et al.*, Mapping changes in cortical activity during sleep in the first 4 years of life. *Journal of*
Sleep Research **25**, 381-389 (2016).
57. V. Jaramillo *et al.*, Characterization of overnight slow-wave slope changes across development in an age-,
amplitude-, and region-dependent manner. *Sleep* **43**, zsaa038 (2020).
58. M. Grigg-Damberger *et al.*, The visual scoring of sleep and arousal in infants and children. *Journal of*
Clinical Sleep Medicine **3**, 201-240 (2007).
59. J. L. Moore, D. Z. Carvalho, E. K. St Louis, C. Bazil, Sleep and epilepsy: a focused review of
pathophysiology, clinical syndromes, co-morbidities, and therapy. *Neurotherapeutics* **18**, 170-180 (2021).
60. J. Carpay *et al.*, The diagnostic yield of a second EEG after partial sleep deprivation: a prospective study in
children with newly diagnosed seizures. *Epilepsia* **38**, 595-599 (1997).
61. A. Buchmann *et al.*, EEG sleep slow-wave activity as a mirror of cortical maturation. *Cerebral Cortex* **21**,
607-615 (2011).
62. J. W. Britton *et al.*, Electroencephalography (EEG): an introductory text and atlas of normal and abnormal
findings in adults, children, and infants. (2016).
63. L. Cornelissen, S.-E. Kim, P. L. Purdon, E. N. Brown, C. B. Berde, Age-dependent electroencephalogram
(EEG) patterns during sevoflurane general anesthesia in infants. *elife* **4**, e06513 (2015).
64. P. Laoprasert, *Atlas of Pediatric EEG*. (McGraw-Hill, 2011), vol. 1.
65. J. M. Lee *et al.*, A prospective study of age-dependent changes in propofol-induced electroencephalogram
oscillations in children. *Anesthesiology* **127**, 293-306 (2017).
66. T. Hahn *et al.*, From 'loose fitting'to high-performance, uncertainty-aware brain-age modelling. *Brain* **144**,
e31-e31 (2021).
67. J. Zhang *et al.*, Moving towards vertically integrated artificial intelligence development. *NPJ digital*
medicine **5**, 1-9 (2022).
68. J. Zhang *et al.*, An interactive dashboard to track themes, development maturity, and global equity in
clinical artificial intelligence research. *The Lancet Digital Health* **4**, e212-e213 (2022).
69. P. C. M. Koolschijn, E. A. Crone, Sex differences and structural brain maturation from childhood to early
adulthood. *Developmental cognitive neuroscience* **5**, 106-118 (2013).
70. R. K. Lenroot, J. N. Giedd, Sex differences in the adolescent brain. *Brain and cognition* **72**, 46-55 (2010).
71. A. Zhang, L. Xing, J. Zou, J. C. Wu, Shifting machine learning for healthcare from development to
deployment and from models to data. *Nature Biomedical Engineering*, 1-16 (2022).
72. R. B. Berry *et al.*, The AASM manual for the scoring of sleep and associated events. *Rules, Terminology*
and Technical Specifications, Darien, Illinois, American Academy of Sleep Medicine **176**, 2012 (2012).
73. S. Ghirardo, A. Amaddeo, L. Griffon, S. Khirani, B. Fauroux, Central apnea and periodic breathing in
children with underlying conditions. *Journal of Sleep Research* **30**, e13388 (2021).
74. K. He, X. Zhang, S. Ren, J. Sun, in *Proceedings of the IEEE conference on computer vision and pattern*
recognition. (2016), pp. 770-778.
75. C. E. Rasmussen, C. Williams, Gaussian processes for machine learning. *MIT Press. Cambridge, MA* **32**,
68 (2006).
76. C. Szegedy *et al.*, in *Proceedings of the IEEE conference on computer vision and pattern recognition*.
(2015), pp. 1-9.
77. E. Winter, The shapley value. *Handbook of game theory with economic applications* **3**, 2025-2054 (2002).
78. M. D. Stasinopoulos, R. A. Rigby, G. Z. Heller, V. Voudouris, F. De Bastiani, *Flexible regression and*
smoothing: using GAMLSS in R. (CRC Press, 2017).
79. J. B. Burbidge, L. Magee, A. L. Robb, Alternative transformations to handle extreme values of the
dependent variable. *Journal of the American Statistical Association* **83**, 123-127 (1988).
80. H. W. Lilliefors, On the Kolmogorov-Smirnov test for normality with mean and variance unknown.
Journal of the American statistical Association **62**, 399-402 (1967).

Acknowledgements

We would like to acknowledge and thank Dr Paula Sanz-Leon for technical assistance with the deployment of functional brain age software provided on our GitHub repository. For Dataset 1, we thank Mr Jarmo Mäki and other EEG technicians within the HUS diagnostic center in annotating EEGs, as well as Mr Eero Ahtola for assisting with the preprocessing of EEG data. For Dataset 2, we thank all technicians, nurses and doctors involved with data collection, overnight annotations and clinical assessment of polysomnography recordings conducted at the Respiratory and Sleep Medicine Department located within the Queensland Children's hospital.

923 **Funding:**
924
925   National Health and Medical Research Council (NHMRC) Grant no. 2002135
926
927   The HUS Children's Hospital/HUS diagnostic center research funds
928
929   Finnish Academy (335788, 332017),
930
931   Finnish Pediatric Foundation (Lastentautiensäätiö)
932
933   Sigrid Juselius Foundation,
934
935   QIMR Berghofer Medical Research Institute, Early Career Seed Funding grant.
936
937 **Author contributions:**
938
Conceptualization: KKI, JAR, MW, SV, NJS

Methodology: KKI, MW, LL, SV, NJS

Investigation: KKI, SV, NJS

Visualization: KKI, NJS

Funding acquisition: MW, SV, NJS

Project administration: KKI, LL, SV, NJS

Supervision: LL, SV, NJS

Writing – original draft: KKI, JAR, MW, LL, SV, NJS

Writing – review & editing: KKI, JAR, MW, JC, SJV, AK, LMH, LL, SV, NJS
948

Competing interests: None

Data and materials availability:

Trained Res-NN and GPR models for age prediction are available on our GitHub repository,

accessible at: https://github.com/brain-modelling-group/functional-brain-age. Data used to train

and validate the FBA will be made available to researchers on reasonable request to S.V., N.J.S.

K.I.: data sharing is subject to a material transfer agreement, approved by the legal departments

of the requesting researcher and by all legal departments of the institutions that provided data

and ethical clearances for the study. EEG feature matrices (z-scored) and subjects ages

(quantized by 3 monthly intervals for ages 1 month to 4 years, and 1 year intervals for ages 5 and

beyond) for each EEG recording in our training dataset (D1, 18 channel and 2 channel versions)

and external validation set (D2, 2 channel version) are also available in our GitHub repository.

These files have been pseudonymized.

MATLAB and Python code for the estimation of functional brain age by Res-NN based methods

on 18- and 2-channel EEG are available on our GitHub repository.

964
965

Supplementary Material

Supplementary Table 1 – Performance of individual EEG features in predicting age following Gaussian process regression (GPR) and 10-fold cross-validation of training data (D1). Results are based on 60 second EEG epochs acquired in the bipolar montage. Amplitude metrics (ID1 to ID7) denote the 5th, 50th (median), 95th centiles based on the absolute value of the Hilbert transform of the EEG (amplitude envelope), along with four central moments of the signal: mean, standard deviation, skewness and kurtosis. Frequency metrics (ID8 to ID19) summarize frequency domain metrics based on the power spectrum density of the EEG signal, and included features such as peak frequency, signal amplitude at this peak, along with mean frequency, bandwidth and relative spectral power in common EEG oscillatory bands. Informational metrics (ID20 to ID32) summarized various non-linear and statistical properties of the EEG. SNLEO stands for the smoothed non-linear energy operator.

		GPR			Shapley	
Feature ID	EEG Features	R^2	RMSE	MAE	Value	Rank
1	5th percentile Amplitude Envelope	0.802	1.698	1.312	0.524	1
2	50th percentile amplitude envelope	0.802	1.700	1.341	0.246	21
3	95th percentile amplitude envelope	0.755	1.832	1.532	0.069	14
4	Mean amplitude envelope	0.773	1.784	1.460	0.187	22
5	Standard deviation amplitude envelope	0.742	1.865	1.577	0.051	26
6	Skewness amplitude envelope	0.183	1.664	3.021	0.113	7
7	Kurtosis amplitude envelope	0.307	2.002	2.714	0.311	31
8	Peak Frequency (PSD)	0.084	1.146	3.298	0.110	17
9	Power in peak frequency	0.145	1.537	3.236	0.044	2
10	Signal amplitude of peak frequency component	0.781	1.762	1.500	0.185	13
11	Mean frequency	0.555	2.111	2.200	0.023	19
12	Bandwidth	0.581	2.101	2.079	0.159	4
13	Relative spectral power (delta 1, 0.5-2Hz)	0.194	1.699	3.153	0.231	10
14	Relative spectral power (delta 2, 2-4Hz)	0.401	2.091	2.322	0.401	15
15	Relative spectral power (theta, 4-8Hz)	0.253	1.842	2.890	0.181	20
16	Relative spectral power (alpha, 8-12Hz)	0.590	2.100	1.978	0.036	12
17	Relative spectral power (beta, 12-30Hz)	0.620	2.073	1.951	0.262	30
18	Total power in all bands	0.763	1.809	1.488	0.075	25
19	Spectral slope (decay in power spectrum)	0.572	2.109	2.093	0.204	6
20	Sample Entropy	0.547	2.119	2.233	0.172	32
21	Fractal Dimension (Higuchi)	0.573	2.101	2.175	0.523	8
22	Spectral entropy	0.568	2.110	2.169	0.372	18
23	Spectral difference	0.786	1.735	1.438	0.031	24
24	Hjorth 1 (Activity)	0.768	1.795	1.478	0.073	3

25	Hjorth 2 (Mobility)	0.803	1.693	1.443	0.115	27
26	Hjorth 3 (Complexity)	0.827	1.613	1.324	0.361	5
27	SNLEO mean	0.506	2.150	2.063	0.068	9
28	SNLEO standard deviation	0.408	2.090	2.402	0.007	16
29	Burst shape skewness (all bursts)	0.073	1.124	3.328	0.024	23
30	Burst shape kurtosis (all bursts)	0.207	1.712	2.912	0.118	29
31	Burst duration (mean)	0.677	1.989	1.801	0.305	11
32	Burst duration (standard deviation)	0.536	2.124	2.299	0.112	28

976
977

Supplementary Figure 1 – Comparative performance of FBA estimation from a GPR model with the Res-NN **a.** The Res-NN model (R^2=0.96, MAE=0.57 years) with error bounds of ±1.5 years and **b.** The residual error (PAD), in years. **c.** GPR results based on a multivariable model of EEG features (R^2=0.93, MAE=0.79 years). **d.** Residual error (PAD), in years, for the GPR model. Individual EEG recordings (blue filled circles) are plotted across ages.

984
985
Supplementary Figure 2 - *The general architecture of the residual neural network used in this work.* For inception

networks, we used elements of the inceptionv3 network in MATLAB with a final regression layer. The scale of this

pre-trained network was too large to accurately train on the dataset used in our work, so we tested architectural

aspects of the network rather than the entire network. Similarly to the residual neural network we added variability

by changing the temporal filter width (FW), filter channel depth (FD) and filter number (FN) within the

convolutional layers. We used the file generate_networks_v2.m to generate networks with different configurations

and architectures (see the GitHub page for more details).

993
994
995

996
997

Supplementary Figure 3 – *Individual EEG features versus age (all subjects from training dataset D1).* As per Supplementary Table 1, each EEG feature and their respective ID are based on 60 epochs from the bipolar montage. Feature types are indicated by colors, where amplitude features (reddish hue), frequency domain features (green hue) and informational metrics (blue hue). A spline is fitted across the median values derived across 2 yearly age bins.

1003

1004

1005   ***Supplementary Figure 4** – Relating individual EEG features to the late network layer of the Res-NN model.* For

1006   each individual feature we used GPR to assess the strength of correlation between individual EEG features with

1007   network activation layers (here the late layer, i.e., layer 61 from 2-channel EEG data from D1 is shown) to derive a

1008   predicted feature value based on UMAP1 and UMAP2 dimensions derived from this training phase. The strength of

1009   correlation (R^2 value) indicates how individual EEG features and predicted individual EEG features are linked by

UMAP representations of the Res-NN model, following 10-fold cross-validation. The color bar indicates the relative range of the EEG feature pertaining to its minimum and maximum values (e.g. for feature ID4, low mean amplitude versus high mean amplitude).

Supplementary Figure 5 - *Validation of the FBA across sites.* **a.** Differences in MAE between D1 and D2: cross-validation (train and test D1 CrossVal, indicated by blue circles) versus external validation on D2 (train D1 test D2, red circles). The MAE across age is higher when testing D1's model on D2. The MAE is shown with error bars indicating the standard deviation. **b.** Mapping EEG features onto the late layer network UMAP (layer 61) derived from 2-channel D1 data; here we show the EEG median amplitude as an exemplar. The R^2 value indicates the strength of correlation between UMAP values and feature value, based on a GPR prediction. **c.** Differences between EEG features and site across age. EEG features (IDs 1 to 32, see also *Supplementary Table 1* for feature names) were ordered by Shapley values (highest to lowest) to indicate the relative contribution of the feature to the overall

model. Red and blue colors indicate significant age biases following multiple comparisons correction (Bonferroni's

method). A positive age bias percentage (red) indicates EEG feature values being 'older' in D2 compared to D1 and

a negative age bias percentage (blue) indicates EEG feature values being 'younger' in D2 compared to D1.

Approximately 37% of EEG features/age bins combinations differed significantly across age and sites with 191 out

of 512 hypothesis tests meeting significance at $p<0.05$ following correction for multiple comparisons. See methods

for further details. **d.** Fractal dimension and relative delta power (0.5 to 2 Hz) are exemplar EEG features that show

site related differences (D1=blue; D2=maroon).

1031
1032

1033 ***Supplementary Table 2** – Overall performance of FBA across datasets.* The performances of the FBA, using a GPR
1034 model, across all training, test, and combined datasets. MAE, wMAE and RMSE are shown. CI is confidence
1035 interval, [a]only N2 from D1 used, [b]10-fold cross-validation.

Train	# channels, # recordings, # epochs	Test	# channels, # recordings, # epochs	MAE (in years; 95CI)	wMAE (in years; 95CI)	RMSE (in years)
D1	19, 1062, 30798	D1[b]	19, 1062, 30798	0.79 (0.74 – 0.84)	1.10 (0.91 – 1.29)	1.09
D1	2, 1062, 20178	D1[b]	2, 1062, 20178	0.93 (0.88 – 0.99)	1.25 (1.02 – 1.51)	1.24
D1[a]	2, 1062, 20178	D2	2, 723, 13737	1.96 (1.83 – 2.06)	2.55 (2.17 – 2.85)	1.35
D2	2, 723, 13737	D2[b]	2, 723, 13737	1.25 (1.17 – 1.31)	1.34 (1.11 – 1.57)	1.41
D1+D2	2, 1792, 33915	D1+D2[b]	2, 1792, 33915	1.08 (1.04 – 1.12)	1.30 (1.14 – 1.47)	1.36

1036

1037

1038

1039

1040   *Supplementary Figure 6 – FBA growth charts stratified into infant and childhood age groups.* **a.** FBA charts for

1041   infants (0 to 2 years) and **b.** children and adolescents (2 to 18 years) were generated to offer a magnified view of the

1042   FBA with respect to chronological age. In both age-stratified versions of the chart, over 95% of the data (blue dotted

1043   points) are captured between the 3rd and 97th centiles estimated. The 3rd (red), 15th (yellow), 50th (green), 85th

1044   (purple) and 97th (blue) centiles are indicated.

1045

1046

1047
1048
1049   *Supplementary Figure 7 -* The effect of epoch length and montage on cross-validated RMSE on a random selection

1050   of 7 network architectures. An epoch length of 60s and the bipolar montage were selected *a priori*. Different EEG

1051   epoch durations (30 s, 60 s, 120 s, 300 s) and EEG montages (referential, bipolar, average, current source density)

1052   were tested. The root mean square error between predicted age (FBA) and age across all testing data from a 10-fold

1053   cross-validation was used to determine the optimal selection, with a minimum RMSE indicating the optimal results.

1054
1055

1056
1057
Supplementary Figure 8 - The effect of solver and mini-batch size on cross-validated RMSE on a random selection

of network architectures, epoch length, and montage ($n = 8$ per group for solver and $n = 7$ per group for mini-batch

size). For training the neural networks, two hyper-parameters of training were selected based on an initial run on

dataset D1 (Solver Type and Mini-batch size). A random selection of network architectures were made and the

network was trained and tested within a 10-fold cross-validation with the RMSE on the accumulated left-out test

data used to select the hyper-parameters. The ADAM solver and a mini-batch size of 128 samples were selected.

1064

Supplementary Note 1 - **The effect of data balancing on NN results**

Balancing the dataset is an important component of constructing viable training datasets. In this case, the number of

recording per age bin should be equal for the best general regression results; increased diversity of data (typically by

adding more data) also provides an improved fit. Furthermore, the training data distribution should best represent the

distribution of the larger population. We show the effect of data balancing by training 10 randomly selected network

architectures with training datasets containing different levels of heterogeneity with respect to age (***Supplementary***

Figure 9). To ensure the number of training points is equal we extract more epochs per recording as the dataset

becomes more balanced by changing the overlap of epoch extraction. We select three levels of balance: balanced

(uniform distribution of subjects across age) extracts at most 40 recordings from each age group (ages are grouped at

yearly intervals), Unbalanced1 (more samples at early ages) extracts at most 80 recordings from each age group, and

Unbalanced2 (most samples at early ages) extracts at most 160 recordings from each age group. The RMSE and

weighted RMSE (an age adjusted RMSE) are calculated on all EEG recordings that were left out during the 10-fold

cross-validation (an unbalanced cohort).

1078

1079
1080
1081   ***Supplementary Figure 9*** *- The general effects of data balancing on cross-validated RMSE for 18-channel data (D1*

1082   *datasets).* The left plot is the RMSE and the right plot is the wRMSE which is an RMSE averaged across age bins

1083   (yearly) rather than per EEG recording. In both cases, the increased diversity in recordings associated with

1084   unbalanced training dataset offers lower errors, although the reduction in error is smaller when considering wRMSE.

1085
Supplementary Figure 10 *- The general effects of data network architecture type (Residual vs Inception) on cross-*

validated RMSE for 18-channel data (D1 datasets). Once general EEG parameters and network training hyper-

parameters were initialized, we tested two general architectures based on residual neural networks and inception

based neural networks. We randomly selected 10 networks based on both architectures and computed RMSE. Based

on these parameter evaluations, residual networks were selected as the most suitable candidate architecture for

estimating age.

1092
1093

Supplementary Note 2 – Testing various of network parameters (18 channel FBA for D1)

We also tested parameters of filter width, filter depth, network depth and filter number within the residual network.

The optimal combination was selected using the internal validation data, averaged across all folds. The latter was

performed to ensure only 1 network architecture was used across fold. A fixed grid optimization search across was

performed across an array of network parameters. For the 18-channel FBA based on dataset D1, this resulted in 180

different network architectures (***Supplementary Figure 8***). The general trends are shown in Figure 4 with an optimal

combination of network parameters selected: a filter width of 9 samples ($n = 45$ per group), filter depth of 6 channels

($n = 36$ per group), network depth of 3 ($n = 60$ per group), and a filter number of 16 ($n = 60$ per group): a total of

867,601 learnable parameters.

1103
1104   ***Supplementary Figure 11 -*** The general effects of residual network parameters on cross-validated RMSE for 18-

1105   channel data (D1 dataset). Internal single fold cross-validation averaged across all 10 training folds selected. Note

1106   that, as filter depth of 1 channel had a considerably higher RMSE than other filter depths, RMSE points associated

1107   with a 1 channel network were removed before generating the remaining boxplots for visualization purposes.

1108
1109   ***Supplementary Figure 12 -*** *The general effects of residual network parameters on cross-validated RMSE for a 2-*

1110   *channel FBA based on dataset D1.* This resulted in 60 different network architectures. The optimal combination of

1111   network parameters selected were a filter width of 9 samples ($n = 18$ per group), filter depth of 2 channels ($n = 30$

1112   per group), network depth of 2 ($n = 30$ per group), and a filter number of 2 ($n = 20$ per group): a total of 906,657

1113   learnable parameters.

1114
1115 ***Supplementary Figure 13 -*** *The general effects of residual network parameters on cross-validated RMSE for 2-*

1116 *channel data (D2 dataset).* This resulted in 90 different network architectures. The optimal parameters were a filter

1117 width of 9 samples ($n = 18$ per group), filter depth of 2 samples ($n = 45$), network depth of 2 ($n = 30$ per group), and

1118 a filter number of 32 ($n = 30$ per group): a total of 906,657 learnable parameters.

1119
Supplementary Figure 14 - *The general effects of network parameters on cross-validated RMSE for 2-channel data*

(D1 and D2 combined). This resulted in 45 different network architectures. The RMSE of an internal single fold

cross-validation, average across all 10 training folds selected a filter width of 9 samples ($n = 9$ per group), a network

depth of 2 ($n = 45$ per group), network depth of 2 ($n = 15$ per group), and a filter number of 32 ($n = 15$ per group): a

total of 906,657 learnable parameters. Note, we did not vary the filter depth/channel number as previous

experiments on D1 and D2 alone showed that 2 channels was optimal.

1126

1127

Supplementary Figure 15 - *Summary plot of GPR model performance*. **a.** The mean absolute error (MAE) of the

GPR (dark blue triangles, left axis) across 2 yearly age bins and the relative accuracy of PAD (in blue, right axis)

across 2 yearly age bins. For both plots, the mean and standard error of the mean (SEM) are plotted to reflect the

sample distribution within age bins. **b.** Differences in MAE of D1 following cross-validation (D1 CrossVal) versus

external validation on D2 (D1 test on D2) for GPR. Here differences in sites are more noticeable for younger and

older age groups. **c.** Change in MAE across EEG channel locations, with average MAE shown and error bars

indicated by standard deviation. **d.** Temporal transitions in the EEG across N1 (light pink) and N2 (light purple).

Similar to the Res-NN, the lowest MAE is observed during a transition between sleep N1 and N2 stages. Error bars

denote standard deviation for each epoch.

1137

1138

1139 ***Supplementary Table 3** – Performance of FBA across sleep stages in the validation dataset (D2).* The performances
1140 of the FBA following 10-fold cross-validation, using either a Res-NN or GPR model, in the D2 dataset. MAE,
1141 wMAE and RMSE are shown. N.B. N1 was excluded due to inconsistent availability across ages in D2.

Sleep stage	# channels, # recordings, # epochs	Res-NN		GPR	
		MAE (in years)	wMAE (in years)	MAE	wMAE
N3	2, 723, 13737	1.46	1.76	1.75	2.00
REM	2, 714, 13339	1.55	1.98	1.68	1.89
WAKE	2, 564, 8810	1.89	2.53	2.06	2.61
N2	2, 723, 13737	1.46	1.73	1.25	1.34

1142
1143

1144

1145 ***Supplementary Table 4*** *– Performance summary across test folds for Res-NN and GPR models*. Within each of the

1146 10 test folds, the mean absolute error (MAE) is summarized for both model approaches.

1147

1148

1149

1150

Folds	Res-NN MAE	GPR MAE
	Test D1	*Test D1*
1	0.63	0.72
2	0.52	0.73
3	0.61	0.73
4	0.52	0.78
5	0.61	0.75
6	0.55	0.76
7	0.50	0.98
8	0.58	0.80
9	0.62	0.86
10	0.53	0.80
Fold-wise Average	0.57	0.79
	Test D2	*Test D2*
1	1.69	1.20
2	1.43	1.25
3	1.63	1.25
4	1.40	1.15
5	1.43	1.34
6	1.63	1.11
7	1.39	1.21
8	1.35	1.34
9	1.32	1.20
10	1.51	1.33
Fold-wise Average	1.46	1.24
	Test D1 + D2	*Test D1 + D2*
1	1.15	1.07
2	0.97	1.05
3	1.03	0.91
4	0.95	1.14
5	0.96	1.21
6	1.06	1.16
7	0.89	1.11
8	1.36	1.04
9	1.00	1.13
10	1.11	1.03
Fold-wise Average	1.05	1.09