1

Tgfbr1 controls developmental plasticity between the hindlimb and external genitalia by remodeling their regulatory landscape

Anastasiia Lozovska[1], Artemis G. Korovesi[1], André Dias[1#], Alexandre Lopes[1], Donald A. Fowler[1], Gabriel G. Martins[1], Ana Nóvoa[1], and Moisés Mallo[1*]

1 Instituto Gulbenkian de Ciência, Rua da Quinta Grande 6, 2780-156 Oeiras, Portugal

Current address: Department of Experimental and Health Sciences, Universitat Pompeu Fabra, Barcelona, Spain.

* Lead author: mallo@igc.gulbenkian.pt

Key words: Tgfbr1, hindlimb, external genitalia, chromatin conformation, mouse

The hindlimb and external genitalia of present-day tetrapods are thought to derive from an ancestral common primordium that evolved to generate a wide diversity of structures adapted for efficient locomotion and mating in the ecological niche conquered by the species [1]. We show that despite long evolutionary distance from the ancestral condition, the early primordium of the mouse external genitalia preserved the capacity to take hindlimb fates. In the absence of *Tgfbr1*, the pericloacal mesoderm generates an extra pair of hindlimbs at the expense of the external genitalia. It has been shown that the hindlimb and the genital primordia share many of their key regulatory factors [2–5]. *Tgfbr1* controls the response to those factors acting in a pioneer-like mode to modulate the accessibility status of regulatory elements that control the gene regulatory networks leading to the formation of genital or hindlimb structures. Our work uncovers a remarkable tissue plasticity with potential implications in the evolution of the hindlimb/genital area of tetrapods, and identifies a novel mechanism for *Tgfbr1* activity that might also contribute to the control of other physiological or pathological processes.

The hindlimbs and the external genitalia both develop at the posterior end of the trunk
as an integral component of the trunk to tail transition [6]. Formation of the external
genitalia results from reciprocal interactions between the pericloacal mesoderm and
the endodermal cloaca that generate the genital tubercle (GT), the precursor of the
external genitalia, while also organizing the exit channels of the digestive and
excretory systems with the formation of the rectus/anus, the bladder and the urethra
[2,7–10]. The hindlimbs emerge from the somatic layer of the lateral plate mesoderm [11]
close to the position of the developing GT. Although the mature hindlimb and external
genitalia are anatomically very different, molecular studies indicate that they might be
closely connected developmentally, as they share many of their regulatory processes
[2–5]. Indeed, it has been reported that in some tetrapod species hindlimbs and genitals
share a common primordium, which might represent the ancestral condition [1]. In
mammals, it has been suggested that the GT became separated from the hindlimb by
the posterior displacement of the cloaca relative to the hindlimb bud, that provides
signals promoting genital fates to cells originating from the ventral and tail bud
mesenchyme [1].

Genetic analyses showed that *Tgfbr1* plays a key role during the induction stage of
these structures. *Tgfbr1* null mutant embryos fail to induce hindlimb or cloacal
structures, as shown by the absence of early markers for these tissues [12]. Conversely,
premature activation of Tgfbr1 signaling in the axial progenitors resulted in earlier
induction of these structures, as estimated by gene marker expression and the more
anterior position of the hindlimb buds, adjacent to the forelimbs [6]. Given the essential
role of Tgfbr1 signaling in the induction of both hindlimb and GT primordia despite
generating morphologically and functionally distinct structures, we wanted to
understand whether it also played a role at later stages of hindlimb and/or GT
development.

We used a conditional approach to inactivate *Tgfbr1* after the hindlimb and GT
primordia were laid down relying on the *Cdx2-Cre^{ERT}* transgenic driver [6], a *Tgfbr1* null
allele [12] and either *Tgfbr1^{flox}*, carrying LoxP sites flanking Exon 3 [13], or *Tgfbr1^{3ex3-flox}*,
containing a triplicated Exon 3 and associated LoxP sites that codes for a hypomorphic
receptor (Supplementary Fig. 1A-D). Experiments using either *Tgfbr1* floxed allele
gave similar phenotypes, but *Tgfbr1^{3ex3-flox}* turned out to facilitate the analysis of the
conditional *Tgfbr1* mutants, as *Tgfbr1^{3ex3-flox/-}:Cdx2Cre^{ERT+/0}* embryos required half of
the dose of tamoxifen than *Tgfbr1^{flox/-}:Cdx2Cre^{ERT+/0}* embryos to obtain the same

mutant phenotype, thus reducing the frequency of tamoxifen-derived embryo
miscarriage. Our tamoxifen treatment scheme produced *Tgfbr1* inactivation in the
tissues caudal from the forelimb bud, as confirmed by RT-qPCR analyses on RNA
extracted from the posterior part of E10,5 embryos, including the hindlimbs
(Supplementary Fig. 1E). Tamoxifen-treated embryos and fetuses with either the
Tgfbr1^{flox/-}:Cdx2Cre^{ERT+/0} or the *Tgfbr1^{3ex3-flox/-}:Cdx2Cre^{ERT+/0}* genotype will be
hereafter referred to as *Tgfbr1-cKO*.

The most prominent feature of *Tgfbr1-cKO* mutants was a duplication of their
hindlimbs [4/5 at embryonic age (E)16.5] (Fig. 1A, B). The fifth fetus analyzed at this
stage lacked overt hindlimb duplication but still contained small protrusions posterior
to the hindlimbs, a phenotype likely derived from less efficient recombination. The limb
identity of the duplicated structures was confirmed by the presence of skeletal
structures that, although variable in morphology, were clearly identified as belonging
to limbs (Fig. 1C; Supplementary Video 1). Another prominent phenotype of *Tgfbr1-*
cKO fetuses was the presence of an omphalocele (Fig. 1B). This phenotype, but not
the duplicated hindlimbs, was also observed in *Tgfb2*/*Tgfb3* double mutants [14]. This
indicates both that Tgfbr1 might mediate Tgfb2 and Tgfb3 activity in the body wall and
that different ligands should trigger Tgfbr1 signaling in the body wall and the
pericloacal/hindlimb mesoderm. *Tgfbr1-cKO* mutants lacked kidneys (Fig. 1D-E';
Supplementary Videos 2 and 3), a trait also observed in mutant fetuses for *Gdf11* [6,15],
another of the ligands shown to activate Tgfbr1 signaling during embryonic
development [16]. The cloacal-derived tissues were also strongly compromised in these
mutant embryos (Fig. 1D-E'; Supplementary Videos 2 and 3). The gut failed to
generate normal rectal-anal structures. Instead, the caudal end of the intestinal tube
merged with a structure that could be identified as a small bladder connected to a
hypomorphic urethra, thus resembling a persistent cloaca. The external genitalia were
also virtually absent from *Tgfbr1-cKO* fetuses. These phenotypes are likely to result
from interference with the functional interactions between the cloacal endoderm and
the pericloacal mesoderm driving normal morphogenesis of the cloacal region [2,3,5,9,17].
Indeed, endodermal malformations in this area were already evident in *Tgfbr1-cKO*
mutants at E10.5-E11.5, including an enlarged cloaca lacking features associated with
normal septation, and often containing protrusions entering the intra-cloacal lumen
(Supplementary Fig. 2D, D').

Morphological alterations in the hindlimb/pericloacal area of *Tgfbr1-cKO* embryos

were already visible at mid-gestation (Fig. 2). *Fgf8* expression in E10.0 embryos

revealed a posterior and medial expansion of the apical ectodermal ridge (AER) of the

hindlimb bud almost reaching the ventral midline of the embryo (Fig. 2A, A'),

suggesting an extension of the hindlimb field into the peri-cloacal mesoderm of *Tgfbr1-*

cKO embryos. This was associated with several expression changes in the cloacal

region. For instance, hindlimb specific markers, like *Pitx1* and *Lin28a* were ectopically

activated in the pericloacal region (Fig. 2B-C'). Conversely, *Tbx5* expression, normally

observed in the pericloacal mesoderm but not in the hindlimb, could not be detected

in this region of *Tgfbr1-cKO* embryos (Fig 2G, G'), and *Fgf8* expression was absent

from the cloacal endoderm of the mutant embryos. Together these observations

suggest a change of identity in the pericloacal mesoderm, becoming incorporated into

the hindlimb field instead of entering its normal GT developmental fate. Additional

observations were consistent with this hypothesis. At E11.0 when the presence of two

independent buds started to become apparent, thus anticipating the duplicated

hindlimb, the more caudal of these buds seemed to emerge from the pericloacal area.

The limb identity of this caudal bud was further supported by the expression of limb

dorso-ventral patterning markers, including *En1* and *Lmx1b* [18,19], which followed

typical limb patterns, although they were not as well defined as in control hindlimb

buds (Supplementary Fig. 2A-b'). Expression of genes active in both the hindlimb and

the pericloacal region/GT, was also consistent with the pericloacal tissues taking a

hindlimb fate. For instance, *Wnt5a*, which in wild type embryos is expressed in limb

buds beneath the AER and in the pericloacal region and GT, was not detected next to

the cloacal endoderm of *Tgfbr1-cKO* embryos, being instead observed beneath the

AER in the two buds of the extended hindlimb domain (Fig. 2E, E'). Also, *Hand2* and

Bmp4 were expressed in the mutants as a continuous domain with the more posterior

hindlimb bud invading the peri-cloacal tissue, instead of showing the separation

between the hindlimb and GT region observed in wild type embryos (Fig. 2F, F';

Supplementary Fig. 2C, C'). Additional support for the conversion of pericloacal

mesoderm into hindlimb identity was provided by genes showing dynamic expression

patterns in these structures. In wild type embryos, *Isl1* is initially activated in the

hindlimb and GT primordia, becoming later downregulated in the hindlimb while

keeping strong expression in the GT [20](Fig 2H). In *Tgbfr1-cKO* mutants *Isl1* showed

residual expression next to the expanded endodermal cloaca (note that *Isl1* is also

expressed in the cloacal endoderm [21]) but became downregulated in the adjacent

mesoderm in the area likely to generate the extra hindlimb bud (Fig 2H'). Conversely,

Fgf10, an essential gene for limb induction [22], whose expression in the GT is observed

only at later stages [23], was strongly activated already at E10,5 following limb-like

patterns, reaching mesodermal tissue adjacent to the endoderm (Fig. 2D, D').

It has been shown that regulation of GT development relies to a large extent on signals

from the cloacal endoderm, with Shh playing a central role in this process [17,24]. The

observation that *Wnt5a*, a known Shh target in the GT [17], was not activated in the

pericloacal mesoderm of *Tgfbr1-cKO* despite strong endodermal *Shh* expression (Fig.

3A, A', a, a'; Supplementary Fig. 2D, D', d, d'), suggested that either this mesenchyme

became refractory to Shh activity or that it changed its response profile. The

expression of the Shh target *Gli1* was consistent with the second possibility, as it

showed equivalent patterns in the pericloacal mesenchyme in both wild type and

Tgfbr1-cKO embryos (Fig. 3B, B'), thus indicating that this tissue still responds to the

Shh signal from the endoderm.

We explored the mechanisms modulating the pericloacal mesoderm response to Shh

(and most likely also to other regulatory factors) by comparing the genomic

accessibility profiles in the extra hindlimbs and pericloacal region of *Tgfbr1-cKO* with

those of hindlimb buds and GT of control embryos by ATAC-seq [25]. The global

chromatin accessibility patterns in the extra hindlimb of the mutant embryos were

closer to those obtained from the control GT than to those from control hindlimbs (Fig.

3C; Supplementary Fig. 3A), consistent with the idea of its pericloacal origin. Despite

this, we identified two changes in accessibility patterns fitting with their potential

involvement in the control of cell fate decisions of the pericloacal mesoderm. Pattern

1 included 4922 regions that in control tissues were accessible in the GT but not in the

hindlimb, and that lost accessibility in the tissues from *Tgfbr1-cKO* embryos (Fig. 3D,

E). These elements could represent enhancers involved in the activation of genes

expressed in the pericloacal region of wild type embryos but inactive in the *Tgfbr1-*

cKO mutants. Pattern 2 contains 2013 chromatin regions representing elements that

in control embryos were accessible in the hindlimb but not in the GT and became

accessible in the extra hindlimb of the mutant embryos (Fig 4A, A'). These elements

could be involved in activating the limb program in the pericloacal mesoderm of *Tgfbr1-*

cKO embryos.

We identified pattern 1 elements within the *Tbx5, Isl1* and *Wnt5a* genomic regions,

three of the pericloacal mesoderm markers down-regulated in the pericloacal region

of *Tgfbr1-cKO* mutants (Fig. 3F-G'; Supplementary Fig. 3B, B'). These elements

mapped to highly conserved areas, suggesting their possible involvement in regulatory

processes. When tested using a transgenic reporter assay, one of the elements

associated with *Tbx5* showed activity in the pericloacal region (n=6/14) (Fig 3G, G').

Interestingly, it did not activate overt forelimb expression (Supplementary Fig. 3D), the

major expression domain of this gene [26], further indicating specificity for the pericloacal

region. Similarly, the element identified downstream of *Isl1* also reproduced to a large

extent *Isl1* expression in the GT when tested in transgenic reporter assays (n=5/12).

An element associated with *Wnt5a* was also able to activate expression in the GT,

although much less frequently as the other elements (n=2/18) (Supplementary Fig.

3B, B'). This element was also active in the tail, but it did not promote expression in

the hindlimb. Together, these observations are consistent with these elements indeed

representing enhancers potentially involved in the expression of the relevant genes in

the pericloacal area.

We found pattern 2 elements in regions associated with genes playing essential roles

during the earliest stages of limb development, which could thus play a relevant role

in promoting limb fates from the pericloacal mesoderm (Fig. 4A, A'). As for pattern 1

elements, these regions were also highly conserved among vertebrates. We found

pattern 2 elements in the *Fgf10* genomic region (Supplementary Fig. 3C), a gene

activated in the mesenchyme adjacent to the cloaca entering hindlimb fate in the

mutant embryo (Fig. 2D, D'). Interestingly, published ChIP-seq data from forelimb buds

[27,28] showed binding of Gli3 and Hoxa13 to one of these elements, located within an

intronic region of this gene (Supplementary Fig. 3C), indicating that it might respond

to Shh and/or Hoxa13 activities in the developing limb buds. The lack of accessibility

of this element in the GT of wild type embryos might thus suggest that Tgfbr1 renders

this enhancer blind to endodermal Shh and/or the strong pericloacal Hoxa13 (and

maybe also Hoxd13) expression [29], consistent with absent *Fgf10* at early stages in this

tissue. Rather surprisingly, this element failed to generate reporter activity in

transgenic embryos (n=10). We still do not understand whether this reflects real lack

of activity or the absence of proper genomic context for their activity.

Another potentially relevant pattern 2 element was located within the *Fmn1* locus,

matching the position of one of the enhancers (element 7 in Malkmus et al., 2021)

controlling *Grem1* in the limb buds (Supplementary Fig. 3E) that had been shown to

respond to Shh activity from the ZPA [30,31]. In wild type embryos *Grem1* expression is

not detected in the pericloacal mesoderm or in the GT (Fig. 4D), indicating that in these

tissues *Tgfbr1* might render the *Grem1* regulatory elements blind to endodermal Shh.

The locked configuration of element 7 in the GT (Fig. 4C) is consistent with this

hypothesis. Conversely, the gain of accessibility of element 7 in the extra hindlimb

suggests that in the absence of *Tgfbr1* the pericloacal mesoderm could become

responsive to the endodermal Shh. Considering the capacity of *Grem1* to expand the

expression of AER markers [32–34], this gene is a candidate to contribute to the observed

AER extension in the hindlimb primordium of *Tgfbr1-cKO* embryos. *Grem1* expression

analyses in *Tgfbr1-cKO* embryos revealed unexpected complexities. Some features

of the expression pattern were consistent with *Grem1* becoming responsive to

endodermal Shh, as expression was detected in the caudal-most end of the expanded

hindlimb next to the endoderm (Fig. 4D'). Rather surprisingly, however, *Grem1*

expression was strongly reduced or absent throughout the mesoderm of the extended

hindlimb bud of *Tgfbr1-cKO* embryos (Fig. 4D'). *Grem1* limb expression depends on

Shh activity from the zone of polarizing activity (ZPA) [30,35,36], suggesting that the

absent *Grem1* expression resulted from the abnormal *Shh* expression observed in the

hindlimb buds of *Tgfbr1-cKO* mutants, which followed spotty patterns beneath the AER

instead of forming a well-defined ZPA in the posterior limb bud mesenchyme (Fig. 3A,

A').

We explored whether this abnormal *Shh* expression resulted from a functional impact

of the absence of Tgfbr1 activity on the ZRS, the regulatory element controlling *Shh*

expression in the ZPA [37]. As expected, this region was accessible in wild type

hindlimbs (Fig. 4E). Conversely, it was in a locked configuration in the wild type GT,

consistent with the absence of the *Shh* expression in the pericloacal mesoderm or the

GT, despite the presence of relevant activators of this enhancer, like Hand2, Hoxa13

and Hoxd13 [4,29,38,39] (the endodermal *Shh* expression is under the control of a different

enhancer[40]). Remarkably, the ZRS was also non-accessible in the hindlimb region of

Tgfbr1-cKO embryos, consistent with the observed abnormal *Shh* limb expression.

Therefore, Tgfbr1 seems to also control ZRS accessibility to their regulatory

transcription factors. Interestingly, this enhancer is part of an additional pattern

(pattern 3) consisting of elements requiring Tgfbr1 activity to become accessible in the

limb bud. Analysis of the ATAC-seq data revealed that 13471 elements fit the criteria

of this pattern (Fig. 4B, B'). The complementary tissue specificities of patterns 1 and

3 indicates context dependency for Tgfbr1 activity in the modulation of genomic

configuration.

Discussion

The capacity of the pericloacal mesoderm of mouse embryos to generate both external

genitalia and hindlimb structures reveals a remarkable degree of developmental

plasticity in this tissue. This finding provides a new dimension to the developmental

connection between the hindlimb and the GT, previously suggested by the

considerable overlap in their regulatory networks [2–5]. It has been shown that the paired

external genitalia of snakes develop at least to some extent from mesodermal tissues

in the prospective hindlimb domain. It had been suggested that the proximity of this

mesoderm to the cloacal signals could play a role in this process [1]. However, such

proximity was also observed in lizards that develop both hemipenises and hindlimbs

[1]. It will be therefore interesting to determine whether a mechanism related to the

control of developmental plasticity uncovered in our work plays a role in the absence

of hindlimbs in snakes. In addition, understanding the relative weight of the different

factors whose expression was lost or ectopically activated in the pericloacal region of

the *Tgfbr1-cKO* mutants might help to uncover the core regulatory network able to

promote specific and stable fates from an intrinsically plastic tissue. This will require

generating different combinations of gain and loss of function experiments.

An important finding from our work is that *Tgfbr1* controls pericloacal mesoderm

plasticity by modulating its regulatory landscape. It has been reported that BMP

signaling can affect the chromatin status of Smad-responsive enhancers by interacting

with chromatin remodelers like Foxh1 or Smarca4, eventually rendering these

regulatory elements competent to convey the Smad transcriptional activity [41,42]. What

we have observed in the pericloacal/hindlimb region does not easily fit within this

mechanism. For instance, pattern 2 elements are rendered non-accessible by *Tgfbr1*,

compatible with a repressive rather than potentially activating function. This could

result from interference with a positive activity provided by other members of the

Tgfb/BMP family on those enhancers. However, at least for the *Grem1* enhancer 7,

this is not a likely explanation, as ChIP-seq analyses failed to detect Smad4 binding

on limb tissues, where *Grem1* is normally expressed [30]. In addition, we failed to detect

enrichment of Smad binding to enhancers rendered accessible by *Tgfbr1* using a

HINT-ATAC transcription factor footprinting analysis [43] on the ATAC-seq profiles
(Supplementary Table 1). Even considering the limitations of the bioinformatic
approach, these observations suggest that effectors of the Tgbfr1/BMP signaling
pathways are not among the main regulators of the activity of those elements.
Consistent with this, Smads have so far not been identified among the factors involved
in the activation of the ZRS enhancer.

Our data suggest a general model according to which Tgfbr1 signaling acts in a
pioneer-like activity, determining the accessibility of regulatory elements to patterning
transcription factors. As many of the patterning factors regulating hindlimb and GT
development are common to both structures, the global output of Tgfbr1 activity would
be establishing how their primordia respond to those factors. While extensive analysis
of limb development over the last decades might already guide the identification of
factors interacting with different enhancers involved in limb-related processes,
understanding the control of pericloacal gene expression will require focused ChIP-
seq profiles from this tissue and integrating them with the ATAC-seq data. This
information will also provide a direct estimation of the involvement of effectors of the
Tgfβ/Bmp pathways in these processes.

It should be noted, however, that although the global output resembles some
hallmarks of a pioneer activity, the mechanism is unlikely to follow routes similar to
those described for classical pioneer factors [44], as the Tgfbr1-mediated chromatin
remodeling process does not seem to involve Smad binding to the regulated elements,
and controls accessibility in opposite ways for different enhancers. Identifying the
mechanism of this activity, and whether similar or different processes operate for the
different patterns in the regulatory landscape, will be essential to fully understand how
Tgfbr1 signaling regulates morphogenetic processes. In this regard, it will be important
to determine whether a similar pioneer-like mechanism also operates in other
physiological and pathological processes under the control of members of the
Tgfβ/BMP signaling family.

1. Tschopp, P. *et al.* A relative shift in cloacal location repositions external genitalia in
amniote evolution. *Nature* **516**, 391–394 (2014).
2. Yamada, G. *et al.* Molecular genetic cascades for external genitalia formation: An
emerging organogenesis program. *Developmental Dynamics* **235**, 1738–1752 (2006).

3.   Cohn, M. J. Development of the external genitalia: Conserved and divergent
mechanisms of appendage patterning. *Developmental Dynamics* **240**, 1108–1115
(2011).

4.   Kondo, T., Zakany, J., Innis, J. W. & Duboule, D. Of fingers, toes and penises [1]. *Nature*
390, 29 (1997).

5.   Lin, C. *et al.* Delineating a Conserved Genetic Cassette Promoting Outgrowth of Body
Appendages. *PLoS Genet* **9**, e1003231 (2013).

6.   Jurberg, A. D., Aires, R., Varela-Lasheras, I., Nóvoa, A. & Mallo, M. Switching axial
progenitors from producing trunk to tail tissues in vertebrate embryos. *Dev Cell* **25**,
451–462 (2013).

7.   Seifert, A. W., Harfe, B. D. & Cohn, M. J. Cell lineage analysis demonstrates an
endodermal origin of the distal urethra and perineum. *Dev Biol* **318**, 143–152 (2008).

8.   Miyagawa, S. *et al.* Disruption of the temporally regulated cloaca endodermal β-
catenin signaling causes anorectal malformations. *Cell Death Differ* **21**, 990–997
(2014).

9.   Miyagawa, S. *et al.* Dosage-dependent hedgehog signals integrated with Wnt/β-
catenin signaling regulate external genitalia formation as an appendicular program.
Development **136**, 3969–3978 (2009).

10. Huang, Y. C., Chen, F. & Li, X. Clarification of mammalian cloacal morphogenesis using
high-resolution episcopic microscopy. *Dev Biol* **409**, 106–113 (2016).

11. Tickle, C. How the embryo makes a limb: Determination, polarity and identity. *J Anat*
227, 418–430 (2015).

12. Dias, A. *et al.* A Tgfbr1/Snai1-dependent developmental module at the core of
vertebrate axial elongation. *Elife* **9**, e56615 (2020).

13. Pineault, K. M., Novoa, A., Lozovska, A., Wellik, D. M. & Mallo, M. Two CRISPR/Cas9-
mediated methods for targeting complex insertions, deletions, or replacements in
mouse. *MethodsX* **6**, 2088–2100 (2019).

14. Dünker, N. & Krieglstein, K. Tgfβ2-/-Tgfβ3-/- double knockout mice display severe
midline fusion defects and early embryonic lethality. *Anat Embryol (Berl)* **206**, 73–83
(2002).

15. McPherron, A. C., Lawle, A. M. & Lee, S.-J. Regulation of anterior/posterior patterning
of the axial skeleton by growth/differentiation factor 11. *Nat Genet* **22**, 260–264
(1999).

16. Andersson, O., Reissmann, E. & Ibáñez, C. F. Growth differentiation factor 11 signals
through the transforming growth factor-beta receptor ALK5 to regionalize the
anterior-posterior axis. *EMBO Rep* **7**, 831–837 (2006).

17. Perriton, C. L., Powles, N., Chiang, C., Maconochie, M. K. & Cohn, M. J. Sonic
hedgehog signaling from the urethral epithelium controls external genital
development. *Dev Biol* **247**, 26–46 (2002).

18. Davis, C. A., Holmyard, D. P., Millen, K. J. & Joyner, A. L. Examining pattern formation in
mouse, chicken and frog embryos with an En-specific antiserum. *Development* **111**,
287–298 (1991).

19. Riddle, R. D. *et al.* Induction of the LIM Homeobox Gene Lmx7 by WNT7a Establishes
Dorsoventral Pattern in the Vertebrate Limb. *Cell* **83**, 631–640 (1995).

20. Yang, L. *et al.* Isl1 Cre reveals a common Bmp pathway in heart and limb development.
Development **133**, 1575–1585 (2006).

21.   Su, T. *et al.* LIM homeodomain transcription factor Isl1 affects urethral epithelium
353   differentiation and apoptosis via Shh. *Cell Death Dis* **10**, 713 (2019).

354   22.   Min, H. *et al.* Fgf-10 is required for both limb and lung development and exhibits
355   striking functional similarity to Drosophila branchless. *Genes Dev* **12**, 3156–3161
356   (1998).

357   23.   Satoh, Y. *et al.* Regulation of external genitalia development by concerted actions of
358   FGF ligands and FGF receptors. *Anat Embryol (Berl)* **208**, 479–486 (2004).

359   24.   Haraguchi, R. *et al.* Unique functions of sonic hedgehog signaling during external
360   genitalia development. *Development* **128**, 4241–4250 (2001).

361   25.   Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and enables
362   interrogation of frozen tissues. *Nat Methods* **14**, 959–962 (2017).

363   26.   Rallis, C. *et al.* Tbx5 is required for forelimb bud formation and continued outgrowth.
364   *Development* **130**, 2741–2751 (2003).

365   27.   Sheth, R. *et al.* Distal Limb Patterning Requires Modulation of cis-Regulatory Activities
366   by HOX13. *Cell Rep* **17**, 2913–2926 (2016).

367   28.   Lex, R. K. *et al.* GLI transcriptional repression regulates tissue-specific enhancer
368   activity in response to hedgehog signaling. *Elife* **9**, e50670 (2020).

369   29.   Scott, V., Morgan, E. A. & Stadler, H. S. Genitourinary functions of Hoxa13 and Hoxd13.
370   *Journal of Biochemistry* vol. 137 671–676 Preprint at
371   https://doi.org/10.1093/jb/mvi086 (2005).

372   30.   Malkmus, J. *et al.* Spatial regulation by multiple Gremlin1 enhancers provides digit
373   development with cis-regulatory robustness and evolutionary plasticity. *Nat Commun*
374   **12**, 5557 (2021).

375   31.   Zhu, J., Patel, R., Trofka, A., Harfe, B. D. & Mackem, S. Sonic hedgehog is not a limb
376   morphogen but acts as a trigger to specify all digits in mice. *Dev Cell* **57**, 2048–2062
377   (2022).

378   32.   Michos, O. *et al.* Gremlin-mediated BMP antagonism induces the epithelial-
379   mesenchymal feedback signaling controlling metanephric kidney and limb
380   organogenesis. *Development* **131**, 3401–3410 (2004).

381   33.   Nissim, S., Hasso, S. M., Fallon, J. F. & Tabin, C. J. Regulation of Gremlin expression in
382   the posterior limb bud. *Dev Biol* **299**, 12–21 (2006).

383   34.   Farin, H. F. *et al.* Tbx2 Terminates Shh/Fgf Signaling in the Developing Mouse Limb Bud
384   by Direct Repression of Gremlin1. *PLoS Genet* **9**, e1003467 (2013).

385   35.   Zuniga, A. *et al.* Mouse limb deformity mutations disrupt a global control region
386   within the large regulatory landscape required for Gremlin expression. *Genes Dev* **18**,
387   1553–1564 (2004).

388   36.   Li, Q. *et al.* A Gli silencer is required for robust repression of gremlin in the vertebrate
389   limb bud. *Development (Cambridge)* **141**, 1906–1914 (2014).

390   37.   Lettice, L. A. *et al.* A long-range Shh enhancer regulates expression in the developing
391   limb and fin and is associated with preaxial polydactyly. *Hum Mol Genet* **12**, 1725–
392   1735 (2003).

393   38.   Galli, A. *et al.* Distinct roles of Hand2 in initiating polarity and posterior Shh expression
394   during the onset of mouse limb bud development. *PLoS Genet* **6**, e1000901 (2010).

395   39.   Leal, F. & Cohn, M. J. Loss and Re-emergence of Legs in Snakes by Modular Evolution
396   of Sonic hedgehog and HOXD Report Loss and Re-emergence of Legs in Snakes by
397   Modular Evolution of Sonic hedgehog and HOXD Enhancers. *Current Biology* **26**, 1–8
398   (2016).

40. Sagai, T. *et al.* A cluster of three long-range enhancers directs regional Shh expression
in the epithelial linings. *Development* **136**, 1665–1674 (2009).
41. Coda, D. M. *et al.* A network of transcription factors governs the dynamics of
NODAL/Activin transcriptional responses. *J Cell Sci* **135**, jjcs259972 (2022).
42. Coda, D. M. *et al.* Distinct modes of SMAD2 chromatin binding and remodeling shape
the transcriptional response to NODAL/Activin signaling. *Elife* **6**, 1–31 (2017).
43. Li, Z. *et al.* Identification of transcription factor binding sites using ATAC-seq. *Genome*
Biol **20**, 45 (2019).
44. Zaret, K. S. & Carroll, J. S. Pioneer transcription factors: Establishing competence for
gene expression. *Genes Dev* **25**, 2227–2241 (2011).
409

Acknowledgements

We would like to thank the members of the Mallo lab, especially Ana Casaca for continuous support at different stages of this project, the IGC mouse facility for their help with animal housing and genomics facility for helping with the ATAC-seq procedures. This project was funded by Fundação para a Ciência e a Tecnologia (FCT) grants 2022.01629.PTDC to MM, PTDC/BII-BTI/32375/2017 to GGM, and PhD fellowships PD/BD/128437/2017 to AL and PD/BD/128426/2017 to AD, by the PhD fellowship 36/8l-D/21 from Fundação Calouste Gulbenkian to AK, and the research infrastructures PPBI-POCI-01-0145-FEDER-022122 to the Advanced Imaging Facility, and Congento LISBOA-01-0145-FEDER-022170 to the animal facility, both co-financed by Lisboa Lisboa 2020/FEDER and FCT (Portugal).

Author contribution

Conceptualization: A.L. and M.M.; Formal Analysis: A.L. and A.K.; Funding Acquisition: M.M.; Investigation: A.L., A.K. and M.M.; Methodology: A.L., A.D. G.G.M. A.N., A.Lo. and D.F.; Writing – Original Draft Preparation: A.L. and M.M.; Writing – Review & Editing: A.L., A.K. and M.M.

Competing interest declaration

The authors declare no competing interests.

Figure 1. Malformations in the E16,5 *Tgfbr1-cKO* fetuses. (A, B) Images of the fixed WT (A) and *Tgfbr1-cKO* (B) fetuses. The mutant fetus shows the presence of an omphalocele (asterisk) and hindlimb duplication. (C) 3D reconstruction of the limb skeleton of a *Tgfbr1-cKO* fetus obtained by OPT and after segmentation of the limb skeleton. Extra hindlimbs are in magenta. Ossification shown in yellow. (D–E') 3D reconstruction of organs and the excretory outlets of the control (D, D') and *Tgfbr1-cKO* (E, E') fetuses. Images were obtained by OPT, followed by segmentation of the relevant structures. D' E' show virtual sections of the segmented 3D specimen. Asterisk in E' shows the gut-bladder connection in the mutant. K: kidney; G: gut; Bl: bladder; Ov: ovary Ut: uterus; V: vagina; Ur: urethra.

443
Supplementary Figure 1. Generation on the *Tgfbr1-cKO*.(A) Scheme showing
Tgfbr1^{flox} and *Tgfbr1$^{3ex3-flox}$* alleles. Highlighted is the region containing exons 2, 3
and 4 (colored boxes) and the LoxP sites (black triangles). Also represented is the
generation of a mRNA with a triplicated exon 3 after splicing of the transcript from the
Tgfbr1$^{3ex3-flox}$ allele. (B, B') Skeletal staining of a E17.5 wild type (B) and *Tgfbr1^{3ex3-}*
$^{flox/-}$ fetus (B'), showing the presence of additional thoracic (T) and lumbar (L)
vertebrae, resembling a *Gdf11* mutant phenotype. (C) RT-PCR of the *Tgfbr1* coding
region from wild type (1,5 kb) and *Tgfbr1$^{3ex3-flox/3exe-flox}$* (2 kb) embryos. (D). Western
blot analysis of protein extracts from 293T cells transfected with mammalian
expression vectors carrying either the Tgfbr1^{3ex3} or the Tgfbr1 coding regions,
showing that the *Tgfbr1^{3ex3}* transcript produces a protein with a higher molecular
mass than its wild type counterpart. (E) RT-qPCR showing recombination efficiency
in the *Tgfbr1$^{flox/-}$:Cdx2Cre$^{ERT+/0}$* embryos. Normalized *Tgfbr1* expression level in
Tgfbr1$^{flox/-}$:Cdx2Cre$^{ERT+/0}$ is not different from *Tgfbr1$^{-/-}$*. Level of normalized
expression in *Tgfbr1$^{+/flox}$:Cdx2Cre$^{ERT+/0}$* equals that of *Tgfbr1$^{flox/-}$* and half of
Tgfbr1$^{flox/+}$ control, consistent with it carrying only one functioning allele. NS:
nonsignificant, **p<0.01, ***p<0.001
461
462

463
464
Figure 2. The pericloacal mesenchyme of the *Tgfbr1-cKO* embryos adopts hindlimb
fate. Ventral views of the hindlimb/cloacal region of wild type (A-H) or *Tgfbr1-cKO*
embryos (A'-H') stained for *Fgf8* (A, A'), *Lin28a* (B, B'), *Pitx1* (, C'), *Fgf10* (D, D'),
Wnt5a (E, E'), *Hand2* (F, F'), *Tbx5* (G, G') or *Isl1* (H, H'). *Fgf8* expression show
posterior and medial elongation of the AER (black arrow) and the absence of
expression in the cloacal endoderm (black arrowhead). Hindlimb markers *Lin28a*,
Pitx1 and *Fgf10* extend into the pericloacal region in E10,5 *Tgfbr1-cKO* embryos
(white arrows). *Wnt5a* and *Hand2* switch their expression profiles in the pericloacal
mesoderm (yellow arrowheads), now following a limb-like pattern. In addition,
pericloacal expression of *Wnt5a*, *Tbx5* and *Isl1* is lost in the mutant embryos (yellow
arrows).
476

Supplementary Figure 2. Molecular characterization of the *Tgfbr1-cKO* embryos. (A-b') Dorsal-ventral polarity of the *Tgfbr1-cKO* mutant's hindlimb. Expression of the ventral limb marker *En1* in E10.5 control (A, a) and *Tgfbr1-cKO* (A', a') embryos. (A, A') show lateral views of the hindlimb region in whole mounted embryos; (a, a') show transversal sections through the region indicated in A, A'. Black arrowheads show *En1* expression in the ventral ectoderm of the limb bud. Expression of the dorsal mesodermal limb marker *Lmx1b* in E10.5 control (B, b) and *Tgfbr1-cKO* (B', b') embryos. (B, B') show lateral views of the hindlimb region in whole mounted embryos; (b, b') show transversal sections through the region indicated in B, B'. Yellow arrowheads show *Lmx1b* expression in the dorsal mesenchyme of the limb bud. (C, C') Ventral views of E10.5 wild type (C) and *Tgfbr1-cKO* (C') embryos stained for *Bmp4* showing the extension of the signal into the pericloacal region in the mutant embryos (white arrow). (D-d') Expression of *Shh* in E11,5 control (D, d) and *Tgfbr1-cKO* (D', d') embryos. (D, D') show lateral views of whole mounted embryos; (d, d') show transversal section through the region indicated in D, D' to show the enlarged cloaca (black arrows) with presence of the intralumenal protrusions in the mutant embryo (asterisk), still keeping endodermal *Shh* expression.

497
Figure 3. Tgfbr1 regulates the chromatin landscape in the pericloacal region. (A–B')
The pericloacal mesenchyme of *Tgfbr1-cKO* embryos responds to Shh signaling
from the cloacal endoderm. *Shh* expression in E11,5 control (A, a) and *Tgfbr1-cKO*
(A', a') embryos. (A, A') show ventral views of the stained embryos: (a, a') show
sagittal sections through cloacal (CI) region. *Gli1* expression in E10,5 control (B) and
Tgfbr1-cKO (B') embryos. (C) PCA analysis reveals closer proximity of the chromatin
accessibility pattern of the *Tgfbr1-cKO* extra hindlimb to that of *Tgfbr1-cKO* and
control GT than to the control hindlimb. (D, E) Chromatin regions following pattern 1.
(D) Venn diagram showing interception of the regions tested more accessible in
control GT than in other tissues by pairwise comparisons [FDR<0.001, logFC>1.5].
(E) Heatmap showing log2 normalized counts across samples of the regions
following pattern 1. (F, G') ATAC-seq tracks showing accessible chromatin in control
GT, but not in other samples in regions 24 kb downstream of the *Isl1* TSS (F) and 93
kb upstream of the *Tbx5* TSS (G). The regions highlighted with the blue shade are
characterized by high conservation level among placental animals (bottom track). (F',
G') The highlighted elements drive b-galactosidase expression in the GT region
(white arrows). Hoxa13 and Gli3 ChiP-Seq tracks from forelimb buds (GSE81356
and GSE133710, respectively) are shown in yellow and magenta, respectively. cHL:

control hindlimb; mExHL: mutant extra hindlimb; cGT: control GT; mGT: mutant GT;
CR: conserved region.
518

519
Supplementary Figure 3. (A) Distance heatmap of the chromatin accessibility
patterns showing that the *Tgfbr1-cKO* extra hindlimb clusters closer to mutant and
control GT than to control limb. (B) ATAC-seq tracks showing a chromatin region
(highlighted with a blue shadow) accessible in control GT, but not in other tissues of
a region 93 kb upstream of the *Wnt5a* TSS. This highly conserved among placental
animals (bottom track). (B') The highlighted element in (B) drives b-galactosidase
expression in the GT region (white arrow) in a transgenic reporter assay. (C) A region
located within intron 1 of the *Fgf10* gene (highlighted with a blue shadow), is
accessible in the hindlimb samples of the control and *Tgfbr1-cKO* embryos, but not
in GT samples. ChIP data from forelimb buds show that this genomic region is bound
by Hoxa13 and Gli3. (D) The putative *Tbx5* GT enhancer shown in Figure 3 does not
drive b-galactosidase expression in the forelimb (black arrowhead, forelimb outlined
with the dashed line). (E) b-galactosidase expression driven by the *Grem1* enhancer
in Figure 4 using a transgenic reporter assay (n=2/8) (limbs outlined with the dashed
line). CR: conserved region.
535
536

Figure 4. (A, A') Chromatin regions following pattern 2. (A) Venn diagram showing
interception between the regions more accessible in mutant extra limb than in mutant
GT [FDR<0.001, logFC>1.5] and more accessible in control hindlimb than in control
GT [FDR<0.001, logFC>1.5], but at the same time not differentially accessible
between mutant extra hindlimb and control hindlimb [FDR>0.01, logFC<0.5]. (A')
Heatmap showing log2 normalized counts from genomic regions following pattern 2
across the samples. (B, B') Chromatin regions following pattern 3. (B) Venn diagram
showing interception between regions tested more accessible in control hindlimb
than in other tissues by pairwise comparisons [FDR<0.001, logFC>1.5]. (E) Heatmap
showing log2 normalized counts for the chromatin regions following pattern 3 across
samples. (C) ATAC-Seq profiles showing a *Grem1* enhancer that gains accessibility
in the mutant extra hindlimb (highlighted with the blue shadow). Also shown are
ChiP-Seq tracks of ChIP seq data from wild type forelimbs for Gli3 (yellow) and
Hoxa13 (magenta) ((GSE81356 and GSE133710, respectively), as well as the
conservation in placental mammals. (D, D') *Grem1* expression in E10.5 wild type (D)
or *Tgfbr1-cKO* (D') embryos showing the absence of pericloacal expression in wild
type embryos and ectopic activation in the caudal-most region of the pericloacal
mesoderm of the mutant embryo (white arrow). *Grem1* is expressed in the
developing limb bud of the control embryo, but it is almost non detectable in the

developing limb buds of the mutant embryo (white arrowheads). (E) ATAC-seq data through the ZRS (highlighted with the blue shadow), showing that it is accessible in the control limb but not in any of the other samples. cHL: control hindlimb; mExHL: mutant extra hindlimb; cGT: control GT; mGT: mutant GT.

Supplementary movie 1. OPT image from a E16.5 *Tgfbr1-cKO* fetus highlighting the limbs. The extra hindlimbs are shown in magenta.

Supplementary movie 2. OPT image from a E16.5 wild type fetus highlighting the caudal end of the intestine (red), the kidneys (cold green), the ovaries and uterus (bright green), the vagina (yellow), the urinary bladder (dark blue) and the urethra (light blue).

Supplementary movie 3. OPT image from a E16.5 *Tgfbr1-cKO* fetus highlighting the caudal end of the intestine (red), the urinary bladder (dark blue) and the urethra (light blue).

Supplementary Table 1. Transcription factor footprint analysis of pattern 1 elements. Column A: number of pattern 1 regions with at least 1 occurrence of the specified transcription factor; column B: number of pattern1 regions with no occurrence of the specified transcription factor; C: number of background regions with at least 1 occurrence of the specified transcription factor; D: number of background regions with no occurrence of the specified transcription factor.

Materials and methods

Ethics statement

All animal procedures were performed in accordance with Portuguese (Portaria 1005/92) and European (directive 2010/63/EU) legislations and guidance on animal use in bioscience research. The project was reviewed and approved by the Ethics Committee of "Instituto Gulbenkian de Ciência" and by the Portuguese National Entity "Direcção Geral de Alimentação Veterinária" (license reference: 014308).

Mice

Generation of the *Tgfbr1^Flox^* mouse lines was previously reported [13]. *Tgfbr1^3ex3-flox^*

resulted from the same experiments in which *Tgfbr1^flox^* was generated, identified upon

sequencing through the Tgfbr1 locus which revealed the presence of three exon 3

together with LoxP sites in the introns between the triplicated exons and in those

connecting with exon 2 and exon 4. Characterization of mice homozygous for the

*Tgfbr1^3ex3-flox^* allele revealed the presence of a transcript containing the three exon 3

in frame with the flanking exons 2 and 4. Expression of this mRNA in HEK293T cells

confirmed that it can generate a protein product with the expected size of Tgfbr1

containing a triplication of exon 3 (supplementary Fig. 1). The *Cdx2-cre^ERT^* mice were

previously reported [6]. Mice were genotyped from ear of digit biopsies. Samples were

incubated in 50 μL of the PBND buffer (50 mM KCl, 10 mM Tris-HCl, pH 8.3, 2.5 mM

MgCl$_2$, 0.1 mg/mL gelatin, 0.45% NP40, 0.45% Tween 20) supplemented with 100

µg/mL of proteinase K at 55˚C overnight. Proteinase K was inactivated at 95˚C for 15

minutes and 1 μL of genomic DNA was used for genotyping by PCR. *Cdx2-*

*cre^ERT^::Tgfbr1^+/-^* mice were genotyped for the *Cre* allele and *Tgfbr1* null allele;

*Tgfbr1^flox/+^* and *Tgfbr1^3ex3-flox/+^* mice were genotyped with primers surrounding 3'-LoxP

(primers listed in supplementary table 2).

Tgfbr1-cKO (*Cdx2-cre^ERT^::Tgfbr1^flox/-^* or *Cdx2-cre^ERT^::Tgfbr1^3ex3-flox/-^*) embryos were

obtained by crossing *Tgfbr1^flox/flox^* or *Tgfbr1^3ex3-flox/+^* females with *Tgfbr1^+/-^:Cdx2-cre^ERT^*

males. Noon of the day of the plug was considered E0.5. To induce recombination

pregnant females were treated with tamoxifen (Sigma, T5648) dissolved in corn oil.

Tamoxifen was administered at 0.1 mg per gram of body weight by oral gavage either

once at E6.75 when *Tgfbr1^3ex3-flox^* females were involved, or twice, at E6.75 and E7.25

when *Tgfbr1^flox^* females were involved.

Embryos were obtained by cesarean section and processed for further analyses (see

below). They were genotyped from their yolk sacs for Cre, *Tgfbr1^-^* and *Tgfbr1^flox^* when

analyzing *Tgfbr1-cKO* embryos or for β-gal when analyzing the reporter transgenic

embryos (primers listed in supplementary table 2). Yolk sacs were collected in 50 μL

of yolk sac lysis buffer (50 mM KCl, 10 mM Tris-HCl, pH8.3, 2 mM MgCl$_2$, 0.45%

Tween-20, 0.45% NP40) supplemented with 100 µg/mL of proteinase K and incubated

at 55˚C overnight. Samples were heat-deactivated as previously described and used

for PCR.

Transgenic embryos

To generate transgenic constructs, the relevant enhancers (genomic coordinates shown in supplementary table 3) were amplified by PCR (primers in supplementary table 2) from genomic DNA and cloned into a vector containing the adenovirus major late promoter, the coding region of the β-galactosidase gene and the SV40 polyadenylation signal [6]. The enhancers were confirmed by direct sequencing. The constructs were isolated from the plasmid backbone, gel-purified using the NZYGelpure (NZYTech #MB01102) and eluted from the columns with 50 μL of the kit's elution buffer. The purified constructs were diluted in microinjection buffer (10 mM Tris.HCl, 0.25 mM EDTA, pH 7.5) at 2 ng/μL and microinjected into the pronucleus of fertilized FVB/N oocytes according to standard protocols [45]. Microinjected oocytes were transferred into the uteri of pseudopregnant NMRI females and embryos were recovered at E10.5 or E11.5 and stained for β-galactosidase activity.

RT-qPCR

To assess recombination efficiency, the posterior part of the E10.5 *Tgfbr1-cKO* embryos, including their hindlimbs, were dissected in PBS (137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, and 1.8 mM KH_2PO_4) on ice. Whole E10.5 *Tgfbr1$^{-/-}$* embryos were used as controls. RNA was extracted from fresh tissue using TRI reagent (Sigma, #T9424) according to manufacturer's protocol. 200 ng of RNA from each sample was used to prepare complementary DNA using a random hexamer mix (NZYTech #MB12901) and following the protocol of the NZY Reverse Transcriptase enzyme (NZYTech #MB12401). All cDNA samples were diluted 1:5 and 1 ml was used in quantitative PCR (qPCR) reactions with iTaq Universal SYBR Green Supermix (Bio-Rad, #1725124), according to manufacturer's instruction and using the Applied Biosystems QuantStudio 7 flex qPCR System. Primers were localized in exon 3 and exon 4 of *Tgfbr1* mRNA (listed in supplementary table 2), so that in case of successful recombination the product is not amplified.

PCR product concentration in each sample was calculated using the standard curve with following formula *Quantity = 10^(mean Ct-b/m)* (where "m" is slope and "b" is an intercept of the standard curve), and *Tgfbr1* expression level was normalized to *b-Actin* expression. Linear model was fitted (Tgfbr1 normalized expression ~ Genotype) with R built-in function (lm). Analysis of variance (ANOVA) was performed on the linear

model with the built-in function in R (anova). Significance levels between groups of

samples were assessed by Tukey method using the "glht" function from "multicomp"

package in R. Differences were considered significant at *p<0.05, **p<0.01 and

***p<0.001. Boxplot was generated with "ggplot2" and "ggdignif" packages in R.

Cell culture and transfection

HEK 293T cells (ATCC #CRL-3216) were cultured in Dulbecco's modified Eagle's

medium (DMEM, Life Technologies) supplemented with 10% fetal bovine serum (FBS,

Life Technologies) and 1% penicillin/streptomycin at 37 °C in a 5% CO_2 atmosphere.

Cells were seeded in 35 mm plates and when they reached 70% confluency, the

medium was changed to transfection medium (DMEM supplemented with 10% FBS).

Expression vectors were constructed by amplifying the coding region of the *Tgfbr1*

transcript isolated from wild type or *Tgfbr1^3ex3-flox^* homozygous embryos by RT-PCR

(primers in supplementary table 2) and cloned into the pRK5 expression vector.

Constructs were verified by sequencing. 3 μg of the plasmids were transfected into

the 393T cells using Lipofectamine 2000 (Thermo Fisher Scientific #11668027)

according to the manufacturer's instructions. After 24 hours incubation, cells were

processed for western blot analysis

Protein extraction and western blot

The transfected cells were washed twice with ice-cold PBS and then 150 μL of lysis

buffer (50 mM Tris-HCl, 150 mM NaCl and 1% Triton X-100, pH8.0) was added to

each plate and incubated on ice for 10 minutes. Lysates were scraped into

microcentrifuge tubes on ice and centrifuged at 20000 rcf for 10 min at 4˚C. The

supernatant was collected, frozen on dry ice and stored at -80˚C. Protein lysates were

mixed (2:1) with 3X loading buffer (150 mM Tris.HCl, pH 6.8, 6% SDS, 30% glycerol,

7.5% β-mercaptoethanol, 0.03% bromophenol blue). Samples were incubated for 15

minutes at 65˚C and resolved by SDS-PAGE in a 10% polyacrylamide gel. Proteins

were transferred to PVDF membranes in 20% methanol, 25 mM Tris, 200 mM glycine

at 200 mA for 1 hour. The membranes were blocked in blocking solution [5% dry milk

dissolved in PBS containing 0.1% Tween-20 (PBT)] for 1 hour at room temperature

and then incubated with primary antibodies overnight at 4˚C. Primary antibodies were

anti-Tgfbr1 (Sigma-Aldrich #HPA056473, 1:1000 in blocking solution) and anti-actin

(Abcam #ab179467, 1:1000 in blocking solution). Membranes were then washed in

PBT and incubated with HRP conjugated-anti-rabbit IgG (GE Healthcare # NA9340,

1:5000 in blocking solution) at room temperature for 1 hour and washed twice in PBT.

Signals were developed by chemiluminescence using Amersham ECL Prime Western

Blotting Detection Reagent (GE Healthcare #RPN2232) and images were captured

using the GE Amersham Imager 600.

Skeletal preparation

E17,5 fetuses were recovered from pregnant females by cesarean section and

dissected from the extraembryonic membranes, eviscerated, the skin removed and

then fixed in 100% ethanol for 2 days. Cartilages are then stained by incubation with

450 mg/l of alcian blue (Sigma, #A5268) in 80% ethanol/20% acetic acid for one day.

Fetuses were then postfixed in 100% ethanol overnight. Tissue was cleared by

incubation in 2% KOH for 6 hours, followed by staining of ossified bones with a 50 mg/l

of alizarin red S (Sigma, #130-22-3) solution in 2% KOH for 3 hours. Specimens were

then further incubated in 2% KOH until tissues were fully cleared. Skeletons were

stored in 25% glycerol in water. All incubations were performed at room temperature

with rolling. Genotyping was performed on gut tissue by incubating in Laird's buffer

(100 mM Tris-HCl pH 8.5, 5 mM EDTA, 0.2% SDS, 200 mM NaCl) containing 10 μg/ml

proteinase K at 55ºC overnight. Genomic DNA was then recovered by precipitation

with isopropanol (1:1, vol:vol) and transferred to TE pH 8.0. Genotyping was then

performed by PCR using oligos specified in supplementary table 2.

Optical projection tomography

Optical projection tomography (OPT) was used to image E16.5 fetuses as previously

described [46] with minor modifications. Briefly, fetuses were recovered by cesarean

section in ice cold PBS, washed several times in PBS and fixed in 4%

paraformaldehyde (PFA) in PBS at 4ºC for several days. After several washes with

demineralized water fetuses were dehydrated by sequential incubation in

demineralized water with increasing concentrations of methanol (10% increases) and

then twice in 100% methanol. Fetuses were then bleached in a three day process with

a sequence of 2.5%, 5% and 10% hydrogen peroxide in methanol at room

temperature. Fetuses were then rehydrated through a reverse

methanol/demineralized water series, and embedded in a 0,7% low-melting agarose. Clearing was done with a 1:2 solution of Benzoic Alcohol:Benzyl Benzoate (BABB) using a BABB/methanol series (25% BABB increases). 100% BABBwas introduced on day 3 and replaced every day for the next 4 days, until the fetuses became completely transparent. Anatomical datasets were then obtained using a custom built OPT scanner and procedures; briefly, green auto-fluorescence was acquired on 1600 sequential angles for a full revolution, the raw dataset was pre-processed with a custom-built ImageJ macro, and then back-projection reconstructed using SkyScan's nrecon as in [47], and then post-processed to reduce noise and enhance contrast in ImageJ. 3D reconstruction and visualization and manual segmentation of the limb skeleton and internal organs was performed using the Amira software (Thermo Fisher)

Whole mount in situ hybridization and sectioning

Embryos were fixed in 4% PFA overnight, then dehydrated through 25%, 50% and 75% series of methanol in PBT (PBS, 0,1% Tween 20), then incubated in 100% methanol. Embryos were then rehydrated through a reverse methanol/PBT series and incubated three times in PBT for at least 5 minutes each at room temperature. Embryos were then bleached for 1 hour in 6% hydrogen peroxide in PBT and permeabilized in 10 μg/mL of proteinase K in PBT for a time period that depended on the embryo size. The reaction was then quenched with a 2 mg/mL solution of glycine in PBT, washed twice in PBT and postfixed in a 4% PFA and 0,2% glutaraldehyde mix for 20 minutes, followed by two washes in PBT. Hybridization was performed at 65°C overnight in hybridization solution (50% formamide, 1.3 x SSC pH 5.5 [20 x SSC is 3M NaCl, 300 mM sodium citrate], 5 mM EDTA, 0.2 % Tween 20, 50 µg/mL yeast tRNA, 100 µg/mL heparin) containing the relevant digoxigenin-labelled antisense RNA probes. The probes used in this work are listed in supplementary table 2. RNA probes were transcribed in vitro from the vector for 1.5 hours at 37°C with relevant RNA polymerase and the DIG RNA Labeling Mix (Roche #11277073910). The reaction product was verified in 0,8% agarose gel, precipitated with ethanol in the presence of 0.3M sodium acetate, pH 5.3 and resuspended in 100 μL of TE pH 8.0. Probes were diluted in hybridization solution (6 μL in 1 ml of solution). After hybridization, embryos were washed twice with hybridization solution without tRNA, heparin, at 65°C for 30 min each, then in a 1:1 mix of hybridization solution and TBST (25 mM Tris.HCl, pH

8.0, 140 mM NaCl, 2.7 mM KCl, 0.1% Tween 20) for 30 minutes at 65ºC, and finally

washed three times with TBST at room temperature. Embryos were then equilibrated

in MABT (100 mM Maleic acid, 150 mM NaCl, 0.1 % Tween-20, pH 7.5) and blocked

in MABT blocking buffer [MABT containing 1% blocking reagent (Roche

#11096176001)] with 10% sheep serum for 2,5 hours at room temperature. Embryos

were then incubated overnight at 4ºC with a 1:2000 dilution of alkaline phosphatase-

conjugated anti-digoxigenin antibody (Roche #11093274910) in MABT blocking buffer

with 1% sheep serum. After extensive washes with MABT at room temperature,

embryos were equilibrated in NTMT buffer (100 mM Tris HCl, pH 9.5, 50 mM $MgCl_2$,

100 mM NaCl, 0.1% Tween-20) and developed with a 1:50 dilution of a NBT/BCIP

solution (Roche #11681451001) in NTMT at room temperature in the dark. The

reaction was stopped with PBT, and embryos postfixed with 4% PFA overnight at room

temperature, and stored in PBT. At least 3 embryos were used per probe. Pictures

were taken with a SWIFTCAM SC1803 connected to a stereoscope. Stained embryos

were embedded in a mix containing 0,45% gelatin, 27% bovine serum albumin (BSA),

18% sucrose, jellified with 1,75% glutaraldehyde and sectioned at 35 μm on a Leica

Vibratome VT 1000 S.

β-galactosidase staining staining

Embryos were dissected out in ice cold PBS and fixed in 4% PFA at 4ºC for 30

minutes. They were then washed three times in β-gal wash buffer (PBS plus 0.02%

Tween 20) for 10 minutes at room temperature and β-galactosidase activity developed

in PBS containing 5 mM $K_3Fe(CN)_6$, 5 mM $K_4Fe(CN)_6$, 2 mM $MgCl_2$, 0.02% Tween 20,

0.4 mg/ml X-gal (Promega #V3941) overnight in the dark at 37ºC. The reaction was

stopped with β-gal wash buffer, embryos postfixed in 4% PFA overnight at room

temperature and stored in PBS.

ATAC-Seq

ATAC-seq was performed as in [25] with minor modifications. E11.25 mouse embryos

were dissected on ice in DMEM High Glucose medium (Biowest #L0102-500)

containing 10% FBS (this will be referred to as medium in the rest of the protocol).

Hindlimbs and genital tubercles were dissected out and collected on ice-cold media.

Single cell suspension was prepared by treating the tissue with 500 μL of Accutase

(Sigma #A6964-500ML) for 30 minutes at 37˚C with shaking at 600 rpm. Single cells

were pelleted at 6000 rcf for 5 minutes at 4˚C, resuspended in 200 μL of media and

counted. 50000 viable cells from each sample were used for nuclei extraction. Cells

were incubated with 50 μL of ATAC resuspension buffer (10 mM Tris HCl pH 7.4, 10

mM NaCl, 3 mM MgCl$_2$, 0.1% NP-40, 0.1% Tween-20 (Sigma-Aldrich, P7949), 0.01%

Digitonin (Target Mol #282T2721-1ml/10mM in DMSO) for 3 minutes on ice. Lysis was

quenched by adding 1 ml of ATAC resuspension buffer without NP-40 and Digitonin.

Nuclei were pelleted by centrifugation at 500 rcf for 10 minutes at 4˚C. The pellet was

then resuspended in 50 μL of transposition buffer [for 50 μL: 25 μL 2x TD buffer, 2.5

μL Tn5 transposition enzyme (100 nM final) (Illumina #15028212), 16.5 μL PBS, 0.5

μL 1% digitonin, 0.5 μL 10% Tween-20, 5 μL H$_2$O] and incubated at 37˚C for 30

minutes with shaking at 1000 rpm. The DNA was purified with Qiagen MinElute PCR

purification kit (Qiagen #50928004) and eluted in 20 μL of the kit's elution buffer.

Libraries were amplified by PCR with NEBNext High-Fidelity 2X PCR Master Mix (New

England Biolabs #174M0541S) for 9 cycles in 96-well Thermal cycler (Biorad) and

purified with Qiagen MinElute PCR purification kit. Tagmentation efficiency was

assessed on TapeStation 4200 (Agilent). Double size selection to remove primer

dimers and fragments exceeding 1 kb was performed using SPRIselect beads

(Beckman Coulter). Another quality control was performed with High Sensitivity DNA

assay using the Fragment Analyzer (Agilent). The 4nM libraries pool was sequenced

with Illumina NextSeq 2000 (100 cycles, Pair-end 50 bp). Two biological replicates

were performed per tissue.

The data obtained in this work has been deposited in the GEO repository, (accession

number GSE231592).

ATAC-Seq bioinformatic and statistical analysis

Data bioinformatic analysis was performed on the Galaxy server [48]. Raw sequencing

fastq files for each library were assessed for quality, adapter content and duplication

rates with FastQC (v0.11.9)

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc). Adapters were trimmed

and reads with length < 20 bp were removed using cutadapt (v1.16.5)[49] (3´ adapter

sequence: CTGTCTCTTATACACATCT). The trimmed reads were aligned to the

mouse reference genome (GRCm38/mm10 Dec. 2011) using Bowtie2 (v2.4.2)[50]

[parameters paired-end options (-X/–maxins 1000, --fr, --dovetail), --very-sensitive].

The aligned reads were filtered using BamTools Filter (v2.4.1) [51] (parameters --

isProperPair true, --mapQuality ≥ 30, and --reference!=chrM) and the duplicate reads

were removed using Picard MarkDuplicates (v2.18.2.2)

(http://broadinstitute.github.io/picard/). The resulting BAM files were converted to BED

and the reads that overlap to the blacklisted regions listed in

https://www.encodeproject.org/files/ENCFF547MET/ were removed using Bedtools

(v2.30.0) [52]. The filtered BED files were used as inputs for peak calling. Peaks were

called from individual replicates using MACS2 callpeak (v2.1.1) [53] (parameters --

format single-end BED, --nomodel, --extsize 200, --shift -100, --qvalue 0.05).

BAM files were then converted to BigWig with "deeptools" [54] normalizing to 1x effective

genome size (2652783500 for GRCm38/mm10). BigWig files were used for data

visualization in IGV with the addition of the conservation scoring by phyloP

(phylogenetic p-values)[55] for 60 vertebrate genomes from the UCSC genome browser

(mm10.60way.phyloP60way.bw file downloaded from

http://hgdownload.cse.ucsc.edu/goldenpath/mm10/phyloP60way/) and ChIP-seq data

from Hoxa13 (GSE81356) [27] and Gli3 (GSE133710) [28].

The raw count matrix was created by merging peaks across samples as in [56,57].

DESeq2 object was built from the raw count matrix and blind dispersion estimated with

the "vst" function from the "DESeq2" package in R [58]. Top 2500 most variable peaks

estimated with the "rowVars" function in R were selected for the principal component

analysis (PCA) and sample distance matrix. PCA was performed using the built-in

"prcomp" function in R. Sample distance matrix was made using the built-in "dist"

function in R. For statistical analysis raw reads were analyzed using EdgeR [59].

ANOVA-like test was performed on all samples and pairwise comparisons between

the samples were made by analyzing individual contrasts. Venn diagrams were

created from the lists of selected differentially accessible peaks with "eulerr" package

in R. Heat Maps were made from dataframes containing log2 normalized counts

across samples with the "pheatmap" package (https://CRAN.R-

project.org/package=pheatmap.)

Footprinting analysis of ATAC-Seq data

We further used the program RGT HINT-ATAC [43] and motif matching to identify, plot

and compare transcription factors (TF) footprints in the different samples. In detail, the

BAM files containing the filtered aligned reads for each biological replicate were

merged and used as a matrix for the footprinting analysis for the regions corresponding

to the peaks called by MACS2 (BED file) (described above). A subset of the BAM and

BED files corresponding to the regions that were found accessible only in GT (pattern

1 peaks) were used for the footprint analysis and then tested for motif matching to the

Hocomoco database[60] for finding the motif-predicted binding sites (mpbs). Motif

enrichment analysis was performed on the footprinting results for the peaks of pattern

1, using as a background the footprinting results from all the peaks called in GT.

Additional references

45.   Hogan, B., Beddington, R., Constantini, F. & Lacy, E. *Manipulating the mouse embryo: a*
laboratory manual. (Cold Spring Harbour Laboratory Press, 1994).
46.   Dias, A., Martins, G. G., Lopes, A. & Mallo, M. Three and Four-Dimensional
Visualization and Analysis Approaches to Study Vertebrate Axial Elongation and
Segmentation. *Journal of Visualized Experiments* **168**, e62086 (2021).
47.   Martins, G. G. *et al.* Optical projection tomography. in *Imaging Modalities for*
Biological and Preclinical Research: A Compendium, Volume 1: Part I: Ex vivo biological
imaging (eds. Walter, A., Mannheim, J. G. & Caruana, C. J.) i.2.f1-i.2.f11 (IOP
Publishing, 2021). doi:10.1088/978-0-7503-3059-6ch12.
48.   Afgan, E. *et al.* The Galaxy platform for accessible, reproducible and collaborative
biomedical analyses: 2016 update. *Nucleic Acids Res* **44**, W3–W10 (2016).
49.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet J* **17**, 10–12 (2011).
50.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**, 357–359 (2012).
51.   Barnett, D. W., Garrison, E. K., Quinlan, A. R., Stïrmberg, M. P. & Marth, G. T. Bamtools:
A C++ API and toolkit for analyzing and managing BAM files. *Bioinformatics* **27**, 1691–
1692 (2011).
52.   Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).
53.   Feng, J., Liu, T., Qin, B., Zhang, Y. & Liu, X. S. Identifying ChIP-seq enrichment using
MACS. *Nat Protoc* **7**, 1728–1740 (2012).
54.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res* **44**, W160–W165 (2016).
55.   Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral
substitution rates on mammalian phylogenies. *Genome Res* **20**, 110–121 (2010).
56.   Rausch, T., Hsi-Yang Fritz, M., Korbel, J. O. & Benes, V. Alfred: Interactive multi-sample
BAM alignment statistics, feature counting and feature annotation for long- and short-
read sequencing. *Bioinformatics* **35**, 2489–2491 (2019).
57.   Erarslan-Uysal, B. *et al.* Chromatin accessibility landscape of pediatric T-lymphoblastic
leukemia and human T-cell precursors. *EMBO Mol Med* **12**, e12104 (2020).
58.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

59.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–
140 (2009).
60.   Kulakovskiy, I. V. *et al.* HOCOMOCO: Towards a complete collection of transcription
factor binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic*
Acids Res **46**, D252–D259 (2018).
61.   Jones, C. M., Lyons, K. M. & Hogan, B. L. M. Involvement of Bone Morphogenetic
Protein-4 (BMP-4) and Vgr-1 in morphogenesis and neurogenesis in the mouse.
Development **111**, 531–542 (1991).
62.   Davis, C. A. & Joyner, A. L. Expression patterns of the homeo box-containing genes and
the proto-oncogene int-1 diverge during mouse development. *Genes Dev* **2**, 1736–
1744 (1988).
63.   Crossley, P. H. & Martin, G. R. The mouse Fgf8 gene encodes a family of polypeptides
and is expressed in regions that direct outgrowth and patterning in the developing
embryo. *Development* **121**, 439–451 (1995).
64.   Srivastava, D. *et al.* Regulation of cardiac mesodermal and neural crest development by
the bHLH transcription factor, dHAND. *Nat Genet* **16**, 154–160 (1997).
65.   Aires, R. *et al.* Tail Bud Progenitor Activity Relies on a Network Comprising Gdf11,
Lin28, and Hox13 Genes. *Dev Cell* **48**, 383–395 (2019).
66.   Szeto, D. P. *et al.* Role of the Bicoid-related homeodomain factor Pitx1 in specifying
hindlimb morphogenesis and pituitary development. *Genes Dev* **13**, 484–494 (1999).
67.   Echelard, Y. *et al.* Sonic hedgehog, a member of a family of putative signaling
molecules, is implicated in the regulation of CNS polarity. *Cell* **75**, 1417–1430 (1993).
68.   Chapman, D. L. *et al.* Expression of the T-box family genes, Tbx1-Tbx5, during early
mouse development. *Developmental Dynamics* **206**, 379–390 (1996).
69.   Yamaguchi, T. P., Bradley, A., McMahon, A. P. & Jones, S. A Wnt5a pathway underlies
outgrowth of multiple structures in the vertebrate embryo. *Development* **126**, 1211–
1223 (1999).

Supplementary table 2. Oligonucleotides used in this work.

Genotyping primers		
Tgfbr1-flox	Forward	TTGAGCTTGCTGTCTGACTGGATAG
	Reverse	CTGTGGTTGGCAGGCATGTG
Cre	Forward	CGAGTGATGAGGTTCGCAAG
	Reverse	CCTGATCCTGGCAATTTCGGCT
Tgfbr1-null	Forward	CTACTGTGTTTCAAATGGGAGGGC
	Reverse	GGCCTGTCGGATCCTATCATC
β-galactosidase	Forward	AGCAGTTTTTCCAGTTCCGTTTATC
	Reverse	AGCGGCGTCAGCAGTTGTTTTTTAT
Cloning of regulatory elements		
CR-Isl1	Forward	GACTCGAGTCTGTGATACAAAACAATATATC
	Reverse	GAGGATCCAATCTTTCAAAGACATGGAGGG
CR-Tbx5	Forward	GACTCGAGCTTTATGTATCTGAGCACACTG

	Reverse	GAAGATCTTCCTTCAACAAACCAT CCAC
CR-Fgf10	Forward	CAGCTCGAGACTAGTAATCTGAGGTGGTTGCTCACTC
	Reverse	CATCCATGGCTATGGAAGGCTTATGTATCAC
Grem1-Enh	Forward	GACGTCGACACTAGTGTCATGTGTTCCTGTGATTCTG
	Reverse	GAGCCTGTATAAGAAGTTCAGGGCCA TGGCAG
Other primers		
qPCR-Tgfbr1	Forward	ACCGTGTGCCAAATGAAGAGG
	Reverse	CATCTAGATCTTGTAACACAATGGTCCTGGC
qPCR-Actin	Forward	ATGAAGATCCTGACCGAGCG
	Reverse	TACTTGCGCTCAGGAGGAGC
Tgfbr1-CDS	Forward	GAGAATTCCGGGCCACAAACAGTGGC
	Reverse	GAGTCGACCATTTTGATGCCTTCCTGT TGGC
Oligonucleotides for generating in situ probes		
Lmx1b	Forward	CTGCTGTGCAAGGGTGACTATGAG
	Reverse	GAGGCAAAGTAGGAGCTCTGCATG
Fgf10	Forward	TTCCTAGAAGTTATGGATGTTG
	Reverse	GTACTGCATCCACCAACAGTG
Grem1	Forward	AATGAATCGCACCGCATACAC
	Reverse	GACTAATACGACTCACTATAGGGAAGCAACTGCTGGTTCTTCTG
In situ probes		
Bmp4	Jones et al 1991, ref [61]	
En1	Davis and Joyner, 1988, ref [62]	
Fgf10	Whole coding region cloned into TOPO. Oligonucleotides listed above	
Fgf8	Crossley and Martin, 1995, ref [63]	
Grem1	PCR fragment with the whole coding region and T7 promoter at the 3' end. Oligonucleotides listed above	
Hand2	Srivastava et al, 1997, ref [64]	
Isl1	Jurberg et al 2013, ref [6]	
Lin28a	Aires et al., 2019, ref [65]	
Lmx1b	PCR fragment encompassing nucleotides 504 to 1202 of the mRNA cloned into pKS bluescript. Oligonucleotides listed above.	
Pitx1	Szeto et al, 1999, ref [66]	
Shh	Echelard et al, 1993, ref [67]	
Tbx5	Chapman et al, 1996, ref [68]	
Wnt5a	Yamaguchi et al, 1999, ref [69]	

931 **Supplementary table 3.** Coordinates of the genomic elements used in the reporter

932 assays (from the GRCm38/mm10 reference genome).

Element	Coordinates
CR-Isl1	chr13:116,284,024-116,286,017
CR-Tbx5	chr5:119,700,950-119,703,188
CR-Fgf10	chr14:28,412,202-28,414,123
Grem1 enhancer	chr2:113,580,726-113,582,264